

ACIDIFICATION OF LAKES IN CANADA BY ACID
PRECIPITATION AND THE RESULTING
EFFECTS ON FISHES

RICHARD J. BEAMISH, Pacific Biological Station Fisheries and Marine Service, Department of the Environment, P.O. Box 100, Nanaimo, British Columbia, V9R 5K6.

ABSTRACT

There are several areas in Canada that could receive acidic fallout. Only in the area that produces Canada's largest single source of sulfur oxides has there been appreciable research. In the Sudbury region of Ontario, Canada, fallout of sulfur oxides has been shown to be responsible for damage to vegetation, lakes and fishes.

The acidic fallout has been shown to effect a rate of acidification in many lakes that over several decades has resulted in the extinction of many species of fishes. Fishes exhibit profound differences in acid tolerance but show some similarities in their physiological response to levels within the range of their individual susceptibilities. Prior to extinction most females of a particular species did not release their ova to be fertilized. The failure of females to spawn was coincident with an inability to maintain normal serum calcium levels. In some species growth was reduced despite an adequate supply of preferred food items. High concentrations of acid were considered to be the principle factor stressing the fish populations. Elevated concentrations of some heavy metals may add to the stress caused by high concentrations of acid.

As a result of damage in one area to trees, lakes and fishes by the fallout of sulfur oxides the Canadian Government started court proceedings against the companies alleged to be responsible for the pollution. After 4 years the suit was settled out of court just days before the trial was scheduled to commence.

ACID PRECIPITATION IN CANADA

It is the intention of this paper to discuss the effects of acid precipitation on Canadian lakes and fishes. However, it is only in the Sudbury, Ontario area that such effects have been well documented. The principle source of acid entering the Sudbury lakes comes from local nickel mining and smelting industries. In 1969 it was estimated that 2.6 million metric tons of sulfur oxide were released into the atmosphere from the Sudbury area (Beamish and Harvey, 1972). In 1970 the emissions were estimated to be equivalent to half of Canada's total emissions (EPS, 1973) or about 10% of the total United States of America emissions (Beamish, 1974a). In fact the Sudbury area may have the dubious distinction of being the largest single source of sulfur dioxide in the world. The second largest point source (about 700,000 metric tons) in Canada is at Noranda, Quebec about 138 km north of Sudbury (EPS, 1973) (Figure 1). Excluding the major cities of Toronto




Figure 1. Summary of principle sources of sulfur oxide emissions in Canada (from EPS, 1973).

and Montreal and the important emission areas of Sudbury and Noranda there are only four or five other point sources ranging in annual emissions of 200,000 to 400,000 metric tons that can be considered to be potential sources of acid precipitation to nearby lakes. In one of these areas, near Flin Flon, Manitoba a survey in 1973 of 27 lakes in the vicinity of the smelter revealed only one lake with a pH below 6.5 (Van Loon and Beamish, 1976). The one acid lake (pH 4.0-5.0) received acid effluent in the spring from a holding area. At present there is no other information that identifies other point sources of sulfur dioxide in Canada as causing the acidification of lakes.

Similarly there is no published study at present that identifies long-range transportation of acid (from many point sources) as affecting Canadian lakes. Several studies have reported that fallout of acid has occurred and with hindsight long-range transport of acid may have been involved. On April 9, 1970 Barica and Armstrong (1971) recorded a snow at the Experimental Lakes Area (E.L.A.), in northwestern Ontario that contained excessive particulate material and sulfate concentrations five to 10 times greater than measured in previous samples. Since there are no major industrial areas in the vicinity of E.L.A. it appears that this fallout could have originated somewhere in the north central United States. Precipitation may also be the principle source of nutrients to the E.L.A. Lakes (Schindler, 1971). In Halifax County, Nova Scotia acid lakes were found (Gorham, 1957) in which the acidity could not be completely accounted for by natural processes. Gorham felt that rain was a major source of SO_4 for some of these lake waters as "suggested by recent analyses of Herman (unpublished), who found an average of 0.96 ppm total sulfur, equivalent to 2.9 ppm SO_4 , in rain collected over a 2-year period at the Dominion Experimental Station, Kentville, Nova Scotia." This is one of the earliest records of acid precipitation in Canada and it is possible that the source of this acid was either from the Halifax area or from combined areas on the east coast of North America.

ACID PRECIPITATION IN THE SUDBURY AREA

The effects of acid fallout on plants and soils in the Sudbury area have been demonstrated in numerous studies over the past 30 years (McCallum, 1944; Linzon, 1958, 1966, 1971; Gorham and Gordon, 1960a, 1960b; McGovern and Basillie, 1972, 1973; Whitby and Hutchinson, 1974; Le Blanc et al., 1972; Stokes et al., 1973; and others). However, the effects of Sudbury emissions on lakes have not been as extensively studied. Studies undertaken in the early 1960's did show a deterioration in water quality of lakes and ponds in the immediate Sudbury area (Gorham and Gordon, 1960a, 1963) and recently lakes located in the La Cloche Mountains some 65 km southwest of the smelters have been found to be extremely acidic (Beamish and Harvey, 1972).

There is good evidence that indicates the Sudbury smelters are the principle source of acid entering the La Cloche Mountain lakes. As previously mentioned it is known that extremely large quantities of SO_2 are released annually into the atmosphere around Sudbury. A summary of the average percentage of wind hours as recorded at Sudbury, Ontario from 1961 to 1973 (Department of Transport, monthly record of meteorological observations in Canada) indicates the prevailing wind directions are northeast and southwest (Figure 2a). By comparing the mean ground




Figure 2a. Average percentage of time wind blew in each of 16 compass directions. Length of arrows indicates the average percentage of wind-hours that were recorded for each direction from 1961 to 1972. Data obtained from Department of Transport, Monthly meteorological observations in Canada.

sulfur dioxide concentrations from 1964 to 1968 (Figure 2b redrawn from Dreisinger and McGovern, 1970) it is apparent that sulfur dioxide falls out in decreasing concentrations from the smelters in the direction of the prevailing winds. A similar fallout pattern occurs for nickel, iron and copper (McGovern and Basillie, 1973; Kramer, 1973a). It is also known that the Sudbury smelters release large quantities of heavy metals into the atmosphere. Stack emission reports prepared by one of the companies indicate that approximately 14,500 tons of iron, 2,000 tons of nickel and 1,800 tons of copper have been released annually into the atmosphere over the last decade (Falkowski, 1973).

2B




Figure 2b. Average ground concentrations of sulfur dioxide in the Sudbury area from 1964 to 1968 (Figure reproduced from Dreisinger and McGovern, 1970).

A study of the dispersion of the Sudbury smelter plumes showed the plumes from the four main smelters combined to form a plume approximately 25 km wide and contained sulfur dioxide concentrations ranging from 0.3 to 0.02 ppm at a distance of 40 km from Sudbury (Whaley and Lee, 1971). Whaley and Lee (1974) used the plume dispersion study to show that high ground-level sulfur dioxide concentrations measured at a site 36 km southwest of the Sudbury smelters during fumigation conditions could be attributed to Sudbury smelter emissions. These authors computed ground-level sulfur dioxide levels using the plume dispersion study and Sudbury smelter production figures, and were able to obtain a good correlation between observed and predicted ground-level sulfur dioxide concentration over the period 1961-1971.

In the immediate Sudbury area the pH of precipitation has been reported to average about 4.5 with an extreme low of 3.2 (Kramer, 1973a). In the La Cloche Mountains, the average pH of 18 precipitation events from 1972 to 1973 was 4.3 (Beamish and Van Loon, 1975) and the average pH of monthly precipitation collections for this period was 4.4. The lowest pH measured was 2.9 (Beamish and Harvey, 1972). As mentioned, precipitation in the Sudbury area also contains high concentrations of nickel and copper (Beamish and Van Loon, 1975; Kramer, 1973b). Kramer (1973b) claimed that most of northern Ontario has been subjected to rates of fall of nickel that are 10 times the rate observed in the Continental United States. The consistent observation that substances or byproducts of substances known to be emitted from the smelters, occur in precipitation; the evidence that concentrations decrease with distance from the smelters in the direction of the prevailing winds; the

calculations showing that the plume disperses over great distances and ground-level concentrations can be related to concentrations of sulfur dioxide in the plume; the effective "labelling" of this plume by the high levels of nickel emitted by the industries and the anomalously high nickel concentrations in precipitation leave little doubt that fallout of substances from the plume is occurring in the Sudbury area.

It can be shown that this fallout has altered the water quality of lakes in the Sudbury area. Harvey (1975a) has measured the sulfate concentration of over 100 lakes and found concentrations were highest close to the smelters and decreased with distance from the smelters. Sulfate ions were partially balanced to hydrogen ions indicating that part of the sulfate that was entering the lake from the atmosphere was in the form of sulfuric acid. A survey of 150 lakes in the La Cloche Mountain area indicated that some 33 lakes had a pH of 4.5 or less and one-half of the lakes had a pH of 5.5 or lower (Beamish and Harvey, 1972). From a study of the pH change in one lake over a period of 5 years it was shown that the lake was acidifying at a rate of approximately 0.13 pH units per year (Beamish et al., 1975a). This compares with estimated average decrease of 0.16 pH units for 22 La Cloche Mountain Lakes for which determinations measured in 1971 could be compared to readings taken 10 years earlier (Beamish and Harvey, 1972).

A detailed examination of the precipitation loading of acid to a small lake in the La Cloche Mountains from 1972 to 1973 showed that this 21 ha lake received approximately 2100 kg of sulfuric acid from precipitation in 1972, that would be capable of depressing the pH below its January 1972 value (Beamish and Van Loon, 1975). A titration curve, calculated using lake water obtained during an earlier period of higher pH, predicted that the addition of this much acid would reduce the pH of the lake from 5.2 to 4.8. The actual measured change was from 5.2 to 4.7. In 1973, 600 kg H_2SO_4 were added to the lake from precipitation that would be capable of depressing the pH below the January 1973 concentration. The titration curve for the lake predicted a pH change from 4.8 to 4.74. By January 1973, the lake pH had risen from 4.7 to 4.8 and there was no net change in pH throughout the year. Thus the pH of this lake during this 2-year period was closely related to the acid content of the precipitation.

This study and others (Beamish, 1974a; Beamish et al., 1975a, 1975c) also indicated that lakes in the La Cloche Mountains contained high concentrations of nickel and copper when compared to concentrations measured in lakes remote from major areas of industrialization. The remote area used for a comparison was part of the Canadian Shield as were the La Cloche Mountain lakes (Beamish et al., 1975b).

It has been shown that the Sudbury plume was responsible for the deposition of sulfur dioxide, acid and nickel. It has also been shown that many La Cloche Mountain lakes were acid, contained elevated concentrations of nickel and probably responded to atmospheric loadings of acid as closely as the one lake studied. Therefore, there can be little

doubt that industry in Sudbury has been a major contributor to the atmospheric pollution that has altered the water quality of these lakes.

CHANGES IN WATER CHEMISTRY

Changes in water chemistry of Canadian lakes as a result of atmospheric inputs of acid have been described in only a few reports. Also it is difficult to examine changes in lake chemistry when the lake chemistry was not described before the change occurred. However, by comparing the chemistry of acidified lakes with the chemistry of unpolluted lakes from a similar geological area it is possible to gain some insight into the changes that have occurred.

As might be expected, sulfate concentrations have been shown to be high in many acidified lakes (Gorham and Gordon, 1960a; Harvey 1975a; Beamish and Van Loon, 1975). The high sulfate concentrations were found in association with low pH and by balancing anions and cations it was possible to demonstrate that part of the sulfate ion is balanced with hydrogen ion, indicating the pH results from the presence of sulfuric acid (Beamish, 1974a). A comparison of the major ions in four of the acid lakes with unpolluted lakes (Table I) (see also Beamish, 1974a) indicated that the ionic composition was similar except for hydrogen and sulfate ions. Hydrogen ions were two to three orders of magnitude greater in the acidic lake than in the E.L.A. lakes and sulfate represented approximately 90% of the anions compared to an average of about 40% for the E.L.A. lakes.

Calcium concentrations were slightly higher in the acid La Cloche Mountain lakes (average 2.6 mg/l, Table II) than the average (2.0 mg/l) for the 102 E.L.A. lakes (Tables I). Gorham and Gordon (1960) reported that calcium in ponds near Sudbury smelters appeared to decrease between pH 6 and pH 5 while below pH 5 calcium concentrations appeared to increase. There is some historic data of unknown analytical quality for the La Cloche lakes that suggests the calcium concentrations were at least double their present values. The historic data is to some extent corroborated by the observation that the less acidic Kakakise Lake (Table II) has a calcium concentration almost double the values of the nearby acid lakes. Since calcium appears to be very important to the survival of fishes (Beamish et al., 1975) more information concerning the effect of acidification on the calcium content of lakes may be necessary for a thorough understanding of the response of fishes to acidification.

It is apparent that bicarbonate alkalinity (as mg/l CaCO_3) will be greatly reduced in any lake of low pH. Alkalinity of the acidic La Cloche Mountain Lakes was less than 0.5 mg/l which was at the limit of detection. By comparison alkalinity of the E.L.A. lake averaged 11.4 mg/l (Table I).

Table I. Chemical composition of four acid lakes and non-polluted lakes in a remote area of N.W. Ontario. For details of water chemistry see references.
Range in brackets.

	O.S.A. ^a	Muriel ^b	George ^{cc}	Lumsden ^{ccc}	102 lakes E.L.A. N.W. Ontario ^{****}
pH	4.5 (4.4-4.9)	4.7 (4.5-4.8)	5.0 (4.8-5.3)	4.8 (4.7-5.3)	6.5
Sulfate mg/l	15 (14-17)	15 (14-15)	14.3 (12-16)	10.7 (10.0-12.6)	3.8
Chloride mg/l	1 (<1-1)	<1 (<1-1)	1.0 (0.6-1.6)	0.7 (0.5-1.2)	0.8
Sodium mg/l	0.6 (0.6-0.8)	0.6 (0.6-0.7)	0.8 (0.7-1.0)	0.4 (0.4-0.8)	0.9
Calcium mg/l	3.2 (2.9-3.7)	3.4 (3.1-3.4)	3.2 (1.8-4.0)	2.2 (1.4-2.9)	2.0
Magnesium mg/l	0.9 (0.7-1.0)	0.9 (0.7-0.9)	1.0 (0.9-1.6)	0.7 (0.5-1.0)	0.6
Potassium mg/l	0.4 (0.3-0.5)	0.3 (0.3-0.4)	0.5 (0.3-0.6)	0.3 (0.2-0.4)	0.3
Hardness (as CaCO ₃) mg/l	11 (10-14)	11 (11-12)	12.3 (7.9-16.4)	8.4 (5.6-11.3)	7.5
Total dissolved solids mg/l	23 (7-59)	16 (15-16)	28 (10-49)	18 (10-25)	-
Total suspended solids mg/l	1 (0-2)	1 (0-1)	<1 (<1-4)	<1 (<1-4)	2
Bicarbonate alkalinity (as CaCO ₃) mg/l	0.5	<0.5	<0.5	<0.5	10
Conductivity μ mho/cm ²	48 (46-53)	32 (8-47)	40 (35-45)	32 (27-35)	19
Iron μ g/l	23 (18-29)	40 (34-46)	27 (7-45)	35 (13-77)	143
Nickel μ g/l	11 (8-13)	12 (10-15)	10 (5-15)	8 (6-9)	<3
Copper μ g/l	4 (2-6)	2 (2-4)	2 (1-8)	4 (2-16)	2
Zinc μ g/l	36 (30-48)	29 (25-31)	24 (8-45)	32 (17-35)	<1
Cadmium μ g/l	<1	<1	<0.5 (<0.5)	0.4 (0.2-0.5)	<0.1
Lead μ g/l	4 (2-6)	1 (0-3)	<1 (<1)	2 (1-3)	<1
Manganese μ g/l	-	-	220 (160-260)	258 (210-310)	3

^aBeamish, 1974a.

^{cc}Beamish et al., 1975a.

^{ccc}Beamish and Van Loon, 1975.

^{****}Beamish et al., 1975b.

Heavy metal concentrations in the La Cloche lakes appeared to be greatly altered during the acidification process. As previously mentioned nickel and to a lesser extent copper concentrations were higher in the acidic lakes than in the E.L.A. lakes. Nickel concentrations ranged from 8 to 12 μ g/l, many times higher than found in most lakes (Table I; Beamish, 1974a). Copper concentrations averaged 2 to 5 μ g/l but were only slightly higher than found in the E.L.A. lakes. As previously discussed the higher concentrations of nickel (and copper) appear to result from precipitation loadings (Beamish and Van Loon, 1975; Kramer, 1973a and b). Zinc concentrations (25-46 μ g/l) were rather normal when compared to an average concentration of 49 μ g/l obtained for over 1,500 samples from waters within the United States of America (Kopp and Kroner, 1969). They were, however, much higher than the average value for E.L.A. lakes. Zinc concentrations were similar in the acidic La Cloche Mountain lakes and lakes of similar pH in Sweden (Beamish and Van Loon, 1975; Hörnström et al., 1973). Manganese concentrations were also high in both the La Cloche lakes and Swedish lakes. While the higher levels of zinc may be partly due to precipitation loading and increasing solubility at lower pH, the higher concentrations of

Table II. Calcium concentrations (mg/l) for acidic lakes
in the La Cloche Mountains.

Lake	pH		Ca		Lake	pH	Ca
	average	(range)	average	(range)			
George	5.0	(4.8-5.3)	3.2	(0.8-4.0)	1A*	4.4	2.2
Lumsden	4.8	(4.7-5.3)	2.2	(1.4-2.9)	2A	4.8	2.2
O.S.A.	4.5	(4.4-4.9)	3.2	(2.9-3.7)	3A	4.1	2.0
Muriel	4.7	(4.5-4.8)	3.4	(3.1-3.4)	4A	4.4	2.1
Kakakisc	5.6		4.1		5A	4.5	1.1
A.Y. Jackson	4.4		2.9		6A	4.5	2.3

*Not named. See Beamish et al. 1974a for locations.

manganese appear to be the result of increased solubility (Beamish and Van Loon, 1975).

EFFECTS OF LAKE ACIDIFICATION ON FISHES

Extensive fishing operations have documented the absence of fish populations for several of the acidic La Cloche Mountain lakes (Beamish and Harvey, 1972; Beamish, 1974a; Beamish et al., 1975a). Less intensive netting operations enabled a greater number of lakes to be examined and it was found that 28 of 67 lakes in the La Cloche Mountain region had lost the majority of their fishes (Harvey, 1975b).

In Lumsden Lake (Beamish and Harvey, 1972), O.S.A. and Muriel lakes (Beamish, 1974a) and George Lake (Beamish et al., 1975a) the condition of the ovaries of female fish were examined before, during and after the lake pH was reduced to levels that affected various species. It was found that most species which had reproduced normally each year, stopped reproduction when the pH of the lake reached a certain level. The pH at which reproduction ceased varied among species. Failure to reproduce did not appear to result from egg to fry mortalities as most females did not release ova to be fertilized. Ovaries were mature prior to the normal spawning period but few, if any fish, were found with spawned ovaries after the normal spawning period. Ova became flaccid and watery and fish appeared to resorb material from ovaries. Some old fish were found in this condition (Beamish, 1974a) suggesting that the development of ova but the failure to spawn may continue in some species for many years. With the absence of recruitment, the numbers in a

population appear to decrease rapidly in response to the natural mortality rate which may be increased because of the added stress of the low pH. Mortalities due to acid stress would be expected to be related to the lake conditions, species tolerance, and to the rate of acidification. It could not be determined if some loss in production due to partial mortalities of eggs or fry occurred prior to the complete failure of females to release ova.

Because different species responded to different levels of acidification and because reproduction appeared to be the most sensitive major physiological process that was critical to the survival of the species, it was necessary to determine the approximate pH at which species ceased reproduction. The term "critically acidic" was adopted and defined to be the pH which was sufficient to inhibit reproduction of a particular species (Beamish, 1974a). A lake was then considered to be critically acidic at a pH which was sufficient to inhibit the reproduction of the most acid sensitive species in the lake. Fish populations from the acidic La Cloche Mountain lakes appeared to be affected in the pH range of 4.5 to perhaps as high as 6.0 (Beamish et al, 1975a). The most sensitive species (Table III), smallmouth bass, walleye, lake trout, were also the most desired sportfish. Smallmouth bass appeared to be the most sensitive species in the area as they had disappeared from most lakes before this study commenced in 1966. Local residents reported this species to be abundant in several lakes prior to the late 1950's. It is possible that this species ceased reproduction at pH values above 6.0. Walleye is another important sportfish that appears to be very susceptible to additions of acid. This species disappeared by the mid 1960's from most lakes in the La Cloche Mountains. Lake trout were of intermediate tolerance, becoming extinct at pH values below 5.0 and ceasing reproduction at about 5.5. However, in one lake the occasional female lake trout was observed to have spawned when the lake pH was 5.2 to 5.0. The titration curve for several dilute lakes (Beamish and Van Loon, 1975) indicates that equal additions of acid in the range of pH from about 5.5 to 5.0 cause greater decreases in pH than at other pH values. Thus fish of intermediate susceptibility, lake trout, trout perch, brown bullhead, white sucker, and northern pike (Table III) may exhibit a shortened "long-term" response to changes in lake chemistry. Cyprinids (minnows) were the most resistant species. The lake chub was observed to reproduce at pH of 4.5. Yellow perch, lake herring and rockbass also were tolerant of low pH values.

The minimum pH of 6.0 established for aquatic organisms by the National Technical Advisory Committee (1968) would probably be acceptable for most fishes. However, if smallmouth bass and walleye were desired species as they are in most areas then the acceptable pH may be at some value greater than 6.0. The pH range of 5-9 considered by the EIFAC Working Party on Water Quality (1968) not to be directly lethal for fishes should be ignored as it has been shown that in the La Cloche lakes two to four important species were extinct before pH 5.0 was reached.

Table III. Approximate pH at which fish in the La Cloche Mountain lakes stopped reproduction.

pH	Species	Family
6.0+ to 5.5	Smallmouth bass	Centrarchidae
	<u>Micropterus dolomieu</u>	
	Walleye	Percidae
	<u>Stizostedion vitreum</u>	
	Burbot	Gadidae
5.5 to 5.2	<u>Lota lota</u>	
	Lake trout	Salmonidae
	<u>Salvelinus namaycush</u>	
	Troutperch	Percopsidae
	<u>Percopsis omiscomaycus</u>	
5.2 to 4.7	Brown bullhead	Ictaluridae
	<u>Ictalurus nebulosus</u>	
	White sucker	Catostomidae
	<u>Catostomus commersoni</u>	
	Rock bass	Centrarchidae
4.7 to 4.5	<u>Ambloplites rupestris</u>	
	Lake herring	Salmonidae
	<u>Coregonus artedii</u>	
	Yellow perch	Percidae
	<u>Perca flavescens</u>	
	Lake chub	Cyprinidae
	<u>Coxesius plumbeus</u>	

The extinction of fishes did not appear to be an indirect effect caused by an earlier loss of food items. Predators such as smallmouth bass, walleye lake trout, and northern pike all became extinct while major prey species such as lake herring and yellow perch and cyprinids remained abundant. White suckers in Lumsden Lake were observed to become extinct even though approximately 10 times the biomass of preferred benthic food items was present (Beamish, 1974b). A decrease in feeding intensity was observed for white suckers held for several months in sublethal concentrations of acid (Beamish, 1972) and it is possible white suckers and other species responded to acid stress in the natural

environment in a similar manner by reducing feeding or food utilization or both. Changes in plankton composition of the La Cloche lakes have been reported (Sprules, 1975) but may not be important as the species most dependent upon plankton for food (lake herring) was one of the last species to become extinct (Beamish, 1974b).

Growth of some species did appear to be affected by the decreasing pH. Between 1966 and 1968 in Lumsden Lake the mean size of similar aged white suckers decreased as the population was becoming extinct (Beamish, 1974b). In 1972 and 1973 white suckers in George Lake were found to be significantly smaller than similar aged fish sampled earlier (Beamish et al., 1975a). Many of the remaining lake trout and northern pike in George Lake in 1972 and 1973 were observed to be emaciated and generally in poor condition. The changes in growth of these species occurred during the period that the fish first stopped reproducing. During this same period, yellow perch increased in numbers then increased in mean size (Beamish et al., 1975a). An increase in size as a lake became more acidic has also been reported for fishes from acid lakes in other areas (Almer, 1972; Schofield, 1975). Since there is a wide range of tolerance to pH among fishes one might expect there is also a wide range of pH values in which growth of a species may be altered. Therefore it is important not only to identify changes in growth but to relate these changes to other physiological changes that may be occurring such as the failure to reproduce. If species are observed to increase in mean size at a specific age and the species have ceased to reproduce then such an observation would indicate that the response of white suckers and perhaps lake trout and northern pike in the La Cloche lakes may be characteristic only of the species or of the area.

It was found that female fish from the acid lakes had abnormally low serum calcium concentrations during the period of ovarian maturation (Beamish et al., 1975a). The cause of this reduction in serum calcium of females was not determined but because changes occurred when pH was depressed to critical levels it did appear that the changes in normal calcium metabolism have resulted from the increased acid content of the lake and these changes caused or resulted from the reproductive failures. Another indication that normal calcium metabolism might be altered by the changes in water quality was found in the increased occurrence of deformed white suckers in George Lake (Beamish et al., 1975a). The occurrence of deformities in fish during periods of sublethal acid stress has also been described in laboratory acid toxicity tests (Mount, 1973; Beamish, 1972). The similar response of white suckers in an acidifying acid lake and from a controlled laboratory experiment is accepted as corroboration that this species was responding to the stress of increased acidity.

OTHER EXPLANATIONS FOR OBSERVED
RESPONSE OF FISHES

It has been assumed that the cause of the reproductive failures and the ultimate extinction of the fish populations were a result of the increasing acid content of the lake. Another possibility would be that some unknown very toxic chemical was causing the observed responses. Such a possibility always is present in research and while there appears to be a more reasonable explanation for the observed responses, the presence of some unknown agent should not be completely discounted. A more probable alternative would be that the fish were responding to heavy metal toxicity, increased in effect because of the low pH. In the La Cloche lakes only zinc concentrations were sufficiently high (20-40 $\mu\text{g}/\text{l}$) to be potentially hazardous to fishes (Canadian Department of the Environment, 1972). The observed zinc concentrations were well below incipient LC50 values considered representative for most salmonid fishes in water of a hardness similar to the La Cloche lakes (Canadian Department of the Environment, 1972). The zinc concentrations were well within the range of "safe" concentrations of 30 $\mu\text{g}/\text{l}$ which had no effect and 180 $\mu\text{g}/\text{l}$ which caused an 83% reduction in egg production of fathead minnows (*Pimephales promelas*) in hard water (Brungs, 1969). Zinc concentrations were also lower than concentrations considered to be "safe" for *Daphnia magna* (Biesinger and Christensen, 1972). Manganese and nickel concentrations were high in all acid lakes, however, they were well below concentrations that caused a minimum of reproductive impairment to *Daphnia magna* (Biesinger and Christensen, 1972). Nickel concentrations were much lower than the 380 $\mu\text{g}/\text{l}$ considered "safe" for fathead minnows in hard water (Pickering, 1974). Copper concentrations ranged from 2 to 4 $\mu\text{g}/\text{l}$ and were much lower than the value of 17 $\mu\text{g}/\text{l}$ that did not affect survival, growth or reproduction of adult brook trout (*Salvelinus fontinalis*) in soft water (McKim and Benoit, 1971). Cadmium ($< 1 \mu\text{g}/\text{l}$) was well below the range of 4.5 to 37 $\mu\text{g}/\text{l}$ that had no adverse effect on survival, growth or reproduction of fathead minnows in hard water (Pickering and Gast, 1972).

These tests of toxicity were performed at higher pH values and it is possible that the "safe" levels are too high for the acidic La Cloche Mountain lakes. Conversely heavy metal concentrations probably increased gradually as the lakes acidified, allowing fish to acclimate to the increasing concentrations. Schofield (1965) found that only brook trout that were acclimated to higher zinc concentrations could be introduced into an acidic lake with high zinc concentration. Also, if heavy metal concentrations increased as the lake acidified then the loss of the more sensitive species occurred at even lower concentrations than presently found.

In contrast to the lack of a good relationship between concentrations of heavy metals in the lakes and the concentrations observed to affect fishes in controlled laboratory experiments, there was a good correlation between pH and the observed responses of fishes. Mount (1973) showed that the survival of fathead minnows in hard water was not

affected at pH 7.5 to 4.5 but at pH 5.9 and lower, egg production and hatchability were reduced. At pH 5.2 to 4.5 females did not spawn. It was in the range of pH 5.2 to 4.7 that five species in George Lake were first observed to stop reproducing (Beamish et al., 1975a). The occurrence of decreased egg production and reduced hatchability were not studied in the acidic La Cloche Mountain lakes and may have influenced total fish production. Water quality standards allowing pH values lower than pH 6 were not considered acceptable for fathead minnows. Mount (1973) recommended that a minimum pH of 6.5 be maintained for fish populations.

In George Lake five species were observed to cease reproduction after pH dropped from 5.5 to 4.7 in the early spring of 1971, followed by a brief winter rise to 5.2, and a fall again to 4.8 in the summer of 1972 (Beamish et al., 1975a). This is an increase from .003 to .020 mg/l of hydrogen ion in 1971 and a change from .006 to .016 mg/l in 1972. It also appears that small increases in copper, nickel, and manganese may have occurred when the George Lake pH decreased from about 5.5 to 4.7 (Table IV). Zinc may have doubled in concentration from about 15 µg/l to 30 µg/l. Iron determinations were so variable that comparisons could not be made. There was no significant correlation ($p = 0.05$) between pH and copper, nickel, or manganese from February 1972 to November 1973 (Table IV). The decrease in zinc concentrations was significantly correlated ($p = 0.05$) with increasing pH. A comparison of heavy metal concentrations in George Lake with concentrations from a nearby less acidic lake (Kakakise Lake, Table IV) indicates that heavy metal concentrations may have doubled in George Lake when the pH dropped from 5.5 to 4.7. However, the more than sixfold increase in hydrogen ion that lowered the pH into a range known to prevent fish from spawning (Mount, 1973) appeared to be a more important stress than the heavy metal increases that still were below concentrations considered to be safe for fishes. Thus there were small changes in heavy metal concentrations in relation to the very large increase in hydrogen ion concentration during the time many species in the lake first ceased spawning. Finally, there is a substantial literature describing the levels of pH that are lethal to fishes (EIFAC, 1969; McKee and Wolf, 1963; Daudoroff and Katz, 1950; and others). While such literature in general does not consider the wide range of sensitivities exhibited by fishes and the effects of prolonged sublethal concentrations of acid and resulting reproductive failures, it is unanimous in its agreement that pH values below 5.0 are harmful to fishes.

It is possible that synergistic effects of acid and heavy metals, or additive effects of heavy metals and acid may cause stress at a slightly higher pH than if heavy metals were not present. A fish may be responding to an acid stress modified by heavy metal toxicity but there does not appear to be any justification for considering that the failure of reproduction and ultimate extinction of fish populations has resulted primarily from the toxic effect of heavy metals. It is important to consider that the elevated concentrations of heavy metals such as zinc,

Table IV. Heavy metal concentrations and 3 m pH values* for George and Kakakise lakes.

Date	Cu	Ni	Cd	Zn	Pb	Fe	Mn	pH
George Lake								
February 1972	<2.5	-	<1	27	1.2	41	-	-
July	3	7	nd	20	1	19	260	5.0
August	3	11	-	31	3	20	260	4.9
September	4	8	nd	26	3	19	200	5.1
October	4	9	nd	27	nd	22	200	5.0
December	2	14	nd	20	nd	12	210	5.2
January 1973	3	9	0.3	32	1	7	-	4.8
February	2	10	<.2	24	nd	27	-	5.0
April	3	5	nd	21	nd	33	210	5.0
May	3	5	nd	21	nd	33	200	5.0
June	1	9	.5	11	<2	6	-	5.0
August	<1	9	<.5	25	3	32	220	5.1
November	1	9	.3	13	2	22	240	5.3
Kakakise Lake								
November 1972	<1	5	.2	10	<2	8	150	-
February 1973	2	6	<.2	12	nd	8	190	5.7

*See Beamish et al. (1975a) for methods of chemical analysis.

- indicates no measurement; nd indicates not detected.

manganese, and perhaps other substances may occur because of increased weathering and solubility due to acidification of soils and lakes. If this is true then it is practical to think of fish populations as being stressed by an association of factors of which acid is the principle stressing agent as well as controlling the concentrations of the less important factors.

COURT ACTION

In 1970 a court action was initiated by the Federal Department of Indian Affairs in cooperation with the Department of Justice. It was alleged that the major mining and smelting companies in Sudbury had

caused damage to forests and lakes on a nearby Crown Indian Reserve as a result of their sulfur dioxide and heavy metal emissions. For the purposes of the action, special investigations were conducted of the dispersion of the plume of sulfur dioxide as it left the companies stack; the correlation of quantities of sulfur dioxide in the plume with measured ground-level concentrations, the effects of these emissions over the past few decades on forest in the Reserve; and the effects of emissions on lakes and fish populations. After discovery hearings, and subsequent trial postponements at the request of the companies, the action was settled out of court for an undisclosed sum without any liability being assessed. The settlement occurred in October 1974 just days before the trial was scheduled to commence.

Such a settlement while probably being beneficial to the Indian Reserve in that monies became available for the compensation of damages suffered, did not appear to be particularly helpful to those concerned with maintaining and improving the quality of the environment. However, because there was a settlement on the part of the companies, the action was important because it demonstrated that it is possible to provide evidence that at least can encourage major polluters to control the emissions of pollutants into the atmosphere.

REFERENCES

- Almer, B. 1972. Försurningens Inverkan på Fiskbestånd i Västkuſtsjoar. Information fran Sötvattenslaboratoriet. DROTTNINGHOLM. 12: 45p.
- Barica, J. and F. A. J. Armstrong. 1971. Contribution by snow to the nutrient budget of some small northwest Ontario lakes. LIMNOL. OCEANOGR. 16: 891-899.
- Beamish, R. J. 1972. Lethal pH for the white sucker *Catostomus commersoni* (Lacépède). TRANS. AMER. FISH. SOC. 101: 355-358.
- _____. 1974 a. Loss of fish populations from unexploited remote lakes in Ontario, Canada as a consequence of atmospheric fallout of acid. WATER RES. 8: 85-95.
- _____. 1974 b. Growth and survival of white suckers (*Catostomus commersoni*) in an acidified lake. J. FISH RES. BOARD CAN. 31: 49;54.
- Beamish, R. J. and H. H. Harvey. 1972. Acidification of the La Cloche Mountain Lakes, Ontario, and resulting fish mortalities. J. FISH RES. BOARD CAN. 29: 1131-1143.

- Beamish, R. J., W. L. Lockhart, J. C. Van Loon, and H. H. Harvey. 1975a Longterm acidification of a lake and resulting effects on fishes. *AMBIO* 4: 92-102.
- Beamish, R. J., L. Milanese, and G. A. McFarlane. 1975b. A fish and chemical survey of 110 ELA lakes with appended reports on whitefish ageing errors and northwestern Ontario bait fishery. *FISH. RES. BOARD CAN. TECH. REP.* (In press.)
- Beamish, R. J., G. A. McFarlane, J. C. Van Loon, and J. Lichwa. 1975c. An examination of the possible effects of Sudbury nickel mining and smelting operations on fishes and the water chemistry of lakes within the Whitefish Lake Indian Reserve. *FISH RES. BOARD CAN. TECH. REP.* (In press.)
- Beamish, R. J. and J. C. Van Loon. 1975. PRECIPITATION LOADING OF ACID, HEAVY METALS AND OTHER SUBSTANCES TO A SMALL ACID LAKE NEAR SUDBURY, ONTARIO, CANADA. (Manuscript in preparation.)
- Biesinger, K. E. and G. M. Christensen. 1972. Effects of various metals on survival growth, reproduction and metabolism of *Daphnia magna*. *J. FISH RES. BOARD CAN.* 29: 1691-1700.
- Brungs, W. A. 1969. Chronic toxicity of zinc to the fathead minnow. *Pimephales promelas* Rafinesque. *TRANS. AMER. FISH. SOC.* 98: 272-279.
- Canada Department of the Environment. 1972. GUIDELINES FOR WATER QUALITY OBJECTIVES AND STANDARDS, A PRELIMINARY REPORT. Inland Waters Branch Publication No. 67, 156 p.
- Daudoroff, P. and M. Katz. 1950. Critical review of literature on the toxicity of industrial wastes and their components to fish. 1. Alkalis, acids and inorganic gases. *SEWAGE IND. WASTES* 22: 1432-1458.
- Dreisinger, B. R. and P. C. McGovern. 1970. MONITORING ATMOSPHERIC SULPHUR DIOXIDE AND CORRELATING ITS EFFECTS ON CROPS AND FOREST IN THE SUDBURY AREA. In *Proc. Impact of Air Pollution on Vegetation Conference*. Toronto, Ont. 23p.
- EIFAC Working Party on Water Quality. 1968. Water quality criteria for European freshwater fish. Rept. on extreme pH values and inland fisheries. *EIFAC Tech. Pap.* 4: 24 p. Also in *WATER RES.* 1969. 3: 593-611.

- EPS. 1973. AIR POLLUTION CONTROL DIRECTORATE, Environmental Protection Service, Department of the Environment. A nation-wide inventory of air pollutant emissions 1970. Rept. EPS 3-AP-73-2, Ottawa, Canada. 123p + app.
- Falkowski, P. 1973. The Inco stack engineering triumph or biological disaster. STEEL LABOR 38(10): 2.
- Gorham, E. 1957. The chemical composition of lake waters in Halifax Co., Nova Scotia. LIMNOL. OCEANOGR. 2: 12-21.
- Gorham, E. and A. G. Gordon. 1960a. The influence of smelter fumes upon the chemical composition of lake waters near Sudbury, Ontario, and upon the surrounding vegetation. CAN J. BOT. 38: 477-487.
- _____. 1960b. Some effects of smelter pollution northeast of Falconbridge, Ontario. CAN. J. BOT. 38: 307-312.
- _____. 1963. Some effects of smelter pollution upon aquatic vegetation near Sudbury, Ontario. CAN. J. BOT. 41: 371-378.
- Harvey, H. H. 1975a. Unpublished manuscript.
- Hörnström, E., C. Ekström, U. Miller, and W. Dickson. 1973. Försurningens inverkan på västkustsjöar. Information från sötvattenslaboratoriet. DROTNINGHOLM. 4: 97p.
- Kopp, J. F. and R. C. Kroner. 1969. TRACE METALS IN WATERS OF THE UNITED STATES (Oct. 1, 1962-Sept. 30, 1967). U. S. Dep. Inter. Fed. Water Pollut. Contr. Admin. Div. Pollut. Surveillance Cincinnati, Ohio. 28 p. + app.
- Kramer, J. R. 1973a. FATE OF ATMOSPHERIC SULPHUR DIOXIDE AND RELATED SUBSTANCES AS INDICATED BY CHEMISTRY OF PRECIPITATION. Unpubl. Rept. J. R. Kramer, Dept. Geology, McMaster Univ. Hamilton, Ontario, 143 p.
- _____. 1973b. Atmospheric composition and precipitation of the Sudbury region. ALTERNATIVES 2: 18-25.
- LeBlanc, F., D. N. Rao, and G. Comeau. 1972. The epiphytic vegetation of *Populus Balsamifera* and its significance as an air pollution indicator in Sudbury, Ontario. CAN J. BOT. 50: 519-528.
- Linzon, S. N. 1958. THE INFLUENCE OF SMELTER FUMES ON THE GROWTH OF WHITE PINE IN THE SUDBURY REGION. Ont. Dep. Lands For., Ont. Dep. Mines, Toronto, Ont. 45 p.

- _____. 1966. Damage to eastern white pine by sulphur dioxide, semimature tissue needle blight, and ozone. J. AIR POLLUT. CONTROL ASSOC. 16: 140-144.
- _____. 1971. Economic effects of sulfur dioxide on forest growth. J. AIR POLLUT. CONTROL ASSOC. 21: 81-86.
- McCallum, A. W. 1944. REPORT ON A PRELIMINARY EXAMINATION OF THE SUDBURY SMELTER AREA (Mimeographed report) Ont. Dep. Mines, Toronto, Ont.
- McGovern, P. C. and D. Basillie. 1972. SULPHUR DIOXIDE LEVELS AND ENVIRONMENTAL STUDIES IN THE SUDBURY AREA DURING 1971. Air Quality Branch, Min. of the Environ. 33 p.
- McGovern, P. C. and D. Basillie. 1973. SULPHUR DIOXIDE (1972) - HEAVY METAL (1971) LEVELS AND VEGETATIVE EFFECTS IN THE SUDBURY AREA. Air Management Branch, Ont. Min. of the Environ. 50 p.
- McKee, J. E. and H. W. Wolf. 1963. WATER QUALITY CRITERIA. State Calif. Water Qual. Control Board, Sacramento Publ. 3A: 548 p.
- McKim, J. M. and D. A. Benoit. 1971. Effects of long-term exposures to copper on survival growth, and reproduction of brook trout (*Salvelinus fontinalis*). J. FISH. RES. BOARD CAN. 28: 655-662.
- Mount, D. I. 1973. Chronic effect of low pH on fathead minnow survival, growth and reproduction. WATER RES. 7: 987-993.
- National Technical Advisory Committee. 1968. WATER QUALITY CRITERIA. Report to the Secretary of the Interior. Fed. Water Pollut. Control Admin. Washington, D. C. 234 p.
- Pickering, Q. H. 1974. Chronic toxicity of nickel to the fathead minnow (*Pimephales promelas*) Rafinesque. J. WATER POLL. CONTROL FED. 46: 760-765.
- Pickering, Q. H. and M. Gast. 1972. Acute and chronic toxicity of cadmium to the fathead minnow (*Pimephales promelas*). J. FISH RES. BOARD CAN. 29: 1099-1106.
- Schindler, D. W. 1971. A hypothesis to explain differences and similarities among lakes in the Experimental Lakes Area, northwestern Ontario. J. FISH RES. BOARD CAN. 28: 295-301.

- Schofield, C. L. 1965. Water quality in relation to survival of brook trout, *Salvelinus fontinalis* (Mitchill). TRANS. AMER. FISH. SOC. 94: 227-235.
- _____. 1975. LAKE ACIDIFICATION IN THE ADIRONDACK MOUNTAINS OF NEW YORK: CAUSES AND CONSEQUENCES PROC. SYMPOSIUM ACID PRECIPITATION AND THE FOREST ECOSYSTEM, (In press.)
- Sprules, W. G. 1975. Midsummer crustacean zooplankton communities in acid-stressed lakes. J. FISH. RES. BOARD CAN. 32: 389-395.
- Stokes, P. M., J. C. Hutchinson, and K. Krauter. 1973. Heavy metal tolerance in algae isolated from polluted lakes near Sudbury, Ontario smelters. WATER POLLUT. RES. IN CANADA, 1973. Proc. 8th Sym. Univ. Waterloo.
- Whaley, H. and G. K. Lee. 1971. THE DISPERSION OF SMELTER PLUMES IN THE SUDBURY AREA. Canada Dep. Energy, Mines and Resources, Ottawa. Mines Branch Investigation Rept. IR 71-37: 20p.
- _____. 1974. PLUME DISPERSION COMPUTATIONS FOR INCO AND FALCONBRIDGE STACK EMISSIONS UNDER LIMITED MIXING CONDITIONS. Canada Dep. Energy, Mines and Resources, Ottawa. Mines Branch Investigations Rept. IR 74-3: 21p.
- Whitby, L. M. and J. C. Hutchinson. 1974. Heavy metal pollution in the Sudbury mining and smelting region of Canada, II Soil toxicity tests. ENVIRON. CONSERVATION 1: 191-200.
- Van Loon, J. C. and R. J. Beamish. 1976. HEAVY METAL CONTAMINATION OF FLIN FLON AREA LAKES AND ITS RELATION TO FISH POPULATIONS. (Manuscript in preparation.)