

North American populations of *Entoleuca mammata* are genetically more variable than populations in Europe

Risto KASANEN^{1*}, Jarkko HANTULA¹, Michael OSTRY², Jean PINON³ and Timo KURKELA¹

¹Finnish Forest Research Institute, Vantaa Research Station, P.O. Box 18, FIN-01301 Vantaa, Finland.

²North Central Research Station, 1561 Lindig Avenue, St Paul, MN 55108, USA.

³UR Pathologie forestière INRA, F-54280 Champenoux, France.

E-mail: risto.kasanen@metla.fi

Received 20 December 2003; accepted 13 April 2004.

Entoleuca mammata (syn. *Hypoxyton mammatum*) is a damaging pathogen of *Populus tremuloides* and *P. grandidentata* in North America and *P. tremula* in Europe, where the fungus occurs only sporadically in alpine regions and Scandinavia. It has been hypothesized that *E. mammata* was introduced to Europe from North America. In this study, *E. mammata* isolates collected from Europe and North America were compared by a sequence analysis of two DNA markers derived from DNA fingerprints. The objective of the study was to elucidate the relationship between North American and European *E. mammata* populations by testing two hypotheses: (1) North American and European isolates are conspecific; and (2) the fungus was introduced between continents causing both a founder effect and a genetic bottleneck. North American populations were found to be more polymorphic, but no major phylogenetic differences between fungal isolates collected from different continents were found. This result combined with the historical observations of the disease in Europe implies that *E. mammata* was introduced to Europe several centuries ago.

INTRODUCTION

Entoleuca mammata (syn. *Hypoxyton mammatum*) is a damaging pathogen of *Populus tremuloides* and to a lesser extent *P. grandidentata* in North America. In Europe, *Populus alba*, *P. trichocarpa*, *P. tremula* and hybrid aspen (*P. tremula* × *P. tremuloides*) are susceptible to the pathogen (Pinon 1979, Terrason, Pinon & Ride 1988, Mazzaglia *et al.* 2001). The taxonomy of *Hypoxyton* has been subject to several rearrangements. Miller (1961) suggested that several species described in the nineteenth century, were synonyms. These included *H. mammiiformis*, *H. pauperatum* and *H. macrosporum* which were collected in northern Scandinavia (Lapland). Miller (1961) gave a relatively broad definition of the genus, while Rogers & Ju (1996) defined a stricter structure of *Hypoxyton*. Based on their definition of *Hypoxyton*, *H. mammatum* was moved to the genus *Entoleuca*, the type species of which is *E. callimorpha* (Rogers & Ju 1996). Later Mazzaglia *et al.* (2001) evaluated genetic relationships among *Hypoxyton* and related genera using 5.8S and ITS sequences of rDNA, but since they did not include *E. callimorpha*

in the analysis, the transfer is not yet confirmed by molecular evidence.

In North America, the disease is prevalent throughout the north-east, the Great Lakes region, and north-western prairies. Incidence is fairly low in the central Rocky Mountains, but the disease is noticeably absent in the northern Rocky Mountains and in Alaska despite the abundance of aspen and collections of the fungus (Manion & Griffin 1986). In the Lake States where *P. tremuloides* is the most important pulpwood species *E. mammata* is considered the major cause of aspen mortality.

Recently, large-scale plantations of aspen and hybrid aspen have been established in Nordic and Baltic countries as the forest industry has become interested in aspen fibre for pulp production. As the cultivation of native and hybrid aspen increases, fungal diseases of aspen may gain in both economic and ecological importance. Potentially, *E. mammata* could hinder the development of aspen cultivation. Serious damage in hybrid aspen trials caused by *E. mammata* has been reported in Scandinavia (Illstedt & Gullberg 1993). However, the report did not describe the basis for the identification of the pathogen, so the exact cause of the damage is uncertain.

* Corresponding author.

E. mammata was thought to be restricted to North America, even though European *P. alba* and *P. tremula* were known to be susceptible (Manion & Griffin 1986). The disease was first observed in France in 1975, while the fungus was still a quarantine pest in Europe. At the time, it was thought that the pathogen might have been introduced to Europe in the 1950s on infected flower branches of *P. tremuloides* and *P. grandidentata* used in hybrid aspen breeding (Pinon 1979). However, the geographical distribution of the fungus in France makes it unlikely that the fungus was a recent introduction. The disease was present at several isolated locations in the French Alps where *P. tremula* had been collected for breeding purposes, but where no potentially infected plant material had been introduced. In France, the fungus is widely distributed and damaging in mountain stands, but it is relatively rare and localized in the plains. The distribution of the fungus may depend on suitable hosts. In France two races of *P. tremula* exist, one in mountain regions and the other in the plains. The mountain race is found in Jura, in the Alps (350–1850 m a.s.l.) and in the Pyrénées.

Several European *E. mammata* herbarium specimens examined by Miller (1961) had been collected in the nineteenth century, confirming that *E. mammata* has been present in Europe longer than previously believed. Few occurrences of *E. mammata* are represented in Fennoscandian herbaria (Miller 1961), and the fungus apparently occurs only sporadically in aspen stands. Although the fungus can cause serious mortality of aspen in individual stands, no serious outbreaks have been recorded in Europe. This could be interpreted as a balance between host and pathogen that is usually considered a result of coexistence and coevolution. Based on the information of historical abundance and geographic range of the fungus in Europe, *E. mammata* could be considered a native species. However, the relationship of European and North American populations of the fungus has not been examined, and molecular studies of the population structure of this worldwide pathogen are lacking. The history of *E. mammata* in Europe is consequently poorly known, and open to several interpretations.

The objective of this investigation was to elucidate the relationship between North American, central European and Finnish isolates of *E. mammata* by studying the genetic structure of populations to test the following hypotheses: (1) North American and European isolates are conspecific; and (2) the fungus was introduced between continents causing both a founder effect and a genetic bottleneck. According to hypothesis (1), identical alleles should be found in both continents and no phylogenetic grouping of alleles based on the geographic origin of the isolates should be detected. Hypothesis (2) was tested by measuring genetic differentiation between populations and by analysing genetic variability between continents.

MATERIALS AND METHODS

Fungal isolates

Entoleuca mammata samples were collected from Europe (EU) from 20 locations in France, Finland, Switzerland and Italy (Table 1). The host species included one *Salix*, two *Populus* species, and a *Populus* hybrid. The collections were made during 1975–2002. The French isolates were collected from several locations across the known distributional area of *E. mammata*. The Finnish populations were collected from two natural aspen stands in Eastern Finland. The size of the sampling area was approx. 5 ha in both stands, which were separated by 10 km. In North America (NA), two populations located in Minnesota were collected in 2001 and 2002. The 2001 population was from along a trail set in natural forest over three miles, and in 2002 the sampling was performed throughout a hybrid aspen plantation several acres in size. The NA populations were geographically separated by over 100 miles. In total, 63 samples were subjected to analysis. Most of the isolations of fungal hyphae were made from the canker margin. Pieces of wood (approx. 10 × 5 × 2 mm) were surface-sterilized by rinsing in 4% NaOCl, 75% ethanol and finally sterile water (10 s each) and incubated on 1.5% malt agar. As fungal growth occurred, the isolates were sub-cultured on 1.5% malt agar. In a few cases, single-ascospore isolations were made by casting spores from moistened perithecia onto 1.5% malt agar (Table 1).

Molecular markers

DNA isolations were made as described by Vainio, Korhonen & Hantula (1998). The fungal tissue was homogenized in 1.5 ml disposable plastic tubes by grinding with a glass rod and quartz sand. Briefly, the DNA isolation procedure included two phenol-chloroform (1:1) extractions, chloroform-phenol-isoamyl alcohol extractions, precipitation with polyethylene glycol (PEG) and drying. The DNA was resuspended into 10 mM Tris HCl buffer containing 1 mM EDTA.

Random amplified microsatellite markers DBV(CAT)₅ and HBH(GAG)₅ (where B = C/G/T, D = A/G/T, H = A/C/T and V = A/C/G) (RAMS; Hantula *et al.* 1996) were screened in order to find single loci that were amplifiable from a subset of European samples. In the RAMS method, the primers are attached to distal ends of microsatellites, a procedure also utilized in ISSR-PCR (inter short-sequence repeat PCR) (Zhou *et al.* 1999). The PCR protocol by Hantula, Dusabenyagasaki & Hamelin (1996) was used. Next, two RAMS markers signed as *Hgag* (819 bp) and *Hcat* (327 bp) were cloned with TOPO TA cloning kit (Invitrogen, Carlsbad, CA). The sequences were determined with a Li-Cor IR2 DNA sequencer, using M13 universal forward and reverse primers. Based on the sequences obtained, two pairs of primers were designed to amplify fragments. The primer sequences

Table 1. Origin of *Entoleuca mammata* isolates analysed. All isolates are cultured *in vitro* and deposited in the Finnish Forest Research Institute (Vantaa).

Isolate number	Host	Locality	Year of isolation	Isolation
Population samples from Finland				
H (5 isolates)	<i>Populus tremula</i>	Punkaharju	2000	CM ^b
K (6 isolates)	<i>P. tremula</i>	Punkaharju	2000	CM
Population samples from North America				
US (14 isolates)	<i>P. tremuloides</i>	Minnesota	2001	CM
U2 (13 isolates)	<i>P. tremuloides</i>	Minnesota	2002	CM
Isolates from France				
JP22	<i>P. tremula</i>	Chorges (Hautes-Alpes)	nk ^a	CM
JP24	<i>P. tremula</i>	Alps (Les Guibertes)	1975 or 76	
JP26	<i>P. tremula</i>	Alps (Les Guibertes)	1975 or 76	CM
JP28	<i>P. tremula</i> × <i>tremuloides</i>	Velaine-sous-Amance (Lorraine)	1975	CM
JP37	<i>P. tremula</i>	Alps (Le Revard)	1976	CM
JP38	<i>P. tremula</i>	Alps (Le Revard)	1976	CM
JP39	<i>P. tremula</i>	Alps (Albion)	1976	CM
JP41	<i>P. tremula</i>	Alps (Revest-du-Bion)	1977	CM
JP43	<i>P. tremula</i>	Alps (Couloubron)	1977	CM
JP44	<i>P. tremula</i>	Jura	1977	CM
JP84	<i>P. tremula</i>	Lorraine (Ludres)	1978	monospore
JP86	<i>P. tremula</i>	Lorraine (Ludres)	1978	monospore
JP87	<i>P. tremula</i>	Lorraine (Ludres)	1978	monospore
JP88	<i>P. tremula</i>	Lorraine (Ludres)	1978	monospore
JP164	<i>P. trichocarpa</i>	Orléans	Around 1978	CM
JP178	<i>P. tremula</i>	nk	nk	monospore
JP180	<i>P. tremula</i>	nk	nk	monospore
JP182	<i>P. trichocarpa</i>	West (Nantes)	1985	CM
JP187	<i>P. trichocarpa</i>	West (Nantes)	1985	CM
JP192	<i>P. trichocarpa</i> 'Fritzi Pauley'	(Centre) Nogent-sur-Vernisson	1986	CM
JP193	<i>P. trichocarpa</i> 'Fritzi Pauley'	(Centre) Nogent-sur-Vernisson	1986	CM
Isolates from other countries				
JP21	<i>P. tremula</i>	Italy (around Cuneo)	1975	CM
JP181	<i>P. tremula</i>	Italy (around Cuneo)	1985	CM
JP323	<i>Salix</i> sp.	Switzerland	1991	CM
JP271	<i>P. tremula</i>	Andorra, Fontenada	1992	CM

^a nk, not known. ^b CM, canker margin.

were: *HgagF*; 5' gga gat gaa cag cga gga ct, *HgagR*; 5' ggc gaa act gag gag agt ca, *HcatF*; 5' aaa tcc act ccc aaa tgc aa and *HgagR*; 5' ccc tac ctc tag gcc acc tc. These fragments were amplified from all isolates, purified with Hi-Pure PCR Product Purification Kit (Roche, Mannheim) and sequenced with the primers used for amplification. The sequencing was conducted with a LI-COR IR2 automated sequencer according to the manufacturer's recommendations except that annealing temperature of 61 °C was applied.

Testing the hypothesis of conspecificity

To support the hypothesis of conspecificity (1) of the fungus in North America (NA) and Europe (EU), identical alleles should be found from both continents and no phylogenetic grouping of alleles based on geographic origin of isolates should be detected. The alleles were determined by sequencing. To study phylogenetic structures within sequence data, a distance matrix was computed from nucleotide sequences using Kimura's 2-parameter genetic distance model (Kimura 1980). The UPGMA algorithm was used to calculate phylogeny from the distance matrix. Prior to analysis the data was bootstrapped with SEQBOOT

program within the PHYLIP software package. The calculations were made with DNADIST and NEIGHBOR programs. Consensus trees were constructed by CONSENSE program (Felsenstein 1989). For the out-group we chose the sample JP323 collected from *Salix* sp. in Switzerland because in preliminary analyses the sample was shown to form a distant and separate branch in phylogenies. Phenograms were constructed with the TREEDRAW program (Page 1996).

Testing the introduction hypothesis

The introduction hypothesis (2) was tested by measuring genetic differentiation of the fungal populations between continents and analysing genetic variability within populations. In order to measure genetic differentiation caused by a hypothetical founder effect between North American and European populations, we calculated the F_{st} analogue defined by Weir & Cockerham (1984) and G_{st} value (Nei 1987). All calculations were made with FSTAT software (Goudet 2000). As FSTAT has been originally designed for diploid data sets, haploid fungal genotypes were encoded as diploid homozygotes, as suggested by Goudet (2000).

		3	5	7	9	11	13	15	17	19	21	23	25	27	29	31	33	35	37	39	41	43	45	47	49	51	53	55	57	59	61	63	65	67	69	71	73	75	77	79	81	83	85	87	89	91	93	95	97	99	101	103	105	107	109	111	113	115	117	119	121	123	125	127	129	131	133	135	137	139	141	143	145	147	149	151	153	155	157	159	161	163	165	167	169	171	173	175	177	179	181	183	185	187	189	191	193	195	197	199	201	203	205	207	209	211	213	215	217	219	221	223	225	227	229	231	233	235	237	239	241	243	245	247	249	251	253	255	257	259	261	263	265	267	269	271	273	275	277	279	281	283	285	287	289	291	293	295	297	299	301	303	305	307	309	311	313	315	317	319	321	323	325	327	329	331	333	335	337	339	341	343	345	347	349	351	353	355	357	359	361	363	365	367	369	371	373	375	377	379	381	383	385	387	389	391	393	395	397	399	401	403	405	407	409	411	413	415	417	419	421	423	425	427	429	431	433	435	437	439	441	443	445	447	449	451	453	455	457	459	461	463	465	467	469	471	473	475	477	479	481	483	485	487	489	491	493	495	497	499	501	503	505	507	509	511	513	515	517	519	521	523	525	527	529	531	533	535	537	539	541	543	545	547	549	551	553	555	557	559	561	563	565	567	569	571	573	575	577	579	581	583	585	587	589	591	593	595	597	599	601	603	605	607	609	611	613	615	617	619	621	623	625	627	629	631	633	635	637	639	641	643	645	647	649	651	653	655	657	659	661	663	665	667	669	671	673	675	677	679	681	683	685	687	689	691	693	695	697	699	701	703	705	707	709	711	713	715	717	719	721	723	725	727	729	731	733	735	737	739	741	743	745	747	749	751	753	755	757	759	761	763	765	767	769	771	773	775	777	779	781	783	785	787	789	791	793	795	797	799	801	803	805	807	809	811	813	815	817	819	821	823	825	827	829	831	833	835	837	839	841	843	845	847	849	851	853	855	857	859	861	863	865	867	869	871	873	875	877	879	881	883	885	887	889	891	893	895	897	899	901	903	905	907	909	911	913	915	917	919	921	923	925	927	929	931	933	935	937	939	941	943	945	947	949	951	953	955	957	959	961	963	965	967	969	971	973	975	977	979	981	983	985	987	989	991	993	995	997	999	1001	1003	1005	1007	1009	1011	1013	1015	1017	1019	1021	1023	1025	1027	1029	1031	1033	1035	1037	1039	1041	1043	1045	1047	1049	1051	1053	1055	1057	1059	1061	1063	1065	1067	1069	1071	1073	1075	1077	1079	1081	1083	1085	1087	1089	1091	1093	1095	1097	1099	1101	1103	1105	1107	1109	1111	1113	1115	1117	1119	1121	1123	1125	1127	1129	1131	1133	1135	1137	1139	1141	1143	1145	1147	1149	1151	1153	1155	1157	1159	1161	1163	1165	1167	1169	1171	1173	1175	1177	1179	1181	1183	1185	1187	1189	1191	1193	1195	1197	1199	1201	1203	1205	1207	1209	1211	1213	1215	1217	1219	1221	1223	1225	1227	1229	1231	1233	1235	1237	1239	1241	1243	1245	1247	1249	1251	1253	1255	1257	1259	1261	1263	1265	1267	1269	1271	1273	1275	1277	1279	1281	1283	1285	1287	1289	1291	1293	1295	1297	1299	1301	1303	1305	1307	1309	1311	1313	1315	1317	1319	1321	1323	1325	1327	1329	1331	1333	1335	1337	1339	1341	1343	1345	1347	1349	1351	1353	1355	1357	1359	1361	1363	1365	1367	1369	1371	1373	1375	1377	1379	1381	1383	1385	1387	1389	1391	1393	1395	1397	1399	1401	1403	1405	1407	1409	1411	1413	1415	1417	1419	1421	1423	1425	1427	1429	1431	1433	1435	1437	1439	1441	1443	1445	1447	1449	1451	1453	1455	1457	1459	1461	1463	1465	1467	1469	1471	1473	1475	1477	1479	1481	1483	1485	1487	1489	1491	1493	1495	1497	1499	1501	1503	1505	1507	1509	1511	1513	1515	1517	1519	1521	1523	1525	1527	1529	1531	1533	1535	1537	1539	1541	1543	1545	1547	1549	1551	1553	1555	1557	1559	1561	1563	1565	1567	1569	1571	1573	1575	1577	1579	1581	1583	1585	1587	1589	1591	1593	1595	1597	1599	1601	1603	1605	1607	1609	1611	1613	1615	1617	1619	1621	1623	1625	1627	1629	1631	1633	1635	1637	1639	1641	1643	1645	1647	1649	1651	1653	1655	1657	1659	1661	1663	1665	1667	1669	1671	1673	1675	1677	1679	1681	1683	1685	1687	1689	1691	1693	1695	1697	1699	1701	1703	1705	1707	1709	1711	1713	1715	1717	1719	1721	1723	1725	1727	1729	1731	1733	1735	1737	1739	1741	1743	1745	1747	1749	1751	1753	1755	1757	1759	1761	1763	1765	1767	1769	1771	1773	1775	1777	1779	1781	1783	1785	1787	1789	1791	1793	1795	1797	1799	1801	1803	1805	1807	1809	1811	1813	1815	1817	1819	1821	1823	1825	1827	1829	1831	1833	1835	1837	1839	1841	1843	1845	1847	1849	1851	1853	1855	1857	1859	1861	1863	1865	1867	1869	1871	1873	1875	1877	1879	1881	1883	1885	1887	1889	1891	1893	1895	1897	1899	1901	1903	1905	1907	1909	1911	1913	1915	1917	1919	1921	1923	1925	1927	1929	1931	1933	1935	1937	1939	1941	1943	1945	1947	1949	1951	1953	1955	1957	1959	1961	1963	1965	1967	1969	1971	1973	1975	1977	1979	1981	1983	1985	1987	1989	1991	1993	1995	1997	1999	2001	2003	2005	2007	2009	2011	2013	2015	2017	2019	2021	2023	2025	2027	2029	2031	2033	2035	2037	2039	2041	2043	2045	2047	2049	2051	2053	2055	2057	2059	2061	2063	2065	2067	2069	2071	2073	2075	2077	2079	2081	2083	2085	2087	2089	2091	2093	2095	2097	2099	2101	2103	2105	2107	2109	2111	2113	2115	2117	2119	2121	2123	2125	2127	2129	2131	2133	2135	2137	2139	2141	2143	2145	2147	2149	2151	2153	2155	2157	2159	2161	2163	2165	2167	2169	2171	2173	2175	2177	2179	2181	2183	2185	2187	2189	2191	2193	2195	2197	2199	2201	2203	2205	2207	2209	2211	2213	2215	2217	2219	2221	2223	2225	2227	2229	2231	2233	2235	2237	2239	2241	2243	2245	2247	2249	2251	2253	2255	2257	2259	2261	2263	2265	2267	2269	2271	2273	2275	2277	2279	2281	2283	2285	2287	2289	2291	2293	2295	2297	2299	2301	2303	2305	2307	2309	2311	2313	2315	2317	2319	2321	2323	2325	2327	2329	2331	2333	2335	2337	2339	2341	2343	2345	2347	2349	2351	2353	2355	2357	2359	2361	2363	2365	2367	2369	2371	2373	2375	2377	2379	2381	2383	2385	2387	2389	2391	2393	2395	2397	2399	2401	2403	2405	2407	2409	2411	2413	2415	2417	2419	2421	2423	2425	2427	2429	2431	2433	2435	2437	2439	2441	2443	2445	2447	2449	2451	2453	2455	2457	2459	2461	2463	2465	2467	2469	2471	2473	2475	2477	2479	2481	2483	2485	2487	2489	2491	2493	2495	2497	2499	2501	2503	2505	2507	2509	2511	2513	2515	2517	2519	2521	2523	2525	2527	2529	2531	2533	2535	2537	2539	2541	2543	2545	2547	2549	2551	2553	2555	2557	2559	2561	2563	2565	2567	2569	2571	2573	2575	2577	2579	2581	2583	2585	2587	2589	2591	2593	2595	2597	2599	2601	2603	2605	2607	2609	2611	2613	2615	2617	2619	2621	2623	2625	2627	2629	2631	2633	2635	2637	2639	2641	2643	2645	2647	2649	2651	2653	2655	2657	2659	2661	2663	2665	2667	2669	2671	2673	2675	2677	2679	2681	2683	2685	2687	2689	2691	2693	2695	2697	2699	2701	2703	2705	2707	2709	2711	2713	2715	2717	2719	2721	2723	2725	2727	2729	2731	2733	2735	2737	2739	2741	2743	2745	2747	2749	2751	2753	2755	2757	2759	2761	2763	2765	2767	2769	2771	2773	2775	2777	2779	2781	2783	2785	2787	2789	2791	2793	2795	2797	2799	2801	2803	2805	2807	2809	2811	2813	2815	2817	2819	2821	2823	2825	2827	2829	2831	2833	2835	2837	2839	2841	2843	2845	2847	2849	2851	
--	--	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	--

		38	40	88	96	100	106	119	123	160	180	189	191	192	201	203	204	208	216	217	218	223	227	232	233	235	236	240	242	248	249	250	253	254	916	
U7	:	C	G	A	A	T	-	G	A	C	C	A	A	T	T	C	C	C	T	G	G	C	T	C	C	A	G	C	-	T	-	C	-	-	T	AY210717
U14	:	C	G	A	G	T	-	G	A	C	G	A	A	T	C	C	C	C	T	G	G	C	T	C	C	A	A	C	-	T	-	C	-	-	T	AY210718
U9	:	T	G	A	A	T	-	C	A	C	G	A	A	T	T	T	C	C	T	G	G	C	T	C	C	A	G	C	-	T	-	C	T	-	T	AY210719
U13	:	C	G	A	A	T	-	G	A	C	G	A	A	T	T	C	C	C	T	G	G	C	T	C	T	A	A	C	-	T	-	C	-	-	T	AY210720
K6	:	C	G	A	A	T	-	G	A	C	C	G	A	T	T	C	C	T	T	G	G	T	T	C	C	A	G	C	-	T	-	C	T	-	T	AY210721
U5	:	C	G	A	G	T	-	A	A	C	G	A	A	T	T	C	C	C	T	G	G	C	T	C	C	A	A	C	-	T	-	C	-	-	T	AY210722
U3	:	C	G	A	G	T	-	G	A	C	C	G	A	T	T	C	C	T	T	G	G	C	T	C	C	A	G	C	-	T	-	C	T	-	T	AY210723
H1	:	C	G	A	G	T	-	G	A	C	C	A	A	T	T	C	C	T	T	G	G	C	T	C	C	A	G	C	-	T	-	C	T	-	T	AY210724
U1	:	C	G	A	A	C	-	G	A	C	C	G	A	T	T	C	C	T	T	G	G	C	T	C	C	A	G	C	-	C	-	C	T	-	T	AY210725
U2-4	:	C	G	A	A	T	-	G	T	T	C	G	A	T	T	C	C	T	T	G	G	C	T	C	C	A	G	C	-	T	-	T	T	-	T	AY210726
U2-1	:	C	G	A	G	T	-	G	A	C	C	G	A	T	T	C	C	T	C	G	G	C	T	C	C	A	G	C	-	T	-	C	T	-	T	AY210727
U2-11:		C	G	C	A	T	-	G	A	C	G	A	A	T	T	C	C	C	T	G	G	C	T	C	C	A	A	C	-	T	-	C	-	-	T	AY210728
U2-12:		C	A	A	A	T	-	G	A	C	C	G	A	T	T	C	C	T	T	G	G	C	T	C	C	A	G	C	-	T	-	C	T	-	T	AY210729
U2-7	:	C	G	A	A	T	G	G	A	C	G	A	A	T	T	C	C	T	T	G	G	C	T	C	C	A	G	C	C	T	T	C	-	-	T	AY210730
U2-9	:	C	G	A	A	T	-	G	A	C	C	G	A	T	T	C	C	T	-	-	-	C	T	C	C	A	A	C	-	T	-	C	-	-	T	AY210731
U2-8	:	C	G	A	A	T	-	G	A	C	C	A	A	T	T	C	C	T	T	G	G	C	T	C	C	A	G	C	-	T	-	C	-	-	T	AY210732
JP87	:	C	G	A	A	T	-	G	A	C	C	G	A	T	T	C	C	T	T	G	G	C	T	C	C	A	G	C	-	T	-	C	T	-	G	AY210733
JP44	:	C	G	A	A	T	-	G	A	C	C	G	A	T	T	C	C	T	T	G	G	C	T	C	C	A	G	C	-	T	-	C	T	-	T	AY210734
JP187:		C	G	A	A	T	-	G	A	C	C	A	A	C	T	C	C	T	T	G	G	T	T	T	C	A	G	C	-	T	-	C	-	-	T	AY210735
JP323:		C	G	A	A	T	-	G	A	C	C	A	A	C	T	C	G	T	T	A	G	T	G	C	C	A	G	C	-	T	-	C	-	-	T	AY210736
U8	:	C	G	A	G	T	-	G	A	C	C	A	A	T	T	C	C	T	T	G	G	C	T	C	C	C	G	C	-	T	-	C	-	-	T	AY210737
JP21	:	C	G	A	A	T	-	G	A	C	C	G	A	T	T	C	C	T	T	G	G	C	T	C	C	A	G	T	-	T	-	C	T	*	T	AY210738

Fig. 2. Polymorphic sites of alleles of marker locus *Hgag* (819) bp. Non-polymorphic sites are deleted from the sequences. Complete sequences can be obtained from GenBank (<http://www.ncbi.nlm.nih.gov>; accession numbers are given in the end of each line). The nucleotides used for artificial loci are in bold. The sample JP21 contained a 590 bp insert which is marked as *.

Table 2. Distribution of alleles in marker loci.

Marker	<i>n</i>	Number of alleles	Number of alleles occurring in USA/number of isolates	Number of alleles occurring in Europe/number of isolates	Number of common alleles
<i>Hcat</i>	63	28	21/27	9/36	2
<i>Hgag</i>	63	22	17/27	8/36	3

Introduction hypothesis

Allelic richness defined by Nei's diversity was considerably higher in NA populations than in EU populations (Table 3). Rarefaction curves and the Chao-estimator both suggest that the allelic and haplotype richness is higher in North America than in Europe (Fig. 5, Table 3).

The G_{st} and F_{st} values calculated over *Hgag* and *Hcat* loci were 0.160 and 0.193, respectively (Table 4). Thus the analysis implies that there is population differentiation between NA and EU samples. Compared to the allele numbers at these loci (28 and 22) the allele numbers in artificial loci were lower; the number of alleles in *gag5*, *cat7*, *gagL* and *catL* were six, eight, five and seven, respectively. The G_{st} and F_{st} values calculated over the four artificial loci were 0.150 and 0.187, respectively, which are almost identical with those calculated for *Hgag* and *Hcat* loci (Table 4). Higher G_{st} and F_{st} values were obtained from artificial loci based on point mutations than from those based only on the length. This was obviously due to identical length of alleles with considerably different sequences


(homoplasy). This indicates that the high number of alleles is not the source of the high F_{st} and G_{st} values obtained for *Hgag* and *Hcat*.

DISCUSSION

This is the first molecular analysis of the genetic polymorphism in *Entoleuca mammata* populations. The number of alleles observed in the two loci was relatively high, 22 and 28 alleles per locus. The resolution of the marker system enabled us to identify a unique genotype for all NA samples ($n=27$) with the two markers. A clear methodological conclusion can be drawn: sequence analysis of individual loci obtained originally from genetic fingerprinting is a powerful method of analysing variation. A methodological benefit over genetic fingerprinting systems such as RAMS (Hantula *et al.* 1996) and RAPD (Williams *et al.* 1990) is that the homology of the loci studied between samples is simultaneously verified in sequencing. This is especially valuable if sampling and fungal isolation covers a wide geographical area (different continents) and the fungus has a highly variable *in vitro* morphology such as *E. mammata* (Bagga & Smalley 1974).

The hypothesis of conspecificity of North American and European *E. mammata* was supported by the observation of several identical alleles in the two loci on both continents. The conspecificity was also supported by the lack of phylogenetic grouping of alleles according to their origin, and the absence of significant nodes in the phylogenies.

The second question addressed in this investigation was the direction of introduction. Theoretically,




H and K are Finnish isolates; U, North American isolates; JP, European isolates.

* Isolate JP44 was identical to isolates JP22, JP271, JP28, JP39, JP41, JP26, JP37, JP38, JP43, JP84, JP86, JP88, JP329, JP178, JP180, JP192, JP28, JP193, U2-13 and U11.

** Isolate H1 was identical to H3, H7, H10, H11, K3, K4, K7, U4, U10, U12, U2-5, JP24 and JP164.

Fig. 3. Unrooted UPGMA consensus tree of *Entoleuca mammata* isolates. The sequence data from *HgaI* marker locus was bootstrapped (100×) and using Kimura's two-parameter genetic distance was calculated (9). Bootstrap values for each node are given. The alignment and the tree can be obtained from TreeBase (<http://www.treebase.org>; reference numbers: S1066, M1817).

organisms introduced from one location to another should have reduced genetic diversity compared to the original population. Good examples of such effects are the reduced diversity of the European race of *Gremmeniella abietina* in North America (Hamelin, Lecours & Laflamme 1998), and the clonal spread of the Irish potato famine fungus, *Phytophthora infestans* (Goodwin, Cohen & Fry 1994). Therefore, low variation can be used as a fingerprint for introduced species. In this study the genetic diversity of *E. mammata* was estimated using Nei's diversity, which was lower in EU than in NA. As this difference could have been due to differences in sample sizes, we calculated rarefaction curves and Chao-estimates, both of which supported our hypothesis. This was also in accordance



H and K are Finnish isolates; U, North American isolates; JP, European isolates.

* Isolate K11 was identical with isolates K8, U9, U2-12, JP24, JP28, JP87, JP178, JP180, JP193 and JP271.

** Isolate U2-7 was identical with U2-8, JP39, JP43, JP44, JP84, JP86, JP88, JP164 and JP192.


Fig. 4. Unrooted UPGMA consensus tree of *Entoleuca mammata* isolates. The sequence data from *Hcat* marker locus was bootstrapped (100×) and using Kimura's two-parameter genetic distance was calculated (9). Bootstrap values for each node are given. The alignment and the tree can be obtained from TreeBase (<http://www.treebase.org>; reference numbers: S1066, M1818).

with our being able to identify a unique genotype for all NA-isolates, whereas groups of identical genotypes were observed among EU-isolates. All these analyses (Nei's diversity, rarefaction, Chao-estimator, and distribution of genotypes) support a hypothesis suggesting introduction of *E. mammata* from NA to EU.

If an introduction from NA to EU had taken place, genetic differentiation should have occurred due to a founder effect and random drift. In accordance with this, both F_{st} and G_{st} values suggested that *E. mammata* populations in the two continents would be genetically differentiated. As it was unclear whether the very high number of alleles per locus could cause bias in the analyses of population differentiation, we tested this possibility by calculating the same indices (G_{st} , F_{st}) from the measured (total) data and from an artificial

Table 3. Expected number of alleles (Chao-estimators) and Nei's genetic diversities calculated for the two loci.

Marker	Chao-estimates		Nei's diversity		All samples
	North America	Europe	North America	Europe	
Hcat	76	14	0.94	0.81	0.90
Hgag	29	15	0.92	0.58	0.81

**Fig. 5.** Richness estimated using the rarefaction method, $E(g_n)$. Richness estimates for haplotypes; North America (USA), Finland, and Central Europe (EU).

data set (created by including only a subset of total variation to reach a more 'normal' number of alleles/locus). Since the F_{st} and G_{st} values for both total data and artificial data sets were in agreement, we concluded that the high number of alleles/locus did not affect the analysis. Therefore, our results indicate true genetic difference between populations of *E. mammatata* in the two continents and support the hypothesis that *E. mammatata* was introduced from NA to EU.

The number of shared alleles in Europe and North America is relatively small at both loci. This is, however, expected in hypervariable loci when the sampling is not sufficient to cover most of the alleles occurring in the populations (as was the case during this study). In short, assuming that there are no large differences in allele frequencies, the expected numbers of shared alleles (if in EU only a subset of NA alleles occurred) can be calculated from the following equation:

$$ALL_{shared} = \frac{EU_{est}}{NA_{est}} \times EU_{obs}$$

where ALL_{shared} is the expected number of alleles occurring both in EU and NA, EU_{est} is the estimated total number of alleles in Europe, NA_{est} is the estimated total number of alleles in NA and EU_{obs} is the observed number of alleles in EU (Table 2). For EU_{est}

Table 4. Comparison of F_{st} and G_{st} values calculated for the original loci Hgag and Hcat and averaged over artificial loci gag5, cat7 and gagL, catL which presented lower diversity with 6, 8, 5 and 7 alleles for locus, respectively.

Marker	F_{st}	G_{st}
Hcat & Hgag	0.193	0.160
gag5 & cat7	0.285	0.233
gagL & catL	0.088	0.067
Over all artificial loci	0.187	0.150

and NA_{est} we used the Chao estimators obtained for European and North American populations, respectively (Table 3).

For Hcat-locus the Chao estimators were 14 and 76 alleles in EU and NA, respectively, and in Europe altogether nine alleles were observed. Based on these figures, 1.7 ($9 \times 14 / 76$) shared alleles were expected to be found in our data set. This figure is very close to the two shared alleles actually observed. For Hgag-locus the expected number of common alleles in EU and NA was 4.1 ($8 \times 15 / 29$), which also is close to the three common alleles actually observed. Therefore, these calculations support the introduction hypothesis.

Before concluding that *E. mammatata* was introduced by man from North America to Europe, we must consider an alternative explanation for the reduced diversity in Europe. Since the sampling of European isolates was conducted over 27 yr, it is possible (although unlikely) that population shifts occurred over this period. Another possibility is the proportion of latent infections (Chapela 1989), which were not subject to sampling; theoretically this could have skewed the sampling because only aggressive (canker-causing) isolates have been collected. However, our experience is that if a canker is 'contained' or limited by callus growth it is mostly due to resistance of the genotype of the host clone rather than differences in aggressiveness of the isolates. We also consider that selection of aggressiveness should not have had an effect on the results obtained from selectively neutral molecular markers, assuming that the markers and aggressiveness are not genetically linked. If these distant possibilities are excluded, the diversity of European population would reduce if *E. mammatata* was native in both North America and Europe and the European population would have gone through a genetic bottleneck after the separation of the two continents. Such a bottleneck could have been caused by glaciations if the Mediterranean sea and the mountains in southern Europe would have hindered the escape of fungi from ice. The outcome of such a bottleneck would have been an acceleration of neutral marker evolution due to fixation of new mutations in small populations. This should have been seen in our data, because unique alleles in Europe would have reduced the number of alleles shared by NA and EU. This, however, was not the case (see above), and therefore we favor the introduction hypothesis over the subdivision + bottleneck

hypothesis. This conclusion is also in accordance with the literature, as there are several native fungi in NA and EU that have evolved to genetic dissimilarity. Examples include *Gremmeniella abietina* (Dusabenyagani, Laflamme & Hamelin 2002), *Heterobasidion annosum* IS-group S (Johannesson & Stenlid 2003), and *Phlebia gigantea* (Vainio & Hantula 2000).

In NA populations, we observed a high degree of polymorphism which is consistent with the assumption that *E. mammata* is a common, variable and native pathogen in NA. The identification of a unique genotype for each NA isolate strongly suggests that *E. mammata* is an actively outcrossing fungus lacking asexual spread but disseminated primarily by ascospores (Griffin *et al.* 1992). Observations of a complex interaction of *E. mammata* with several wounding agents; woodpeckers and insects such as the poplar borers *Saperda inornata* and *Oberea schaumii*, the periodical cicada *Magicicada septendecim*, the dogday cicada *Tibicen linnei* and the treehopper *Telamona tremulata* (Ostry & Anderson 1998) also suggest a long-term coevolution of the host-pathogen system in North America.

In speculating on possible introduction events, the host range of *E. mammata* must be taken into consideration. The fungus has a wide host range of world-wide genera of trees, including *Acer*, *Alnus*, *Carpinus*, *Fagus*, *Picea*, *Populus*, *Pyrus*, *Salix*, *Sorbus*, and *Ulmus* (Miller 1961). It is therefore possible that the fungus was introduced with some tree species other than poplar or aspen, perhaps some seedlings brought in for ornamental or gardening purposes.

It is possible that in Europe the distribution of the fungus is actually wider than recognized and may include hosts other than *Populus* sp. For example, several species of *Salix* are common, widespread and susceptible (Miller 1961). Being an economically overlooked tree species, there are no reports of the disease on *Salix* in Europe, but herbarium samples have been collected. Unfortunately only a few limited host range studies have been conducted by cross-inoculating hosts, and these have been done only in North America (French, Hodges & Froyd 1969, French & Juzwik 1984). Although little data from other tree species are available, sample JP323 collected from *Salix* sp. (Switzerland) had the most dissimilar alleles at both loci, suggesting that the pathogen in the two tree genera might be distinct although closely related.

Examples of introduced fungi include ones carried by humans and long-distance dispersal by air and other vectors (Brown & Hovmoller 2002). In many cases introduced pathogens such as *Cryptonectria parasitica*, the cause of chestnut blight, and more recently *Phytophthora ramorum*, the cause of sudden oak death, have caused severe outbreaks in new environments or on new hosts (Beattie & Diller 1954, Rizzo *et al.* 2002). Compared to those examples, *E. mammata* has so far caused only minor damage to European aspen stands, although in local situations it can be devastating. In

North America, where the pathogen causes more damage, a number of wounding agents (insects and birds) and various environmental variables are believed to play critical roles in spreading the pathogen (Ostry & Anderson 1998). Although the importance of wounding agents or vectors has not been studied in Europe it is possible that an absence of some factor could be limiting the ability of *E. mammata* to spread and cause damage.

In this study we tested two hypotheses: conspecificity of *E. mammata* in NA and EU, and the hypothesis of introduction. The hypothesis of conspecificity was supported by the observation of identical alleles and absence of phylogenetic differentiation. Supporting the introduction hypothesis, analyses of the distribution of genotypes, degree of genetic variation and genetic differentiation all imply that *E. mammata* is a native pathogen in North America and an introduced one in Europe. The introduction may have occurred several centuries ago, and the actual introduction event remains unknown. The degree of genetic variation is relatively high in Europe but considerably less than in North America. This suggests that the introduction was not a single event but most likely occurred in several locations during the last centuries.

ACKNOWLEDGEMENTS

We thank Egbert Beuker for field observations of damage to aspen, Neil Anderson for reviewing the manuscript, and Arlette Schipfer for her participation in survey and isolations. This study was financially supported by the Metsäliitto Group.

REFERENCES

- Bagga, D. K. & Smalley, E. B. (1974) Variation of *Hypoxyylon pruinaum* in cultural morphology and virulence. *Phytopathology* **64**: 663–667.
- Beattie, R. K. & Diller, J. D. (1954) Fifty years of chestnut blight in America. *Journal of Forestry* **52**: 323–329.
- Brown, J. K. M. & Hovmoller, M. S. (2002) Aerial dispersal of pathogens on the global and continental scales and its impact on plant disease. *Science* **297**: 537–541.
- Chao, A. (1987) Estimating the population size for capture-recapture data with unequal catchability. *Biometrics* **43**: 783–791.
- Chapela, I. H. (1989) Fungi in healthy stems and branches of American beech and aspen: a comparative study. *New Phytologist* **113**: 65–75.
- Dusabenyagani, M., Laflamme, G. & Hamelin, R. C. (2002) Nucleotide polymorphisms in three genes support host and geographic speciation in tree pathogens belonging to *Gremmeniella* spp. *Canadian Journal of Botany* **80**: 1151–1159.
- Felsenstein, J. (1989) PHYLIP – phylogeny inference package (Version 3.2). *Cladistics* **5**: 164–166.
- French, D. W., Hodges, C. S. & Froyd, J. D. (1969) Pathogenicity and taxonomy of *Hypoxyylon mammatum*. *Canadian Journal of Botany* **47**: 223–226.
- French, D. W. & Juzwik, J. (1984) *Hypoxyylon* canker of *Salix* sp. *Phytopathology* **74**: 840.
- Goodwin, S. B., Cohen, B. A. & Fry, W. E. (1994) Panglobal distribution of a single clonal lineage of the Irish potato famine fungus. *Proceedings of the National Academy of Sciences, USA* **91**: 11591–11595.

- Goudet, J. (2000) *FSTAT: a program to estimate and test gene diversities and fixation indices*. Version 2.9.3. <http://www.unil.ch/izea/software/fstat.html>.
- Griffin, D. H., Quinn, K. E., Gilbert, G. S., Wang, C. J. K. & Rosemarin, S. (1992) The role of ascospores and conidia as propagules in the disease cycle of *Hypoxylon mammatum*. *Phytopathology* **82**: 114–119.
- Grünwald, N. J., Goodwin, S. B., Milgroom, M. G. & Fry, W. E. (2003) Analysis of genotypic diversity data for populations of microorganisms. *Phytopathology* **93**: 738–746.
- Hamelin, R. C., Lecours, N. & Laflamme, G. (1998) Molecular evidence of two distinct introductions of the European race of *Gremmeniella abietina* to North America. *Phytopathology* **88**: 582–588.
- Hantula, J., Dusabenyagasaki, M. & Hamelin, R. C. (1996) Random amplified microsatellites (RAMS) – a novel method for characterizing genetic variation within fungi. *European Journal of Forest Pathology* **26**: 159–166.
- Illstedt, B. & Gullberg, U. (1993) Genetic variation in a 26-year old hybrid aspen trial in Southern Sweden. *Scandinavian Journal of Forest Research* **8**: 185–192.
- Johannesson, H. & Stenlid, J. (2003) Molecular markers reveal genetic isolation and phylogeography of the S and F intersterility groups of the wood-decay fungus *Heterobasidion annosum*. *Molecular Phylogenetics and Evolution* **29**: 94–101.
- Kimura, M. (1980) A simple model for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* **16**: 111–120.
- Manion, P. D. & Griffin, D. H. (1986) Sixty-five years of research on *Hypoxylon* canker of aspen. *Plant Disease* **70**: 803–808.
- Mazzaglia, A., Anselmi, N., Vicario, S. & Vannini, A. (2001) Sequence analysis of the 5.8S rDNA and ITS regions in evaluating genetic relationship among some species of *Hypoxylon* and related genera. *Mycological Research* **105**: 670–675.
- Miller, J. H. (1961) *A Monograph of the World Species of Hypoxylon*. University of Georgia Press, Athens.
- Nei, M. (1973) Analysis of gene diversity in subdivided populations. *Proceedings of the National Academy of Sciences, USA* **70**: 3321–3323.
- Nei, M. (1987) *Molecular Evolutionary Genetics*. Columbia University Press, New York.
- Ostry, M. E. & Anderson, N. A. (1998) Interactions of insects, woodpeckers and *Hypoxylon* canker on aspen. *USDA Forest Service Research Paper* 331.
- Page, R. D. M. (1996) TREEVIEW: an application to display phylogenetic trees on personal computers. *Computer Applications in the Biosciences* **12**: 357–358.
- Pinon, J. (1979) Origine et principaux caractères des souches françaises d'*Hypoxylon mammatum* (Wahl.) Miller. *European Journal of Forest Pathology* **9**: 129–142.
- Rizzo, D. M., Garbelotto, M., Davidson, J. M., Slaughter, G. W. & Koike, S. T. (2002) *Phytophthora ramorum* as the cause of extensive mortality of *Quercus* spp. and *Lithocarpus densiflorus* in California. *Plant Disease* **86**: 205–214.
- Rogers, J. D. & Ju, Y.-M. (1996) *Entoleuca mammatata* comb. nov. for *Hypoxylon mammatum* and the genus *Entoleuca*. *Mycotaxon* **59**: 441–448.
- Simberloff, D. S. (1978) Use of rarefaction and related methods in ecology. In *Biological Data in Water Pollution Assessment: quantitative and statistical analyses* (K. L. Dickson, J. Cairns & R. J. Livingston, eds): 150–165. American Society for Testing and Materials, Philadelphia.
- Terrasson, D., Pinon, J. & Ridé, M. (1988) L'infection naturelle de *Populus trichocarpa* par *Hypoxylon mammatum*. *Revue Forests Française* **40**: 126–130.
- Vainio, E. J. & Hantula, J. (2000) Genetic differentiation between European and North American population of *Phlebiopsis gigantea*. *Mycologia* **92**: 436–446.
- Vainio, E. J., Korhonen, K. & Hantula, J. (1998) Genetic variation in *Phlebiopsis gigantea* as determined with random amplified microsatellite (RAMS) markers. *Mycological Research* **102**: 187–192.
- Weir, B. S. & Cockerham, C. C. (1984) Estimating F-statistics for the analysis of population structure. *Evolution* **38**: 1358–1370.
- Williams, J. G. K., Kubelik, A. R., Livak, K. J., Rafalski, J. A. & Tingey, S. V. (1990) DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research* **18**: 6531–6535.
- Zhou, Z. H., Miwa, M. & Hogetsu, T. (1999) Analysis of genetic structure of a *Suillus grevillei* population in a *Larix kaempferi* stand by polymorphism of inter-simple sequence repeat (ISSR). *New Phytologist* **144**: 55–63.

Corresponding Editor: G. W. Griffith