

Reproduction of a Woodwasp, *Urocerus japonicus* (Hymenoptera: Siricidae) Using No Maternal Symbiotic Fungus

Hideshi Fukuda

Unit of Environmental Systems, Faculty of Social and Informational Sciences,
Nihon Fukushi University Higashiehaemi-cho, Handa, Aichi Japan

Abstract

Most woodwasps (Siricidae) are symbiotically associated with the specific fungus, *Amylostereum* spp. Female adults inoculate the fungus during their oviposition in sapwood of the host trees (Morgan 1968). Woodwasp larvae can digest sapwood with low nutritional quality with the aid of symbiosis (Kukor and Martin 1983). In the earlier study, we clarified that a woodwasp with no fungal symbionts, *Xeris spectrum* can utilize the fungal symbionts of other woodwasp species without possessing any symbiotic fungi of its own (Fukuda and Hijii 1997). Moreover, the larvae of fungus-carrying woodwasp species cannot develop on living trees. The female adults oviposit selectively on fresh trees that are presumed to be suitable for fungus propagation, because the conditions of the wood at the time of oviposition affect propagation of the fungus (Fukuda 2002). Whether fungus-carrying woodwasps can develop using no maternal symbiotic fungus or not has never been studied. Thus, I conducted fungus-isolation and oviposition experiments to evaluate the preference and performance of *U. japonicus* on fungus-inoculated trees.

Experiments were conducted from 1999 to 2001. In July 1999, the first year, new female adults of *U. japonicus* were allowed to oviposit on living trees of *Cryptomeria japonica* (oviposited trees). In October, the symbiotic fungus of *U. japonicus* was artificially inoculated on living trees of *C. japonica* (inoculated trees). In November, oviposited trees, inoculated trees and control (not oviposited and not inoculated) trees were felled and a part of the inoculated trees were bucked to 2 m. In July 2000, the second year, new female adults were allowed to oviposit on each tree. At the same time, fungi were isolated from both oviposited and inoculated trees. In 2001, the third year, I counted the number of new adults that emerged from each tree where oviposition occurred in the 2nd year, and I also counted oviposition holes on each tree.

The inoculated symbiotic fungus propagated on both oviposited trees and inoculated trees 8 months after tree-felling, oviposition period of the next year. Moreover, on inoculated trees, especially on bucked trees, the symbiotic fungus was distributed widely in the wood. The oviposition by *U. japonicus* was higher on oviposited trees and inoculated trees than on control trees; moreover, the oviposition was higher on inoculated trees than on oviposited trees. On inoculated trees, many oviposition holes were observed near vertical lines from inoculated positions, where the symbiotic fungus would propagate vigorously. Next generation adults emerged from inoculated trees, whereas no adults emerged from oviposited and control trees. The number of emerged adults was especially high on bucked trees. From these results, I demonstrated that a fungus-carrying woodwasp species, *U. japonicus*, can reproduce using no maternal symbiotic fungus. These results indicate that both *X. spectrum*, a woodwasp species that has no maternal symbiont, and *U. japonicus* a fungus-carrying woodwasp species, can utilize the *Amylostereum* fungi which had already propagated in the wood. This information is important for clarifying the coevolution between woodwasps and *Amylostereum* fungi.

Key Words: *Amylostereum* fungus, fungus-isolation, oviposition preference, survival rate, *Urocerus japonicus*

References Cited

- Fukuda, H. 2000.** Trunk sapwood utilization by woodwasps. *Japanese Journal of Ecology* 52:75-80.
- Fukuda H. and Hijii N. 1997.** Reproductive strategy of a woodwasp with no fungal symbionts, *Xeris spectrum* (Hymenoptera: Siricidae). *Oecologia* 112:551-556.
- Kukor J.J. and Martin M.M. 1983.** Acquisition of digestive enzymes by siricid woodwasps from their fungal symbiont. *Science* 220:1161-1163.
- Morgan F.D. 1968.** Bionomics of Siricidae. *Annual Review of Entomology* 13: 239-256.