



**Forest  
Service**


**Northern  
Research Station**

**Resource Bulletin  
NRS-14**



# **Iowa's Forest Resources in 2005**

**Susan J. Crocker, Gray J. Brand, and Aron Flickinger**



---

## **Abstract**

This report presents Iowa's annual inventory results for 2005. Estimates show that Iowa has more than 2.8 million acres of forest land. Total live-tree volume on forest land is 4.0 billion cubic feet (ft<sup>3</sup>). Ninety-eight percent of forest land is classified as timberland. Oak/hickory is the predominant forest-type group, representing 54 percent of timberland area. Growing-stock volume has risen to an estimated 3.1 billion ft<sup>3</sup>. Live-tree aboveground biomass is 112 million dry tons. In general, Iowa's forests are healthy. However, diseases, including white oak decline and Dutch elm disease, and the threat of invasion from exotic species such as emerald ash borer and nonnative plants are a concern when assessing the health of Iowa's forests.

---

Manuscript received for publication 21 December 2006

Published by:  
USDA FOREST SERVICE  
NORTHERN RESEARCH STATION  
11 CAMPUS BLVD, SUITE 200  
NEWTOWN SQUARE PA 19073-3294

March 2007

Visit our homepage at: <http://www.nrs.fs.fed.us/>

# Iowa's Forest Resources, 2005

## INTRODUCTION

The forest resources of Iowa are annually inventoried by the Northern Research Station's Forest Inventory and Analysis (NRS-FIA) program. Annual inventory is based on data from a set of field plots that are systematically located across the entire state. Each year, a new panel consisting of one-fifth of the total field plots is measured and combined with four previously measured panels. From these "moving average" estimates, annual inventories are available for each year since 2003, providing yearly estimates for such variables as area, volume, and biomass. Estimates for components of change (growth, removals, and mortality) are based on remeasured plots. For this report, estimates of growth, removals, and mortality were calculated using two panels of plots measured in 1999 and 2000, and then remeasured in 2004 and 2005.

Reported estimates are based on sampling and estimation techniques of Bechtold and Patterson (2005). All of the tables in this report and many others can be generated at the Mapmaker Program at <http://nrs.fs.fed.us/fia/data-tools/mapping-tools/default.asp>.

## RESULTS

### Area

There are approximately 35.7 million acres of land in Iowa (U.S. Census Bureau 2006). Eight percent of the state's total land area, or more than 2.8 million acres, is forest land<sup>1</sup> (Table 1).

Forest land has three components:

1. Timberland<sup>2</sup>—land that is not restricted from harvesting by statute, administrative

regulation, or designation and is capable of growing trees at a rate of 20 cubic feet (ft<sup>3</sup>) per acre per year.

2. Reserved forest land—land that is restricted from harvesting by statute, administrative regulation, or designation (e.g., state parks, national parks and lakeshores, and federal wilderness areas).
3. Other forest land—land that is not capable of growing trees at a rate of 20 ft<sup>3</sup>/ac/yr and is not restricted from harvesting.

Timberland totals 2.8 million acres and accounts for 98 percent of forest land (Table 2). The remaining 2 percent (55.2 thousand acres) of forest land is classified as reserved or other forest land.

Timberland area declined in the late 1950's and continued to do so through the early 1970's. A record low was recorded in 1974 when timberland was an estimated 1.4 million acres (Fig. 1). Since the 1974 inventory, the area of Iowa timberland has steadily increased. This rebound coincided with the decline of the farm economy in the 1970's and 1980's. During this time, pastures and marginal agricultural lands reverted to timberland. Timberland area also increased as a result of the re-establishment of trees in riparian zones and the action of initiatives such as the Conservation Reserve Program. In 2003, the area of timberland surpassed the acreage present in 1954 (Fig. 1). The expansion of timberland continues as the 2005 estimate shows an increase in total area since 2003.

### THE AUTHORS


SUSAN J. CROCKER and GARY J. BRAND are, respectively, a Forester and Research Forester with the USDA Forest Service's Northern Research Station, St. Paul, Minnesota.

ARON FLICKINGER is a Forester with the Iowa Department of Natural Resources, Bureau of Forestry.

<sup>1</sup> Forest land is land that is at least 10-percent stocked with trees of any size, or that formerly had such tree cover and is not currently developed for a nonforest use. The minimum area for classification of forest land is 1 acre. In addition, strips of timber must have a crown width of at least 120 feet.

<sup>2</sup> Timberland may not be equivalent to the area actually available for commercial timber harvesting or other access. The actual availability of land for various uses depends on owner decisions that consider economic, environmental, and social factors.

Figure 1.—Area of timberland by inventory year, Iowa, 1954-2005 (the sampling error associated with each inventory estimate represents a 67-percent confidence interval and is depicted by the vertical line at the top of each bar).



Private owners hold nearly 9 of every 10 acres of Iowa timberland (Table 2). The remaining area (312 thousand acres) is publicly owned. Public ownership ensures that people will have access to forest recreation opportunities and helps protect wildlife habitat and watershed quality.


Ninety-six percent of Iowa's timberland is composed of hardwoods (Table 2). The state's timberland is primarily oak/hickory (54 percent), though elm/ash/cottonwood (24 percent) and maple/basswood<sup>3</sup> forest-type groups

(14 percent) are also large components of overall species composition (Table 3, Fig. 2). In the oak/hickory forest-type group, the white oak/red oak/hickory forest type (555 thousand acres) and the mixed upland hardwoods forest type (510 thousand acres) predominate. Within the elm/ash/cottonwood group, the sugarberry/ hackberry/elm/green ash forest type (324 thousand acres) occupies the largest area. The sugar maple/yellow birch<sup>4</sup> (148 thousand acres) and hard maple/basswood (149 thousand acres) forest types dominate the maple/basswood forest-type group (Table 3).

<sup>3</sup> The maple/basswood forest-type group is regionally specific and equivalent to the maple/beech/birch forest-type group found in the core tables. Forest-type groupings in the core tables are nationally standardized to maintain consistency and allow comparisons across states.

<sup>4</sup> The sugar maple/yellow birch forest type is equivalent to the sugar maple/beech/yellow birch forest-type group found in the core tables. Forest types in the core tables are nationally standardized to maintain consistency and allow comparisons across states.

Figure 2.—Percentage of timberland by forest-type group, Iowa, 2001-2005 (the maple/basswood forest-type group is regionally specific and is equivalent to the maple/beech/birch forest-type group; Iowa is outside of the native range of beech).



Softwood forest types make up only 1 percent of total timberland area. Eastern redcedar is the principal coniferous species in Iowa forests. Although the eastern redcedar forest type accounts for only 1 percent (33.1 thousand acres) of total timberland area, this species is often mixed with hardwoods in the eastern redcedar/hardwood forest type. Overall, eastern redcedar has a significant presence as it is found on an estimated 95.1 thousand acres of Iowa timberland (Table 3). The remaining 3 percent of timberland is nonstocked.<sup>5</sup>

Stand-size class is a measure of the average diameter of the dominant trees in a stand and reflects a stand's size structure. There are three classes: sawtimber—large trees, softwoods at least 9 inches in diameter at breast height (d.b.h.) and hardwoods at least 11 inches in d.b.h.; poletimber—medium trees, 5 inches in d.b.h. to sawtimber size; and sapling/seedling—small trees, 1 to 5 inches in d.b.h., or live trees less than 1 inch in d.b.h. that are expected to survive. In 2005, most of Iowa's timberland area was composed of large diameter trees. Sawtimber stands accounted for 64 percent of timberland area (Fig. 3, Table 3). Poletimber and sapling/seedling stands repre-

sented 21 and 12 percent, respectively; remaining timberland was nonstocked.

The considerable area of sawtimber indicates a mature forest. However, the relatively high proportion of sapling/seedling acreage in the eastern redcedar and eastern redcedar/hardwood forest types suggests a growing eastern redcedar resource.

## Volume

Total net volume of live trees on forest land is 4.0 billion ft<sup>3</sup>, or 1,424 ft<sup>3</sup>/ac of forest land (Table 4). Net volume is gross volume less deductions for rot, sweep, or other defects affecting use for timber products, and is computed from a 1-foot stump to a 4-inch diameter top (outside the bark) for live trees at least 5 inches in d.b.h. Eighty-six percent of live volume is on privately owned forest land (Table 4). Virtually all (99 percent) of the volume of live trees on forest land is in hardwoods; 31 percent of live-tree volume is in oak species, primarily bur, white, and northern red oaks (Table 4).

Ninety-eight percent of live-tree volume is on timberland (Table 5); 77 percent of this volume is in growing-stock trees. Growing-stock volume has traditionally been used to ascertain wood volume that is useful for commercial

<sup>5</sup> Nonstocked land is timberland that is less than 10-percent stocked with live trees.




Figure 3.—Stand-size class as a percentage of total timberland area, Iowa, 2001-2005.

purposes. It is the amount of solid wood in commercial trees 5 inches in d.b.h. or larger, measured from a 1-foot stump to a minimum 4-inch diameter top, with deductions for poor form or defect. Excluded are rough, rotten, and dead trees, and trees of noncommercial species.

Growing-stock volume on Iowa's timberland has been on the rise since 1974, gradually increasing each year since 2003 (Fig. 4). Nearly doubling from 1.6 billion ft<sup>3</sup> in 1990, the estimate of growing-stock volume for 2005 is 3.1 billion ft<sup>3</sup> (Fig. 4). This increase reflects a forest in which larger trees continue to increase in volume.

Slightly more than half of total growing-stock volume is in the oak/hickory forest-type group (Table 6). The oak/hickory, elm/ash/cottonwood, and maple/basswood forest-type groups account for 99 percent of all growing-stock volume. Growing-stock volume present in softwood forest types is 5.7 million ft<sup>3</sup>, or less than 0.2 percent of total growing-stock volume. However, growing-stock volume for all softwood species is 18.1 million ft<sup>3</sup> (slightly more than 0.5 percent of Iowa's total growing-stock volume) because softwoods sometimes make up a portion of hardwood stands.


Seventy-eight percent of total growing-stock volume is in trees that are 11 inches or larger in d.b.h. (Table 7). Twenty-four percent of total growing-stock volume is in trees that are greater than or equal to 21 inches in d.b.h.; 38

percent of this volume is in various oak species. Sixty-six percent of eastern cottonwood growing-stock volume is in trees 21 inches d.b.h. or larger. Additionally, cottonwood volume is 30 percent of total growing-stock volume in trees 21 inches or larger in d.b.h. (Table 7).

Sawtimber volume is the volume of the sawlog portion of growing-stock trees of sawtimber size expressed in board feet (International 1/4-inch rule). Iowa sawtimber volume totals 11.3 billion board feet (Table 8). Virtually all of Iowa sawtimber volume is in hardwood trees; less than 1 percent of sawtimber volume comes from softwoods. Six species account for 62 percent of the sawtimber volume in Iowa: eastern cottonwood (1.4 billion board feet); silver maple, white oak, and bur oak (1.2 billion board feet each); northern red oak (1.0 billion board feet); and black walnut (864 million board feet) (Table 8). Of total sawtimber volume, 29 percent is in trees 21 inches or larger in d.b.h. Eastern cottonwood accounts for 31 percent of volume of sawtimber trees 21 inches or larger in d.b.h. (Table 8).

Twenty-three percent of live-tree volume on timberland (930 million ft<sup>3</sup>) is in cull trees (Table 5). Cull trees are unsuitable for use as wood products due to poor form, rot, defect, or because they are considered an undesirable species. Cull volume is often used for commercial purposes. For instance, rough trees are sometimes harvested for chipping or making pallets. Salvable dead trees, standing or down

Figure 4.—Growing-stock volume on timberland, Iowa, 1954-2005 (the sampling error associated with each inventory estimate represents a 67-percent confidence interval and is depicted by the vertical line at the top of each bar).



dead trees that are considered merchantable by regional standards, make up an estimated 57.4 million ft<sup>3</sup> of wood volume (Table 5). Salvable dead trees have commercial applications and are an important source of firewood. They also play an important role in overall species diversity, providing habitat for a wealth of wildlife species, including cavity nesting birds and mammals that require den sites.

### **Biomass**

Biomass estimates are becoming increasingly important in analyses of questions related to carbon sequestration, wood fiber availability for fuels, and assessment of fuel loads in forest stands. Live-tree aboveground biomass was estimated for growing-stock and nongrowing-stock trees, and live trees 1 to 5 inches in d.b.h. (Table 9). In 2005, live-tree aboveground biomass on Iowa's timberland was an estimated 112 million dry tons (an average of 39 dry tons/ac of timberland). Ninety-nine percent or 111 million dry tons of biomass was in hardwood species. Sixty-nine percent of live-tree aboveground biomass was in growing-stock trees, 26 percent in nongrowing-stock trees, and 5 percent in trees 1 to 5 inches in d.b.h. For both growing-stock and nongrowing-stock trees, nearly 75 percent of total aboveground biomass was in the boles of trees; the remainder was in stumps, tops, and limbs (Table 9).

### **Growth, Removals, and Mortality**

Estimates of growth, removals, and mortality provide information on changes and trends in forest resources over time.

Net growth (gross growth minus mortality) of growing stock on Iowa's timberland increased by an average of 141 million ft<sup>3</sup>/yr (Table 10). Virtually all (99 percent or 139 million ft<sup>3</sup>) of annual net growth was in hardwoods. The majority of hardwood growth occurred in the select white oaks (24.7 million ft<sup>3</sup>/yr) and other eastern soft hardwoods species groups (23.3 million ft<sup>3</sup>/yr) (Table 10). Softwoods accounted for the remaining 1 percent of net growth. Seventy-nine percent of net annual growth for all species groups was on private land (Table 10).

Average annual removals of growing stock on timberland totaled 18.2 million ft<sup>3</sup>/yr, or 13 percent of annual net growth (Table 11).

Eighty-five percent of annual removals occurred within five species groups: black walnut (4.2 million ft<sup>3</sup>/yr), select red oaks (3.3 million ft<sup>3</sup>/yr), select white oaks (3.2 million ft<sup>3</sup>/yr), other eastern soft hardwoods (2.4 million ft<sup>3</sup>/yr), and other red oaks (2.3 million ft<sup>3</sup>/yr) (Table 11). The reported estimates may not yet provide a true depiction of removals because only two inventory panels have been remeasured. The remeasurement of additional panels will yield more definitive results.

Making up 16 percent of annual net growth, average annual mortality of growing stock on timberland was 23.1 million ft<sup>3</sup>/yr (Table 12). Nearly 100 percent of total mortality was among hardwoods. The other eastern soft hardwoods group had the highest mortality at 11.9 million ft<sup>3</sup>/yr, or 51 percent of total mortality (Table 12). Most of the mortality in the other eastern soft hardwoods group is likely due to a statewide outbreak of Dutch elm disease. The oak species groups accounted for 20 percent of total average annual mortality; within oak species groups, select white oaks had the highest mortality at 3.1 million ft<sup>3</sup>/yr (Table 12). Oak mortality in Iowa is generally caused by oak wilt and white oak decline.

## **FOREST HEALTH**

The following information about the pathogens and insects affecting Iowa's forests was obtained from staff at the Iowa Department of Natural Resources (DNR) and the Forest Service's Northeastern Area, State and Private Forestry. Additional information was gathered from the national Forest Health Monitoring (FHM) program webpage (<http://fhm.fs.fed.us/>), 2005 Insect and Disease Conditions Report (<http://www.na.fs.fed.us/fhp/pcond/>), and Central States Forest Health Watch newsletter (<http://www.na.fs.fed.us/fhp/fhw/csflhw/>). Several issues of concern in 2005 are highlighted below. For more information on the health of Iowa's forests, contact the Iowa DNR.

## Dutch Elm Disease

Dutch elm disease (DED) continues to affect the structure and health of Iowa forests. Aerial surveys conducted along the Des Moines, Iowa, Cedar, and Mississippi Rivers detected DED on 1,362 acres in 27 counties in central and western Iowa. The highest incidences were in Black Hawk, Louisa, Clinton, and Buchanan Counties. DED affects all species of elm, but American elm is particularly susceptible; trees are often killed before reaching saw-timber size, so maturing stands present a future health risk.

DED is caused by the fungi *Ophiostoma ulmi* and *O. novo-ulmi*. Infected tree branches clog water-conducting tissues in an attempt to stop fungal growth and as a result, prevent water from reaching the crown. Leaves then wilt and die, causing tree death. The DED fungus is spread to healthy trees in two ways. Long-distance dispersal occurs when elm bark beetles pick up fungal spores in diseased trees and carry them to healthy trees. Spores are deposited as beetles feed in twig crotches or as they bore through the inner bark and sapwood. Local spread occurs through root grafts, which allow the fungus to readily move between neighboring trees. For more information on DED, see Haugen (1998) or [http://www.na.fs.fed.us/spfo/pubs/howtos/ht\\_ded/ht\\_ded.htm](http://www.na.fs.fed.us/spfo/pubs/howtos/ht_ded/ht_ded.htm).

## Gypsy Moth

Since its introduction to North America in 1869, gypsy moth has become widespread in the Eastern United States. Its distribution stretches south from Massachusetts to Virginia and west to Wisconsin and Illinois (USDA Animal and Plant Health Inspection Service 2006). A pest of numerous species of trees and shrubs, including oaks, basswood, cottonwood, and willow, gypsy moth defoliates nearly 1 million acres a year (McManus et al. 1992). Iowa does not have an established gypsy moth population, but concern is increasing as recent surveys have identified established moth populations in neighboring Wisconsin. In response to the growing threat of gypsy moth invasion, monitoring has been intensified through the collaborative efforts of the Iowa Department of Agriculture and Land

Stewardship, USDA Animal and Plant Health Inspection Service, Iowa DNR, Eastern Iowa City Foresters, and numerous volunteers. In 2005, approximately 5,000 insect traps placed throughout the state yielded 4 moths. The 2005 trap catch represents a dramatic decrease since the 2003 capture of 159 moths and the fewest moths captured in Iowa in 16 years. Decreases in the number of gypsy moth captured in Iowa are likely due to a combination of environmental factors and the recent success of the national "Slow the Spread" program in Wisconsin and Illinois.

## Oak Wilt

A significant contributor to oak mortality, especially in southeastern Iowa, oak wilt has been identified as the most serious disease affecting Central States forests. All species of oak are susceptible to oak wilt, but the disease occurs more frequently and progresses more rapidly in red oak species. Oak wilt is caused by the fungus *Ceratocystis fagacearum*. As with DED, the fungus enters the vascular system of trees. In an attempt to block fungal growth, the tree plugs water-conducting tissues, which disrupts the translocation of water from the roots to the crown. This disruption causes the foliage to wilt and die. Rapid progression of the disease results in tree mortality within a year of initial infection. As there is no cure for oak wilt, prevention and early detection are important means of sustaining tree health.

*C. fagacearum* can spread through root grafts or be transported from a diseased tree to a healthy tree by sap-feeding beetles carrying fungal spores. Spread of the fungus through the root system is inhibited by mechanically breaking natural root grafts that have formed between diseased and healthy trees. Overland spread can be reduced by removing dead or dying oaks before fungal mats, which contain spores, are produced. Effective fungal mats only form during the first spring following tree death, and only if there is adequate moisture in the bark; for this reason, dead trees should be removed and treated before fungal mats form. Standing dead trees left longer than 1 year can be left without posing further risk of fungal spread. Injury and fresh pruning

wounds attract sap-feeding beetles, so unnecessary pruning, and pruning between April 15 and July 1 should be avoided. Additional information on the biology and control of oak wilt can be found in O'Brien et al. (2000), available at [http://www.na.fs.fed.us/spfo/pubs/howtos/ht\\_oakwilt/toc.htm](http://www.na.fs.fed.us/spfo/pubs/howtos/ht_oakwilt/toc.htm).

### **White Oak Decline**

In recent years there has been an increase in the amount of mature standing-dead white oak in northeastern Iowa woodlands. The cause of this white oak decline and mortality has yet to be identified. Researchers from the Iowa DNR and the Forest Service are investigating this decline complex. Aerial surveys conducted in 2005 revealed that the combined area affected by white oak decline and oak wilt totaled 1,422 acres.

### **Sudden Oak Death**

*Phytophthora ramorum* is the causal agent of sudden oak death (SOD). Hosts of *P. ramorum* include many species of oaks, Douglas-fir, *Rhododendron* spp., and a variety of other trees and shrubs. On oaks, *P. ramorum* causes bleeding cankers along the stem. Cankered trees can survive for several years after infection. However, mortality occurs within weeks of the onset of crown dieback (O'Brien et al. 2002). SOD was identified in the United States in 1995. While transported via nursery stock to states outside of the West Coast, established populations of *P. ramorum* are known to occur only in California and southwestern Oregon. Because transportation of infected nursery stock is a major threat to central and eastern hardwood forests (O'Brien et al. 2002), the Iowa DNR targeted nurseries that have received California nursery stock that may have been infected, and established transects in forested areas to monitor for established populations. All samples collected in Iowa tested negative for the SOD pathogen. For more information, please visit the California Oak Mortality Task Force webpage, [www.sudenoakdeath.org](http://www.sudenoakdeath.org).

### **Emerald Ash Borer**

Emerald ash borer (EAB), an exotic inner bark-boring beetle, was discovered in the United States in 2002. From its introduction point in southeastern Michigan, this insect has spread throughout Lower Michigan and into Ohio and Indiana; in 2006, during the writing of this report, EAB was positively identified in Maryland and northeastern Illinois. EAB attacks and kills all species of ash, including green, white, and black ash. Larval feeding and gallery production in the phloem and outer sapwood sever the flow of nutrients and girdle the tree, causing tree death. Mortality occurs within 1 to 3 years of initial infestation. Since its discovery, EAB has killed millions of ash trees. Natural spread of EAB has been enhanced by human-assisted transportation of infested firewood and nursery stock. Detection monitoring programs for EAB were initiated in Iowa in 2004. Continued in 2005, visual surveys conducted in all 99 Iowa counties and trap tree inspection failed to detect EAB. More information about EAB can be found at <http://www.emeraldashborer.info>.

### **Invasive Plants**

Invasive plant species continue to threaten the health of Iowa forests. Garlic mustard, buckthorn, bush honeysuckle, and multiflora rose are the state's primary woodland invaders. In the battle for limited resources, invasive species often outcompete native plants and thus achieve greater reproductive success. Invasive species are a major concern because they threaten overall ecological diversity and alter natural communities. Efforts to monitor the presence and severity of invasive species in Iowa are under way.

## **SUMMARY**

The area of Iowa timberland is increasing. Growing-stock volume is also rising, having nearly doubled since 1990. While Iowa forests are reasonably healthy, the dispersal of exotic insects and fungal pathogens continues to threaten forest health.

## APPENDIX

### Accuracy of the Inventory

Sampling errors measure the uncertainty in estimates derived from a portion of a population rather than from the population as a whole. The following are sampling errors for the estimates of statewide totals in this report. These sampling errors represent a 67-percent confidence interval.

| Table no. | Variable                                                             | Estimate   | Sampling error |
|-----------|----------------------------------------------------------------------|------------|----------------|
|           |                                                                      |            | Percent        |
| 1         | Area of forest land (thousand acres)                                 | 2,879      | 2.67           |
| 2         | Area of timberland (thousand acres)                                  | 2,824      | 2.74           |
| 4         | All-live volume on forest land (thousand ft <sup>3</sup> )           | 4,099,569  | 4.10           |
| 5         | All-live volume on timberland (thousand ft <sup>3</sup> )            | 4,045,090  | 4.17           |
| 6         | Growing-stock volume on timberland (thousand ft <sup>3</sup> )       | 3,114,499  | 4.84           |
| 8         | Sawtimber volume on timberland (thousand board feet)                 | 11,380,405 | 5.58           |
| 9         | All-live aboveground biomass on timberland (thousand dry tons)       | 112,598    | 3.72           |
| 10        | Growing-stock growth on timberland (thousand ft <sup>3</sup> /yr)    | 141,249    | 14.66          |
| 11        | Growing-stock removals on timberland (thousand ft <sup>3</sup> /yr)  | 18,281     | 38.34          |
| 12        | Growing-stock mortality on timberland (thousand ft <sup>3</sup> /yr) | 23,179     | 18.88          |

#### Inventory Methods

With the NRS-FIA annual inventory system, about one-fifth of all field plots are measured each year. The first annual inventory cycle was completed in 2003. Measurements from the 2003 inventory were summarized in Leatherberry et al. (2005); for a more detailed analysis of the data see Leatherberry et al. (2006). Each panel measured after 2003 represents a complete sample of the forest resources of Iowa. Therefore, changes in estimates of area, volume, and biomass can be compared for 2003, 2004, and 2005.

The 2005 Iowa forest inventory includes a total sample of 6,067 plots over the entire state from 2001 to 2005. All of the plots provide information for estimates of current conditions (area, volume, and biomass), but only the remeasured plots provide information for estimates of change (growth, removals, and mortality).

Current estimates for growth, removals, and mortality are based on remeasurement of plots established in the first and second panels of the annual inventory (1999 and 2000). Over the next 3 years, the remaining panels will be

remeasured and estimates of change will be annually updated with new measurements. The smaller number of sample plots currently available for estimation of change components is reflected in the sampling errors presented in this report. Sampling errors for growing-stock growth, removals, and mortality on timberland (14.66 percent, 38.34 percent, and 18.88 percent, respectively) are considerably higher than those for timberland area and growing-stock volume on timberland (2.74 percent and 4.84 percent, respectively).

The annual inventory system includes new remote sensing technology, and a new sampling design and plot configuration. The advent of remote-sensing technology has allowed NRS-FIA to use classifications of Multi-Resolution Land Characterization data and other available remote sensing and GIS products to stratify the total area of the state and improve the precision of estimates.

Additionally, new algorithms have been used to assign forest type and stand-size class to each condition observed on a plot. These algorithms are employed nationwide by FIA to increase consistency among states.

## **Inventory Phases**

The annual inventory system is based on a three-phase inventory. In the first phase, classified satellite images are used to stratify the state and aerial photographs are used to select plots for measurement. The second phase involves measuring of a traditional FIA suite of mensurational variables (basic tree and stand attributes), while the third phase focuses on measuring a suite of variables related to forest health.

The only plots that could not be measured were on: (1) private land where field personnel were unable to obtain permission from the owner to measure the field plot, and (2) plots that were inaccessible because of a hazard or danger to field personnel. The methods used in preparing this report were adjusted to account for sites where access was denied or hazardous.

### **Phase 1**

This inventory used a classification of satellite imagery and ancillary data for stratification. FIA used the imagery to form two initial strata: forest and nonforest. Pixels within 60 m (2-pixel widths) of a forest/nonforest boundary formed two additional strata: forest edge and nonforest edge. Forest pixels within 60 m on the forest side of a forest/nonforest boundary were classified in a forest edge stratum. Pixels within 60 m of the boundary on the nonforest side were classified in a nonforest edge stratum. In addition, all strata were divided into public or private ownership based on information available in the Protected Lands Database (DellaSala et al. 2001). The estimated population total for a variable is the sum across all strata of the product of each stratum's area (from the pixel count) and the variable's mean per unit area (from plot measurements) for the stratum.

### **Phase 2**

Phase 2 of the inventory consisted of measuring 6,067 field plots throughout the state. Current FIA precision standards for annual inventories require a sampling intensity of one plot for approximately every 6,000 acres for

phase 2 plots. FIA has divided the entire area of the United States into nonoverlapping hexagons, each of which contains 5,937 acres (McRoberts 1999). An array of field plots was established by selecting one plot from each hexagon based on the following rules: (1) if an FHM plot (Mangold 1998) fell within a hexagon, it was selected; (2) if no FHM plot fell within a hexagon, the existing NRS-FIA plot from the most recent periodic inventory nearest the hexagon center was selected; and (3) if neither FHM nor existing NRS-FIA plots fell within the hexagon, a new NRS-FIA plot was established in the hexagon (McRoberts 1999). This array of plots is designated the federal base sample and is considered an equal probability sample; its measurement is funded by the Federal government. Of the 6,067 phase 2 field plots measured for the 2005 annual inventory, 618 plots contained forest land.

The total federal base sample was systematically divided into five interpenetrating, nonoverlapping subsamples or panels. Each year, the plots in a single panel are measured; panels are selected on a 5-year, rotating basis (McRoberts 1999). For estimation purposes, the measurement of each panel of plots can be considered an independent systematic sample of all land in a state. Field crews measure vegetation on plots forested at the time of the last inventory and on plots currently classified as forest by trained photointerpreters using aerial photos or digital orthoquads.

### **Phase 3**

One of every 16 phase 2 plots is also a phase 3 plot. Using a sampling intensity of one plot for approximately every 96,000 acres, phase 3 plots are measured with the full suite of FHM vegetative and health variables (Mangold 1998) collected as well as the full array of measures associated with phase 2 plots. Phase 3 plots must be measured between June 1 and August 30 to accommodate the additional measurement of nonwoody, understory vegetation; ground cover; soils; and other variables. The five panels of measurements that make up this inventory include 374 phase 3 plots, of which 40 contained forest land.

The new national FIA plot configuration (fig. 5) was first used for data collection in Iowa in 1999, the first annual inventory year, and will continue to be used in subsequent years. The national plot configuration requires mapping forest conditions on each plot.

The overall plot layout for the new configuration consists of four subplots. The centers of subplots 2, 3, and 4 are located 120 feet from the center of subplot 1. The azimuths to subplots 2, 3, and 4 are 0, 120, and 240 degrees, respectively, from the center of subplot 1.

Trees with a d.b.h. 5 inches or larger are measured on a 24-foot-radius (1/24-acre) circular subplot. All trees with a d.b.h. 1 inch or larger but less than 5 inches are measured on a 6.8-foot-radius (1/300-acre) circular microplot located 12 feet east of the center of each of the four subplots. Seedlings (trees less than 1 inch in d.b.h. and at least 6 inches tall for softwood species and 12 inches tall for hardwood species) are counted but not individually measured on this same microplot. Forest conditions on all subplots are recorded. Factors that differentiate forest conditions are changes in forest type, stand-size class, land use, ownership, and density. Each condition that occurs anywhere on any subplot is identified, described, and mapped if the area of the condition meets or exceeds 1 acre in size.

Field plot measurements are combined with phase 1 estimates in the compilation process and table production. The number of tables presented here is limited. However, other tabular data can be generated at <http://nrs.fs.fed.us/fia/data-tools/mapping-tools/default.asp>.

For additional information, contact:  
Program Manager, Forest Inventory and Analysis, Northern Research Station, 1992 Folwell Avenue, St. Paul, MN 55108 or Bureau Chief of Forestry, Iowa Department of Natural Resources, Bureau of Forestry, Wallace State Office Building, 502 East 9th Street, Des Moines, IA 50319-0034 (<http://www.iowadnr.com>).




Figure 5.—Current NRS-FIA field plot design.

## LITERATURE CITED

- Bechtold, W.A.; Patterson, P.L., eds. 2005. **The enhanced Forest Inventory and Analysis program—national sampling design and estimation procedures**. Gen. Tech. Rep. SRS-80. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 85 p.
- DellaSala, D.A.; Staus, N.L.; Strittholt, J.R.; Hackman, A.; Iacobelli, A. 2001. **An updated protected areas database for the United States and Canada**. *Natural Areas Journal*. 21(2): 124-135.
- Haugen, L. 1998. **How to identify, prevent, and control Dutch elm disease**. NA-PR-07-98. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area, State and Private Forestry. 26 p. [http://www.na.fs.fed.us/spfo/pubs/howtos/ht\\_ded/ht\\_ded.htm](http://www.na.fs.fed.us/spfo/pubs/howtos/ht_ded/ht_ded.htm)
- Leatherberry, E.C.; Roussopoulos, S.M.; Spencer, J.S., Jr. 1992. **An analysis of Iowa's forest resources, 1990**. Resour. Bull. NC-142. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 67 p.

- Leatherberry, E.C.; Brand, G.J.; Pennington, S. 2005. **Iowa's forest resources in 2003**. Resour. Bull. NC-240. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Research Station. 31 p.
- Leatherberry, E.C.; Pennington, S.; Flickinger, A.; Perry, C.H.; Jepson, E.; Woodall, C.W. 2006. **Iowa's forests, 1999-2003, part A**. Resour. Bull. NC-266A. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Research Station. 84 p.
- Mangold, R.D. 1998. **Forest health monitoring field methods guide (National 1998)**. Research Triangle Park, NC: U.S. Department of Agriculture, Forest Service, National Forest Health Monitoring Program. 429 p.
- McManus, M.; Schneeberger, N.; Reardon, R.; Mason, G. 1992. **Gypsy moth**. For. Insect and Dis. Leaflet. 162. Washington, DC: U.S. Department of Agriculture, Forest Service. 13 p.
- McRoberts, R.E. 1999. **Joint annual forest inventory and monitoring system, the North Central perspective**. Journal of Forestry. 97(12): 27-31.
- O'Brien, J.G.; Mielke, M.E.; Oak, S.; Moltzan, B. 2002. **Sudden oak death: pest alert**. NA-PR-02-02. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area, State and Private Forestry. 2 p.
- O'Brien, J.G.; Mielke, M.E.; Starkey, D.; Juzwik, J. 2000. **How to identify, prevent, and control oak wilt**. NA-PR-03-00. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area, State and Private Forestry. 28 p.
- Spencer, J.S., Jr; Jakes, P.J. 1980. **Iowa forest resources, 1974**. Resour. Bull. NC-52. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 90 p.
- Thornton, P.L.; Morgan, J.T. 1959. **The forest resources of Iowa**. For. Surv. Rel. 22. Columbus, OH: U.S. Department of Agriculture, Forest Service, Central States Forest Experiment Station. 46 p.
- U.S. Census Bureau. 2006. **State and county quick facts**. Last revised: 8 Jun 2006. Washington, DC: U.S. Census Bureau. <http://quickfacts.census.gov/qfd/states/19000.html>.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service. 2006. **Gypsy Moth—European gypsy moth, background**. <http://www.aphis.usda.gov/ppq/ispm/gm/egm-background.html>.

## TABLE TITLES

Table 1.—*Area of forest land by forest-type group, forest type, and owner category, Iowa, 2001-2005*

Table 2.—*Area of timberland by major forest-type group, stand origin, and owner category, Iowa, 2001-2005*

Table 3.—*Area of timberland by forest-type group, forest type, and stand-size class, Iowa, 2001-2005*

Table 4.—*Net volume of all live trees on forest land by species group, species, and owner category, Iowa, 2001-2005*

Table 5.—*Net volume of all live trees and salvageable dead trees on timberland by class of timber and softwood/hardwood species category, Iowa, 2001-2005*

Table 6.—*Net volume of growing stock on timberland by forest-type group, forest type, and softwood/hardwood species category, Iowa, 2001-2005*

Table 7.—*Net volume of growing stock on timberland by species group, species, and diameter class, Iowa, 2001-2005*

Table 8.—*Net volume of sawtimber on timberland by species group, species, and diameter class, Iowa, 2001-2005*

Table 9.—*All live aboveground tree biomass on timberland by owner category, softwood/hardwood species category, and tree biomass component, Iowa, 2001-2005*

Table 10.—*Average annual net growth of growing stock on timberland by species group and owner category, Iowa, 1999-2000 to 2004-2005*

Table 11.—*Average annual removals of growing stock on timberland by species group and owner category, Iowa, 1999-2000 to 2004-2005*

Table 12.—*Average annual mortality of growing stock on timberland by species group and owner category, Iowa, 1999-2000 to 2004-2005*

## TABLES

Table 1. -- Area of forest land by forest type group, forest type, and owner category, Iowa, 2001-2005

(In thousand acres)

| Forest type group/<br>forest type        | Owner category |              |                       |
|------------------------------------------|----------------|--------------|-----------------------|
|                                          | All<br>owners  | Public       | Unidentified<br>owner |
| <b>Softwood type groups</b>              |                |              |                       |
| Pinyon / juniper group                   |                |              |                       |
| Eastern redcedar                         | 39.7           | 1.8          | 38.0                  |
| All forest types                         | 39.7           | 1.8          | 38.0                  |
| All softwood groups                      | 39.7           | 1.8          | 38.0                  |
| <b>Hardwood type groups</b>              |                |              |                       |
| Oak / pine group                         |                |              |                       |
| Eastern redcedar / hardwood              | 67.9           | 9.7          | 58.2                  |
| All forest types                         | 67.9           | 9.7          | 58.2                  |
| Oak / hickory group                      |                |              |                       |
| Post oak / blackjack oak                 | 11.2           | --           | 11.2                  |
| White oak / red oak / hickory            | 589.7          | 51.7         | 518.0                 |
| White oak                                | 159.8          | 35.9         | 123.9                 |
| Northern red oak                         | 48.8           | 18.6         | 30.2                  |
| Bur oak                                  | 163.3          | 2.5          | 160.8                 |
| Black walnut                             | 90.5           | 16.3         | 74.2                  |
| Black locust                             | 11.7           | --           | 11.7                  |
| Mixed upland hardwoods                   | 509.6          | 18.1         | 491.6                 |
| All forest types                         | 1,564.6        | 143.0        | 1,421.6               |
| Elm / ash / cottonwood group             |                |              |                       |
| Elm / ash / cottonwood group             | 1.5            | --           | 1.5                   |
| Black ash / American elm / red maple     | 11.5           | 3.6          | 7.8                   |
| River birch / sycamore                   | 36.8           | --           | 36.8                  |
| Cottonwood                               | 80.8           | 16.0         | 64.8                  |
| Willow                                   | 18.2           | 5.7          | 12.4                  |
| Sycamore / pecan / American elm          | 8.7            | 1.7          | 7.0                   |
| Sugarberry / hackberry / elm / green ash | 323.7          | 29.4         | 294.4                 |
| Silver maple / American elm              | 182.1          | 32.7         | 149.4                 |
| Cottonwood / willow                      | 24.0           | --           | 24.0                  |
| All forest types                         | 687.1          | 89.1         | 598.0                 |
| Maple / beech / birch group              |                |              |                       |
| Sugar maple / beech / yellow birch       | 148.3          | 13.0         | 135.3                 |
| Black cherry                             | 17.8           | 9.0          | 8.7                   |
| Cherry / ash / yellow-poplar             | 11.1           | --           | 11.1                  |
| Hard maple / basswood                    | 149.1          | 25.9         | 123.2                 |
| Elm / ash / locust                       | 84.9           | 16.9         | 67.9                  |
| All forest types                         | 411.2          | 64.9         | 346.3                 |
| Aspen / birch group                      |                |              |                       |
| Aspen                                    | 13.0           | --           | 13.0                  |
| All forest types                         | 13.0           | --           | 13.0                  |
| Exotic hardwoods group                   |                |              |                       |
| Other exotic hardwoods                   | 9.9            | 4.8          | 5.1                   |
| All forest types                         | 9.9            | 4.8          | 5.1                   |
| All hardwood groups                      | 2,753.8        | 311.6        | 2,442.2               |
| Nonstocked                               | 85.4           | 13.8         | 71.6                  |
| <b>All forest groups</b>                 | <b>2,878.9</b> | <b>327.2</b> | <b>2,551.8</b>        |

(Table 1 continued on next page)

(Table 1 continued)

| Forest type group/<br>forest type | Owner category |        |                       |
|-----------------------------------|----------------|--------|-----------------------|
|                                   | All<br>owners  | Public | Private               |
| Hardwood type groups              |                |        | Unidentified<br>owner |

All table cells without observations in the inventory sample are indicated by --. Table value of 0.0 indicates the acres round to less than 0.1 thousand acres. Columns and rows may not add to their totals due to rounding.

Table 2. -- Area of timberland by major forest type group, stand origin, and owner category, Iowa, 2001-2005

(In thousand acres)

| Major forest type group<br>and stand origin | Owner category |        |                                  |
|---------------------------------------------|----------------|--------|----------------------------------|
|                                             | All<br>owners  | Public | Private<br>Unidentified<br>owner |
| <b>Softwood type groups</b>                 |                |        |                                  |
| Natural                                     | 33.1           | 1.8    | 31.3                             |
| All softwood types                          | 33.1           | 1.8    | 31.3                             |
| <b>Hardwood type groups</b>                 |                |        |                                  |
| Natural                                     | 2,697.7        | 299.9  | 2,397.7                          |
| Planted                                     | 20.0           | 3.2    | 16.8                             |
| All hardwood types                          | 2,717.7        | 303.2  | 2,414.5                          |
| Nonstocked                                  | 72.9           | 7.3    | 65.6                             |
| <b>All groups</b>                           | 2,823.7        | 312.3  | 2,511.4                          |

All table cells without observations in the inventory sample are indicated by --. Table value of 0.0 indicates the acres round to less than 0.1 thousand acres. Columns and rows may not add to their totals due to rounding.

Table 3. -- Area of timberland by forest type group, forest type, and stand-size class, Iowa, 2001-2005

(In thousand acres)

| Forest type group/<br>forest type        | Stand-size class |                |              |                      |                 |
|------------------------------------------|------------------|----------------|--------------|----------------------|-----------------|
|                                          | All<br>stands    | Sawtimber      | Poletimber   | Sapling-<br>seedling | Non-<br>stocked |
| <b>Softwood type groups</b>              |                  |                |              |                      |                 |
| Pinyon / juniper group                   |                  |                |              |                      |                 |
| Eastern redcedar                         | 33.1             | 1.8            | 5.3          | 26.0                 | --              |
| All forest types                         | 33.1             | 1.8            | 5.3          | 26.0                 | --              |
| All softwood groups                      | 33.1             | 1.8            | 5.3          | 26.0                 | --              |
| <b>Hardwood type groups</b>              |                  |                |              |                      |                 |
| Oak / pine group                         |                  |                |              |                      |                 |
| Eastern redcedar / hardwood              | 62.0             | 20.7           | 9.5          | 31.8                 | --              |
| All forest types                         | 62.0             | 20.7           | 9.5          | 31.8                 | --              |
| Oak / hickory group                      |                  |                |              |                      |                 |
| Post oak / blackjack oak                 | 11.2             | 9.8            | 1.4          | --                   | --              |
| White oak / red oak / hickory            | 555.2            | 381.9          | 105.9        | 67.4                 | --              |
| White oak                                | 153.3            | 143.3          | 10.0         | --                   | --              |
| Northern red oak                         | 48.8             | 48.8           | --           | --                   | --              |
| Bur oak                                  | 154.0            | 130.4          | 14.5         | 9.1                  | --              |
| Black walnut                             | 90.5             | 67.8           | 8.3          | 14.4                 | --              |
| Black locust                             | 11.7             | 5.1            | 6.6          | --                   | --              |
| Mixed upland hardwoods                   | 509.6            | 257.3          | 172.8        | 79.6                 | --              |
| All forest types                         | 1,534.5          | 1,044.4        | 319.6        | 170.4                | --              |
| Elm / ash / cottonwood group             |                  |                |              |                      |                 |
| Elm / ash / cottonwood group             | 1.5              | --             | 1.5          | --                   | --              |
| Black ash / American elm / red maple     | 11.5             | 8.7            | 2.2          | 0.6                  | --              |
| River birch / sycamore                   | 36.8             | 25.0           | 11.8         | --                   | --              |
| Cottonwood                               | 80.8             | 69.2           | 4.7          | 6.9                  | --              |
| Willow                                   | 18.2             | 8.8            | 6.5          | 2.8                  | --              |
| Sycamore / pecan / American elm          | 8.7              | 8.4            | 0.3          | --                   | --              |
| Sugarberry / hackberry / elm / green ash | 323.7            | 178.0          | 117.0        | 28.8                 | --              |
| Silver maple / American elm              | 182.1            | 162.8          | 13.3         | 6.0                  | --              |
| Cottonwood / willow                      | 24.0             | 13.3           | 10.7         | --                   | --              |
| All forest types                         | 687.1            | 474.1          | 167.9        | 45.1                 | --              |
| Maple / beech / birch group              |                  |                |              |                      |                 |
| Sugar maple / beech / yellow birch       | 148.3            | 79.6           | 44.2         | 24.5                 | --              |
| Black cherry                             | 17.8             | 7.6            | 1.5          | 8.8                  | --              |
| Cherry / ash / yellow-poplar             | 11.1             | --             | --           | 11.1                 | --              |
| Hard maple / basswood                    | 149.1            | 133.7          | 14.0         | 1.4                  | --              |
| Elm / ash / locust                       | 84.9             | 38.9           | 23.5         | 22.4                 | --              |
| All forest types                         | 411.2            | 259.7          | 83.2         | 68.2                 | --              |
| Aspen / birch group                      |                  |                |              |                      |                 |
| Aspen                                    | 13.0             | 13.0           | --           | --                   | --              |
| All forest types                         | 13.0             | 13.0           | --           | --                   | --              |
| Exotic hardwoods group                   |                  |                |              |                      |                 |
| Other exotic hardwoods                   | 9.9              | 5.1            | --           | 4.8                  | --              |
| All forest types                         | 9.9              | 5.1            | --           | 4.8                  | --              |
| All hardwood groups                      | 2,717.7          | 1,817.0        | 580.3        | 320.3                | --              |
| Nonstocked                               | 72.9             | --             | --           | --                   | 72.9            |
| <b>All forest groups</b>                 | <b>2,823.7</b>   | <b>1,818.8</b> | <b>585.6</b> | <b>346.3</b>         | <b>72.9</b>     |

All table cells without observations in the inventory sample are indicated by --. Table value of 0.0 indicates the acres round to less than 0.1 thousand acres. Columns and rows may not add to their totals due to rounding.

Table 4. -- Net volume of all live trees on forest land by species group, species, and owner category, Iowa, 2001-2005

(In thousand cubic feet)

| Species group/<br>species   | Owner category |        |         |                       |
|-----------------------------|----------------|--------|---------|-----------------------|
|                             | All<br>owners  | Public | Private | Unidentified<br>owner |
| <b>Softwoods</b>            |                |        |         |                       |
| Other yellow pines          |                |        |         |                       |
| Scotch pine                 | 150            | --     | 150     | --                    |
| All species                 | 150            | --     | 150     | --                    |
| Eastern white and red pines |                |        |         |                       |
| Red pine                    | 849            | 393    | 455     | --                    |
| Eastern white pine          | 236            | --     | 236     | --                    |
| All species                 | 1,085          | 393    | 692     | --                    |
| Other eastern softwoods     |                |        |         |                       |
| Eastern redcedar            | 38,845         | 5,982  | 32,862  | --                    |
| Larch spp.                  | 460            | --     | 460     | --                    |
| Tamarack (native)           | 247            | --     | 247     | --                    |
| All species                 | 39,552         | 5,982  | 33,569  | --                    |
| Ponderosa and Jeffrey pines |                |        |         |                       |
| Ponderosa pine              | 3,461          | 552    | 2,910   | --                    |
| All species                 | 3,461          | 552    | 2,910   | --                    |
| Total softwoods             | 44,248         | 6,927  | 37,321  | --                    |
| <b>Hardwoods</b>            |                |        |         |                       |
| Select white oaks           |                |        |         |                       |
| White oak                   | 369,496        | 72,095 | 297,400 | --                    |
| Swamp white oak             | 4,301          | 69     | 4,232   | --                    |
| Bur oak                     | 468,143        | 12,770 | 455,373 | --                    |
| Chinkapin oak               | 3,933          | 1,957  | 1,977   | --                    |
| All species                 | 845,872        | 86,891 | 758,982 | --                    |
| Select red oaks             |                |        |         |                       |
| Northern red oak            | 272,706        | 64,261 | 208,445 | --                    |
| All species                 | 272,706        | 64,261 | 208,445 | --                    |
| Other white oaks            |                |        |         |                       |
| Post oak                    | 683            | --     | 683     | --                    |
| All species                 | 683            | --     | 683     | --                    |
| Other red oaks              |                |        |         |                       |
| Northern pin oak            | 30,641         | 2,336  | 28,305  | --                    |
| Shingle oak                 | 34,882         | 80     | 34,802  | --                    |
| Pin oak                     | 21,755         | 8,548  | 13,207  | --                    |
| Black oak                   | 86,911         | 11,179 | 75,732  | --                    |
| All species                 | 174,190        | 22,144 | 152,046 | --                    |
| Hickory                     |                |        |         |                       |
| Bitternut hickory           | 81,450         | 7,306  | 74,144  | --                    |
| Pignut hickory              | 2,251          | 1,092  | 1,159   | --                    |
| Shagbark hickory            | 162,354        | 15,755 | 146,599 | --                    |
| Mockernut hickory           | 2,326          | 604    | 1,722   | --                    |
| All species                 | 248,381        | 24,757 | 223,624 | --                    |
| Hard maple                  |                |        |         |                       |
| Black maple                 | 61,381         | 4,425  | 56,955  | --                    |
| Sugar maple                 | 37,395         | 12,905 | 24,489  | --                    |
| All species                 | 98,775         | 17,331 | 81,445  | --                    |

(Table 4 continued on next page)

(Table 4 continued)

| Species group/<br>species    | Owner category |        |         |                       |
|------------------------------|----------------|--------|---------|-----------------------|
|                              | All<br>owners  | Public | Private | Unidentified<br>owner |
| <b>Hardwoods</b>             |                |        |         |                       |
| Soft maple                   |                |        |         |                       |
| Red maple                    | 665            | 665    | --      | --                    |
| Silver maple                 | 445,353        | 81,305 | 364,048 | --                    |
| All species                  | 446,018        | 81,970 | 364,048 | --                    |
| Ash                          |                |        |         |                       |
| White ash                    | 57,288         | 4,260  | 53,028  | --                    |
| Black ash                    | 4,408          | 1,208  | 3,201   | --                    |
| Green ash                    | 73,452         | 17,909 | 55,543  | --                    |
| All species                  | 135,149        | 23,377 | 111,772 | --                    |
| Cottonwood and aspen         |                |        |         |                       |
| Eastern cottonwood           | 338,956        | 90,314 | 248,641 | --                    |
| Bigtooth aspen               | 14,976         | --     | 14,976  | --                    |
| Quaking aspen                | 3,715          | --     | 3,715   | --                    |
| All species                  | 357,647        | 90,314 | 267,333 | --                    |
| Basswood                     |                |        |         |                       |
| American basswood            | 174,115        | 29,814 | 144,300 | --                    |
| All species                  | 174,115        | 29,814 | 144,300 | --                    |
| Black walnut                 |                |        |         |                       |
| Black walnut                 | 249,351        | 25,285 | 224,065 | --                    |
| All species                  | 249,351        | 25,285 | 224,065 | --                    |
| Other eastern soft hardwoods |                |        |         |                       |
| Boxelder                     | 118,751        | 6,720  | 112,030 | --                    |
| Ohio buckeye                 | 9,918          | 126    | 9,792   | --                    |
| River birch                  | 4,545          | --     | 4,545   | --                    |
| Paper birch                  | 2,918          | --     | 2,918   | --                    |
| Northern catalpa             | 815            | --     | 815     | --                    |
| Hackberry                    | 185,000        | 9,086  | 175,913 | --                    |
| Butternut                    | 778            | --     | 778     | --                    |
| American sycamore            | 3,776          | 313    | 3,463   | --                    |
| Black cherry                 | 61,268         | 8,247  | 53,021  | --                    |
| Black willow                 | 50,929         | 2,653  | 48,276  | --                    |
| White willow                 | 475            | --     | 475     | --                    |
| American elm                 | 240,013        | 27,224 | 212,790 | --                    |
| Siberian elm                 | 9,361          | 1,048  | 8,314   | --                    |
| Slippery elm                 | 106,662        | 10,679 | 95,983  | --                    |
| All species                  | 795,210        | 66,096 | 729,113 | --                    |
| Other eastern hard hardwoods |                |        |         |                       |
| Honeylocust                  | 125,026        | 17,710 | 107,316 | --                    |
| White mulberry               | 460            | --     | 460     | --                    |
| Red mulberry                 | 40,937         | 6,343  | 34,595  | --                    |
| Black locust                 | 42,674         | 6,530  | 36,143  | --                    |
| All species                  | 209,097        | 30,583 | 178,514 | --                    |

(Table 4 continued on next page)

(Table 4 continued)

| Species group/<br>species       | Owner category |         |           |                       |
|---------------------------------|----------------|---------|-----------|-----------------------|
|                                 | All<br>owners  | Public  | Private   | Unidentified<br>owner |
| <b>Hardwoods</b>                |                |         |           |                       |
| Eastern noncommercial hardwoods |                |         |           |                       |
| Ailanthus                       | 162            | --      | 162       | --                    |
| Serviceberry spp.               | 117            | 117     | --        | --                    |
| Eastern redbud                  | 564            | 23      | 540       | --                    |
| Hawthorn spp.                   | 917            | --      | 917       | --                    |
| Cockspur hawthorn               | 788            | --      | 788       | --                    |
| Downy hawthorn                  | 69             | --      | 69        | --                    |
| Osage-orange                    | 27,634         | 175     | 27,459    | --                    |
| Apple spp.                      | 336            | --      | 336       | --                    |
| Eastern hophornbeam             | 14,791         | 1,382   | 13,409    | --                    |
| American plum                   | 260            | --      | 260       | --                    |
| Willow spp.                     | 239            | 239     | --        | --                    |
| Peachleaf willow                | 243            | --      | 243       | --                    |
| Russian-olive                   | 2,008          | --      | 2,008     | --                    |
| All species                     | 48,128         | 1,936   | 46,192    | --                    |
| Total hardwoods                 | 4,055,320      | 564,759 | 3,490,561 | --                    |
| <b>All species groups</b>       | 4,099,569      | 571,687 | 3,527,882 | --                    |

All table cells without observations in the inventory sample are indicated by --. Table value of 0 indicates the volume rounds to less than 1 thousand cubic feet. Columns and rows may not add to their totals due to rounding.

Table 5. -- Net volume of all live trees and salvageable dead trees on timberland by class of timber and softwood/hardwood species category, Iowa, 2001-2005

(In thousand cubic feet)

| Class of timber                   | All species      | Softwood species | Hardwood species |
|-----------------------------------|------------------|------------------|------------------|
| <b>Live trees</b>                 |                  |                  |                  |
| <b>Growing-stock trees</b>        |                  |                  |                  |
| Sawtimber                         |                  |                  |                  |
| Saw log portion                   | 2,165,489        | 16,219           | 2,149,270        |
| Upper stem portion                | 292,547          | 2,360            | 290,187          |
| Total                             | 2,458,036        | 18,579           | 2,439,457        |
| Poletimber                        | 656,464          | 10,473           | 645,991          |
| <b>All growing-stock trees</b>    | <b>3,114,499</b> | <b>29,051</b>    | <b>3,085,448</b> |
| <b>Cull trees</b>                 |                  |                  |                  |
| Rough trees <sup>1</sup>          |                  |                  |                  |
| Sawtimber size                    | 616,258          | 8,088            | 608,170          |
| Poletimber size                   | 242,471          | 6,902            | 235,569          |
| Total                             | 858,729          | 14,990           | 843,740          |
| Rotten trees <sup>1</sup>         |                  |                  |                  |
| Sawtimber size                    | 60,830           | --               | 60,830           |
| Poletimber size                   | 11,032           | 116              | 10,916           |
| Total                             | 71,862           | 116              | 71,745           |
| <b>All live trees</b>             | <b>930,591</b>   | <b>15,106</b>    | <b>915,485</b>   |
| <b>All live trees</b>             | <b>4,045,090</b> | <b>44,157</b>    | <b>4,000,933</b> |
| <b>Salvageable dead trees</b>     |                  |                  |                  |
| Sawtimber size                    | 35,763           | 654              | 35,109           |
| Poletimber size                   | 21,676           | 295              | 21,380           |
| <b>All salvageable dead trees</b> | <b>57,438</b>    | <b>949</b>       | <b>56,489</b>    |
| <b>All classes</b>                | <b>4,102,528</b> | <b>45,106</b>    | <b>4,057,422</b> |

All table cells without observations in the inventory sample are indicated by --. Table value of 0 indicates the volume rounds to less than 1 thousand cubic feet. Columns and rows may not add to their totals due to rounding.

<sup>1</sup> Includes noncommercial species.

Table 6. -- Net volume of growing stock on timberland by forest type group, forest type, and softwood/hardwood species category, Iowa, 2001-2005

(In thousand cubic feet)

| Forest type group/<br>forest type        | All<br>species | Softwood<br>species | Hardwood<br>species |
|------------------------------------------|----------------|---------------------|---------------------|
| <b>Softwood type groups</b>              |                |                     |                     |
| Pinyon / juniper group                   |                |                     |                     |
| Eastern redcedar                         | 5,702          | 3,006               | 2,696               |
| All forest types                         | 5,702          | 3,006               | 2,696               |
| All softwood groups                      | 5,702          | 3,006               | 2,696               |
| <b>Hardwood type groups</b>              |                |                     |                     |
| Oak / pine group                         |                |                     |                     |
| Eastern redcedar / hardwood              | 12,438         | 6,262               | 6,176               |
| All forest types                         | 12,438         | 6,262               | 6,176               |
| Oak / hickory group                      |                |                     |                     |
| Post oak / blackjack oak                 | 22,565         | --                  | 22,565              |
| White oak / red oak / hickory            | 647,179        | 6,781               | 640,398             |
| White oak                                | 282,454        | 520                 | 281,933             |
| Northern red oak                         | 84,942         | 117                 | 84,824              |
| Bur oak                                  | 203,655        | --                  | 203,655             |
| Black walnut                             | 89,429         | 108                 | 89,321              |
| Black locust                             | 3,336          | --                  | 3,336               |
| Mixed upland hardwoods                   | 352,011        | 3,420               | 348,591             |
| All forest types                         | 1,685,569      | 10,946              | 1,674,623           |
| Elm / ash / cottonwood group             |                |                     |                     |
| Black ash / American elm / red maple     | 16,646         | --                  | 16,646              |
| River birch / sycamore                   | 33,969         | --                  | 33,969              |
| Cottonwood                               | 257,220        | 552                 | 256,668             |
| Willow                                   | 4,788          | --                  | 4,788               |
| Sycamore / pecan / American elm          | 9,549          | --                  | 9,549               |
| Sugarberry / hackberry / elm / green ash | 213,819        | 1,173               | 212,646             |
| Silver maple / American elm              | 348,635        | 2,665               | 345,971             |
| Cottonwood / willow                      | 41,974         | --                  | 41,974              |
| All forest types                         | 926,600        | 4,390               | 922,211             |
| Maple / beech / birch group              |                |                     |                     |
| Sugar maple / beech / yellow birch       | 133,531        | 2,977               | 130,554             |
| Black cherry                             | 9,460          | --                  | 9,460               |
| Cherry / ash / yellow-poplar             | 1,400          | --                  | 1,400               |
| Hard maple / basswood                    | 259,185        | 747                 | 258,438             |
| Elm / ash / locust                       | 62,182         | --                  | 62,182              |
| All forest types                         | 465,759        | 3,724               | 462,035             |
| Aspen / birch group                      |                |                     |                     |
| Aspen                                    | 13,830         | 170                 | 13,660              |
| All forest types                         | 13,830         | 170                 | 13,660              |

(Table 6 continued on next page)

(Table 6 continued)

| Forest type group/<br>forest type | All<br>species | Softwood<br>species | Hardwood<br>species |
|-----------------------------------|----------------|---------------------|---------------------|
| <b>Hardwood type groups</b>       |                |                     |                     |
| Exotic hardwoods group            |                |                     |                     |
| Other exotic hardwoods            | 2,370          | 146                 | 2,224               |
| All forest types                  | 2,370          | 146                 | 2,224               |
| All hardwood groups               | 3,106,566      | 25,638              | 3,080,928           |
| Nonstocked                        | 2,231          | 407                 | 1,824               |
| <b>All forest groups</b>          | 3,114,499      | 29,051              | 3,085,448           |

All table cells without observations in the inventory sample are indicated by --. Table value of 0 indicates the volume rounds to less than 1 thousand cubic feet. Columns and rows may not add to their totals due to rounding.

Table 7. -- Net volume of growing stock on timberland by species group, species, and diameter class, Iowa, 2001-2005

(In thousand cubic feet)

| Species group/<br>species   | All<br>classes | Diameter class (inches at breast height) |         |          |           |           |           |           |           |           |         |    | 29.0+ |
|-----------------------------|----------------|------------------------------------------|---------|----------|-----------|-----------|-----------|-----------|-----------|-----------|---------|----|-------|
|                             |                | 5.0-5.9                                  | 7.0-8.9 | 9.0-10.9 | 11.0-12.9 | 13.0-14.9 | 15.0-16.9 | 17.0-18.9 | 19.0-20.9 | 21.0-28.9 |         |    |       |
| Softwoods                   |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Other yellow pines          |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Scotch pine                 | 150            | --                                       | 150     | --       | --        | --        | --        | --        | --        | --        | --      | -- | --    |
| All species                 | 150            | --                                       | 150     | --       | --        | --        | --        | --        | --        | --        | --      | -- | --    |
| Eastern white and red pines |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Red pine                    | 849            | --                                       | 142     | 251      | 455       | --        | --        | --        | --        | --        | --      | -- | --    |
| All species                 | 849            | --                                       | 142     | 251      | 455       | --        | --        | --        | --        | --        | --      | -- | --    |
| Other eastern softwoods     |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Eastern redcedar            | 24,221         | 5,459                                    | 4,477   | 5,702    | 4,020     | 3,548     | 1,015     | --        | --        | --        | --      | -- | --    |
| Larch spp.                  | 371            | --                                       | --      | 371      | --        | --        | --        | --        | --        | --        | --      | -- | --    |
| All species                 | 24,591         | 5,459                                    | 4,477   | 6,072    | 4,020     | 3,548     | 1,015     | --        | --        | --        | --      | -- | --    |
| Ponderosa and Jeffrey pines |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Ponderosa pine              | 3,461          | 108                                      | 137     | 576      | 436       | 2,205     | --        | --        | --        | --        | --      | -- | --    |
| All species                 | 3,461          | 108                                      | 137     | 576      | 436       | 2,205     | --        | --        | --        | --        | --      | -- | --    |
| Total softwoods             | 29,051         | 5,567                                    | 4,906   | 6,899    | 4,912     | 5,733     | 1,015     | --        | --        | --        | --      | -- | --    |
| Hardwoods                   |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Select white oaks           |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| White oak                   | 311,299        | 4,757                                    | 11,576  | 30,487   | 42,819    | 46,598    | 35,631    | 39,374    | 40,521    | 58,536    | --      | -- | --    |
| Swamp white oak             | 3,621          | 378                                      | 228     | 426      | 618       | --        | --        | 1,971     | --        | --        | --      | -- | --    |
| Bar oak                     | 301,710        | 4,328                                    | 10,478  | 14,091   | 23,288    | 26,602    | 20,100    | 34,487    | 43,470    | 85,133    | 39,732  | -- | --    |
| Chinkapin oak               | 1,744          | 261                                      | 725     | 274      | 484       | --        | --        | --        | --        | --        | --      | -- | --    |
| All species                 | 618,733        | 9,724                                    | 23,007  | 45,277   | 67,209    | 73,200    | 55,731    | 75,383    | 83,992    | 144,668   | 39,732  | -- | --    |
| Select red oaks             |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Northern red oak            | 240,748        | 4,344                                    | 3,737   | 13,360   | 21,825    | 29,164    | 33,252    | 34,627    | 19,592    | 63,162    | 17,684  | -- | --    |
| All species                 | 240,748        | 4,344                                    | 3,737   | 13,360   | 21,825    | 29,164    | 33,252    | 34,627    | 19,592    | 63,162    | 17,684  | -- | --    |
| Other white oaks            |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Post oak                    | 683            | --                                       | 365     | 318      | --        | --        | --        | --        | --        | --        | --      | -- | --    |
| All species                 | 683            | --                                       | 365     | 318      | --        | --        | --        | --        | --        | --        | --      | -- | --    |
| Other red oaks              |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Northern pin oak            | 26,488         | 588                                      | 770     | 1,462    | 5,058     | 1,668     | 5,075     | 2,692     | 1,885     | 7,279     | --      | -- | --    |
| Shingle oak                 | 27,692         | 3,638                                    | 4,016   | 5,833    | 1,677     | 3,049     | 2,499     | 5,104     | 1,876     | --        | --      | -- | --    |
| Pin oak                     | 21,755         | 431                                      | 624     | 327      | 1,652     | --        | --        | 5,068     | 2,578     | 3,671     | 7,404   | -- | --    |
| Black oak                   | 70,805         | 1,720                                    | 3,969   | 7,060    | 8,673     | 17,025    | 11,977    | 5,657     | 9,969     | 4,754     | --      | -- | --    |
| All species                 | 146,740        | 6,377                                    | 9,380   | 14,683   | 17,060    | 21,742    | 19,551    | 18,321    | 16,319    | 15,703    | 7,404   | -- | --    |
| Hickory                     |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Bitternut hickory           | 68,453         | 7,424                                    | 12,625  | 11,298   | 11,504    | 10,794    | 2,547     | 5,922     | 2,414     | 3,924     | --      | -- | --    |
| Pignut hickory              | 2,003          | --                                       | 467     | 444      | --        | 1,092     | --        | --        | --        | --        | --      | -- | --    |
| Shagbark hickory            | 147,051        | 14,281                                   | 21,635  | 25,500   | 26,903    | 23,480    | 15,111    | 7,254     | 6,693     | 6,195     | --      | -- | --    |
| Mockernut hickory           | 2,326          | 662                                      | --      | 462      | 598       | 604       | --        | --        | --        | --        | --      | -- | --    |
| All species                 | 219,833        | 22,367                                   | 34,727  | 37,703   | 39,005    | 35,970    | 17,658    | 13,177    | 9,108     | 10,119    | --      | -- | --    |
| Hard maple                  |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Black maple                 | 47,714         | 2,735                                    | 3,377   | 7,825    | 9,368     | 6,227     | 7,535     | 7,610     | --        | 3,036     | --      | -- | --    |
| Sugar maple                 | 31,711         | 1,990                                    | 3,938   | 2,193    | 620       | 1,900     | 3,522     | 5,703     | 2,770     | 9,076     | --      | -- | --    |
| All species                 | 79,425         | 4,724                                    | 7,315   | 10,018   | 9,988     | 8,127     | 11,057    | 13,313    | 2,770     | 12,113    | --      | -- | --    |
| Soft maple                  |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Red maple                   | 665            | --                                       | --      | 665      | --        | --        | --        | --        | --        | --        | --      | -- | --    |
| Silver maple                | 318,606        | 4,805                                    | 11,179  | 21,952   | 30,925    | 45,481    | 41,274    | 34,984    | 42,305    | 48,236    | 37,465  | -- | --    |
| All species                 | 319,271        | 4,805                                    | 11,179  | 22,616   | 30,925    | 45,481    | 41,274    | 34,984    | 42,305    | 48,236    | 37,465  | -- | --    |
| Ash                         |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| White ash                   | 46,591         | 2,215                                    | 6,351   | 9,338    | 3,847     | 10,574    | 4,608     | --        | 2,350     | --        | 7,308   | -- | --    |
| Black ash                   | 3,272          | 445                                      | 619     | --       | --        | 1,052     | --        | 1,156     | --        | --        | --      | -- | --    |
| Green ash                   | 60,050         | 5,426                                    | 6,940   | 9,572    | 5,953     | 11,052    | 9,547     | 7,099     | 4,460     | --        | --      | -- | --    |
| All species                 | 109,913        | 8,087                                    | 13,910  | 18,909   | 9,800     | 22,678    | 14,156    | 8,255     | 6,810     | --        | 7,308   | -- | --    |
| Cottonwood and aspen        |                |                                          |         |          |           |           |           |           |           |           |         |    |       |
| Eastern cottonwood          | 326,117        | 550                                      | 2,548   | 7,642    | 13,797    | 18,444    | 7,562     | 21,062    | 27,048    | 100,087   | 127,376 | -- | --    |
| Bigtooth aspen              | 14,976         | 516                                      | 484     | 2,008    | 3,830     | 950       | 5,008     | --        | 2,180     | --        | --      | -- | --    |
| Quaking aspen               | 3,551          | 197                                      | 1,638   | 1,018    | 697       | --        | --        | --        | --        | --        | --      | -- | --    |
| All species                 | 344,644        | 1,264                                    | 4,671   | 10,669   | 18,324    | 19,393    | 12,570    | 21,062    | 29,228    | 100,087   | 127,376 | -- | --    |

(Table 7 continued)

| Species group/<br>species    | All<br>classes | Diameter class (inches at breast height) |         |          |           |           |           |           |           |           |         |
|------------------------------|----------------|------------------------------------------|---------|----------|-----------|-----------|-----------|-----------|-----------|-----------|---------|
|                              |                | 5.0-6.9                                  | 7.0-8.9 | 9.0-10.9 | 11.0-12.9 | 13.0-14.9 | 15.0-16.9 | 17.0-18.9 | 19.0-20.9 | 21.0-28.9 | 29.0+   |
| Hardwoods                    |                |                                          |         |          |           |           |           |           |           |           |         |
| Basswood                     |                |                                          |         |          |           |           |           |           |           |           |         |
| American basswood            | 154,591        | 6,206                                    | 11,093  | 14,498   | 13,558    | 15,921    | 17,946    | 28,751    | 6,298     | 40,320    | --      |
| All species                  | 154,591        | 6,206                                    | 11,093  | 14,498   | 13,558    | 15,921    | 17,946    | 28,751    | 6,298     | 40,320    | --      |
| Black walnut                 |                |                                          |         |          |           |           |           |           |           |           |         |
| Black walnut                 | 219,447        | 6,710                                    | 11,214  | 18,385   | 29,204    | 39,005    | 47,198    | 33,159    | 11,495    | 23,077    | --      |
| All species                  | 219,447        | 6,710                                    | 11,214  | 18,385   | 29,204    | 39,005    | 47,198    | 33,159    | 11,495    | 23,077    | --      |
| Other eastern soft hardwoods |                |                                          |         |          |           |           |           |           |           |           |         |
| Boxelder                     | 26,773         | 3,142                                    | 3,976   | 6,469    | 4,340     | 2,529     | 3,592     | --        | --        | 2,725     | --      |
| Ohio buckeye                 | 6,506          | 626                                      | 385     | --       | 1,174     | 963       | 1,179     | --        | 2,179     | --        | --      |
| River birch                  | 3,779          | 395                                      | 1,079   | 294      | --        | 798       | 1,214     | --        | --        | --        | --      |
| Paper birch                  | 2,844          | 253                                      | 146     | 284      | --        | 920       | 1,242     | --        | --        | --        | --      |
| Hackberry                    | 156,418        | 10,126                                   | 15,338  | 23,720   | 15,491    | 24,990    | 21,607    | 15,933    | 12,338    | 16,476    | --      |
| Butternut                    | 357            | 62                                       | 294     | --       | --        | --        | --        | --        | --        | --        | --      |
| American sycamore            | 3,776          | 333                                      | 313     | --       | --        | --        | --        | --        | --        | 2,778     | --      |
| Black cherry                 | 48,604         | 3,775                                    | 5,413   | 7,477    | 6,391     | 6,047     | 5,541     | 4,813     | 4,147     | --        | --      |
| Black willow                 | 26,352         | 551                                      | 853     | 1,579    | 540       | 1,949     | 1,492     | 5,102     | 2,315     | 11,972    | --      |
| White willow                 | 142            | --                                       | 142     | --       | --        | --        | --        | --        | --        | --        | --      |
| American elm                 | 167,848        | 24,450                                   | 30,580  | 27,953   | 18,318    | 20,608    | 18,902    | 11,737    | 5,490     | 9,810     | --      |
| Siberian elm                 | 3,324          | 451                                      | 332     | 557      | --        | 803       | 1,182     | --        | --        | --        | --      |
| Slippery elm                 | 79,984         | 8,897                                    | 9,686   | 11,770   | 11,951    | 15,519    | 6,091     | 6,594     | 2,176     | 7,299     | --      |
| All species                  | 521,706        | 53,060                                   | 68,776  | 80,415   | 58,205    | 75,125    | 62,041    | 44,179    | 28,845    | 51,060    | --      |
| Other eastern hard hardwoods |                |                                          |         |          |           |           |           |           |           |           |         |
| Honeylocust                  | 71,790         | 4,335                                    | 5,998   | 7,778    | 8,709     | 14,010    | 12,668    | 11,254    | 7,086     | --        | --      |
| Red mulberry                 | 4,274          | 1,449                                    | 706     | 1,056    | 1,063     | --        | --        | --        | --        | --        | --      |
| Black locust                 | 34,010         | 2,117                                    | 3,108   | 5,551    | 3,053     | 6,083     | 6,519     | 2,453     | 2,393     | 2,733     | --      |
| All species                  | 110,073        | 7,901                                    | 9,812   | 14,385   | 12,826    | 20,093    | 19,187    | 13,708    | 9,429     | 2,733     | --      |
| Total hardwoods              | 3,085,448      | 135,568                                  | 209,186 | 301,237  | 327,931   | 403,899   | 351,619   | 339,570   | 266,190   | 511,277   | 236,971 |
| All species groups           | 3,114,499      | 141,135                                  | 214,092 | 308,136  | 332,843   | 411,652   | 352,634   | 339,570   | 266,190   | 511,277   | 236,971 |

All table cells without observations in the inventory sample are indicated by --. Table value of 0 indicates the volume rounds to less than 1 thousand cubic feet. Columns and rows may not add to their totals due to rounding.

Table 8. — Net volume of sawtimber on timberland by species group, species, and diameter class, Iowa, 2001-2005

(In thousand board feet)<sup>1</sup>

| Species group/<br>species   | All<br>classes | 9.0-10.9 | 11.0-12.9 | 13.0-14.9 | 15.0-16.9 | 17.0-18.9 | 19.0-20.9 | 21.0-28.9 | 29.0+   |
|-----------------------------|----------------|----------|-----------|-----------|-----------|-----------|-----------|-----------|---------|
| Eastern white and red pines |                |          |           |           |           |           |           |           |         |
| Red pine                    | 3,432          | 1,247    | 2,185     | --        | --        | --        | --        | --        | --      |
| All species                 | 3,432          | 1,247    | 2,185     | --        | --        | --        | --        | --        | --      |
| Other eastern softwoods     |                |          |           |           |           |           |           |           |         |
| Eastern redcedar            | 73,715         | 31,357   | 20,511    | 17,165    | 4,683     | --        | --        | --        | --      |
| Larch spp.                  | 1,773          | 1,773    | --        | --        | --        | --        | --        | --        | --      |
| All species                 | 75,488         | 33,130   | 20,511    | 17,165    | 4,683     | --        | --        | --        | --      |
| Ponderosa and Jeffrey pines |                |          |           |           |           |           |           |           |         |
| Ponderosa pine              | 16,009         | 2,764    | 2,141     | 11,103    | --        | --        | --        | --        | --      |
| All species                 | 16,009         | 2,764    | 2,141     | 11,103    | --        | --        | --        | --        | --      |
| Total softwoods             |                |          |           |           |           |           |           |           |         |
|                             | 94,928         | 37,142   | 24,836    | 28,268    | 4,683     | --        | --        | --        | --      |
| Hardwoods                   |                |          |           |           |           |           |           |           |         |
| Select white oaks           |                |          |           |           |           |           |           |           |         |
| White oak                   | 1,231,641      | --       | 210,065   | 224,323   | 169,053   | 183,131   | 184,572   | 260,497   | --      |
| Swamp white oak             | 12,297         | --       | 3,056     | --        | --        | 9,242     | --        | --        | --      |
| Bur oak                     | 1,208,429      | --       | 113,536   | 127,636   | 94,437    | 159,915   | 197,087   | 364,187   | 151,631 |
| Chinkapin oak               | 2,410          | --       | 2,410     | --        | --        | --        | --        | --        | --      |
| All species                 | 2,454,777      | --       | 329,067   | 351,958   | 263,490   | 352,287   | 381,660   | 624,684   | 151,631 |
| Select red oaks             |                |          |           |           |           |           |           |           |         |
| Northern red oak            | 1,069,718      | --       | 108,055   | 145,510   | 166,420   | 172,770   | 97,079    | 301,990   | 77,893  |
| All species                 | 1,069,718      | --       | 108,055   | 145,510   | 166,420   | 172,770   | 97,079    | 301,990   | 77,893  |
| Other red oaks              |                |          |           |           |           |           |           |           |         |
| Northern pin oak            | 114,957        | --       | 25,089    | 8,298     | 25,147    | 13,424    | 9,258     | 33,740    | --      |
| Shingle oak                 | 69,885         | --       | 8,252     | 15,119    | 12,317    | 25,112    | 9,065     | --        | --      |
| Pin oak                     | 91,044         | --       | 8,131     | --        | --        | 24,809    | 12,332    | 16,906    | 28,867  |
| Black oak                   | 287,073        | --       | 43,105    | 84,661    | 59,386    | 27,937    | 48,970    | 23,013    | --      |
| All species                 | 562,959        | --       | 84,577    | 108,078   | 96,850    | 91,283    | 79,646    | 73,659    | 28,867  |
| Hickory                     |                |          |           |           |           |           |           |           |         |
| Bitternut hickory           | 184,657        | --       | 57,266    | 53,799    | 12,669    | 29,471    | 11,941    | 19,511    | --      |
| Pignut hickory              | 5,446          | --       | --        | 5,446     | --        | --        | --        | --        | --      |
| Shagbark hickory            | 415,074        | --       | 130,019   | 113,809   | 73,280    | 35,237    | 32,547    | 30,183    | --      |
| Mockernut hickory           | 5,807          | --       | 2,885     | 2,922     | --        | --        | --        | --        | --      |
| All species                 | 610,985        | --       | 190,171   | 175,976   | 85,949    | 64,708    | 44,488    | 49,694    | --      |
| Hard maple                  |                |          |           |           |           |           |           |           |         |
| Black maple                 | 160,554        | --       | 44,646    | 29,731    | 35,811    | 36,249    | --        | 14,117    | --      |
| Sugar maple                 | 109,498        | --       | 2,917     | 9,124     | 16,844    | 26,910    | 12,894    | 40,809    | --      |
| All species                 | 270,053        | --       | 47,563    | 38,855    | 52,655    | 63,159    | 12,894    | 54,926    | --      |
| Soft maple                  |                |          |           |           |           |           |           |           |         |
| Silver maple                | 1,208,038      | --       | 134,143   | 199,156   | 181,269   | 153,500   | 184,833   | 206,736   | 148,401 |
| All species                 | 1,208,038      | --       | 134,143   | 199,156   | 181,269   | 153,500   | 184,833   | 206,736   | 148,401 |
| Ash                         |                |          |           |           |           |           |           |           |         |
| White ash                   | 128,966        | --       | 17,349    | 49,069    | 21,736    | --        | 11,211    | --        | 29,600  |
| Black ash                   | 10,385         | --       | --        | 4,895     | --        | 5,490     | --        | --        | --      |
| Green ash                   | 178,194        | --       | 26,941    | 51,295    | 44,921    | 33,752    | 21,285    | --        | --      |
| All species                 | 317,545        | --       | 44,290    | 105,260   | 66,657    | 39,242    | 32,496    | --        | 29,600  |
| Cottonwood and aspen        |                |          |           |           |           |           |           |           |         |
| Eastern cottonwood          | 1,483,155      | --       | 64,769    | 89,641    | 37,393    | 106,125   | 136,605   | 497,888   | 550,734 |
| Bigtooth aspen              | 57,790         | --       | 17,697    | 4,716     | 24,737    | --        | 10,640    | --        | --      |
| Quaking aspen               | 3,389          | --       | 3,389     | --        | --        | --        | --        | --        | --      |
| All species                 | 1,544,335      | --       | 85,855    | 94,358    | 62,129    | 106,125   | 147,245   | 497,888   | 550,734 |
| Basswood                    |                |          |           |           |           |           |           |           |         |
| American basswood           | 604,691        | --       | 68,583    | 80,156    | 89,704    | 142,299   | 30,897    | 193,053   | --      |
| All species                 | 604,691        | --       | 68,583    | 80,156    | 89,704    | 142,299   | 30,897    | 193,053   | --      |
| Black walnut                |                |          |           |           |           |           |           |           |         |
| Black walnut                | 864,925        | --       | 140,702   | 188,666   | 226,140   | 155,901   | 53,324    | 100,192   | --      |
| All species                 | 864,925        | --       | 140,702   | 188,666   | 226,140   | 155,901   | 53,324    | 100,192   | --      |

Table 8 continued on next page)

(Table 8 continued on next page)

(Table 8 continued)

| Species group/<br>species    | All<br>classes | Diameter class (inches at breast height) |           |           |           |           |           |           |         | 29.0+ |
|------------------------------|----------------|------------------------------------------|-----------|-----------|-----------|-----------|-----------|-----------|---------|-------|
|                              |                | 9.0-10.9                                 | 11.0-12.9 | 13.0-14.9 | 15.0-16.9 | 17.0-18.9 | 19.0-20.9 | 21.0-28.9 |         |       |
| Hardwoods                    |                |                                          |           |           |           |           |           |           |         |       |
| Other eastern soft hardwoods |                |                                          |           |           |           |           |           |           |         |       |
| Boxelder                     | 59,933         | --                                       | 19,979    | 11,624    | 16,392    | --        | --        | 11,958    | --      |       |
| Ohio buckeye                 | 24,927         | --                                       | 5,381     | 4,404     | 5,398     | --        | 9,744     | --        | --      |       |
| River birch                  | 9,235          | --                                       | --        | 3,650     | 5,584     | --        | --        | --        | --      |       |
| Paper birch                  | 9,949          | --                                       | --        | 4,247     | 5,701     | --        | --        | --        | --      |       |
| Hackberry                    | 472,033        | --                                       | 72,069    | 113,802   | 96,295    | 70,426    | 54,205    | 65,236    | --      |       |
| American sycamore            | 12,443         | --                                       | --        | --        | --        | --        | --        | 12,443    | --      |       |
| Black cherry                 | 122,759        | --                                       | 29,487    | 27,735    | 25,333    | 21,724    | 18,479    | --        | --      |       |
| Black willow                 | 98,070         | --                                       | 2,360     | 8,561     | 6,692     | 22,774    | 10,225    | 47,468    | --      |       |
| American elm                 | 380,399        | --                                       | 85,990    | 94,418    | 84,897    | 52,158    | 23,365    | 39,552    | --      |       |
| Siberian elm                 | 9,036          | --                                       | --        | 3,658     | 5,378     | --        | --        | --        | --      |       |
| Slippery elm                 | 224,198        | --                                       | 55,594    | 71,816    | 27,449    | 28,965    | 9,384     | 30,991    | --      |       |
| All species                  | 1,423,000      | --                                       | 270,859   | 343,915   | 279,110   | 196,047   | 125,421   | 207,647   | --      |       |
| Other eastern hard hardwoods |                |                                          |           |           |           |           |           |           |         |       |
| Honeylocust                  | 243,781        | --                                       | 39,894    | 64,071    | 57,692    | 50,839    | 31,284    | --        | --      |       |
| Red mulberry                 | 4,929          | --                                       | 4,929     | --        | --        | --        | --        | --        | --      |       |
| Black locust                 | 105,740        | --                                       | 14,073    | 27,963    | 29,798    | 11,187    | 10,654    | 12,065    | --      |       |
| All species                  | 354,451        | --                                       | 58,897    | 92,034    | 87,491    | 62,027    | 41,938    | 12,065    | --      |       |
| Total hardwoods              | 11,285,476     | --                                       | 1,562,762 | 1,923,922 | 1,657,864 | 1,599,349 | 1,231,921 | 2,322,532 | 987,127 |       |
| All species groups           | 11,380,405     | 37,142                                   | 1,587,598 | 1,952,189 | 1,682,547 | 1,599,349 | 1,231,921 | 2,322,532 | 987,127 |       |

All table cells without observations in the inventory sample are indicated by --. Table value of 0 indicates the volume rounds to less than 1 thousand board feet. Columns and rows may not add to their totals due to rounding.

<sup>1</sup> International 1/4-inch rule.

Table 9. -- All live aboveground tree biomass on timberland by owner category, softwood/hardwood species category, and tree biomass component, Iowa, 2001-2005

(In thousand dry tons)

| Owner category and<br>softwood/hardwood<br>category | All<br>components | All live<br>1-5 inch trees | Tree biomass component |        |                            |                         |        |                            |
|-----------------------------------------------------|-------------------|----------------------------|------------------------|--------|----------------------------|-------------------------|--------|----------------------------|
|                                                     |                   |                            | Growing-stock trees    |        |                            | Non-growing-stock trees |        |                            |
|                                                     |                   |                            | Total                  | Boles  | Stumps, tops,<br>and limbs | Total                   | Boles  | Stumps, tops,<br>and limbs |
| Public                                              |                   |                            |                        |        |                            |                         |        |                            |
| Softwoods                                           | 210               | 62                         | 85                     | 62     | 23                         | 64                      | 46     | 18                         |
| Hardwoods                                           | 14,636            | 842                        | 11,172                 | 8,301  | 2,871                      | 2,622                   | 1,910  | 711                        |
| Total                                               | 14,846            | 904                        | 11,257                 | 8,363  | 2,893                      | 2,686                   | 1,956  | 730                        |
| Private                                             |                   |                            |                        |        |                            |                         |        |                            |
| Softwoods                                           | 916               | 105                        | 515                    | 374    | 141                        | 296                     | 210    | 86                         |
| Hardwoods                                           | 96,836            | 5,194                      | 65,873                 | 48,286 | 17,588                     | 25,768                  | 18,934 | 6,834                      |
| Total                                               | 97,752            | 5,300                      | 66,388                 | 48,660 | 17,729                     | 26,064                  | 19,144 | 6,920                      |
| All ownerships                                      |                   |                            |                        |        |                            |                         |        |                            |
| Softwoods                                           | 1,126             | 167                        | 600                    | 436    | 164                        | 360                     | 256    | 104                        |
| Hardwoods                                           | 111,472           | 6,036                      | 77,046                 | 56,587 | 20,458                     | 28,390                  | 20,844 | 7,546                      |
| Total                                               | 112,598           | 6,203                      | 77,645                 | 57,023 | 20,622                     | 28,750                  | 21,100 | 7,650                      |

All table cells without observations in the inventory sample are indicated by --. Table value of 0 indicates the aboveground tree biomass rounds to less than 1 thousand dry tons. Columns and rows may not add to their totals due to rounding.

Table 10. -- Average annual net growth of growing stock on timberland by species group and owner category, Iowa, 1999-2000 to 2004-2005

(In thousand cubic feet per year)

| Species group                | Owner category |        |                    |
|------------------------------|----------------|--------|--------------------|
|                              | All owners     | Public | Unidentified owner |
| <b>Softwoods</b>             |                |        |                    |
| Other yellow pines           | 68             | --     | 68                 |
| Other eastern softwoods      | 1,385          | 35     | 1,350              |
| Total softwoods              | 1,452          | 35     | 1,418              |
| <b>Hardwoods</b>             |                |        |                    |
| Select white oaks            | 24,747         | 2,389  | 22,358             |
| Select red oaks              | 7,924          | 1,198  | 6,726              |
| Other red oaks               | 6,043          | 1,248  | 4,796              |
| Hickory                      | 6,245          | 1,050  | 5,194              |
| Hard maple                   | 2,965          | 163    | 2,802              |
| Soft maple                   | 15,877         | 7,568  | 8,309              |
| Ash                          | 6,696          | 1,415  | 5,281              |
| Cottonwood and aspen         | 12,204         | 1,778  | 10,426             |
| Basswood                     | 6,385          | 2,925  | 3,460              |
| Black walnut                 | 15,694         | 730    | 14,965             |
| Other eastern soft hardwoods | 23,331         | 5,855  | 17,476             |
| Other eastern hard hardwoods | 11,686         | 3,356  | 8,330              |
| Total hardwoods              | 139,797        | 29,674 | 110,123            |
| <b>All species groups</b>    | 141,249        | 29,709 | 111,540            |

All table cells without observations in the inventory sample are indicated by --. Table value of 0 indicates the volume rounds to less than 1 thousand cubic feet. Columns and rows may not add to their totals due to rounding.

Table 11. -- Average annual removals of growing stock on timberland by species group and owner category, Iowa, 1999-2000 to 2004-2005

(In thousand cubic feet per year)

| Species group                | Owner category |        |         |                    |
|------------------------------|----------------|--------|---------|--------------------|
|                              | All owners     | Public | Private | Unidentified owner |
| <b>Softwoods</b>             |                |        |         |                    |
| Other eastern softwoods      | 86             | --     | 86      | --                 |
| Total softwoods              | 86             | --     | 86      | --                 |
| <b>Hardwoods</b>             |                |        |         |                    |
| Select white oaks            | 3,201          | --     | 3,201   | --                 |
| Select red oaks              | 3,325          | --     | 3,325   | --                 |
| Other red oaks               | 2,388          | --     | 2,388   | --                 |
| Hickory                      | 336            | --     | 336     | --                 |
| Ash                          | 447            | --     | 447     | --                 |
| Basswood                     | 1,406          | --     | 1,406   | --                 |
| Black walnut                 | 4,207          | --     | 4,207   | --                 |
| Other eastern soft hardwoods | 2,468          | --     | 2,468   | --                 |
| Other eastern hard hardwoods | 417            | --     | 417     | --                 |
| Total hardwoods              | 18,195         | --     | 18,195  | --                 |
| <b>All species groups</b>    | 18,281         | --     | 18,281  | --                 |

All table cells without observations in the inventory sample are indicated by --. Table value of 0 indicates the volume rounds to less than 1 thousand cubic feet. Columns and rows may not add to their totals due to rounding.

Table 12. -- Average annual mortality of growing stock on timberland by species group and owner category, Iowa, 1999-2000 to 2004-2005

(In thousand cubic feet per year)

| Species group                | Owner category |        |         |                    |
|------------------------------|----------------|--------|---------|--------------------|
|                              | All owners     | Public | Private | Unidentified owner |
| <b>Softwoods</b>             |                |        |         |                    |
| Other eastern softwoods      | 33             | --     | 33      | --                 |
| Total softwoods              | 33             | --     | 33      | --                 |
| <b>Hardwoods</b>             |                |        |         |                    |
| Select white oaks            | 3,104          | 946    | 2,158   | --                 |
| Select red oaks              | 941            | --     | 941     | --                 |
| Other red oaks               | 604            | --     | 604     | --                 |
| Hickory                      | 1,194          | --     | 1,194   | --                 |
| Soft maple                   | 42             | --     | 42      | --                 |
| Ash                          | 505            | 147    | 358     | --                 |
| Cottonwood and aspen         | 2,352          | --     | 2,352   | --                 |
| Basswood                     | 2,071          | --     | 2,071   | --                 |
| Black walnut                 | 293            | --     | 293     | --                 |
| Other eastern soft hardwoods | 11,946         | 168    | 11,777  | --                 |
| Other eastern hard hardwoods | 94             | --     | 94      | --                 |
| Total hardwoods              | 23,146         | 1,261  | 21,885  | --                 |
| <b>All species groups</b>    | 23,179         | 1,261  | 21,917  | --                 |

All table cells without observations in the inventory sample are indicated by --. Table value of 0 indicates the volume rounds to less than 1 thousand cubic feet. Columns and rows may not add to their totals due to rounding.

Crocker, Susan J.; Brand, Gary J.; Flickinger, Aron.

2007. **Iowa's forest resources, 2005.** Resour. Bull. NRS-14. Newtown Square, PA: U.S.

Department of Agriculture, Forest Service, Northern Research Station. 31 p

Report presents Iowa's annual inventory results for 2005. Estimates show that Iowa has more than 2.8 million acres of forest land. Total live-tree volume on forest land is 4.0 billion cubic feet (ft<sup>3</sup>). Ninety-eight percent of forest land is classified as timberland.

Oak/hickory is the predominant forest-type group, representing 54 percent of timberland area. Growing-stock volume has risen to an estimated 3.1 billion ft<sup>3</sup>. Live-tree aboveground biomass is 112 million dry tons. In general, Iowa's forests are healthy. However, diseases, including white oak decline and Dutch elm disease, and the threat of invasion from exotic species such as emerald ash borer and nonnative plants are a concern when assessing the health of Iowa's forests.

**KEY WORDS:** Annual inventory, forest land, timberland, forest type, volume, biomass, growth, removals, mortality, forest health, Iowa

---

---

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, SW, Washington, DC 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

---

---



*Capitalizing on the strengths of existing science capacity in the Northeast and Midwest to attain a more integrated cohesive landscape scale research program*