



United States  
Department of  
Agriculture

Forest Service


Pacific Northwest  
Research Station

Research Paper  
PNW-RP-465  
May 1993



# Response of the Engraver Beetle, *Ips Perturbatus*, to Semiochemicals in White Spruce Stands of Interior Alaska

Richard A. Werner





This publication reports research involving pesticides. It does not contain recommendations for their use, nor does it imply that the uses discussed here have been registered. All uses of pesticides must be registered by appropriate State and Federal agencies before they can be recommended.

**CAUTION:** Pesticides can be injurious to humans, domestic animals, desirable plants, and fish or other wildlife—if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended practices for the disposal of surplus pesticides and pesticide containers.

## **Author**

**RICHARD A. WERNER** is supervisory research entomologist, Pacific Northwest Research Station, Institute of Northern Forestry, 308 Tanana Drive, Fairbanks, Alaska 99775-5500.

## Abstract

**Werner, Richard A. 1993.** Response of the engraver beetle, *Ips perturbatus*, to semiochemicals in white spruce stands of interior Alaska. Res. Pap. PNW-RP-465. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 9 p.

Field tests on the efficacy of various scolytid bark beetle pheromones to attract *Ips perturbatus* (Eichhoff) were conducted from 1977 through 1992 in stands of white spruce (*Picea glauca* (Moench) Voss) in interior Alaska. Several pheromones attracted high numbers of *I. perturbatus* and species of the predator *Thanasimus* to baited funnel traps. Test results also indicated that attacks by *I. perturbatus* may be deterred by certain semiochemicals.

**Keywords:** Bark beetles, *Ips perturbatus*, semiochemicals, pheromones, aggregation pheromones, antiaggregation pheromones, insect management, white spruce, *Picea glauca*, Alaska (interior).

## Summary

Data from 7 years of field tests on the efficacy of scolytid bark beetle pheromones to attract *I. perturbatus* are summarized. High numbers of beetles were caught in traps baited with racemic ipsdienol (+50:-50). The addition of *exo*-brevicomin to racemic ipsdienol increased the number of beetles caught by 97 percent in 1992 field tests. Racemic ipsenol consistently caught few beetles in baited traps. In 1987 when beetle populations were high, field tests using beetle deterrents showed that the addition of 2-methyl-3-buten-2-ol and racemic ipsenol to racemic ipsdienol in funnel traps reduced the number of beetles caught by 92 and 38 percent, respectively. The addition of either 2-methyl-3-buten-2-ol or racemic ipsenol to racemic ipsdienol reduced beetle catch by 88 percent in 1992 field tests when beetle populations were low. In 1991 field tests, species of the predator *Thanasimus* were caught by the enantiomers of ipsdienol tested alone or in combinations; however, the combination of racemic ipsdienol + (+) ipsdienol (+97:-3) + (-) ipsdienol (-3:+97) caught the most *Thanasimus*. Future research will concentrate on field tests to deter attacks by *I. perturbatus* with semiochemicals.

## Contents

|   |                                               |
|---|-----------------------------------------------|
| 1 | <b>Introduction</b>                           |
| 2 | <b>Materials and Methods</b>                  |
| 2 | Study Area Characteristics and Trap Placement |
| 3 | Experimental Design                           |
| 3 | Statistical Analyses                          |
| 3 | <b>Results and Discussion</b>                 |
| 4 | <b>Conclusions</b>                            |
| 5 | <b>Metric and English Units of Measure</b>    |
| 5 | <b>References</b>                             |
| 8 | <b>Appendix</b>                               |

## Introduction

The engraver beetle (*Ips perturbatus* (Eichhoff)) (Coleoptera: Scolytidae) usually is not a primary killer of spruce trees in Alaska but prefers to attack and colonize white spruce (*Picea glauca* (Moench) Voss) stressed by such disturbance as fire, periodic flooding, drought, logging, ice and snow damage, or wind. Wind-thrown and ice- or snow-damaged trees and logging slash are ideal breeding habitats (Holsten and Werner 1987). In most years, endemic populations infest individual standing trees, but during warm, dry summers after mild winters with above-average snowfall, beetle populations increase to such high levels that live, standing trees are infested and killed (Werner 1988). The average number of hectares infested annually in interior Alaska from 1978 to 1985 was 5060; however, high beetle populations killed trees over 6475 hectares in the Fairbanks area alone in 1986 and 1987 after a wildfire in 1983 and excessive ice and snow damage in 1985 (Holsten 1986). A record snowfall in the Fairbanks area in 1991 caused breakage of tree boles and tops in stands of white spruce. Observations of these trees during summer 1991 noted high levels of infestation by *I. perturbatus*. These populations reached epidemic levels during summer 1992 because winter temperatures and snowfall were above average. These recurring problems with epidemic populations of *I. perturbatus* prompted renewed research to develop new management strategies in addition to those described by Werner (1988).

The production of aggregation pheromones by pioneer beetles has been described for many scolytid species (Borden 1982). Species of *Ips* beetles aggregate on selected hosts in response to pheromones produced by males in the initial construction of egg galleries. Identification of pheromones from species of *Ips* started in 1966 when three terpene alcohols were identified as components of the pheromone for *I. paraconfusus* Lanier (then *I. confusus* LeConte) (Silverstein and others 1966a, 1966b; Wood and others 1966). These compounds were (1) 2-methyl-6-methylene-7-octen-4-ol (ipsenol), (2) *cis*-verbenol, and (3) 2-methyl-6-methylene-2,7-octadien-4-ol (ipsdienol). Mori (1975, 1976) describes the absolute configuration of these compounds as (–) ipsenol and (+) ipsdienol. Neither compound was active alone, but a combination of ipsenol with either of the other two was active. These compounds and *trans*-verbenol are present in various combinations in several species of *Ips* (Renwick and Vite 1972, Vite and others 1972). Ipsdienol, ipsenol, and lanierone are the only biologically active, insect-produced compounds known from *Ips* (Bakke 1970, Teale and others). Monoterpenes of the host tree serve as precursors of the pheromones (Bakke 1975; Hughes 1973, 1974; Vite and others 1972; Vite and others 1974). These same compounds were implicated as pheromones of *I. confusus* (Young and others 1973). Vite and Renwick (1971) identified ipsenol as an aggregation pheromone for *I. grandicollis* (Eichhoff), and *I. calligraphus* (Germar) responds to a combination of ipsdienol and *cis*-verbenol along with host volatiles (Renwick and Vite 1972). Bakke and others (1977) identified ipsdienol, *cis*-verbenol, and 2-methyl-3-butene-2-ol (methyl butenol) as the aggregation pheromones of *I. typographus* (L.). *Ips pini* (Say) responds to the two enantiomers of ipsdienol (Birch and others 1980, Lanier and others 1980, Stewart 1975); whereas, *I. latidens* (LeConte) and *I. paraconfusus* Lanier respond to ipsenol (Lanier and Wood 1975).

**Materials and  
Methods  
Study Area  
Characteristics and  
Trap Placement**

The use of semiochemicals, bioactive message-bearing chemicals, to manipulate bark beetle populations has gained considerable interest during the last decade because of minimum environmental impact. This use includes synthetic host attractants and pheromones to either prevent attack or reduce the attack of bark beetles to a level below the threshold density required for the development of brood trees. This type of strategy has several advantages. First, the use of traditional insecticides could be minimized. Secondly, development of beetle resistance to treatment should be negligible because pheromones are part of the communication system of bark beetles. Lastly, there would be little direct mortality to parasites and predators as occurs with insecticide treatment.

Several objectives were developed and tested from 1977 through 1992:

1. To identify which scolytid bark beetle pheromones were attractive to *I. perturbatus* in interior Alaska.
2. To determine the efficacy of various enantiomers of ipsdienol and ipsenol to attract *I. perturbatus* to baited traps.
3. To determine the efficacy of several semiochemicals to deter attacks on white spruce stands, slash, and log decks by *I. perturbatus*.

Mature white spruce stands 300 m in elevation in the Bonanza Creek Experimental Forest, 40 km west of Fairbanks, Alaska, were used for the studies. Trees were 165 years old with an average diameter of 30 cm, an average height of 36 m, and a stand density of 875 trees per ha. Shrub cover was sparse with only *Alnus crispa* (Ait.) Pursh and *Viburnum edule* (Michaux) Raf. occupying the sites. The climate of the area is strongly continental with an average temperature of -3.5 °C (range -50 to +35 °C). The average temperature during beetle flight is 16 °C. A different site was selected each year for each study to reduce the bias of unequal population size.

Semiochemicals were released from perforated polyethylene vials attached to cylindrical wire mesh sticky traps (Kline and others 1974) in the field tests from 1977 through 1980. Traps measured 45 cm in length and 25 cm in diameter and were constructed of 6.35-mm mesh wire screen. Traps were set on iron posts 1.5 m aboveground. The studies from 1985 through 1992 used slow-release formulations of semiochemicals dispersed from 12-unit Lindgren<sup>1</sup> multiple funnel traps (Lindgren 1983). Funnel traps were suspended from a nylon rope tied between two nonhost trees or white spruce <7.6 cm diameter at breast height so the collection container of the trap was 0.3 m aboveground. All traps were placed in a grid pattern at 30-m intervals. Beetles were collected from the traps weekly from mid-May to early July. The insects caught were placed in glass vials with 70-percent ethanol and taken to the laboratory for identification and total beetle counts. The various semiochemicals and their respective release rates used in the different studies reported in this paper are shown in tables 1 through 4. Semiochemicals were purchased commercially and releasers prepared in the laboratory for the studies from 1977 through 1980; studies thereafter used slow-release formulations of semiochemicals. All semiochemicals used in the field tests had a chemical purity above 98.6 percent.

---

<sup>1</sup> The use of trade or firm names in this publication is for reader information and does not imply endorsement by the U.S. Department of Agriculture of any product or service.

|                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|-------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Experimental Design</b>    | Treatments for all field tests were replicated five times in a randomized complete block design. Traps were rotated randomly each week to reduce the effects of host terpene attraction from adjacent live white spruce trees and spillover of beetles into these trees. Treatments consisted of semiochemicals in traps with unbaited traps as controls.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Statistical Analyses</b>   | Data were tested for homogeneity and, if nonhomogenous, were transformed by using a parametric test to overcome variations due to zero counts. Total numbers of <i>Ips</i> and <i>Thanasimus</i> beetles caught for each of the treatments were analyzed by using ANOVA. Differences between means were tested by using Tukey's (1953) studentized range (HSD) test ( $P = 0.05$ ).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Results and Discussion</b> | <p>Field tests from 1977 through 1980 compared beetle response to racemic ipsdienol (+50:-50) and racemic ipsenol (+50:-50) with response to pheromones from other scolytids (table 1). Sticky, cylindrical, wire traps baited with racemic ipsdienol caught significantly more <i>I. perturbatus</i> than did other treatments. Racemic ipsdienol caught 58 to 64 percent more beetles than other scolytid pheromones did during years when beetle populations were low (table 1). Few beetles were caught in traps baited with racemic ipsenol, and no beetles were caught with <i>cis</i>-verbenol or <i>trans</i>-verbenol.</p> <p>Further field tests were conducted in 1987 with commercial baits in slow-release formulations. These tests compared the efficacy of racemic ipsdienol (0.4 mg per day), racemic ipsenol (0.4 mg per day), and 2-methyl-3-buten-2-ol (17-20 mg per day). In these tests, Lindgren funnel traps baited with racemic ipsdienol caught 84 percent more beetles than racemic ipsenol alone and 62 percent more beetles than the combination of racemic ipsdienol and racemic ipsenol (table 2). Racemic ipsenol repeatedly caught few beetles in replicated field tests from 1977 to 1992. The addition of racemic ipsenol and 2-methyl-3-buten-2-ol to racemic ipsdienol reduced beetle catch by 38 and 92 percent, respectively.</p> <p>As a result of the 1987 field tests, racemic ipsdienol released at a rate of 0.4 mg per day was recommended as a bait in funnel traps and on trap trees to trap out populations of <i>I. perturbatus</i> before and after logging operations in interior Alaska (Werner 1988). The recommendation stated that traps should be spaced at 30-m intervals when used in barrier trapping around clearcuts and selective cuts and around log decks, and at 60-m intervals when used in preharvest and postharvest trapping. Baited trap trees should be spaced at 60-m intervals in stands before timber harvest and trap decks at 30-m intervals along rights-of-way clearings.</p> <p>Racemic ipsdienol released at a rate of 0.4 mg per day was used to bait trap trees and log decks after logging operations in interior Alaska from 1987 to 1991; however, the number of beetles trapped was low compared to the number of beetles caught in Lindgren funnel traps baited with racemic ipsdienol at a release rate of 0.2 mg per day, which was used for monitoring populations of <i>Ips</i>. The difference in beetle catch could be related to the different release rates or the different spacings between traps and between trap trees or log decks.</p> |

An additional field test was done in 1991 to further compare the efficacy of the enantiomers of ipsdienol; that is, racemic ipsdienol, (+) ipsdienol, and (-) ipsdienol. Racemic ipsdienol caught 98 percent more beetles than (+) ipsdienol and 64 percent more beetles than (-) ipsdienol (table 3). All enantiomers of ipsdienol and combinations except (-) ipsdienol alone attracted *Thanasimus*; however, the combination of racemic ipsdienol + (+) ipsdienol + (-) ipsdienol caught the most beetles (25 percent of 538 *Thanasimus* caught). The 1991 field test also investigated the response of recently emerged adults to the enantiomers of ipsdienol before these adults entered their overwintering sites. *Ips perturbatus* adults were caught from May 12 through June 25 with peak flight the week of May 26. No newly emerged *I. perturbatus* adults were caught during August and September, which indicates no response to the ipsdienol enantiomers at this stage of development. This study confirmed previous results that racemic ipsdienol at a release rate of 0.2 mg per day is more attractive to *I. perturbatus* than the other enantiomers or combinations of enantiomers.

Based on results from the 1987 field tests, a field test was conducted in 1992 to determine if certain semiochemicals deterred the response of *I. perturbatus* to racemic ipsdienol. The addition of various semiochemicals to racemic ipsdienol reduced beetle catch in funnel traps as follows: 2-methyl-3-buten-2-ol and racemic ipsenol by 88 percent, 3-methylcyclohex-2-enone (MCH) by 31 percent, and verbenone by 19 percent; however, the addition of *exo*-brevicomin increased beetle catch by 97 percent (table 4). Birch and Wood (1975) found that *I. pini* and *I. paraconfusus* are not cross-attractive although they attack the same host trees at the same time. Ipsenol from male *I. paraconfusus* inhibits the response of *I. pini* to male *I. pini*; whereas, linalool from male *I. pini* reduces the catch of *I. paraconfusus* in response to male *I. paraconfusus*. Borden and others (1991) found that verbenone + ipsenol reduced the number of logs attacked by *I. pini* by 66.7 percent and a 98.8 percent reduction in attack density. Livingston<sup>2</sup> also found in 1992 field bioassays that the addition of racemic ipsenol to the aggregation pheromones racemic ipsdienol + lanierone reduced the number of *I. pini* caught by 70 percent. The addition of racemic ipsenol + verbenone in combination to racemic ipsdienol + lanierone decreased trap catch by 95 percent.

## Conclusions

Seven years of field testing on the efficacy of various semiochemicals on *I. perturbatus* was completed in 1992. *Ips perturbatus* were attracted to Lindgren funnel traps baited with racemic ipsdienol alone at release rates of 0.2 and 0.4 mg per day and *exo*-brevicomin + racemic ipsdienol at a release rate of 0.2 mg per day. Racemic ipsdienol (0.2 mg per day) currently is recommended for use in Lindgren funnel traps, and trap trees and log decks. The addition of 2-methyl-3-buten-2-ol (3.8 mg per day) and racemic ipsenol (0.2 mg per day) to racemic ipsdienol in funnel traps reduced the number of beetles caught compared to traps baited with racemic ipsdienol alone.

<sup>2</sup> Personal communication, December 8, 1992, R. Ladd  
Livingston, Idaho Department of Lands, Coeur d'Alene, ID  
83814-0670.



Parent and new brood adults did not respond to baited pheromone traps upon emergence from host trees in August but did respond the following spring when beetles emerged from overwintering sites. Species of the predator *Thanasimus* were caught by the enantiomers of ipsdienol tested alone or in combinations; however, the combination of racemic ipsdienol + (+) ipsdienol + (-) ipsdienol caught the most *Thanasimus*. Future research will concentrate on field tests to deter attacks by *I. perturbatus*.

## Metric and English Units of Measure

| When you know:   | Multiply by:   | To find:   |
|------------------|----------------|------------|
| Celsius (°C)     | 1.8 and add 32 | Fahrenheit |
| Centimeters (cm) | 2.54           | Inches     |
| Hectares (ha)    | 2.47           | Acres      |
| Kilometers (km)  | 0.621          | Miles      |
| Meters (m)       | 3.281          | Feet       |
| Millimeters (mm) | 0.254          | Inches     |

## References

- Bakke, A. 1970.** Evidence of a population aggregating pheromone in *Ips typographus* (Coleoptera: Scolytidae). Contributions Boyce Thompson Institute for Plant Research, Professional Papers. 24: 309-310.
- Bakke, Alf. 1975.** Aggregation pheromone in the bark beetle *Ips duplicatus* (Sahlberg). Norwegian Journal of Entomology. 22: 67-69.
- Bakke, A.; Froyen, P.; Skattebol, L. 1977.** Field response to a new pheromonal compound isolated from *Ips typographus*. Naturwissenschaften. 64: 98.
- Birch, M.C.; Light, D.M.; Wood, D.L. [and others]. 1980.** Pheromone attraction and allomonal interruption of *Ips pini* in California by the two enantiomers of ipsdienol. Journal of Chemical Ecology. 6: 703-717.
- Birch, M.C.; Wood, D.L. 1975.** Mutual inhibition of the attractant pheromone response by two species of *Ips* (Coleoptera: Scolytidae). Journal of Chemical Ecology. 1: 101-113.
- Borden, J.H. 1982.** Aggregation pheromones. In: Mitton, J.B.; Sturgeon, K.B., eds. Bark beetles in North American conifers. Austin, TX: University of Texas Press: 74-139.
- Borden, J.; Devlin, D.R.; Miller, D.R. 1991.** Synomones of two sympatric species deter attack by the pine engraver, *Ips pini* (Coleoptera: Scolytidae). Canadian Journal of Forest Research. 22: 381-387.
- Holsten, E.H. 1986.** Preliminary evaluation of an *Ips perturbatus* outbreak in the Bonanza Creek Experimental Forest, Fairbanks, Alaska. Biological Evaluation R10-86-1. Anchorage, AK: U.S. Department of Agriculture, Forest Service, Alaska Region. 8 p.
- Holsten, Edward H.; Werner, Richard A. [1987].** Engraver beetles in Alaska forests [Leaflet]. [Portland, OR]: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. [Not paged].
- Hughes, P.R. 1973.** *Dendroctonus*: Production of pheromones and related compounds in response to host monoterpenes. Zeitschrift für Angewandte Entomologie. 95: 294-312.

- Hughes, P.R. 1974.** Myrcene: a precursor of pheromones in *Ips* beetles. *Journal of Insect Physiology*. 20: 1271-1275.
- Kilne, L.N.; Schmitz, R.F.; Rudinsky, J.A. [and others]. 1974.** Repression of spruce beetle (Coleoptera: Scolytidae) attraction by methylcyclohexenone in Idaho. *Canadian Entomologist*. 106: 485-491.
- Lanier, Gerald N.; Classon, Alf.; Stewart, Thomas [and others]. 1980.** *Ips pini*: the basis for interpopulational differences in pheromone biology. *Journal of Chemical Ecology*. 6: 677-687.
- Lanier, G.N.; Wood, D.L. 1975.** Specificity of response to pheromones in the genus *Ips* (Coleoptera: Scolytidae). *Journal of Chemical Ecology*. 1: 9-23.
- Lindgren, B.S. 1983.** A multiple funnel trap for scolytid beetles (Coleoptera) *Canadian Entomologist*. 115: 299-302.
- Mori, K. 1975.** Synthesis and absolute configuration of (-)-ipsenol (2-methyl-6-methylene-7-octen-4-ol), the pheromone of *Ips paraconfusus* Lanier. *Tetrahedron*: 21: 87-2190.
- Mori, K. 1976.** Absolute configuration of (+)-ipsdienol, the pheromone of *Ips paraconfusus* Lanier, as determined by the synthesis of its (R)-(-) isomer. *Tetrahedron Letters*: 1609-1612.
- Renwick, J.A.A.; Vite, J.P. 1972.** Pheromones and host volatiles that govern aggregation of the six-spined engraver beetle, *Ips calligraphus*. *Journal of Insect Physiology*. 18: 1215-1219.
- Silverstein, Robert M.; Rodin, J. Otto.; Wood, David L. 1966a.** Sex attractants in frass produced by male *Ips confusus* in ponderosa pine. *Science*. 154: 509-510.
- Silverstein, R.M.; Rodin, J.O.; Wood, D.L. [and others]. 1966b.** Identification of two new terpene alcohols from frass produced by *Ips confusus* in ponderosa pine. *Tetrahedron*. 22: 1929-1936.
- Stewart, T.E. 1975.** Volatiles isolated from *Ips pini*: isolation, identification enantiomer composition, biological activity, and enantiometric composition of other insect pheromone alcohols and bicyclic ketols. Syracuse, NY: State University of New York. 135 p. M.S. thesis.
- Tukey, J.W. 1953.** The problem of multiple comparisons. Princeton, NJ: Princeton University. 396 p. Unpublished manuscript. On file with: Princeton University, Firestone Library, Princeton, NJ 08544.
- Vite, J.P.; Bakke, A.; Renwick, J.A.A. 1972.** Pheromones in *Ips* (Coleoptera: Scolytidae): occurrence and production. *Canadian Entomologist*. 104: 1967-1975.
- Vite, J.P.; Bakke, A.; Hughes, P.R. 1974.** Ein Populationslockstoff des zwölfzahnigen Kiefernborstenkäfers *Ips sexdentatus*. *Naturwissenschaften*. 8: 365-366.
- Vite, J.P.; Renwick, J.A.A. 1971.** Population aggregating pheromone in the bark beetle, *Ips grandicollis*. *Journal of Insect Physiology*. 17: 1699-1704.

- Werner, Richard A. 1988.** Recommendations for suppression of an *Ips perturbatus* outbreak in interior Alaska using integrated control. In: Integrated control of scolytid bark beetles: Proceedings of the IUFRO working party and 17th international congress of entomology symposium; 1988 July 4; Vancouver, BC. Blacksburg, VA: Virginia Polytechnic Institute and State University Press: 189-196.
- Wood, D.L.; Browne, L.E.; Silverstein, R.M. [and others]. 1966.** Sex pheromones of bark beetles—I. Mass-production, bioassay, source, and isolation of the sex pheromone of *Ips confusus* (LeC.). Journal of Insect Physiology. 12: 523-536.
- Young, J.C.; Silverstein, R.M.; Birch, M.C. 1973.** Aggregation pheromone of the beetle *Ips confusus*: isolation and identification. Journal of Insect Physiology. 19: 2273-2277.

## Appendix

**Table 1—Total number of *Ips perturbatus* caught in cylindrical sticky traps by synthetic scolytid pheromones, Fairbanks, Alaska, 1977-80**

| Treatment                                             | Beetles caught <sup>a</sup> |      |      |      |
|-------------------------------------------------------|-----------------------------|------|------|------|
|                                                       | 1977                        | 1978 | 1979 | 1980 |
|                                                       | ----- Number -----          |      |      |      |
| Frontalin                                             | 3d                          | 105d | 92d  | 397b |
| Frontalin + alpha pinene (1:1)                        | 2d                          | 234d | 187c | 243d |
| Alpha pinene                                          | 0d                          | 66e  | 32e  | 143d |
| Seudenol                                              | 10d                         | 85e  | 101d | 374c |
| Seudenol + alpha pinene                               | 3d                          | 166d | 218c | 423b |
| Cis-verbenol                                          | 0d                          | 0f   | 0e   | —    |
| Trans-verbenol                                        | 0d                          | 0f   | 0e   | —    |
| Racemic ipsenol                                       | 125c                        | 144d | 236c | 429b |
| Racemic ipsenol + alpha pinene                        | 101c                        | 384c | 54d  | 207d |
| Racemic ipsdienol                                     | 423a                        | 995a | 564a | 721a |
| Racemic ipsdienol + alpha pinene                      | 112c                        | 618b | 144d | 311c |
| Racemic ipsenol + racemic ipsdienol                   | 231b                        | 704b | 377b | 660a |
| Racemic ipsenol + racemic ipsdienol<br>+ alpha pinene | 183b                        | 466c | 266c | 312c |
| Sulcatol + alpha pinene                               | 99c                         | 156d | 88d  | 49e  |
| Unbaited control                                      | 0d                          | 25f  | 15e  | 49e  |

— = no data were collected.

<sup>a</sup> Values followed by the same letter within columns are not significantly different ( $P < 0.05$ , Tukey's [1953] studentized range test [HSD]).

**Table 2—Effect of 2-methyl-3-buten-2-ol and racemic ipsenol on the response of *Ips perturbatus* to racemic ipsdienol dispersed from Lindgren funnel traps, Fairbanks, Alaska, 1987**

| Treatment                                                      | Release rate | Beetles caught <sup>a</sup> |
|----------------------------------------------------------------|--------------|-----------------------------|
|                                                                | Mg per day   | Mean number $\pm$ SD        |
| Racemic ipsdienol                                              | 0.4          | 2,286 $\pm$ 276a            |
| Racemic ipsenol                                                | .4           | 373 $\pm$ 46c               |
| 2-methyl-3-buten-2-ol                                          | 17-20        | 86 $\pm$ 11d                |
| Racemic ipsdienol + racemic ipsenol                            | .4           | 1,420 $\pm$ 116b            |
| Racemic ipsdienol + 2-methyl-3-buten-2-ol                      | .4           | 185 $\pm$ 23c               |
| Racemic ipsenol + 2-methyl-3-buten-2-ol                        | .4           | 248 $\pm$ 35c               |
| Racemic ipsdienol + racemic ipsenol<br>+ 2-methyl-3-buten-2-ol | .4           | 1,517 $\pm$ 154b            |
| Unbaited control                                               | —            | 384 $\pm$ 61c               |

<sup>a</sup> Values followed by the same letter are not significantly different ( $P < 0.05$ , Tukey's [1953] studentized range test [HSD]).

**Table 3—Number of *Ips perturbatus* and *Thanasimus* spp. caught in Lindgren funnel traps by combinations of enantiomers of ipsdienol, Fairbanks, Alaska, 1991**

| Treatment                                            | Beetles caught <sup>a</sup> |                   |
|------------------------------------------------------|-----------------------------|-------------------|
|                                                      | <i>I. perturbatus</i>       | <i>Thanasimus</i> |
| --- Mean number ± SD ---                             |                             |                   |
| Racemic ipsdienol                                    | 7.30 ± 35.83a               | 3.12 ± 2.9b       |
| (+) ipsdienol (97:3)                                 | .13 ± 0.47d                 | 2.64 ± 1.2b       |
| (-) ipsdienol (3:97)                                 | 2.62 ± 11.25c               | .56 ± 0.32c       |
| Racemic ipsdienol + (+) ipsdienol                    | .75 ± 2.17d                 | 2.20 ± 2.6b       |
| Racemic ipsdienol + (-) ipsdienol                    | 4.30 ± 11.31b               | 2.36 ± 3.2b       |
| (+) ipsdienol + (-) ipsdienol                        | 1.98 ± 4.10c                | 2.56 ± 1.0b       |
| Racemic ipsdienol + (+) ipsdienol<br>+ (-) ipsdienol | .75 ± 1.35d                 | 5.52 ± 4.1a       |
| Unbaited control                                     | .73 ± 2.76d                 | .64 ± 0.29c       |

<sup>a</sup> Release rates were 0.2 mg per day. Values followed by the same letter are not significantly different ( $P < 0.05$ , Tukey's [1953] studentized range test [HSD]).

**Table 4—Effect of various semiochemicals on the response of *Ips perturbatus* to racemic ipsdienol dispersed from Lindgren funnel traps, Fairbanks, Alaska, 1992**

| Treatment                                 | Release rate | Beetles caught <sup>a</sup> |
|-------------------------------------------|--------------|-----------------------------|
|                                           | Mg/day       | Mean number ± SD            |
| (-) verbenone (+13:-87)                   | 5.0          | 0.00 ± 0.00a                |
| MCH                                       | 5.0          | .00 ± 0.00a                 |
| 2-methyl-3-buten-2-ol                     | 3.8          | .25 ± 0.05a                 |
| Racemic ipsenol                           | .2           | .25 ± 0.05a                 |
| Exo-brevicomin                            | .2           | 2.00 ± 0.50ab               |
| Racemic ipsdienol                         | .2           | 6.50 ± 1.65ab               |
| Racemic ipsdienol + 2-methyl-3-buten-2-ol | —            | .75 ± 0.15a                 |
| Racemic ipsdienol + racemic ipsenol       | .4           | .75 ± 0.15a                 |
| Racemic ipsdienol + MCH                   | —            | 4.50 ± 1.00ab               |
| Racemic ipsdienol + verbenone             | —            | 5.25 ± 1.25ab               |
| Racemic ipsdienol + exo-brevicomin        | —            | 12.75 ± 2.55b               |
| Unbaited control                          | —            | .00 ± 0.00a                 |

<sup>a</sup> Values followed by the same letter are not significantly different ( $P < 0.05$ , Tukey's [1953] studentized range test [HSD]).

**Werner, Richard A. 1993.** Response of the engraver beetle, *Ips perturbatus*, to semiochemicals in white spruce stands of interior Alaska. Res. Pap. PNW-RP-465. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 9 p.

Field tests on the efficacy of various scolytid bark beetle pheromones to attract *Ips perturbatus* (Eichhoff) were conducted from 1977 through 1992 in stands of white spruce (*Picea glauca* (Moench) Voss) in interior Alaska. Several pheromones attracted high numbers of *I. perturbatus* and species of the predator *Thanasimus* to baited funnel traps. Test results also indicated that attacks by *I. perturbatus* may be deterred by certain semiochemicals.

**Keywords:** Bark beetles, *Ips perturbatus*, semiochemicals, pheromones, aggregation pheromones, antiaggregation pheromones, insect management, white spruce, *Picea glauca*, Alaska (interior).

The Forest Service of the U.S. Department of Agriculture is dedicated to the principle of multiple use management of the Nation's forest resources for sustained yields of wood, water, forage, wildlife, and recreation. Through forestry research, cooperation with the States and private forest owners, and management of the National Forests and National Grasslands, it strives—as directed by Congress—to provide increasingly greater service to a growing Nation.

The U.S. Department of Agriculture is an Equal Opportunity Employer. Applicants for all Department programs will be given equal consideration without regard to age, race, color, sex, religion, or national origin.

Pacific Northwest Research Station  
333 S.W. First Avenue  
P.O. Box 3890  
Portland, Oregon 97208-3890

---

U.S. Department of Agriculture  
Pacific Northwest Research Station  
333 S.W. First Avenue  
P.O. Box 3890  
Portland, OR 97208

---

Official Business  
Penalty for Private Use, \$300

**do NOT detach label**