

Sonderdruck aus:

Silvae Genetica

Provenance Variation in *Eucalyptus camaldulensis* Dehnh. in California

By B. M. EMERY and F. T. LEDIG



J. D. Sauerländer's Verlag, Frankfurt a. M.

Silvae Genetica 36 (1987)

Silvae Genetica
is edited from the
**Bundesforschungsanstalt für Forst- und
Holzwirtschaft Hamburg**
in collaboration with

W. T. Adams
Oregon State University
College of Forestry
Corvallis, Oregon 97331-5704
U. S. A.

M. R. Ahuja
Institut für Forstgenetik
und Forstpflanzenzüchtung
Siekerlandstraße 2
D-2070 Großhansdorf 2-Schmalenbeck
Bundesrepublik Deutschland

M. Arbez
Laboratoire d'Amélioration
des Arbres Forestiers de Bordeaux
Recherches Forestières
Domaine de l'Hermitage-Pierroton
33510 Cestas Principal
France

R. D. Barnes
University of Oxford
Dept. of Agricultural
and Forest Sciences
South Parks Road
Oxford OX1 3RB
England

W. H. G. Barrett
Fiplasto Forestal SA
Alsina 756 - Piso 8°
1087 Buenos Aires
Argentina

I. R. Brown
University of Aberdeen
Dept. of Forestry
St. Machar Drive
Old Aberdeen AB9 2UU
Scotland

T. Eguiluz
Centro de Genética Forestal A.C.
División de Ciencias Forestales
Universidad Autónoma Chapingo
Chapingo, México C.P. 56230

K. G. Eldridge
CSIRO
Division of Forest Research
P. O. Box 4008
Canberra A. C. T. 2600
Australia

D. P. Fowler
Maritimes Forest Research Centre
Canadian Forestry Service
Tree Improvement
P. O. Box 4000
Fredericton, N. B. E3B 5P7
Canada

Editorial Office: G. Heinrich Melchior

Address: Siekerlandstraße 2
D-2070 Grosshansdorf 2
(Schmalenbeck),
Federal Republic of Germany

Rules for publishing a manuscript in *Silvae Genetica* can be requested from the publishing house or from the editorial office.

Silvae Genetica
herausgegeben von der
**Bundesforschungsanstalt für Forst- und
Holzwirtschaft Hamburg**
unter Mitwirkung von

M. Giertych
Instytut Dendrologii
Polskiej Akademii Nauk
63-120 Kórnik
ul. Parkowa 5
Poland

M. Hagman
Forest Research Institute
Rillitie 10
SF-01300 Vantaa 30
Finland

H. H. Hattemer
Institut für Forstgenetik
und Forstpflanzenzüchtung,
Forstliche Biometrie und Informatik
der Universität Göttingen
Büsgenweg 2
D-3400 Göttingen-Weende
Bundesrepublik Deutschland

R. C. Kellison
North Carolina State University
School of Forest Resources
Dept. of Forestry
Box 5488
Raleigh, N. C. 27650
U. S. A.

F. T. Ledig
USDA Forest Service
Pacific Southwest Forest
and Range Experiment Station
P. O. Box 245
Berkeley, California 94701
U. S. A.

D. Lindgren
The Swedish University
of Agricultural Sciences
The Faculty of Forestry
Dept. of Forest Genetics
and Plant Physiology
S-901 83 Umeå
Sweden

C. Mátyás
Hungarian Forest Research Institute,
Kámon Arboretum
H-9707 Szombathely
Hungary

J. B. Mitton
University of Colorado
Environmental, Population
and Organismic Biology
122 Ramaley
Campus Box B-334
Boulder, Colorado 80309-0334
U. S. A.

H. J. Muhs
Institut für Forstgenetik
und Forstpflanzenzüchtung
Siekerlandstraße 2
D-2070 Großhansdorf 2-Schmalenbeck
Bundesrepublik Deutschland

Schriftleitung: G. Heinrich Melchior

Adresse: Siekerlandstraße 2
D-2070 Großhansdorf 2
(Schmalenbeck),
Bundesrepublik Deutschland

Richtlinien zur Veröffentlichung eines Manuskriptes in *Silvae Genetica* können von der Redaktion oder vom Verlag angefordert werden.

Silvae Genetica
édité par
**Bundesforschungsanstalt für Forst- und
Holzwirtschaft Hamburg**
avec la coopération de

G. Namkoong
USDA Forest Service
Southeastern Forest Experiment Station
North Carolina State University
Dept. of Genetics
P. O. Box 7614
Raleigh, N. C. 27695
U. S. A.

D. G. Nikles
Division of Technical Services (R & U)
Dept. of Forestry
P. O. Box 631, 80 Meiers Road
Indooroopilly 4068
Queensland
Australia

K. Ohba
University of Tsukuba,
Institute of Agriculture and Forestry,
Sakura-mura,
Ibaraki-ken 305,
Japan

M. Quijada R.
Universidad de los Andes
Facultad de Ciencias Forestales
Instituto de Silvicultura
Apartado No. 305
Mérida
Venezuela

H. Roulund
Royal Veterinary
and Agricultural University
Arboretum
DK-2970 Hørsholm
Denmark

C. J. A. Shelbourne
Forest Research Institute
Private Bag
Rotorua
New Zealand

R. R. Silen
USDA Forest Service
Pacific Northwest Forest
and Range Experiment Station
Forestry Sciences Laboratory
3200 Jefferson Way
Corvallis, Oregon 97331
U. S. A.

R. F. Stettler
University of Washington
Biological Sciences Division
College of Forest Resources
115 Anderson Hall
Seattle, Washington 98195
U. S. A.

Maison d'Édition: G. Heinrich Melchior

Adresse: Siekerlandstraße 2
D-2070 Grosshansdorf 2
(Schmalenbeck)
République fédérale d'Allemagne

Des instructions pour publier un manuscrit dans *Silvae Genetica* peuvent être demandées à la rédaction ou à la maison d'édition.

Provenance Variation in *Eucalyptus camaldulensis* Dehnh. in California*)

By B. M. EMERY and F. T. LEDIG¹⁾

Institute of Forest Genetics,
Pacific Southwest Forest and Range Experiment Station,
U.S.D.A. Forest Service, Box 245,
Berkeley, California 94701, USA

(Received 19th March 1986)

Summary

In California, the Lake Albacutya provenance of river red gum was clearly superior in volume growth to 22 other provenances collected throughout the range of the species in Australia. It had at least 2.5 times the volume of the plantation mean at 5.5 years, consistent with its performance in other countries with Mediterranean climates like California's. Other provenances from southeastern Australia, particularly those in the Murray River basin, also performed well.

The Murray River trees tended to be more crooked and have more forking than those from other areas, and also were among the most heavily flowering in the plantation. The Lake Albacutya trees were somewhat straighter and less forked than those from the Murray River drainage proper.

All successful provenances had a high rainfall pattern correlation (RPC), a statistic defined as the correlation of monthly precipitation at provenance origin with precipitation at the planting site. Cluster analysis on monthly precipitation, mean maximum temperature, and length of frost season linked provenances from the Murray River drainage and Lake Albacutya. Successful provenances had an $RPC > 0$, long frost periods > 120 days, and low summer mean maximum temperatures $< 34^{\circ}\text{C}$. When the planting site was included in the cluster analysis, it also joined the Lake Albacutya and Murray River group. RPC and the multivariate techniques provide a basis for objectively recognizing homoclimes.

Key words: geographic variation, *Eucalyptus camaldulensis*, Lake Albacutya, biomass fuel, stem form.

Zusammenfassung

In einem Provenienzversuch in Kalifornien (USA) mit 23 Herkunft von *Eucalyptus camaldulensis* aus dem natürlichen Verbreitungsgebiet der Art in Australien, war die Herkunft Lake Albacutya den 22 anderen im Wachstum eindeutig überlegen. Diese hatte im Alter von 5½ Jahren mindestens 2,5 mal soviel Holz produziert wie das Versuchsflächenmittel ausmachte. Dies entspricht zugleich der Wachstumsleistung beim Versuchsanbau in anderen Ländern, die ein ähnliches Klima haben wie Kalifornien. Andere Herkunft aus Südost-Australien, besonders solche aus dem Murray-River-Becken, zeigten ebenfalls eine gute Leistung.

Die Bäume vom Murray River tendierten zu einer mehr gekrümmten und Zwiesel-Stammform als die aus anderen Regionen, und zählten außerdem zu den blühfreudigsten der Versuchsflächen. Die Lake Albacutya-Bäume waren etwas geradschäftiger und weniger verzweigt als die vom Murray River.

Alle erfolgreichen Herkunft zeigten eine enge Korrelation mit dem Niederschlagsmuster (RPC), eine Statistik, die

als eine Korrelation zwischen dem monatlichen Niederschlag am Ursprungsort der Herkunft und dem Niederschlag am Versuchsflächen-Standort definiert ist. Eine Cluster-Analyse des monatlichen Niederschlages, des mittleren Temperatur-Maximums und der Länge der Frostperiode verband die Herkunft vom Murray-River-Becken und vom Lake Albacutya. Erfolgreiche Herkunft hatten eine RPC von über 0, lange Frostperioden über 120 Tage und niedrige maximale Sommerdurchschnittstemperaturen von unter 34°C . Wenn der Standort der Versuchsfläche in die Cluster-Analyse mit einbezogen wurde, verband dies ebenfalls die Lake Albacutya- und die Murray River-Gruppe. Die RPC und das multivariate Verfahren liefern eine Basis, um Zonen gleicher Klimabedingungen objektiv erkennen zu können.

Introduction

River red gum (*Eucalyptus camaldulensis* DEHNH.) has been introduced throughout the Mediterranean basin and various parts of Africa and southern Asia to produce wood for fuel, posts, poles, construction timbers, and hardboard flooring (FAO 1979). Its popularity can be attributed to its rapid growth in subhumid climates that are characterized by extended dry seasons and moderate to high temperatures. In the subhumid coastal zone of California there are considerable acreages of land not currently in productive use, but suitable for the establishment of eucalypt plantations. Demand for biomass fuel for home heating and for cogenerating plants is growing, and eucalypts, such as river red gum, could fill a real need.

Species trials established in 1964 by the U.S. Forest Service at Concord, California, indicated that river red gum grew well and suffered no noticeable damage from drought or frost (KING and KRUGMAN, 1980). Even better growth might be expected because provenances of river red gum are known to differ considerably in performance (ELDRIDGE, 1975). Only one seed source was included in the initial trials at Concord. Therefore, a provenance trial was initiated in 1976 in order to better evaluate adaptability and growth. Presented here are the 5.5 year results of that provenance trial and recommendations for seed sources for commercial planting in central California. Six to ten years is the anticipated rotation for eucalypt fuel biomass production in California, so results at 5.5 years can be extrapolated to rotation age with little risk.

Materials and Methods

Seed from 23 natural stands of river red gum were obtained from the Forestry and Timber Bureau (now CSIRO Division of Forest Research), Canberra, Australia. The stands covered a wide range of latitude, longitude, elevation, and climate (Table 1). Seed were stratified for 60 days, and in June 1975 were germinated in flats under a mist system. They were transplanted into Tinus²⁾ containers (20 cm deep, volume of 340 cc) at the cotyledon stage, and outplanted between February 18 to 26, 1976.

*) Based on a thesis by the senior author presented to North Carolina State University in partial fulfillment of the requirements for the degree of Master of Science.

1) To whom correspondence should be directed.

2) Trade names and commercial enterprises or products are mentioned solely for information. No endorsement by the U.S. Department of Agriculture is implied.

Table 1. — Geographic and climatic data for river red gum provenances.

Provenance ¹	Australian Seed Lot No.	IFG ² Lot No.	Number of Trees ³	Latitude (°S)	Longitude (°E)	Elevation (m)	Natural Region	Average Max. Temp. of Warmest Month (°C)	Average Min. Temp. of Coldest Month (°C)	Length of Frost Period (days)	Total Annual Rainfall (mm)	Rainfall Pattern Correlation ⁴ (r)
A Lake Albacutya, Vict	S 6845	L	10	35°50'	142°00'	60	SUB-HUMID	30.7	3.8	150	419	0.88
B Hamilton, Vict	S 6844	K	10	37°16'	142°17'	305	TEMPERATE	26.3	4.4	125	617	0.87
C Nathalia, Vict	S 6846	M	10	36°00'	145°10'	150	SUB-HUMID	30.4	4.6	138	431	0.80
D Angaston, SA	S 6965-74	AC	10	34°30'	139°00'	610	TEMPERATE	25.1	4.8	149	1190	0.94
E Darlington Pt., NSW	S 6990	AF	8	34°34'	146°01'	90	SUB-HUMID	31.8	3.6	153	407	0.37
F Forbes, NSW	S 6956	X	10	33°20'	147°59'	240	SUB-HUMID	33.3	3.5	133	557	0.00
G Port Lincoln, SA	S 6975-79	AD	5	34°40'	135°50'	30	TEMPERATE	25.2	8.0	0	484	0.96
H Petford, Qld	S 6953	U	5	17°20'	144°57'	520	SUB-MONSOON	34.8	9.3	9	786	-0.76
I Mundiwindi, WA	S 7037	AK	4	23°05'	120°08'	305	ARID	38.1	5.2	65	263	-0.67
J +9 mi. S. Agnew, WA	S 7106	AR	2	28°30'	121°00'	455	ARID	35.9	5.7	76	220	-0.33
K Todd River, Alice Springs, NT	S 6788	I	10	23°38'	133°35'	580	ARID	35.2	3.8	102	267	-0.87
L Onslow, WA	S 7030	AI	3	22°02'	115°01'	30	ARID	35.7	10.8	0	264	-0.05
M Adavale Rd., Quilpie, Qld	S 6963	AB	3	26°35'	144°18'	190	SEMI-ARID	36.8	5.4	92	502	-0.81
N Pentland, Qld	S 6945	R	1	20°35'	145°27'	455	SEMI-ARID	36.1	11.9	7	486	-0.75
O Hughenden, Qld	S 6949	T	5	20°43'	144°21'	395	SEMI-ARID	37.4	8.1	7	486	-0.75
P Wiluna, WA	S 7046	AM	10	26°34'	120°03'	490	ARID	37.2	4.9	24	225	-0.21
Q Temple Bar Creek, Alice Springs, NT	S 6716	H	1	23°50'	133°50'	550	ARID	35.2	3.8	102	267	-0.87
R 20 mi. S. Agnew, WA	S 7052	AO	5	28°15'	120°40'	335	ARID	35.9	5.2	76	220	-0.33
S Charleville Rd., Quilpie, Qld	S 6871	P	10	26°52'	144°49'	225	SEMI-ARID	36.8	5.4	92	502	-0.81
T Thargomindah, Qld	S 6872	Q	4	27°58'	143°43'	120	ARID	36.7	5.8	37	272	-0.82
U Three Springs, WA	S 7084	AQ	4	29°20'	115°35'	305	SUB-HUMID	35.4	7.0	0	394	0.87
V Katherine, NT	S 6869	N	10	14°25'	132°15'	110	MONSOON	38.0	13.2	0	951	-0.83
W Newcastle Waters, NT	S 7080	AP	10	17°30'	133°30'	215	SUB-MONSOON	39.1	11.1	0	456	-0.78

¹) State Abbreviations: Victoria (Vict), South Australia (SA), New South Wales (NSW), Queensland (Qld), Western Australia (WA), Northern Territory (NT).

²) Institute of Forest Genetics lot number.

³) Number of trees represented in seed collection.

⁴) The correlation coefficient between mean monthly precipitation at Concord and at the provenance origin. Provenance with a positive or zero correlation have winter rainfall maximum, those with negative coefficients, a summer maximum.

The climate of the planting site at Concord, although located only 32 km east of San Francisco, is typical of many dry, inland areas of California. It is separated from the coast by a range of low hills. The long-term mean annual precipitation is 366 mm, almost all during the winter months (U.S. Department of Commerce, 1964). During the

decade 1968 to 1978, precipitation ranged from 150 to 500 mm per year (U.S. Department of Commerce, 1969—1978). From May through September less than 20 mm of rain falls, with peak rainfall of 83 mm in January alone. The mean annual temperature is 14.2° C, ranging from 6.9° C for the coldest month to 20.7° C for the warmest (U.S. Department of Commerce, 1964). The mean daily minimum for January, the coldest month, is 2.9° C and the lowest records are about -7.2° C. The mean daily maximum for July, the warmest month, is 32.9° C and the highest recorded temperatures are about 45.6° C. The elevation of the planting site is 62 m, and soils tend to be deep, well-drained clay or loam of good agricultural quality.

The site was disced after the autumn rains began in 1975 and rototilled just prior to planting in February, 1976. Planting holes were prepared with a power-driven augur. Spacing was 3.05 m × 3.05 m. Grass in the first 9 blocks was controlled with simazine ³⁾, sprayed just after planting and again in September 1977. Block 10, later dropped from the analysis, was sprayed with devrinol. Open-ended, cardboard milk cartons were placed over seedlings to reduce damage from drying winds and from rodents. The area was disced in March or April of 1977, 1979, and 1980 for fire protection. Seedlings were hand-irrigated with about 4 liters of water per seedling at the time of planting and monthly in the first year during the dry season.

³⁾ This publication does not represent a recommendation for the pesticides reported, nor does it imply that they have been registered by the appropriate government agency.

The year of planting and the following year were two of the worst drought years on record in California.

The planting design was a randomized complete block with 10 blocks. In each block, each provenance was represented by a 4-tree row plot. Two of the 10 blocks were affected by competing herbaceous vegetation and had relatively high mortality (28%) compared to the others (7%). These two blocks were omitted from the analysis.

Survival and second-year heights were recorded in 1977. In June of 1981, 5.5 years after planting, measurements were taken on height and diameter at breast height, using a height-pole and diameter-tape. Relative volume was estimated by $V = (\text{diameter})^2 \times (\text{height})$, an index rather than a true measure of yield. Straightness was scored from 1 to 5. In scoring, an attempt was made to distribute the scores more or less normally; the average score, 3, occurs about 40% of the time, scores 2 and 4, 20% of the time each, and scores 1 and 5, 10% of the time each. Reproductive output was determined by estimating the proportion of the crown that supported capsules or flowers: if less than one-third of the crown supported capsules or flowers, it was considered to be light, one-third to two-thirds was medium, and over two-thirds was heavy. Forking was recorded in terms of the number of stems forked at the base and those forked higher in the tree. Basal forking is, perhaps, an indication of dieback from frost damage in the year of establishment.

Statistical analyses were made on the plot means with the Statistical Analysis System computing package (HELWIG and COUNCIL, 1979). The analysis of variance and Duncan's

Table 2. — Provenance means for relative volume, height at 1.5 years and 5.5 years, dbh, straightness, and forking.

Provenance	Volume (cu m)	Height 1.5 (m)	Height 5.5 (m)	DBH (cm)	Straightness	Trees forked at base (%)	Trees forked above base (%)
A Lake Albacutya	.0684 a	1.32 abc	7.17 a	9.5 a	2.4 ghij	3.1 c	34.4 abode
B Hamilton	.0570 ab	1.23 bcde	5.97 bc	8.9 ab	2.6 efghij	3.1 c	52.0 ab
C Nathalia	.0463 abc	1.58 ab	5.78 bcd	7.8 abc	2.9 cdefg	0.0 c	47.6 abc
D Angaston	.0462 abc	1.15 bce	5.85 bcd	8.5 ab	2.4 ghij	6.3 bc	40.5 abc
E Darlington Point	.0421 bcd	1.26 bcde	6.32 ab	7.6 abcd	2.6 efghij	12.5 abc	40.5 abc
F Forbes	.0297 cde	0.98 cdef	5.40 bcde	7.2 bcde	2.5 efghij	9.4 abc	43.8 abc
G Port Lincoln	.0270 cde	0.91 cdef	4.94 cdef	7.1 bcde	2.1 j	10.4 abc	58.0 a
H Felford	.0253 cde	1.77 a	5.99 bc	6.2 cdef	3.2 bc	0.0 c	30.0 abcde
I Mundiwindi	.0230 cde	0.91 cdef	4.98 cdef	6.0 cdef	3.4 ab	12.5 abc	22.9 bcde
J Agnew	.0224 cde	1.02 cdef	4.85 def	6.1 cdef	3.0 bcdef	15.7 abc	20.9 cde
K Alice Springs	.0215 cde	1.10 bcdef	4.54 ef	5.7 cdef	2.8 cdefgh	18.8 abc	7.3 de
L Onslow	.0201 de	1.02 cdef	4.36 ef	6.0 cdef	2.5 fghij	24.9 abc	4.1 e
M Quilpie	.0200 de	0.89 cdef	5.19 cdef	5.8 cdef	3.1 bcd	9.4 abc	25.0 bcde
N Pentland	.0194 de	1.10 bcdef	4.85 def	5.8 cdef	2.6 efghi	3.1 c	52.0 ab
O Hughenden	.0191 de	1.20 bcdef	4.79 def	5.8 cdef	2.8 cdefgh	15.6 abc	41.6 abc
P Wiluna	.0180 de	0.72 f	4.99 cdef	5.5 cdef	3.8 a	6.3 bc	31.3 abcde
Q Alice Springs	.0173 de	0.81 def	4.89 def	5.3 ef	3.0 bcde	6.3 bc	35.4 abcd
R Agnew	.0170 de	1.26 bcde	4.82 def	5.5 def	2.9 cdef	16.6 abc	27.0 bcde
S Quilpie	.0148 e	0.97 cdef	4.55 ef	5.5 def	2.2 ij	13.5 abc	32.1 abcde
T Thargomindah	.0146 e	1.04 cdef	4.18 f	5.1 ef	2.3 hij	25.0 abc	21.9 bcde
U Three Springs	.0124 e	0.77 ef	4.29 f	5.2 ef	2.7 defgh	6.3 bc	25.0 bcde
V Katherine	.0107 e	0.98 cdef	4.36 ef	4.6 f	3.0 bcde	6.3 bc	25.0 bcde
W Newcastle Waters	.0098 e	0.88 cdef	4.14 f	4.3 f	2.7 defgh	28.0 abac	27.1 bcde
Error Std. Sev.	.0221	0.56	0.92	1.9	0.6	16.8	25.4




Figure 1. — Relative volume of 23 provenances of river red gum grown in California. Letters (A, B, . . . , W) refer to provenances listed in Table 1.

New Multiple Range Test were used to separate and rank provenances. SAS stepwise regression and cluster analysis (PROC CLUSTER) were used to describe provenance performance based on climatic variation at the provenance origin.

Results

Provenance mean height at 5.5 years ranged from 4.1 to 7.2 m, and relative volume from 0.010 to 0.068 cu m. The largest tree was 12.1 m tall and 19.8 cm dbh. Significant differences were found among provenances for volume, diameter, height, straightness, forking, and flowering (Table 2 and Fig. 1)

Despite extensive overlap in volume growth, as indicated by Duncan's New Multiple Range Test, there seems to be a major gap that separates provenances from the Murray River system and adjacent areas in southeastern Australia, which rank as the top five, from the rest of Australia, which do poorly in comparison (Table 2). The top five provenances are Lake Albacutya, Hamilton, and Nathalia in Victoria, Angaston in South Australia, and Darlington Point in New South Wales. In the ranking of provenance means for diameter growth, the same five provenances are joined in the top group by two other provenances that are in or near the Murray River basin, one to the northeast at Forbes, New South Wales, and the other to the west at Port Lincoln, South Australia (Table 2). Provenances from more distant areas are inferior and do not differ significantly from each other. The provenances also fall into two groups for height, but there is more overlap (Table 2). Though trees from Port Lincoln are superior in diameter growth, they are not sufficiently superior in height growth

to be included in the top group for volume production. The Petford provenance is superior in height growth but only average in diameter growth, resulting in only mediocre volume production. The Forbes provenance is marginally superior for both diameter and height but its mean volume production is sufficiently low to exclude it from the top group.

Two provenances from Western Australia, Wiluna and Mundiwindi, were the straightest, followed by the Petford and Quilpie sources from Queensland and the Katherine and Alice Springs sources from the Northern Territory. Although the provenances from southeastern Australia were superior for growth traits, they fall within the most crooked group, except for Lake Albacutya. It was average with regard to straightness, ranking 9 out of 23.

The provenances from southeastern Australia as a whole tend to have 12.5% or less basal forking. Eight of the remaining 17 provenances have more than 12.5% basal forking (Table 2). Basal forking may indicate susceptibility of young seedlings to freezing, and its low incidence in the provenances from southeastern Australia is reasonable because they originated from the latitudes with the coldest climates (Table 1). However, this group surpasses most other provenances in forking above the base (greater than 34.4% forked).




Figure 2. — Flowering classes of river red gum at 5.5 years of age. Letters (A, B, . . . , W) refer to provenances listed in Table 1.

Table 3. — Correlations among growth and form characteristics in river red gum, based on provenance means (in parentheses) and trees within provenances.

	Dbh	Height 1.5	Height 5.5	Straightness	Flowering	Basal Forking	Forking Above Base
Volume	0.88	0.66	0.79	-0.06	-0.37	-0.50	0.44
Dbh		---	(0.93) 0.66	(0.33) -0.08	(-0.68) -0.34	(0) -0.04	(0.35) 0.12
Height 1.5			(0.59)	-0.13	-0.29	---	---
Height 5.5				(0.06) -0.22	(-0.51) -0.33	(0) -0.05	(0.04) -0.02
Straightness					(-0.53) 0	(0) 0.13	(0.84) 0.24
Flowering						(0) 0.02	(-0.65) -0.02
Basal Forking							(0) -0.03

Flowering was moderate to heavy in over 80% of the trees from southeastern Australia. Three provenances from Western Australia also had more than 80% of their trees in the moderate to heavy flowering category, while provenances Thargomindah and Quilpie from Queensland had less than 40% (Fig. 2).

Provenance means for height at 5.5 years were fairly well correlated with height after two growing seasons ($r = 0.59$; Table 3). All five of the provenances from southeastern Australia ranked near the top after only two seasons. Provenances from Agnew in Western Australia, Hughendon and Pentland in Queensland, and Alice Springs in the Northern Territory fell from the upper ranks over time, while those from Port Lincoln and Forbes improved significantly. The tallest provenance at 1.5 years was from Petford, but it subsequently dropped to third place.

Correlations among provenance means revealed only weak correlations between stem straightness or forking on the one hand, and growth on the other. The tendency was for negative associations. The combined association of crooked stems and forking with growth meant that the Murray River group had relatively poor form. The most heavily flowering provenances tended to be slower-growing. Correlations among individuals within provenances

were generally similar in sign to those among provenance means but much weaker, suggesting few barriers to simultaneously selecting for superior growth and good form.

Discussion

Provenances from the Murray and Darling River system and peripheral areas of Victoria, South Australia, and New South Wales had superior growth in our California planting, as well as on sites with summer drought in Greece (PANETOS, 1970), Israel (KARSCHON, 1974; MORESHET, 1981), Italy (GIORDANO, 1974), MOROCCO (DESTREMAU *et al.*, 1973), Rhodesia (BARRETT and CARTER, 1976), Zambia, Spain, Algeria, and Portugal (LACAZE, 1977). Even in Pakistan, with a monsoon climate but with a winter precipitation maximum, results from an irrigated trial reported by SIDDQUI *et al.* (1979) are similar to those in California. For 9 seed sources common to both tests, the correlation between relative volume in California and volume per hectare per year at 10 years of age in Pakistan was $r = 0.73$ ($P < 0.05$; Fig. 3).

The Lake Albacutya provenance is consistently outstanding in Mediterranean climates like California's, which may be attributed to its adaptation to a low summer rainfall regime. The Lake Albacutya population is located around the margin of a lake bed that is usually dry.

In our trial, the Lake Albacutya provenance had 20% greater volume than the Hamilton, Victoria provenance, the next best, which is on the Glenelg River system that flows southward away from the Murray River. The Lake Albacutya provenance was 161% better than the mean of all provenances and was 6 times or 598% better than the worst provenance, Newcastle Waters, Northern Territory. Even in the two replications omitted because of weed competition, the Lake Albacutya provenance performed outstandingly well. When those two replications were included, the Lake Albacutya provenance had 63% more volume than the next best provenance and was 200% better than the plantation mean. River red gum was recommended for California based on the growth of a seed source of unknown origin received from Brazil (KING and KRUGMAN, 1980). After 6 years at Concord, height of that provenance was 6.2 m and dbh was 6.6 cm compared to 7.2 m and 9.5 cm for the Lake Albacutya provenance after only 5.5 years, which included two of the severest drought years in California's recorded history.

It might be anticipated that the best provenances for California are those whose climate-of-origin correspond most closely in pattern to that at Concord. We calculated the correlation between monthly rainfall at a meteorological




Figure 3. — Correlation between volume of river red gum provenances at 5.5 years in California and volume growth to 10 years in Pakistan. Letters (D, E, . . . , W) refer to provenances listed in Table 1.

station representing each of the 23 provenance origins (HALL, 1972) with monthly rainfall at Concord during the decade 1969–1978 (U.S. Department of Commerce, 1969–1978), keeping in mind the opposition of the seasons in the Northern and Southern Hemispheres; i.e., January in California corresponds to July in Australia, February with August, etc. A simple correlation coefficient was then calculated from the 12 pairs of observations for each provenance and the Concord planting site. The correlations express the similarity between the rainfall pattern at Concord and the pattern at each of the 23 provenance locations. The correlations ranged from 0.96 between Concord and Port Lincoln, South Australia, a summer dry locality, to –0.87 between Concord and Alice Springs, Northern Territory, which has a summer rainfall maximum (Table 1). These rainfall pattern correlations (RPC) explained much of the variation among provenances in volume (Table 4). Nevertheless, the relationship did not appear smooth

Table 4. — Correlations between provenance growth and form characteristics and the climate at provenance origin¹⁾.

	Volume	Dbh	Height 5.5	Basal Forking	Forking Above Base
Latitude	0.72***	0.79***	0.58**	-0.31	0.47*
Longitude	0.42*	0.40	0.51*	--	--
Elevation	-0.12	-0.09	0.05	--	--
Mean Maximum Temperature	-0.75***	-0.83***	-0.60**	0.44*	-0.66**
Mean Minimum Temperature	-0.46*	-0.50*	-0.46	--	--
Length of Precipitation	0.69***	0.70***	0.67***	-0.29	0.22
RPC	0.72***	0.76***	0.54***	-0.41	0.45

¹⁾ Significance indicated by: ***, $P < 0.01$; **, $0.01 < P < 0.05$; *, $0.05 < P < 0.10$.




Figure 4. — Relationship of provenance mean volume of river red gum at 5.5 years in California to rainfall pattern correlation, a statistic correlating monthly rainfall at provenance origin with rainfall at the planting site. Letters (A, B, . . . , W) refer to provenances listed in Table 1.




Figure 5. — River red gum locations in Australia, clustered by rainfall amount and pattern. Letters (A, B, . . . , W) refer to provenances listed in Table 1.

(Fig. 4). Provenances from areas with an RPC of zero or less did uniformly poorly. Provenances from areas with an RPC of circa 0.8, ranged from poor to exceptionally good, with no obvious relationship between volume and RPC. The poor performance of provenances such as Three Springs, Western Australia, with an RPC of 0.87, must be explained by other factors.

Cluster analysis was used to examine the 23 provenances for evidence of regions of similar rainfall patterns (Fig. 5). The degree of similarity among provenances was expressed by pairwise Euclidean distances that were based on the monthly rainfall values. It was arbitrarily decided to terminate the clustering process at six clusters because these clusters represented six easily described rainfall regions (Fig. 5), climates with a winter rainfall maximum and high, medium, or low total precipitation, and those with a summer maximum and either high, medium, or low total precipitation. Summer rainfall predominates in northern latitudes and winter rainfall in southern latitudes; the highest rainfall is found along the coast in both regions, while the lowest rainfall is found inland. Monthly rainfall for the provenances within each cluster were averaged and plotted over the twelve months (Fig. 6) for comparison with the rainfall pattern at Concord. The top seven provenances for growth at Concord were all in the clusters with highly positive RPC values, the three winter rainfall clusters.

Growth was also correlated strongly with provenance latitude, length of the frost period, and mean maximum temperature in the warmest month (Figs. 7–9, Table 4). However, the relationships with latitude and frost period, in particular, do not appear smooth (Figs. 7, 8). Provenances that did best in Concord came from temperate latitudes in Australia ($> 32^{\circ}$ S) with mainly winter rainfall ($RPC > 0$), long frost periods (> 120 days), and low summer maximum temperatures ($< 34^{\circ}$ C). For comparison, the planting site is located at $38^{\circ}01'$ N, has a frost period of 125 days (U.S. Department of Commerce, 1974) and mean maximum temperature of the warmest month is 30.3° C (U.S. Department of Commerce, 1964).

It is apparent that the occurrence of winter rainfall and the length of the frost period at the provenance origin,




Figure 6. — Monthly rainfall for six river red gum provenance clusters and the Concord, California planting site. Letters (A, B, . . . , W) refer to provenances listed in Table 1.




Figure 7. — Relationship of river red gum provenance mean volume at 5.5 years in California with latitude of origin. Letters (A, B, . . . , W) refer to provenances listed in Table 1.

when combined, account for most of the observed provenance variation in growth response at Concord, although both maximum temperature and latitude are predictive. Based on rainfall pattern correlation alone, the Three Springs, Western Australia provenance should do well at Concord. Its poor growth is rationalized with reference to the complete lack of frost in its native habitat.

It is reasonable to expect a greater degree of frost tolerance among the more southerly provenances, because some of the northern provenances never experience frost in their native regions. Frost is a definite factor at Concord; in December 1972, temperatures dropped below freezing for nine consecutive nights, reaching a low of -9°C (U.S. Department of Commerce, 1969—1978). PRYOR and BYRNE (1969) found that seedlings of southern provenances of river red gum were more frost resistant than northern provenances, but AWE and SHEPHERD (1975) claimed to find the opposite relationship.

It is conceivable that the southeast Australian provenances, which are adapted to striking deep roots rapidly, were successful in California because they were able to exploit deep moisture during the dry season. Provenances from Lake Albacutya and Nathalia, Victoria (Murray River drainage) germinated more vigorously than northern provenances when tested on a soil brought to field capacity and then left with no additional application of moisture, simulating a winter rainfall pattern (AWE *et al.*, 1976). They




Figure 8. — Relationship of river red gum provenance mean volume at 5.5 years in California with length of frost period at the provenance origin. Letters (A, B, . . . , W) refer to provenances listed in Table 1.

also produced the greatest root dry weight and number of roots (AWE *et al.*, 1976; MORESHET, 1981).

The provenance from Three Springs, Western Australia is adapted to the same rainfall pattern as the southeast Australian provenances, and yet it ranks third from the bottom in growth response. This may be due to the fact that there is no frost period in Three Springs, although the percentage of forking at and above the base, common signs of frost damage, are quite low and do not indicate that it has experienced cold damage. Trees from this provenance show an affinity for Western Australian flooded gum (*E. rudis* ENDL.) and may be hybrids, accounting for their deviant behavior. Similarly, the Petford, Queensland provenance, which performs much better in height growth than expected for a provenance of its latitude, length of frost period, and rainfall pattern, shows evidence of introgression with forest red gum (*E. tereticornis* SM.), according to PRYOR and BYRNE (1969).

Paradoxically, the Port Lincoln source, which is native to a frost-free climate, performs as well as most of the other winter rainfall provenances in diameter growth. The Port Lincoln provenance had a high root:shoot ratio, even higher than seedlings from Lake Albacutya, when grown in Israel (SANDS, 1981), a factor that, perhaps, contributed to its success at Concord. The Port Lincoln provenance is unique: it consists of just a few trees growing on a limestone outcrop, an unusual site for the species. It grows well on soils of high pH in Pakistan (QUADRI, 1983) and in Italy (LACAZE, 1977; TURNBULL, 1973). It deserves attention for its potential on calcareous soils.

River red gum may not be the ideal species for all sites in California. For example, it was outproduced by manna gum (*E. viminalis* DEANE et MAID.) on some sites (STANDIFORD and DONALDSON, 1982). However, when river red gum is planted, the Lake Albacutya provenance should be used, based on our results and its consistently superior perfor-




Figure 9. — Correlation of river red gum provenance mean volume at 5.5 years in California with mean maximum temperature in the warmest month at the provenance origin. Letters (A, B, . . . , W) refer to provenances listed in Table 1.

mance in countries which, like California, have Mediterranean climates. It may also prove particularly valuable on saline soils (SANDS, 1981). Its tendency to crookedness should be little deterrent for its use as fuel biomass. Other provenances should be introduced from the Murray River, Wimmera River, and adjacent drainages to increase the genetic base. More intensive sampling of this area may reveal provenances of equal or better performance than those already tested. Mother-tree identities should be retained in future testing to facilitate selection within provenance collections. Gains can almost certainly be made by selection within provenances, since KARSCHON (1974) has already demonstrated the existence of within-stand variation. Incidentally, the concurrence of results among countries with Mediterranean climates suggests that progress could be made by exchange of selected materials.

Acknowledgements

This study was supported in large part by the U.S. Department of the Navy. We thank J. W. TURNBULL, CSIRO Division of Forest Research, Canberra for his help in supplying seed of river red gum and reviewing this manuscript, J. P. KING, who designed and initiated the field trial, and to K. G. ELDRIDGE and Y. ZOHAR for helpful reviews.

Literature Cited

- AWE, J. O. and SHEPHERD, K. R.: Provenance variation in frost resistance in *Eucalyptus camaldulensis* DEHN. Aust. For. 38: 26—33 (1975). — AWE, J. O., SHEPHERD, K. R. and FLORENCE, R. G.: Root development in provenances of *Eucalyptus camaldulensis* DEHN. Aust. For. 39: 201—209 (1976). — BARRETT, R. L. and CARTER, D. T.: *Eucalyptus camaldulensis* provenance trials in Rhodesia. 2. Later results. Rhod. Bull. For. Res. 2 (pt. 2). 39 pp. (1976). — DESTREMAU, D. X., JOLLY, H. and KORBA, M.: Information on the comparative trial of *Eucalyptus camaldulensis* at Sidi Slimane. Ann. Rech. For. Maroc. 13: 121—154 (1973). — ELDRIDGE, K. G.: An annotated bibliography of genetic variation in *Eucalyptus camaldulensis*. Trop. For. Pap. No. 8. Dep. For., Commonw. For. Inst., Univ. Oxford and CSIRO, Div. For. Res., Canberra (1975). — FAO: Eucalypts for planting. FAO, U.N., Rome (1979). — GIORDANO, E.: Osserva-

Silvae Genetica is an international Journal, which continues *Zeitschrift für Forstgenetik und Forstpflanzenzüchtung* (Journal of Forest Genetics and Forest Tree Breeding) founded by W. LANGNER in 1951. In *Silvae Genetica* original articles, short notes, reviews, announcements and reports on meetings and congresses on Forest Tree Breeding and Genetics and related fields are published in German, English or French.

Reproduction of contributions is not permitted; reproduction of illustrations is permitted only with the approval of the author and the publisher.

Authors should send two copies of their manuscript to one of the regional co-editors, generally on close to the institute from which the article originates. Authors are however free to select an appropriate co-editor who should be a native speaker of the language of the paper and familiar with the subject matter.

Original articles, including tables, illustrations and literature must not exceed six printed pages (equivalent to 24 typed pages with 30 lines at double spacing and with a 5 cm border). Longer papers can only be accepted in exceptional cases; additional pages will be charged to the author at a price of DM 220,— per page. The same applies to extra costs for special plates. — In preparing the manuscript authors are asked to conform to the following arrangements. The title page should contain the following information: a) Title of the paper; b) Christian and surnames of the authors; c) Institute where the article has been written; d) if necessary, the present address of the authors as footnote; e) up to 10 key words to facilitate referencing of the paper; and on a separate special page f) a summary in the original language of the paper and one in German (which can, if necessary, be revised in the editorial office).

For financial and technical reasons tables, figures and mathematical terms or formulae must be submitted separately from the text and ready for photoprinting. This applies especially to mathematical formulae which should be numbered and brought together in the finished state into one or few groups.

Manuscripts ready for printing will be registered the month they reach the editorial office in Grosshansdorf-Schmalenbeck. They will be published in the order of their receipt. Their rearrangement for technical reasons is the prerogative of the editorial manager.

Short notes may be up to one page in length (about 4 pages of typescript) and should also be classified by some key words. They will be published as soon as possible.

Announcements and reports on meetings and congresses should be made early and quote the organizer, subject, possible working groups, the place and period of the meeting. Reports should summarize briefly events of importance to forest tree breeders and forest geneticists and indicate points asked for in announcements.

Galley proofs of original articles and short notes will be sent to authors for correction. They must be returned with the least possible delay. The only corrections permitted are those arising during typesetting. No amendments of the manuscript are permitted. Corrections on announcements, reports, and reviews will be made in the editorial office.

Reprints: Authors obtain, free of charge, up to 30 reprints of their articles. Addi-

Silvae Genetica ist eine internationale Zeitschrift; sie setzt die im Jahre 1951 von W. LANGNER begründete *Zeitschrift für Forstgenetik und Forstpflanzenzüchtung* fort. *Silvae Genetica* veröffentlicht Originalarbeiten, Kurzmittelungen, Besprechungen sowie Hinweise und Berichte über Tagungen und Kongresse über Genetik und Züchtung sowie verwandte Fachgebiete in deutscher, englischer oder französischer Sprache, soweit sie für Forstgenetik und Forstpflanzenzüchtung von Bedeutung sind.

Der Nachdruck der Beiträge ist nicht gestattet; der Nachdruck von Abbildungen nur mit Genehmigung des Verfassers und des Verlags.

Manuskripte werden in zweifacher Ausfertigung im allgemeinen an einen der regionalen Mitarbeiter in der Nähe der Institution erbeten, aus der der Artikel stammt. Autoren sind jedoch frei, einen entsprechenden Mitarbeiter zur Durchsicht ihrer Arbeit zu wählen. Er sollte die Sprache, in der der Artikel abgefaßt ist, als Muttersprache beherrschen und mit dem bearbeiteten Problem vertraut sein.

Originalarbeiten einschließlich Tabellen Abbildungen und Literaturhinweisen sollen nicht über 6 Druckseiten hinausgehen. (Dieser Umfang entspricht ungefähr 24 Schreibmaschinenseiten mit 30 Zeilen mit doppeltem Abstand und 5 cm Rand). Längere Arbeiten können nur in Ausnahmefällen angenommen werden. Zusätzliche Seiten werden dem Autor zum Preis von 220,— DM pro Seite in Rechnung gestellt. Das gleiche gilt für spezielle Druckverfahren. — Die Anordnung der Manuskripte soll in der für *Silvae Genetica* üblichen Form geschehen. Die Titelseite soll die folgenden Informationen aufweisen: a) Titel der Arbeit, b) Vor- und Zunamen der Autoren, c) Institut, in dem die Arbeit entstanden ist, d) wenn notwendig, die gegenwärtige Anschrift der Verfasser als Fußnote, e) bis zu 10 Schlagworte zur Charakterisierung des Inhalts der Arbeit und zur leichteren bibliographischen Erfassung und außerdem auf besonderem Blatt f) eine Zusammenfassung in der Originalsprache der Arbeit und eine weitere in deutsch. Letztere kann in der Redaktion überarbeitet werden.

Aus finanziellen und technischen Gründen sollen Tabellen, Abbildungen und mathematische Ausdrücke oder Formeln getrennt vom Text als druckfertige Klischee-Vorlagen für den Photo-Druck eingereicht werden. Dies gilt besonders auch für im Text vorkommende mathematische Ausdrücke, die mit entsprechender Kennzeichnung (Numerierung) in einer oder wenigen Klischee-Vorlagen zusammengefaßt werden sollen.

Druckfertige Manuskripte werden mit dem Monat ihres Eingangs in die Schriftleitung in Großhansdorf-Schmalenbeck registriert und in der Reihenfolge ihres Eingangs veröffentlicht. Aus technischen Gründen kann der Redakteur jedoch eine andere Entscheidung treffen.

Kurzmittelungen können bis zu einer Druckseite (etwa 4 Schreibmaschinenseiten) lang sein und sollten ebenfalls mit einigen Schlagworten charakterisiert werden. Sie erscheinen baldmöglichst außer der Reihe.

Ankündigungen und Berichte über Tagungen und Veranstaltungen sollten frühzeitig geschehen und den Veranstalter, das Thema, ev. Arbeitsgruppen, den Ort und den Zeitraum der Veranstaltung nennen. — Berichte sollen die für den Forstpflanzenzüchter und Forstgenetiker stattgefundenen wichtigen Ereignisse in gedrängter Form mit Hinweis auf die unter Ankündigungen geforderten Punkte zusammenfassen.

Silvae Genetica est un Périodique international qui est la suite de *Zeitschrift für Forstgenetik und Forstpflanzenzüchtung* (Journal of Forest Genetics and Forest Tree Breeding) fondé en 1951 par W. LANGNER et publie des articles originaux, en allemand, anglais ou français, ainsi que des Notes et des mises au point annonces et des compte-rendus de Réunions et de Congrès sur l'Amélioration des Arbres Forestiers, la Génétique et les disciplines voisines. La reproduction des articles n'est pas autorisée, la reproduction des illustrations n'est autorisée qu'avec l'approbation de l'auteur et de l'éditeur.

Les auteurs doivent envoyer deux exemplaires de leur manuscrit à l'un des co-éditeurs régionaux, en général le plus proche de l'Institut dont l'auteur de l'article fait partie. Toutefois, les auteurs sont libres de choisir un co-éditeur de même langue que celle de l'article, et à qui le sujet de l'article soit familier.

Les articles originaux y compris les tableaux, les figures et la bibliographie, ne doit pas dépasser 6 pages imprimées (soit 24 pages dactylographiées de 30 lignes, à double interligne et avec une marge de 5 cm).

Les articles plus longs ne peuvent être acceptés que dans des cas exceptionnels; les pages supplémentaires sont facturées à l'auteur au prix de 220 DM la page. Les mêmes conditions s'appliquent aux frais supplémentaires pour les clichés spéciaux. Pour la préparation du manuscrit, il est demandé aux auteurs de se conformer aux indications suivantes: La page de titre doit contenir les informations suivantes a) Titre de l'article; b) Nom et prénoms des auteurs; c) Institut auquel l'auteur appartient; d) si nécessaire, l'adresse actuelle des auteurs en note infrapaginale; e) des mots clés permettant de faciliter la classification de l'article — dans le limite de 10; et, sur une page séparée f) un résumé dans la langue originale de l'article — un résumé en allemand (qui peut, si nécessaire, être revu par le Bureau de rédaction).

Pour des raisons financières et techniques, les tableaux, les figures et les termes ou formules mathématiques doivent être présentés séparés du texte et prêts à l'impression-photo. Ceci s'applique particulièrement aux formules mathématiques qui doivent être numérotées, et rassemblées dans leur état définitif en un ou plusieurs groupes.

Les manuscrits prêts à l'impression seront enregistrés dans le mois de leur arrivée au Bureau de Rédaction de Grosshansdorf-Schmalenbeck. Ils sont publiés dans l'ordre chronologique de leur arrivée au Bureau de Rédaction. Le Rédacteur peut toutefois modifier cet ordre pour des raisons techniques.

Les notes peuvent avoir jusqu'à 1 page (environ 4 pages dactylographiées) et doivent être indexées par quelques mots clés. Elles seront publiées aussi rapidement que possible.

Annonces et Compte-rendus de Réunions et de Congrès. Les annonces doivent être communiquées très tôt et indiquer l'Organisateur, le sujet, les éventuels résumer brièvement les faits importants pour Améliorateurs forestiers et les Généticiens forestiers, et comporter les mêmes indications que les Annonces de Réunions et Congrès.

Les épreuves d'imprimerie des articles originaux et des notes seront envoyées aux auteurs pour correction. Elles doivent être retournées dans le délai le plus bref pos-

tional reprints may be purchased if ordered in advance from the publisher.

Publication schedule: 6 numbers each year.

Subscription: *Silvae Genetica* may be ordered through book-sellers in Germany and other countries, or directly from the publisher. Subscriptions are effective for a complete volume and continue in force unless terminated following delivery of the last number of a volume.

Price of subscription for the 6 numbers of a volume, DM 248,—; for students DM 198,40; postal expenses are extra.

This journal is covered by Biological Abstracts, Biosciences Information Service of Biological Abstracts, Chemical Abstracts, Current Contents (Series Agriculture, Biology and Environmental Sciences) of Institute for Scientific Information, Forestry Abstracts, Institute for Scientific Information of the Academy of Sciences of the USSR, Cambridge Scientific Abstracts, the Internationale Bibliographie der Zeitschriftenliteratur and Internationale Bibliographie der Rezensionen.

Advertisements: Inquires about the size and price of advertisements should be addressed to the publisher.

Druckfahnen von Originalarbeiten und Kurzmittellungen werden den Autoren zur Korrektur zugesandt. Sie sollen unter Verwendung der international üblichen Symbole korrigiert und möglichst umgehend an die Schriftleitung zurückgeschickt werden. Dabei sind nur solche Fehler zu korrigieren erlaubt, die auf fehlerhaften Satz zurückzuführen sind. Änderungen des Manuskripts sind nicht gestattet. Die Korrekturen von Ankündigungen, Berichten und Besprechungen werden in der Schriftleitung durchgeführt.

Sonderdrucke: Die Verfasser erhalten von ihren Arbeiten bis zu 30 Sonderdrucke kostenlos. Bei rechtzeitiger Bestellung bestehen für weitere Sonderdrucke Bezugsmöglichkeiten gegen Berechnung.

Erscheinungsweise: 6 Hefte im Jahr.

Bezugsmöglichkeiten: *Silvae Genetica* kann durch den in- und ausländischen Buchhandel oder direkt vom Verlag bezogen werden. Das Abonnement läuft weiter, wenn nicht unmittelbar nach der Lieferung des Schlußheftes eines Bandes eine Abbestellung erfolgt.

Bezugspreis für die 6 Hefte des Bandes beträgt DM 248,— (für Studenten DM 198,40) zuzüglich Versandkosten.

Anzeigen: Anfragen über Preise und Größe von Anzeigen werden an den Verlag erbeten.

sible. Les seules corrections permises sont celle qui ont rapport à la typographie. Aucune modification du manuscrit n'est pas permise. Les corrections des annonces, compte-rendus et mises au point sont fait dans le Bureau de Rédaction.

Tirés à part: Les auteurs peuvent obtenir gratuitement jusqu'aux 30 tirés à part de leur article. Les quantités supplémentaires peuvent être achetées par commande à l'avance auprès de l'éditeur.

Périodicité de publication: 6 fascicules par an.

Abonnement: On peut s'abonner à *Silvae Genetica* auprès des libraires en Allemagne ou dans les autres pays, ou bien directement auprès de l'éditeur. L'abonnement est valable pour un volume complet, et se poursuit automatiquement, à moins que l'abonné ne demande qu'il se termine après le dernier numéro du volume en cours.

Prix de l'abonnement: pour les 6 numéros d'un volume: 248,— DM — pour les étudiants: 198,40 DM. Les frais d'expédition sont en sus. **Publicité:** Les demandes d'informations concernant les prix et la dimension des annonces publicitaires doivent être adressées à l'éditeur.

Neuerscheinung:

Einführung in die Genetik

für Studierende der Forstwissenschaft

VON H. H. HATTEMER UND F. BERGMANN

268 Seiten mit 86 Abbildungen und 59 Tabellen

Kartonierte DM 22,80

Der erste Teil des Buches stellt zunächst kurz die molekularen und zytologischen Grundlagen der Genetik vor, behandelt die verschiedenen Formen der Mutation als Quelle genetischer Variation und beschäftigt sich eingehend mit den Methoden der genetischen Analyse, um die Zusammenhänge zwischen dem variablen Phänotyp und seinem kontrollierenden Genotyp aufzudecken.

Der zweite, der elementaren Populationsgenetik gewidmete Teil, erfährt seine Ausrichtung an der Evolutionsbiologie. Schwerpunktmäßig werden die Sexualsysteme behandelt, die für die Reproduktion von Populationen grundlegende Bedeutung besitzen und den Rahmen für die anschließend dargestellten Paarungssysteme (auch Inzuchtssysteme) schaffen. Auf die Behandlung von Populationen endlicher Größe mit ihren eigenen Gesetzmäßigkeiten folgt eine Darstellung der klassischen Selektionstheorie. Den Schluß des Buches bildet eine kurze Betrachtung aktueller Fragen auf dem Grenzgebiet zwischen Waldbau und Forstgenetik, welche die bestehenden Vorschriften über forstliches Vermehrungsgut nicht ausschließt.

Die von MENDEL konzipierte Methodik der genetischen Analyse wird fast ausschließlich an Beispielen der biochemischen Genetik erläutert. Der Leser wird dadurch unmittelbar an die Grundlagen eines der heute wesentlichen Anwendungsgebiete herangeführt; populationsgenetische Untersuchungen anhand von Isoenzym-Genloci. Die zahlreichen, vom Text abgesetzten Beispiele und Übungsaufgaben sind fast alle dem forstlichen Bereich entnommen, woraus sich auch die vorwiegend angesprochene Zielgruppe erklärt. Im übrigen stellt das Buch eine ausgewogene Verbindung des auch in Lehrbüchern für Biologen vermittelten Stoffs mit der für den Forstmann vorrangigen Populationsgenetik dar.

Anstatt einer Beschreibung aktueller Anwendungsprobleme aus der Forstpflanzenzüchtung oder der Erhaltung forstlicher Genressourcen vermittelt das Buch die heute so dringend erforderliche Einführung in die hierfür benötigten Grundlagen der Genetik und gibt eine kurze Darstellung einiger wesentlicher Ergebnisse forstgenetischer Forschung.

J. D. SAUERLÄNDER'S VERLAG, FRANKFURT AM MAIN