

NEWLY DISCOVERED POPULATIONS OF SALAMANDERS FROM SISKIYOU COUNTY CALIFORNIA REPRESENT A SPECIES DISTINCT FROM *PLETHODON STORMI*

LOUISE S. MEAD^{1,5,6}, DAVID R. CLAYTON², RICHARD S. NAUMAN³,
DEANNA H. OLSON³, AND MICHAEL E. PFRENDER⁴

¹Department of Zoology, 3029 Cordley Hall, Oregon State University, Corvallis, OR 97331, USA

²U.S. Fish and Wildlife Service, Roseburg Field Office, 2900 NW Stewart Parkway, Roseburg, OR 97470, USA

³USDA Forest Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis, OR 97331, USA

⁴Department of Biology, Utah State University, 5305 Old Main Hill Road, Logan, UT 84322, USA

ABSTRACT: *Plethodon stormi* and *Plethodon elongatus* are two closely related species of plethodontid salamanders that are restricted to the Klamath Province of northwestern California and southwestern Oregon. Discovery of three localities south of the Klamath River, in the Scott River drainage, not assignable to either *P. elongatus* or *P. stormi*, motivated closer examination of this complex. We describe molecular (mitochondrial DNA) and morphological variation among specimens collected from the three newly discovered populations and compare these to populations of *P. elongatus* and *P. stormi* from Siskiyou County, California and Jackson and Josephine Counties, Oregon. Analyses of mitochondrial sequence data from the ATPase 6 and cytochrome b genes recovered clades corresponding to *P. elongatus*, *P. stormi* and the Scott River populations. Multivariate analyses indicate that Scott River drainage animals are morphologically distinct from *P. elongatus* and *P. stormi*. Because both genetic and morphological data indicate that the Scott River populations are distinctive, we provide a description of a new species, *Plethodon asupak*, to reflect the evolutionary history of this group and facilitate species management.

Key words: California; Mitochondrial DNA; Morphological variation; New species; Plethodontidae; *Plethodon asupak*; *Plethodon elongatus*; *Plethodon stormi*; Siskiyou County

PLETHODON STORMI and *Plethodon elongatus* are closely related species of plethodontid

salamanders (Highton and Larson, 1979; Mahoney, 2001, 2004) with restricted ranges in northern California and southern Oregon (Petranka, 1998). Species identification, however, continues to be problematic in Siskiyou County, California and the taxonomic status of *P. stormi* is in need of clarification. Some

⁵ PRESENT ADDRESS: Section of Ecology and Evolution, 2320 Storer Hall, University of California, Davis, CA 95616, USA.

⁶ CORRESPONDENCE: e-mail, lmead@ucdavis.edu

authors treat *P. stormi* as a subspecies of *P. elongatus* (Stebbins, 1985, 2003) whereas others treat *P. stormi* as a full species (Brodie, 1970; Highton and Brame, 1965; Nussbaum, 1974; Nussbaum et al., 1983; Petranka, 1998). Variation in body length and pigmentation along an east-west cline from coastal *P. elongatus* to interior *P. stormi* (Bury, 1973, 1999) suggests *Plethodon* along the Klamath River between Seattle Creek and Seiad Valley may be hybrids. These observations are concordant with allozyme data indicating *P. elongatus* and *P. stormi* may be merging upon secondary contact (R. Highton, personal communication), but discordant with a study of mitochondrial DNA variation indicating *P. stormi* extends across this range (Mahoney, 2004). Finally, recently discovered populations of *Plethodon* from near the confluence of the Klamath and Scott Rivers (hereafter referred to as "Scott River" populations) are not readily assignable based on morphology to either *P. elongatus* or *P. stormi* and raise the question of a previously unknown species or a second region of hybridization.

The taxonomic treatment of *P. stormi* from Siskiyou County and the newly discovered Scott River animals depends on the criteria used to define groups and the choice of species concept. Shared characters, whether molecular, biochemical or morphological, can indicate common ancestry, hybridization or convergence. Different views regarding the implications of genetic divergence and complex patterns of gene flow at species boundaries can also lead to different interpretations of available data (Highton, 1998, 2000; Wake and Schneider, 1998). Interpretation of geographic patterns of genetic and phenotypic variation has considerable impact on conservation. Federal agencies manage large portions of the ranges of both *P. elongatus* (65%) and *P. stormi* (84%) (Nauman and Olson, 1999). In addition, the California State Endangered Species Act provides protection for *P. stormi* but not *P. elongatus* on private lands. Because *P. elongatus* has a much larger range than *P. stormi*, species protections are greater for *P. stormi* than *P. elongatus* on federal, state and private lands. The lack of reliable field characters and continuing taxonomic uncertainty of the specific status of populations, particularly in Siskiyou County,

California, has caused difficulties in land management planning. We use mtDNA sequence data and morphological characteristics to clarify the status of the Scott River populations and refine the distribution of *P. stormi* and *P. elongatus*.

MATERIALS AND METHODS

Species and Populations Studied

This study included populations of *P. elongatus* and *P. stormi* from a total of 38 localities (Table 1, Fig. 1) in Siskiyou County, California and Jackson and Josephine Counties, Oregon. In addition, we included samples from three recently discovered populations in the Scott River drainage south and east of the known range of *P. stormi*. We collected tail tips from individuals in the field and preserved tissue in 95% ethanol. Animals used for the morphological study were transported to Oregon State University and euthanized by topical application of a solution of 20% ethyl-P amino benzoate dissolved in polyethylene glycol.

Samples used to assess molecular variation included partial sequences of the mitochondrial gene ATPase 6 for 38 *P. elongatus* representing 17 populations, 43 *P. stormi*, representing 21 populations and 18 samples of a potential new species from the three Scott River populations (Appendix I). This region of the mitochondrial genome has proven useful for fine scale phylogeography (Vitt et al., 1997), and was previously used by Mahoney (2004) to examine broad phylogeographical patterns in *P. elongatus* and *P. stormi*. We also obtained a 370 bp region of the mtDNA cytochrome *b* gene for 36 of these individuals (*P. elongatus*: $n = 16$; *P. stormi*: $n = 14$; Scott River populations: $n = 6$) to estimate pairwise differences and calculate percent sequence divergence within and between the major clades identified in our phylogenetic analysis. We used cytochrome *b* for these estimates to compare divergence estimates to other plethodontids (Jackman et al., 1997; Mahoney, 2004; Mead, 2001; Moritz et al., 1992).

We surveyed morphological variation among populations of *Plethodon elongatus*, *P. stormi*, and the Scott River populations. We collected salamanders from 13 localities (Table 1). We examined a total of 15 adult Scott River

TABLE 1.—Localities for samples used in this study * Indicates sample(s) from this locality includes individuals used in morphological analysis.

Number	Locality	County	State	Latitude	Longitude
1	Rainie Falls	Josephine	OR	42.647	–123.608
2	Graves Creek	Josephine	OR	42.651	–123.583
3	Near Fiddler Gulch	Josephine	OR	42.255	–123.760
4	Powell Creek	Josephine	OR	42.275	–123.80
5	Cave Creek	Josephine	OR	42.109	–123.428
6	Ferris Gulch	Jackson	OR	42.238	–123.218
7	Nine Mile Creek*	Jackson	OR	42.139	–123.163
8	Grouse Creek	Jackson	OR	42.148	–122.998
9	Carberry Creek	Jackson	OR	42.040	–123.181
10	China Gulch	Jackson	OR	42.062	–123.157
11	Gravel Pit	Jackson	OR	42.097	–123.018
12	Yellow Jacket	Jackson	OR	42.017	–122.948
13	East Fork Indian	Siskiyou	CA	41.937	–123.411
14	Thompson Creek	Siskiyou	CA	41.935	–123.349
15	Horse Camp Trail*	Siskiyou	CA	41.972	–123.183
16	Applegate Bridge*	Siskiyou	CA	41.984	–123.176
17	Hutton Guard Station	Siskiyou	CA	42.000	–123.140
18	Joe Creek	Siskiyou	CA	41.995	–123.130
19	Elliot Creek	Siskiyou	CA	41.986	–123.031
20	Seiad Creek	Siskiyou	CA	41.905	–123.141
21	Horse Creek Northern*	Siskiyou	CA	41.900	–123.104
22	Horse Creek Southern*	Siskiyou	CA	41.881	–123.090
23	Seattle Creek	Siskiyou	CA	41.843	–123.301
24	Joe Miles	Siskiyou	CA	41.813	–123.294
25	Evans Mountain*	Siskiyou	CA	41.833	–123.265
26	West Grider	Siskiyou	CA	41.848	–123.231
27	Walker Gulch	Siskiyou	CA	41.827	–123.143
28	Muck-a-Muck*	Siskiyou	CA	41.774	–123.031
29	Mill Creek*	Siskiyou	CA	41.743	–122.958
30	Ottley Gulch*	Siskiyou	CA	41.771	–123.336
31	Grider Creek	Siskiyou	CA	41.787	–123.205
32	Clear Creek*	Siskiyou	CA	41.727	–123.519
33	Elk Creek/Twin Creek*	Siskiyou	CA	41.726	–123.365
34	Seiad Valley (Walker Creek)	Siskiyou	CA	41.750	–123.175
35	Independence River Access	Siskiyou	CA	41.596	–123.397
36	Dillon Creek	Siskiyou	CA	41.573	–123.543
37	Dobbins Creek*	Siskiyou	CA	41.539	–123.538
38	Sandy Bar Creek	Siskiyou	CA	41.487	–123.513
39	Somes Bar (2 mi. NW)	Siskiyou	CA	41.402	–123.515
40	Sawyers Bar	Siskiyou	CA	41.297	–123.064
41	DeadMan East	Siskiyou	CA	41.894	–123.435

drainage animals from two populations, 22 adult *P. elongatus* from four populations, and 19 adult *P. stormi* from seven populations (Appendix I). Thirty-five of these individuals also were used in molecular analyses described above. Six sampled *P. elongatus* and *P. stormi* populations (nos. 24, 25, 30, 32, 33, and 37) occurred across the region reported by Bury (1973) as showing clinal variation. Voucher specimens were deposited at the University of Michigan Museum of Zoology. In addition, two juveniles (one from population 28 and one from population 29) were examined for the type series description.

DNA Extraction, Amplification,
Sequencing and Analysis

We extracted total genomic DNA from frozen and ethanol preserved tissue following a standard proK digestion and phenol/chloroform/isoamyl alcohol extraction (Hillis et al., 1996). The polymerase chain reaction was used to amplify a 667 base-pair mitochondrial DNA segment containing a portion of the ATPase 6 gene using primers L9252 (5'-AACCTGACCATGAACCTAAGCT-3') and H9923 (5'-TAGGAGTGTGCTTGGTGTGC-CAT-3') (Vitt et al., 1997). Amplifications were




FIG. 1.—Map showing sampled populations included in analyses. The shade and shape of markers for populations in all figures correspond to the clades resolved in our phylogenetic analyses as follows: filled circles: clade I; filled triangles: clade II; black stars: clade III; unfilled squares: clade IV.

performed in 50 μ l reactions using 1 μ l of genomic DNA, 5 μ l 10 $\times$ PCR buffer, 5 μ l 8mM dNTP mix, 5 μ l of each 2 μ M primer, 3 μ l of 25 mM $MgCl_2$, 0.2 μ l Taq DNA polymerase (Life Technologies, Carlsbad, CA) and water to final volume. Reactions consisted of an initial denaturation step followed by 40 cycles of 92 C for 45 s, 48 C for 45 s, and 72 C for 90 s. QIAquick PCR purification kit (Qiagen, Valencia, CA) was used to purify the PCR product. The DNA sequence of purified PCR products was obtained using an Applied Biosystems Inc. (Foster City, CA) ABI 377 automated DNA sequencer using Big-Dye Terminator Cycle Sequencing. PCR fragments were sequenced in both directions to check accuracy. The same protocol was used to obtain a 370 bp region of the cytochrome *b* gene using primers MVZ15 (5'-GAAGTAATGGCCACACWWTACG-3') and CYTB2 (5'-CCCCTCAGAATGATATTGTCCTCA-3') (Moritz et al., 1992) and the following PCR reaction protocol: an initial denaturation step followed by 35 cycles of 96 C for 30 s, 54 C for 60 s, and 72 C for 90 s.

Mitochondrial DNA sequences (Appendix I) were aligned by eye using the programs SEQED (Applied Biosystems Inc., Foster City, CA) and a Clustal W algorithm implemented in MegAlign (DNASTAR Inc., Madison, WI). We analyzed ATPase 6 and cytochrome *b* gene sequences separately. Sequence variation was calculated as the percent of variable sites in pair-wise comparisons of unique haplotypes (genetic variants) for both ATPase 6 and cytochrome *b* sequences.

We examined ATPase 6 variation using maximum parsimony (MP) and maximum likelihood (ML) implemented in the computer program PAUP* version 4.0b-10 (Swofford, 1999) and Bayesian estimation (MB) performed with MrBayes version 3 (Huelsenbeck and Ronquist, 2001). Because ML analyses are computationally time consuming with large datasets, we used a subset of haplotypes for the ML analysis only, randomly subsampling a total of 31 individuals from among the unique haplotypes in each major clade (I, II, III, and IV). A model of nucleotide substitution was selected by the hierarchical likelihood ratio tests implemented by Modeltest version 3.0 (Posada and Crandall, 1998). This analysis selected the GTR+I+ Γ substitution model

(Yang, 1994) with empirical base frequencies, 6 rate parameters ([A-C] = 0.86187, [A-G] = 4.09176, [A-T] = 0.56107, [C-G] = 0.46517, [C-T] = 6.79798, [G-T] = 1.0), proportion of invariant sites equal to 0.3323, and a gamma distribution shape parameter equal to 0.8874. For the ML and MP analyses heuristic searches were conducted with 10 random additions and branch swapping by tree bisection rearrangement. Support was assessed using nonparametric bootstrap with 100 pseudoreplicates under the heuristic method. We used the mtDNA sequences of *P. dunni* and *P. vandykei* as outgroups based on previous analysis of molecular variation within the genus *Plethodon* (Mahoney, 2001).

Bayesian analyses were conducted with a random starting tree, run 5.0×10^6 generations and sampled every 100 generations. Base frequencies and initial starting values for specific nucleotide model parameters (including base frequencies, rate matrix, gamma shape distribution, and proportion of invariant sites) were estimated from the data assuming a general time reversible model + I + Γ . We determined stationarity of the Markov chain as the point when sampled log likelihood values plotted against generation time reached a stable mean equilibrium value. Data sampled from generations preceding this point were discarded (Huelsenbeck and Ronquist, 2001). We estimated nodal posterior probabilities, mean log likelihood scores, and a summary phylogeny using all data collected at stationarity.

Morphological Analysis

Morphological measurements were made to the nearest 0.1 mm using digital calipers. The following measurements were taken from each specimen: standard length from the snout to the posterior margin of the vent (SVL); head length from anterior end of the snout to posterior end of gular fold (HL); head width between angle of jaws (HW); interocular distance, measured between medial margins of eyelids (IO); internarial distance (IN), measured between the medial margins of the nares; eye-naris distance, measured between the medial and anterior margin of the eye to the medial and posterior margin of the nares (EN); forelimb length, measured from the anterior point of contact between the forelimb and body to the tip of the third digit (FLB);

hindlimb length, measured from the anterior point of contact between the hindlimb and the body to the tip of the third digit (HLB). We also recorded the number of vomerine teeth (VM) and the number of premaxillary and maxillary teeth (PMM) using a dissecting microscope, and the number of costal grooves (COS) and intercostal folds between adpressed limbs (INCOS). We determined sex of specimens by direct examination of the gonads.

Statistical analyses were performed using SAS version 8e (SAS Institute, Cary, NC). Only sexually mature specimens were used for statistical analyses. To determine if variances were the same within each of the groups defined by the independent variable (SPECIES), we used Levene's tests for homogeneity of variances between traits among species. We assigned each individual to a species according to phylogenetic results (*P. stormi* = clade I and II; *P. elongatus* = clade IV; Scott River = clade III). We also performed a one-way ANOVA to examine the total variation in response (dependent) variables (SVL, HL, HW, IO, IN, EN, FLB, HLB, VM, PMM, COS, INCOS) due to the effects of the classification (independent) variable (SPECIES). Additionally, a variable was chosen (SEX), which separates the data set into groups of observations.

We used univariate Analysis of Covariance (ANCOVA) with SVL as the covariate to test for sexual dimorphism. Each species was analyzed separately. SAS was used to perform an ANOVA with HL, HW, IO, IN, EN, FLB, HLB, PMM, VM, COS, and INCOS as dependent variables, SEX as the classification variable and SVL as the covariate. SVL was used as a covariate to adjust for any effects of size on the dependent variables, thereby reducing the variance of the error term. An interaction between SEX and SVL was also specified to account for variation in the dependent variable not explained by only main effects and covariates.

In all multivariate analyses, we analyzed males and females separately. We performed a Canonical Discriminant Analysis (CDA) on z-transformed data. The z-transformation generates scores that equalize differences in magnitudes among variables by transforming the raw score to a value that corresponds to the deviation from its distribution's mean,

expressed in units of its standard deviation. CDA is a type of discriminant function analysis used to discriminate between two or more naturally occurring groups by deriving functions that optimally combine variables such that the first function provides the most overall discrimination among groups, the second provides second most, and so on. Computation of two canonical variables was specified in the CDA based on linear functions using Mahalanobis distances to compute squared distances between class means based on pooled within-class covariance matrices. We also examined how well we could predict group membership using classification functions. Classification criteria were based on the pooled covariance matrix (yielding a linear function) and accounted for prior probabilities of group membership. We set prior probabilities proportional to the sample sizes.

RESULTS

Our survey revealed 51 unique haplotypes. Identical haplotypes were found in samples from nine populations and three haplotypes were shared across populations. Of the 285 variable sites in our alignment, 176 were parsimony informative. Thirty-six individuals from 33 populations were sequenced for the cytochrome *b* gene, yielding 31 unique haplotypes.

Phylogenetic Analyses

The equally weighted maximum parsimony analysis using all haplotypes resulted in one most parsimonious tree of 435 steps (Fig. 2A). Phylogenetic reconstruction of the subset of unique haplotypes using maximum likelihood resulted in a single ML tree with a likelihood score of -3401.149 (Fig. 2B). The tree resulting from the Bayesian analysis also exhibited similar topological structure to the MP tree in Fig. 2A. The Markov chain appeared to reach stationarity by the 1×10^5 generation. We used a conservative burn-in 1×10^6 generations, resulting in 4.0×10^6 generations considered for estimations of mean likelihood score (-4901.37). Posterior probabilities were comparable to bootstrap support values for most nodes (Fig. 2A).

All our analyses resolved four monophyletic groups. One cluster of haplotypes, correspond-




FIG. 2.—A. Single tree retained from maximum parsimony analysis for ATPase 6 sequences. Tree was rooted with *P. vandykei* and *P. dunni*. Bootstrap values are given above each branch. Bayesian posterior probabilities are given below each branch. Four large haplotype clades are designated by roman numerals. Each haplotype includes the population number from which it was sampled followed by individual identification number. B. Strict bootstrap consensus tree for maximum likelihood analysis.

ing to clade I (Fig. 2), comprised eight unique haplotypes distributed from the Applegate River drainage in Oregon to a region in Siskiyou County, California, just south of the Applegate River drainage and north of the

Klamath River (Fig. 1). Clade I received 75–76% bootstrap support upon resampling of data. This cluster of haplotypes was most closely related to the monophyletic group of 12 unique haplotypes (clade II, Fig. 2) found

TABLE 2.—Range and average of percent sequence divergence at mtDNA cytochrome *b* gene for within (on diagonal) and between (above diagonal) clade comparisons and ATPase 6 gene between clades (below diagonal). Sample sizes in top row are for cytochrome *b* and in first column are for ATPase 6.

	Clade I <i>P. stormi</i> <i>n</i> = 5	Clade II <i>P. stormi</i> <i>n</i> = 9	Clade III Scott River <i>n</i> = 6	Clade IV <i>P. elongatus</i> <i>n</i> = 16
Clade I <i>P. stormi</i> <i>n</i> = 25	0.0–0.73 (0.317)	1.83–2.93 (2.22)	10.9–11.7 (11.5)	8.06–9.89 (8.62)
Clade II <i>P. stormi</i> <i>n</i> = 18	0.225–4.5 (3.19)	0.0–2.56 (1.01)	10.6–13.2 (11.68)	6.59–9.15 (7.59)
Clade III Scott River <i>n</i> = 18	11.7–13.9 (12.7)	11.9–13.5 (12.9)	0.0–1.09 (0.58)	11.35–14.28 (12.85)
Clade IV <i>P. elongatus</i> <i>n</i> = 38	7.7–11.7 (9.5)	8.4–12.5 (10.4)	11.24–13.19 (13.0)	0.15–6.49 (3.34)

in seven populations in northern California between Indian Creek and Seiad Creek on the north side of the Klamath River and between Elk Creek and Grider Creek on the south side of the Klamath River (Fig. 1). Bootstrap support for this clade ranged from 93–95%. Clades I and II formed a monophyletic group which received 96–98% bootstrap support in all analyses (Fig. 2) and appeared as the sister group to *P. elongatus* (clade IV, Fig. 2).

Samples from the Scott River populations appeared as a basal monophyletic group (clade III, Fig. 2) of six unique haplotypes (representing 18 sampled individuals). Bootstrap support for clade III was strong, ranging from 93–100% in all analyses (Fig. 2). Placement of this clade as the sister group to a clade containing *P. stormi* and *P. elongatus* also received strong support (92–100%) in MP and ML analyses. Posterior probability of this node in Bayesian analysis was 0.56 (Fig. 2).

Genetic Diversity

The major clades obtained in our analysis showed variable levels of within-clade divergence, with average percent sequence divergence in cytochrome *b* ranging from 0.3% to 3.34% (Table 2). Between-clade estimates of average percent sequence ranged from 2.22 between *P. stormi* clades I and II to 12.85 between clade III (Scott River) and clade IV (*P. elongatus*). These values were similar to estimates of divergence obtained using ATPase 6 sequences (Table 2). Between-clade estimates of average percent sequence for ATPase 6 ranged from 3.19 between clade I and II (*P. stormi*) to 13.0 between clade III (Scott River) and clade IV (*P. elongatus*) (Table 2).

Morphological Analysis

Morphometric measurements and tooth counts appear in Table 3. The ranges of most

morphological measurements show overlap among species. We found no significant heterogeneity in variances among groups (Appendix II). The one-way ANOVA indicated differences among species in SVL, HL, HW, IO, IN, FLB, HLB, COS, INCOS for males and SVL, HL, HW, IO, EN, IN, FLB, HLB, COS, and INCOS for females (Appendix II). Because many morphological measurements vary with body size and sex, we also used analysis of covariance with snout-vent length (SVL) as the covariate to test for sexual dimorphism within species. For characters that differed in intercept but not slope we also performed a test of the least-square adjusted means (Appendix III). Most characters did show differences between the sexes. However, once differences attributable to SVL were accounted for, within *P. elongatus*, there was significant sexual dimorphism for HW and VM. *Plethodon stormi* exhibited significant sexual dimorphism for INCOS. Among specimens collected from Scott River only PMM and INCOS varied sexually (Appendix III).

Canonical discriminant analysis for males identified SVL, HW, and FLB as contributing most to discrimination among species along the first canonical axis (Table 4), which accounted for 96.7% of the total dispersion ($F = 4.99$, $P = 0.0058$). The second canonical axis produced loadings that were much more uniform (Table 4) although HLB and INCOS provided some discrimination among species. For females, the first variable explained 97.7% of the dispersion among species with SVL, IN, and FLB contributing most to CAN1 ($F = 4.94$, $P = 0.002$). Along the second canonical axis HLB and SVL contributed to variation among species (Table 4). Ordination of individual specimens on the first two canonical axes indicates good separation based on individual canonical scores

TABLE 3.—Comparison of morphometric variation and number of teeth in adults of *P. elongatus*, *P. stormi*, and Scott River populations. Mean $\pm$ SE (above) and range (below); all measurements in mm. Modal numbers are given for costal grooves and intercostals folds.

Variable	<i>P. elongatus</i>		<i>P. stormi</i>		Scott River	
	Males n = 9	Females n = 13	Males n = 6	Females n = 13	Males n = 7	Females n = 8
Snout-vent length	50.2 $\pm$ 3.42 (30.6–63.0)	47.7 $\pm$ 3.7 (27.5–64.9)	62.9 $\pm$ 2.2 (53.5–67.9)	58.2 $\pm$ 2.2 (48.6–72.3)	60.7 $\pm$ 3.0 (48.4–71.3)	67.2 $\pm$ 2.0 (54.9–72.0)
Tail length	41.4 $\pm$ 3.82 (25.9–62.0)	49.7 $\pm$ 5.1 (27.5–65.0)	54.0 $\pm$ 3.7 (46.3–59.3)	52.9 $\pm$ 3.0 (37.0–48.0)	50.6 $\pm$ 2.3 (34.9–58.9)	57.8 $\pm$ 2.0 (48.8–51.5)
Head length	12.1 $\pm$ 0.72 (7.5–14.4)	11.3 $\pm$ 0.7 (7.8–13.8)	14.5 $\pm$ 0.4 (12.4–15.3)	13.7 $\pm$ 0.5 (11.3–15.8)	14.4 $\pm$ 0.8 (11.7–17.4)	15.3 $\pm$ 0.8 (11.4–17.9)
Head width	7.1 $\pm$ 0.42 (4.7–8.7)	6.5 $\pm$ 0.3 (4.9–8.0)	8.7 $\pm$ 0.3 (7.4–9.9)	8.1 $\pm$ 0.3 (6.8–10.0)	8.9 $\pm$ 0.4 (7.3–10.3)	9.6 $\pm$ 0.4 (8.0–10.9)
Interocular distance	2.2 $\pm$ 0.1 (1.6–2.7)	1.9 $\pm$ 0.1 (1.2–2.6)	2.6 $\pm$ 0.1 (2.3–2.9)	2.4 $\pm$ 0.1 (1.6–3.1)	2.9 $\pm$ 0.1 (2.4–3.2)	2.8 $\pm$ 0.1 (2.4–3.3)
Internarial distance	2.0 $\pm$ 0.1 (1.3–2.6)	1.9 $\pm$ 0.1 (1.3–2.6)	2.6 $\pm$ 0.1 (1.9–2.9)	2.4 $\pm$ 0.1 (1.9–2.7)	2.9 $\pm$ 0.1 (2.6–3.5)	2.9 $\pm$ 0.1 (2.7–3.2)
Eye-nostril distance	1.5 $\pm$ 0.1 (0.9–2.1)	1.3 $\pm$ 0.1 (0.9–2.6)	1.9 $\pm$ 0.1 (1.4–2.2)	1.8 $\pm$ 0.1 (1.4–2.0)	1.8 $\pm$ 0.1 (1.3–2.2)	2.0 $\pm$ 0.1 (1.7–2.6)
Forelimb length	8.8 $\pm$ 0.5 (5.7–10.8)	8.6 $\pm$ 0.5 (6.0–11.2)	11.5 $\pm$ 0.5 (9.7–12.5)	10.9 $\pm$ 0.3 (9.7–13.3)	12.5 $\pm$ 0.4 (10.7–14.3)	12.9 $\pm$ 0.2 (11.6–13.7)
Hindlimb length	10.4 $\pm$ 0.7 (6.2–13.2)	10.0 $\pm$ 0.6 (5.9–12.9)	13.8 $\pm$ 0.4 (12.3–14.8)	13.1 $\pm$ 0.3 (11.6–15.1)	14.3 $\pm$ 0.6 (11.7–16.0)	15.5 $\pm$ 0.4 (13.1–16.4)
Maxillary-premaxillary teeth	46.7 $\pm$ 1.8 (40–55)	50.8 $\pm$ 1.2 (45–56)	43.5 $\pm$ 3.1 (31–52)	54.8 $\pm$ 1.9 (45–67)	50.3 $\pm$ 2.9 (44–65)	59.8 $\pm$ 2.7 (42–67)
Vomerine teeth	14.1 $\pm$ 0.9 (9–18)	11.6 $\pm$ 0.5 (9–14)	12.2 $\pm$ 0.7 (10–14)	13.2 $\pm$ 0.6 (9–17)	13.2 $\pm$ 0.7 (10–15)	13.4 $\pm$ 0.7 (11–16)
Costal grooves	18 (17–18)	18 (17–19)	17 (17–18)	17 (16–18)	16 (16–17)	17 (16–17)
Intercostal folds	6 (5–7)	6 (4–7)	5 (5–6)	5 (4–6)	3 (2–3)	3 (3–5)

(Fig. 3). Males from all three groups clearly separated along CAN1 (Fig. 3, top). Fore- and hindlimb length, head width, and number of intercostal folds between adpressed limbs provide good diagnostic differences between groups. Females also showed good separation on the first canonical axis (Fig. 3, bottom), indicating measurements of fore- and hindlimb length, internarial distance and in-

tercostal folds are important diagnostic differences between groups.

Linear discriminant functions generated for the classification analysis indicated SVL, HW, FLB, and HLB were most important in discriminating all three groups for males. Additionally, HL discriminated *P. stormi* and INCOS and EN discriminated individuals from Scott River. For females, IN and INCOS best discriminated all three groups. Snout-vent length (SVL) was important in discriminating both *P. stormi* and Scott River populations. Examination of posterior probability of group membership indicated the functions succeeded in differentiating the three groups. Classification of males and females was entirely accurate.

TABLE 4.—Canonical variate coefficients for the Canonical Discriminant Analysis of adult males and females.

Variable	Males		Females	
	CAN 1	CAN 2	CAN 1	CAN 2
Snout-vent length	-7.825	-2.412	2.667	-4.798
Head length	1.835	-0.797	0.348	-0.789
Head width	4.529	-0.567	0.116	1.658
Interocular distance	0.629	-1.661	-0.321	-0.003
Internarial distance	-0.235	0.352	3.182	-1.933
Eye-nostril distance	-2.238	1.125	-0.133	-0.282
Forelimb length	5.047	2.108	-1.187	1.361
Hindlimb length	1.215	3.572	-1.009	4.189
Vomerine teeth	0.656	0.203	0.421	-0.121
Premaxillary-maxillary	-0.214	1.266	0.914	-0.412
Costal grooves	-0.651	-0.752	-0.229	-0.347
Intercostal folds	-2.202	2.524	-3.859	1.375

DISCUSSION

Molecular and morphological variation support recognition of three distinct groups: *P. elongatus*, *P. stormi*, and the Scott River populations. Molecular variation in mitochondrial DNA detect four distinct clades distributed across Jackson, Josephine and Siskiyou counties. Clades I and II correspond to *P.*




FIG. 3.—Projections of individuals on the two canonical variate axes derived from morphological measurements on adults. *Plethodon elongatus*: open squares; *P. stormi*: filled circles (clade I) and triangles (clade II); Scott River: filled stars.

stormi, clade IV to *P. elongatus*, and clade III to Scott River populations. These results are consistent with those of Mahoney (2004). *Plethodon elongatus* and *P. stormi* are differentiated from one another by 7–10% divergence at both ATPase 6 and cytochrome *b*. Similar estimates were obtained from ATPase 6, cytochrome *b*, and ND4 by Mahoney (2004). Furthermore, populations from Scott River are genetically distinct, exhibiting 10 to

14% divergence (based on uncorrected percent difference at the cytochrome *b* mitochondrial gene) from both *P. stormi* and *P. elongatus*. Levels of divergence between Scott River animals and *P. elongatus* and *P. stormi* are higher than the divergence between *P. elongatus* and *P. stormi* and comparable to divergence between reproductively isolated members of the *Ensatina eschscholtzii* complex (Moritz et al., 1992), morphologically

distinct species of desmognathine salamanders (Mead, 2001), and genera within both the Bolitoglossini and Plethodontini tribes (Jackman et al., 1997). Morphological analyses further reinforce genetic data and support recognizing three distinct groups distributed throughout Siskiyou County, California. Groups show significant morphological variation in a number of characters, with *P. elongatus* and *P. stormi* appearing more similar to one another than either is to Scott River populations. Canonical discriminant analysis provides good discrimination between *P. elongatus*, *P. stormi*, and Scott River populations (Fig. 3). Furthermore, morphological characters traditionally used to distinguish *P. elongatus* and *P. stormi* can also be used to identify Scott River individuals (see below).

Phylogeography

Our analyses also reveal a major phylogeographic subdivision within *P. stormi*. One clade is distributed throughout the Applegate River valley in southern Oregon and extends south into northern Siskiyou County, California. A second clade (*P. stormi* clade II) is distributed across part of Siskiyou County, California north and south of the Klamath River from east of Indian Creek to west of Seiad Creek. The range of *P. stormi* clade II corresponds to the area where there has been disagreement in field identification of *Plethodon*. Historically, individuals found in Siskiyou County, California between Indian Creek and Seiad Valley were referred to *P. elongatus* (e.g., specimens from Seattle Creek, Siskiyou County, California Museum of Vertebrate Zoology nos. 181501–12, collected 23 April 1983), although Brodie (1970) and Bury (1973) reported individuals sampled from this region exhibited characteristics intermediate between *P. stormi* and *P. elongatus*. Our mitochondrial DNA phylogeny indicates Seattle Creek (population 23) as well as all other samples east of Indian Creek (populations 13 and 14) and nearby populations north and immediately south of the Klamath River (populations 23, 24, 25 and 26) are *P. stormi* (clade II). All phylogenetic analyses support clade II as most similar and most closely related to *P. stormi* clade I. Multivariate analyses of morphological data also show clade II males cluster as *P. stormi* (Fig. 3). Female *P. stormi* from clade II

are more intermediate morphologically between *P. stormi* I and *P. elongatus* and may explain why some have interpreted the clinal variation in morphological characters along the Klamath River to represent a broad region of hybridization.

It is difficult to decipher the historical events that have led to the observed pattern of divergence since the geological history of the Klamath-Siskiyou region is poorly understood (Atwater, 1970; Earnst, 1981). However, high levels of population subdivision and Pliocene level divergences have been found in other vertebrate taxa from this region (Loiron et al., 2000; Pfrender et al., 2004). The existence of two divergent *P. stormi* clades and the finding of no mixing of *P. elongatus* and *P. stormi* haplotypes in populations distributed in the region of Seattle Creek strongly supports the scenario that *P. stormi* II diverged from *P. stormi* clade I and represents an isolated group of populations that are intermediate in color and morphology between *P. elongatus* and *P. stormi*, rather than a broad region of hybridization. We focused on populations of *Plethodon* in Siskiyou County, California and Jackson and Josephine Counties, Oregon, however, and did not include known localities of *P. elongatus* further south, in Humboldt County, California or further north in Coos and Curry Counties, Oregon. Current understanding of the molecular phylogenetics of *P. elongatus* and *P. stormi*, inclusive of all *P. elongatus*, are that *P. stormi* and *P. elongatus* form two distinct, monophyletic groups (Mahoney, 2004).

Molecular data also confirm a zone of parapatry between *P. elongatus* and *P. stormi*. In Siskiyou County, California *P. stormi* clade II haplotypes occur just east of Indian Creek (populations 13 and 14) and a haplotype clustering within *P. elongatus* occurs just west of Indian Creek (population 41). The distribution of haplotypes reinforces the possibility of an extremely narrow region of secondary contact across Indian Creek (Mahoney, 2004). Plethodontid salamanders are known to exhibit complex patterns of gene flow across species boundaries (e.g., Mead and Tilley, 2000; Mead et al., 2001; Tilley and Mahoney, 1996; Wake, 1997; Wake and Schneider, 1998; Wake and Jockusch, 2000) and molecular variation at mitochondrial genes should be

interpreted cautiously. Examination of the contact zone between *P. elongatus* and *P. stormi* using nuclear molecular markers is currently in progress (DeGross et al., 2002, 2004).

Scott River

Potential gene flow between *P. elongatus* and *P. stormi* does not influence our conclusions about the unique nature of the Scott River animals. Phylogenetic analyses reinforce that the group of populations along the Scott River are an independent lineage, distinct from *P. elongatus* and *P. stormi*. As noted earlier, costal grooves and intercostals folds between adpressed limbs differentiate *P. elongatus* and *P. stormi* (Highton and Brame, 1965). Scott River animals also differ from *P. elongatus* by having a distinct number of costal grooves and from both *P. elongatus* and *P. stormi* by having one less intercostal fold between adpressed limbs. Studies indicate number of costal grooves correlates with the number of vertebrae (Highton, 1957) and therefore Scott River animals may also differ from *P. elongatus* in number of vertebrae. Relative limb lengths (i.e., FLB/SVL) also indicated animals from Scott River are more robust, with longer limbs and corresponding fewer intercostal folds between adpressed limbs. Small sample sizes for both *P. stormi* and Scott River animals could exaggerate differences. However, variation within *P. elongatus* indicates larger sample sizes would not significantly affect measurable differences between Scott River animals and *P. stormi* or *P. elongatus*, assuming levels of variation are comparable in each group.

It is clear from molecular and morphological data that individuals sampled from Scott River represent a unique, basal lineage with a high degree of divergence from both *P. stormi* and *P. elongatus*. We believe the level of morphological and molecular divergence and basal position of this group suggests an evolutionarily distinct lineage that warrants specific status. Our data on genetic divergence indicate these lineages show sufficient divergence so that they no longer form a cohesive, interbreeding group. As currently known, this new group may have the most restricted range of any species of western *Plethodon* salamander.

Conservation

Our assessment of population structure and the degree of genetic cohesion between lineages has important implications for biodiversity conservation. The rarity, restricted distributions, and ecological sensitivity to land management practices and natural disturbances place these species at risk. We have confirmed that *P. elongatus* and *P. stormi* are distinct evolutionary lineages, potentially having been on different evolutionary trajectories. We have also detected a new, unique basal lineage corresponding to the Scott River populations. Our phylogeographic analyses suggest a long history of fragmentation and restricted gene flow among these three lineages.

Taxonomy

The discovery of cryptic species has become a recurring pattern in phylogenetic studies across numerous taxonomic groups (e.g., Dawood et al., 2002; Masta et al., 2002; Mayer and von Helversen, 2001; Wilke and Pfenninger, 2002). Given concordance between morphology and molecules, there are two alternatives for treatment of the Scott River populations. First, all three groups can be collapsed into a single species, *P. elongatus*. This option would create a single species exhibiting extensive genetic and morphological variation, with more variation across its range than many other plethodontid salamanders (Chippendale et al., 2000; Mead et al., 2001; Moritz et al., 1992). Alternatively, the Scott River populations can be described as a new species, concordant with the existence of distinct mtDNA clades that correspond to geographically defined groups of populations, suggesting a history of independent evolution. We choose to recognize groups that are clearly on independent evolutionary trajectories. We therefore treat *Plethodon* from Scott River as a new species as we believe this option best represents the evolutionary history of the group, recognizes the unique nature of the Scott River animals, and provides the best concordance between taxonomy and our current understanding of species boundaries.

Plethodon asupak, sp. nov.

Holotype.—UMMZ (University of Michigan Museum of Zoology) 262026, MEP field tag

0040, an adult female 70.7 mm SVL (snout to posterior margin of vent), collected at Muck-a-Muck Creek (41.774 N, 123.031 W) above Scott Bar, Siskiyou County California at the confluence of the Scott and Klamath Rivers on April 2 2001 by Louise S. Mead, Richard S. Nauman, Doug DeGross, and Alyssa Zastopil.

Paratypes.—UMMZ 232002 and 232003, MEP field tags 0038 and 0045, same location as holotype; UMMZ 232004 and 232005, field tag MEP 0070 and MEP 0072, collected from Mill Creek, Siskiyou County, California (41.743 N, 122.958 W).

Diagnosis.—*Plethodon asupak* is a medium size salamander of the western *Plethodon* group that closely resembles *P. elongatus* and *P. stormi* but typically is more robust, with a wider head and longer limbs. *Plethodon asupak* is distinguished from *P. elongatus* by having a modal number of 17 costal grooves (*P. elongatus* = 18) and from *P. elongatus* and *P. stormi* by 2.5–3.5 intercostal folds between adpressed limbs (*P. elongatus* = 5–6; *P. stormi* = 4–5). The body of *P. asupak* is elongate with the tail slightly more than 80% of snout to vent length (mean TL/SVL for males: 0.83; females: 0.85) compared to 85–90% for *P. elongatus* and *P. stormi* (mean TL/SVL for *P. elongatus* males: 0.87; females: 0.91; and for *P. stormi* males: 0.90; females: 0.88). *Plethodon asupak* has relatively long fore- and hindlimbs. Males exhibit larger limb/SVL ratios compared to these measures in both *P. elongatus* and *P. stormi*: Ranges of FLB/SVL for males—*P. asupak*: 0.19–0.22; *P. elongatus*: 0.15–0.18; *P. stormi*: 0.16–0.19. Ranges of HLB/SVL for males—*P. asupak*: 0.22–0.25; *P. elongatus*: 0.17–0.22; *P. stormi*: 0.19–0.22.

Description of holotype.—An adult female (UMMZ 232026) with enlarged ova measuring 2–3 mm. Measurements before preservation (in mm): SVL, 57; head length (snout to posterior end of gular fold), 17.9; head width between angle of jaws, 10.2; interorbital distance, 2.4; internarial distance, 2.8; eye to naris distance, 2.2; forelimb length (measured from the anterior point of contact between the forelimb and body to the tip of the third digit), 12.5; hindlimb length (measured from the anterior point of contact between the hindlimb and the body to the tip of the third digit), 15.9; tail length, 62.8 (no evidence of regeneration). There are 17 costal grooves and 3 intercostal

folds between adpressed limbs. There are a total of 60 premaxillary and maxillary teeth and a total of 15 vomerine teeth.

Coloration.—Color in life: Lateral portions of the body are chocolate brown, composed of brown and black pigmentation. The dorsal region of the head and body are distinguished from the lateral surfaces by brown and bronze pigmentation. The coloring of this dorsal stripe extends from the head to the tip of the tail. White and yellow flecks cover most parts of the body but are concentrated on the sides and limbs. The venter is mottled, with light gray patches distributed across a dark gray to purplish background. White flecking is also present on venter, particularly in gular region. The degree of mottling varies. The chin is gray with some mottling. Eyes are black with gold flecking on upper and lower surface (degree of gold varies).

Juveniles have two orange to reddish-brown stripes that extend from just posterior of the eyes toward the tail. The two stripes fuse into a single stripe just posterior to the vent region. Concentrated black pigment lines the margins of the dorsal stripes and is replaced by brown pigment along the lateral portions of the body. White to yellow flecks also appear across the entire body, concentrated on the lateral portions of the trunk, the dorsal surface of the head, and the limbs. Venter is dark gray to purple with white flecking.

Variation in the type series.—The type series consists of one adult female (MEP 0040, UMMZ 232026) with enlarged ova ca. 2–3 mm in diameter; a mature male (MEP0045, UMMZ 232003), a mature female with enlarged ova (MEP 0070, UMMZ 232004) and 2 unsexed juveniles (MEP 0038, UMMZ 232002 and MEP 0072, UMMZ 232005) (Appendix IV). All paratypes classified as mature adults resemble the type-specimen in coloration, presence of dorsal stripe, and gold flecking on eye. The degree of white flecks on dorsal and lateral regions of the head and body vary. The degree of mottling on the ventral side also varies among individuals. All mature adults have 17 costal grooves. Both mature females have 3 intercostal folds between adpressed limbs. The adult male has 2 intercostal folds between adpressed limbs. The two juveniles both exhibit the characteristic paired dorsal stripe, varied degrees of white

flecking, 16–17 costal grooves, and 2–3 intercostal folds.

Etymology.—The word Asupak is the Shasta Indian name for Scott Bar, the area near the confluence of the Scott River and the Klamath River, and the type locality for *Plethodon asupak*. The Scott River and Valley area was first visited by various bands the Shasta Indians and eventually inhabited by the Karuk Tribe. The Karuk refer to indicator species in their understanding of nature and viewed salamanders as having the specific function of water purifier and as an omen of good luck (Hillman and Salter, 1997).

Acknowledgments.—S. Cuenca, D. Degross, and H. Welsh provided valuable tissue samples. K. Freed assisted with molecular laboratory work. We especially thank T. Titus for motivating early investigations in this system and providing stimulating discussions on plethodontid salamanders. Kevin de Queiroz and Meredith Mahoney, along with The Evolution and Natural History Group at Utah State University provided useful comments on the manuscript. This work was funded by an USDA Forest Service Pacific Northwest Research Station research grant (PNW 01-CR-11261952-212) to S. J. Arnold and D. H. Olson and U.S. Fish and Wildlife Service grant (1448-13420-01-M105) to SJA.

LITERATURE CITED

- ATWATER, T. 1970. Implications of plate tectonics for the Cenozoic tectonic evolution of western North America. *Bulletin of the Geologic Society of America* 81: 3513–3536.
- BRODIE, E. D., JR. 1970. Western salamanders of the genus *Plethodon*: systematics and geographic variation. *Herpetologica* 26:468–516.
- BURY, R. B. 1973. Western *Plethodon*: systematics and biogeographic relationships of the *elongatus* group. *Hiss News-Journal* 1:56–57.
- . 1999. Evolution, zoogeography and management of the Del Norte and Siskiyou Mountain salamanders. *Northwestern Naturalist* 80:121.
- CHIPPENDALE, P. T., A. H. PRICE, J. J. WIENS, AND D. M. HILLIS. 2000. Phylogenetic relationships and systematic revision of central Texas hemidactyliine plethodontid salamanders. *Herpetological Monographs* 14:1–80.
- DAWOOD, A., A. CHANNING, AND J. P. BOGART. 2002. A molecular phylogeny of the frog genus *Tomopterna* in Southern Africa: examining species boundaries with mitochondrial 12S rRNA sequence data. *Molecular Phylogenetics Evolution* 22:407–413.
- DEGROSS, D., L. S. MEAD, AND S. J. ARNOLD. 2002. Assessing gene flow across contact zones between *Plethodon elongatus* and *Plethodon stormi* using microsatellite markers. *Northwestern Naturalist* 84:97.
- . 2004. Novel tetranucleotide microsatellite markers from the Del Norte salamander (*Plethodon elongatus*) with application to its sister species the Siskiyou Mountain salamander (*P. stormi*). *Molecular Ecology Notes*: in press.
- EARNST, W. G. 1981. *The Geotectonic Development of California*. Prentice Hall, Englewood Cliffs, New Jersey, U.S.A.
- HIGHTON, R. 1957. Correlating costal grooves with trunk vertebrae in salamanders. *Copeia* 1957:107–109.
- . 1998. Is *Ensatina eschscholtzii* a ring-species? *Herpetologica* 54:254–278.
- . 2000. Detecting cryptic species using allozyme data. Pp. 215–241. *In* R. C. Bruce, R. G. Jaeger, and L. D. Houck (Eds.), *The Biology of Plethodontid Salamanders*. Plenum Press, New York, New York, U.S.A.
- HIGHTON, R., AND A. H. BRAME. 1965. *Plethodon stormi* species nov. *Amphibia: Urodela: Plethodontidae*. Pilot Register of Zoology, Card No. 20.
- HIGHTON, R., AND A. LARSON. 1979. The genetic relationships of the salamanders of the genus *Plethodon*. *Systematic Zoology* 28:579–599.
- HILLIS, D. M., B. K. MABLE, A. LARSON, S. K. DAVIS, AND E. A. ZIMMER. 1996. Nucleic acids. IV. Sequencing and cloning. Pp. 321–381. *In* D. M. Hillis, C. Moritz, and B. K. Mable (Eds.), *Molecular Systematics*, 2nd ed. Sinauer Associates, Sunderland, Massachusetts, U.S.A.
- HILMAN, L., AND J. F. SALTER. 1997. Environmental management: American Indian knowledge and the problem of sustainability. *Forests, Trees and People* 34:1–11.
- HUELSENBECK, J., AND F. RONQUIST. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17:754–755.
- JACKMAN, T. R., G. APPLEBAUM, AND D. B. WAKE. 1997. Phylogenetic relationships of bolitoglossine salamanders: a demonstration of the effects of combining morphological and molecular data sets. *Molecular Biology and Evolution* 14:883–891.
- LORION, C. M., D. F. MARKEL, S. REID, AND M. DOCKER. 2000. Redescription of the presumed-extinct Miller Lake lamprey, *Lampetra minima*. *Copeia* 2000:1019–1028.
- MAHONEY, M. J. 2001. Molecular systematics of *Plethodon* and *Aneides* (Caudata: Plethodontidae: Plethodontini): phylogenetic analysis of an old and rapid radiation. *Molecular Phylogenetics and Evolution* 18:174–188.
- . 2004. Molecular systematics and phylogeography of the *Plethodon elongatus* species group: combining phylogenetic and population genetic methods to investigate species history. *Molecular Ecology* 13: 149–166.
- MASTA, S. E., B. K. SULLIVAN, T. LAMB, AND E. J. ROUTMAN. 2002. Molecular systematics, hybridization, and phylogeography of the *Bufo americanus* complex in Eastern North America. *Molecular Phylogenetics and Evolution* 24:302–314.
- MAYER, F., AND O. VON HELVERSEN. 2001. Cryptic diversity in European bats. *Proceedings of the Royal Society of London B Biological Sciences* 268:1825–1832.
- MEAD, L. S. 2001. Patterns and processes of speciation in desmognathine salamanders. Ph.D. Dissertation, University of Massachusetts, Amherst, Massachusetts, U.S.A.
- MEAD, L. S., AND S. G. TILLEY. 2000. Ethological Isolation and variation in allozymes and dorsolateral pattern between parapatric forms in the *Desmognathus ochrophaeus* complex. Pp. 181–198. *In* R. C. Bruce, R. G. Jaeger, and L. D. Houck (Eds.), *The Biology of Plethodontid Salamanders*. Plenum Press, New York, New York, U.S.A.

MEAD, L. S., S. G. TILLEY, AND L. A. KATZ. 2001. Genetic structure of the Blue Ridge Dusky Salamander (*Desmognathus orestes*): inferences from mitochondrial DNA, allozymes, and behavior. *Evolution* 55: 2287–2302.

MORITZ, C., C. J. SCHNEIDER, AND D. B. WAKE. 1992. Evolutionary relationships within the *Ensatina eschscholtzii* complex confirm the ring species interpretation. *Systematic Biology* 41:273–291.

NAUMAN, R. S., AND D. H. OLSON. 1999. Survey and manage salamander known sites. Pp. 43–78. In D. H. Olson (Ed.), *Survey Protocols for Amphibians under the Survey and Manage Provision of the Northwest Forest Plan*. Available at : <http://www.or.blm.gov/surveyandmanage/SP/Amphibians99/protoch.pdf>

NUSSBAUM, R. A. 1974. The distributional ecology and life history of the Siskiyou Mountains salamander, *Plethodon stormi*, in relation to the potential impact of the proposed Applegate Reservoir on this species. U.S. Army Corps of Engineers, Portland Division, Portland, Oregon, U.S.A.

NUSSBAUM, R. A., J. E. D. BRODIE, AND R. M. STORM. 1983. *Amphibians and Reptiles of the Pacific Northwest*. University of Idaho Press, Moscow, Idaho, U.S.A.

PETRANKA, J. W. 1998. *Salamanders of the United States and Canada*. Smithsonian Institution Press, Washington, D.C., U.S.A.

PREFENDER, M. E., J. HICKS, AND M. LYNCH. 2004. Biogeographic patterns and current distribution of molecular-genetic variation among populations of speckled dace, *Rhinichthys osculus* (Girard). *Molecular Phylogenetics and Evolution* 26:125–132.

POSADA, D., AND K. CRANDALL. 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics* 14:2000.

STEBBINS, R. C. 1985. *Western Reptiles and Amphibians*, 2nd ed. Houghton Mifflin Company, Boston, Massachusetts, U.S.A.

———. 2003. *Western Reptiles and Amphibians*, 3rd ed. Houghton Mifflin Company, Boston, Massachusetts, U.S.A.

SWOFFORD, D. L. 1999. *PAUP*: Phylogenetic Analysis Using Parsimony*. Version 4.0beta4b. Sinauer Associates, Sunderland, Massachusetts, U.S.A.

TILLEY, S. G., AND M. J. MAHONEY. 1996. Patterns of genetic differentiation in salamanders of the *Desmognathus ochrophaeus* complex (Amphibia: Plethodontidae). *Herpetological Monographs* 10:1–42.

VITT, L. J., J. P. CALDWELL, P. A. ZANI, AND T. A. TITUS. 1997. The role of habitat shift in the evolution of lizard morphology: evidence from tropical *Tropidurus*. *Proceedings of the National Academy of Sciences. U.S.A.* 94:3828–3832.

WAKE, D. B. 1997. Incipient species formation in salamanders of the *Ensatina* complex. *Proceedings of the National Academy of Sciences of the U.S.A.* 94:7761–7767.

WAKE, D. B., AND C. J. SCHNEIDER. 1998. Taxonomy of the plethodontid salamander genus *Ensatina*. *Herpetologica* 54:279–298.

WAKE, D. B., AND E. L. JOCKUSCH. 2000. Detecting species borders using diverse data sets: examples from plethodontid salamanders in California. Pp. 95–120. In R. C. Bruce, R. G. Jaeger, and L. D. Houck (Eds.), *The Biology of Plethodontid Salamanders*. Plenum Press, New York, New York, U.S.A.

WILKE, T., AND M. PFENNINGER. 2002. Separating historic events from recurrent processes in cryptic species: phylogeography of mud snails (*Hydrobia* spp.). *Molecular Ecology* 11:1439–1451.

YANG, Z. 1994. Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites: approximate methods. *Journal of Molecular Evolution* 39:306–399.

Accepted: 8 November 2004
Associate Editor: Kevin de Queiroz

APPENDIX I Population, sample identification number, GenBank accession numbers, indication as to whether individual used for morphological analysis, and University of Michigan Museum of Zoology catalog number for all specimens examined.

Population number	Sample ID	ATPase 6	Cytochrome b	Morphology	UMMZ Catalog Number	Species	mtDNA Clade	State	County	Locality
1	RAF0245	AY688208	AY688284			<i>Plethodon elongatus</i>	IV	OR	Josephine	Rainie Falls
2	GVC0244	AY688212	AY688279			<i>Plethodon elongatus</i>	IV	OR	Josephine	Graves Creek
3	FID0275	AY688204	AY688275			<i>Plethodon elongatus</i>	IV	OR	Josephine	Near Fiddler Gulch
4	POW0260	AY688209	AY688283			<i>Plethodon elongatus</i>	IV	OR	Josephine	Powell Creek
4	POW0266	AY688210				<i>Plethodon elongatus</i>	IV	OR	Josephine	Powell Creek
5	CAV0270	AY688203	AY688291			<i>Plethodon elongatus</i>	IV	OR	Josephine	Cave Creek
6	FRG0216	AY688196	AY688276			<i>Plethodon stormi</i>	I	OR	Jackson	Ferris Gulch
7	NMC0074	AY688157		x		<i>Plethodon stormi</i>	I	OR	Jackson	Nile Mile Creek
7	NMC0075	AY688158		x		<i>Plethodon stormi</i>	I	OR	Jackson	Nile Mile Creek
7	NMC0076	AY688159				<i>Plethodon stormi</i>	I	OR	Jackson	Nile Mile Creek
7	NMC0077	AY688160				<i>Plethodon stormi</i>	I	OR	Jackson	Nile Mile Creek
7	NMC0223	AY688202				<i>Plethodon stormi</i>	I	OR	Jackson	Nine Mile Creek

APPENDIX I
Continued.

Population number	Sample ID	ATPase 6	Cytochrome b	Morphology	UMMZ Catalog Number	Species	mtDNA Clade	State	County	Locality
8	GCR1239	AY688114				<i>Plethodon stormi</i>	I	OR	Jackson	Grouse Creek
9	CBC0235	AY688194	AY688267			<i>Plethodon stormi</i>	I	OR	Jackson	Carberry Creek
10	CHG0214	AY688195	AY688268			<i>Plethodon stormi</i>	I	OR	Jackson	China Gulch
11	GRA0236	AY688200	AY688277			<i>Plethodon stormi</i>	I	OR	Jackson	Gravel Pit
12	YJK0237	AY688207				<i>Plethodon stormi</i>	I	OR	Jackson	Yellow Jacket
13	EFT0020	AY688123	AY688272			<i>Plethodon stormi</i>	II	CA	Siskiyou	East Fork Indian
14	TCR0019	AY688122	AY688289			<i>Plethodon stormi</i>	II	CA	Siskiyou	Thompson Creek
15	HCT0107	AY688144	AY688280	x	232092	<i>Plethodon stormi</i>	I	CA	Siskiyou	Horse Camp Trail
16	ABR1211	AY688129				<i>Plethodon stormi</i>	I	CA	Siskiyou	Applegate Bridge
16	ABR0108	AY688130		x	232093	<i>Plethodon stormi</i>	I	CA	Siskiyou	Applegate Bridge
16	ABR0109	AY688131			232094	<i>Plethodon stormi</i>	I	CA	Siskiyou	Applegate Bridge
16	ABR0161			x	232099	<i>Plethodon stormi</i>	I	CA	Siskiyou	Applegate Bridge
16	ABR0162			x	232100	<i>Plethodon stormi</i>	I	CA	Siskiyou	Applegate Bridge
16	ABR0163			x	232101	<i>Plethodon stormi</i>	I	CA	Siskiyou	Applegate Bridge
17	HGS8010	AY688115				<i>Plethodon stormi</i>	I	CA	Siskiyou	Hutton Guard Station
18	JCR0224	AY688205	AY688302			<i>Plethodon stormi</i>	I	CA	Siskiyou	Joe Creek
19	ELI0217	AY688169	AY688274			<i>Plethodon stormi</i>	I	CA	Siskiyou	Eliot Creek
19	ELI0233	AY688206				<i>Plethodon stormi</i>	I	CA	Siskiyou	Eliot Creek
20	SDC0078	AY688192	AY688286		232078	<i>Plethodon stormi</i>	I	CA	Siskiyou	Seiad Creek
21	HCR0081	AY688132	AY688293	x	232081	<i>Plethodon stormi</i>	I	CA	Siskiyou	Horse Creek Northern
21	HCR0082	AY688133		x	232082	<i>Plethodon stormi</i>	I	CA	Siskiyou	Horse Creek Northern
21	HCR0083	AY688134		x	232083	<i>Plethodon stormi</i>	I	CA	Siskiyou	Horse Creek Northern
21	HCR0084	AY688135		x	232084	<i>Plethodon stormi</i>	I	CA	Siskiyou	Horse Creek Northern
21	HCR0085			x	232085	<i>Plethodon stormi</i>	I	CA	Siskiyou	Horse Creek Northern
22	HCR0079	AY688136		x	232079	<i>Plethodon stormi</i>	I	CA	Siskiyou	Horse Creek Southern
22	HCR0080			x	232080	<i>Plethodon stormi</i>	I	CA	Siskiyou	Horse Creek Southern
23	STC0001	AY688120				<i>Plethodon stormi</i>	II	CA	Siskiyou	Seattle Creek
23	STC0002	AY688121				<i>Plethodon stormi</i>	II	CA	Siskiyou	Seattle Creek
24	JMC0004	AY688118	AY688297			<i>Plethodon stormi</i>	II	CA	Siskiyou	Joe Miles Creek
24	JMC0005	AY688119				<i>Plethodon stormi</i>	II	CA	Siskiyou	Joe Miles Creek
24	JMC0091	AY688145		x	232087	<i>Plethodon stormi</i>	II	CA	Siskiyou	Joe Miles Creek
24	JMC0092	AY688182		x	232088	<i>Plethodon stormi</i>	II	CA	Siskiyou	Joe Miles Creek
24	JMC0093	AY688183			232089	<i>Plethodon stormi</i>	II	CA	Siskiyou	Joe Miles Creek
24	JMC0094	AY688184			232090	<i>Plethodon stormi</i>	II	CA	Siskiyou	Joe Miles Creek
25	EVM0051	AY688142		x	232077	<i>Plethodon stormi</i>	II	CA	Siskiyou	Evans Mountain
25	EVM0095	AY688143		x	232091	<i>Plethodon stormi</i>	II	CA	Siskiyou	Evans Mountain
25	EVM0119	AY688165		x	232095	<i>Plethodon stormi</i>	II	CA	Siskiyou	Evans Mountain
25	EVM0120	AY688179			232096	<i>Plethodon stormi</i>	II	CA	Siskiyou	Evans Mountain
25	EVM0121	AY688180	AY688301		232097	<i>Plethodon stormi</i>	II	CA	Siskiyou	Evans Mountain
25	EVM0122	AY688181			232098	<i>Plethodon stormi</i>	II	CA	Siskiyou	Evans Mountain
26	WGR0007	AY688116	AY688294			<i>Plethodon stormi</i>	II	CA	Siskiyou	West Grider Ridge
26	WGR0008	AY688117				<i>Plethodon stormi</i>	II	CA	Siskiyou	West Grider Ridge
27	WAG8013	AY688126				Scott River	III	CA	Siskiyou	Walker Gultch
27	WAG0014	AY688127	AY688290			Scott River	III	CA	Siskiyou	Walker Gultch
27	WAG0015	AY688128	AY688292			Scott River	III	CA	Siskiyou	Walker Gultch
28	MMC0012	AY688147	AY688299			Scott River	III	CA	Siskiyou	Muck-A-Muck Creek

APPENDIX I
Continued.

Population number	Sample ID	ATPase 6	Cytochrome b	Morphology	UMMZ Catalog Number	Species	mtDNA Clade	State	County	Locality
28	MMC0013	AY688148				Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0049	AY688150		x	232026	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0050	AY688154		x	232027	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0053			x	232008	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0054			x	232009	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0055			x	232010	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0056			x	232011	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0057				232002	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0058			x	232012	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0059			x	232001	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0060	AY688149			232014	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0062	AY688151			232015	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0063	AY688155	AY688300		232016	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0064	AY688156		x	232003	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0065	AY688152			232017	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0067			x	232019	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0068	AY688153		x	232020	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
28	MMC0072			x	232024	Scott River	III	CA	Siskiyou	Muck-A-Muck Creek
29	SFM0016	AY688124	AY688287			Scott River	III	CA	Siskiyou	Mill Creek
29	MCR0087	AY688185		x	232006	Scott River	III	CA	Siskiyou	Mill Creek
29	MCR0088	AY688201		x	232004	Scott River	III	CA	Siskiyou	Mill Creek
29	MCR0089	AY688146	AY688296	x	232007	Scott River	III	CA	Siskiyou	Mill Creek
29	MCR0090	AY688186			232005	Scott River	III	CA	Siskiyou	Mill Creek
30	OTG0099	AY688161		x	232032	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Ottley Gulch
30	OTG0100	AY688164	AY688282	x	232033	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Ottley Gulch
30	OTG0127	AY688187			232036	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Ottley Gulch
30	OTG0128	AY688188		x	232037	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Ottley Gulch
30	OTG0130	AY688189			232039	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Ottley Gulch
31	GRI0147	AY688190	AY688278			<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Grider Creek
32	CLC0101	AY688137			232034	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Clear Creek
32	CLC0102	AY688138	AY688269	x	232035	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Clear Creek
32	CLC0146	AY688166			232055	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Clear Creek
33	ELC0097	AY688140	AY688273	x	232030	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
33	ELC0098	AY688141		x	232031	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
33	ELC0131	AY688172		x	232040	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
33	ELC0132			x	232041	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
33	ELC0133	AY688173		x	232042	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
33	ELC0134	AY688174		x	232043	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
33	ELC0135	AY688175		x	232044	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
33	ELC0136	AY688176		x	232045	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
33	ELC0137	AY688177		x	232046	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek

APPENDIX I
Continued.

Population number	Sample ID	ATPase 6	Cytochrome b	Morphology	UMMZ Catalog Number	Species	mtDNA Clade	State	County	Locality
33	ELC0138	AY688178		x	232047	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
33	ELC0139			x	232048	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
33	ELC0140			x	232049	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
33	ELC0141			x	232050	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Elk Creek
34	SDV0110	AY688193	AY688298			<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Seiad Valley
35	IRA0022	AY688211	AY688281			<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Independence RA
36	DLC0117	AY688139	AY688270			<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Dillon Creek
37	DOC0052			x	232028	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Dobbins Creek
37	DOC0096	AY688167			232029	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Dobbins Creek
37	DOC0148	AY688168	AY688271	x	232056	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Dobbins Creek
37	DOC0149			x	232057	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Dobbins Creek
37	DOC0150	AY688170		x	232058	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Dobbins Creek
37	DOC0151			x	232059	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Dobbins Creek
37	DOC0152	AY688171			232060	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Dobbins Creek
37	DOC0153	AY688197			232061	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Dobbins Creek
37	DOC0154	AY688198			232062	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Dobbins Creek
37	DOC0155	AY688199			232063	<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Dobbins Creek
38	SBC0115	AY688162				<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Sandy Bar Creek
39	SMB0116	AY688163	AY688288			<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Somes Bar
40	SAW0205	AY688191	AY688285			<i>Plethodon elongatus</i>	IV	CA	Siskiyou	Sawyers Bar
41	DEE0283		AY688295			<i>Plethodon elongatus</i>	IV	CA	Siskiyou	DeadMan East
		AY688125				<i>Plethodon dunni</i>		OR	Douglas	South of Canyonville
		AY688213	AY688303			<i>Plethodon vandykei</i>		WA	Lewis	Willapa Hills

APPENDIX II Tests for differences among species (*Plethodon elongatus*, *P. stormi*, and Scott River populations) and homogeneity of variance.

Variable	ANOVA among species		Levene's test for homogeneity of variance	
	Males	Females	Males	Females
Snout-vent length				
F =	4.99	9.44	1.15	3.18
P =	0.018	0.0006	0.33	0.053
Head length				
F =	4.15	9.2	0.99	0.51
P =	0.03	0.0007	0.39	0.60
Head width				
F =	7.26	18.81	0.6	0.56
P =	0.0045	<0.0001	0.48	0.57
Interocular distance				
F =	10.80	11.15	2.37	1.07
P =	0.0007	0.0002	0.12	0.35
Internarial distance				
F =	11.24	22.74	0.29	1.99
P =	0.0006	<0.0001	0.75	0.15
Eye-nostril distance				
F =	3.19	7.20	1.36	1.39
P =	0.063	0.0027	0.28	0.26
Forelimb length				
F =	16.09	23.01	0.64	2.76
P =	<0.001	<0.0001	0.53	0.78
Hindlimb length				
F =	12.38	22.06	1.53	2.83
P =	0.0004	<0.0001	0.24	0.07

APPENDIX II
Continued.

Variable	ANOVA among species		Levene's test for homogeneity of variance	
	Males	Females	Males	Females
Max-premax. teeth				
<i>F</i> =	1.54	3.23	0.47	0.54
<i>P</i> =	0.2422	0.058	0.63	0.59
Vomerine teeth				
<i>F</i> =	1.39	1.34	1.80	0.37
<i>P</i> =	0.2767	0.2825	0.19	0.69
Costal grooves				
<i>F</i> =	10.45	5.66	0.62	0.99
<i>P</i> =	0.0009	0.008	0.54	0.38
Intercostal folds				
<i>F</i> =	91.81	9.64	0.31	0.63
<i>P</i> =	<0.0001	0.0006	0.73	0.54

APPENDIX III Test for sexual dimorphism using ANCOVA.

Variable	ANOVA between sexes within species with SVL as covariate				Least-square adjusted means For ANCOVA					
					<i>P. elongatus</i>		<i>P. stormi</i>		Scott River	
	<i>P. elongatus</i>	<i>P. stormi</i>	Scott River		M	F	M	F	M	F
Head length				Adj.mean	11.66	11.44	13.65	13.62	14.65	14.91
<i>F</i> =	102.6	67.92	0.65	SE	0.2	0.16	0.29	0.15	0.93	0.9
<i>P</i> =	<0.0001	<0.0001	0.59	<i>t</i> -test	0.39		0.91		0.11	
				Slope	NS		NS		NS	
				Intercept	NS		NS		NS	
									X	
Head width				Adj.mean	6.81	6.52	8.12	8.05	9.23	9.13
<i>F</i> =	197.03	27.44	19.45	SE	0.07	0.06	0.29	0.15	0.19	0.19
<i>P</i> =	<0.0001	<0.0001	0.0001	<i>t</i> -test	0.008		0.83		0.6	
	^Sex × SVL			Slope	0.0005		NS		NS	
				Intercept	X		NS		NS	
Interocular distance				Adj.mean	2.1	1.95	2.52	2.39	2.99	2.69
<i>F</i> =	9.11	1.63	2.03	SE	0.09	0.07	0.23	0.12	0.11	0.11
<i>P</i> =	0.0006	0.2245	0.168	<i>t</i> -test	0.22		0.64		0.09	
				Slope	NS		NS		NS	
				Intercept	NS		NS		NS	
Internarial distance				Adj.mean	1.92	1.97	2.31	2.37	3.02	2.85
<i>F</i> =	19.29	8.31	4.08	SE	0.07	0.06	0.13	0.06	0.09	0.09
<i>P</i> =	<0.0001	0.0017	0.035	<i>t</i> -test	0.58		0.71		0.2	
				Slope	NS		NS		NS	
				Intercept	NS		NS		NS	
Eye–nostril distance				Adj.mean	1.43	1.45	1.67	1.76	1.87	1.96
<i>F</i> =	6.30	23.73	1.00	SE	0.11	0.09	0.07	0.03	0.12	0.12
<i>P</i> =	0.0038	<0.0001	0.43	<i>t</i> -test	0.87		0.28		0.58	
				Slope	NS		NS		NS	
				Intercept	NS		NS		NS	
									X	
Forelimb length				Adj.mean	8.51	8.7	10.63	10.89	12.96	12.67
<i>F</i> =	74.22	23.55	13.06	SE	0.16	0.13	0.36	0.18	0.21	0.2
<i>P</i> =	<0.0001	<0.0001	0.0006	<i>t</i> -test	0.38		0.54		0.33	
				Slope	NS		NS		NS	
				Intercept	NS		NS		NS	
Hindlimb length				Adj.mean	10.02	10.29	13.12	13	14.95	15.02
<i>F</i> =	78.78	39.11	20.86	SE	0.22	0.17	0.33	0.17	0.25	0.24
<i>P</i> =	<0.0001	<0.0001	<0.0001	<i>t</i> -test	0.53		0.77		0.85	
				Slope	NS		NS		NS	
				Intercept	NS		NS		NS	

APPENDIX III
Continued.

ANCOVA between sexes within species with SVL as covariate					Least-square adjusted means For ANCOVA					
					<i>P. elongatus</i>		<i>P. stormi</i>		Scott River	
Variable	<i>P. elongatus</i>	<i>P. stormi</i>	Scott River		M	F	M	F	M	F
Max.-premax.teeth				Adj.mean	46.03	49.9	47.2	54.6	47.21	60.4
<i>F</i> =	2.33	3.61	5.09	SE	1.68	1.67	4.38	2.32	2.72	2.65
<i>P</i> =	0.1357	0.0429	0.019	<i>t</i> -test	0.14		0.16		0.005	
				Slope	NS		NS		NS	
				Intercept	NS		NS		0.023	
							X			
Vomerine teeth				Adj.mean	13.87	11.68	13.07	13.02	13	13.22
<i>F</i> =	2.58	0.89	0.016	SE	0.83	0.83	1.24	0.66	0.9	0.79
<i>P</i> =	0.1118	0.4721	0.9212	<i>t</i> -test	0.092		0.97		0.86	
				Slope	NS		NS		NS	
				Intercept	0.032		NS		NS	
							X		X	
Costal grooves				Adj.mean	17.7	17.7	17.54	17.1	16.53	16.7
<i>F</i> =	0.79	0.93	0.97	SE	0.2	0.16	0.35	0.17	0.22	0.21
<i>P</i> =	0.5155	0.4513	0.44	<i>t</i> -test	0.95		0.3		0.49	
				Slope	NS		NS		NS	
				Intercept	NS		NS		NS	
							X		X	
Intercostal folds				Adj.mean	5.76	5.55	5.48	4.82	2.68	3.5
<i>F</i> =	3.82	11.69	7.92	SE	0.18	0.14	0.33	0.16	0.23	0.22
<i>P</i> =	0.0268	0.0003	0.0043	<i>t</i> -test	0.36		0.09		0.026	
	^ Sex × SVL			Slope	NS		0.028		NS	
				Intercept	NS		X		0.026	

^ Sex × SVL indicates there is no significant linear model (no significant relationship with SVL).

x Indicates a significant difference in slope and no comparison is made.

APPENDIX IV Morphological measurements (mm) for specimens examined and included in type series.

Field tag	MEP 0040 Holotype	MEP 0038	MEP 0045	MEP 0070	MEP 0072
Sample ID	MMC0049	MMC0057	MMC0064	MCR0088	MCR0090
UMMZ	232026	232002	232003	232004	232005
Locality	Muck-A-Muck	Muck-A-Muck	Muck-A-Muck	Mill Creek	Mill Creek
	Siskiyou Co., CA	Siskiyou Co., CA	Siskiyou Co., CA	Siskiyou Co., CA	Siskiyou Co., CA
Sex	F	J	M	F	J
SVL	70.7	31.3	54.2	66.3	41.5
TL	62.8	26.8	46.5	53.6	30.1
HL	17.9	8.5	13.8	15.2	10.7
HW	10.2	5.3	8.4	8.7	6.5
IO	2.4	1.6	2.8	3.1	2
IN	2.8	1.5	2.6	2.9	1.7
EN	2.2	0.9	2	1.9	1.6
FLB	12.5	7.3	12	12.5	8.6
HLB	15.9	8	13.4	15.6	9.6
PMM	60	49	52		
VM	8, 7	5, 5	7, 8		
COS	17	16	17	17	17
INCOS	3	2	2	3	2