

COMPARATIVE AUTECOLOGICAL CHARACTERISTICS OF NORTHWESTERN TREE SPECIES... A LITERATURE REVIEW

BY DON MINORE

FILE COPY

EDITOR'S
FILE COPY



ABSTRACT

This report is a compilation of autecological information previously scattered about in several hundred publications. It includes a comparison of the tolerances, traits, and attributes of native northwestern tree species. The species are ranked with respect to 69 environmental factors, phenotypic characteristics, and physical parameters. These rankings, with the literature references from which they were derived, should be useful aids to species selection, management, and research in the Pacific Northwest.

KEYWORDS: Autecology (plant), native plants, Pacific Northwest.





CONTENTS

INTRODUCTION	1
LIGHT	3
Shade Tolerance	3
Light Intensity	5
Photoperiod	6
TEMPERATURE	6
Frost Tolerance	6
Heat Tolerance	8
Mean Temperature Conditions	8
MOISTURE	9
Drought Tolerance	10
Excess Moisture	12
Optimum Moisture	13
NUTRIENTS	15
Macronutrient Concentrations	15
Micronutrient Concentrations	17
Nutrient Deficiency Tolerances	17
Nutrient Excesses	19
Ammonium Vs. Nitrate Nitrogen	19
Mycorrhizal Relationships	20
GROWTH	22
Seedling Root Growth Rate	22
Shoot Growth Period	24
Radial Growth Period	24
REPRODUCTION	25
Vegetative Reproductive	25
Seed Production	25
Seed Dissemination	25
Seed Crop Size	27
Seed Crop Frequency	28
Seed Soundness	28
Seed Flight	29
Seed Durability and Longevity	29
Seed Germination	29
Seed Stratification Requirements	32
Germination Environments	32
Seedbeds	32
PHYSICAL CHARACTERISTICS	34
Needles	35
Crowns	35
Litter Fall	40
Wood	40
Bark	41
Slash	41
Moisture Content	42
Foliage Flammability and Epiphyte Receptivity	42
Fire Resistance	42
Roots	43
Windthrow Resistance	44
Dense Soil Tolerance	44
Mechanical Damage Resistance	45

ANIMAL RELATIONSHIPS	45
INSECT RESISTANCE	46
DISEASE RESISTANCE	47
CHEMICALS	49
Herbicides	50
Salt	50
Sulfur Dioxide	50
Fluorides	50
Ozone	51
MISSING COMPARISONS	51
UNKNOWN ATTRIBUTES	54
RESEARCH NEEDS	55
LITERATURE CITED	56



INTRODUCTION

More than 40 native tree species are found in the Pacific Northwest¹ (table 1). Often differing almost as much as do the wide range of environments in which they occur, these species comprise an immensely valuable reservoir of growth traits, physiological tolerances, and physical attributes. By carefully selecting the proper species or combinations of species, foresters and forest researchers can often select biological raw material to fit their management and research needs.

Unfortunately, it is difficult to select biological raw material if one does not have some way of comparing this material in terms of the needs to be filled. Much is known about northwestern tree species, but most of this knowledge is scattered about in numerous publications that often are unavailable or unknown to the foresters and researchers who need them. This report is a collection of such scattered knowledge gleaned from available publications.


Published autecological information was included here whenever references contained comparisons or two or more northwestern tree species. Conclusions obtained by comparing the absolute values resulting from separate observations and experiments have been avoided. As Jarvis (1963) wrote, "The conditions of experiments are never the same as those occurring naturally, so that difficulties arise in applying the results to explain situations observed in the field. However, by doing similar experiments with several species under similar conditions, physiological differences in response between species may be found and these differences may be of ecological significance."

Differences, not absolute values, are presented in this report. Species

responses to differing conditions are complex and difficult to compare, however, even when only relative differences between species are investigated. Four major problems are involved: (1) Genetic variation within the species being compared may be almost as great as the variation among species, and differing ecotypic responses may obscure species comparisons; (2) Responses to differences in one condition may be confounded by responses to differences in other, associated conditions; (3) Species occupying widely diverse environments are difficult to compare directly, and young seedlings often must be grown under artificial conditions if these species are to be compared; (4) Young seedlings and older trees do not always respond similarly, and conclusions based upon seedling growth may not be valid for more mature trees. These problems are well summarized by Krajina (1955) and Spurr (1964, p. 165).

Nevertheless, northwestern tree species have been compared by many different investigators in many different environments. If carefully evaluated, these comparisons should be useful aids in selecting species for various environmental conditions. When all of them are listed, the published comparisons should also be useful indicators of unknown relationships and research needs.

Although all of the comparisons cited here involve species listed in table 1, some of these comparisons did not occur in British Columbia, Washington, Oregon, or Idaho. They were all useful, however, and comparisons that occurred elsewhere were not rejected when species characteristics were evaluated.



¹British Columbia, Washington, Oregon, and Idaho.

Table 1--Northwestern tree species, with their scientific and common names^{1/}

Scientific name	Common name
<i>Abies amabilis</i> (Dougl.) Forbes	Pacific silver fir
<i>Abies concolor</i> (Gord. and Glend.) Lindl.	White fir
<i>Abies grandis</i> (Dougl.) Lindl.	Grand fir
<i>Abies lasiocarpa</i> (Hook.) Nutt.	Subalpine fir
<i>Abies magnifica</i> A. Murr.	California red fir
<i>Abies magnifica</i> var. <i>shastensis</i> Lemm.	Shasta red fir
<i>Abies procera</i> Rehder	Noble fir
<i>Acer macrophyllum</i> Pursh	Bigleaf maple
<i>Acer negundo</i> L.	Boxelder
<i>Alnus rubra</i> Bong.	Red alder
<i>Arbutus menziesii</i> Pursh	Pacific madrone
<i>Betula papyrifera</i> Marsh.	Paper birch
<i>Castanopsis chrysophylla</i> (Dougl.) A.DC.	Golden chinkapin
<i>Chamaecyparis lawsoniana</i> Parl.	Port-Orford-cedar
<i>Chamaecyparis nootkantensis</i> (D. Don) Spach	Alaska-cedar
<i>Fraxinus latifolia</i> Benth.	Oregon ash
<i>Juniperus occidentalis</i> Hook.	Western juniper
<i>Juniperus scopulorum</i> Sarg.	Rocky mountain juniper
<i>Larix laricina</i> (Du Roi) K. Koch	Tamarack
<i>Larix lyalli</i> Parl.	Subalpine larch
<i>Larix occidentalis</i> Nutt.	Western larch
<i>Libocedrus decurrens</i> Torr.	Incense-cedar
<i>Lithocarpus densiflora</i> (H. and A.) Rehd.	Tanoak
<i>Picea breweriana</i> Wats.	Brewer spruce
<i>Picea engelmannii</i> Parry ex Engelm.	Engelmann spruce
<i>Picea glauca</i> (Moench) Voss	White spruce
<i>Picea mariana</i> (Mill.) B.S.P.	Black spruce
<i>Picea sitchensis</i> (Bong.) Carr.	Sitka spruce
<i>Pinus albicaulis</i> Engelm.	Whitebark pine
<i>Pinus attenuata</i> Lemm.	Knobcone pine
<i>Pinus contorta</i> Dougl. ex Loud.	Lodgepole pine
<i>Pinus flexilis</i> James	Limber pine
<i>Pinus jeffreyi</i> Grev. and Balf.	Jeffrey pine
<i>Pinus lambertiana</i> Dougl.	Sugar pine
<i>Pinus monticola</i> Dougl. ex D. Don	Western white pine
<i>Pinus ponderosa</i> Dougl. ex Loud.	Ponderosa pine
<i>Populus balsamifera</i> L.	Balsam poplar
<i>Populus tremuloides</i> Michx.	Quaking aspen
<i>Populus trichocarpa</i> T. and G. ex Hook.	Black cottonwood
<i>Pseudotsuga menziesii</i> (Mirbel) Franco	Douglas-fir
<i>Quercus chrysolepis</i> Liebm.	Canyon live oak
<i>Quercus garryana</i> Dougl.	Oregon white oak
<i>Quercus kelloggii</i> Newb.	California black oak
<i>Sequoia sempervirens</i> (D. Don) Endl.	Redwood
<i>Taxus brevifolia</i> Nutt.	Pacific yew
<i>Thuja plicata</i> Donn	Western redcedar
<i>Tsuga heterophylla</i> (Raf.) Sarg.	Western hemlock
<i>Tsuga mertensiana</i> (Bong.) Carr.	Mountain hemlock
<i>Umbellularia californica</i> (H. and A.) Nutt.	California-laurel

^{1/} Nomenclature is according to Garrison et al. (1976) and Little (1953). For the sake of clarity, scientific names are used throughout this report.

LIGHT

SHADE TOLERANCE

Shade tolerance will be used here in the restricted sense advocated by Shirley (1943), who defined it as "...the capacity of a tree species to survive in light of low intensity." Proceeding from shade tolerant to intolerant, species are discussed below and ranked in table 2.

Krajina (1965) listed *Abies amabilis* as the most tolerant coniferous tree species in the Pacific Northwest at low elevations and placed it in the same highly tolerant category as *Tsuga mertensiana* and *Chamaecyparis nootkatensis* at subalpine altitudes. Franklin (1964) considered *A. amabilis* to be even more tolerant than *T. mertensiana* in the Washington Cascade Range, also finding it to be more tolerant than *Tsuga heterophylla* in most true fir - hemlock forests. In coastal British Columbia, Schmidt (1957) found *A. amabilis* to at least equal *T. heterophylla* and *Thuja*

plicata in shade tolerance, probably being the most tolerant of all native coastal conifers. This ranking is supported by Hanzlik (1932), who noted that *A. amabilis* is more tolerant than *Tsuga heterophylla* on the western and southern slopes of the Olympic Mountains. It conflicts with the observations of Buckland et al. (1949), who noted that *A. amabilis* appeared to be less tolerant than *T. heterophylla* on Vancouver Island. Working with the collected observations of others on a national scale, without the local experience of the authorities cited above, Sudworth (1908) and Baker (1949) ranked *Abies amabilis* as less shade tolerant than *Taxus brevifolia*, *Thuja plicata*, *Chamaecyparis nootkatensis*, and *Tsuga heterophylla*. Their rankings probably are less accurate than those of Schmidt (1957), Franklin (1964), and Krajina (1965).

The above statements were based upon field observations of seral and climax stands as they existed in nature, where shade tolerance was assumed to have been the dominant

Table 2--Comparative shade tolerance of northwestern tree species^{1/}

<i>Abies amabilis</i> , <i>Tsuga heterophylla</i> , <i>Taxus brevifolia</i>
<i>Thuja plicata</i> , <i>Chamaecyparis nootkatensis</i> , <i>Tsuga mertensiana</i>
<i>Abies lasiocarpa</i>
<i>Abies grandis</i>
<i>Picea sitchensis</i>
<i>Picea glauca</i>
<i>Abies concolor</i> , <i>Picea breweriana</i>
<i>Abies magnifica</i> var. <i>shastensis</i>
<i>Chamaecyparis lawsoniana</i>
<i>Pinus monticola</i>
<i>Pseudotsuga menziesii</i>
<i>Pinus lambertiana</i>
<i>Libocedrus decurrens</i> , <i>Abies procera</i> , <i>Picea engelmannii</i> , <i>Alnus rubra</i>
<i>Pinus contorta</i>
<i>Larix occidentalis</i>
<i>Pinus ponderosa</i>
<i>Juniperus occidentalis</i> , <i>Quercus kelloggii</i>

^{1/}Compiled from information published by Larsen (1930), Haig et al. (1941), Baker (1949), Buckland et al. (1949), Wood (1955), Edwards (1957), Schmidt (1957), Tackle (1959), Franklin (1964), Franklin and Mitchell (1967), Hodges and Scott (1968), Krajina (1965, 1970), Krueger and Ruth (1969), Thornburgh (1969), Eis (1970), Tinus (1970), Brix (1972), Emmingham and Waring (1973), Franklin and Dyrness (1973), Wali and Krajina (1973), Ronco (1975), Edmonds (1975), Waring et al. (1975). More shade tolerant species are listed above less tolerant ones. Species on the same line are not necessarily equal, but data are insufficient for their separation.

factor determining species relationships. Other factors may have been involved, however, and successional status cannot always be equated with relative shade tolerance. When he measured seedlings growing under low-light conditions, Thornburgh (1969) found *Tsuga heterophylla* photosynthesis and height growth rates to be higher than those of *Abies amabilis*. When all field and laboratory observations are considered, it is difficult to compare the shade tolerances of *Abies amabilis*, *Tsuga heterophylla*, and *Taxus brevifolia*. Nevertheless, these three species are the most shade-tolerant trees in the Pacific Northwest.

Tsuga mertensiana is slightly less tolerant than *Abies amabilis* in British Columbia (Krajina 1970). Larsen (1930) also rated it as slightly less tolerant than *Thuja plicata* and *Taxus brevifolia* in Montana and northern Idaho--a rating supported by Baker's nationwide poll of professional opinion in 1949. *Tsuga mertensiana* is the most tolerant species present at high elevations in most of southern Oregon (Franklin 1964), but it may be less shade tolerant than *Abies concolor* or *Picea breweriana* in the Siskiyou Mountains (Waring et al. 1975).

Although Sudworth (1908) noted that *Chamaecyparis nootkatensis* is less shade tolerant than *Thuja plicata*, Krajina (1965, 1970) considered it to be slightly more tolerant. Both species probably are about as shade tolerant as *Tsuga mertensiana* in most environments. Referring to studies of seedling mortality, Harris and Farr (1974) concluded that *Thuja plicata* is less shade tolerant than *Tsuga heterophylla* or *Picea sitchensis* in southeastern Alaska. Their seedling mortality may not have been a result of shade intolerance, but *T. plicata* seems to be less tolerant in coastal Alaska than it is inland or farther south. Larsen (1930) and Haig et al. (1941) rated *T. plicata* as even more shade tolerant than *Tsuga heterophylla* in the Rocky Mountains.

Abies lasiocarpa is less tolerant than *A. amabilis* or *Tsuga mertensiana* (Franklin and Mitchell 1967). Despite the contrary opinions of

Sudworth (1908) and Haig et al. (1941), *Abies lasiocarpa* seems to be considerably more shade tolerant than *Picea engelmannii* (Larsen 1930, Baker 1949, LeBarron and Jemison 1953, Smith and Clark 1960, Krajina 1965). *Abies lasiocarpa* also is more shade tolerant than *Picea glauca*, *Pinus contorta* (Wali and Krajina 1973), or *Abies grandis* (Larsen 1930, Baker 1949, Schmidt 1957).

Abies grandis seems to be more shade tolerant than *Pseudotsuga menziesii*, *Pinus monticola*, or *Picea engelmannii* (Sudworth 1908, Larsen 1930, Haig et al. 1941, and Baker 1949). Photosynthetic rates measured by Hodges and Scott (1968) also indicate that it is more shade tolerant than *Abies procera* or *Picea sitchensis*.

Picea sitchensis is more shade tolerant than *Pseudotsuga menziesii* in its native habitat (Wood 1955). *P. sitchensis* also is more shade tolerant than *P. glauca* when grown in controlled environments (Brix 1972).

Although Baker's (1949) poll and Franklin and Dyrness' (1973) tolerance table placed *Picea sitchensis*, *Abies grandis*, and *A. concolor* in the same "tolerant" category, Krajina's (1965) ranking of *A. concolor* as considerably less tolerant than the other two species probably is correct. *A. concolor* is shade tolerant, however, for Emmingham and Waring (1973) found both *A. concolor* and *A. magnifica* var. *shastensis* at low-light levels where *Pseudotsuga menziesii* was absent in southwest Oregon. Waring et al. (1975) also found the *Abies* species to be more tolerant than *P. menziesii* and observed that *A. concolor* and *Picea breweriana* are slightly more shade tolerant than *A. magnifica* var. *shastensis*.

Abies magnifica var. *shastensis* is more shade tolerant than *Pinus lambertiana* and *P. monticola* (Krajina 1965, Waring et al. 1975). *Pinus lambertiana*, *P. monticola*, and *Pseudotsuga menziesii* are all given an intermediate tolerance rating by Baker (1949) and Waring et al. (1975); but Krajina (1965) ranked *P. lambertiana* as less tolerant than *Pseudotsuga menziesii*, *P. monticola* as more tolerant. Krajina's ranking seems more appropriate for environments in which these species occur together.

Chamaecyparis lawsoniana is even more shade tolerant than *Pinus monticola*; and it also ranks above *Pseudotsuga menziesii*, *Pinus lambertiana*, *P. contorta*, *Libocedrus decurrens*, *Abies procera*, *Picea engelmannii*, *Alnus rubra*, and *Larix occidentalis* (Baker 1949, James and Hayes 1954, Franklin and Dyrness 1973). Franklin and Dyrness (1973) place it in the same tolerant category as *Lithocarpus densiflora*, *Acer macrophyllum*, and *Sequoia sempervirens*. Zobel and Hawk² found *C. lawsoniana*, *Tsuga heterophylla*, and *Pinus monticola* in the same shady microhabitats.

Hodges and Scott (1968) found that the photosynthetic rate of *Abies procera* was less than that of *Pseudotsuga menziesii* in low-light conditions, greater than *P. menziesii* in high light. These data support the relative tolerance ratings of Baker (1949), who characterized *P. menziesii* as "intermediate," *A. procera* as "intolerant." They apparently conflict with Krajina's (1970) listing of both *A. magnifica* var. *shastensis* and *A. procera* as more shade tolerant than *P. menziesii*. Interacting light and moisture effects may be responsible for conflicts such as this. Atzet and Waring (1970) found that the minimum light requirement for *P. menziesii* increased where moisture became limiting.

Although Baker (1949) and Franklin and Dyrness (1973) listed *Picea engelmannii* as more tolerant than either *Pseudotsuga menziesii* or *Abies procera*, Tinus' (1970) measurements indicate that *P. engelmannii* and *Alnus rubra* probably are less shade tolerant than *P. menziesii*. Where subjective ratings conflict with objective measurements, the measurements have been given precedence here. Tinus' results were used in constructing table 2.

Edmonds (1975), describing the photosynthesis measurements of Helms and Rutter, rated *Libocedrus decurrens* as less tolerant than *Pseudotsuga menziesii*, more tolerant than *Pinus ponderosa*. Again, these measurements conflict with the estimated *Pseudotsuga* - *Libocedrus* ratings compiled by Baker (1949), Krajina (1965), and Franklin and Dyrness (1973); but

photosynthetic measurements probably are more reliable than estimates of relative tolerance.

Tinus (1970) and Ronco (1975) measured lower light saturation points for *Picea engelmannii* than for *Pinus contorta*. These measured indications of greater *P. engelmannii* shade tolerances are supported by the conclusions of several observers (Larsen 1930, Haig et al. 1941, Baker 1949, Krajina 1965, and Franklin and Dyrness 1973). The *Pinus contorta* - *Pinus ponderosa* tolerance difference is less distinctive, but *P. contorta* is more shade tolerant than *P. ponderosa* (Larsen 1930, Krajina 1965, Kerr 1913, and Haig et al. 1941).

Although Mosher (1965) implied that *Larix occidentalis* was more shade tolerant than *Pinus contorta* in a northeastern Washington stand, both Baker (1949) and Tackle (1959) considered *L. occidentalis* to be less tolerant. The shade tolerance of *L. occidentalis* has most often been placed somewhere between the tolerances of *Pinus contorta* and *P. ponderosa* (Larsen 1930, Haig et al. 1941, Krajina 1965).

Larsen (1930) and Krajina (1965) considered *Juniperus occidentalis* to be even less shade tolerant than *Pinus ponderosa*. Edwards (1957) also found *Quercus kelloggii* to be less tolerant than *P. ponderosa*. Krajina (1965) listed *Larix lyalli* and *Pinus albicaulis* as the most intolerant subalpine species.

LIGHT INTENSITY

Many species comparisons in the literature provide supplementary light data in addition to shade tolerance information. Emmingham and Waring (1973) recorded maximum leader growth of *Pseudotsuga menziesii*, *Abies concolor*, and *A. magnifica* var. *shastensis* at 100-percent light intensity in southwestern Oregon. Fairbairn and Neustein (1970) noted a similar response for *Picea sitchensis* grown under full sunlight in England but found that *Pseudotsuga menziesii*, *Abies grandis*, and *Tsuga heterophylla* attained maximum height growth at lower light levels. The English environment may have been responsible for this difference.

Controlled environments were utilized by Tinus (1970) in comparing the photosynthesis of several species under increasing light intensities.

² Zobel, Donald B., and Glenn M. Hawk. The environment of *Chamaecyparis lawsoniana*. Unpublished manuscript.

He found that *Picea sitchensis*, *Tsuga heterophylla*, and *Pseudotsuga menziesii* were light saturated at lower light intensities than *Picea engelmannii* and *Alnus rubra*. Krueger and Ruth (1969) found that *Alnus rubra*'s photosynthetic rate also was higher than those of *Picea sitchensis*, *Tsuga heterophylla*, and *Pseudotsuga menziesii* at high light intensities. Clark and Lister (1975) measured three times as much chlorophyll per unit fresh weight in *Alnus rubra* as in *P. menziesii* or *Picea sitchensis* and found *A. rubra* photosynthesis to be more efficient in blue light. The *Alnus rubra* seedling certainly seems to be a marvelously effective photosynthetic mechanism.

Hellmers (1964) measured growth distribution effects in *Sequoia sempervirens* and *Abies magnifica* seedlings grown under different light intensities. Although light intensity scarcely affected growth distribution in *S. sempervirens*, high light intensities produced the greatest basal growth in *A. magnifica*.

When Pharis et al. (1967) made diurnal comparisons of *Pseudotsuga menziesii* and *Pinus ponderosa*, they found that *P. menziesii* photosynthetic rates changed more rapidly than those of *P. ponderosa*, increasing faster in the morning and declining sooner and further in the afternoon.

PHOTOPERIOD

Owston's 1974 observations of the effects of shortened photoperiod indicate that *Picea sitchensis* and coastal *Pseudotsuga menziesii* are more sensitive than *Abies procera* and *Pinus ponderosa* to shortened photoperiods. *Picea sitchensis* seems to have a longer optimum photoperiod than *Thuja plicata* (Malcolm and Caldwell 1971). When Vaartaja (1959) used the ratio of height growth under very long days to height growth under very short days in species comparisons, he found $Picea engelmannii \geq P. sitchensis \geq Pinus contorta \geq P. ponderosa \geq Pseudotsuga menziesii \geq Thuja plicata \geq Tsuga mertensiana$. In general, the more northerly a species, the greater was its photoperiodic sensitivity.

TEMPERATURE

Genetic variation within species often makes the comparison of species temperature responses difficult. Different geographic races of the same species differ--sometimes in unexpected ways. For example, northern races and those from inland locations usually are more frost tolerant than their southern and coastal relatives (Haller 1961) Sakai and Okada 1971, Sakai and Weiser 1973). Nevertheless, Conkle et al. (1967) observed more winter injury in *Abies concolor* seedlings from northern locations than in those from southern seed sources. Within-species variation should be kept in mind when pondering the species comparisons presented here. Species temperature responses are considered under three categories: frost tolerance, heat tolerance, and mean temperature conditions.

FROST TOLERANCE

Frost tolerance comparisons are most abundant in the literature. They are summarized in table 3.

The freezing resistances of dormant buds and twigs measured by Sakai and Weiser (1973) show that *Pinus contorta* and *Pinus monticola* are the most frost-resistant northwestern tree species while in the dormant state. Their data indicate *Picea engelmannii* to be the next most resistant, followed by *Populus trichocarpa*, *Alnus rubra*, *Abies lasiocarpa* and *A. concolor*, *Larix occidentalis*, *Pinus ponderosa*, *P. jeffreyi*, *Picea sitchensis*, *Chamaecyparis nootkatensis*, *Thuja plicata* and *Abies grandis*, *A. amabilis*, *Pseudotsuga menziesii*, *Tsuga mertensiana*, and *T. heterophylla*...in approximately that order. This frost resistance ranking, derived from the comparison of frost damage to dormant twigs artificially hardened before freezing in the laboratory, somewhat resembles the less complete but more realistic ranking that can be created by compiling literature references to frost damage in actively growing trees.

Table 3--Comparative frost tolerances of northwestern tree species^{1/}

Pinus contorta, *P. monticola*
Picea engelmannii
Populus trichocarpa
Pinus ponderosa, *P. jeffreyi*, *Abies magnifica* var. *shastensis*,
Tsuga mertensiana
Libocedrus decurrens, *Picea breweriana*
Pinus lambertiana
Abies concolor, *A. lasiocarpa*
Abies grandis, *A. amabilis*
Picea sitchensis
Pseudotsuga menziesii
Thuja plicata
Tsuga heterophylla
Arbutus menziesii

^{1/}Compiled from information published by Day (1928), Schubert (1955), Duffield (1956), Daubenmire (1957), Wagener (1960), Haller (1961), Stein (1963), Fowells and Stark (1965), Krajina (1970), Cochran and Berntsen (1973), Sakai and Weiser (1973), Waring et al. (1975), and Timmis (undated). Tolerant species are listed above intolerant ones. Species in the same group are not necessarily equal, but data are insufficient for their separation.

Young *Pinus contorta* seedlings are more frost tolerant than young *P. ponderosa* seedlings (Berntsen 1967, Cochran and Berntsen 1973), and a late spring frost in central Oregon damaged *P. contorta* megasporangiate strobili much less than *P. ponderosa* strobili at the same stage of development (Sorensen and Miles 1974).

Pinus jeffreyi may be slightly more frost tolerant than *P. ponderosa* near the California coast (Haller 1959, 1961), but the two species appear equally frost tolerant elsewhere (Wagener 1960, Haller 1961). Frost tolerances of *Abies magnifica* var. *shastensis* and *Tsuga mertensiana* equal that of *P. jeffreyi* (Waring et al. 1975).

Waring et al. noted that *Picea breweriana* is less tolerant of cold temperatures than *Pinus jeffreyi*, more tolerant than *Abies grandis*, *Pseudotsuga menziesii*, or *Pinus jeffreyi*. As *Libocedrus decurrens* is less frost tolerant than *Pinus ponderosa* but more tolerant than *Abies grandis*, *Pseudotsuga menziesii*, or *Pinus lambertiana* (Stein 1963, Fowells and Stark 1965), it may be ranked with *Picea breweriana* in this respect.

Schubert (1955) observed that *Pinus lambertiana* was more susceptible to freezing than *P. ponderosa* and *P. jeffreyi*. *P. lambertiana* survival also was poorer than *Libocedrus decurrens* survival, but better than that of *Abies concolor* in the frost mortality measurements made by Fowells and Stark (1965) in California. Stein's (1963) southwestern Oregon observations were similar--frost damage to *Pinus lambertiana* was more severe than to *Libocedrus decurrens*, but less severe than that to *Abies grandis* or *Pseudotsuga menziesii*. *P. menziesii* is less frost tolerant than *Picea glauca* in the interior of British Columbia (Cayford and Bickerstaff 1968).

A November 1955 cold wave in western Washington damaged the *Abies* spp. less than *Picea sitchensis*, and *P. sitchensis* was less damaged than *Pseudotsuga menziesii* (Duffield 1956). Day (1928) also found *Picea sitchensis* to be more frost resistant than *Pseudotsuga menziesii* in England. Sakai and Okada (1971) found the dormant frost resistance of *Picea sitchensis* to be similar to that of *Chamaecyparis lawsoniana*, but Duffield (1956) observed more damage to *C. lawsoniana*

than to *P. sitchensis* after the November freeze. Similarly, the excellent dormant frost tolerance of *Alnus rubra* reported by Sakai and Weiser (1973) was not evident in November, 1955, when *Alnus* and *Arbutus* trees were extensively killed by unusually cold temperatures (Daubenmire 1957). As Sakai and Weiser measured only the frost tolerance of detached dormant twigs, Duffield's and Daubenmire's observations of entire trees probably are more pertinent. Both Duffield (1956) and Daubenmire (1957) found *Pseudotsuga menziesii* more frost tolerant than *Thuja plicata*, and *T. plicata* more tolerant than *Tsuga heterophylla*. These observations apply only to the shoots, however, for Van Eerden (1974) found *Pseudotsuga* roots to be more susceptible to low temperatures than *Tsuga* roots. *Arbutus menziesii* is "...the least frost resistant tree native to Canada" (Krajina 1970).

HEAT TOLERANCE

Levitt (1951) stated that frost, heat, and drought resistances are basically similar, with resistance to one accompanied by resistance to the other two factors. Although this seems to be true in a general sense, relative resistances to heat injury are not the same as resistances to frost injury when species rankings from the literature are compiled and compared. Unfortunately, only a few species have been compared for heat tolerance.

Pinus lambertiana is more heat resistant than *P. ponderosa* or *Pseudotsuga menziesii* (Cochran 1963). The heat tolerance of *Pinus jeffreyi* is at least equal to that of *P. ponderosa* (Haller 1959), but no *P. jeffreyi* - *P. lambertiana* comparisons are available. *Pseudotsuga menziesii*, *Pinus monticola*, and *Larix occidentalis* seedlings are more heat resistant than *Tsuga heterophylla* or *Thuja plicata* seedlings (Haig 1936). *Pseudotsuga* seedlings also are more resistant than those of *Abies grandis* (Baker 1929), which is more susceptible to heat killing than other species (Schubert and Adams 1971). *Abies lasiocarpa* is

able to establish itself in hotter conditions than *Picea engelmannii* - *p. glauca* hybrids in the Rocky Mountains (Day 1964).

MEAN TEMPERATURE CONDITIONS

In addition to comparing frost and heat responses, it often is advantageous to compare species responses to mean temperatures. This may be done by comparing the published relative occurrence and/or growth of species observed under identical temperatures...for several different temperature conditions.

Quercus kelloggii, *Quercus garryana*, and *Arbutus menziesii* occur in the warmest northwestern environments (Cleary and Waring 1969, Zobel et al. 1976). In the eastern Siskiyou Mountains of southern Oregon, *Pinus ponderosa* also often occurs in hot environments (Waring 1969).

Frost tolerance notwithstanding, *Pinus contorta* does not seem to grow any better than *P. ponderosa* under low night temperatures (Cochran 1972). Its boreal range extends farther north, however, than that of *P. ponderosa* (Harlow and Harrar 1950). Although it demands a lower average temperature than *P. ponderosa* and *Pseudotsuga menziesii* in the Rocky Mountains, *Pinus contorta* probably requires a higher temperature than *Picea engelmannii* or *Abies lasiocarpa* (Tackle 1959). Kaufmann and Eckard (1977) recently suggested that *Picea engelmannii* establishment may occur more rapidly than that of *Pinus contorta* in cool environments.

In the central Oregon Cascades *Pseudotsuga menziesii*, *Pinus lambertiana*, *Acer macrophyllum*, and *Libocedrus decurrens* are most important where environments are slightly cooler than those typical of *Arbutus*'s center of importance (Zobel et al. 1976). In controlled environments, *Pseudotsuga menziesii* grows less well than *Pinus ponderosa* under high temperature conditions (see Steinbrenner and Rediske 1964, Pharis et al. 1967). Nevertheless, high temperatures seem to be less detrimental to *P. menziesii* than they are to *Tsuga heterophylla* (Brix 1971). A

small diurnal temperature variation appears to be beneficial for *P. menziesii*, but variation is not required for *Sequoia sempervirens* (Hellmers and Sundahl 1959).

When Oregon Cascade environments cooler than those typical of *Pseudotsuga menziesii*-dominated stands are considered, *Thuja plicata*, *Tsuga heterophylla*, *Abies grandis*, *A. procera*, and *A. amabilis* become more important as temperatures drop (Zobel et al. 1976). *Abies magnifica* var. *shastensis* and *Tsuga mertensiana* occupy the coolest forested environments in the eastern Siskiyou Mountains (Cleary and Waring 1969).

Farther north, timberline for *Abies amabilis* in British Columbia is about 150 m (492 ft) lower than timberline for *Tsuga mertensiana*, *Abies lasiocarpa*, and *Pinus albicaulis* (Krajina 1970). Using controlled temperatures, Brix (1972) found *Picea glauca* to be more productive than *P. sitchensis* under high temperature - high light conditions.

A complete ranking of the mean temperature relations of northwestern tree species has not been attempted, but approximate relationships based upon available information are summarized in table 4. The approximate accuracy of field observation is associated with a Rocky Mountain

ranking of species susceptibility to "red belt" injury (Mason 1915):

Juniperus sp. (most resistant)
Abies lasiocarpa
Picea engelmannii
Pinus contorta
Pseudotsuga menziesii
Pinus ponderosa (least resistant)

Red belt injury apparently results from crown desiccation that occurs when high needle transpiration is accompanied by frozen ground conditions that prevent water uptake. As such, it is a combination of temperature and moisture phenomena.

MOISTURE

Moisture comparisons among species may be considered in three categories: tolerance of deficient moisture, tolerance of excess moisture, and optimum moisture conditions. If quantity of published information is any indication of the relative importance of these categories, ability to endure deficient moisture (drought tolerance) is most important.



Table 4--Mean temperature relationships among northwestern tree species^{1/}

Pinus ponderosa, *Quercus kelloggii*, *Quercus garryana*,
Arbutus menziesii
Pseudotsuga menziesii, *Pinus lambertiana*, *Libocedrus decurrens*,
Acer macrophyllum
Thuja plicata
Tsuga heterophylla
Abies grandis
Abies amabilis, *A. procera*, *A. magnifica* var. *shastensis*
Tsuga mertensiana, *Abies lasiocarpa*, *Pinus albicaulis*
Pinus contorta
Picea mariana, *P. glauca*, *Larix laricina*

^{1/}Compiled from information published by Hustich (1953), Steinbrenner and Rediske (1964), Pharis et al. (1967), Waring (1969), Cleary and Waring (1969), Krajina (1970), Brix (1971), and Zobel et al. (1976). Species adapted to warm temperatures are listed above those adapted to cool temperatures. Species in the same group are not necessarily equal, but data are insufficient for their separation.

DROUGHT TOLERANCE

Dry conditions may be endured by trees; or they may be avoided (by rapid and deep root growth, for example). In either case, several physiological attributes contribute to drought tolerance. Unfortunately, related attributes do not always coincide with overall tolerance and different researchers do not always agree. This is particularly true of the drought endurance associated with stomatal closure. Hodges (1967) found that *Abies procera* stomata closed more completely than *Pseudotsuga menziesii* stomata at night...and *P. menziesii* stomata closed more completely than *Tsuga heterophylla* and *Picea sitchensis*. *Abies concolor* stomata did not close at all. Hodges and Scott (1968) concluded that *A. procera* controlled internal moisture better than the other species. Running (1973, 1976) noted that *A. procera* showed little stomatal control, however, whereas *Pseudotsuga menziesii* and *A. concolor* appeared to have good control. When the data and conclusions of different researchers differ so, it is difficult to integrate them. In this case, characteristics of the environments in which the species normally occur make Running's conclusions seem more appropriate than those of Hodges and Scott.

Lopushinsky (1969) also studied stomatal closure. He found that *Pinus ponderosa*, *P. contorta*, and *Picea engelmannii* stomata closed sooner than those of *Pseudotsuga menziesii* and *Abies grandis*. Wambolt (1973), in a study of conifer water potential, concurred with Lopushinsky in ranking *Pinus ponderosa* stomatal control as more sensitive than that of *P. contorta*, *Picea engelmannii*, and *Pseudotsuga menziesii*.

Barker (1973) observed less stomatal control in *Abies concolor* than in *Pinus ponderosa* under comparable environmental conditions, but Rutter (1977), working with foliar samples exposed to full sunlight in the field, recorded more stomatal control in *A. concolor*. He found *Libocedrus decurrens* stomata to be more sensitive to increasing moisture stress than those of *A. concolor* or *P. ponderosa*

and attributed its drought tolerance to stomatal control, suggesting that the long tap root of *P. ponderosa* allows it to avoid moisture stress rather than tolerate it. Rutter worked with day-to-day fluctuations that resulted in few common environmental conditions for the three species studied, however; and he did not attempt to account for within tree, between tree, or seasonal differences in behavior. Perhaps these unmeasured variables affect Rutter's *Pinus ponderosa* observations. Cleary (1971) observed an abrupt decline in *P. ponderosa* photosynthesis at moderate plant moisture stress (15-20 atm), a gradual decline in *Pseudotsuga menziesii* photosynthesis over a much wider range (<8->22 atm). Both declines probably were associated with stomatal closure.

Stomatal closure has not been studied in all northwestern tree species, but drought tolerance as such has been widely studied and observed (table 5). This may be just as well, for as Parker (1969) observed, whole seedlings involve other adaptations that may be more important than leaf resistance alone.

Franklin and Dyrness (1973) ranked *Quercus garryana* as the most moisture-stress tolerant southwestern Oregon major tree species, followed by *Quercus kelloggii*, *Pinus ponderosa*, and *Arbutus menziesii*. They did not rank *P. jeffreyi*, but Stone (1957), studying the effect of artificial dew on the survival of four western conifers, observed that unwatered seedlings of *P. jeffreyi* in greenhouse cans survived longer than unwatered *P. ponderosa*, *Libocedrus decurrens*, or *Abies concolor* seedlings. When he applied artificial dew in the form of a nightly water spray, *A. concolor* still died first, followed by the pines (together) and finally *L. decurrens*. Stone qualified his results by pointing out that greenhouse behavior may not duplicate that in the field, where differences in relative root growth rates among species might alter the relative drought resistance observed in greenhouse containers. Nevertheless, Haller (1959) noted that *Pinus jeffreyi* is more tolerant of aridity

Table 5--Comparative drought tolerances of
northwestern tree species^{1/}

Quercus garryana
Quercus kelloggii
Pinus jeffreyi
Pinus ponderosa
Pinus contorta, *Libocedrus decurrens*, *Arbutus menziesii*
Pseudotsuga menziesii
Picea engelmannii
Abies grandis
Pinus lambertiana, *Larix occidentalis*
Abies lasiocarpa, *Thuja plicata*, *Pinus monticola*
Abies concolor, *Picea breweriana*
Chamaecyparis lawsoniana
Tsuga heterophylla, *Picea sitchensis*
Abies amabilis
Abies magnifica var. *shastensis*, *Tsuga mertensiana*

^{1/}Compiled from information published by Daubenmire (1943), Tarrant (1953), Stone (1957), Haller (1959), Cochran (1963), Pharis (1966), Fraser and Cordes (1967), Kotar (1972), Franklin and Dyrness (1973), and Edmonds (1975). Drought tolerant species are listed above intolerant ones. Where species occur on the same line, information is insufficient to separate them.

than *P. ponderosa* in California. Tarrant (1953), comparing the characteristics of *P. ponderosa* and *P. contorta* in a literature review, attributed more drought tolerance to *P. ponderosa*. *P. ponderosa* also has a greater physiological capacity to exploit hot, dry environmental conditions than *L. decurrens* (Edmonds 1975) or *Pseudotsuga menziesii* (Daubenmire 1943, Daubenmire and Deters 1947, Parker 1951, and Cochran 1963).

Further comparisons involving *P. menziesii* are numerous. Marshall (1931), measuring critical soil moisture and survival, observed that *P. menziesii*, *Abies grandis*, and *Thuja plicata* survived with less soil moisture than that required by *Pinus monticola*, *Tsuga heterophylla*, or *Picea glauca*. Although transpiration rates declined faster in *Picea engelmannii* than in *Pseudotsuga menziesii* and *Abies grandis* when potted seedlings were subjected to moderate soil moisture stresses by Lopushinsky and Klock (1974), Daubenmire (1943) observed that *Pseudotsuga menziesii* survived longer than *Picea engelmannii* and *P. engelmannii* survived longer than *Abies lasiocarpa* or *Thuja*

plicata when his potted seedlings were maintained at or below the wilting point for 2, 4, 6, 8, and 10 days. Using diurnal variations in sap pressure and dormancy as their criteria. Fraser and Cordes (1967) rated *Pseudotsuga menziesii* more able to avoid moisture stress than *Tsuga heterophylla* or *Picea sitchensis*.

When Parker (1951) cut leaves from several tree species and measured moisture retention, *Pseudotsuga menziesii* leaves retained a higher moisture content than those of *Picea engelmannii*, *Thuja plicata*, *Abies grandis*, or *Pinus monticola*. Measuring foliage moisture content and soil moisture stress, Pharis (1966) ranked *Pseudotsuga menziesii* and *Libocedrus decurrens* above *A. grandis* and *A. grandis* above *Pinus lambertiana* in drought resistance. In considering these comparisons, it is important to remember that *P. menziesii* trees from different geographic localities... and even from different aspects... may differ in drought resistance (Pharis and Ferrell 1966, Ferrell and Woodward 1966). Similar variation probably occurs in other species.

Pinus lambertiana seems to be more drought tolerant than *P. monticola* in southwestern Oregon

(Franklin and Dyrness 1973); and *P. monticola* appears to be less able to adapt to inadequate soil moisture than *Pseudotsuga menziesii*, *Abies grandis*, *Larix occidentalis*, or *Thuja plicata* in the western white pine type (Leaphart and Wicker 1966). Southwestern Oregon *Pinus lambertiana* also is more drought tolerant than *Picea breweriana*, which is quite tolerant of soil moisture stress but intolerant of evaporative demand (Waring et al. 1975). Waring et al. indicate that *Abies concolor* in southwestern Oregon slightly exceeds *P. breweriana* in ability to tolerate evaporative demand but is a little less tolerant of soil moisture stress. *Tsuga mertensiana* and *Abies magnifica* var. *shastensis* are much less tolerant of soil moisture deficiencies.

Franklin and Dyrness (1973) ranked *Chamaecyparis lawsoniana*, *Abies magnifica* var. *shastensis*, *Tsuga mertensiana*, and *T. heterophylla* below *Abies concolor* in decreasing order of moisture stress tolerance in southwestern Oregon. As Kotar (1972) found *T. heterophylla* to be more drought tolerant than *Abies amabilis* when seedlings of both species were subjected to high moisture stresses in the laboratory and as Lowry's 1972 laboratory measurements showed *A. amabilis* to be more drought tolerant than *Tsuga mertensiana*, Franklin and Dyrness's *Tsuga* rankings probably should be reversed. *Tsuga heterophylla* seems to be more drought tolerant than *T. mertensiana*. Puritch (1973) in comparing the effect of water stress on photosynthesis, respiration, and transpiration of four *Abies* species, found *A. grandis* to be most stress tolerant, followed by *A. lasiocarpa*, *A. amabilis*, and *A. balsamea*--in that order.

EXCESS MOISTURE

Excess moisture usually is the result of flooding or the presence of a shallow water table in nature. Observations of flood mortality, through sometimes confounded by other factors, often provide the only available species comparisons. Brink (1954), observing the results

of a 1948 spring flood in the lower Fraser River Valley, found *Picea sitchensis*, *Pinus contorta*, and *Thuja plicata* mortality to be lower than that of *Tsuga heterophylla*. *T. heterophylla* mortality was lower than that of *Pseudotsuga menziesii* or *Alnus rubra*. Krajina (1970) observed that *Populus trichocarpa*, *P. tremuloides*, *Abies grandis*, *A. lasiocarpa*, *Thuja plicata*, *Picea sitchensis*, *P. engelmannii*, *P. glauca*, and *Tsuga heterophylla* survive on floodplain habitats where *Pseudotsuga menziesii* cannot grow. *Sequoia sempervirens* and *Umbellularia californica* survive flooding that kills *P. menziesii*, *A. grandis*, and *Lithocarpus densiflora* (Stone and Vasey 1968).

Controlled flooding experiments indicate that *Pinus contorta* is more tolerant of oxygen deficiency than is *Picea sitchensis* (Boggie 1974, Crawford and Baines 1977), surviving better when seedlings of both species are artificially flooded (Minore (1968). Minore also found *Pinus contorta* and *Thuja plicata* seedlings to be more flood tolerant than those of *Alnus rubra* and *Tsuga heterophylla*, which were more tolerant than *Pseudotsuga menziesii* seedlings. Gill (1970), however, emphasized in his excellent flooding review that it probably is not permissible to extrapolate from seedling performance to mature trees when dealing with flood tolerance. Physical size may be very important.

Soil characteristics apparently influence species tolerance of shallow water tables in nature. Howell (1931) found *Pinus contorta* in the saturated soil over hard pan areas in California and claimed that *P. ponderosa* could not endure such excess moisture. Like Howell, Tarrant (1953) concluded that *P. contorta* was very tolerant of continuous high soil moisture but *P. ponderosa* was intolerant. In contrast, Stephens (1965) observed that *P. ponderosa* thrived on imperfectly drained soils with clay subsoils developed from Eocene to Miocene volcanic rocks in the western Cascade Mountains of Oregon, growing better than *Pseudotsuga menziesii* or *Pinus lambertiana* on these moist, fine-textured substrates. The Eocene to Miocene

parent materials seem to be uniquely effective in favoring *P. ponderosa* on wet sites. More than moisture must be involved, for Cochran (1972) obtained no significant mortality in either *P. ponderosa* or *P. contorta* when he subjected seedlings to carefully controlled shallow water-table conditions in a greenhouse. Haller (1959) indicated that *P. jeffreyi* is at least as tolerant of high moisture as *P. ponderosa*.

Minore and Smith (1971), reporting on the occurrence and growth of four tree species in swamps and stream bottoms on the Olympic Peninsula, ranked *Alnus rubra* and *Thuja plicata* as most tolerant of shallow water tables, followed by *Picea sitchensis* and (least tolerant) *Tsuga heterophylla*. Absence of *Pseudotsuga menziesii* in these wet habitats indicated even less tolerance than *T. heterophylla*. When Minore (1970) compared the seedling growth of eight species over artificial water tables, he found *Pinus contorta*, *Thuja plicata*, *Alnus rubra*, and *Picea sitchensis* to be more tolerant of shallow water tables than *Pseudotsuga menziesii*. These and other excess-moisture tolerance relationships are summarized in table 6.

OPTIMUM MOISTURE

Optimum moisture conditions are difficult to determine in controlled growth chamber or greenhouse environ-

ments. Other environmental factors influence species growth, and the resulting interactions often are difficult to identify and analyze. Where optimum conditions are defined on the basis of natural range comparisons in the field, further difficulties arise. Competition from other species may influence the species range under consideration, and the observed moisture range may not be the optimum range. Nevertheless, the results of controlled-environment experiments where moisture conditions for the maximum growth of two or more species are compared would provide useful information, and species occurrences along moisture gradients in nature probably reflect their moisture optima in a relative sense.

Controlled-environment experiments involving moisture conditions for maximum growth seem to be almost non-existent. Steinbrenner and Rediske (1964), however, did find that *Pseudotsuga menziesii* seedlings were more responsive to high levels of moisture and humidity than *Pinus ponderosa* seedlings.

A few investigators have observed species presence along measured moisture gradients in the field. Waring and Major (1964) ranked 10 northwestern species in coastal northern California by assigning "ecological optimum" numbers along a scale of 0 (dry) to 100 (wet): *Libocedrus decurrens* (on serpentine soil)--0, *Quercus kelloggii* and *Q. garryana*--10, *Pseudotsuga menziesii*--

Table 6--Comparative tolerances of northwestern tree species to excess moisture^{1/}

<i>Pinus contorta</i> , <i>Thuja plicata</i> , <i>Tsuga mertensiana</i> , <i>Chamaecyparis nootkatensis</i> , <i>Abies lasiocarpa</i> , <i>Sequoia sempervirens</i> , <i>Populus trichocarpa</i> , <i>Umbellularia californica</i>
<i>Alnus rubra</i>
<i>Pinus ponderosa</i> , <i>P. jeffreyi</i> , <i>Picea sitchensis</i> , <i>Abies grandis</i>
<i>Tsuga heterophylla</i>
<i>Pseudotsuga menziesii</i>

^{1/}Compiled from information published by Howell (1931), Brink (1954), Wood (1955), Haller (1959), Stone and Vasey (1968), Minore (1968, 1970), Krajina (1970), Minore and Smith (1971), Boggie (1974), and Crawford and Baines (1977). Moisture tolerant species are listed above intolerant ones. Where species occur in the same group, information is insufficient to separate them.

17, *Arbutus menziesii*--20, *Lithocarpus densiflora*--25, *Sequoia sempervirens*--62, *Alnus rubra*--78, *Acer macrophyllum*--85, and *Populus trichocarpa*--100. Waring (1969) also positioned major forest vegetation types of the eastern Siskiyou Mountains of southwestern Oregon along a gradient from dry (*Quercus kelloggii*) to wet (*Abies magnifica* var. *shastensis* and *Tsuga mertensiana*). Farther north, in the central western Cascade Range of Oregon, Zobel, et al. (1976) ranked tree species distribution along a plant moisture stress gradient. In order, the species are: *Arbutus menziesii* (dry), *Libocedrus decurrens* and *Pinus lambertiana*, *Acer macrophyllum* and *Pseudotsuga menziesii*, *Tsuga heterophylla* and *Thuja plicata*, *Abies grandis*, *Abies procera* and *A. amabilis*, and *Tsuga mertensiana* (wet).

Most comparisons of optimum moisture conditions among northwestern tree species consist of somewhat subjective statements based upon careful observation and field experience. Species that grow in moist conditions often are compared to *Pseudotsuga menziesii* in these statements. For example, *Tsuga heterophylla*, *Thuja plicata*, *Abies procera*, *A. amabilis*, and *Picea*

sitchensis all require more moisture than *Pseudotsuga menziesii* (see Munger 1940, Forest Soils Committee of the Douglas-fir Region 1957, and Aller 1956). Comparisons among two or three species provide more detail and help to refine these rather broad contrasts. They have been combined whenever possible and summarized in table 7.

Day (1957) suggested that *Thuja plicata* requires more moisture than *Picea sitchensis* or *Tsuga heterophylla*. Habeck (1968) observed *T. heterophylla* dominating *Thuja plicata* on well-drained slopes, *T. plicata* dominating *T. heterophylla* on poorly drained depressions and wet ravines in Glacier National Park. Wood (1955) noted that *Picea sitchensis* dominated *T. heterophylla* on moist sites but not on dry ones in southeastern Alaska. Combining the observations of these three authors, one can rank the three species in order: *Thuja plicata* (very moist), *Picea sitchensis*, and *Tsuga heterophylla* (less moist). Furthermore, the information presented by Zobel et al. (1976) indicates that *Thuja plicata* prefers slightly drier habitats than *Tsuga mertensiana*, *Abies amabilis*, and *A. procera*. Zobel and Hawk's (see footnote 2, page 5) observations suggest that

Table 7--Optimum moisture relationships among northwestern tree species^{1/}

<i>Tsuga mertensiana</i> , <i>Abies amabilis</i> , <i>A. procera</i> , <i>A. magnifica</i> var. <i>shastensis</i> , <i>A. concolor</i> , <i>Populus trichocarpa</i>
<i>Abies grandis</i>
<i>Thuja plicata</i>
<i>Picea sitchensis</i>
<i>Tsuga heterophylla</i>
<i>Acer macrophyllum</i>
<i>Alnus rubra</i>
<i>Pseudotsuga menziesii</i>
<i>Pinus ponderosa</i>
<i>Pinus lambertiana</i> , <i>Libocedrus decurrens</i>
<i>Arbutus menziesii</i>
<i>Quercus garryana</i> , <i>Quercus kelloggii</i>

^{1/}Compiled from information published by Munger (1940), Wood (1955), Day (1957), Forest Soils Committee (1957), Steinbrenner and Rediske (1964), Thilenius (1964), Aller (1956), Habeck (1968), Newton et al. (1968), Waring and Major (1964), and Zobel et al. (1976). Species with moist optimums are listed above those with dry optimums. Where species occur in the same group, information is insufficient to separate them.

Chamaecyparis lawsoniana may require more moisture than *Pseudotsuga menziesii*. Combining and comparing observations made in different areas in this way introduces an unknown amount of ecotypic variation, but species comparisons often are not possible otherwise.

Abies lasiocarpa requires moister habitats than *Picea engelmannii* (Smith and Clark 1960, Krajina 1970) or *Pinus contorta* (Wali and Krajina 1973). Further examination of the literature indicates that *P. engelmannii* requires more moisture than *P. contorta*, and *P. contorta* requires more moisture than *Pseudotsuga menziesii* or *Pinus ponderosa* (Tackle 1959). *Pinus contorta* requires less drainage than *P. ponderosa* (Kerr 1913), and it makes better growth than *P. ponderosa* on wet sites (Youngberg and Dyrness 1959). Unfortunately, information relating *Pinus contorta*, *Abies lasiocarpa*, and *Picea engelmannii* to other species is lacking.

Some information is available for northwestern hardwoods. According to the previously cited ecological optima of Waring and Major (1964), *Acer macrophyllum* prefers slightly more moisture than *Alnus rubra*. Both of these species appear to be suited to slightly more moist sites than *Pseudotsuga menziesii* (Thilenius 1964, Newton et al. 1968). *Lithocarpus densiflora* requires more moisture than *Arbutus menziesii*, *Quercus garryana*, or *Q. kelloggii* (Roy 1962).

NUTRIENTS

MACRONUTRIENT CONCENTRATIONS

Relative foliage, wood, bark, and litter concentrations of the mineral elements necessary for plant growth apparently differ among northwestern tree species. These concentrations vary somewhat with soils and season, so concentration differences may or may not reflect relative species nutrient requirements. Nevertheless, they do influence the environments in which they occur. Different species contribute differing kinds and quantities of nutrient material to their surroundings. Some of these difference have been extracted

from the literature and ranked as relative macronutrient concentrations in table 8.

Working on several soils, Daubenmire (1953) measured the nutrient content of freshly fallen foliage from 12 northern Rocky Mountain species--species that also occur in the Pacific Northwest. He suggested that the high calcium content of *Thuja plicata* litter was at least partly responsible for the higher pH of soil measured under this species. Surprisingly, the even higher calcium content of *Abies grandis* litter produced no such effect. Alban (1969) measured higher soil pH, exchangeable calcium, cation-exchange capacity, base saturation, and organic matter under old *Thuja plicata* than under old *Tsuga heterophylla* trees.

Tarrant et al. (1951) also worked on several soils when they measured nutrients in the annual litter fall of 10 northwestern tree species, but Webber (1973) studied young trees on a single soil type. He measured higher calcium contents in *Thuja plicata* than in either *Tsuga heterophylla* or *Pseudotsuga menziesii*. Webber's data, like those recorded for young seedlings by Lavender (1962) but unlike the information published by Tarrant et al. (1951), Daubenmire (1953), and Ovington (1956), show *Tsuga heterophylla* to have more calcium than *Pseudotsuga menziesii*. Soil diversity and ages of the trees, foliage, and litter sampled may be responsible for these differences. When mature trees are analyzed, calcium concentrations seem to be higher in *Pseudotsuga menziesii* than in *Tsuga heterophylla*.

Regardless of soil or age, there seems to be little doubt of the high calcium characteristic of *Thuja plicata*. Beaton, Moss et al. (1965) found calcium content to be higher in *T. plicata* foliage than in the foliage of forest grown *Abies lasiocarpa*, *A. amabilis*, *Pseudotsuga menziesii*, *Tsuga heterophylla*, *Picea sitchensis*, or *P. engelmannii*; and Ovington (1956) found both *Thuja plicata* and *Chamaecyparis lawsoniana* leaves to be richer in calcium than those of *Pseudotsuga menziesii* or *Tsuga heterophylla*. The *Thuja* and *Chamaecyparis* leaves and litter layers also had higher

Table 8--Relative macronutrient concentrations in northwestern tree species^{1/}

Nitrogen	Phosphorus	Potassium	Calcium	Magnesium	Sulfur
<i>Alnus rubra</i>	<i>Larix occidentalis</i>	<i>Populus tremuloides</i>	<i>Abies grandis</i>	<i>Picea sitchensis</i>	<i>Pseudotsuga menziesii</i>
<i>Abies amabilis</i>	<i>Populus tremuloides</i>	<i>Acer macrophyllum</i>	<i>Populus tremuloides</i>	<i>Pinus ponderosa</i> , <i>P. contorta</i> , <i>P. monticola</i>	<i>Tsuga heterophylla</i>
<i>Picea sitchensis</i>	<i>Pseudotsuga menziesii</i>	<i>Alnus rubra</i>	<i>Thuja plicata</i> , <i>Abies amabilis</i>		
<i>Acer macrophyllum</i>	<i>Alnus rubra</i>	<i>Pinus ponderosa</i>	<i>Acer macrophyllum</i>	<i>Abies amabilis</i> , <i>Tsuga heterophylla</i>	<i>Thuja plicata</i>
<i>Pinus contorta</i>	<i>Tsuga heterophylla</i> , <i>Abies lasiocarpa</i>	<i>Pseudotsuga menziesii</i> , <i>Larix occidentalis</i>	<i>Pseudotsuga menziesii</i>		
<i>Populus tremuloides</i>	<i>Abies amabilis</i> , <i>Pinus ponderosa</i> , <i>P. monticola</i> , <i>P. albicaulis</i> , <i>Abies grandis</i> , <i>Picea engelmannii</i> , <i>Acer macrophyllum</i>	<i>Abies grandis</i>	<i>Picea engelmannii</i>	<i>Thuja plicata</i>	
<i>Abies lasiocarpa</i>		<i>Thuja plicata</i>	<i>Abies lasiocarpa</i>	<i>Pseudotsuga menziesii</i>	
<i>Abies grandis</i>		<i>Pinus contorta</i>	<i>Abies amabilis</i>	<i>Acer macrophyllum</i>	
<i>Pseudotsuga menziesii</i>		<i>Tsuga heterophylla</i>	<i>Alnus rubra</i>	<i>Alnus rubra</i>	
<i>Picea engelmannii</i>		<i>Abies magnifica</i> var. <i>shastensis</i> , <i>Picea engelmannii</i>	<i>Pinus monticola</i>		
<i>Pinus ponderosa</i>		<i>Pinus monticola</i>	<i>Pinus contorta</i> , <i>Tsuga heterophylla</i>		
<i>Tsuga heterophylla</i>		<i>Pinus monticola</i>	<i>Picea sitchensis</i>		
<i>Pinus albicaulis</i>		<i>Abies amabilis</i> , <i>Picea sitchensis</i>	<i>Larix occidentalis</i>		
<i>Larix occidentalis</i>		<i>Pinus albicaulis</i>	<i>Pinus ponderosa</i>		
<i>Thuja plicata</i>			<i>Pinus albicaulis</i>		

^{1/}Compiled from foliage, wood, bark, and litter data published by Tarrant et al. (1951), Daubenmire (1953), Ovington (1956), Wollum and Youngberg (1964), Beaton, Moss et al. (1965), van den Driessche (1969), and Webber (1973). Species are ranked from high concentration (top) to low concentration (bottom) in these lists. Species in the same group are not necessarily equal, but data are insufficient for species comparisons within groups.

pH values. *Chamaecyparis lawsoniana* leaf nutrient concentrations are similar to those found in *Thuja plicata*, except for higher magnesium (Zobel and Hawk undated).

Although *Thuja plicata* sapwood has more calcium than *Pseudotsuga menziesii*, it does not differ significantly from *Tsuga heterophylla* or *Alnus rubra* with respect to sapwood calcium content (Radwan 1969). Radwan's data show that *A. rubra* sapwood has more nitrogen and magnesium, but less iron than that of *Thuja plicata*, *Tsuga heterophylla*, or *Pseudotsuga menziesii*. Relative sapwood and foliage concentrations probably differ when species are compared, for the data of Tarrant et al. (1951) show *A. rubra* foliage to have less magnesium than other species.

Tarrant et al. (1951), Daubenmire (1953), and Wollum and Youngberg (1964) all found *Pinus ponderosa* litter to be lower than *Pseudotsuga menziesii* litter in calcium and nitrogen contents. *Thuja plicata* also ranked below *P. menziesii* with respect to nitrogen concentration. Indeed, published data indicate that *T. plicata* may have the lowest nitrogen content of any northwestern tree species (Tarrant et al. 1951, Daubenmire 1953, Beaton, Moss et al. 1965, and Webber 1973). In contrast, *Alnus rubra* may be top ranked for nitrogen content (Tarrant et al. 1951, Wollum and Youngberg 1964). *Abies amabilis*, *Acer macrophyllum*, *Picea sitchensis*, and *Pinus contorta* also have high nitrogen contents. *Picea sitchensis* seedlings, though higher in nitrogen concentration, have less phosphorus than *Pseudotsuga menziesii* seedlings of similar age when grown in sand culture (van den Driessche 1969).

Larix occidentalis seems to have the highest phosphorus concentration, *Pinus contorta* the lowest (Tarrant et al. 1951, Daubenmire 1953). *Populus tremuloides* heads the list for potassium; *Pinus albicaulis* places last (Daubenmire 1953). *Thuja plicata* foliage apparently has less sulfur than the foliage of *Abies lasiocarpa*, *A. amabilis*, *Pinus contorta*, *Picea engelmannii*, *P. sitchensis*, *Pseudotsuga menziesii*, or *Tsuga heterophylla*; and *T. heterophylla* foliage seems to have slightly

less sulfur than that of *P. menziesii* (Beaton, Moss et al. 1965).

MICRONUTRIENT CONCENTRATIONS

Micronutrient levels in foliage often are influenced by "luxury consumption," and may reflect soil differences rather than species differences. When Landis (1976) compared foliage nutrient levels in "ideal" seedlings of three Rocky Mountain tree species growing in the same Colorado nursery, however, he found manganese concentrations to be higher in *Picea engelmannii* than in *Pinus contorta* or *P. ponderosa*. Zinc and boron concentrations were higher in *P. contorta* than in *P. ponderosa* or *Picea engelmannii* in these nursery seedlings. Beaton, Brown, et al. (1965), measured micronutrient concentrations in the foliage of three British Columbia conifers. They found that manganese and aluminum levels tended to be higher in *Tsuga heterophylla* needles than in those of *Pseudotsuga menziesii*, which had much more silica than *T. heterophylla*. *P. menziesii* also may have more silica (and less potassium) than *Arbutus menziesii* (Young 1974). Unfortunately, Young's analyses were based on only one tree of each species, and replicated *Pseudotsuga*-*Arbutus* comparisons are not available.

NUTRIENT DEFICIENCY TOLERANCES

Species tolerances of nutrient deficiencies or excesses and differing species growth responses to various nutrient conditions probably are of more immediate concern than species nutrient concentrations. Relative tolerances of nutrient deficiencies are listed in table 9.

Nitrogen deficiencies are best tolerated by *Picea sitchensis*, least tolerated by *Thuja plicata*, *Pinus monticola*, and *P. ponderosa* (Krajina 1970). *Pseudotsuga menziesii* var. *menziesii* also is intolerant of nitrogen deficiencies (Murison 1961). Nitrogen stimulates its shoot growth in relation to root growth but does just the opposite for *Thuja plicata*, decreasing top growth in relation to root growth (Smith et al. 1968).

Table 9--Relative tolerance of macronutrient deficiencies in northwestern tree species^{1/}

Nitrogen	Phosphorus	Potassium	Calcium	Magnesium	Sulfur
<i>Picea sitchensis</i>	<i>Pinus ponderosa</i> ,	<i>Picea engelmannii</i>	<i>Tsuga hetero-</i>	<i>Tsuga hetero-</i>	<i>Picea engelmannii</i>
<i>Tsuga heterophylla</i>	<i>P. monticola</i> ,	<i>Pseudotsuga</i>	<i>phylla</i>	<i>phylla</i>	<i>Pseudotsuga</i>
<i>Pseudotsuga</i>	<i>Tsuga hetero-</i>	<i>menziesii</i> var.	<i>Picea sitchensis</i>	<i>Pseudotsuga</i>	<i>menziesii</i> var.
<i>menziesii</i>	<i>phylla</i> , <i>Thuja</i>	<i>menziesii</i>	<i>Pseudotsuga</i>	<i>menziesii</i> var.	<i>glauca</i>
<i>Thuja plicata</i> ,	<i>plicata</i>	<i>Pinus monticola</i>	<i>menziesii</i>	<i>menziesii</i>	<i>Pinus monticola</i>
<i>Pinus monticola</i> ,	<i>Pseudotsuga</i>	<i>Pinus ponderosa</i>	<i>Pinus monticola</i> ,	<i>Pinus monticola</i>	<i>Pseudotsuga</i>
<i>P. ponderosa</i>	<i>menziesii</i> var.	<i>Pseudotsuga</i>	<i>Thuja plicata</i> ,	<i>Pseudotsuga</i>	<i>menziesii</i> var.
	<i>menziesii</i>	<i>menziesii</i> var.	<i>Chamaecyparis</i>	<i>menziesii</i> var.	<i>menziesii</i>
	<i>Pseudotsuga</i>	<i>glauca</i>	<i>nootkatensis</i> ,	<i>glauca</i>	<i>Pinus ponderosa</i>
	<i>glauca</i>		<i>Pinus ponderosa</i>	<i>Picea sitchensis</i>	
	<i>Picea sitchensis</i>			<i>Thuja plicata</i> ,	
				<i>Chamaecyparis</i>	
				<i>nootkatensis</i>	

^{1/} Most of the information used in compiling this table was obtained from Krajina (1959, 1970). Tolerant species are listed above intolerant species. Information is insufficient to separate species grouped together.

Picea sitchensis growth is stimulated more than *Pseudotsuga menziesii* growth by high nitrogen levels (van den Driessche 1968), and *P. sitchensis* should surpass *P. menziesii* on high-nitrogen, coastal sites.

Pseudotsuga menziesii is less tolerant of phosphorus deficiencies than *Pinus ponderosa*, *P. monticola*, *Tsuga heterophylla*, or *Thuja plicata* (Krajina 1970). It is more tolerant of low phosphorus than *Picea sitchensis* (van den Driessche 1968, Krajina 1970) or *P. engelmannii* (Murison and Krajina, summarized in Krajina 1970).

Pseudotsuga menziesii var. *menziesii* tolerates phosphorus, potassium, and magnesium deficiencies better than *P. menziesii* var. *glauca*, but is less tolerant of low sulfur (Krajina 1970). Krajina and his co-workers have shown *Tsuga heterophylla* to be more tolerant of calcium and magnesium deficiencies than *P. menziesii*, while *Thuja plicata* is less tolerant. Using a "replacable calcium gradient" Waring and Major (1964) ranked *Libocedrus decurrens* as more tolerant of calcium deficiencies than either *P. menziesii* or *Arbutus menziesii*.

Although comparative species responses to nutrient deficiencies apparently vary with the deficient nutrient, several authors have compared the general tolerance of two or more northwestern tree species to all nutrient deficiencies.

In general, *Tsuga heterophylla* seems to be more tolerant of soil nutrient deficiencies than *Pseudotsuga menziesii* (Krajina 1970, Heilman and Ekuan 1973), *Abies amabilis* (Krajina 1970), or *A. grandis* (Aldhous and Low 1974). Indeed, *T. heterophylla*'s soil nutrient requirements are among the lowest known for any coniferous tree (Krajina 1970). Although they often grow on ultramafic soils and are not directly compared with *T. heterophylla* in the literature, *Picea breweriana* and *Pinus jeffreyi* may be comparable to it with respect to nutrient deficiencies--they are more tolerant of low fertility than *Abies concolor*, *A. magnifica* var. *shastensis*, *Pinus lambertiana*, *Pseudotsuga menziesii*, or *Tsuga mertensiana* (Waring et al. 1975).

Pseudotsuga menziesii (Gessel et al. 1951) and *Abies amabilis* (Krajina 1970) tolerate poorer nutrient conditions than *Thuja plicata*. *Pinus lambertiana* is more sensitive than *Pseudotsuga menziesii* to poor supplies of calcium and magnesium (Krajina 1970), but it is better able to utilize infertile ultrabasic soils than *Abies magnifica* var. *shastensis* or *Tsuga mertensiana* (Waring 1970). Infertile soils also are better utilized by *Pinus contorta* than by *Abies lasiocarpa* (Wali and Krajina 1973).

NUTRIENT EXCESSES

Nutrient excesses may be just as limiting as deficiencies, but they occur less frequently. Along the Pacific Coast, *Picea sitchensis* is able to tolerate the excess magnesium in sea spray better than *Tsuga heterophylla* and other conifers (Krajina 1970). Applications of lime are tolerated by *Pseudotsuga menziesii* better than by *T. heterophylla* (Heilman and Ekuan 1973), and *Thuja plicata* is more tolerant of excess calcium than other northwestern trees (Aldhous and Low 1974). *Thuja plicata* also is more tolerant of alkaline soils than *Picea sitchensis*, *Tsuga heterophylla*, or *Pinus contorta* (Benzian 1965). Acidic soils are tolerated better by *Tsuga heterophylla* than by *Pseudotsuga menziesii* (Heilman and Ekuan 1973).

AMMONIUM VS. NITRATE NITROGEN

In addition to nutrient quantities (deficiencies, optimal amounts, or excesses), nutrient qualities and chemical formulations are important when comparing the survival and growth of tree species. Nitrogen is particularly noteworthy in this respect. Ammonium nitrogen is tolerated by seedlings of *Tsuga heterophylla* (Taylor 1935, Krajina 1970 and 1971, Madoc-Jones 1970, Heilman and Ekuan 1973) and *Pinus contorta* (Madoc-Jones 1970, Krajina 1971). Van den Driessche (1971) found that *Pseudotsuga menziesii* seedlings grew better with ammonium nitrogen than with nitrate nitrogen;

but Krajina et al. (1973) recorded contrary results...their *P. menziesii* and *Pinus contorta* seedlings grew better when supplied with nitrate nitrogen. Mycorrhizal differences may be responsible for these differing results. Without mycorrhizae, nitrates are relatively poor sources of nitrogen for *Pseudotsuga menziesii* seedlings, and growth is better with ammonium nitrate (Li et al. 1972). *Alnus rubra* grows better than *P. menziesii* without mycorrhizae or ammonium, for *A. rubra* roots are capable of reducing nitrate. *P. menziesii* roots are not (Li et al. 1972).

A mixture of both ammonium and nitrate nitrogen appears to be best for *Picea engelmannii* (Krajina 1970), *P. sitchensis* (van den Driessche 1971), *Thuja plicata* and *Tsuga heterophylla* (Krajina et al. 1973). *Abies amabilis*, *A. grandis*, and *Chamaecyparis nootkatensis* seem to prefer nitrate nitrogen.

MYCORRHIZAL RELATIONSHIPS

A review of autecological characteristics would be incomplete without further mention of mycorrhizae, for most woody plants depend upon mycorrhizal fungi for absorption of nutrients from the soil (Trappe 1977). Many species of mycorrhizal fungi are involved. Only three mycorrhizal types occur on northwestern trees, however--the vesicular-arbuscular mycorrhizae, the ectomycorrhizae + ectendomycorrhizae, and the ericoid

mycorrhizae.³ Individual trees usually have several species of mycorrhizal fungi on their roots (Trappe 1977), but these species tend to be all of the same mycorrhizal type...and different tree species support different types (table 10).

Trappe (1977) demonstrated with *Pinus ponderosa*, *Tsuga heterophylla* and *Pseudotsuga menziesii* that different host species can react differently to a given mycorrhizal fungus. In Trappe's experiment, inoculation with pure cultures of four mycorrhizal fungi increased *T. heterophylla* seedling weights, but not the weights of *P. ponderosa* or *Pseudotsuga* seedlings. Similar data on the mycorrhizal responses of other northwestern tree species have been published, but they are too scanty and variable as yet to provide much useful information. The autecological characteristics of mycorrhizal fungi need further study before they can be related to the autecological characteristics of northwestern tree species in more detail.

³James M. Trappe, Forestry Sciences Laboratory, Corvallis, Oreg. Personal communication, June 1978.



Table 10--Types of mycorrhizae associated with northwestern tree species^{1/}

Vesicular-arbuscular mycorrhizae	Ectomycorrhizae + ectendomycorrhizae	Ericoid mycorrhizae
<i>Acer macrophyllum</i>	<i>Abies amabilis</i>	<i>Arbutus menziesii</i>
<i>Acer negundo</i>	<i>Abies concolor</i>	
	<i>Abies grandis</i>	
<i>Chamaecyparis</i>	<i>Abies lasiocarpa</i>	
<i>lawsoniana</i>	<i>Abies magnifica</i>	
<i>Chamaecyparis</i>	<i>Abies magnifica</i> var.	
<i>nootkatensis</i>	<i>shastensis</i>	
	<i>Abies procera</i>	
<i>Fraxinus oregona</i>		
	<i>Alnus rubra</i>	
<i>Juniperus occidentalis</i>		
<i>Juniperus scopulorum</i>	<i>Betula papyrifera</i>	
<i>Libocedrus decurrens</i>	<i>Castanopsis chrysophylla</i>	
<i>Populus balsamifera</i>	<i>Larix laricina</i>	
<i>Populus tremuloides</i>	<i>Larix lyalli</i>	
<i>Populus trichocarpa</i>	<i>Larix occidentalis</i>	
<i>Sequoia sempervirens</i>	<i>Lithocarpus densiflora</i>	
<i>Taxus brevifolia</i>	<i>Picea breweriana</i>	
	<i>Picea engelmannii</i>	
<i>Thuja plicata</i>	<i>Picea glauca</i>	
	<i>Picea mariana</i>	
<i>Umbellularia</i>	<i>Picea sitchensis</i>	
<i>californica</i>		
	<i>Pinus albicaulis</i>	
	<i>Pinus attenuata</i>	
	<i>Pinus contorta</i>	
	<i>Pinus jeffreyi</i>	
	<i>Pinus lambertiana</i>	
	<i>Pinus monticola</i>	
	<i>Pinus ponderosa</i>	
	<i>Populus balsamifera</i>	
	<i>Populus tremuloides</i>	
	<i>Populus trichocarpa</i>	
	<i>Pseudotsuga menziesii</i>	
	<i>Quercus chrysolepis</i>	
	<i>Quercus garryana</i>	
	<i>Quercus kelloggii</i>	
	<i>Tsuga heterophylla</i>	
	<i>Tsuga mertensiana</i>	

^{1/}James M. Trappe, Forestry Sciences Laboratory, Corvallis, Oreg.
Personal communication, June 1978.

GROWTH

Light, temperature, moisture, nutrients, and other environmental factors all influence growth rates; and species growth rates should not be treated as separate entities. Growth is the result of genetic-environmental interactions that vary with both species and environment. One can compare species under given environmental conditions and derive rankings based upon species growth in those conditions. For example, Godman (1949) found that *Picea sitchensis* diameter growth outstripped that of *Tsuga heterophylla* in coastal Alaska, and Johnstone (1976) measured faster juvenile height growth of *Pinus contorta* than of *Picea glauca* in west-central Alberta. Unfortunately, such local comparisons reflect differential species reactions to particular light, temperature, moisture, and/or nutrient conditions. They may not reflect species growth rates as such, for different species often grow differently in different environments.

Several examples illustrate the interactions between growth rate and environment. Walters, Soos, and Ker (1961) observed that *Thuja plicata* seedlings reached breast height faster than *Pseudotsuga menziesii* seedlings on good sites in British Columbia. On poor sites the species ranking was reversed, with *P. menziesii* seedlings growing fastest and *T. plicata* seedlings slowest. Walters and Haddock (1966) concluded that inherent species vigor is better expressed on high quality sites than on poor sites, and they rated *Tsuga heterophylla* as faster growing than *Thuja plicata* or *Abies amabilis*. Nevertheless, growth-site interactions are confusing.

Stein (1963) planted *Pinus ponderosa*, *P. lambertiana*, *Libocedrus decurrens*, *Pseudotsuga menziesii*, and *Abies grandis* seeds from the same seed sources on three kinds of soil in southwestern Oregon. The resulting seedling height growth comparisons differed with soil type...*Libocedrus decurrens* ranked first on one soil, fourth on another after one growing season. When Deitschman and Green (1965) related

Pinus monticola site index and tree heights of several associated species in northern Idaho, they found that species height rankings varied with both site index and stand age. *Pseudotsuga menziesii* was taller than *Pinus monticola*, *Tsuga heterophylla*, or *Abies grandis* in 30-year-old stands of site index 40, shorter than these three species in 30-year-old stands of site index 80.

Clearly, ranking growth rates as such is impractical. Nevertheless, species do grow differently. Roots develop more rapidly in some than in others, seasonal shoot growth periods vary in length, mature sizes differ, and the species have different life spans. As recorded in the literature, these differences are summarized in table 11.

SEEDLING ROOT GROWTH RATE

Pinus ponderosa seedlings characteristically develop deep, sturdy root systems more rapidly than their associates (Pearson 1924, Stein 1955 and 1963, Tarrant 1953). *P. lambertiana* seedlings also develop root systems very rapidly. *Libocedrus decurrens* (Stein 1963) and *Pinus contorta* (Tarrant 1953) root development is somewhat slower.

Pseudotsuga menziesii, *Abies grandis*, *Pinus monticola*, and *Larix occidentalis* may be ranked next (see Haig et al. 1941, Stein 1955 and 1963). Seedling root growth rates in these species are faster than those of *Thuja plicata* and *Tsuga heterophylla* (Haig et al. 1941). As *Tsuga heterophylla* root penetration is poorer than that of *Thuja plicata* (Boyd 1959), it probably is also slower. *Tsuga heterophylla* probably is not the slowest species, however, for Ruth (1968a) found *Picea sitchensis* roots to be even shorter than those of *Tsuga heterophylla* after the first growing season.

Picea engelmannii seedlings produce roots more slowly than *Pseudotsuga menziesii* (Pearson 1924) or *Abies lasiocarpa* (Smith 1955) seedlings, and *Abies grandis* root growth is less than that for *P. menziesii* (Stein 1963, but

Table 11--Relative root growth rates, shoot growth periods, mature sizes, and longevity of northwestern tree species^{1/}

Seedling root growth rate ^{2/}	Shoot growth period ^{3/}	Size at maturity ^{4/}		Longevity ^{4/}
		Diameter	Height	
<i>Pinus ponderosa</i> , <i>P. lambertiana</i>	<i>Thuja plicata</i> , <i>Libocedrus decurrens</i>	<i>Sequoia semper-virens</i>	<i>Sequoia semper-virens</i>	<i>Sequoia semper-virens</i> , <i>Thuja plicata</i> , <i>Chamaecyparis nootkatensis</i>
<i>Libocedrus decurrens</i> , <i>Pinus contorta</i>	<i>Pinus ponderosa</i> , <i>P. jeffreyi</i> , <i>Chamaecyparis lawsoniana</i>	<i>Thuja plicata</i> <i>Picea sitchensis</i>	<i>Pseudotsuga menziesii</i> <i>Picea sitchensis</i>	<i>Picea sitchensis</i>
<i>Pseudotsuga menziesii</i> , <i>Pinus monticola</i> , <i>Larix occidentalis</i>	<i>Tsuga heterophylla</i> <i>Picea sitchensis</i> , <i>P. glauca</i>	<i>Pseudotsuga menziesii</i> <i>Chamaecyparis lawsoniana</i> <i>Larix occidentalis</i>	<i>Abies procera</i> <i>Tsuga heterophylla</i> <i>Thuja plicata</i>	<i>Pseudotsuga menziesii</i> <i>Larix occidentalis</i> <i>Pinus ponderosa</i>
<i>Thuja plicata</i>	<i>Sequoia sempervirens</i>	<i>Abies concolor</i> , <i>A. procera</i> , <i>Chamaecyparis nootkatensis</i>	<i>Chamaecyparis lawsoniana</i> , <i>Pinus monticola</i>	<i>Chamaecyparis lawsoniana</i> , <i>Libocedrus decurrens</i> , <i>Picea engelmannii</i> , <i>Quercus garryana</i>
<i>Tsuga heterophylla</i>	<i>Pinus contorta</i>	<i>Abies grandis</i>	<i>Abies grandis</i>	<i>Picea engelmannii</i> , <i>Quercus garryana</i>
<i>Picea sitchensis</i>	<i>Abies grandis</i>	<i>Abies magnifica</i> , <i>Pinus lambertiana</i>	<i>Abies amabilis</i> , <i>Pinus lambertiana</i>	<i>Abies amabilis</i> , <i>A. procera</i> , <i>Pinus lambertiana</i> , <i>P. monticola</i> , <i>Tsuga heterophylla</i> , <i>T. mertensiana</i>
	<i>Pseudotsuga menziesii</i>	<i>Libocedrus decurrens</i> , <i>Tsuga heterophylla</i>	<i>Larix occidentalis</i>	
	<i>Abies procera</i>	<i>Pinus monticola</i>	<i>Picea engelmannii</i>	
	<i>Abies amabilis</i>	<i>Abies grandis</i> , <i>Pinus ponderosa</i>	<i>Abies concolor</i>	<i>Abies concolor</i> , <i>A. grandis</i> , <i>A. magnifica</i> , <i>Acer macrophyllum</i>
	<i>Abies concolor</i> , <i>Pinus lambertiana</i>	<i>Abies amabilis</i>	<i>Libocedrus decurrens</i>	
		<i>Picea engelmannii</i>		
		<i>Tsuga mertensiana</i>	<i>Pinus ponderosa</i>	<i>Abies lasiocarpa</i> , <i>Pinus contorta</i>
		<i>Populus trichocarpa</i>	<i>Chamaecyparis nootkatensis</i> , <i>Alnus rubra</i>	<i>Populus trichocarpa</i>
		<i>Lithocarpus densiflora</i>	<i>Abies lasiocarpa</i> , <i>Pinus contorta</i> , <i>Tsuga mertensiana</i> , <i>Populus trichocarpa</i>	<i>Lithocarpus densiflora</i>
		<i>Quercus garryana</i>		<i>Alnus rubra</i>
		<i>Alnus rubra</i>		
		<i>Abies lasiocarpa</i>		
		<i>Pinus contorta</i> , <i>Acer macrophyllum</i>	<i>Lithocarpus densiflora</i>	
			<i>Quercus garryana</i>	

^{1/}Fast growing, large or long-lived species are at the tops of these columns; slow growing, small or short-lived species are at the bottoms. Species with long growth periods are ranked above those with short periods. Information is insufficient to separate species grouped together.

^{2/}From Pearson (1924), Haig et al. (1941), Stein (1955, 1963), Tarrant (1953), Boyd (1959), and Ruth (1968a). Rankings were based upon total root length comparisons of equal-age seedlings. Seedling ages varied from 3 months to 2 years.

^{3/}From Fowells (1941), Buckland (1956), Walters and Soos (1963), Mitchell (1965), and Williams (1968).

^{4/}From Franklin, Jerry F., and C. T. Dyrness (1973).

these species have not been compared enough with others to permit ranking in table 11.

SHOOT GROWTH PERIOD

In a given locality, some species begin shoot growth earlier in the spring than others. For example, *Pseudotsuga menziesii* breaks bud before *Quercus garryana* in the Willamette Valley of western Oregon (Thilenius 1964), and low-elevation *P. menziesii* trees begin growing several days ahead of associated *Abies grandis* trees in northeastern Oregon (Wickman 1976). Shoot growth initiation dates are extremely variable, however, both within and among species.

Relative growing season lengths apparently are more constant when species are compared...long-season species tend to have longer growing periods than short-season ones whenever they are grown together. Fowells (1941) measured the height growth periods of six northwestern species for 8 years in the Sierra Nevada of California. *Libocedrus decurrens* had the longest growing season, followed by *Pinus ponderosa* and *P. jeffreyi*, *P. contorta*, *Abies concolor*, and *Pinus lambertiana*.

Thuja plicata shoots grow for a longer period than those of *Pinus contorta*, and *P. contorta* has a longer growing season than *Tsuga heterophylla* in the Cascade Mountains of Oregon (Williams 1968). Williams observed that *Tsuga heterophylla*'s growing season exceeded those of *Abies lasiocarpa*, *A. procera*, *A. amabilis*, *Tsuga mertensiana*, and *Pseudotsuga menziesii*. Similar observations were made by Walters and Soos (1963) in southwestern British Columbia. Like Williams, they found that *Thuja plicata* had the longest growing season. *Tsuga heterophylla* had a longer season than *Pseudotsuga menziesii*, which grew for a longer period than *Pinus monticola*. Buckland (1956), also working in southwestern British Columbia, noted that *Pseudotsuga menziesii* and *Abies grandis* ceased shoot growth earlier and more abruptly than *Tsuga heterophylla* and *Thuja plicata*.

Perhaps the most extensive study of shoot growth periods was accomplished by Mitchell (1965), in England. He compared 25 conifer species for four to six seasons. *Thuja plicata* led the northwestern species for a mean length of growing season, followed in order by *Chamaecyparis lawsoniana*, *Tsuga heterophylla*, *Picea sitchensis*, *P. glauca*, *Sequoia sempervirens*, *Pinus contorta*, *Abies grandis*, *Pseudotsuga menziesii*, *Abies procera*, *A. amabilis* and *A. concolor*.

RADIAL GROWTH PERIOD

Godman and Gregory (1953) found that the radial growth of *Picea sitchensis* near Juneau, Alaska occurred over a longer period than that of *Tsuga heterophylla*. This longer radial growth period may contribute to the greater size of *P. sitchensis*, but similar growth period-tree size relationships are not consistently evident among other species. Although the radial growth period of *Alnus rubra* is shorter than the growth periods of the larger *Pseudotsuga menziesii* and *Thuja plicata* (Reukema 1965), the radial growth period of *Abies concolor* is significantly shorter than growth periods of the smaller *Pinus ponderosa* and *Libocedrus decurrens* (Fowells 1941)...and *Sequoia sempervirens* begins growth later than *Abies concolor* in northern California (Bawcom et al. 1961). Growth rate probably is more important than length of growth period in determining mature tree sizes. Sometimes these growth rates are somewhat erratic. While comparing the radial growth of *Abies grandis*, *Larix occidentalis*, *Pinus ponderosa*, and *Thuja plicata* in northern Idaho, Daubenmire (1946) recorded a second September growth spurt. This second spurt occurred only in *Thuja plicata* at low elevations, however, and Daubenmire's radial growth observations may have been affected by the shrinkage associated with drought-caused dehydration.



REPRODUCTION

VEGETATIVE REPRODUCTION

Unlike all other northwestern conifers, *Sequoia sempervirens* reproduces vegetatively by producing vigorous stump sprouts (Harlow and Harrar 1950). *Picea sitchensis*, *Tsuga heterophylla*, and *T. mertensiana* occasionally reproduce by branch-layering (Cooper 1931). *Thuja plicata*, *Chamaecyparis nootkatensis*, *Picea mariana*, *Larix laricina*, and *Abies lasiocarpa* also reproduce vegetatively by developing adventitious roots on low-hanging limbs, broken-off branches, and fallen live boles (Hustich 1953, Schmidt 1955). This vegetative regeneration is as important as reproduction from seed in high density *Thuja plicata* forests (Schmidt 1955), but it seldom occurs in *Abies lasiocarpa* (Cooper 1911). Although northwestern hardwood species (*Alnus rubra*, *Acer* spp., *Quercus* spp. etc.) also develop adventitious roots and stump sprouts, reproduction of northwestern tree species usually involves the production, dissemination, and germination of seeds.

SEED PRODUCTION

Seed production begins at younger ages in some species than in others. The species ranking in table 12 illustrates this in a general way. More detailed comparisons are available for several of these species, but they are difficult to include in an overall comparison. For example, *Abies lasiocarpa* begins to bear seed earlier in life than *Picea engelmannii* (Le Barron and Jemison 1953), but additional information that would relate these species to others in table 12 is lacking.

Additional reproductive differences become evident when seed producing trees of different species are compared. Some develop strobili earlier than others in the spring, and seed dissemination does not occur simultaneously in the fall (table 12). Studying the effects of elevation and climatic factors on production and dispersal of coniferous tree pollen on Vancouver

Island, Ebell and Schmidt (1959) found that anthesis occurred first in *Thuja plicata* and *Chamaecyparis nootkatensis*, last in *Tsuga mertensiana*. *Pseudotsuga menziesii* and *T. heterophylla* produced pollen earlier than the *Abies* species on Vancouver Island. Pollen production for *Abies amabilis*, *A. grandis*, and *A. lasiocarpa* occurs somewhat earlier than for *A. procera* in the Cascades (Franklin and Ritchie 1970).

SEED DISSEMINATION

Seed dissemination occurs during a period of several months for most species. Nevertheless, northwestern species differ rather conspicuously when initial seed dissemination dates, rates, and peak periods are compared. *Picea sitchensis* seeds mature earlier than *Tsuga heterophylla* seeds in southeastern Alaska (Harris 1969).

In the somewhat similar environments of western British Columbia, *Abies grandis* and *A. amabilis* seedfalls begin before the seedfall of *Thuja plicata* (British Columbia Forest Service 1950, Garman 1951, 1955, and Hetherington 1965). Then *Thuja plicata*, *Pseudotsuga menziesii*, and *Tsuga heterophylla* begin to drop their seeds, with *P. menziesii* slightly ahead of *T. plicata*, and *T. heterophylla* beginning last (British Columbia Forest Service 1950). *Abies grandis* seed dispersal is completed first, followed by almost all of the *Thuja plicata* seeds, then *P. menziesii*, and finally *T. heterophylla* (British Columbia Forest Service 1950, Garman 1951, 1955). A few *Thuja plicata* seeds must remain in the cones for many months, however, for Pickford (1929) observed *T. plicata* seedfall in July and August of the year following initial seed dissemination.

Farther south, *Abies amabilis* seedfall precedes that of *A. procera* in the Cascade Range (Franklin and Ritchie 1970). *Pinus monticola* seed dissemination occurs before that of *Abies grandis* in the western white pine type, and *A. grandis* seedfall is followed by *Larix occidentalis* and *P. menziesii*. The last species to disseminate seed in the western white pine type are *Thuja plicata* and *Tsuga heterophylla*.

Table 12--Comparisons of minimum seed bearing ages, anthesis order, and^{1/}
initial seed dissemination times of northwestern tree species^{2/}

Minimum seed bearing ages ^{2/}	Anthesis ^{3/}	Initial seed dissemination ^{4/}
<i>Arbutus menziesii</i>	<i>Thuja plicata</i> , <i>Chamaecyparis nootkatensis</i>	<i>Populus tremuloides</i> , <i>Populus trichocarpa</i>
<i>Pinus contorta</i>	<i>Pseudotsuga menziesii</i>	<i>Pinus monticola</i> , <i>P. lambertiana</i> , <i>P. ponderosa</i>
<i>Pinus jeffreyi</i> , <i>Alnus rubra</i> , <i>Pseudotsuga menziesii</i> var. <i>menziesii</i> , <i>Sequoia sempervirens</i> , <i>Acer macrophyllum</i> , <i>Populus trichocarpa</i>	<i>Tsuga heterophylla</i>	<i>Abies grandis</i> , <i>A. amabilis</i>
<i>Abies procera</i> , <i>Pinus ponderosa</i> , <i>Pinus monticola</i> , <i>Chamaecyparis lawsoniana</i> , <i>Populus tremuloides</i>	<i>Abies grandis</i>	<i>Pseudotsuga menziesii</i> , <i>Abies concolor</i> , <i>Larix occidentalis</i> , <i>Libocedrus decurrens</i> , <i>Pinus contorta</i> , <i>P. jeffreyi</i> , <i>Picea engelmannii</i> , <i>Chamaecyparis lawsoniana</i>
	<i>Abies amabilis</i>	
	<i>Pinus contorta</i>	
	<i>Pinus monticola</i>	
<i>Thuja plicata</i> , <i>Picea sitchensis</i> , <i>Abies grandis</i> , <i>Pseudotsuga menziesii</i> var. <i>glauca</i>	<i>Abies lasiocarpa</i>	<i>Abies magnifica</i> var. <i>shastensis</i> , <i>Thuja plicata</i>
<i>Tsuga heterophylla</i> , <i>T. mertensiana</i> , <i>Larix occidentalis</i> , <i>Pinus albicaulis</i>	<i>Abies procera</i> , <i>Tsuga mertensiana</i>	<i>Abies lasiocarpa</i> , <i>A. procera</i> , <i>Picea sitchensis</i>
		<i>Chamaecyparis nootkatensis</i> , <i>Tsuga heterophylla</i>
<i>Larix lyalli</i> , <i>Abies amabilis</i>		
<i>Quercus kelloggii</i> , <i>Picea glauca</i>		
<i>Abies magnifica</i> , <i>A. magnifica</i> var. <i>shastensis</i> , <i>A. concolor</i>		
<i>Pinus lambertiana</i>		

^{1/} The species are ranked from youngest and earliest (top) to oldest and latest (bottom). Where species are absent or occur in the same group, information is insufficient to include or separate them.

^{2/} Compiled from information obtained from For. Serv. Agric. Handb. No. 450 (1974).

^{3/} Compiled from the data of Ebell and Schmidt (1959) and Franklin and Ritchie (1970).

^{4/} Compiled from information published by Pickford (1929), Haig et al. (1941), British Columbia For. Serv. (1950), Garman (1951, 1955), Hetherington (1965), Gashwiler (1969), Harris (1969), Franklin and Ritchie (1970), For. Serv. Agric. Handb. No. 450 (1974), and Wang (1974).



(Haig et al. 1941). *Thuja plicata* and *Tsuga heterophylla* seedfall also occur later than that of *P. menziesii* in west-central Oregon (Gashwiler 1969).

Early *Abies amabilis* seed dissemination may be the results of active cone disintegration--scales literally tear themselves off the cone axis as they distort in drying. *A. grandis* and *A. lasiocarpa* cones also disintegrate, but less actively; *A. procera* cones remain intact until wind action or branch movement disturbs them (Franklin and Ritchie 1970). No obvious cone structure or modification seems to be responsible; but *Tsuga heterophylla* releases seed more slowly than most northwestern trees, regardless of location (Haig et al. 1941, British Columbia Forest Service 1950, Garman 1951 and 1955, Hetherington 1965, Gashwiler 1969, and Harris 1969).

SEED CROP SIZE

Quantity of seeds produced in a seed crop probably is more important than when those seeds are disseminated. Unfortunately, seed crop sizes vary with seasons as well as with species. As a result, observers comparing species in different years often rank species differently. Seed production also varies with locations and with tree age to further complicate species comparisons. For example, Gregory (1957, 1958, 1959) and Harris (1960, 1962) recorded larger average cone crops for *Picea sitchensis* than for *Tsuga heterophylla* during most years in Alaska. In contrast, Ruth and Berntsen (1955) found that *T. heterophylla* "...was a more prolific seed producer..." than *P. sitchensis* in coastal Oregon. Similar observer differences occur with other species. They were reconciled whenever possible in constructing table 13. Where reconciliation was unnecessary (e.g.,

Table 13--Comparative seed crop sizes and frequencies of northwestern tree species^{1/}

Seed crop size ^{2/}	Seed crop frequency ^{3/}
<i>Thuja plicata</i> , <i>Tsuga heterophylla</i> , <i>Chamaecyparis nootkatensis</i> , <i>C.</i> <i>lawsoniana</i> , <i>Picea sitchensis</i> , <i>Larix occidentalis</i>	<i>Acer macrophyllum</i> , <i>A. negundo</i> , <i>Populus trichocarpa</i>
<i>Pseudotsuga menziesii</i> , <i>Pinus</i> <i>contorta</i> , <i>P. jeffreyi</i>	<i>Pinus monticola</i> , <i>P. contorta</i>
<i>Pinus ponderosa</i>	<i>Tsuga mertensiana</i> , <i>T. heterophylla</i> , <i>Abies lasiocarpa</i> , <i>A. grandis</i> , <i>A.</i> <i>amabilis</i> , <i>Picea engelmannii</i> , <i>Thuja plicata</i>
<i>Pinus lambertiana</i>	<i>Pseudotsuga menziesii</i> , <i>Larix</i> <i>occidentalis</i>
<i>Abies concolor</i>	<i>Pinus ponderosa</i> , <i>Picea mariana</i> , <i>Populus tremuloides</i>
<i>Abies magnifica</i>	

^{1/}Species with the largest or most frequent seed crops are at the top, those with smallest or least frequent crops are at the bottom. Where species are omitted or occur in the same group, information is insufficient to include or separate them.

^{2/}Compiled from information published by Boe (1953), Garman (1955), Fowells and Schubert (1956), Gashwiler (1969), Gordon (1970), and Zobel (see footnote 2, page 5).

^{3/}Compiled from information published by Boe (1954), Franklin (1968), Franklin et al. (1974), Wang (1974), and Dahms and Barrett (1975).

Garman 1955, Gashwiler 1969), species were ranked without further comment. Where reconciliation was impractical or where sufficient information was lacking, species were discussed, but not included in table 13.

Sequoia sempervirens is a prolific seed producer. Its seed crops are larger than those of *Pinus ponderosa* (Schubert and Adams 1971). Comparing trees of similar size in northern California, Fowells and Schubert (1956) found that *Pinus ponderosa* produced more cones than *P. lambertiana*. When large trees were compared, both *Pinus* species had more cones than *Abies concolor*. Nevertheless, *Abies concolor* produces more sound seeds than *A. magnifica* (Gordon 1970). Schubert and Adams (1971) ranked *Libocedrus decurrens* last in seed production, below *A. magnifica*.

Comparisons with other species are lacking, but Dobbs (1972) found seed production in *Picea glauca* stands to be much higher than production in *P. engelmannii* stands in interior British Columbia. Southward, in northwestern Montana, Boe (1953) counted many more *Larix occidentalis* seeds than *Pseudotsuga menziesii* seeds at the edge of a mixed *Larix-Pseudotsuga* stand. Unfortunately, such stand comparisons are affected by the number of trees in each species that make up the stand. Zobel⁴ calculated numbers of seeds per square meter of basal area for each species in his sample stands. Measuring seed crops for 2 seed years, he found *Tsuga heterophylla* to be a better seed producer than *Chamaecyparis lawsoniana*, which produced more seeds than either *Abies concolor* or *Pseudotsuga menziesii* when compared on a basal area basis. When Franklin et al. (1974) compared upper slope species with respect to the number of seeds produced per tree over a 12-year period, they ranked them as follows:

Abies procera (most seeds)
Picea engelmannii
Tsuga mertensiana
Abies grandis, *A. concolor*
Abies amabilis
Abies magnifica var. *shastensis*
Pinus monticola (fewest seeds).

SEED CROP FREQUENCY

Although seasonal and location variation probably account for most of the differences in relative seed quantities recorded by different observers for the same species, variation in seed crop frequency (table 13) also must have affected their data and conclusions. Some species, *Pinus monticola* (Franklin 1968, Franklin et al. 1974) for example, produce seeds quite regularly. Others, like *Thuja plicata* (Gashwiler 1969), are erratic and produce a large crop one year and no crop at all for the next year or two.

Although most species seem to be rather intermediate with respect to seed crop frequency, *Tsuga heterophylla* tends to be more frequent than *Picea sitchensis* (Meyer 1937). In Montana, *Pinus contorta* seed crops are more frequent than *Larix occidentalis* crops, which are more frequent than those of *P. ponderosa* (Boe 1954). *Pinus contorta* seed crops also are more frequent than *P. ponderosa* crops in Oregon (Dahms and Barrett 1975). In interior British Columbia, where *Picea engelmannii* and *P. glauca* hybridize, good *P. engelmannii* seed crops are more frequent than good *P. glauca* crops (Dobbs 1972). Wang (1974) published a general summary of seed crop frequencies for 21 northwestern tree species.

SEED SOUNDNESS

Efficacy of any seed crop, regardless of its frequency, is dependent upon sound seeds in that crop. Few species comparisons are available; but *Pinus lambertiana*, *P. ponderosa*, and *Libocedrus decurrens* seed crops have higher proportions of sound seed than *Abies concolor* crops (Fowells and

⁴Zobel, Donald B. Seed production in forests of *Chamaecyparis lawsoniana*. Unpublished manuscript.

Schubert 1956). Gashwiler (1969) found more filled seed in *Thuja plicata* than in *Tsuga heterophylla* or *Pseudotsuga menziesii* during the 12 years he monitored seed fall in west-central Oregon.

SEED FLIGHT

Several important characteristics of seed fall (speed and dispersal distance, for example) are functions of seed weight and structure. Average seed weights of northwestern species are listed in table 14. Seed structures vary greatly. Rather than comparing them as such, several investigators have measured seed flight to compare species seed distribution directly.

Siggins (1933) seems to have done the definitive work in seed flight measurements. After dropping seeds of 13 western conifers down an elevator shaft in the campanile, University of California, Berkeley, he concluded that *Tsuga heterophylla* seeds fell the slowest, *Pinus lambertiana* seeds the fastest. In a less-precise experiment, Isaac (1930) released seeds from a box kite and a pilot balloon under varying wind conditions. He measured the flights of *Tsuga heterophylla*, *Pinus monticola*, *Pseudotsuga menziesii*, *Abies procera*, *Thuja plicata*, and *Pinus ponderosa* seeds over snow covered fields and found them to be in the order given--*Tsuga heterophylla* seeds flying farthest, *Thuja plicata* and *Pinus ponderosa* seeds landing closest to the release point. Pickford (1929) and Gashwiler (1969) also concluded that *Tsuga heterophylla* seeds fly farther than *Thuja plicata* seeds.

In British Columbia and south-eastern Alaska, *Tsuga heterophylla* seed flights may be shorter than the flights of *Thuja plicata* (Garman 1951) or *Picea sitchensis* (Godman 1953b) seeds. These northern conclusions were based upon indirect observations rather than direct measurements, however, and unequal consumption by animals or different tree heights and densities may have influenced the seed numbers counted on the ground.

Using seed trap transects extending outward from clearcut edges, Carkin et al. (1978) found that *Abies procera* seed fall declined more rapidly than that of *A. amabilis* in the Washington and Oregon Cascade Range. *Picea engelmannii* seeds fly further than *P. glauca* seeds (Dobbs 1972). Other comparisons are summarized in table 14.

SEED DURABILITY AND LONGEVITY

Both long and short seed flights separate seeds from their parent cones and expose them to environmental stresses that must be endured until germination is accomplished. *Pinus monticola* (Hofmann 1925), *P. contorta*, and *P. ponderosa* (Wright 1931) seeds are more resistant to high temperatures than *Pseudotsuga menziesii* seeds. When seeds are stored for long periods under cooler temperature conditions, the *Pinus* species again seem to be more resistant and long-lived (Isaac 1940 and 1943, Schubert 1954, Holmes and Buszewicz 1958). Other species longevity comparisons are summarized in table 15.

SEED GERMINATION

Given similar environmental conditions, *Pseudotsuga menziesii* seeds usually are easier to germinate than *Pinus monticola* or *P. lambertiana* seeds (Allen and Bientjes 1954). Similar differences have been noted by other workers, using other species. For example, Dobbs (1972) found *Picea engelmannii* seeds to have "...on the average, higher germinative energy and higher germinative capacity..." than *P. glauca* seeds. Germination rates and stratification requirements are compared in table 16.

Species germination rates have been compared more than other germination attributes. *Abies procera* seeds germinated faster than *A. magnifica* var. *shastensis* seeds when covered with 1/2 inch (1.3 cm) of mineral soil, and both of these *Abies* species germinated before adjacent *Pinus*

Table 14--Comparative seed weights and seed flight distances of
northwestern tree species

Seed weights ^{1/}	Seed flight distances ^{2/}
<i>Populus tremuloides</i>	<i>Tsuga heterophylla</i>
<i>Alnus rubra</i>	<i>Pinus contorta</i>
<i>Thuja plicata</i>	<i>Picea sitchensis</i>
<i>Tsuga heterophylla</i>	<i>Pseudotsuga menziesii</i>
<i>Arbutus menziesii</i>	<i>Abies amabilis</i>
<i>Picea glauca</i>	<i>Pinus ponderosa</i>
<i>Picea sitchensis</i> , <i>Chamaecyparis</i> <i>lawsoniana</i>	<i>Thuja plicata</i>
<i>Larix lyalli</i>	<i>Abies grandis</i> , <i>A. concolor</i>
<i>Larix occidentalis</i>	<i>Libocedrus decurrens</i>
<i>Picea engelmannii</i>	<i>Pinus jeffreyi</i>
<i>Sequoia sempervirens</i>	<i>Sequoia sempervirens</i>
<i>Tsuga mertensiana</i>	<i>Pinus lambertiana</i>
<i>Chamaecyparis nootkatensis</i>	
<i>Pinus contorta</i>	
<i>Picea breweriana</i>	
<i>Pseudotsuga menziesii</i>	
<i>Abies lasiocarpa</i>	
<i>Pinus monticola</i>	
<i>Abies grandis</i>	
<i>Libocedrus decurrens</i>	
<i>Taxus brevifolia</i>	
<i>Abies procera</i>	
<i>Pinus ponderosa</i>	
<i>Abies concolor</i> , <i>A. amabilis</i>	
<i>Abies magnifica</i> var. <i>shastensis</i>	
<i>Abies magnifica</i>	
<i>Pinus jeffreyi</i> , <i>Acer macrophyllum</i>	
<i>Pinus albicaulis</i> , <i>P. lambertiana</i>	
<i>Lithocarpus densiflora</i> , <i>Quercus</i> <i>garryana</i> , <i>Q. kelloggi</i>	

^{1/}Derived from data published in For. Serv. Agric. Handb. No. 450 (1974). Species with light seeds are listed above those with heavy seeds. Species on the same line have equal seed weights.

^{2/}Based upon the seedfall measurements of Siggins (1933). Species with long seed flights are listed above those with short flights. Species on the same line have the same flight characteristics. Where species are omitted, data are insufficient for comparison.

monticola seeds in the Cascade Range (Franklin and Hoffman 1968). Further differences are present among both *Abies* and *Pinus* species. Davidson (1959) found *Abies grandis* germination to be faster than that of *A. amabilis*, and Lanquist (1946) observed faster germination in *A. concolor* than in *A. magnifica*. Lanquist's data also showed that *Pinus jeffreyi* seeds germinated faster than those of *P. ponderosa*,

which were far ahead of *P. lambertiana* seeds. His data show *Abies concolor* to germinate slightly faster than *Libocedrus decurrens*--a finding that conflicts with an earlier statement by Show (1930) attributing greater rapidity to the *L. decurrens* germination. As Lanquist sowed the *Abies* and *Libocedrus* seeds at different depths, Show's statement probably is correct. *Pseudotsuga menziesii*

Table 15--Comparative seed longevity of northwestern tree species^{1/}

Pinus ponderosa
Pinus contorta
Pinus jeffreyi
Pinus lambertiana
Pinus monticola
Pseudotsuga menziesii, *Thuja plicata*, *Chamaecyparis lawsoniana*, *Tsuga heterophylla*, *Picea sitchensis*
Larix occidentalis
Pinus albicaulis
Abies procera, *Tsuga mertensiana*, *Picea engelmannii*
Abies concolor, *A. grandis*, *A. lasiocarpa*, *A. magnifica*, *Libocedrus decurrens*, *Sequoia sempervirens*

^{1/}Compiled from data published by Schubert (1954) and Holmes and Buszewicz (1958). Species with more longevity are listed above those with less. Information is insufficient to separate species listed in the same group.

Table 16--Comparative germination attributes of northwestern tree species^{1/}

Germination rate ^{2/}	Stratification required ^{3/}
<i>Pinus jeffreyi</i> , <i>P. ponderosa</i> , <i>Abies procera</i> <i>Pseudotsuga menziesii</i> , <i>Abies grandis</i> , <i>A. concolor</i> , <i>A. magnifica</i> var. <i>shastensis</i> , <i>Libocedrus decurrens</i> <i>Pinus lambertiana</i> , <i>P. monticola</i> , <i>Picea engelmannii</i>	<i>Picea sitchensis</i> , <i>P. engelmannii</i> , <i>P. glauca</i> , <i>Thuja plicata</i> , <i>Chamaecyparis lawsoniana</i> , <i>Tsuga heterophylla</i> , <i>Pinus ponderosa</i> , <i>P. jeffreyi</i> <i>Abies magnifica</i> var. <i>shastensis</i> , <i>A. grandis</i> , <i>A. concolor</i> , <i>A. procera</i> , <i>A. lasiocarpa</i> , <i>A. amabilis</i> , <i>Pseudotsuga menziesii</i> , <i>Pinus contorta</i> <i>Pinus lambertiana</i> , <i>P. monticola</i> , <i>Acer</i> species, <i>Taxus brevifolia</i>

^{1/}Species germinating faster and those requiring less stratification are listed above slower, stratification-dependent species. Where species occur in the same group, information is insufficient for further separation.

^{2/}From Lanquist (1946), Murison (1959), Stein (1963), and Franklin and Hoffman (1968).

^{3/}From Heit (1968), Kotar (1972), and For. Serv. Agric. Handb. No. 450 (1974).

germination is slower than that of *Pinus ponderosa* (Lanquist 1946, Murison 1959), faster than that of *Tsuga heterophylla* (Gashwiler 1969) or *Pinus lambertiana* (Lanquist

1946, Murison 1959, Stein 1963). Murison (1959) found *Pinus ponderosa* seed germination to be faster than *Pseudotsuga menziesii* var. *glauca*, which was faster than *P. menziesii*

var. *menziesii*. Both *Pseudotsuga* varieties germinated faster than *Pinus lambertiana* after a 6-week stratification period. The earliest germination of Murison's stratified *Picea engelmannii* seeds occurred slightly later than that of his stratified *Pinus monticola*.

SEED STRATIFICATION REQUIREMENTS

Seed stratification is essential for the prompt germination of some northwestern tree species, unnecessary for others. Stratification requirements appear to be highest for *Pinus lambertiana*, *P. monticola*, *Taxus brevifolia*, and *Acer* species. *Abies grandis*, *A. procera*, *Pseudotsuga menziesii*, *Picea breweriana*, and stored *Pinus ponderosa* seeds also benefit from a moist, cool pregermination treatment; but *Thuja plicata*, *Picea engelmannii*, *P. sitchensis*, and fresh *Pinus ponderosa* seeds do not require prechilling (Heit 1968, Forest Service, USDA 1974). Although *Abies amabilis* germination occurs faster after stratification for 8 weeks at 4°C, *Tsuga heterophylla* germination is retarded by this treatment (Kotar 1972).

GERMINATION ENVIRONMENTS

Soil attributes have been related to the germination of several northwestern tree species. *Thuja plicata* germination is best in alkaline soil; but *Tsuga heterophylla* germination is depressed there, reaching its optimum under acid conditions (Thrupp 1939). Thrupp, working under poorly controlled greenhouse conditions, concluded that magnesium retards *Picea sitchensis* and *Thuja plicata* germination but accelerates the germination of *Pseudotsuga menziesii* and *Tsuga heterophylla*. The germination of *P. menziesii* is not as sensitive to soil moisture stress as that of *Picea engelmannii* (Brayshaw 1970). Soil moisture may have influenced the results published by Garman (1955). He noted that maximum germination of *Tsuga heterophylla*, *Picea sitchensis*, and *Thuja plicata* occurred in 75-percent shade, whereas maximum

Pseudotsuga menziesii germination occurred in only 25-percent shade. Moisture's importance certainly was demonstrated by Ackerman (1957), who sowed *Picea glauca* and *Pinus contorta* seeds on both trenched and untrenched seedbeds in Alberta. Where the trenching removed root competition of the residual stand, *P. contorta* germination and survival were benefited but *P. glauca* was not affected.

SEEDBEDS

Seed germination and subsequent seedling survival are greatly affected by seedbed conditions. Although good seedbeds (e.g., mineral soil) tend to be favorable for all northwestern tree species, some species apparently are able to utilize good seedbeds more efficiently than others. Conversely, some species are able to tolerate poor seedbeds better than others. These general seedbed-species relationships are shown for several northwestern trees in table 17.

Lack of available moisture probably contributes to the poor seedbed characteristics of moss, litter, and duff. Surface moss dries out rapidly. *Tsuga heterophylla* seeds apparently germinate better than *Thuja plicata* or *Picea sitchensis* seeds in this moss (Godman 1953a, Phelps 1973)...and the *T. plicata* survival is better than that of either *Tsuga* or *Picea* in southeastern Alaska (Godman 1953a).

Moisture associated with shaded conditions probably influenced the germination and survival of Sierra Nevada conifers on the seedbeds subjectively compared by Stark (1965). Stark concluded that *Libocedrus decurrens* germinated best on bare soil in half shade, but *Pinus lambertiana* germination was best on light litter in full sunlight. Survival of both species was judged to be highest where litter occurred.

In greenhouse and laboratory studies, seedbed differences persist even when moisture differences are minimized. An ash seedbed stimulated *Pseudotsuga menziesii* germination but reduced the germination of *Tsuga heterophylla* and *Thuja plicata* when Jablanczy (1963) compared ash

Table 17--Comparative seedbed suitability of northwestern tree species^{1/}

Organic seedbeds		Mineral soil seedbeds		Burned seedbeds
Coastal	Interior	Coastal	Interior	
<i>Tsuga heterophylla</i>	<i>Picea glauca</i> ,	<i>Alnus rubra</i>	<i>Larix occidentalis</i> ,	<i>Pseudotsuga</i>
<i>Thuja plicata</i>	<i>Pseudotsuga</i>	<i>Picea sitchensis</i>	<i>Picea</i>	<i>menziesii</i> , <i>Abies</i>
<i>Picea sitchensis</i>	<i>menziesii</i>	<i>Thuja plicata</i>	<i>engelmannii</i>	<i>grandis</i> , <i>Pinus</i>
<i>Alnus rubra</i>	<i>Abies lasiocarpa</i> ,	<i>Tsuga heterophylla</i>	<i>Pinus contorta</i> ,	<i>ponderosa</i>
	<i>Pinus contorta</i>		<i>Abies lasiocarpa</i>	<i>Tsuga heterophylla</i> ,
	<i>Larix occidentalis</i> ,		<i>Pseudotsuga</i>	<i>Larix occidentalis</i> ,
	<i>Picea</i>		<i>menziesii</i> , <i>Picea</i>	<i>Picea engelmannii</i> ,
	<i>engelmannii</i>		<i>glauca</i>	<i>Pinus contorta</i>
				<i>Thuja plicata</i> , <i>Pinus</i>
				<i>monticola</i> , <i>Abies</i>
				<i>lasiocarpa</i>

^{1/} Compiled from information published by Fisher (1935), Taylor (1935), Godman (1953a), Bever (1954), Muri (1955), Smith (1955), Day (1957), Smith and Clark (1960), Jablanczy (1963), Soos and Walters (1963), Ruth (1968b), Boyd and Deitschman (1969), Schmidt (1969), Phelps (1973), and Ryker (1975). Species in upper groups are better suited to the seedbed than those in the lower groups. Data are insufficient for species comparisons within groups.

Table 18--Comparative physical characteristics

Cotyledon number ^{2/}	Needle longevity ^{3/}	Needle volume ^{3/}	Needle weight ^{3/}	Total crown weight ^{4/}
<i>Pinus lambertiana</i>	<i>Abies grandis</i>	<i>Pinus monticola</i>	<i>Pinus contorta</i>	<i>Pinus ponderosa</i>
<i>Pinus jeffreyi</i>	<i>Picea sitchensis</i>	<i>Pinus contorta</i>	<i>Pinus monticola</i>	<i>Abies grandis</i> , <i>A.</i> <i>concolor</i> , <i>A.</i> <i>lasiocarpa</i> ,
<i>Pinus ponderosa</i>	<i>Abies amabilis</i>	<i>Abies grandis</i>	<i>Abies grandis</i>	<i>Pseudotsuga</i> <i>menziesii</i> ,
<i>Pinus monticola</i>	<i>Pseudotsuga</i> <i>menziesii</i>	<i>Picea sitchensis</i>	<i>Picea sitchensis</i>	<i>Picea</i> <i>engelmannii</i>
<i>Pinus attenuata</i> , <i>Pseudotsuga</i> <i>menziesii</i> , <i>Abies</i> <i>magnifica</i> var. <i>shastensis</i> , <i>A.</i> <i>concolor</i>	<i>Pinus monticola</i>	<i>Pseudotsuga</i> <i>menziesii</i>	<i>Abies amabilis</i>	
<i>Picea engelmannii</i>	<i>Tsuga heterophylla</i>	<i>Tsuga</i> <i>heterophylla</i>	<i>Pseudotsuga</i> <i>menziesii</i>	<i>Tsuga heterophylla</i> ,
<i>Abies amabilis</i> , <i>A.</i> <i>grandis</i> , <i>Larix</i> <i>occidentalis</i> , <i>Picea breweriana</i>	<i>Pinus ponderosa</i>		<i>Tsuga</i> <i>heterophylla</i>	<i>Thuja plicata</i> ,
	<i>Pinus contorta</i>			<i>Pinus contorta</i> ,
				<i>P. albicaulis</i> ,
				<i>P. lambertiana</i>
				<i>Larix occidentalis</i> ,
				<i>Pinus monticola</i> ,
				<i>Libocedrus</i> <i>decurrens</i>
<i>Pinus contorta</i>				
<i>Picea sitchensis</i> ,				
<i>Abies procera</i>				
<i>Abies lasiocarpa</i>				
<i>Tsuga mertensiana</i>				
<i>Tsuga heterophylla</i>				
<i>Chamaecyparis</i> <i>nootkatensis</i> ,				
<i>C. lawsoniana</i> ,				
<i>Thuja plicata</i> ,				
<i>Taxus brevifolia</i> ,				
<i>Libocedrus</i> <i>decurrens</i>				

^{1/}Species in the same group are not necessarily equal, but data are insufficient for species comparisons within groups.

^{2/}Compiled from data published by Franklin (1961). Species with many cotyledons are listed above those with few.

^{3/}Based on data published by Smith (1970). Species with the largest, heaviest, and most persistent needles are listed above those with smaller, lighter, and shorter-lived needles.

^{4/}From Chandler (1960) and Brown (1976). Species with heavy crowns are listed above those with light crowns.

of northwestern tree species^{1/} (continued)

Crown length ^{5/}	Crown width ^{6/}	Crown density ^{7/}	Annual litter fall ^{8/}	Heartwood specific gravity ^{9/}
<i>Picea engelmannii</i> , <i>Abies lasiocarpa</i>	<i>Pinus ponderosa</i> , <i>Tsuga hetero- phylla</i>	<i>Pinus albicaulis</i> <i>Abies lasiocarpa</i>	<i>Thuja plicata</i> <i>Pseudotsuga menziesii</i> (350 years old)	<i>Quercus garryana</i> <i>Taxus brevifolia</i>
<i>Tsuga heterophylla</i> , <i>T. mertensiana</i> , <i>Thuja plicata</i> , <i>Abies grandis</i> , <i>A. concolor</i>	<i>Thuja plicata</i> <i>Pseudotsuga menziesii</i>	<i>Picea engelmannii</i> <i>Abies grandis</i> , <i>Tsuga hetero- phylla</i> , <i>Pseudotsuga menziesii</i> , <i>Pinus</i>	<i>Abies amabilis</i> <i>Acer macrophyllum</i> <i>Pinus monticola</i>	<i>Larix occidentalis</i> <i>Fraxinus oregona</i> <i>Pseudotsuga menziesii</i> var. <i>menziesii</i>
<i>Picea sitchensis</i> , <i>Pinus ponderosa</i> , <i>P. contorta</i> , <i>Pseudotsuga menziesii</i> , <i>Abies procera</i> , <i>Pinus monticola</i>	<i>Abies grandis</i> <i>Picea engelmannii</i> <i>Pinus monticola</i> , <i>P. contorta</i> , <i>Larix occidentalis</i>	<i>ponderosa</i> , <i>P. contorta</i> , <i>P. monticola</i> <i>Larix occidentalis</i>	<i>Alnus rubra</i> <i>Tsuga heterophylla</i> <i>Pseudotsuga menziesii</i> (100 years old)	<i>Acer macrophyllum</i> <i>Tsuga mertensiana</i> <i>Pseudotsuga menziesii</i> var. <i>glauca</i> , <i>Sequoia sempervirens</i> , <i>Chamaecyparis lawsoniana</i>
<i>Larix occidentalis</i>		<i>Thuja plicata</i>	<i>Picea sitchensis</i> <i>Pinus ponderosa</i> <i>Pinus contorta</i>	<i>Tsuga heterophylla</i> , <i>Pinus contorta</i> , <i>P. ponderosa</i> <i>Picea sitchensis</i> , <i>P. glauca</i> , <i>Pinus jeffreyi</i> , <i>P. monticola</i> , <i>P. lambertiana</i> , <i>Abies magnifica</i> , <i>A. concolor</i> , <i>A. grandis</i> , <i>A. procera</i> , <i>A. amabilis</i> , <i>Alnus rubra</i> , <i>Populus tremuloides</i> , <i>Libocedrus decurrens</i> <i>Populus trichocarpa</i> , <i>Picea engelmannii</i> , <i>Abies lasiocarpa</i> , <i>Thuja plicata</i>

^{5/}From Flint (1925), Starker (1934), and Fahnestock (1960). Species with long crowns are listed above those with shorter crowns.

^{6/}From Fahnestock (1960). Species with wide crowns are listed above those with narrow crowns.

^{7/}From Brown's (1978) measurements of foliage plus branchwood bulk densities in dominant trees. Species with dense crowns are listed above those with less-dense crowns.

^{8/}From Tarrant et al. (1951). The heaviest litter fall is listed at the top, the lightest at the bottom.

^{9/}Compiled from data published by Markwardt (1931), Markwardt and Wilson (1935), USDA Forest Products Laboratory (1955, 1965), Born (1966), and Farr (1973). Species with high specific gravities are listed above those with low specific gravities.

Table 18--Comparative physical characteristics

Branchwood specific gravity ^{10/}	Sapwood thickness ^{11/}	Bark thickness ^{12/}	Foliage flammability ^{13/}	Epiphyte receptivity ^{14/}
<i>Pseudotsuga menziesii</i>	<i>Pinus ponderosa</i>	<i>Larix</i>	<i>Picea engelmannii</i> ,	<i>Alnus rubra</i>
<i>Pinus contorta</i>	<i>Pinus contorta</i>	<i>occidentalis</i> ,	<i>P. sitchensis</i> ,	<i>Picea sitchensis</i> ,
<i>Abies grandis</i> ,	<i>Pseudotsuga menziesii</i> ,	<i>Pseudotsuga menziesii</i>	<i>Pseudotsuga menziesii</i> , <i>Thuja</i>	<i>Abies procera</i>
<i>Larix occidentalis</i>	<i>Picea engelmannii</i>	<i>Pinus ponderosa</i>	<i>plicata</i> , <i>Tsuga heterophylla</i> ,	<i>Tsuga heterophylla</i> , <i>Picea engelmannii</i>
	<i>Thuja plicata</i>	<i>Abies grandis</i> ,	<i>T. mertensiana</i> ,	
	<i>Larix occidentalis</i>	<i>A. concolor</i> ,	<i>Abies procera</i> ,	<i>Pseudotsuga menziesii</i> , <i>Larix occidentalis</i>
		<i>A. procera</i>	<i>A. lasiocarpa</i>	<i>Pinus monticola</i>
		<i>Pinus monticola</i> ,	<i>Abies grandis</i> , <i>A. concolor</i> , <i>Pinus monticola</i>	<i>Thuja plicata</i>
		<i>Tsuga heterophylla</i> , <i>T. mertensiana</i>	<i>Pinus contorta</i>	<i>Populus trichocarpa</i>
		<i>Thuja plicata</i>	<i>Larix occidentalis</i> ,	<i>Acer macrophyllum</i>
		<i>Picea sitchensis</i> ,	<i>Pinus ponderosa</i>	<i>Chamaecyparis nootkatensis</i> ,
		<i>P. engelmannii</i> ,		<i>Taxus brevifolia</i>
		<i>Pinus contorta</i> ,		<i>Abies lasiocarpa</i>
		<i>Abies lasiocarpa</i> ,		<i>Abies grandis</i> ,
		<i>Chamaecyparis nootkatensis</i>		<i>A. concolor</i>
				<i>Populus tremuloides</i>
				<i>Pinus contorta</i>
				<i>Quercus garryana</i>
				<i>Arbutus menziesii</i>
				<i>Tsuga mertensiana</i>

^{10/} From Ryan and Pickford (1978). Species with high specific gravities are listed above those with low specific gravities.

^{11/} From Lassen and Okkonen (1969). Species with thick sapwood are listed above those with thin sapwood.

^{12/} Compiled from information published by Flint (1925), Starker (1934), and Bones (1962). Species with thick bark are listed above those with thin bark.

^{13/} Compiled from information published by Flint (1925) and Starker (1934). Species with highly flammable foliage are listed above those with less flammable foliage.

^{14/} Compiled from information published by Flint (1925), Starker (1934), Coleman et al. (1956), and Pechanec and Franklin (1968). Species supporting more epiphytes are listed above those supporting less.

of northwestern tree species^{1/}

Fire resistance ^{15/}		Rooting depth ^{16/}	Windthrow resistance ^{17/}	Growth in dense soils ^{18/}	Snow damage resistance ^{19/}
Coastal	Interior				
<i>Pseudotsuga menziesii</i>	<i>Larix occidentalis</i>	<i>Pinus ponderosa</i>	<i>Pseudotsuga menziesii</i> , <i>Thuja plicata</i>	<i>Larix occidentalis</i> , <i>Pinus monticola</i> , <i>P. ponderosa</i> , <i>P. lambertiana</i>	<i>Tsuga mertensiana</i> , <i>T. heterophylla</i> , <i>Picea breweriana</i> , <i>Abies magnifica</i> var. <i>shastensis</i>
<i>Abies grandis</i> , <i>A. concolor</i>	<i>Pinus ponderosa</i> , <i>Pseudotsuga menziesii</i>	<i>Pinus lambertiana</i> , <i>Libocedrus decurrens</i>	<i>Picea sitchensis</i>	<i>Pseudotsuga menziesii</i> , <i>Pinus contorta</i>	<i>Pinus monticola</i> , <i>Abies amabilis</i>
<i>Tsuga mertensiana</i>	<i>Pinus lambertiana</i> , <i>Abies grandis</i> , <i>A. concolor</i>	<i>Pseudotsuga menziesii</i> , <i>Abies grandis</i> , <i>Pinus monticola</i> , <i>Larix occidentalis</i>	<i>Tsuga heterophylla</i> <i>Abies amabilis</i>	<i>Abies amabilis</i> <i>Picea sitchensis</i> , <i>Tsuga heterophylla</i> , <i>Thuja plicata</i>	<i>Abies procera</i> , <i>A. concolor</i> , <i>Pinus jeffreyi</i>
<i>Abies procera</i>	<i>Libocedrus decurrens</i> , <i>Pinus monticola</i> , <i>Thuja plicata</i> , <i>Tsuga mertensiana</i>	<i>Thuja plicata</i> <i>Tsuga heterophylla</i>			<i>Pseudotsuga menziesii</i> , <i>Pinus lambertiana</i> , <i>P. ponderosa</i>
<i>Pinus monticola</i>	<i>Pinus contorta</i> , <i>Picea engelmannii</i> , <i>Tsuga heterophylla</i>				
<i>Pinus contorta</i>	<i>Abies lasiocarpa</i>				

^{15/}Based upon observations by Kerr (1913), Flint (1925), Starker (1934), Boyd (1959), Tackle (1959), and Wallis et al. (1974). Fire resistant species are listed above less resistant species.

^{16/}From Haig (1936) and Stein (1963). Deep-rooted species are listed above shallow-rooted species.

^{17/}From Boyce (1929), Ruth and Yoder (1953), and Gratkowski (1956). Resistant species are listed above non-resistant species.

^{18/}From Leaphart (1958), Stephens (1965), and Minore et al. (1969). Species capable of growing in dense soils are listed above those that are not.

^{19/}From Williams (1966), Kangur (1973), and Waring et al. (1975). Resistant species are listed above non-resistant species.

Conflicting references are not a problem with crown widths and crown densities, for there seem to be very few sources of this information. Fahnestock (1960) described *Pinus ponderosa* and *Tsuga heterophylla* crowns as "wide;" *Pinus monticola*, *P. contorta*, and *Larix occidentalis* crowns as "narrow." His intermediate descriptions were used in ranking crown widths in table 18. Brown's (1978) measurements of foliage and branchwood bulk densities were used in ranking crown densities. Although he did not rank species in terms of crown width or density, Gratkowski (1956) noted that *Thuja plicata* crowns were less dense than the crowns of *Tsuga heterophylla* and *Abies amabilis*. Fahnestock (1960) observed that the quantity of foliage plus very fine twigs in individual crowns seems to be related to species shade tolerance--the higher the percentage of total crown weight in foliage, the more tolerant is the species.

LITTER FALL

Large, dense crowns do not necessarily produce large quantities of litter. When Tarrant et al. (1951) measured the annual litter fall in pure, well-stocked stands of 10 northwestern tree species, they found that *Thuja plicata* (a species with moderate crown size and density) produced the most litter. A 350-year-old stand of *Pseudotsuga menziesii* dropped more than twice as much litter as a 100-year-old stand, and age differences may have influenced the data published by Tarrant et al. Nevertheless, their general comparison emphasizes the large differences between species...*Thuja plicata* produced almost eight times as much litter as *Pinus contorta*.

WOOD

Although they may not vary annually like litter fall, wood properties of the various species are not at all constant. They vary with latitude, age, and position in the tree. Nevertheless, differences

in wood characteristics among species are of great importance and have been studied extensively. Most of these studies will not be reviewed here. Only two wood properties have been selected for comparison--heartwood specific gravity and sapwood thickness.

Specific gravity values were compared to obtain the species ranking shown in table 18. Not shown are species differences in the way wood specific gravity changes with height in the tree. Okkonen et al. (1972) found that specific gravity decreased as height increased in *Pseudotsuga menziesii*, *Pinus ponderosa*, *Larix occidentalis*, *Abies grandis*, *A. procera*, *A. magnifica*, and *A. amabilis*. It increased with height in *Thuja plicata*, *Populus trichocarpa*, and *Picea engelmannii*. The specific gravity of *Abies concolor* initially decreased, then increased farther up the tree. Progressing outward from pith to bark, tracheid length tends to increase in all conifers--but Anderson's (1951) four-tree sample indicated that this increase is greater in *Pseudotsuga menziesii* than in *Abies concolor* or *A. procera*.

Sapwood thickness increases with increasing tree diameter in western softwoods (Lassen and Okkonen 1969). When Grier and Waring (1974) compared sapwood cross sectional area and foliage mass, they found them to be highly correlated. Furthermore, the correlations varied consistently among the three species investigated. *Pseudotsuga menziesii* had more foliage per unit sapwood than either *Abies procera* or *Pinus ponderosa*. In small trees, *P. ponderosa* had more foliage per unit sapwood than *A. procera*, but in large trees *A. procera* had more foliage for its sapwood than *P. ponderosa*.

The pattern of water conduction in sapwood and heartwood differs among northwestern conifers. It occurs in an ascending right spiral in *Pinus contorta*, *P. ponderosa*, *P. jeffreyi*, *Larix* species, *Picea* species, and *Abies* species (Rudinsky and Vite' 1959). In *Pinus monticola* and *P. lambertiana*, conduction occurs in an ascending left spiral. Rudinsky and Vite' discovered an interlocking zigzag

ascent in *Libocedrus decurrens*, *Sequoia sempervirens*, and *Juniperus* species. In *Pseudotsuga menziesii* and *Tsuga* species, the ascent of water is neither in a complete spiral nor along an interlocking pathway--it winds around the stem in a coherent section. The coherent section does not wind in *Thuja plicata* and *Chamaecyparis lawsoniana*, but goes straight up the tree. Pattern of water conduction probably influences sapwood permeability. *Pseudotsuga menziesii* sapwood is less permeable than that of *Pinus contorta* or *Picea engelmannii* (Markstrom and Hann 1972). In contrast, heartwood permeability to liquids is higher in *P. engelmannii* and *P. menziesii* than in *P. contorta*; permeability to gases is higher in *P. contorta*.

BARK

Bark characteristics are more easily seen and compared than most physical attributes. As a result, bark traits tend to be general knowledge, and only a few comparisons of bark thickness have been published. These are very similar, ranking in the order shown in table 18. As the comparative bark thicknesses indicate, total bark content of the tree is higher for *Pseudotsuga menziesii* than for either *Tsuga heterophylla* or *Thuja plicata* (Smith, Kerr, and Czizmazia 1961). Relatively thin bark and a long cylindrical bole give *Abies procera* a greater volume for its diameter at breast height than associated species, with the possible exception of *Pinus monticola* (Hanzlik 1925).

Resin content of old bark was characterized in general terms by Flint (1925). *Pinus ponderosa*, *P. contorta*, and *P. monticola* barks have the most resin; *Larix occidentalis*, *Abies grandis*, *Tsuga heterophylla*, *T. mertensiana*, and *Thuja plicata* barks have the least. Flint rated *Pseudotsuga menziesii*, *Picea engelmannii*, and *Abies lasiocarpa* as intermediate in bark resin content.

SLASH

Bark resin content is an important determinant of fire intensity. So are the slash characteristics and foliage inflammability of north-western tree species. Foulger et al. (1976) analyzed western softwood logging residues less than .25 inch (.64 cm) in diameter for bark, needle, and solid wood content. They found that *Larix occidentalis* had the highest proportion of solid wood, followed by *Pinus contorta*, *Pseudotsuga menziesii*, *Abies concolor*, *A. lasiocarpa*, *Pinus ponderosa*, and *Picea engelmannii*. Moisture content of the solid wood component of slash also may differ in different species, sometimes in unexpected ways. Scott (1964) found that moisture distribution in large-diameter *Pseudotsuga menziesii* slash was radial, with moisture content increasing from 6 in (15 cm) inside the log out to its surface. The moisture distribution in large *Lithocarpus densiflora* slash on the same clearcut was just the opposite, however, with moisture content decreasing from inside out.

Moisture distribution, branch size, and needle retention all effect the flammability of slash. Olson (1953) made some preliminary tests of slash flammability in the western white pine type. He concluded that fire spread rates were highest in *Larix occidentalis* slash until it lost its needles, followed by *Tsuga heterophylla*, *Pseudotsuga menziesii*, *Pinus monticola*, *Abies grandis*, *Pinus contorta*, *Thuja plicata*, *Picea engelmannii*, and *Pinus ponderosa*. Although it had the slowest spread rate, *P. ponderosa* had the hottest-burning slash. Fahnestock (1960) tested slash quality more extensively. He suggested the following spread-rate rankings:

Fresh Slash

Larix occidentalis (fastest spread)

Pinus monticola, *Thuja plicata*,
Pseudotsuga menziesii, *Tsuga heterophylla*, and *Abies grandis*
(intermediate spread)

Pinus ponderosa and *Picea engelmannii*
(slowest spread)

One-year-old Slash

Pinus monticola, *P. contorta*,
Thuja plicata (fastest spread)

Pinus ponderosa, *Pseudotsuga men-*
ziesii, *Tsuga heterophylla*, and
Picea engelmannii (intermediate
spread)

Abies grandis and *Larix occidentalis*
(slowest spread)

Fahnestock found needle moisture content in fresh slash to be highest in *Larix occidentalis*, lowest in *Pseudotsuga menziesii*. Branchwood moisture in freshly cut slash was highest in *Pinus contorta*, lowest in *Larix occidentalis*.

MOISTURE CONTENT

Moisture content apparently differs in standing trees as well as in slash. Parker (1954) measured more heartwood moisture and total trunk water content in *Pinus ponderosa* and *Abies grandis* than in *Pseudotsuga menziesii* or *Thuja plicata*. When he compared twig moisture content with the foliage weight attached to those twigs, he found that *Pinus ponderosa* also had far more water per unit foliage than *Abies grandis*, *Pseudotsuga menziesii*, or *Thuja plicata*. The seasonal fluctuation of foliage moisture content was studied by Pharis (1967), who noted that the summer moisture content of new foliage was higher in *Pinus ponderosa* and *P. lambertiana* than in *Pseudotsuga menziesii*. Foliage moisture content declined between November and February in *P. menziesii* and *Abies grandis*, but not in *P. lambertiana*, *P. ponderosa*, or *Libocedrus decurrens*.

FOLIAGE FLAMMABILITY AND EPIPHYTE RECEPTIVITY

Foliage moisture probably influences foliage flammability, a factor rated by Flint (1925) in the northern Rocky Mountains and by Starker (1934) in Oregon and Washington. As their species comparisons were quite similar, they were combined in table 18 with only minor modification. Both authors listed

relative quantity of lichen growth by tree species. Coleman et al. (1956) studied the growth of lichens and other epiphytes more intensively, ranking Olympic Peninsula tree species in terms of epiphyte receptivity. They found *Alnus rubra* to be the best epiphyte host, *Tsuga mertensiana* to be the worst host on the Olympic Peninsula of Washington. In Oregon, Pechanec and Franklin (1968) found *A. rubra* to be a better epiphyte host than either *Picea sitchensis* or *Pseudotsuga menziesii*.

FIRE RESISTANCE

Bark characteristics, foliage flammability, and epiphyte populations all contribute to the total flammability of northwestern tree species. This whole-tree flammability and relative species resistance to fire have been compared by several authors. In the interior, Tackle (1959) observed that *Pinus contorta* was more susceptible to fire than other *Pinus* species or *Pseudotsuga menziesii*, but was less susceptible than *Picea engelmannii* and *Abies lasiocarpa*. Kerr (1913) also found that *P. contorta* was more fire-susceptible than *P. ponderosa*. Flint (1925) compared the fire resistance of 11 northern Rocky Mountain conifers, finding *Larix occidentalis* to be most resistant and *Abies lasiocarpa* least resistant. He noted that intolerant trees of the northern Rocky Mountains tend to be highly fire-resistant, while tolerant ones tend to have low resistance.

Relative species fire resistance does not appear to be the same in coastal forests as it is farther inland. *Thuja plicata* is more severely damaged by fire than any of its associates along the coast, but is less susceptible than *Picea engelmannii*, *Tsuga heterophylla*, and *Abies lasiocarpa* in the Inland Empire (Boyd 1959). Therefore, the fire resistance ranking in table 18 was divided into coastal and interior portions. *Pseudotsuga menziesii* seems to be the most fire-resistant coastal species (Starker 1934). Starker listed *Picea sitchensis* as the least fire resistant of 13 Oregon and Washington trees. Farther south, Stone and Vasey (1968) compared the

fire resistance of *Sequoia sempervirens* and its associates, noting that both *S. sempervirens* and *Lithocarpus densiflora* are more fire-resistant than *P. menziesii* and *Abies grandis*.

In the north, a survey of fire-killed timber on a burn in British Columbia showed that 50 percent of the *Thuja plicata*, but only 20 percent of the *Tsuga heterophylla* and only 1 percent of the *Pseudotsuga menziesii* trees were charred enough to degrade mill quality (Wallis, Godfrey, and Richmond 1974). When McNaughton (1944) studied charring and the ignition temperatures of wood, he found that speed of ignition was directly correlated with wood specific gravity. Species with low specific gravity (e.g., *Picea sitchensis*) ignited before species with high specific gravity (e.g., *Larix occidentalis*).

ROOTS

Root charring commonly occurs during severe forest fires, and species rooting characteristics probably are significant determinants of fire resistance. Young *Pinus ponderosa* roots grow deeper than *P. lambertiana* and *Libocedrus decurrens* roots, and the young *P. lambertiana* and *L. decurrens* roots tend to be deeper than *Abies grandis* or *Pseudotsuga menziesii* roots of the same age (Stein 1963). *Thuja plicata* roots are even shallower than those of *A. grandis* and *P. menziesii* (Ross 1932). *Tsuga heterophylla* seedling roots are shallower than the seedling roots of *Abies amabilis* (Scott et al. n.d.). Indeed, *Tsuga heterophylla* roots seem to be the shallowest of all (Haig 1936, Smith 1964). Although not compared directly with the above species, *Abies lasiocarpa* and *Picea glauca* apparently also have shallow roots--shallower than *Pinus contorta* (Eis 1970). *Pinus contorta* tends to be more shallow-rooted than *P. ponderosa* (Gail and Long 1935, Tarrant 1953).

Lateral extent, structure, and composition of species root systems vary even more than depth. *Populus tremuloides* probably has the greatest lateral root extent (Berndt and

Gibbons 1958). *Picea glauca* roots spread laterally more than *Abies lasiocarpa* roots, and *A. lasiocarpa* root systems spread more than those of *Pinus contorta* when root spread was considered in relation to tree height (Eis 1970). When root spread was related to crown width in root spread/crown width ratios, species were ranked as follows (Smith 1964):

Pinus contorta (highest ratio)
Picea sitchensis
Pinus ponderosa
Pseudotsuga menziesii
Tsuga heterophylla
Thuja plicata
Alnus rubra (lowest ratio)

As crown widths of these species differ (table 18), absolute root spread may not always be greater in species with higher root spread/crown width ratios. For example, roots of the wide-crowned *Pinus ponderosa* actually spread more laterally than those of the narrow-crowned *P. contorta* (Tarrant 1953). Nevertheless, *Pseudotsuga menziesii* root spread is wider than that of either *Tsuga heterophylla* or *Thuja plicata* when considered on an absolute basis (Eis 1974). Absolute measurements of the total root masses of windthrown trees of four species (Steinbrenner and Gessel 1956) showed *Pseudotsuga menziesii* to have the most extensive root system, followed in descending order by *Abies procera*, *Tsuga heterophylla*, and *Thuja plicata*.

When *Pseudotsuga menziesii*, *Tsuga heterophylla*, and *Thuja plicata* root distribution and structure are considered further, other differences become evident. *T. plicata* roots are more numerous in the duff than in underlying soil where a heavy duff layer exists (Ross 1932). Ross found that *T. heterophylla* root tips were also present in this duff, but not to the same extent. He found very few *P. menziesii* feeding roots in the duff layer...they were deeper, in the underlying soil. Ross also observed that *T. heterophylla* roots were smaller than those of *P. menziesii*, tending to be more oval in cross section than the circular *Pseudotsuga* roots.

Eis (1974) observed larger root diameters in *Pseudotsuga menziesii* than in either *Tsuga heterophylla* or *Thuja plicata*. After hydraulic

excavation, he found that thin, rope-like roots were most dense on *T. plicata* trees, least dense on *P. menziesii*. In an earlier study Eis (1972) noted that root grafting occurred in all three species, but was most frequent in *T. plicata*, least frequent in *P. menziesii*. The grafts apparently were important aids to the survival of suppressed trees on Vancouver Island.

Farther inland, Eis (1970) studied root growth relationships of juvenile trees in interior British Columbia. For the same age, *Pinus contorta* root systems were denser than those of either *Picea glauca* or *Abies lasiocarpa*. Vertical "sinker" roots were more numerous in *P. contorta*. Its roots were more branched than those of *Picea glauca*. In western Montana, Wyoming, and Idaho, Gail and Long (1935) also studied the rooting relationships of *Pinus contorta* and its associates. They found that *P. contorta* lateral roots grew more horizontal and closer to the soil surface than those of *P. ponderosa*.

The root systems of *Pseudotsuga menziesii* var. *glauca*, *Abies grandis*, *Larix occidentalis*, *Pinus monticola*, and *Thuja plicata* grown in large containers were investigated by Leaphart and Wicker (1966). They found the best taproot development in *L. occidentalis*, followed by *P. menziesii* and *A. grandis*. Taproots in *T. plicata* and *P. monticola* were poorly defined. Fine roots apparently varied inversely with taproot development for *T. plicata* had the most profusely developed fine root system, and *L. occidentalis* had the least profuse system.

Root system strength is a crucial factor in maintaining the stability of steep forested slopes. Burroughs and Thomas (1977) compared the tensile strength of *Pseudotsuga menziesii* var. *menziesii* and *P. menziesii* var. *glauca* roots, finding var. *menziesii* roots to be stronger when fresh. In similar studies, Ziemer and Swanston (1977) found live *Tsuga heterophylla* roots to be stronger than live *Picea sitchensis* roots. O'Laughlin (1973) found no significant difference in tensile strengths of *Pseudotsuga menziesii* and *Thuja plicata* roots.

WINDTHROW RESISTANCE

Root strength, depth, and structure affect the windthrow resistance of northwestern tree species--resistance that has been compared in several publications. The New Zealand Forest Service (1976) rated *Pseudotsuga menziesii* as more windfirm than most other conifers. Steinbrenner and Gessel (1956) also rated *P. menziesii* as most resistant, followed in order by *Abies procera*, *Tsuga heterophylla*, and *Thuja plicata*. Their *P. menziesii*-*T. heterophylla* comparison, based on large samples, is more reliable than their *A. procera* and *T. plicata* rankings, which were based on observations of only two and seven trees, respectively. When Gratkowski (1956) analyzed windthrow by comparing species composition of the stand with species composition of the windthrown trees, he found that short *Thuja plicata* trees growing on dry sites were most resistant, followed by *Pseudotsuga menziesii*, *Tsuga heterophylla*, and (least resistant) *Abies amabilis*. Boyd (1959) tentatively rated *Thuja plicata* fourth in Inland Empire windfirmness, exceeded by *Larix occidentalis*, *Pinus ponderosa*, and *Pseudotsuga menziesii*. Boyce (1929) found both *T. plicata* and *P. menziesii* to be more resistant to windthrow than *Tsuga heterophylla*, *Abies amabilis*, or *Picea sitchensis* on the Olympic Peninsula. In the Oregon Coast Range, *P. sitchensis* is more windfirm than *T. heterophylla* (Ruth and Yoder 1953).

DENSE SOIL TOLERANCE

Soils interact with species characteristics in determining windthrow resistance, and even a windthrow-resistant species may fare poorly on shallow or dense soils that limit root penetration and growth. Northwestern tree species differ with respect to their growth in dense soils, however--some are able to tolerate rather high soil densities, while others cannot.

On soils developed from Eocene to Miocene volcanic rocks in the western Cascade Range, clay subsoils restrict *Pseudotsuga menziesii*, but not *Pinus*

ponderosa or *P. lambertiana* roots (Stephens 1965). Cochran (1963) noted longer lateral roots and greater total root weights in *Pinus lambertiana* than in *P. ponderosa* during the initial stages of seedling development in high density soil. Soil density was one of the parameters utilized by Forristall and Gessel (1955) in comparing the growth of *Thuja plicata*, *Alnus rubra*, *Pseudotsuga menziesii*, and *Tsuga heterophylla* in Washington. Unfortunately, their soil density-growth comparisons probably were confounded by other factors, for the different species grew on plots with differing soil textures and moisture contents.

A uniform soil was artificially compacted to different bulk densities by Minore, Smith, and Woollard (1969) to compare the effects of high soil density on seedling root growth of seven northwestern species. In 2 years, the roots of *Pinus contorta*, *Pseudotsuga menziesii*, *Alnus rubra*, and *Abies amabilis* penetrated soil columns that the roots of *Picea sitchensis*, *Tsuga heterophylla*, and *Thuja plicata* did not. In another study, Leaphart (1958) found that *Larix occidentalis* and *Pinus monticola* roots penetrated hardpan layers more effectively than the roots of *Pseudotsuga menziesii*, *Pinus contorta*, *Abies grandis*, *Picea engelmannii*, *Thuja plicata*, or *Tsuga heterophylla*.

MECHANICAL DAMAGE RESISTANCE

Soil compaction probably is not very important when seedlings are buried by silt. Nevertheless, Hermann and Lavender (1967) observed less damage to *Pinus ponderosa* than to *Pseudotsuga menziesii* when dormant seedlings of both species were buried under silt for more than 8 weeks. Burial under litter that accumulates on top of the winter snow pack is resisted better by *Abies amabilis* than by *Tsuga heterophylla* seedlings (Thornburg 1969).

Damage caused by the snow pack itself varies greatly among species (table 18). *Tsuga heterophylla*, *T. mertensiana*, *Picea breweriana*, and *Abies magnifica* var. *shastensis* seem most resistant to snow damage.

Pinus monticola and *Abies amabilis* are somewhat less resistant but able to tolerate snow weights better than *Abies concolor*, *A. procera*, or *Pinus jeffreyi* (Williams 1966, Waring et al. 1975). *Pseudotsuga menziesii*, *Pinus lambertiana*, and *P. ponderosa* seem to be most subject to snow damage of the species compared.

ANIMAL RELATIONSHIPS

Most animal damage in young stands seems to be caused by gophers, rabbits, or deer. No species comparisons are available for gopher damage, for available data indicate little or no species preference by these rodents. Differences in rabbit damage were noted by Worthington (1955), however, who found *Pseudotsuga menziesii* to be damaged more than *Thuja plicata* and *Tsuga heterophylla*. *Abies grandis* and *Picea sitchensis* were least damaged on Worthington's Olympic Peninsula sites. Staebler et al. (1954) also found *P. menziesii* to be more damaged by rabbits than *T. heterophylla*. The *Chamaecyparis lawsoniana* seedlings on their southwestern Washington plots were browsed by deer, but only after 3 years.

Schubert (1956) observed more deer browsing on *Abies concolor* than on *Pinus lambertiana*, and Stark (1965) found more browsing on *A. concolor* and *Libocedrus decurrens* than on *Pinus ponderosa* and *P. lambertiana*. *Abies magnifica* also seems to be less damaged by deer browsing than *A. concolor* (Gordon 1970). When both rabbit and deer damage are considered, comparative animal damage may be ranked by species groups:

Pinus ponderosa, *P. lambertiana*,
Abies grandis, *A. magnifica*,
and *Picea sitchensis* (least
damaged)

Abies concolor, *Libocedrus*
decurrens, and *Tsuga hetero-*
phylla

Thuja plicata and *Chamaecyparis*
lawsoniana

Pseudotsuga menziesii (most
damaged)

Relative species palatability may influence relative species damage, but highly palatable species may not comprise the majority of an animal's diet. For example, *Thuja plicata* is a highly palatable, preferred deer food; but it occurred in only 20-30 percent of the deer stomachs sampled by Cowan (1945) on Vancouver Island, never amounting to more than 2 percent of the food volume. In contrast, *Pseudotsuga menziesii* occurred in 50-100 percent of Cowan's deer samples and accounted for 25-50 percent of their food volume.

Pseudotsuga menziesii seeds are eaten by rodents in preference to *Abies procera* seeds (Dick 1960). Small mammals and birds also seem to prefer *P. menziesii* seeds to the seeds of *Tsuga heterophylla* (Gashwiler 1970). They prefer *T. heterophylla* to *Thuja plicata* (Gashwiler 1967).

INSECT RESISTANCE

General insect resistance seems to be associated with species longevity. *Thuja plicata* is more insect-resistant than *Tsuga heterophylla* or *Abies amabilis* in coastal British Columbia (Schmidt 1955). It also is more resistant to insect attacks than most of its Inland Empire associates (Boyd 1959). *Thuja plicata* maintains this high insect-resistance rating when more limited insect categories are considered. It is attacked less by woodborers than *Pseudotsuga menziesii*, *Abies grandis*, or *Tsuga heterophylla* (Johnson 1958); and its extracts apparently induce more deaths in western tent caterpillar pupae than extracts of *P. menziesii* (Wellington 1969). *P. menziesii* is attacked by the silver-spotted tiger moth more often than *Thuja plicata*, *Tsuga heterophylla*, *Pinus contorta*, *Picea sitchensis*, *Abies amabilis*, and *A. grandis* in British Columbia (Sellars-St. Clare 1968). *Abies grandis* is damaged more than *P. menziesii* by western spruce budworm attacks, however, even when budworm egg numbers are similar in the upper crowns of these two species (Carolin and Coulter 1975).

The crowns of *Pinus lambertiana* and *P. monticola* seedlings were more seriously infested with woolly aphids than those of *P. albicaulis* when an accidental aphid infestation occurred during a lathhouse blister rust experiment (Hoff and McDonald 1977). Mitchell (1966), comparing *Abies* species in the field, observed that *A. concolor* and *A. procera* were most resistant to balsam woolly aphid. *Abies lasiocarpa* was least resistant (table 19).

Abies lasiocarpa also probably lacks resistance to termite attack, as do *Picea engelmannii*, *Tsuga heterophylla*, *Larix occidentalis*, and *Pseudotsuga menziesii*. Carter and Smythe (1974) found the woods of these species to be better termite food than the woods of *Thuja plicata*, *Sequoia sempervirens*, *Chamaecyparis lawsoniana*, and *Pinus ponderosa*. *Pinus ponderosa* apparently is cutworm resistant as well. When Fowells (1940) and Fowells and Stark (1965) tallied cutworm damage to seedlings in California, they found *Pinus lambertiana* to be least damaged, followed by *P. ponderosa* and *P. jeffreyi* (table 19). The cutworms showed a preference for *Libocedrus decurrens* over *Abies concolor* and the *Pinus* species.

Further comparisons of insect resistance among northwestern species are complicated by host specificity problems. Many serious insect pests attack only one or two tree species, and literature references to comparative species resistances are unavailable for them.




Table 19--Comparative insect resistance of northwestern tree species^{1/}

Cutworms ^{2/}	Wood borers ^{3/}	Termites ^{4/}	Balsam woolly aphid ^{5/}	Woolly aphid ^{6/}
<i>Pinus lambertiana</i>	<i>Thuja plicata</i>	<i>Thuja plicata</i> ,	<i>Abies procera</i> ,	<i>Pinus albicaulis</i>
<i>Pinus ponderosa</i> ,	<i>Pseudotsuga</i>	<i>Chamaecyparis</i>	<i>A. concolor</i>	<i>Pinus monticola</i>
<i>P. jeffreyi</i>	<i>menziesii</i> ,	<i>lawsoniana</i> ,	<i>Abies magnifica</i>	<i>Pinus lambertiana</i>
<i>Abies concolor</i>	<i>Tsuga hetero-</i>	<i>Sequoia</i>	var. <i>shastensis</i>	
<i>Libocedrus</i>	<i>phylla</i> ,	<i>sempervirens</i> ,	<i>Abies grandis</i>	
<i>decurrens</i>	<i>Abies grandis</i>	<i>Pinus</i>	<i>Abies amabilis</i>	
		<i>ponderosa</i>	<i>Abies lasiocarpa</i>	
		<i>Picea engelmannii</i> ,		
		<i>Tsuga hetero-</i>		
		<i>phylla</i> , <i>Larix</i>		
		<i>occidentalis</i>		
		<i>Abies lasiocarpa</i> ,		
		<i>Pseudotsuga</i>		
		<i>menziesii</i>		

^{1/}Resistant species are listed above susceptible ones. Species in the same group are not necessarily equal, but data are insufficient for species comparisons within groups.


^{2/}From Fowells (1940) and Fowells and Stark (1965).

^{3/}From Johnson (1958).

^{4/}From Carter and Smythe (1974).

^{5/}From Mitchell (1966).

^{6/}From Hoff and McDonald (1977).



DISEASE RESISTANCE

Comparing disease resistance of northwestern tree species also is complicated by host specificity. Many pathogens, though widespread, are limited to a small number of host species. Other diseases consist of many pathogens attacking many trees, with each pathogenic species preferring a single tree species. The dwarf mistletoes are an example. Smith (1974) inoculated several tree species with several dwarf mistletoe species. He observed that each had a principal host on which it was particularly infectious, but they were not strictly host-specific. The greatest mistletoe seed retention occurred on the branches of *Picea engelmannii*, *P. sitchensis*, and *P. glauca*. The poorest occurred on *Pinus ponderosa* and *Larix occidentalis*. In contrast, the best mistletoe seed germination

was on *Tsuga heterophylla*, *Pseudotsuga menziesii*, *Pinus ponderosa*, and *Abies grandis*; the poorest was on the *Picea* species.

Host specificity sometimes is evident within a single genus. For example, northwestern *Pinus* species exhibit some differential resistance to attack by rust fungi. *Pinus contorta* is more resistant to comandra rust than *P. ponderosa* (Bergdahl and French 1976). *P. monticola*, though susceptible, is less susceptible to white pine blister rust than other northwestern five-needled *Pinus* species. *Pinus albicaulis* is most susceptible and *P. lambertiana* is intermediate (Childs and Bedwell 1948).

Unfortunately, all northwestern *Pinus* species seem to be equally susceptible to annosus root rot (Bega 1962). In California they die soon after *Fomes annosus* invades and rapidly kills the cambium.

This does not happen in other California conifers. Instead, the fungus causes a chronic decay condition which the *Abies* species and other non-*Pinus* conifers appear to tolerate for several years (Bega and Smith 1966). *Pinus* species may be more tolerant in Britain than they are in California, for Aldhous and Low (1974) subjectively rated the pines as being more resistant than other conifers to *Fomes* butt rot. These other conifers do possess varying degrees of susceptibility to annosus root and butt rot, however, wherever they are grown (table 20). *Tsuga heterophylla* and *Picea sitchensis*

are less resistant than *Abies grandis*, *A. procera*, and *Pseudotsuga menziesii* in Britain, where *Thuja plicata* is relatively susceptible (Aldhous and Low 1974). In British Columbia *Pseudotsuga menziesii* roots are more resistant than *Tsuga heterophylla* roots, and *Thuja plicata* stumps are more resistant to *Fomes* spore infection than the stumps of other conifer species (Wallis and Ginns 1968).

Tsuga heterophylla seedlings are more susceptible than *Pseudotsuga menziesii* seedlings to *Rhizina undulata*, a root pathogen that can be serious where slash has been burned (Morgan 1973).

Table 20--Comparative decay resistance of northwestern tree species^{1/}

Annosus root and butt rot ^{2/}	Laminated root rot ^{3/}	Decay in living trees ^{4/}	Decay in dead wood ^{5/}
<i>Pseudotsuga menziesii</i> , <i>Abies procera</i> , <i>A. grandis</i>	<i>Acer macrophyllum</i> , <i>Alnus rubra</i> , <i>Arbutus menziesii</i> , <i>Lithocarpus densiflora</i> , <i>Populus trichocarpa</i> , <i>P. tremuloides</i>	<i>Pinus ponderosa</i> <i>Pinus monticola</i> , <i>P. lambertiana</i>	<i>Thuja plicata</i> , <i>Sequoia sempervirens</i> , <i>Taxus brevifolia</i>
<i>Tsuga heterophylla</i> , <i>Picea sitchensis</i>	<i>Libocedrus decurrens</i> , <i>Pinus ponderosa</i> , <i>Thuja plicata</i>	<i>Pinus contorta</i> <i>Sequoia sempervirens</i> <i>Picea sitchensis</i> <i>Pseudotsuga menziesii</i>	<i>Larix occidentalis</i> <i>Pseudotsuga menziesii</i> <i>Pinus ponderosa</i> , <i>P. lambertiana</i> , <i>Picea sitchensis</i>
	<i>Pinus contorta</i> , <i>P. monticola</i>	<i>Abies</i> species <i>Tsuga heterophylla</i>	<i>Pinus monticola</i> , <i>P. contorta</i> , <i>Abies procera</i> , <i>A. concolor</i> , <i>A. amabilis</i> , <i>Tsuga heterophylla</i> , <i>Picea engelmannii</i>
	<i>Abies lasiocarpa</i> , <i>Larix occidentalis</i> , <i>Picea engelmannii</i> , <i>P. sitchensis</i> , <i>Tsuga heterophylla</i>		
	<i>Abies amabilis</i> , <i>A. grandis</i> , <i>Pseudotsuga menziesii</i> , <i>Tsuga mertensiana</i>		

^{1/}Resistant species are listed above susceptible ones. Species in the same group are not necessarily equal, but data are insufficient for species comparisons within groups.

^{2/}From Aldhous and Low (1974) and Wallis and Ginns (1968).

^{3/}From Hadfield and Johnson (undated).

^{4/}From Buckland et al. (1949), Wagener and Davidson (1954), and Farr et al. (1976).

^{5/}From Boyce (1929), Englerth and Scheffer (1954), USDA Forest Products Laboratory (1955), and Clark (1957).

When Vaartaja (1957) germinated seven northwestern tree species on agar cultures of a damping-off fungus (*Phytophthora cactorum*), he found *Thuja plicata* to be much more resistant than *Larix laricina*, and *L. laricina* was more resistant than *Picea glauca*, *P. mariana*, *P. sitchensis*, *Pinus contorta*, or *P. ponderosa*.

Another fungus, the laminated root rot caused by *Phellinus* (*Poria*) *weirii*, has been studied extensively. It occurs on many northwestern species. *Thuja plicata*, *Alnus rubra*, and *Acer macrophyllum* have more resistance to this fungus than *Pseudotsuga menziesii* (Wallis and Reynolds 1967). *Phellinus* infected roots have profuse ectotrophic mycelium in *Pseudotsuga menziesii* and *Tsuga heterophylla*; limited mycelium in *Pinus monticola*, *P. contorta*, and *P. ponderosa*; and almost no mycelium in *Thuja plicata* (Wallis 1976). Hadfield and Johnson (undated) compared host susceptibility in the summary publication used as a source for the species comparisons listed in table 20.

Table 20 summarizes the general decay information available in several literature sources. Living *Picea sitchensis* trees less than 400 years old are relatively free of rot in southeastern Alaska; *Tsuga heterophylla* trees are not (Farr et al. 1976). *Abies amabilis* less than 350 years old on Vancouver Island also is more resistant to infection by wood-rotting fungi than *Tsuga heterophylla* (Buckland et al. 1949). When heart rot in *Abies* species is compared with rot in other genera, it is less prevalent than in *Tsuga*, but more prevalent than in *Pseudotsuga* or *Picea* (Wagener and Davidson 1954). Within genera, Wagener and Davidson's data indicate that living *Pinus ponderosa* has less heart rot than *P. monticola* and *P. lambertiana*.

Pinus ponderosa and *P. lambertiana* were more decay-resistant than *P. monticola*, *P. contorta*, *Abies procera*, *A. concolor*, *Tsuga heterophylla*, and *Picea engelmannii* when Clark (1957) exposed heartwood samples to wood-rotting fungi in the laboratory. Clark found *Pseudotsuga menziesii* heartwood to be even more resistant than the *Pinus* species. In similar laboratory

tests, Englerth and Scheffer (1954) found *Larix occidentalis* to be more decay-resistant than *Pseudotsuga menziesii* but less resistant than *Thuja plicata*. *Thuja plicata* was also found to be more decay-resistant than the wood of other species by Boyce (1929) and Wallis et al. (1974). Boyce, studying the deterioration of windthrown timber on the Olympic Peninsula, recorded more decay in *Abies amabilis* and *Tsuga heterophylla* than in *Picea sitchensis*, more in *P. sitchensis* than in *Pseudotsuga menziesii*.

When Childs (1939) studied the deterioration of slash on clearcut areas in the Douglas-fir region, he found the rate of sapwood decay to be about the same in *Pseudotsuga menziesii*, *Picea sitchensis*, and *Tsuga heterophylla* slash. *P. menziesii* heartwood decayed more slowly than the other two species, however, and heartwood decay in *Picea sitchensis* was slower than in *Tsuga heterophylla*. Similar reports were reported by Roff and Eades (1959) and by Ruth and Harris (1975). *Tsuga heterophylla* and *Abies amabilis* heartwood decay faster than the heartwoods of *Picea sitchensis* and *Pseudotsuga menziesii*. *Thuja plicata* is slowest to decay.

When root decay rates are compared, species ranking changes slightly. As one would expect, *Thuja plicata* roots lose their tensile strength more slowly than *Pseudotsuga menziesii* roots (O'Laughlin 1973). *Picea sitchensis* roots decay faster than *Tsuga heterophylla* roots in southeastern Alaska, however (Ziemer and Swanston 1977), reversing the relative heartwood decay rates of these two species.

CHEMICALS

Most chemically oriented species comparisons in the literature refer to chemical damage, not benefits. For example, smoke from slash burning apparently damages *Pseudotsuga menziesii* more than *Pinus lambertiana* or *P. ponderosa*, which are damaged more than *Libocedrus decurrens* or *Abies grandis* (Stein 1963). Nevertheless, some chemical effects are beneficial. Germination rates may be accelerated by applying hydrogen

peroxide to seeds as a substitute for lengthy stratification. *Pseudotsuga menziesii* and *Larix occidentalis* are benefitted more than *Abies lasiocarpa* by this treatment (Shearer and Tackle 1960). *P. menziesii* also benefits more than *Pinus lambertiana* or *P. ponderosa* when seeds are treated with hydrogen peroxide (Stein 1965).

HERBICIDES

Pseudotsuga menziesii is more resistant to injury by herbicides used in brush control than *Pinus ponderosa* (Newton 1963), and it is less damaged than *Alnus rubra* by dormant sprays (Gratkowski 1975). Brush control herbicides damage *Tsuga heterophylla* more than *Picea sitchensis* (Krygier and Ruth 1961). *T. heterophylla* also is damaged by weed-control herbicides. When Stewart (1977) tested 18 weed-control herbicides in forest nurseries, he found *T. heterophylla* and *Sequoia sempervirens* to be more sensitive to the herbicides than *Pseudotsuga menziesii*, *Abies concolor*, *A. grandis*, *A. procera*, and *A. magnifica*. *Pinus ponderosa*, *P. contorta*, and *P. lambertiana* seemed to be less sensitive.

SALT

Picea sitchensis is less sensitive to ocean spray than *Tsuga heterophylla* or *Thuja plicata* on the west coast of Vancouver Island (Cordes 1973). Indeed, no other tree is more tolerant of ocean spray than *P. sitchensis* (Krajina 1970). When lower tolerance levels were compared, *Pinus ponderosa* was more salt-tolerant than *Pseudotsuga menziesii* (Monk and Wiebe 1961, Carpenter 1970).

SULFUR DIOXIDE

The sulfur dioxide produced by copper smelters and other large industrial installations has been damaging northwestern trees for many years. Mason (1915) ranked *Abies lasiocarpa* as the most easily killed Rocky Mountain species, fol-

lowed in order of decreasing susceptibility by *Pseudotsuga menziesii*, *Pinus contorta*, *Picea engelmannii*, *Juniperus scopulorum*, and *Pinus flexilis*. Scheffer and Hedgcock (1955) also rated *A. lasiocarpa* as most susceptible to SO₂ injury. They agreed with Mason's other observations, but added *Pinus ponderosa* var. *scopulorum* and ranked it more resistant than *Picea engelmannii* in Montana.

In the upper Columbia River Valley, Scheffer and Hedgcock found *Abies grandis* to be even more sulfur-dioxide-susceptible than *A. lasiocarpa*. They ranked *Thuja plicata* as slightly more resistant than *Abies lasiocarpa* in this locality but found *Pinus ponderosa* to be less SO₂ tolerant than *P. contorta*, *Larix occidentalis*, or *Picea engelmannii* (table 21). Farther west, on the British Columbia coast, *Tsuga heterophylla* is much more tolerant to sulfur dioxide than *Thuja plicata*, with *Abies amabilis* and *Picea sitchensis* resistance intermediate between these extremes (Miller and McBride 1975). *Tsuga mertensiana* and *Populus tremuloides* are sensitive to sulfur dioxide (U.S. Department of Agriculture 1973).

FLUORIDES

Picea glauca and the *Juniperus* species are tolerant of fluorides in the atmosphere (U.S. Department of Agriculture 1973). Produced by aluminum ore reduction plants and phosphate fertilizer factories, these fluorides cause more damage to *Pinus* species than to *Pseudotsuga menziesii* in Montana (Miller and McBride 1975). *P. menziesii* is damaged more than *Abies lasiocarpa*, *Thuja plicata*, and the *Picea* species. In Norway, Horntvedt and Robak (1975) also found *P. menziesii* to be damaged more than several other northwestern species. Their comparisons indicate the following fluoride ranking:

Tsuga heterophylla (least susceptible)
Abies procera
Picea engelmannii
Abies concolor
Pseudotsuga menziesii (most susceptible)

Table 21--Comparative air pollution tolerance of northwestern tree species^{1/}

Sulfur dioxide ^{2/}	Fluorides ^{3/}	Ozone ^{4/}
<i>Taxus brevifolia</i>	<i>Tsuga heterophylla</i>	<i>Pinus lambertiana</i> , <i>Libocedrus decurrens</i>
<i>Juniperus scopulorum</i>	<i>Abies procera</i>	<i>Pinus ponderosa</i> var. <i>scopulorum</i> , <i>P.</i> <i>jeffreyi</i> , <i>P. attenuata</i> , <i>Pseudotsuga menziesii</i> , <i>Abies concolor</i>
<i>Pinus ponderosa</i> var. <i>scopulorum</i>	<i>Picea engelmannii</i>	
<i>Picea engelmannii</i>	<i>Abies concolor</i>	
<i>Larix occidentalis</i>	<i>Pseudotsuga</i> <i>menziesii</i>	
<i>Pinus contorta</i>	<i>Pinus contorta</i>	<i>Pinus ponderosa</i> , <i>P.</i> <i>monticola</i> , <i>Abies</i> <i>magnifica</i>
<i>Pinus ponderosa</i>	<i>Pinus ponderosa</i>	
<i>Pinus monticola</i>	<i>Pinus monticola</i>	
<i>Pseudotsuga</i> <i>menziesii</i>		
<i>Tsuga heterophylla</i>		
<i>Abies amabilis</i> , <i>Picea sitchensis</i>		
<i>Thuja plicata</i>		
<i>Abies lasiocarpa</i>		
<i>Abies grandis</i>		

^{1/}Resistant species are listed above susceptible ones. Species in the same group are not necessarily equal, but data are insufficient for species comparisons within groups.

^{2/}From Mason (1915), Scheffer and Hedgcock (1955), and Miller and McBride (1975).

^{3/}From Horntvedt and Robak (1975) and Miller and McBride (1975).

^{4/}From Miller and Millecan (1972) and Miller and McBride (1975).

OZONE

Ozone damage is serious in the forests near several urban centers in southern California, and it may become serious elsewhere as urban expansion continues. *Pinus ponderosa*, *P. monticola*, and *Abies magnifica* seem to be most sensitive to this pollution; *P. lambertiana* and *Libocedrus decurrens* are most tolerant (Miller and McBride 1975). *Sequoia sempervirens* and *Thuja plicata* also seem to tolerate ozone in the air (U.S. Department of Agriculture 1973). *Pinus ponderosa* var. *scopulorum*, *P. jeffreyi*, *P. attenuata*, *Pseudotsuga menziesii*, and *Abies concolor* appear to be intermediate with respect to ozone tolerance.

MISSING COMPARISONS

The species comparisons presented in this report are summarized in figure 1, with page numbers to provide access to relative species rankings and relationships. Figure 1 should serve as a useful index. In addition, it illustrates existing gaps in our knowledge of comparative autecological attributes. Some species (e.g., *Pseudotsuga menziesii*, *Tsuga heterophylla*, and *Pinus ponderosa*) have been compared with other species for most of the attributes listed. Others (e.g., *Chamaecyparis nootkatensis* and *Abies magnifica* var. *shastensis*) have been compared for less than half of them.

Figure 1--A summary of autecological comparisons for the major northwestern tree species, with page numbers to indicate where the comparisons are found.

Autecological attributes	Major northwestern tree species ^{1/}																					
	AbAm	AbCo	AbGr	AbLa	AbMa-sh	AbPr	AlRu	ChLa	ChNo	LaOc	LiDe	PiEn	PiSi	PiCo	PiJe	PiLa	PiMo	PiPo	PsMe	ThPl	TsHe	TsMe
Air pollution tolerance	50,51	50,51	49,50,51	50,51		50,51				50,51	49,51	50,51	50,51	50,51	51	49,51	51	49,50,51	49,50,51	50,51	50,51	50
Animal damage		45	45					45			45		45			45		45	45,46	45,46	45,46	
Anthesis	25,26		25,26	25,26		25,26			25,26					26			26		25,26	25,26	25,26	25,26
Bark thickness		38	38	38		38,41			38	38		38	38	38			38,41	38	38,41	38,41	38,41	38
Cotyledon number	36	36	36	36	36	36		36	36	36	35,36	36	36	36	36	36	36	36	36	35,36	36	36
Crown density	40		37	37						37		37		37			37	37	37	37,40	37,40	
Crown length		37	37	35,37		37				35,37		35,37	37	37			35,37	35,37	35,37	37	37	37
Crown weight		35,36	36	36						35,36	35,36	36		35,36		36	35,36	35,36	35,36	36	36	
Crown width			37							37,40		37		37,40			37,40	37,40	37	37	37,40	
Decay resistance	48,49	48,49	48	48		48,49	48,49			48,49	48	48,49	48,49	48,49		48,49	48,49	48,49	48,49	48,49	48,49	48
Dense soil tolerance	39,45		45			45				39,45		45	39,45	39,45		39,45	39,45	39,45	39,44,45	39,45	39,45	
Drought tolerance	11,12	10,11,12	10,11,12	11,12	11,12	10		11,12		11,12	10,11	10,11	10,11	10,11	10,11	11,12	11,12	10,11	10,11,12	11,12	10,11,12	11,12
Epiphyte receptivity		38	38	38		38	38,42		38	38		38	38,42	38			38		38,42,43	38	38,43	38,42
Fire resistance		39	39,43	39,42		39				39,42,43	39	39,42	39,42,43	39,42		39	39	39	39,42,43	39,42,43	39,42	39
Foliage flammability		38	38	38		38				38		38	38	38			38	38	38	38	38	38
Frost tolerance	6,7	6,7	6,7	6,7	7		6,8	7	6	6	7	6,7	6,7,8	6,7	6,7	7	6,7	6,7	6,7,8	6,7,8	6,7,8	6,7
Germination rate	30	30,31	30,31		29,31	29,31					30,31	31,32			30,31	30,31,32	30,31,32	30,31	30,31,32		31	
Heat tolerance			8	8						8		8			8	8	8	8	8	8	8	
Herbicide tolerance		50	50			50	50						50	50		50		50	50		50	
Insect resistance	46,47	46,47	46,47	46,47	47	46,47		46,47		46,47	46,47	46,47	46	46	46,47	46,47	46,47	46,47	46,47	46,47	46,47	
Light intensity		5	5		5		6					6	5,6						5,6		5,6	
Litter fall	37						37						37	37,40			37	37	37,40	37,40	37	
Longevity	23	23	23	23		23	23	23	23	23	23	23	23	23		23	23	23	23	23	23	23
Moisture (excess)			12,13	12,13			12,13		13			12	12,13	12,13	13	12		12,13	12,13	12,13	12,13	13
Moisture (optimum)	14	14	14	15	14	14	14	15			13,14	15	14	15		14		13,14,15	13,14,15	14	14	14
Mycorrhizae	21	21	21	21	21	21	20,21	21	21	21	21	21	21	21	21	21	21	20,21	20,21	21	20,21	21

Autecological attributes	Major northwestern tree species ^{1/}																
	AbAm	AbCo	AbGr	AbLa	AbMa-sh	AbPr	AIRu	ChLa	ChNo	LaOc	Lide	PIEn	PISi	PICo	PIle	PIla	PIPo
Needle density							35						35	35		35	35
Needle longevity	36		36										36	36		36	35,36
Needle volume			36										36	36		36	36
Needle weight	36		36										36	36		36	36
Nutrient concentrations	15,16,17		15,16,17				16,17	15,17		16,17		15,16,17	15,16,17	16,17		16	15,16,17,18
Nutrient deficiencies	19	19	19	19	19				18		19	18,19	17,18,19	19	19	17,18,19	17,18,19
Nutrient excesses													19	19		19	19
Root growth rate			22	22						22,23	22,23	22	22,23	22,23		22,23	22,23
Rooting depth	43		39,43	43						39	39,43		43			39,43	39,43
Sapwood thickness						40				38		38					38,40
Seed bearing age	26	26		25	26	26	26	26		26		25	26	26	26	26	26
Seedbeds	34		33,34	33,34			33,34			33,34	32	33,34	32,33,34	33,34		32	33,34
Seed crop frequency	27		27	27						27,28		27,28	28	27,28		27,28	27,28
Seed crop size	27,28	27,28	28		28	28	27,28	27		27,28	28	28	27	27	27	27,28	27,28
Seed dissemination	25,26,27	26	25,26,27	26,27	26	25,26,27	26	26	26	25,26	26	26	25,26	26	25,26,27	25,26,27	25,26,27
Seed flight	29,30	30	30			29				30		29	29,30	30	29,30	29,30	29,30
Seed longevity		31	31	31		31	31	31		31	31	31	31	31	31	31	31
Seed stratification	31,32	31	31,32	31	31	31,32	31	31				31,32	31,32	31	31,32	31,32	31,32
Seed weight	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30
Shade tolerance	3,4	3,4	3,4	3,4	3,4,5	3,4,5	3,5	3,5	3,4	3,5	3,5	3,4,5	3,4,5	3,4,5	3,4,5	3,4,5	3,4,5
Shoot growth period	23,24	23,24	23,24	24		23,24		23,24				23,24	23,24	23,24	23,24	23,24	23,24
Size	23	23,24	23	23		23	23,24	23	23	23	23,24	23	23,24	23	23	23,24	23,24
Slash flammability			41,42							41,42		41,42		41,42		41,42	41,42
Snow damage	39,45	39,45			39,45	39,45								39,45	39,45		39,45
Specific gravity	37,40	37,40	37,38,40	37		37,40	37			37,38,40		37,40	37	37,38	37	37	37,38,40
Temperature (mean)	9		9	8,9	9	9					8,9	8	9	8,9		8,9	8,9
Windthrow resistance	39,44					44				44			39,44			44	39,44


^{1/} Species names are abbreviated as follows: AbAm = *Abies amabilis*; AbCo = *Abies concolor*; AbGr = *Abies grandis*; AbLa = *Abies lasiocarpa*; AbMa-sh = *Abies magnifica* var. *ehussteniae*; AbPr = *Abies procera*; AIRu = *Alnus rubra*; ChLa = *Chamaecyparis lassoniana*; ChNo = *Chamaecyparis nootkatensis*; LaOc = *Larix occidentalis*; Lide = *Lithocedrus decurvens*; PIEn = *Picea engelmannii*; PISi = *Picea sitchensis*; PICO = *Pinus contorta*; PIle = *Pinus jeffreyi*; PIla = *Pinus lambertiana*; PIPo = *Pinus monticola*; PIPo = *Pinus ponderosa*; PSMe = *Pseudotsuga menziesii*; ThP1 = *Thuja plicata*; TSMe = *Tsuga heterophylla*; TSMe = *Tsuga mertensiana*.

As one would expect, widespread species have been compared more than those with limited ranges. *Chamaecyparis lawsoniana* comparisons are particularly scarce. As *C. lawsoniana* is a valuable species seriously threatened by disease, its autecological attributes probably should be related to those of other species. Such autecological comparisons might help pathology researchers and provide information about suitable substitutes if *C. lawsoniana*'s ecological niche must be occupied by other species.

Pinus jeffreyi appears to be another species that seldom has been compared because of its limited range. It may resemble *P. ponderosa* in some cases where blanks appear in figure 1, but probably has several unique attributes that are not matched in other species. Nutrient concentrations and mean temperature comparisons seem particularly worthy of investigation in *P. jeffreyi*.

Limited range probably does not account for the surprising lack of *Alnus rubra* comparisons. Its low commercial value may be responsible. Whatever the reason, *A. rubra* comparisons should be neglected no longer. Root growth rate and rooting depth relationships seem particularly pertinent in light of *A. rubra*'s nitrogen fixing characteristics (Tarrant and Miller 1963, Bollen et al. 1967, Franklin et al. 1968, and Tarrant et al. 1969). Its tolerance of nutrient deficiencies also should be compared with associated species.

Associated species probably should be compared in as many ways as possible, wherever they grow. Considerable amounts of work remain to be done if this is to be accomplished for the autecological attributes listed here. Comparative crown, needle, and root characteristics, litter fall, slash flammability, and windthrow resistance are all sparsely represented in figure 1.



UNKNOWN ATTRIBUTES

The autecological attributes compared in this report are somewhat subjective. They represent attempts to delineate and define discrete variables in the non-discrete continuum of interacting characteristics that define the nature and properties of northwestern tree species as we know them. Others might use somewhat different categories or discuss these attributes under different headings. Nevertheless, the attributes do exist as such, and most have been compared in this report--however one wishes to categorize and name them.

Some attributes have not been compared, however, for they seem to be absent from the literature. As comparable information is lacking, these attributes constitute unknowns...not just for one or two but for most northwestern tree species.

Most northwestern tree species grow in mixed stands, where they compete with other species. Relative ability to compete constitutes an autecological unknown. Unfortunately, it is an unknown that must be qualified by knowing where the competition will occur and in what environment. Comparative competitive abilities probably will remain unknown until they are determined in various environments throughout the natural ranges of the species involved.

Potential species ranges constitute another unknown autecological attribute. Where would each species occur if it possessed a capacity for worldwide distribution and was not limited by competition with other species? A complete knowledge of all other attributes, for all species, may be a prerequisite to answering this question. It probably will remain unanswered for the foreseeable future.

Fortunately, several other attributes are less likely to remain unknown. Study of mycorrhizal relationships is now being accelerated, and detailed comparisons among northwestern species should be available in a few more years. Although the fungus-host relationships may be rather specific, it should soon be possible to rate

northwestern tree species in terms of mycorrhizae dependence and numbers of fungi involved.

Genetic variability and gene-pool characteristics also are being studied for several northwestern tree species. Genetic plasticity may vary greatly among species, and some gene pools probably are larger and more complex than others. Better understanding of these presently unknown genetic attributes may permit species comparisons that are impossible today. It also should improve all present species ratings by facilitating evaluation of within-and-among species variations.

Although coniferous forests selectively filter light (Atzet and Waring 1970), among-species comparisons of light filtering properties and foliage albedo were not found in the literature--apparently because they are little known for northwestern tree species. Like many of the comparisons missing for individual species, these attributes probably will be recorded and compared as the needed research is accomplished.

RESEARCH NEEDS

Additional research is needed to supply the missing information indicated by blank spaces in figure 1 and to describe the unknown attributes discussed above. More than this is needed, however, even for the filled spaces in figure 1 and for well-known autecological attributes. Many of the comparisons cited here were based upon observations rather than measurements, and most occurred under conditions that could have influenced the results by introducing imperceptible extraneous influences. Quantification and refinement of these comparisons through rigorous experimental procedures carried out under controlled conditions should result in more consistent, reliable conclusions.

Quantitative comparison of photosynthetic activity under various light conditions has been accomplished for few species. It should be done for all, at several ambient temperatures. Photoperiodic responses also should be compared to determine

relative light requirements for strobilus formation, anthesis, and dormancy. Again, several ambient temperatures should be used to investigate the light, temperature, and species interactions.

Species-environment interactions probably are as important as species attributes themselves when growth is being compared in the field. These interactions should be studied carefully by establishing replicated species trials in many different field environments. Wherever possible, these field environments should be categorized by quantitative measurements. They should always be classified or described in ways that allow comparison and contrast with other environments.

Growth comparisons in the field often require long-term experiments that may exceed the professional life span of any single researcher. Comparisons of germination rates, stratification requirements, and optimal germination temperatures do not have this inherent disadvantage. Environmental interactions are also important in these short-term comparisons, however, and they should be accomplished in carefully designed experiments that will result in usable interaction data.

Finally, it should be emphasized that two or more species should be tested together and compared whenever possible in autecological experimentation. Even the most well-designed experiment, carried out under highly controlled conditions, is somewhat unique. Exactly identical conditions are impossible to attain again, regardless of the precautions taken. Comparative data involving more than one species may allow further extrapolation of experimental results than that which is possible with a single species, and wider inferences often may be made. Relative comparisons tend to be more widely applicable than absolute, isolated results.



LITERATURE CITED

- Ackerman, R. F.
1957. The effect of various seed-bed treatments on the germination and survival of white spruce and lodgepole pine seedlings. Can. Dep. Northern Affairs and Nat. Resour., For. Branch, For. Res. Div. Tech. Note No. 63, 23 p.
- Alban, David H.
1969. The influence of western hemlock and western redcedar on soil properties. Soil Sci. Soc. Amer. Proc. 33:453-457.
- Aldhous, J. R., and A. J. Low.
1974. The potential of western hemlock, western red cedar, grand fir and noble fir in Britain. For. Comm. Bull. No. 49, 105 p. London.
- Allen, G. S., and W. Bientjes.
1954. Studies on coniferous tree seed at the University of British Columbia. For. Chron. 30(2):183-196.
- Aller, Alvin R.
1956. A taxonomic and ecologic study of the flora of Monument Peak, Oregon. Amer. Midl. Nat. 56:454-472.
- Anderson, Eric A.
1951. Tracheid length variation in conifers as related to distance from pith. J. For. 49:38-42.
- Atzet, Thomas, and R. H. Waring.
1970. Selective filtering of light by coniferous forests and minimum light energy requirements for regeneration. Can. J. Bot. 48(12):2163-2167.
- Baker, Frederick S.
1929. Effect of excessively high temperatures on coniferous reproduction. J. For. 27:949-975.
- Baker, Frederick S.
1949. A revised tolerance table. J. For. 47:179-181.
- Barker, John E.
1973. Diurnal patterns of water potential in *Abies concolor* and *Pinus ponderosa*. Can. J. For. Res. 3(4):556-564.
- Bawcom, Richard H., Robert J. Hubbell, and David M. Burns.
1961. Seasonal diameter growth in trees on Jackson State Forest. Calif. Div. For. State For. Notes No. 6, 5 p. (Mimeo)
- Beaton, J. D., G. Brown, R. C. Speer, I. MacCrae, W. P. T. McGhee, A. Moss, and R. Kosick.
1965. Concentration of micro-nutrients in foliage of three coniferous tree species in British Columbia. Soil Sci. Soc. Amer. Proc. 29:299-302.
- Beaton, J. D., A. Moss, I. MacCrae, J. W. Konkin, W. P. T. McGhee, and R. Kosick.
1965. Observations on foliage nutrient content of several coniferous tree species in British Columbia. For. Chron. 41(2):222-236.
- Bega, Robert V.
1962. Tree killing by *Fomes annosus* in a genetics arboretum. USDA Plant Dis. Rep. 46:107-110.
- Bega, Robert V., and Richard S. Smith, Jr.
1966. Distribution of *Fomes annosus* in natural forests of California. USDA Plant Dis. Rep. 50:832-836.
- Benzian, Blanche.
1965. Experiments on nutrition problems in forest nurseries. For. Comm. Bull. No. 37, Vol. I. 251 p. London.
- Bergdahl, D. R., and D. W. French.
1976. Susceptibility of five pine species, 2 to 36 months of age, to infection by *Cronartium comandrae*. Can. J. For. Res. 6:319-325.
- Berndt, Herbert W., and Robert D. Gibbons.
1958. Root distribution of some native trees and understory plants growing on three sites within ponderosa pine watersheds in Colorado. USDA For. Serv. Pap. No. 37, 14 p. Rocky Mt. For. and Range Exp. Stn., Fort Collins, Colo.
- Berntsen, Carl Martin.
1967. Relative low temperature tolerance of lodgepole and ponderosa pine seedlings. 158 p. Ph.D. Thesis. Oreg. State Univ., Corvallis.
- Bever, Dale N.
1954. Evaluation of factors affecting natural reproduction of forest trees in central western Oregon. Oreg. State Board of For., Res. Bull. No. 3, 49 p.

- Boe, Kenneth N.
1953. Western larch and Douglas-fir seed dispersal into clear-cuttings. USDA For. Serv. Res. Note 129, 3 p. Northern Rocky Mountain For. and Range Exp. Stn., Fort Collins, Colo.
- Boe, Kenneth N.
1954. Periodicity of cone crops for five Montana conifers. Mont. Acad. Sci. Proc. 14, p. 5-9.
- Boggie, Robert.
1974. Response of seedlings of *Pinus contorta* and *Picea sitchensis* to oxygen concentration in culture solution. New Phytol. 73:467-473.
- Bollen, Walter B., Chi-sin Chen, Kuo C. Lu, and Robert F. Tarrant.
1967. Influence of red alder on fertility of a forest soil: microbial and chemical effects. Oreg. For. Res. Lab., Res. Bull. No. 12, 61 p. Oreg. State Univ., Corvallis.
- Bones, James T.
1962. Relating outside- to inside-bark diameter at the top of first 16-foot log for southeast Alaska timber. USDA For. Serv. Tech. Note No. 52, 2 p. Northern For. Exp. Stn., Juneau, Alaska.
- Born, J. David.
1966. Specific gravity of increment cores from interior Alaska trees. USDA For. Serv. Res. Note NOR-19, 4 p. Northern For. Exp. Stn., Juneau, Alaska.
- Boyce, J. S.
1929. Deterioration of wind-thrown timber on the Olympia peninsula, Wash. USDA Tech. Bull. No. 104, 28 p.
- Boyd, Raymond J., Jr.
1959. Silvics of western redcedar. USDA For. Serv. Misc. Publ. No. 20, 14 p. Intermountain For. and Range Exp. Stn., Ogden, Utah.
- Boyd, R. J., and G. H. Deitschman.
1969. Site preparation aids natural regeneration in western larch - Engelmann spruce strip clearcuttings. USDA For. Serv. Res. Pap. INT-64, 10 p. Intermountain For. and Range Exp. Stn., Ogden, Utah.
- Brayshaw, T. C.
1970. The dry forests of southern British Columbia. Syesis 3:17-43.
- Brink, V. C.
1954. Survival of plants under flood in the lower Fraser River Valley, British Columbia. Ecology 35:94-95.
- British Columbia Forest Service.
1950. Silvicultural studies. Rep. B.C. For. Serv. (1949):25-28.
- Brix, H.
1971. Growth response of western hemlock and Douglas-fir seedlings to temperature regimes during day and night. Can. J. Bot. 49(2):289-294.
- Brix, Holger.
1972. Growth response of Sitka spruce and white spruce seedlings to temperature and light intensity. Dep. Environ., Can. For. Serv. Inform. Rep. BC-X-74, 17 p. For. Res. Centre, Victoria, B.C.
- Brown, James K.
1976. Predicting crown weights for 11 Rocky Mountain conifers. Oslo Biomass Studies, IUFRO Congr., June 1976:103-115.
- Brown, James K.
1978. Weight and density of crown of Rocky Mountain conifers. USDA For. Serv. Res. Pap. INT-197, 56 p. Intermountain For. and Range Exp. Stn., Ogden, Utah.
- Buckland, D. C.
1956. Terminal shoot growth of four western conifers for a single growing season. For. Chron. 32:397-399.
- Buckland, D. C., R. E. Foster, and V. J. Nordin.
1949. Studies in forest pathology. VII. Decay in western hemlock and fir in the Franklin River area, British Columbia. Can. J. Res. Sect. C. 27:312-331.
- Burroughs, Edward R., Jr. and Byron R. Thomas.
1977. Declining root strength in Douglas-fir after felling as a factor in slope stability. USDA For. Serv. Res. Pap. INT-190, 27 p. Intermountain For. and Range Exp. Stn., Ogden, Utah.
- Carkin, Richard E., Jerry F. Franklin, Jack Booth, and Clark E. Smith.
1978. Seeding habits of upper-slope tree species IV. Seed flight of noble fir and Pacific silver fir. USDA For. Serv. Res.

- Note PNW-312, 10 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Carolin, V. M., and W. K. Coulter.
1975. Comparison of western spruce budworm populations and damage on grand fir and Douglas-fir trees. USDA For. Serv. Res. Pap. PNW-195, 16 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Carpenter, Edwin D.
1970. Salt tolerance of ornamental plants. Amer. Nurseryman 131(2):12, 54, 71.
- Carter, F. L., and R. V. Smythe.
1974. Feeding and survival responses of *Reticulitermes flavipes* (Kollar) to extractives of wood from eleven coniferous genera. Holzforschung 28(2):41-45.
- Cayford, J. H., and A. Bickerstaff.
1968. Man-made forests in Canada. Can. For. Br. Publ. No. 1240, 68 p.
- Chandler, Craig C.
1960. Slash weight tables for westside mixed conifers. USDA For. Serv. Pac. Southwest For. and Range Exp. Stn. Tech. Pap. No. 48, 21 p. Pac. Southwest For. and Range Exp. Stn., Berkeley, Calif.
- Childs, Thomas W.
1939. Decay of slash on clearcut areas in the Douglas-fir region. J. For. 37(12):955-959.
- Childs, Thomas W., and J. L. Bedwell.
1948. Susceptibility of some white pine species to *Cronartium ribicola* in the Pacific Northwest. J. For. 46:595-599.
- Clark, Joe W.
1957. Comparative decay resistance of some common pines, hemlock, spruce, and true fir. For. Sci. 3:314-320.
- Clark, John B., and Geoffrey R. Lister.
1975. Photosynthetic action spectra of trees. I. Comparative photosynthetic action spectrum of one deciduous and four coniferous tree species as related to the photorespiration and pigment complements. II. The relationship of cuticle structure to the visible and ultraviolet spectral properties of needles from four coniferous species. Plant Phys. 55(2):401-413.
- Cleary, Brian Dennis.
1971. The effect of plant moisture stress on physiology and establishment of planted Douglas-fir and ponderosa pine seedlings. 85 p. (Ph.D. Thesis, on file Oreg. State Univ., Corvallis.)
- Cleary, B. D., and R. H. Waring.
1969. Temperature: collection of data and its analysis for the interpretation of plant growth and distribution. Can. J. Bot. 47(1):167-173.
- Cochran, Patrick Holmes.
1963. First year development of three conifers on surface and exposed subsoil horizons. 69 p. M.S. Thesis. Oreg. State Univ., Corvallis.
- Cochran P. H.
1972. Tolerance of lodgepole and ponderosa pine seeds and seedlings to high water tables. Northwest Sci. 46(4):322-331.
- Cochran, P. H., and Carl M. Berntsen.
1973. Tolerance of lodgepole and ponderosa pine seedlings to low night temperatures. For. Sci. 19(4):272-280.
- Coleman, Babette Brown, Walter C. Muenscher, and Donald R. Charles.
1956. A distributional study of the epiphytic plants of the Olympic Peninsula, Washington. Amer. Midl. Nat. 56(1):54-87.
- Conkle, M. Thompson, W. J. Libby, and J. L. Hamrick III.
1967. Winter injury among white fir seedlings--unusual pattern in seed source study. USDA For. Serv. Res. Note PSW-138, 7 p. Pac. Southwest For. and Range Exp. Stn., Berkeley, Calif.
- Cooper, William S.
1911. Reproduction by layering among conifers. Bot. Gaz. 52:369-379.
- Cooper, William S.
1931. The layering habit in Sitka spruce and the two western hemlocks. Bot. Gaz. 91:441-451.
- Cordes, Lawrence David.
1973. An ecological study of the Sitka spruce forest on the west coast of Vancouver Island. 395 p. Ph.D. Thesis, Univ. of Brit. Col., Vancouver.
- Cowan, Ian McTaggart.
1945. The ecological relationships of the food of the Columbian

- black-tailed deer, *Odocoileus hemionus columbianus* (Richardson), in the coast forest region of southern Vancouver Island, British Columbia. *Ecol. Mongr.* 15(2):109-139.
- Crawford, R. M. M., and Margaret A. Baines.
1977. Tolerance of anoxia and the metabolism of ethanol in tree roots. *New Phytol.* 79(3):519-526.
- Dahms, Walter G., and James W. Barrett.
1975. Seed production of central Oregon ponderosa and lodgepole pines. *USDA For. Serv. Res. Pap. PNW-191*, 13 p. Pac. Northwest For. and Range Exp. Stn., Portland, Ore.
- Daniel, T. W., and Josef Schmidt.
1972. Lethal and nonlethal effects of the organic horizons of forested soils on the germination of seeds from several associated conifer species of the Rocky Mountains. *Can. J. For. Res.* 2(3):179-184.
- Daubenmire, R. F.
1943. Soil temperatures versus drought as a factor determining lower altitudinal limits of trees in the Rocky Mountains. *Bot. Gaz.* 105:1-13.
- Daubenmire, R. F.
1946. Radial growth of trees at different altitudes. *Bot. Gaz.* 107(4):462-467.
- Daubenmire, R.
1953. Nutrient content of leaf litter of trees in the northern Rocky Mountains. *Ecology* 34:786-793.
- Daubenmire, R.
1957. Injury to plants from rapidly dropping temperature in Washington and northern Idaho. *J. For.* 55(8):581-585.
- Daubenmire, R. F., and M. E. Deters.
1947. Comparative studies of growth in deciduous and evergreen trees. *Bot. Gaz.* 190:1-12.
- Davidson, J. G. N.
1959. Ecological requirements of grand fir and Pacific silver fir. 1958 Prog. Rep., N.R.C. Grant No. T-92:12-15.
- Day, R. J.
1964. The microenvironments occupied by spruce and fir regeneration in the Rocky Mountains. *Can. Dep. For. Publ. No. 1037*, 25 p.
- Day, W. R.
1928. Damage by late frost on Douglas fir, Sitka spruce and other conifers. *Forestry* 2:19-30.
- Day, W. R.
1957. Sitka spruce in British Columbia. *Imp. For. Comm. Bull.* No. 28, 110 p.
- Deitschman, Glenn H., and Alan W. Green.
1965. Relations between western white pine site index and tree height of several associated species. *USDA For. Serv. Res. Pap. INT-22*, 28 p. Intermountain For. and Range Exp. Stn., Ogden, Utah.
- Dick, James.
1960. A direct seeding of Pacific silver fir. *Weyerhaeuser Co. For. Res. Note* 33, 4 p.
- Dobbs, Robert C.
1972. Regeneration of white and Engelmann spruce: a literature review with special reference to the British Columbia interior. *Can. For. Serv. British Columbia Inf. Rep. BC-X-69*, 77 p. Pac. For. Res. Centre, Victoria, B.C.
- Duffield, John W.
1956. Damage to western Washington forests from November 1955 cold wave. *USDA For. Serv. Pac. Northwest For. and Range Exp. Stn. Res. Note No. 129*, 5 p. Pac. Northwest For. and Range Exp. Stn., Portland, Ore.
- Ebell, Lorne F., and Ralph L. Schmidt.
1959. Effect of elevation and climatic factors on production and dispersal of coniferous tree pollen. *Proc. Soc. Amer. For.* 1959:39. (Abstract.)
- Edmonds, Robert L. (ed.).
1975. Coniferous forest biome progress report, July 1974-August 1975. *Coniferous For. Biome Intern. Rep.* 162, 97 p. Coll. of For. Resour., Univ. of Wash., Seattle.
- Edwards, Milton B.
1957. California black oak--its management and economic possibilities. *J. For.* 55(7):506-510.

- Eis, S.
1970. Root-growth relationships of juvenile white spruce, alpine fir, and lodgepole pine on three soils in the interior of British Columbia. Can. Dep. Fish., Forestry, Can. For. Serv. Publ. No. 1276, 10 p.
- Eis, S.
1972. Root grafts and their silvicultural implications. Can. J. For. Res. 2:111-120.
- Eis, S.
1974. Root system morphology of western hemlock, western red cedar, and Douglas-fir. Can. J. For. Res. 4(1):28-38.
- Emmingham, W. H., and R. H. Waring.
1973. Conifer growth under different light environments in the Siskiyou Mountains of southwestern Oregon. Northwest Sci. 47(2):88-99.
- Englerth, George H., and Theodore C. Scheffer.
1954. Tests of decay resistance of four western pole species. USDA For. Serv., For. Prod. Lab. Rep. No. 2006, 12 p. Madison, Wisc.
- Fahnestock, George R.
1960. Logging slash flammability. USDA For. Serv. Intermtn. For. and Range Exp. Stn. Res. Pap. No. 58, 67 p. Intermtn. For. and Range Exp. Stn., Ogden, Utah.
- Fairbairn, W. A., and S. A. Neustein.
1970. Study of response of certain coniferous species to light intensity. Forestry 43:57-71.
- Farr, Wilbur A.
1973. Specific gravity of western hemlock and Sitka spruce in southeast Alaska. Wood Sci. 6(1):9-13.
- Farr, Wilbur A., Vernon J. Labau, and Thomas H. Laurent.
1976. Estimation of decay in old-growth western hemlock and Sitka spruce in southeast Alaska. USDA For. Serv. Res. Pap. PNW-204, 24 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Ferrell, William K., and E. Steve Woodward.
1966. Effects of seed origin on drought resistance of Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco). Ecology 47:499-503.
- Fisher, George M.
1935. Comparative germination of tree species on various kinds of surface-soil material in the western white pine type. Ecology 16:606-611.
- Flint, Howard R.
1925. Fire resistance of northern Rocky Mountain conifers. Idaho Forester 7:7-10, 41-43.
- Forest Service, USDA.
1974. Seeds of woody plants in the United States. USDA For. Serv. Agric. Handb. No. 450, 883 p. (C. S. Schopmeyer, Tech. Coord.)
- Forest Soils Committee of the Douglas-fir Region.
1957. An introduction to forest soils of the Douglas-fir region of the Pacific Northwest. 217 p. Seattle, Wash.
- Forristall, Floyd F., and S. P. Gessel.
1955. Soil properties related to forest cover type and productivity on the Lee Forest, Snohomish County, Washington. Soil Sci. Soc. Amer. Proc. 19:384-389.
- Foulger, A. N., Frank Freese, and Joan E. Lengel.
1976. Solid wood content of western softwood logging residues. USDA For. Serv. Res. Pap. FPL-253, 7 p. For. Prod. Lab., Madison, Wisc.
- Fowells, H. A.
1940. Cutworm damage to seedlings in California pine stands. J. For. 38(7):590-591.
- Fowells, H. A.
1941. The period of seasonal growth of ponderosa pine and associated species. J. For. 39:601-608.
- Fowells, H. A., and G. H. Schubert.
1956. Seed crops of forest trees in the pine region of California. U.S. Dep. Agric. Tech. Bull. No. 1150, 48 p.
- Fowells, H. A., and N. B. Stark.
1965. Natural regeneration in relation to environment in the mixed conifer forest type of California. USDA For. Serv. Res. Pap. PSW-24, 14 p. Pac. Southwest For. and Range Exp. Stn., Berkeley, Calif.
- Franklin, Jerry F.
1961. A guide to seedling identification for 25 conifers of the Pacific Northwest. USDA

- For. Serv. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg. 65 p., illus.
- Franklin, Jerry F.
1964. Ecology and silviculture of the true fir-hemlock forests of the Pacific Northwest. Soc. Amer. Foresters Proc. 1964:28-32.
- Franklin, Jerry F.
1968. Cone production by upper slope conifers. USDA For. Serv. Res. Pap. PNW-60, 21 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Franklin, Jerry F., Richard Carkin, and Jack Booth.
1974. Seeding habits of upper-slope tree species. I. A 12-year record of cone production. USDA For. Serv. Res. Note PNW-213, 12 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Franklin, Jerry F., and C. T. Dyrness.
1973. Natural vegetation of Oregon and Washington. USDA For. Serv. Gen. Tech. Rep. PNW-8, 417 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Franklin, Jerry F., C. T. Dyrness, Duane G. Moore, and Robert F. Tarrant.
1968. Chemical soil properties under coastal Oregon stands of alder and conifers. *In* Biology of Alder, J. M. Trappe, J. F. Franklin, R. F. Tarrant, and G. M. Hansen (eds.). Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Franklin, Jerry F., and John Hoffman.
1968. Two tests of white pine, true fir, and Douglas-fir seed-spotting in the Cascade Range. USDA For. Serv. Res. Note PNW-80, 11 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Franklin, Jerry F., and Russel G. Mitchell.
1967. Successional status of subalpine fir in the Cascade Range. USDA For. Serv. Res. Pap. PNW-46, 16 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Franklin, Jerry F., and Gary A. Ritchie.
1970. Phenology of cone and shoot development of noble fir and some associated true firs. For. Sci. 16:356-364.
- Fraser, Bruce E. C., and Lawrence D. Cordes.
1967. Sap pressure of active and dormant plants of three Pacific Northwest conifers. Appendix A. *In* 1967 Progress Rep., Nat. Res. Counc. Grant No. A-92, V. J. Krajina (ed.), Dep. Bot., Univ. British Columbia, p. 4-12.
- Freeland, R. O.
1952. Effect of age of leaves upon the rate of photosynthesis in some conifers. Plant Physiol. 27:685-690.
- Gail, Floyd W., and E. M. Long.
1935. A study of site, root development and transpiration in relation to the distribution of *Pinus contorta*. Ecology 16:88-100.
- Garman, E. H.
1951. Seed production by conifers in the coastal region of British Columbia related to dissemination and regeneration. British Columbia For. Serv. Tech. Publ. T.35, 48 p. Victoria, British Columbia.
- Garman, E. H.
1955. Regeneration problems and their silvicultural significance in the coastal forests of British Columbia. British Columbia For. Serv. Tech. Publ. T.41, 67 p. Victoria, British Columbia.
- Garrison, G. A., J. M. Skovlin, G. E. Poulton, and A. H. Winward.
1976. Northwest plant names and symbols for ecosystem inventory and analysis. USDA For. Serv. Gen. Tech. Rep. PNW-46, 263 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Gashwiler, Jay S.
1967. Conifer seed survival in a western Oregon clearcut. Ecology 48:431-438.
- Gashwiler, Jay S.
1969. Seedfall of three conifers in west-central Oregon. For. Sci. 15(3):290-295.
- Gashwiler, Jay S.
1970. Further study of conifer seed survival in a western Oregon clearcut. Ecology 51:849-854.

- Gessel, Stanley P., Richard B. Walker, and Philip G. Haddock.
1951. Preliminary report on mineral deficiencies in Douglas-fir and western red cedar. Soil Sci. Soc. Amer. Proc. 15:364-369.
- Gill, C. J.
1970. The flooding tolerance of woody species--a review. For. Abstr. 31(4):671-688.
- Godman, R. M.
1949. What kind of trees make the best growth in southeast Alaska. USDA For. Serv. Alaska For. Res. Center Tech. Note No. 2, 1 p.
- Godman, R. M.
1953a. Moss retards regeneration in southeast Alaska. USDA For. Serv. Alaska For. Res. Center Tech. Note No. 18, 1 p.
- Godman, R. M.
1953b. Seed dispersal in southeast Alaska. USDA For. Serv. Alaska For. Res. Center Tech. Note No. 16, 2 p.
- Godman, Richard M., and Robert A. Gregory.
1953. Seasonal distribution of radial and leader growth in the Sitka spruce-western hemlock forests of southeast Alaska. J. For. 53:827-833.
- Gordon, Donald T.
1970. Natural regeneration of white and red fir...influence of several factors. USDA For. Serv. Res. Pap. PSW-58, 32 p. Pac. Southwest For. and Range Exp. Stn., Berkeley, Calif.
- Gratkowski, H. J.
1956. Windthrow around staggered settings in old-growth Douglas-fir. For. Sci. 2(1):60-74.
- Gratkowski, H.
1975. Silvicultural use of herbicides in Pacific Northwest forests. USDA For. Serv. Gen. Tech. Rep. PNW-37, 44 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Gregory, R. A.
1957. 1956 cone crop report for Alaska tree species. USDA For. Serv. Alaska For. Res. Center Tech. Note No. 35, 2 p.
- Gregory, R. A.
1958. 1957 cone crop report for Alaska tree species. USDA For. Serv. Alaska For. Res. Center Tech. Note No. 41, 2 p.
- Gregory, R. A.
1959. 1958 cone crop report for Alaska tree species. USDA For. Serv. Alaska For. Res. Center Tech. Note No. 44, 2 p.
- Grier, Charles C., and Richard H. Waring.
1974. Conifer foliage mass related to sapwood area. For. Sci. 20(3):205-206.
- Habeck, James R.
1968. Forest succession in the Glacier Park cedar-hemlock forests. Ecology 49:872-880.
- Hadfield, James S., and David W. Johnson.
Undated. Laminated root rot. A guide for reducing and preventing losses in Oregon and Washington forests. USDA For. Serv., Pac. Northwest Region. 16 p.
- Haig, Irvine T.
1936. Factors controlling initial establishment of western white pine and associated species. Yale Univ. Sch. For. Bull. No. 41, 149 p.
- Haig, Irvine T., Kenneth P. Davis, and Robert H. Weidman.
1941. Natural regeneration in the western white pine type. USDA For. Serv. Tech. Bull. 767, 99 p. Northern Rocky Mountain For. and Range Exp. Stn., Fort Collins, Colo.
- Haller, John R.
1959. Factors affecting the distribution of ponderosa and jeffrey pines in California. Madrono 15:65-71
- Haller, John R.
1961. Some recent observations on ponderosa, jeffrey, and washoe pines in northeastern California. Madrono 16(4):126-132.
- Hanzlik, E. J.
1925. A preliminary study of the growth of noble fir. J. Agric. Res. 31:929-934.
- Hanzlik, E. J.
1932. Type successions in the Olympic Mountains. J. For. 30:91-93.
- Harlow, William M., and Ellwood S. Harrar.
1950. Textbook of dendrology. McGraw-Hill Book Co., Inc. New York. Third Edition. 555 p.
- Harris, A. S.
1960. 1959 cone crop report for Alaska tree species. USDA For.

- Serv. Alaska For. Res. Center
Tech. Note No. 50, 2 p.
- Harris, A. S.
1962. Cone crops in coastal Alaska--1960 and 1961. USDA For. Serv. Alaska For. Res. Center Tech. Note No. 53, 4 p.
- Harris, A. S.
1969. Ripening and dispersal of a bumper western hemlock-Sitka spruce seed crop in southeast Alaska. USDA For. Serv. Res. Note PNW-105, 11 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Harris, Arland S., and Wilbur A. Farr.
1974. The forest ecosystem of southeast Alaska. 7. Forest ecology and timber management. USDA For. Serv. Gen. Tech. Rep. PNW-25, 109 p., illus. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Heilman, Paul, and Gordon L. Ekuan.
1973. Reponse of Douglas-fir and western hemlock seedlings to lime. For. Sci. 19(3):220-224.
- Heit, C. E.
1968. Thirty-five years' testing of tree and shrub seed. J. For. 66(8):632-634.
- Hellmers, Henry.
1964. Distribution of growth in tree seedling stems as affected by temperature and light. p. 533-547. In The formation of wood in forest trees. Martin H. Zimmerman, (ed.). Academic Press, N.Y. 562 p.
- Hellmers, Henry, and William P. Sundahl.
1959. Response of *Sequoia sempervirens* (D. Don) Endl. and *Pseudotsuga menziesii* (Mirb.) Franco seedlings to temperature. Nature 194(4694):1247-1248. London.
- Hermann, Richard K., and Denis P. Lavender.
1967. Tolerance of coniferous seedlings to silting. J. For. 65(11):824-825.
- Hetherington, J. C.
1965. The dissemination, germination, and survival of seed on the west coast of Vancouver Island from western hemlock and associated species. British Columbia For. Serv. Res. Note 39, 22 p.
- Hodges, John D.
1967. Patterns of photosynthesis under natural environmental conditions. Ecology 48:234-242.
- Hodges, John D., and David R. M. Scott.
1968. Photosynthesis in seedlings of six conifer species under natural environmental conditions. Ecology 49:973-981.
- Hoff, R. J., and G. I. McDonald.
1977. Differential susceptibility of nineteen white pine species to woolly aphid (*Pineus coloradensis*). USDA For. Serv. Res. Note INT-225, 6 p. Intermountain For. and Range Exp. Stn., Ogden, Utah.
- Hofmann, J. V.
1925. Laboratory tests on effect of heat on seeds of noble and silver fir, western white pine, and Douglas-fir. J. Agric. Res. 31:197-199.
- Holmes, G. D., and G. Buszewicz.
1958. The storage of seed of temperate forest tree species. Part II. For. Abstr. 19:455-476.
- Hornqvist, Richard, and Hakon Robak.
1975. Relative susceptibility of eleven conifer species to fluoride air pollution. Norsk Institutt for Skogforskning. Rep. of the Norwegian For. Res. Inst. 32.5:189-206.
- Howell, Joseph.
1931. Clay pans in the western yellow pine type. J. For. 29:962-963.
- Hustich, Ilmari.
1953. The boreal limits of conifers. Arctic 6(2):149-162.
- Isaac, Leo A.
1930. Seed flight in the Douglas-fir region. J. For. 28:492-499.
- Isaac, Leo A.
1940. Life of seed in the forest floor. USDA For. Serv. Res. Note. No. 31, p. 14. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Isaac, Leo A.
1943. Reproductive habits of Douglas-fir. Charles Lathrop Pack Foundation, Washington, D.C. 107 p.
- Jablanczy, Alexander.
1963. Influence of slash burning on the establishment and initial growth of seedlings of Douglas-fir, western hemlock, and western redcedar. Appendix I. In

- Ecology of the forests of the Pacific Northwest. V. J. Krajina (ed.). 1962 progress rep., Nat. Res. Counc. Grant No. T-92, Dep. Biol., Bot., Univ. British Columbia, p. 70-77.
- James, G. A., and G. L. Hayes. 1954. Highlights of a Port Orford-cedar regeneration study. J. For. 52(11):852-855.
- Jarvis, Paul Gordon. 1963. Comparative studies in plant water relations. Abstracts of Uppsala Dissertations in Science 27, 13 p.
- Johnson, Norman E. 1958. Ambrosia beetle infestation of coniferous logs on clear-cuttings in northwestern Oregon. J. For. 56(7):508-511.
- Johnstone, W. D. 1976. Juvenile height growth of white spruce and lodgepole pine following logging and scarification in west-central Alberta. Can. For. Serv. Inform. Rep. NOR-X-171, 10 p. Northern For. Res. Centre, Edmonton, Alberta.
- Kangur, Rudolf. 1973. Snow damage to young western hemlock and Douglas-fir. Oreg. For. Res. Lab. Res. Pap. 21, 11 p. Oreg. State Univ., Corvallis.
- Kaufmann, Merrill R., and Alan N. Eckard. 1977. Water potential and temperature effects on germination of Engelmann spruce and lodgepole pine seeds. For. Sci. 23(1):27-33.
- Kerr, H. S. 1913. Notes on the distribution of lodgepole and yellow pine in the Walker Basin. For. Quart. 11:509-515.
- Kotar, John. 1972. Ecology of *Abies amabilis* in relation to its altitudinal distribution and in contrast to its common associate *Tsuga heterophylla*. 188 p. Ph.D. Thesis. Univ. of Wash., Seattle.
- Krajina, V. J. 1955. Transpiration and shade-tolerance of trees. (Abstract.) Ecol. Soc. Amer. Bull. 36(2):51.
- Krajina, V. J. 1959. Ecological requirements of Douglas-fir, western hemlock, Sitka spruce, and western redcedar. 1958 Progress Rep., N.R.C. Grant, No. T-92, p. 1-5.
- Krajina, V. J., (ed.). 1965. Ecology of western North America. Vol. 1. Univ. of British Columbia, Vancouver. 112 p.
- Krajina, V. J. (ed.). 1970. Ecology of forest trees in British Columbia. In Ecology of western North America 2(1):1-146.
- Krajina, V. J. 1971. $\text{NH}_4:\text{NO}_3$ in nitrogen economy of conifers in Douglas-fir forests of the Pacific Northwest, America. Pac. Sci. Assoc. Twelfth Congr. Proc. 1971:44. (Abstract.)
- Krajina, V. J., Sarah Madoc-Jones, and Gary Mellor. 1973. Ammonium and nitrate in the nitrogen economy of some conifers growing in Douglas-fir communities of the Pacific Northwest of America. Soil Biol. Biochem. 5(1):143-147.
- Krueger, Kenneth W., and Robert H. Ruth. 1969. Comparative photosynthesis of red alder, Douglas-fir, Sitka spruce, and western hemlock seedlings. Can. J. Bot. 47:519-527.
- Krygier, James T., and Robert H. Ruth. 1961. Effect of herbicides on salmonberry and on Sitka spruce and western hemlock seedlings. Weeds 9:416-422.
- Landis, Thomas D. 1976. Foliage nutrient levels for three Rocky Mountain tree species. Tree Planters' Notes 27(2):4-5.
- Lanquist, Karl B. 1946. Tests of seven principal forest tree seeds in northern California. J. For. 44:1063-1066.
- Larsen, J. A. 1930. Forest types of the northern Rocky Mountains and their climatic controls. Ecology 11:631-672.
- Lassen, L. E., and E. A. Okkonen. 1969. Sapwood thickness of Douglas-fir and five other western softwoods. USDA For. Serv. Res. Pap. FPL-124, 16 p. For. Prod. Lab., Madison, Wisc.
- Lavender, Denis Peter. 1962. The growth of seedlings of some coniferous species in a controlled environment. 172 p. Ph.D. thesis, Oreg. State Univ. Corvallis.

- Leaphart, Charles D.
1958. Root characteristics of western white pine and associated tree species in a stand affected with pole blight of white pine. USDA For. Serv. Res. Pap. No. 52, 10 p. Intermountain For. and Range Exp. Stn., Odgen, Utah.
- Leaphart, Charles D., and Ed F. Wicker.
1966. Explanation of pole blight from responses of seedlings grown in modified environments. Can. J. Bot. 44:121-137.
- Le Barron, Russell K., and George M. Jemison.
1953. Ecology and silviculture of the Engelmann spruce-alpine fir type. J. For. 51:349-355.
- Levitt, J.
1951. Frost, drought, and heat resistance. Ann. Rev. Plant Physiol. 2:245-268.
- Li, C. Y., K. C. Lu, J. M. Trappe, and W. B. Bollen.
1972. Nitrate-reducing capacity of roots and nodules of *Alnus rubra* and roots of *Pseudotsuga menziesii*. Plant and Soil 37(2):409-414.
- Little, Elbert L., Jr.
1953. Checklist of native and naturalized trees of the United States (including Alaska). USDA For. Serv. Agric. Handb. No. 41, 472 p.
- Lopushinsky, William.
1969. Stomatal closure in conifer seedlings in response to leaf moisture stress. Bot. Gaz. 130:258-263.
- Lopushinsky, W., and G. O. Klock.
1974. Transpiration of conifer seedlings in relation to soil water potential. For. Sci. 20(2):181-186.
- Lowry, Robert Franklin.
1972. Ecology of subalpine zone tree clumps in the North Cascade Mountains of Washington. Diss. Abstr. B 33(3):1876-1877.
- Madoc-Jones, Sarah.
1970. The nitrogen nutrition of some conifers of British Columbia. Appendix D. 1970 Progress Rep., Nat. Res. Council. Grant No. A-92, p. 16-23. Univ. British Columbia, Bot. Dep., Vancouver.
- Malcolm, D. C., and E. A. Caldwell.
1971. Environmental effects on shoot growth in conifers. Rep. For. Res. For. Comm. 1970/71:141. London.
- Markstrom, Donald C., and Robert A. Hann.
1972. Seasonal variation in wood permeability and stem moisture content of three Rocky Mountain softwoods. USDA For. Serv. Res. Note RM-212, 7 p. Rocky Mountain For. and Range Exp. Stn., Fort Collins, Colo.
- Markwardt, L. J.
1931. The distribution and the mechanical properties of Alaska woods. U.S. Dep. Agric. Tech. Bull. 226, 80 p.
- Markwardt, L. J., and T. R. C. Wilson.
1935. Strength and related properties of wood grown in the United States. U.S. Dep. Agric. Tech. Bull. 479, 99 p.
- Marshall, Robert.
1931. An experimental study of the water relations of seedling conifers with special reference to wilting. Ecol. Mongr. 1:37-98.
- Mason, D. T.
1915. The life history of lodge-pole pine in the Rocky Mountains. U.S. Dep. Agric. Bull. No. 154, 35 p.
- McNaughton, G. C.
1944. Ignition and charring temperatures of wood. USDA For. Serv. For. Prod. Lab. Rep. 1464, 3 p. For. Prod. Lab., Madison, Wisc.
- Mellow, G. E., and E. B. Tregunna.
1972. The relationship between leaf area and leaf dry weight of three conifer species grown on three sources of nitrogen. Can. J. For. Res. 2(3):377-379.
- Meyer, Walter H.
1937. Yield of even-aged stands of Sitka spruce and western hemlock. U.S. Dep. Agric. Tech. Bull. 544, 86 p.
- Miller, Paul R., and Joe R. McBride.
1975. Effects of air pollutants on forests. p. 195-235. In Responses of plants to air pollution. T. T. Kozlowski, and J. B. Mudd (eds.). Academic Press, New York. 383 p.
- Miller, Paul Robert, and Arthur A. Millecan.
1972. Extent of oxidant air pollution damage to some pines and

- other conifers in California. Plant Dis. Rep. 55:555-559.
- Minore, Don.
1968. Effects of artificial flooding on seedling survival and growth of six northwestern tree species. USDA For. Serv. Res. Note PNW-92, 12 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Minore, Don.
1970. Seedling growth of eight northwestern tree species over three water tables. USDA For. Serv. Res. Note PNW-115, 8 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Minore, Don.
1972. Germination and early growth of coastal tree species on organic seed beds. USDA For. Serv. Res. Pap. PNW-135, 18 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Minore, Don, and Clark E. Smith.
1971. Occurrence and growth of four northwestern tree species over shallow water tables. USDA For. Serv. Res. Note PNW-160, 9 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Minore, Don, Clark E. Smith, and Robert F. Woollard.
1969. Effects of high soil density on seedling root growth of seven northwestern tree species. USDA For. Serv. Res. Note PNW-112, 6 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Mitchell, A. F.
1965. The growth in early life of the leading shoot of some conifers. Forestry 38(1):121-136.
- Mitchell, Russell G.
1966. Infestation characteristics of the balsam woolly aphid in the Pacific Northwest. USDA For. Serv. Res. Pap. PNW-35, 18 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oregon.
- Monk, Ralph W., and Herman H. Wiebe.
1961. Salt tolerance and protoplasmic salt hardness of various woody and herbaceous ornamental plants. Plant Physiol. 36(4):478-482.
- Morgan, Paul Dexter.
1973. The occurrence of rhizina root rot in western Washington. M.S. thesis, Univ. of Wash., Seattle. 302 p.
- Mosher, Milton M.
1965. The growth of larch in northeastern Washington. Wash. State Univ., Coll. of Agric. Circ. 456, 6 p.
- Munger, T. T.
1940. The cycle from Douglas-fir to hemlock. Ecology 21:451-459.
- Muri, Glen.
1955. The effect of simulated slash burning on germination, primary survival and top/root ratios of Engelmann spruce and alpine fir. Univ. British Columbia For. Club Res. Note No. 14, 7 p. Vancouver.
- Murison, W. F.
1959. Ecological requirements of ponderosa pine, western white pine, coastal and interior Douglas-fir, and Engelmann spruce. 1958 Progress Rep., N.R.C. Grant No. T-92, p. 6-11.
- Murison, W. F.
1961. Macronutrient deficiency and its effect on coniferous growth. Appendix A. 1960 Progress Rep. N.R.C. Grant No. T-92, Ecology of the forests of the Pacific Northwest, V. J. Krajina (ed.), p. 4-6.
- Newton, Michael.
1963. Some herbicide effects on potted Douglas-fir and ponderosa pine seedlings. J. For. 61:674-676.
- Newton, Michael, B. A. El Hassan, and Jaroslav Zavitkovski.
1968. Role of red alder in western Oregon forest succession. p. 73-74. In Biology of Alder, J. M. Trappe, J. F. Franklin, R. F. Tarrant, and G. M. Hansen (eds.). Northwest Sci. Assoc. Fortieth Ann. Meet., Symp. Proc. 1967.
- New Zealand Forest Service.
1976. Report of Forest Research Institute for 1 January to 31 December 1975. Wellington, New Zealand, p. 34.
- Okkonen, E. A., H. E. Wahlgren, and R. R. Maeglin.
1972. Relationships of specific gravity to tree height in commercially important species. For. Prod. J. 22(7):37-42.

- O'Laughlin, C. L.
1973. A study of tree root strength deterioration following clearfelling. *Can. J. For. Res.* 4(1):107-113.
- Olson, D. S.
1953. Preliminary tests on relative inflammability of logging slash by species in the western white pine type. *For. Wildl. Range Exp. Stn. Res. Note No. 5*, 7 p. Univ. Idaho, Moscow.
- Ovington, J. D.
1956. The composition of tree leaves. *Forestry* 29(1):22-28.
- Owston, Peyton W.
1974. Effects of shortened photoperiod on growth and hardiness of four western conifers. *Third North Amer. For. Biol. Workshop Proc.*, p. 354-355. (Abstract)
- Parker, Johnson.
1951. Moisture retention in leaves of conifers of the northern Rocky Mountains. *Bot. Gaz.* 113:210-216.
- Parker, Johnson.
1954. Available water in stems of some Rocky Mountain conifers. *Bot. Gaz.* 115:380-385.
- Parker, Johnson.
1969. Further studies of drought resistance in woody plants. *Bot. Rev.* 35(4):317-371.
- Pearson, G. A.
1924. Studies in transpiration of coniferous tree seedlings. *Ecology* 5:340-347.
- Pechanec, Anna A., and Jerry F. Franklin.
1968. Comparison of vegetation in adjacent alder, conifer communities. II. Epiphytic, epixylic, and epilithic cryptogams. *In Biology of Alder*, J. M. Trappe, J. F. Franklin, R. F. Tarrant, and G. M. Hansen (eds.). Northwest Sci. Assoc. Fortieth Ann. Meet., Symp. Proc. 1967:85-98.
- Pharis, Richard P.
1966. Comparative drought resistance of five conifers and foliage moisture content as a viability index. *Ecology* 47(2):211-221.
- Pharis, Richard P.
1967. Seasonal fluctuation in the foliage-moisture content of well-watered conifers. *Bot. Gaz.* 128:179-185.
- Pharis, R. P., and William K. Farrell.
1966. Differences in drought resistance between coastal and inland sources of Douglas-fir. *Can. J. Bot.* 44:1651-1659.
- Pharis, R. P., H. Hellmers, and E. Schuurmans.
1967. Kinetics of the daily rate of photosynthesis at low temperatures for two conifers. *Plant Physiol.* 42:525-531.
- Phelps, V. H.
1973. Sitka spruce: a literature review with special reference to British Columbia. *Pac. For. Res. Centre Inform. Rep. BC-X-83*, 39 p. Victoria, Canada.
- Pickford, A. E.
1929. Studies of seed dissemination in British Columbia. *For. Chron.* 5(4):8-16.
- Prochnau, A. E.
1963. Direct seeding experiments with white spruce, alpine fir, Douglas fir, and lodgepole pine in the central interior of British Columbia. *British Columbia For. Serv. Res. Note* 37, 24 p.
- Puritch, George S.
1973. Effect of water stress on photosynthesis, respiration, and transpiration of four *Abies* species. *Can. J. For. Res.* 3:293-298.
- Radwan, M. A.
1969. Chemical composition of the sapwood of four tree species in relation to feeding by the black bear. *For. Sci.* 15(1):11-16.
- Reukema, Donald L.
1965. Seasonal progress of radial growth of Douglas-fir, western redcedar, and red alder. *USDA For. Serv. Res. Pap-26*, 14 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Roff, J. W., and H. W. Eades.
1959. Deterioration of logging residue on the British Columbia coast. *Can. For. Prod. Lab. Tech. Note No. 11*, 38 p.
- Ronco, Frank.
1975. Diagnosis: "sunburned" trees. *J. For.* 73(1):31-35.
- Ross, Charles Robert.
1932. Root development of western conifers. 63 p. M.S. thesis, Univ. Wash., Seattle.

- Roy, Douglass F.
1962. California hardwoods: management practices and problems. J. For. 60(3):184-186.
- Rudinsky, J. A., and J. P. Vité.
1959. Certain ecological and phylogenetic aspects of the pattern of water conduction in conifers. For. Sci. 5(3):259-266.
- Running, Steven William.
1973. Leaf resistance responses in selected conifers interpreted with a model simulating transpiration. 87 p. M.S. Thesis. Oreg. State Univ., Corvallis.
- Running, Steven W.
1976. Environmental control of leaf water conductance in conifers. Can. J. For. Res. 6:104-112.
- Ruth, Robert Harvey.
1968a. Differential effect of solar radiation on seedling establishment under a forest stand. 165 p. Ph.D. Thesis. Oreg. State Univ., Corvallis.
- Ruth, Robert H.
1968b. First-season growth of red alder seedlings under gradients in solar radiation. In Biology of Alder, J. M. Trappe, J. F. Franklin, R. F. Tarrant, and G. M. Hansen (eds.). Northwest Sci. Assoc. Fortieth Annu. Meet., Symp. Proc. 1967:99-105.
- Ruth, Robert H., and Carl M. Berntsen.
1955. A 4-year record of Sitka spruce and western hemlock seed fall. USDA For. Serv. Res. Pap. No. 12, 13 p.
- Ruth, Robert H., and A. S. Harris.
1975. Forest residues in hemlock-spruce forests of the Pacific Northwest and Alaska--a state of knowledge review with recommendations for residue management. USDA For. Serv. Gen. Tech. Rep. PNW-39, 52 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Ruth, Robert H., and Ray A. Yoder.
1953. Reducing wind damage in the forests of the Oregon Coast Range. USDA For. Serv. Pap. No. 7, 30 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Rutter, Mark Russell.
1977. An eco-physiological field study of three Sierra conifers. Ph.D. thesis. Univ. Calif., Berkeley. 116 p.
- Ryan, Kevin C., and Stewart G. Pickford.
1978. Physical properties of woody fuels in the Blue Mountains of Oregon and Washington. USDA For. Serv. Res. Note PNW-315, 10 p. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.
- Ryker, Russell A.
1975. A survey of factors affecting regeneration of Rocky Mountain Douglas-fir. USDA For. Serv. Res. Pap. INT-174, 19 p. Intermountain For. and Range Exp. Stn., Ogden, Utah.
- Sakai, A., and S. Okada.
1971. Freezing resistance of conifers. Silvae Genetica 20(3):91-97.
- Sakai, A., and C. J. Weiser.
1973. Freezing resistance of trees in North America with reference to tree regions. Ecology 54(1):118-126.
- Scheffer, Theodore C., and George G. Hedgcock.
1955. Injury to northwestern trees by sulfur dioxide from smelters. USDA Tech. Bull. 1117, 49 p.
- Schmidt, R. L.
1955. Some aspects of western red cedar regeneration in the coastal forests of British Columbia. British Columbia For. Serv. Res. Note No. 29, 10 p.
- Schmidt, R. L.
1957. The silvics and plant geography of the genus *Abies* in the coastal forests of British Columbia. British Columbia For. Serv. Tech. Publ. No. T-46, 31 p.
- Schmidt, Wyman C.
1969. Seedbed treatments influence seedling development in western larch forests. USDA For. Serv. Res. Note INT-93, 7 p. Intermountain For. and Range Exp. Stn., Ogden, Utah.
- Schubert, G. H.
1954. Viability of various coniferous seeds after cold storage. J. For. 52:446-447.
- Schubert, Gilbert H.
1955. Freezing injury to young sugar pine. J. For. 53:732.

- Schubert, Gilbert H.
1956. Early survival and growth of sugar pine and white fir in clearcut openings. USDA Calif. For. and Range Exp. Stn., For. Res. Note No. 117, 6 p.
- Schubert, Gilbert H., and Ronald S. Adams.
1971. Reforestation practices for conifers in California. Calif. Division For., Sacramento. 359 p.
- Scott, Norman C.
1964. Moisture patterns in Douglas-fir and tanoak slash. USDA For. Serv. Res. Note PSW-55, 5 p. Pac. Southwest For. and Range Exp. Stn., Berkeley, Calif.
- Scott, D. R. M., J. N. Long, and J. Kotar.
Undated. Comparative ecological behavior of western hemlock in the Washington Cascades. p. 26-33. In Western hemlock management conference proceedings, May, 1976. (W. A. Atkinson and R. J. Zasoski, eds.) Univ. of Wash. Coll. of For. Resources Inst. of For. Prod. Contr. No. 34, 317 p.
- Sellars-St. Clare, E.
1968. Silver-spotted tiger moth in British Columbia. Can. For. Insect Dis. Surv. For. Pest Leaflet. No. 5, 6 p.
- Shearer, Raymond C., and David Tackle.
1960. Effects of hydrogen peroxide on germination in three western conifers. USDA For. Serv. Res. Note 80, 4 p. Intermountain For. and Range Exp. Stn., Ogden, Utah.
- Shirley, Hardy L.
1943. Is tolerance the capacity to endure shade? J. For. 41:339-345.
- Show, S. B.
1930. Forest nursery and planting practice in the California pine region. USDA Circ. 92, 75 p.
- Siggins, Howard W.
1933. Distribution and rate of fall of conifer seeds. Journ. Agr. Research 47(2):119-128.
- Smith, J. H. G.
1955. Some factors affecting reproduction of Engelmann spruce and alpine fir. British Columbia For. Serv. Tech. Publ. T-43, 43 p.
- Smith, J. Harry G.
1964. Root spread can be estimated from crown width of Douglas-fir, lodgepole pine, and other British Columbia tree species. For. Chron. 40(4):456-473.
- Smith, J. Harry G.
1970. Weight, size, and persistence of needles of Douglas-fir, western hemlock, western red cedar, and other British Columbia conifers. Univ. British Columbia For. Dep., 27 p. (Mimeographed Rep.)
- Smith, J. Harry G., and M. Bruce Clark.
1960. Growth and survival of Engelmann spruce and alpine fir on seed spots at Bolean Lake, British Columbia, 1954-1959. For. Chron. 36:46-49.
- Smith, J. Harry G., John W. Ker, and Joseph Czizmazia.
1961. Economics of reforestation of Douglas-fir, western hemlock and western red cedar in the Vancouver Forest District. Univ. British Columbia Fac. For. For. Bull. 3, 144 p.
- Smith, J. Harry G., John Walters, and Antal Kozak.
1968. Influences of fertilizers on cone production and growth of young Douglas-fir, western hemlock, and western red cedar on the U.B.C. research forest. Univ. British Columbia Fac. For. For. Bull. 5, 57 p.
- Smith, R. B.
1974. Infection and development of dwarf mistletoes on plantation-grown trees in British Columbia. Can. For. Serv. Rep. BC-X-97, 21 p. Pac. For. Res. Centre, Victoria.
- Soos, J., and J. Walters.
1963. Some factors affecting the mortality of western hemlock and western red cedar germinates and seedlings. Univ. British Columbia Fac. For. Res. Pap. No. 56, 12 p.
- Sorensen, Frank C., and Richard S. Miles.
1974. Differential frost tolerance of ponderosa and lodgepole pine megasporangiate strobili. For. Sci. 20:377-378.
- Spurr, Stephen H.
1964. Forest ecology. The Ronald Press, New York, 352 p.
- Staebler, George R., Paul Lauterbach, and A. W. Moore.
1954. Effect of animal damage on

- a young coniferous plantation in southwest Washington. J. For. 52(10):730-733.
- Stark, N.
1965. Natural regeneration of Sierra Nevada mixed conifers after logging. J. For. 63(6):456-457, 460-461.
- Starker, T. J.
1934. Fire resistance in the forest. J. For. 32:462-467.
- Stein, William I.
1955. Some lessons in artificial regeneration from southwestern Oregon. Northwest Sci. 29:10-22.
- Stein, William Ivo.
1963. Comparative juvenile growth of five western conifers. 194 p., illus. Ph.D. Thesis. Yale Univ., New Haven, Conn.
- Stein, William I.
1965. A field test of Douglas-fir, ponderosa pine, and sugar pine seeds treated with hydrogen peroxide. USDA For. Serv. Tree Planters' Notes 71:25-29.
- Steinbrenner, E. C., and S. P. Gessel.
1956. Windthrow along cutlines in relation to physiography on the McDonald tree farm. Weyerhaeuser Timber Co., For. Res. Note No. 15, 19 p.
- Steinbrenner, E. C., and J. H. Rediske.
1964. Growth of ponderosa pine and Douglas-fir in a controlled environment. Weyerhaeuser Co., For. Pap. 1, 31 p.
- Stephens, F. R.
1965. Ponderosa pine thrives on wet soils in southwestern Oregon. J. For. 63:122-123.
- Stewart, R. E.
1977. Herbicides for weed control in western forest nurseries. Proc., West. Soc. of Weed Sci. Vol 30:78-89.
- Stone, Edward C.
1957. Dew as an ecological factor. I. A review of the literature. II. The effect of artificial dew on the survival of *Pinus ponderosa* and associated species. Ecology 38(3):407-422.
- Stone, Edward C., and Richard B. Vasey.
1968. Preservation of coast redwood on alluvial flats. Science 159(3811):157-161.
- Sudworth, George B.
1908. Forest trees of the Pacific slope. USDA For. Serv., Gov. Printing Office, Wash., D.C. 441 p., illus.
- Sundahl, William E.
1966. Crown and tree weights of madrone, black oak, and tanoak. USDA For. Serv. Res. Note PSW-101, 4 p. Pac. Southwest For. and Range Exp. Stn., Berkeley, Calif.
- Tackle, David.
1959. Silvics of lodgepole pine. USDA For. Serv. Misc. Publ. No. 19, 24 p. Intermountain For. and Range Exp. Stn., Ogden, Utah.
- Tarrant, Robert F.
1953. Soil moisture and the distribution of lodgepole and ponderosa pine: a review of the literature. USDA For. Serv. Pap. No. 8, 10 p. Pac. Northwest For. and Range Exp. Stn., Portland, Ore.
- Tarrant, Robert F., Leo A. Isaac, and Robert F. Chandler, Jr.
1951. Observations on litter fall and foliage nutrient content of some Pacific Northwest tree species. J. For. 49:914-915.
- Tarrant, Robert F., K. C. Lu, W. B. Bollen, and J. F. Franklin.
1969. Nitrogen enrichment of two forest ecosystems by red alder. USDA For. Serv. Res. Pap. PNW-76, 7 p. Pac. Northwest For. and Range Exp. Stn., Portland, Ore.
- Tarrant, Robert F., and Richard E. Miller.
1963. Accumulation of organic matter and soil nitrogen beneath a plantation of red alder and Douglas-fir. Soil Sci. Soc. Amer. Proc. 27:231-234.
- Taylor, Raymond Frank.
1935. Available nitrogen as a factor influencing the occurrence of Sitka spruce and western hemlock seedlings in the forests of southeastern Alaska. Ecology 16:580-602.
- Thilenius, John Frederick.
1964. Synecology of the white-oak (*Quercus garryana*, Doug.) woodlands of the Willamette Valley, Oregon. 167 p. Ph.D. Thesis. Oreg. State Univ., Corvallis.
- Thornburgh, Dale Alden.
1969. Dynamics of the true fir-hemlock forests of the west

- slope of the Washington Cascade Range. 226 p. Ph.D. Thesis. Univ. Wash., Seattle.
- Thrupp, Adrian Cracroft.
1939. Effect of seed-ash characteristics and treatment of seed and soils upon coniferous seed germination. 327 p. Ph.D. Thesis. Univ. Wash., Seattle.
- Timmis, Roger.
Undated. Frost hardiness of western hemlock. p. 118-125. In Western hemlock management conference proceedings. May, 1976 (W. A. Atkinson and R. J. Zasoski, eds.). Univ. Wash. Coll. of For. Resources Inst. of For. Prod. Contr. No. 34, 317 p.
- Tinus, Richard W.
1970. Growing seedlings in controlled environment. West. For. and Conserv. Assoc. West. Reforestation Coord. Comm. Proc. 1970:34-37.
- Trappe, James M.
1977. Selection of fungi for ectomycorrhizal inoculation in nurseries. Ann. Rev. Phytopathol. 15:203-222.
- U.S. Department of Agriculture.
1973. Trees for polluted air. USDA Misc. Publ. No. 1230, 12 p.
- USDA Forest Products Laboratory.
1955. Wood Handbook. U.S. Dep. Agric. Handb. No. 72, 528 p.
- USDA Forest Products Laboratory.
1965. Western wood density survey. Rep. No. 1. USDA For. Serv. Res. Pap. FPL-27, 58 p. For. Prod. Lab., Madison, Wisc.
- Vaartaja, O.
1957. The susceptibility of seedlings of various tree species to *Phytophthora cactorum*. Can. Dep. Agric. Science Serv. For. Biol. Div. Bi-monthly progress report 13(2):2.
- Vaartaja, O.
1959. Evidence of photoperiodic ecotypes in trees. Ecol. Monogr. 29:91-111.
- van den Driessche, R.
1968. A comparison of growth responses of Douglas-fir and Sitka spruce to different nitrogen, phosphorus, and potassium levels in sand culture. Can. J. Bot. 46:531-537.
- van den Driessche, R.
1969. Tissue nutrient concentrations of Douglas-fir and Sitka spruce. British Columbia For. Serv. Res. Note No. 47, 42 p.
- van den Driessche, R.
1971. Response of conifer seedlings to nitrate and ammonium sources of nitrogen. Plant and Soil 34:421-439.
- Van Eerden, E.
1974. Growing season production of western conifers. In North Amer. Containerized For. Tree Seedling Symp. Proc., Great Plains Agric. Counc. Publ. 68:93-103.
- Wagener, Willis W.
1960. A comment on the cold susceptibility of Ponderosa and Jeffrey pine. Madrono 15:217-219.
- Wagener, Willis W., and Ross W. Davidson.
1954. Heart rots in living trees. Bot. Rev. 20(2):61-134.
- Wali, M. K., and V. J. Krajina.
1973. Vegetation-environment relationships of some sub-boreal spruce zone ecosystems in British Columbia. Vegetatio 26(4/6):237-381.
- Wallis, G. W.
1976. Growth characteristics of *Phellinus (Poria) weirii* in soil and on root and other surfaces. Can. J. For. Res. 6(2):229-232.
- Wallis, G. W., and J. H. Ginns, Jr.
1968. Annosus root rot in Douglas-fir and western hemlock in British Columbia. Can. For. Insect Dis. Surv. For. Pest Leaflet. No. 15, 7 p.
- Wallis, G. W., and G. Reynolds.
1967. Poria root rot of Douglas-fir in B.C. Can. For. Insect Dis. Surv. For. Pest Leaflet. No. 3, 8 p.
- Wallis, G. W., J. N. Godfrey, and H. A. Richmond.
1974. Losses in fire killed timber. Can. For. Serv. Inform. Rep. No. BC-X-88, 11 p. Pac. For. Res. Centre, Victoria, British Columbia.
- Walters, J., and P. G. Haddock.
1966. Juvenile height growth of eight coniferous species on five Douglas-fir sites. Univ. British Columbia Fac. For. For. Res. Note No. 75, 17 p.

- Walters, J., and J. Soos.
1963. Shoot growth patterns of some British Columbia conifers. *For. Sci.* 9(1):73-85.
- Walters, J., J. Soos, and J. W. Ker.
1961. Influence of crown class and site quality on growth to breast height of Douglas-fir, western hemlock, and western red cedar. *Univ. British Columbia. Fac. For. Res. Note* No. 37, 4 p.
- Wambolt, Carl L.
1973. Conifer water potential as influenced by stand density and environmental factors. *Can. J. Bot.* 51(12):2333-2337.
- Wang, B. S. P.
1974. Tree-seed storage. *Dep. of the Environment. Can. For. Serv. Publ. No. 1335*, 32 p.
- Waring, R. H.
1969. Forest plants of the eastern Siskiyous: their environmental and vegetational distribution. *Northwest Sci.* 43:1-17.
- Waring, Richard H.
1970. Matching species to site. *In* Regeneration of ponderosa pine. *Symp. Proc.*, 1969. *Oreg. State Univ. Sch. For., Corvallis.* p. 54-61.
- Waring, R. H., W. H. Emmingham, and S. W. Running.
1975. Environmental limits of an endemic spruce, *Picea breweriana*. *Can. J. Bot.* 53(15):1599-1613.
- Waring, R. H., and J. Major.
1964. Some vegetation of the California coastal redwood region in relation to gradients of moisture, nutrients, light, and temperature. *Ecol. Monog.* 34:167-215.
- Webber, Bruce Douglas.
1973. Plant biomass and nutrient distribution in a young *Pseudotsuga menziesii* forest ecosystem. 163 p. *Ph.D. Thesis. Oreg. State Univ., Corvallis.*
- Wellington, W. G.
1969. Effects of three hormonal mimics on mortality, metamorphosis, and reproduction of the western tent caterpillar, *Malacosoma californicum pluviale*. *Can. Entomol.* 101(11):1163-1172.
- Wickman, Boyd E.
1976. Douglas-fir tussock moth egg hatch and larval development in relation to phenology of grand fir and Douglas-fir in northeastern Oregon. *USDA For. Serv. Res. Pap. PNW-206*, 13 p.
- Williams, Carroll B., Jr.
1966. Snow damage to coniferous seedlings and saplings. *USDA For. Serv. Res. Note PNW-40*, 10 p. *Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.*
- Williams, Carroll B., Jr.
1968. Seasonal height growth of upperslope conifers. *USDA For. Serv. Res. Pap. PNW-62*, 7 p. *Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.*
- Wollum, A. G. II, and C. T. Youngberg.
1964. The influence of nitrogen fixation by nonleguminous woody plants on the growth of pine seedlings. *J. For.* 62:316-321.
- Wood, R. F.
1955. Studies of north-west American forests in relation to silviculture in Great Britain. *Bull. For. Comm. No. 25*, 42 p. *London.*
- Worthington, Norman P.
1955. A comparison of conifers planted on the Hemlock Experimental Forest. *USDA For. Serv. Res. Note 111*, 5 p. *Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.*
- Wright, Ernest.
1931. The effect of high temperature on seed germination. *J. For.* 29:679-687.
- Young, R. S.
1974. Composition of ash of arbutus and Douglas-fir. *The Commonwealth For. Rev.* 53(1):49-51.
- Youngberg, C. T., and C. T. Dyrness.
1959. The influence of soils and topography on the occurrence of lodgepole pine in central Oregon. *Northwest Sci.* 33:111-120.
- Ziemer, R. R., and D. N. Swanston.
1977. Root strength changes after logging in southeast Alaska. *USDA For. Serv. Res. Note PNW-306*, 10 p. *Pac. Northwest For. and Range Exp. Stn., Portland, Oreg.*
- Zobel, Donald B., Arthur McKee, Glenn M. Hawk, and C. T. Dyrness.
1976. Relationships of environment to composition, structure, and diversity of forest communities of the central Western Cascades of Oregon. *Ecol. Monog.* 46(2):135-156.

The mission of the PACIFIC NORTHWEST FOREST AND RANGE EXPERIMENT STATION is to provide the knowledge, technology, and alternatives for present and future protection, management, and use of forest, range, and related environments.

Within this overall mission, the Station conducts and stimulates research to facilitate and to accelerate progress toward the following goals:

1. Providing safe and efficient technology for inventory, protection, and use of resources.
2. Developing and evaluating alternative methods and levels of resource management.
3. Achieving optimum sustained resource productivity consistent with maintaining a high quality forest environment.


The area of research encompasses Oregon, Washington, Alaska, and, in some cases, California, Hawaii, the Western States, and the Nation. Results of the research are made available promptly. Project headquarters are at:

Anchorage, Alaska
Fairbanks, Alaska
Juneau, Alaska
Bend, Oregon
Corvallis, Oregon

La Grande, Oregon
Portland, Oregon
Olympia, Washington
Seattle, Washington
Wenatchee, Washington

*Mailing address: Pacific Northwest Forest and Range
Experiment Station
P.O. Box 3141
Portland, Oregon 97208*





The **FOREST SERVICE** of the U.S. Department of Agriculture is dedicated to the principle of multiple use management of the Nation's forest resources for sustained yields of wood, water, forage, wildlife, and recreation. Through forestry research, cooperation with the States and private forest owners, and management of the National Forests and National Grasslands, it strives — as directed by Congress — to provide increasingly greater service to a growing Nation.

The U.S. Department of Agriculture is an Equal Opportunity Employer. Applicants for all Department programs will be given equal consideration without regard to age, race, color, sex, religion, or national origin.