

On the Origin of Fusiform Rust Resistance in Loblolly Pine¹

R.C. Schmidtling², C.D. Nelson², and T.L. Kubisiak²

Abstract:-- Studies of geographic variation in loblolly pine have shown that seed sources from the western (generally west of the Mississippi River) and the northeastern part of the natural distribution are relatively resistant to fusiform rust disease, while those from elsewhere are more susceptible. The greatest problem with rust infection, on the other hand, is in the center of the distribution, exactly where the frequency of resistant genotypes appears to be lowest. One might expect that the frequency of resistant genotypes would be higher, where the disease is more prevalent, due to natural selection. It has been proposed that (1) fusiform rust resistance in loblolly pine in the west originates from hybridization with shortleaf pine. It is well known that shortleaf pine is relatively resistant to fusiform rust, and it is also known that natural hybrids between the two species exist, and they seem to be more common in the west. In the northeast, it has been proposed that (2) hybridization with pond pine is the source of resistance to fusiform rust. Once again, natural hybridization between loblolly and pond pine is known to exist in the northeast, but not much is known about the relative resistance of pond pine to fusiform rust. Allozyme and cortical monoterpene data were used to evaluate these hypotheses and the results suggest that hybridization is not the primary source of fusiform rust resistance either in the west or northeast.

Keywords: *Pinus taeda*, *Cronartium quercuum* f. sp. *fusiforme*, fusiform rust, disease resistance, evolution, hybridization, *Pinus echinata*, *Pinus serotina*.

INTRODUCTION

Fusiform rust (caused by *Cronartium quercuum* (Berk.) Miyabe ex Shirai f. sp. *fusiforme*) is the most damaging disease of southern pines in the southeastern United States (Powers and others 1981) and it causes substantial mortality in severe epidemics. In lesser, more common epidemics the disease causes relatively less mortality but moderate to severe degrade in the quality and strength of the main stem and branches. In loblolly pine (*Pinus taeda* L.), considerable geographic variation in resistance to fusiform rust disease has been found, with resistance decreasing from west to east, except in the northeastern part of the range where trees tend to be nearly as resistant as western seed sources. For the current study we reviewed the available research to evaluate the hypothesis that resistance in western and northeastern sources of loblolly pine originated in related, sympatric species known to be relatively resistant to fusiform rust.

¹ Paper originally presented at the 27th Southern Forest Tree Improvement Conference, June 24-27, 2003, Stillwater, OK and inadvertently omitted from the Proceedings. It is published here (28th Southern Forest Tree Improvement Conference, June 20-23, 2005, Raleigh, NC) as presented in 2003 (08/29/05).

² USDA Forest Service, Southern Institute of Forest Genetics, USDA Forest Service, Saucier, MS 39574.

DISCUSSION

Geographic Variation

In his pioneering study of geographic variation in loblolly pine, Wakeley (1944) found that a Georgia seed source was much more heavily infected with fusiform rust than western sources from Texas, Arkansas and Louisiana. This study was the first evidence for the now well-known superiority in growth and disease resistance of seed sources from Livingston Parish, Louisiana. The results of the Southwide Southern Pine Seed Source Study (SSPSSS) (Wells and Wakeley 1966) confirmed the relative resistance of the western sources to fusiform rust, and also found a great deal of resistance in a seed source from the extreme northeast of the loblolly pine range (i.e., Maryland). The results of these studies and a study by Grigsby (1973) resulted in large-scale planting of Livingston Parish loblolly pine in areas of high rust hazard in Mississippi, Alabama, Georgia, and Florida (Wells 1985).

Significant seed source-by-planting site interaction in fusiform rust resistance was also observed in the SSPSSS, although the resistant sources from the western part of the range plus the Maryland source are clearly separated from the susceptible sources regardless of the infection level (Figure 1). The eastern and western populations of loblolly pine have been considered to be distinctly different, and the isolating effect of the pine-free Mississippi River basin has been proposed as the mechanism that perpetuates these differences (Wells and Wakeley 1966). This is especially true with regard to resistance to fusiform rust. Sources from west of the river are considered resistant, while those from east of the river are considered susceptible. The one important exception has been Livingston Parish loblolly pine, which is from east of the river but is relatively resistant to fusiform rust.

Studies that have sampled the range of loblolly pine more intensively have shown that the variation in fusiform rust resistance is continuous from west to east. In a range-wide study planted in Arkansas, Grigsby (1973) found no distinct separation between western and eastern sources in rust resistance, rather there was a continuous decrease from west to east, reaching a minimum at the longitude of eastern Georgia, then increasing to moderate levels in the Maryland (MD.) and Delaware (DEL.) sources (Figure 2).

Several regional studies with intensive geographic sampling verified the continuous variation model of Figure 2. Across southern Louisiana, Crow (1958) and Dyer and others (1977) found that resistance to fusiform rust disease decreased from west to east in a continuous manner, with no apparent discontinuity at the Mississippi River. Wells and others (1991) also found that variation in rust resistance across the Mississippi River was continuous, and that there was no distinct separation between western and eastern sources in rust resistance. In a study in Georgia, Sluder (1980) showed a decrease in resistance from west to east across Georgia, reaching a probable low point at the longitude of eastern Georgia. The performance of Livingston Parish is not anomalous when the variation is assumed to be continuous.




Figure 1. Fusiform rust infection (% galled trees) in eight seed sources of loblolly pine plotted against average rust infection for each planting (adapted from Wells and Wakeley 1966).

Several other studies verify the pattern of fusiform rust resistance of Figure 2. Wells (1966) compared infection of nine Texas seed sources with a source from Livingston Parish, Louisiana and central Georgia. The Texas sources averaged 12% infected, the Livingston Parish source 10% and the Georgia source 32%. Pait and Draper (1983) included sources from east Texas, Maryland and Livingston Parish as well as Florida sources in several plantings in Florida and south Georgia. In all plantings, the Maryland and east Texas sources suffered less infection than the Florida sources. The performance of the Livingston Parish source was comparable to that of the Texas and Maryland sources in all plantings except for one. Other studies further verify the model of Figure 2, for example, Cole (1973) found only minor differences among sources from Georgia and South Carolina, where only small differences might be expected.

The geographic pattern for fusiform rust resistance is difficult to explain, since the lowest concentration of resistant loblolly genotypes occurs exactly where rust infection levels have been the highest, that is, in central Georgia and adjoining Alabama and South Carolina (Squillace 1976). Since fusiform rust infection often causes reduced growth and mortality, natural selection should favor the more resistant genotypes in areas of high infections (Kareiva 1999).




Figure 2. Fusiform rust infection in loblolly pine seed sources from across the natural range, when planted in south Arkansas, plotted against longitude of the seed source (adapted from Grigsby 1973). Also shown is the average infection of SSPSSS seed sources over 10 plantings (adapted from Wells and Wakeley 1966).

One could argue that infection levels are lower in the west and northeast, because that is where the more resistant genotypes are found. That would logically require greater selection for resistance in these areas now or at some time in the past. Warm temperatures and very high humidity at the time of infection of the pine host is a requirement for successful infection (Snow and Froelich 1968). Currently, however, the climate in the central part of the range is more favorable for infection than in the colder northern portions (Arkansas and Maryland) and drier western portions (Texas and Arkansas) of the loblolly pine range.

One popular explanation for the geographic pattern of fusiform rust resistance in loblolly pine involves hybridization with shortleaf pine (*Pinus echinata* Mill.) in the west and pond pine (*Pinus serotina* Michx.) in the northeast (Wells and Wakeley 1966).

Resistance of Western Sources

The resistance of shortleaf pine to fusiform rust disease is well known. The shortleaf-loblolly hybrid is also quite resistant (Henry and Bercaw 1956) and ample evidence for natural hybridization between loblolly and shortleaf pines exists (Mergen and others 1965). Florence and Hicks (1980) compared putative natural hybrids with loblolly and shortleaf pines and found that

frequencies of the "loblolly" IDH allele in shortleaf pine, especially in the western part of the range (Edwards and Hamrick 1995; Raja and others 1997). The frequency of the loblolly allele was very high in a seed orchard population in southern Arkansas (Schmidtling and Hipkins 2001). Out of 22 clones, four (or 18%) showed evidence of hybridization with loblolly pine using the IDH locus criterion. It is interesting that high frequencies of loblolly IDH alleles occur in populations well north of the current natural range of loblolly pine (Figure 4). This suggests that there may be considerable loblolly pine pollen flow northward in the western part of the natural range.




Figure 4. A map of the natural distribution of shortleaf pine showing the frequency of the "loblolly" IDH allele (adapted from Raja and others 1997 and Edwards and Hamrick 1995).

For resistance to have reached such high levels in western populations of loblolly pine, considerable hybridization would be required, along with selection for the trait. This would presumably result in the inclusion of other traits from shortleaf pine in the western loblolly pine populations. In a morphological study of shortleaf and loblolly pine provenance tests, however,

Wells and others (1977) found that resistant western sources of loblolly pine were no more similar to shortleaf pine than were susceptible eastern sources.

Resistance of Eastern Sources

In the northeastern portion of the loblolly pine range, pond pine is commonly associated with loblolly pine, as is shortleaf pine. There is also some evidence for introgression of pond pine genes into loblolly pine (Kang 1967; Saylor and Kang 1973). Some disagreement is found in the literature regarding the relative fusiform rust resistance of pond pine. In one manual, pond pine is listed as "moderately susceptible" compared to loblolly and slash pines (*Pinus elliottii* Engelm. var. *elliottii*), which are listed as "very susceptible" (USDA Forest Service 1972). In artificial inoculation tests, Powers (1972) found that pond pine was very rust resistant, whereas Hedgcock and Siggers (1949) found that it was nearly as susceptible as loblolly pine.

Data on cortical monoterpenes do not show any obvious relationship between pond pine and loblolly pine (Table 1). Pond pine has very high limonene content, 47.9%, whereas loblolly pine averages only 10.5%. Further, loblolly pine from the northeast has much lower limonene content than the range-wide average (Squillace and Wells 1981). If significant introgression of pond pine genes into loblolly pine was occurring in the northeast, one might expect to find a higher limonene content in northeastern loblolly pine sources.

Table 1. Concentration of major cortical monoterpenes in buds of several southern pine species^a.

Species	Alpha-Pinene	Beta-Pinene	Myrcene	Beta-Phellandrine	Limonene
	----- % -----				
Shortleaf pine	33.2	11.1	7.8	38.0	3.2
Spruce pine	29.9	4.1	11.4	43.6	5.4
Slash pine	56.0	27.0	1.3	0.5	13.0
Loblolly pine	31.1	17.0	19.7	20.4	10.5
Pond pine	10.4	35.7	1.2	1.3	47.9

^a Unpublished data on file at the USDA Forest Service, Southern Institute of Forest Genetics, Saucier, MS.

Some researchers believe that the form of fusiform rust resistance in the northeastern sources differs from that found in the western sources (Squillace and Wells 1981). An independent origin for the resistance of western sources compared to northeastern sources has been proposed based on evolutionary pressures during the Pleistocene (Wells and others 1991). Some evidence for this has been found in artificial inoculation data (Schmidtling unpublished data).

Geographic variation in fusiform rust infection from artificial inoculation tests of seedlings show a different trend than variation in field studies. It is generally acknowledged that results of artificial inoculation tests do not always conform to the results of field trials in loblolly pine (Anderson and Powers 1985). Schmidtling (unpublished data) found that the northeastern

sources tended to be resistant, as they are in field tests, but that the western sources were not, the opposite of their performance in field tests (Figure 5). This leads us to hypothesize that in field tests, several kinds of resistance, for example, physiological, morphological and phenological are expressed, but in artificial inoculation of post-cotyledon-stage seedlings possibly only one kind is expressed. Eastern sources have a relatively higher frequency of this kind of resistance than do the western sources. Thus, the resistance of the two widely separated populations may have different origins.




Figure 5. Infection of loblolly pine seedlings from across the natural range when inoculated with fusiform rust basidiospores. Data are from artificial inoculation trials of seedlings from seed orchard trees (Schmidtling and others 1999). The dotted line is a linear polynomial equation ($Y = X_1 - X_2^2$, where Y = % galled trees, X_1 = longitude and X_2 = latitude) fit to the data shown in this figure.

CONCLUSIONS

Recent introgression of genes for fusiform rust resistance into loblolly pine from shortleaf pine in the west or pond pine in the east seems insufficient to account for the relative resistance of these seed sources. Intensive selection pressure due to optimal conditions for disease development at some time in the distant past is a more likely explanation, although hybridization during this time may have provided the genetic variation necessary for the evolution of the resistance observed today.

Schmidtling (in Wells and others 1991) proposed that loblolly pine retreated southward into two populations during the Wisconsin glaciation—Florida and south Texas or Mexico. The climate at that time was much more humid and favorable for fusiform rust infection in the western population (Texas and Mexico) and in the northern part of the eastern population (Florida) (Webb and others 1987). After the subsequent advance northward at the end of the Wisconsin, the populations merged east of the Mississippi River, creating the basis of the present pattern of geographic variation.

The lack of significant recent hybridization as a factor in resistance to fusiform rust disease may make the study of gene-for-gene interaction in host-pathogen analysis more straightforward. The possible west-east difference in resistance mechanisms also should be explored in detailed host-pathogen genetic studies and possibly exploited in breeding for rust resistance. More detailed molecular analysis of the genomes of the involved host species and the actual resistance genes will be required to completely answer this question.

LITERATURE CITED

Anderson, R.L. and H.R. Powers. 1985. The resistance screening center. In. Proc. IUFRO Rust of Hard Pines Working Party Conf., Athens, GA, pp. 59-66.

Cole, D.E. 1973. Comparisons within and between populations of planted slash and loblolly pine: A seed source study. In: Proc. 12th South. Forest Tree Improv. Conf., Baton Rouge, LA, pp. 277-292.

Crow, A.B. 1958. Fourth-year results from a local geographic seed-source test in planted loblolly pine. Louisiana Ag. Exp. Sta., LSU Forestry Notes No. 21, 2 p.

Dyer, J.M., A.B. Crow and P.W. Hannah. 1977. Performance of loblolly pine from five within-state seed sources on three locations in central Louisiana. Louisiana Ag. Exp. Sta., LSU Forestry Notes No. 120, 3 p.

Edwards, M.A. and J.L. Hamrick. 1995. Genetic variation in shortleaf pine, *Pinus echinata* Mill. (Pinaceae). Forest Genetics 2:21-28.

Florence, L.Z. and R.R. Hicks, Jr. 1980. Further evidence for introgression of *Pinus taeda* with *P. echinata*: Electrophoretic variability and variation in resistance to *Cronartium fusiforme*. Silvae Genetica 29:41-43.

Grigsby, H.C. 1973. South Carolina best of 36 loblolly pine seed sources for southern Arkansas. USDA-Forest Service, Gen. Tech. Rep. SO-89, 10 p.

Hedgcock, G.G. and P.W. Siggers. 1949. A comparison of the pine-oak rusts. USDA Tech. Bull. No. 978, 29 p.

Henry, B.W. and T.E. Bercaw. 1956. Shortleaf-loblolly hybrid pines free of fusiform rust after 5 years' exposure. Journal of Forestry 54:779.

Huneycutt, M. and G.R. Askew. 1989. Electrophoretic identification of loblolly pine-shortleaf pine hybrids. *Silvae Genetica* 38:95-96.

Kang, K.W. 1967. Relationship between loblolly (*Pinus taeda* L.) and pond (*Pinus serotina* Michx.) pines in North Carolina. PhD. Dissertation: North Carolina State University, Raleigh, NC, 145 p. (Diss. Abs. No. 67-7419).

Kareiva, P. 1999. Coevolutionary arms race: Is victory possible? *Proceedings of the National Academy of Sciences (USA)* 96:8-10.

Mergen, F., G.R. Stairs and E.B. Snyder. 1965. Natural and controlled loblolly x shortleaf pine hybrids in Mississippi. *Forest Science* 11:306-323.

Pait, J.A., III, L. Draper, Jr. and R.A. Schmidt. 1985. Third-year comparisons of loblolly and slash pine seed sources for fusiform rust resistance and growth potential in north central Florida. In: *Proc. 18th South. Forest Tree Improv. Conf., Long Beach, MS*, pp. 314-320.

Powers, H.R. Jr. 1972. Testing for pathogenic variability within *Cronartium fusiforme* and *C. quercuum*. In: *Proc. NATO-IUFRO Advanced Study Institute, USDA Misc. Pub. No. 1221*, pp. 505-509.

Powers, H.R. Jr., R.A. Schmidt and G.A. Snow. 1981. Current status and management of fusiform rust on southern pines. *Annual Review of Phytopathology* 69:720-722.

Raja, R.G., C.G. Tauer, R.F. Wittwer and Y. Huang. 1997. Isozyme variation and genetic structure in natural populations of shortleaf pine (*Pinus echinata*). *Canadian Journal of Forestry Research* 27:740-749.

Saylor, L.C. and K.W. Kang. 1973. A study of sympatric populations of *Pinus taeda* L. and *Pinus serotina* Michx. in North Carolina. *Journal of the Elisha Mitchell Science Society* 89:101-110.

Schmidtling, R.C., E. Carroll and T. LaFarge. 1999. Allozyme diversity of selected and natural loblolly pine populations. *Silvae Genetica* 48:35-45.

Schmidtling, R.C. and V. Hipkins. 2001. Factors contributing to genetic variation in ice damage susceptibility in shortleaf pine. In: *Proc. 26th South. For. Tree Improv. Conf., Athens GA*, pp. 60-62.

Sluder, E.R. 1980. A study of geographic variation in loblolly pine in Georgia-- 20th-year results. USDA-Forest Service, Southeastern Forest Experiment Station, Research Paper SE-213, 26 p.

Snow, G.A. and R.C. Froelich. 1968. Daily and seasonal dispersal of basidiospores of *Cronartium fusiforme*. *Phytopathology* 58:1532-1536.

Squillace, A.E. 1976. Geographic patterns of fusiform rust infection in loblolly and slash pine plantations. USDA-Forest Service, Research Note SE-232, 4 p.

Squillace, A.E. and O.O. Wells. 1981. Geographic variation of monoterpenes in cortical oleoresin of loblolly pine. *Silvae Genetica* 30:127-135.

USDA Forest Service. 1972. Insects and diseases of trees in the south. USDA Forest Service, Southeastern Area State & Private Forestry, Atlanta, GA, 81 p.

Wakeley, P.C. 1944. Geographic sources of loblolly pine seed. *Journal of Forestry* 42:23-32.

Webb, T., III, P. J. Bartlein and J.E. Kutzbach. 1987. Climatic changes in eastern North America during the past 18,000 years: Comparisons of pollen data with model results. *The Geology of North America, Vol. K-3, North America and adjacent oceans during the last deglaciation*, The Geological Society of America, pp. 447-462.

Wells, O.O. 1966. Variation in rust resistance among pine seed sources and species in a Mississippi planting. *Forest Science* 12:461-463.

Wells, O.O. 1985. Use of Livingston Parish, Louisiana loblolly pine by forest products industries in the Southeast. *Southern Journal of Applied Forestry* 9:180-185.


Wells, O.O., W.L. Nance and B.A. Thielges. 1977. Variation in needle traits in provenance tests of *Pinus taeda* and *P. echinata*. *Silvae Genetica* 26:125-130.

Wells, O.O., G.L. Switzer and R.C. Schmidtling. 1991. Geographic variation in Mississippi loblolly pine and sweetgum. *Silvae Genetica* 40:105-119.

Wells, O.O. and P.C. Wakeley. 1966. Geographic variation in survival, growth, and fusiform rust infection of planted loblolly pine. *Forest Science Monograph No. 11*, 40 p.

Proceedings

28th Southern Forest Tree Improvement Conference



Sponsored by

The Southern Forest Tree Improvement Committee
&
Department of Forestry and Environmental Resources
North Carolina State University
Raleigh, NC

June 21-23, 2005

Proceeding of the 28th Southern Forest Tree Improvement Conference

Edited by: Steven E. McKeand
Bailian Li
Department of Forestry and Environmental Resources
N.C. State University
Raleigh, NC 27695-8002

The papers and abstracts in these proceeding were submitted by the authors as electronic files. Modifications were made in format to provide for consistency and to allow best possible placement of figures and tables. Responsibility for technical content remains with the respective authors.

Copies of this publication may be obtained from:

The National Technical Information Service
Springfield, Virginia 22161
Phone: 1-800-553-6847
Fax: 1-703-605-6900
<http://www.ntis.gov>

Electronic copies may also be obtained from the NC State University Forestry Extension and Outreach Program (FEOP) web site:

www.ncsu.edu/feop/sftr/proceedings.html

28th Southern Forest Tree Improvement Conference

Plenary session: Celebrating our History / Building a New Legacy

- Moderator: Bailian Li

	<u>Page</u>
Our Roots: The Start of Tree Improvement in the South Bruce Zobel	1
Impacts of Tree Improvement on the Forest Products Industry R.C. Kellison	6
Tree Improvement and Forest Management in the Evolving Landscape of Forest Land Ownership Al Lyons	13
Issues Facing State Tree Improvement Programs Russ Pohl	14
Improving Forest Productivity Through Biotechnology Mark Rutter	15
Clonal Forestry: Out of the Lab, Finally John Pait	16

Genomics & Biotechnology – Moderator: Andy Benowicz

Genomic Resources for the Study of Loblolly Pine and Other Conifers J.F.D. Dean	17
The Cellulose Synthase and Cellulose Synthase-Like Gene Superfamily in the Populus Genome L. Li, S. Suzuki, Y.-H. Sun, and V. Chiang	18
Candidate Gene-Quantitative Phenotype Associations for Resistance to Fusiform Rust and Pitch Canker in Loblolly Pine E.S. Ersoz, et al.	19
Microsatellite DNA Variation within the University/Industry Tree Improvement Cooperatives' Loblolly Pine Founder Population C.D. Nelson, C.S. Echt, and F.E. Bridgwater	20
Identification of a New Retrotransposable Element in Loblolly Pine M.N. Islam-Faridi, et al.	21
Molecular Genetics of Cellulose Synthesis in Developing Wood of Loblolly Pine C. J. Nairn, A. Wood-Jones, W. Lorenz, and J.F. Dean	22

Breeding, Testing, & Selection – Moderator: Richard Bryant

	<u>Page</u>
Forty Years of Genetic Improvement of Shortleaf Pine in Missouri D. P. Gwaze, R. Melick, C. Studyvin, and M. Coggeshall.	23
Family Composition Changes Over Time in a 17-Year-Old Mixed-Family Loblolly Pine Stand J.P. Adams and S.B. Land	38
Heritability and Gain for Early Height Growth and Foliage Retention in Eastern Cottonwood from the Southeast J. P. Jeffreys and S.B. Land	48
Interacting Genes in the Pine-Fusiform Rust Forest Pathosystem H.V. Amerson et al.	60
Genetic Variation in Wood Quality (MOE) of Coastal Douglas-fir R. Johnson and B. Gartner	61
Genetically Improved Eucalypts for Novel Applications and Sites in Florida D.L. Rockwood, G.F. Peter, M.H. Langholtz, B. Becker, A. Clark, III, and J. Bryan	64

Seed & Propagule Production, Clonal Testing – Moderator: Nick Muir

Survival and Promotion of Female and Male Strobili from Topgrafting in Third-Cycle Slash Pine (<i>Pinus elliottii</i> var. <i>elliottii</i>) Breeding Program A.M. Medina, D.A. Huber, T.L. White, and T.A. Martin	76
The Effect of Root Segment Origin, Size, and Orientation in Aspen Rootling Propagation J.S. Brouard, f. Niemi, and L.R. Charleson	80
The Impact of Variable Success of Somatic Embryogenesis Among Elite Crosses on Expected Genetic Gain and Diversity of Selected Varieties T.J. Mullin, M. Lstiburek, J. Pait, and Y.A. El-Kassaby	92
Genetic Variation in MFA, MOE and Wood Density Among Clones of <i>Pinus taeda</i> L. F. Isik, B. Li, B. Goldfarb	95
Genetic Analysis of Early Field Growth of Loblolly Pine Clones and Seedlings from the Same Full-Fib Families B. Baltunis, D. Huber, T. White	96
Accounting for Spatial Variability in Clonal Forestry Trials S.A. Gezan, T.L. White, D.A. Huber	99
Clonal Replacement as a Tool for Seed Orchard Managers C.L. Rosier, S.E. McKeand, E.M. Raley	102

Breeding, Testing, & Selection – Moderator: Greg Powell

	<u>Page</u>
Susceptibility of Loblolly x Slash Pine Interspecific F1 Hybrids to Tip Moth Infestation and Fusiform Rust Infection in a South Mississippi Planting M.T. Highsmith, L.H. Lott, and C.D. Nelson	103
Comparing Parameter Estimation Techniques for Diameter Distributions of Loblolly Pine B.C. Smith, B.P. Bullock, and S.E. McKeand	104
Preliminary Results for Above- and Below-Ground Bio-Sequestration of a Mature F1 Black Spruce Varying in Site and Family Productivity J.E. Major, et al.	107
Expected Genetic Gains and Development Plans for Two Longleaf Pine Third-Generation Seedling Seed Orchards C.D. Nelson, L.H. Lott, and D.P. Gwaze	108
Genetic Variation in Young Fraser Fir Progeny Tests J.L. Emerson, L.J. Frampton, and S.E. McKeand	115
Using Biotechnology to Help Restore the American Chestnut S.A. Merkle and G.M. Andrade	118

Breeding, Testing, & Selection – Moderator: Fred Raley

Total Inside-Bark Volume Estimation for Loblolly Pine (<i>Pinus taeda</i> L.) in Genetic Trials J.R. Sherrill, et al.	123
Selfing Results in Inbreeding Depression of Growth but not of Gas Exchange K. Johnsen, J.E. Major, C. Maier, and D. Barsi	126
Is the <i>cad-n1</i> Allele Associated with Increased Wood Density or Growth in Full-Sib Families of Loblolly Pine? Q. Yu, B. Li, C.D. Nelson, S.E. McKeand, and T.J. Mullin	127
Height and Diameter Growth Differences Among Eight Cherrybark Oak Provenances J.P. Adams, R.J. Rousseau, and J.C. Adams	128
Genetic Variation in Basal Area Increment Phenology and its Correlation with Growth Rate in Loblolly and Slash Pine Families and Clones V. I. Emhart, T.A. Martin, T.L. White, and D.A. Huber	131
On the Origin of Fusiform Rust Resistance in Loblolly Pine R.C. Schmidtling, C.D. Nelson, and T.L. Kubisiak	135

Conservation & Genetic Diversity – Moderator: Gary Hodge

	<u>Page</u>
Seedling Resistance to <i>Phytophthora cinnamomi</i> in the Genus <i>Abies</i> J. Frampton, D.M. Benson, J. Li, A.M. Braham, E.E. Hudson, and K.M. Potter	146
An <i>Ex Situ</i> Gene Conservation Plan for Fraser Fir K.M. Potter and J. Frampton	148
Summary of Important Results from Biological Research Conducted by Camcore Over the Last 25 Years W. S. Dvorak	160
Geographic Variation in Shortleaf Pine - Cortical Monoterpenes R.C. Schmidtling, J.H. Myszewski, and C.E. McDaniel	161
From Tree Improvement to Species Improvement: Restoration and Conservation Efforts on the Forest Service's Southern National Forests B.S. Crane	168

Plenary Session – Changing Wood Products – Moderator: J.B. Jett

A Random Walk Through the History of Breeding for Wood Quality J.P. van Buijtenen T.D. Byram	169
The Outlook for Pine Plantation Management in the South R.C. Abt	170
After 50 Years, Shift Genetic Emphasis Toward the True Tree Value – Pine Sawtimber H.M. Lupold	171
Commercialization of Forest Biotechnology: Economic Targets for Enhanced Global Competitiveness of the U. S. Pulp and Paper Industry G.F. Peter, D.E. White, N. Sicarelli, R. De La Torre, D. Newman	172
What Do We Do to Improve Wood Properties in a Breeding Program? B. Li, F. Isik, and B. Goldfarb	184

Poster Abstracts and Extended Abstracts

	<u>Page</u>
USDA Forest Service Cooperates with the USDA National Center for Genetic Resources Preservation on a Nationwide <i>ex situ</i> Plant Genetic Resources Conservation Plan	
J. Barbour	186
Genetics of Resistance to <i>Phytophthora cinnamomi</i> in Chestnut	
M.E. Bowles and J. Frampton	187
USDA Forest Service, Forest Health Protection, Resistance Screening Center Services	
Carol H. Young	188
Genetic Linkage Map of Nordmann fir: The First in the Genus <i>Abies</i>	
E.E. Hudson, L.J. Frampton, H.V. Amerson, S.A. Garcia, D.M. Benson	189
Towards an Information Concerning Genetic Parameters of European Larch (<i>Larix decidua</i> Mill.) on the Base of Progenies from Diallel Crossing Evaluation at the age of 31 & 34 Years	
J. Frydl, J. Sindelar ¹ , P. Novotny	190
The Role of NAC068, a NAC Domain Protein, in Wood Development	
K. Hunt and S. Covert	191
Genetic Gain and Diversity of Seed Crops under Alternative Management Options in a Clonal Seed Orchard of <i>Pinus thunbergii</i>	
K.S. Kang, D. Lindgren, T.J. Mullin, W.-Y. Choi, S.-U. Han and C.-S. Kim	192
Application of Sprinkler System for Control of Cone Insects in a Conal Seed Orchard of <i>Pinus koraiensis</i>	
K.-Y. Lee, K.-J. Cho, S.-B. Chung and J.-T. Kang	195
Cloning of Mature Black Willow Trees via Shoot Organogenesis	
S. Lyyra, A. Lima and S.A. Merkle	198
Light Quality Treatments Improve Pine Somatic Seedling Production Efficiency	
S.A. Merkle, P.M. Montello, X. Xia, and D.R. Smith	199
A Microsatellite Assessment of Population Architecture and Gene Flow in Fraser Fir	
K. M. Potter, S. Josserand, J. Frampton, and C.D. Nelson	200
Impact of Crop Tree Release on Wood Properties of Pitch x Loblolly Pine Hybrids	
M. L. Jackson, T. R. Fox	201
Expression Patterns of Pinus Defense Genes During Host-Pathogen Interactions	
A.M. Morse, et al.	202
Invertases as Genetic Determinants of Sink Strength	
P.N. Boccock, L.F. Huang, K.E. Koch, J.M. Davis	203
List of Participants	204