

The Use of Plants as Bioindicators of Ozone¹

William J. Manning²

Abstract

*A variety of vascular plant species exhibit typical foliar injury symptoms when exposed to ambient ozone, making them useful as bioindicators of relative air quality for a particular location or region. They are quite useful in areas where mechanical ozone monitors are not available. Bioindicators are often introduced plant species known as sentinels. They are known to be sensitive to ozone and will respond rapidly if they are given special care to ensure ozone uptake and injury. Sentinels are usually genetically-uniform, rapid growing herbaceous annuals. Their rapid and well-characterized response to ozone exposure has made them quite useful. Bel-W3 tobacco (*Nicotiana tabacum* L.) is a sentinel plant bioindicator for ozone that is used all over the world. Detector bioindicators are plant species that are found growing naturally in an area and known to be sensitive to ozone only when conditions are appropriate for ozone uptake and plant injury. Detectors are often slow-growing, determinate perennial plants, shrubs, or trees that respond slowly to ozone, with symptoms occurring quite late in the growing season. Populations of detectors are not genetically uniform and only part of a population may show ozone injury symptoms. Black cherry (*Prunus serotina* L.) is a common detector bioindicator for ozone in North America. A comparison of surveys of sentinel and detector bioindicators in the same area often show different results. From an ecological perspective, visible injury on a detector bioindicator is more significant than visible injury on a sentinel bioindicator. When using plants as bioindicators, careful consideration needs to be given to the nature, requirements, and utility of sentinels and detectors in relation to the relevance and utility of the results obtained.*

Introduction

For more than 50 years, certain plant species have been used as bioindicators because they are sensitive to ozone under ambient conditions (Middleton and others 1950, Noble and Wright 1958). Sensitive individuals exhibit typical foliar injury symptoms when exposed to ambient ozone under conditions appropriate for ozone uptake. These symptoms are considered to be diagnostic or typical as they have been verified in exposure/response studies under experimental conditions (Krupa and Manning 1988). These plants are considered to be reliable biological indicators or bioindicators for ambient ozone. The subjective determination of the intensity or extent of foliar injury of bioindicators is used as an index of relative air quality for ozone for a particular location or region. Ozone has become an air pollution problem in most industrialized nations, resulting in an increased interest in using bioindicator plants on a world-wide basis. According to Guderian and others (1985), injury on Bel-W3 tobacco (*Nicotiana tabacum* L.) is usually the first indication a county or region has developed an ozone problem. The history and use of plants as ozone bioindicators has been extensively reviewed elsewhere (Arndt and others 1987, Burton 1986, deBauer 1972, Feder and Manning 1979, Guderian and others 1985, Heck 1966, Hernandez and deBauer 1989, Manning 1991, Manning and Feder 1980, Posthumus 1982, Stuebing and Jager 1982, Tonneijck and Posthumus 1987, Weintstein and Laurence 1989).

This paper discusses the use of introduced species or sentinels and naturally occurring plants or detectors as bioindicators of ozone exposure.

Sentinels and Detectors

From an ecological perspective, Spellerberg (1991) recognizes two types of plant bioindicators useful in studies designed to detect gaseous air pollutants (*table 1*). Sentinels are non-indigenous plant species, consisting of well-defined selections,

¹ An abbreviated version of this paper was presented at the International Symposium on Air Pollution and Climate Change Effects on Forest Ecosystems, February 5-9, 1996, Riverside, California.

² Phytopathologist, Department of Microbiology, University of Massachusetts, Amherst, MA 01003-5730.

cultivars or clones, that exhibit diagnostic reliable foliar symptoms when exposed to ambient ozone. Sentinels are grown in charcoal-filtered air and then introduced into areas, usually for short time periods. They are quite useful in identifying areas where ozone concentrations and exposures are suspected to be high enough and of a long enough duration to cause foliar injury (Ashmore and others 1980, Bytnerowicz and others 1993, deBauer 1972, Heck and Heagle 1970, Hernandez and deBauer 1989, Jacobson and Feder 1974, Kelleher and Feder 1978, Lipa and Votapkova 1995, Noble and Wright 1958, Oshima 1974, Posthumus 1982). Examples of sentinel bioindicators for ozone include Bel-W3 tobacco (Heck and others 1964, Heck and Heagle 1970, Heggstad 1991) and morning glory (*Ipomoea purpurea*) (Manning 1977, Nouchi and Aoki 1979). Detector plants are indigenous to an area that may exhibit typical foliar injury symptoms to ozone exposure *in situ*. They are useful in assessing the long-term or cumulative effects of ozone. Examples include the sensitive individuals in populations of black cherry (*Prunus serotina*) (Davis and others 1982), Ponderosa (*Pinus ponderosa*) and Jeffrey (*P. Jeffreyi*) pines (Stolte and others 1992), and milkweed (*Asclepias syriaca*) (Duchelle and Skelly 1981).

Table 1 — Types of indicator plant species used to assess relative air quality for ozone (Spellerberg 1991).

Sentinels	Well-defined plants known to be sensitive to ozone are introduced into an area to serve as early warning devices or checks on the efficiency of abatement practices. Examples—Bel-W3 tobacco (<i>Nicotiana tabacum</i>), Morning glory (<i>Ipomoea purpurea</i>)
Detectors	Plants that naturally occur in the area of interest that may exhibit typical foliar responses to ozone. Examples—Black cherry (<i>Prunus serotina</i>), Milkweed (<i>Asclepias syriaca</i>)

Sentinels are the most commonly used bioindicators of ambient ozone. They are usually well-defined, genetically-uniform, herbaceous annuals that grow rapidly. Their response time to ozone is rapid, serving as an early warning of ozone presence. Rapid response, however, may only occur in the early stages of their growth cycles, necessitating frequent re-introduction of new plants. They must be grown in charcoal-filtered air until they are old enough to move to the field. To minimize edaphic factors, they are usually grown in pots of a uniform artificial growing medium. They require water, fertilizer, shading, and often pesticide applications, on a regular basis. Protection from animals and vandals may also be required. If sentinels do not receive the special care they require, they will not respond well to ozone in a typical or relevant manner.

Detectors are native plants selected *in situ* and usually are not given any special cultural care. Usually they are determinate perennial plants, trees, or shrubs that respond slowly and often fairly late in the growing season. Only the sensitive individuals in a population of a detector bioindicator will respond to ozone and only when they experience appropriate edaphic and tropospheric conditions coupled with ozone exposures and concentrations sufficient to cause foliar injury. The distribution of ozone-sensitive genotypes (where there is little phenotypic variation) in a population of a detector bioindicator is usually not well-known. This adds uncertainty in interpreting results from detectors. When large numbers of individuals are present, confidence coefficients increase (Stolte and others 1992).

Bioindicator Plant Response to Ozone

Sentinels and detectors evaluated in the same area often show different results: sentinels may respond but detectors may not. Results from ozone plant bioindicator studies must be interpreted on the basis of a number of interacting biological, cultural, and physical factors (*table 2*). Foliar ozone injury symptoms are only an indication of previous exposure to ozone, when the concentration was high enough for a certain time period, and environmental conditions were conducive to ozone uptake and cellular injury. The visible response is a relative index of air quality for ozone during a previous exposure period. It is not possible to use this response to quantitatively assess air quality for ozone. This is because we do not yet truly understand the relationship between the occurrence and magnitude of plant response and the environmental conditions and concentrations and durations of ambient ozone present when the plant injury response was initiated. If data from a co-located or nearby mechanical ozone monitor is available, however, periods when monitored ozone concentrations have potential biological significance or meaning can be indicated, as reflected by frequency, duration, and intensity of plant responses.

Table 2 — *Biological, cultural, and physical factors affecting responses to ozone (Manning 1991).*

Biological and Cultural		
Factors		Responses
Biological		
Genetic diversity		Homogeneous plants give uniform responses; species, clones, cultivars, and provenances react differently to O ₃
Stage of plant development		Plant developmental stages and leaf age affect responses to O ₃
Cultural		
General cultural practices		Optimal or usual practices result in “typical” responses to O ₃
Growth media		Soilless media allow uniformity and reproductibility, but are less relevant than natural soils.
Nutrients		Optimal levels usually result in optimal injury from O ₃ .
Pesticides		Variable effects, ranging from none to protection, from injury to reduced tolerance to joint effects with O ₃ .
Physical		
Factors		Responses
Air movement		Must be sufficient to alter boundary layer resistance to allow O ₃ uptake.
Light		
Intensity, photoperiod, and quality		Ideal value varies for each plant; less or more than ideal value for each plant reduces O ₃ sensitivity
Temperature		Injury increases in a range from 3 to 30 °C.
Water		
Relative humidity		Controls stomatal opening exchange; uptake of O ₃ and injury should increase as relative humidity increases.
Soil moisture tolerance		Water stress increases O ₃ due to stomatal closure

Previous research has determined some well-defined bioindicators for ambient ozone (table 3). Each of these bioindicators will exhibit typical ozone injury symptoms (table 4). Tobacco cultivar Bel-W3 is the best-defined, best-described, and most widely-used bioindicator for ambient ozone (Heggstad 1991). It has become the standard sentinel bioindicator for ozone and has been used all over the world to detect ambient ozone in areas ranging from local sites to whole countries and continents. Heck and Heagle (1970) used Bel-W3 tobacco to determine incidence of phytotoxic concentrations of ozone around Cincinnati, Ohio. Bel-W3 tobacco plants were used by Manning to demonstrate ozone in Warsaw, Krakow, and forested areas in southern Poland (Bytnerowicz and others 1993). Jacobson and Feder (1974) reported the results of a network of Bel-W3 tobacco plantings in the northeastern United States. Long-range transport of ozone from metropolitan New York to Nantucket Island was demonstrated by response of Bel-W3 tobacco plants on Nantucket (Kelleher and Feder 1978). Ashmore and others (1980) mapped phytotoxic ozone in the British Isles by using networks of Bel-W3 tobacco plants. Schoolchildren in many European countries use Bel-W3 tobacco to assess relative air quality for ozone (Lipa and Votapkova 1995).

Table 3 — Selected bioindicators of ozone.¹

Plant	Latin Name	References
Sentinels		
Blue grass	<i>Poa annua</i>	Noble and Wright 1958
Bean	<i>Phaseolus vulgaris</i>	Oshima 1974, Sanders and others 1992
Clover	<i>Trifolium repens</i>	Heagle and others 1992
	<i>T. subterraneum</i>	Sanders and others 1992
Morning glory	<i>Ipomoea purpurea</i>	Manning 1977, Nouchi and Aoki 1979
Spinach	<i>Spinacea oleraceae</i>	Posthumus 1982
Tobacco	<i>Nicotiana tabacum</i>	Heck and Heagle 1970,
	Bel-W3 (sensitive)	Heaggestad 1991
	Bel-B (tolerant)	
Detectors		
Blackberry	<i>Rubus</i> spp.	Chappelka and others 1986, Manning 1991
Black cherry	<i>Prunus serotina</i>	Chappelka and others 1992, Davis and others 1982
Green ash	<i>Fraxinus pennsylvatica</i>	Davis and Wilhour 1976
Milkweed	<i>Asdepias sytiaca</i>	Duchelle and Skelly 1981
Quaking aspen	<i>Populus tremuloides</i>	Karnosky 1976, Keller 1988
Sassafras	<i>Sassafras albidum</i>	Chappelka 1992
Tulip poplar	<i>Liriodendron tulipifera</i>	Davis and Chappelka 1986, 1992; Davis and Wilhour 1976
White ash	<i>Fraxinus americana</i>	Chappelka 1992, Davis and Wilhour 1976

¹ Not all varieties or individuals in a species will be ozone-sensitive.

Several plant species have cultivars or clones that differ in their response to ambient ozone. The best defined incidence of this are the tobacco cultivars Bel-W3 (ozone-sensitive) and Bel-B (ozone-tolerant). Bel-W3 tobacco responds to ambient ozone at lower concentrations than does Bel-B. When grown together in an area, their comparative responses to ozone can provide a better description of relative air quality for ozone. If both cultivars do not respond, then the air is relatively clean and ozone concentrations are probably low. If Bel-W3 responds, but Bel-B does not, then ozone concentrations are intermediate. If both cultivars respond, then ozone concentrations are higher (Manning and Feder 1980).

Detectors are being used to assess relative long-term air quality in forested and wilderness areas. Study areas are selected and certain species are examined on an annual basis for incidence and severity of foliar ozone injury. In wilderness areas of New Hampshire and Vermont, Manning and others (1991) have surveyed black cherry, white ash, and milkweed. Chappelka and others (1986, 1992) have used black cherry, sassafras, white ash, yellow poplar, and blackberry in the southeastern United States. The USDA Forest Service's Forest Health Monitoring Program uses detector bioindicators in its forest surveys in the U.S. (Conkling and Byers 1993). Detector bioindicators are used in forested regions of the Carpathian Mountains in Poland, Ukraine, Czech Republic, Slovakia, and Rumania (Manning and others, unpublished paper).

EDU (ethylenediurea) is a chemical known to protect plants from ozone injury (Carnahan and others 1978). It can be used to verify the response of an ozone bioindicator in the field. This is especially useful if an ozone monitor is not readily available or there are questions about the nature of the response of the bioindicator. This approach was used in Poland (Bytnerowicz and others 1993). Bel-W3 tobacco plants were outplanted in a number of locations. Half of the plants were treated with EDU before outplanting. The treated plants did not develop any symptoms of foliar injury while the non-treated plants developed varying degrees of typical ozone injury symptoms, verifying that ozone was the cause.

Table 4 — Common symptoms of foliar ozone injury (Krupa and Manning 1988).

Acute injury	
Flecking	Small necrotic areas due to death of palisade cells, metallic or brown, fading to grey or white.
Stippling	Tiny punctate spots where a few palisade cells are dead or injured, may be white, black-red, or red-purple.
Chronic injury	
Pigmentation (Bronzing)	Leaves turn red-brown to brown as phenolic tan pigments accumulate.
Chlorosis	May result from pigmentation or may occur alone as chlorophyll breaks down.
Premature senescence	Early loss of leaves or fruit.

Methodology for Assessing Bioindicator Plants

Great care must be taken in assessing the response of bioindicator plants to ambient ozone. In most cases, assessment of bioindicators involves the subjective determination of the intensity or extent of acute ozone injury symptoms (*table 4*). Leaf injury evaluations should be made at regular intervals, often weekly, and by the same person. The use of a set of color photographs illustrating degrees or categories of severity of injury can help to standardize the evaluation process (Heck 1966; Heck and others 1966, 1969; Oshima 1976). If plants are left in the field for more than 1 week, then new injury on both older and new leaves needs to be estimated and recorded.

The observer should look at each leaf and visually integrate the injured areas of each leaf and then determine the percentage of the total leaf area that has been injured. Depending on the type of plant, extent of injury, and purpose of the study, the evaluation system can be quite simple and uncomplicated (*table 5*). With only six indices, this is a simple system to use. Where more precise information is required, an expanded system is more appropriate (*table 6*).

Table 5—System for evaluation of ozone injury for bean (*Phaseolus vulgaris* L.) (Manning and Feder 1980).

Injury rating	Rating Systems	
	Injury severity index	Percent leaf injury
None	0	0
Slight	1	1-25
Moderate	2	26-50
Moderate-severe	3	51-75
Severe	4	76-99
Complete	5	100

Table 6—The Horsfall-Barratt Scale for assessing foliar injury (Horsfall and Barratt 1945, Horsfall and Cowling 1978).

(Percent severity or class incidence)	
0	0
1	0-3
2	3-6
3	6-12
4	12-25
5	25-50
6	50-75
7	75-88
8	88-94
9	94-97
10	97-100
11	100

Data can be depicted graphically. Plotting weekly injury scores, or number of leaves injured, on a weekly basis against elapsed time gives a cumulative injury curve over time. If data are available from a nearby mechanical ozone monitor, ozone injury can be plotted against cumulative ozone or related to episodes in which thresholds are exceeded (Manning and Feder 1980).

As an alternative to using subjective determination of the intensity or extent of foliar injury as a measure of ozone effects, other investigators have developed bioindicator systems that use shoot biomass response from sequential harvests as a measure of ozone effects. Oshima and others (1976) developed a standardized pot culture system with alfalfa (*Medicago sativa* L.) along an ozone gradient in southern California. Heagle and others (1994) have developed an ozone bioindicator system by using sensitive (NCS) and resistant (NCR) clones of ladino clover (*Trifolium repens* L.) NCS and NCR plants are grown in standardized pot culture. Shoot biomass is removed at 28-day intervals and dry weights are obtained. Ozone impact is determined by obtaining dry weights of harvested shoot biomass and calculating the ratios of NCS to NCR. A ratio of less than one indicates that ambient ozone has had an adverse effect on foliar biomass of NCS.

Summary

Bioindicator plants for ozone can be extremely useful in assessing relative air quality, especially in areas where ozone monitors are not available. If they are not used properly, however, poor quality or misleading results will be obtained. Common factors that produce poor results are poor plant culture for sentinels, failure to adhere to regular evaluation schedules, careless and inaccurate evaluation of symptoms, and the use of more than one person to evaluate plant responses.

Care should be taken in interpreting responses of sentinel bioindicators. As they are well-watered and fertilized, they respond much more frequently than do detectors. Their response indicates the occurrence of periods of ozone exposure when detectors might respond if they were also growing under conditions of sufficient soil moisture and fertility. From an ecological perspective, response of detector bioindicators to ozone is more important than a response by a sentinel bioindicator. A response by a sentinel indicates what could happen under ideal conditions, while a response by a detector indicates what did happen under more realistic conditions.

There is an unfortunate tendency to confuse a bioindicator with a biomonitor. A bioindicator indicates that the system has been affected. A biomonitor should also indicate how much of the causal factor was present and caused the observed effect. Currently, bioindicators of ambient ozone cannot be used as biomonitors. It is not possible to make quantitative inferences about air quality for ozone on the basis of plant symptom expression alone.

With a clear understanding of the strengths and weaknesses of the use of plants as bioindicators of ozone exposure, they can be quite useful in assessing relative air quality, especially in programs like Forest Health Monitoring, in remote areas, and in emerging nations, where technology is lacking.

References

- Arndt, U.; Nobel, W.; Schweizer, B. 1987. **Bioindikatoren: Möglichkeiten, Grenzen und neue Erkenntnisse**. Ulmer, Stuttgart, Germany.
- Ashmore, M. R.; Bell, J. N. B.; Reilly, C. L. 1980. **The distribution of phytotoxic ozone in the British Isles**. Environmental Pollution 1B: 195-216.
- Bennett, J. P.; Stolte, K. W. 1985. **Using vegetation biomonitors to assess air pollution injury in National Parks: milkweed survey**. U.S. National Park Service, Natural Resources Report Series No. 85-1; 16 p.
- Burton, M. A. S. 1986. **Biological monitoring of environmental contaminants**. Tech. Report. London: Monitoring and Assessment Centre; 247 p.
- Bytnerowicz, A.; Manning, W. J.; Grosjean, D.; Chimielewski, W.; Dmuchowski, W.; Grodzinska, K.; Godzik, B. 1993. **Detecting ozone and demonstrating its phytotoxicity in forested areas of Poland: a pilot study**. Environmental Pollution 80: 301-305.
- Carnahan, J. E.; Jenner, E. L.; Wat, E. K. W. 1978. **Prevention of ozone injury to plants by a new protectant chemical**. Phytopathology 68: 1225-1229.
- Chappelka, A. H.; Chevone, B. I.; Brown, H. D. 1986. **Bioindicator survey for ozone injury in Georgia, North Carolina and South Carolina**. Phytopathology 76: 1035.
- Chappelka, A. H.; Hilderbrand, E.; Skelly, J. M.; Mangis, D.; Renfro, J. R. 1992. **Effects of ambient ozone concentrations on mature eastern hardwood trees growing in Great Smoky Mountains National Park and Shenandoah National Park**. Proceedings Annual Meeting, Air and Waste Management Association; Paper 92-150.04.
- Conkling, B. L.; Byers, G. E., eds. 1993. **Forest health monitoring field methods guide**. Las Vegas, NV: U.S. EPA.
- Davis, D. D.; Umback, D. M.; Coppolino, J. B. 1982. **Susceptibility of tree and shrub species and responses of black cherry foliage to ozone**. Plant Disease 65: 904-907.
- Davis, D. D.; Wilhour, R. G. 1976. **Susceptibility of woody plants to sulfur dioxide and photochemical oxidants**. Pub. 600/3-76-102, Corvallis, Oregon: U.S. EPA.
- deBauer, L. I. 1972. **Uso de plantas indicadoras de aeropoluto en la Ciudad de Mexico**. Agrociencia 9 (D): 139-141.
- Duchelle, S. F.; Skelly, J. M. 1981. **Response of common milkweed to oxidant air pollution in the Shenandoah National Park in Virginia**. Plant Disease 65: 661-663.
- Feder, W. A.; Manning, W. J. 1979. **Living plants as indicators and monitors**. In: Heck, W. W.; Krupa, S. V.; Linzon, S. N., eds. Handbook of methodology for the assessment of air pollution effects on vegetation. Pittsburgh, PA: Air Pollution Control Association; 9-1—9-14.
- Guderian, R.; Tingey, D. T.; Rabe, R. 1985. **Effects of photochemical oxidants on plants**. In: Guderian, R., ed. Photochemical oxidants. Berlin: Springer-Verlag; 129-295.
- Heagle, A. S.; Miller, J. E.; Sherrill, B. E. 1994. **A white clover system to estimate effects of tropospheric ozone on plants**. Journal Environmental Quality 23: 613-621.
- Heck, W. W. 1966. **The use of plants as indicators of air pollution**. Air, Water, Pollution International Journal 10: 99-111.
- Heck, W. W.; Fox, F. L.; Brandt, C. S.; Dunning, H. A. 1969. Tobacco a sensitive monitor for photochemical air pollution. U. S. Nat. Air Poll. Contr. Admin. Pub. AP 55.
- Heck, W. W.; Heagle, A. S. 1970. **Measurement of photochemical air pollution with a sensitive monitoring plant**. Journal Air Pollution Control Association 20: 97-99.
- Heggstad, H. E. 1991. **Origin of Bel-W3, Bel-C, and Bel-B tobacco varieties and their use as indicators of ozone**. Environmental Pollution 74: 264-291.
- Hernandez, T. T.; deBauer, L. I. 1989. **La supervivencia Vegetal ante la contaminacion atmosferica**. Edo, Mexico: Colegio de Postgraduados, Chapingo; 79 p.

- Horsfall, J. G.; Barratt, R. W. 1945. **An improved grading system for measuring plant disease.** *Phytopathology* 35: 655.
- Horsfall, J. G.; Cowling, E. B. 1978. **Pathometry: The measurement of plant disease.** In: Horsfall, J. G.; Cowling, E. B., eds. *How disease develops in populations.* Plant Disease, Vol. II. New York: Academic Press; 119-136.
- Jacobson, J. S.; Feder, W. A. 1974. **A regional network for environmental monitoring of atmospheric oxidant concentrations and foliar injury to tobacco plants in the eastern United States.** *Massachusetts Agricultural Experimental Station Bulletin* 604.
- Karnosky, D. F. 1976. **Threshold levels for foliar injury to Populus tremuloides by sulphur dioxide and ozone.** *Canadian Journal Forestry Research* 6: 166-169.
- Kelleher, T. J.; Feder, W. A. 1978. **Phytotoxic concentrations of ozone on Nantucket Island: long-range transport from the Middle Atlantic states over the open ocean confirmed by bioassay with ozone-sensitive tobacco plants.** *Environmental Pollution* 17: 187-194.
- Keller, Th. 1988. **Growth and premature leaf fall in American aspen as bioindications for ozone.** *Environmental Pollution* 52: 183-192.
- Krupa, S. V.; Manning, W. J. 1988. **Atmospheric ozone: formation and effects on vegetation.** *Environmental Pollution* 50: 101-137.
- Lipa, K.; Votapkova, D. 1995. **Projekt Ozone: Zaverena zprova ze sledovani prizemniho ozona v Ceske republice ve dnech 8.3 az 28.6.1995.** TEREZA, Prague, Czech Republic.
- Manning, W. J. 1977. **Morning glory as an indicator plant for oxidant air pollution: cultivar sensitivity.** *Proceedings American Phytopathology Society* 4: 192.
- Manning, W. J. 1991. **Experimental methodology for studying the effects of ozone on crops and trees.** In: Lefohn, A. S., ed. *Surface level ozone exposures and their effects on vegetation.* Chelsea, MA: Lewis Publications; 93-156.
- Manning, W. J. 1993. **Bioindicator plants for assessment of air quality: general considerations and plant responses to ambient ozone.** *Proceedings Annual Meeting, Air and Waste Management Association.* Paper 93-WA-80.01.
- Manning, W. J.; Feder, W. A. 1980. **Biomonitoring air pollutants with plants.** London: Applied Science Pub. Ltd; 142 p.
- Manning, W. J.; Bergman, J. R.; O'Brien, J. T. 1991. **Ozone injury on native vegetation in Class I wilderness areas in New Hampshire and Vermont.** *Proceedings Annual Meeting, Air and Waste Management Association.* Paper 91-144.5.
- Middleton, J. T.; Kendrick, J. B., Jr.; Schwalm, H. W. 1950. **Injury to herbaceous plants by smog or air pollution.** *Plant Disease Reporter* 34: 245-252.
- Miller, P.; Guthrey, D.; Schilling, S.; Carroll, J. 1998. **Ozone injury responses of ponderosa and Jeffrey pine in the Sierra Nevada and San Bernardino mountains in California.** In: Bytnerowicz, A.; Arbaugh, M.; Schilling, S., technical coordinators. *Proceedings of the international symposium on air pollution and climate change effects on forest ecosystems; 1996 February 5-9; Riverside, CA.* Gen. Tech. Rep. PSW-GTR-166. Albany CA: Pacific Southwest Research Station, USDA Forest Service. [this volume].
- Noble, W. M.; Wright, L. A. 1958. **Air pollution with relation to agronomic crops. II. A bio-assay approach to the study of air pollution.** *Agronomy Journal* 50: 551-553.
- Nouchi, I.; Acki, K. 1979. **Morning glory as a photochemical oxidant indicator.** *Environmental Pollution* 18: 289-303.
- Oshima, R. J. 1974. **A viable system of biological indicators for monitoring air pollutants.** *Journal Air Pollution Control Association* 24: 576-578.
- Oshima, R. J.; Poe, M. P.; Braegelmann, P. K.; Baldwin, D. W.; vanWay, V. 1976. **Ozone dosage-crop loss function for alfalfa: a standardized method for assessing crop losses from air pollutants.** *Journal Air Pollution Control Association* 26: 861-865.
- Posthumus, A. C. 1982. **Biological indicators of air pollution.** In: Unsworth, M. H.; Ormrod, D. P., eds. *Effects of gaseous air pollution in agriculture and horticulture.* London: Butterworth; 115-120.
- Sanders, G. E.; Booth, C. E.; Weigel, H. J. 1992. **The use of EDU as a protectant against ozone pollution.** In: Jager, H. J.; Unsworth, M.; De Temmerman, L.; Mathy, P., eds. *Effects of air pollution on agricultural crops in Europe: results of the European open-top chambers project.* Air Pollution Research Report. Brussels: 46, CEC; 359-369.
- Spellerberg, I. F. 1991. **Monitoring ecological change.** Cambridge, UK: Cambridge University Press; 334 p.
- Steubing, L.; Jager, H. J. 1982. **Monitoring of air pollutants by plants: methods and problems.** The Hague: W. Junk Pub.; 161 p.
- Stolte, K. W.; Duriscoe, D. M.; Cook, E. R.; Chine, S. P. 1992. **Methods of assessing response to air pollution.** In: Olson, ed. *The response of western forests to air pollution.* New York: Springer-Verlag.
- Tonneijck, A.; Posthumus, A. C. 1987. **Use of indicator plants for biological monitoring of the effects of the effects of air pollution: the Dutch approach.** *VDR Berichte* 609: 205-216.
- Weinstein, L. H.; Laurence, J. A. 1989. **Indigenous and cultivated plants as bioindicators of air pollution injury.** In: Noble, R. D., eds. *Air pollution effects on vegetation, including forest ecosystems.* *Proceedings Second US-USSR Symposium, Broomall, PA: USDA Forest Service; 201-204.*