

Deployment of Deer-Resistant Western Redcedar (*Thuja plicata*)

John Russell

JOHN RUSSELL

Research Scientist
Research Branch, British Columbia Ministry
of Forests and Range
Cowichan Lake Research Station
Box 335
Mesachie Lake, BC V0R 2N0
Tel: 250.749.6811
E-mail: John.Russell@gov.bc.ca

Russell J. 2008. Deployment of deer-resistant western redcedar (*Thuja plicata*). In: Dumroese RK, Riley LE, technical coordinators. National Proceedings: Forest and Conservation Nursery Associations—2007. Fort Collins (CO): USDA Forest Service, Rocky Mountain Research Station. Proceedings RMRS-P-57:149-153. Available at: http://www.fs.fed.us/rm/pubs/rmrs_p057.html

ABSTRACT

Protecting planted western redcedar (*Thuja plicata*) seedlings from deer browse in the Pacific Northwest and British Columbia is estimated to cost up to CAN\$ 25 million annually. Recent studies linking deer browse and needle monoterpenes has resulted in the initiation of a breeding program for deer-resistant western redcedar at Cowichan Lake Research Station on Vancouver Island. Selections that are not preferred by deer are currently going through non-preferred rapid breeding and testing cycles for enhanced needle monoterpene concentrations. However, planting stock with increased monoterpene concentrations through genetic manipulation selection and breeding alone may not ensure that seedlings will be adequately protected from browse if deployment strategies are not carefully considered. To provide more robust protection to allow seedlings to reach the free-growing stage, interactions must be evaluated of between genetic selection for enhanced monoterpene concentrations, ally enhanced plants with

developmental stage (ontogeny), and nutritional quality, as well as appropriate deployment strategies that allow deer some level of forage choice, could ensure adequate protection from browse.

KEYWORDS

needle monoterpenes, ontogenetic effects,
plant nutrients

Introduction

Deer and elk browsing on newly planted western redcedar (*Thuja plicata*) seedlings in the Pacific Northwest (PNW) can result in delayed regeneration and potential plantation failure. As a result, CAN\$ 20 to 25 million are spent annually in British Columbia to bring a plantation to free-to-grow status using individual tree guards (van Nienjenhuis 2007). Licensees who avoid planting western redcedar incur additional indirect costs, including maladapted or inappropriate species selection, and reduced manufacturing opportunities.

The role of plant secondary metabolites (PSM) in plant defenses has long been recognized (Coley and Barone 1996; Huber and others 2004). Monoterpenes, a group of PSM, have been shown to reduce herbivore predation, including ungulate feeding (for example, Duncan and others 1994; Vourc'h and others 2002a). Plants contain both beneficial and deleterious phytochemicals that impact palatability and diet selection for ungulates (Kimball and Provenza 2003).

Levels of monoterpenes in conifer needles can be impacted by genetics and environmental factors, as well as their interactions (for example, Baradat and others 1972; Vourc'h and others 2002b). Monoterpene concentrations also vary seasonally and with age (for example, Schonwitz and others 1990; Powell and Raffa 1999).

Recent studies linking deer browse and needle monoterpenes in western redcedar (Vourc'h and others 2002b; Russell and Kimball nd) have driven the initiation of a deer-resistance western redcedar breeding program at Cowichan Lake Research Station, Mesachie Lake, British Columbia (CLRS). Deer "non-preferred" selections are currently going through rapid breeding and testing cycles for enhanced needle monoterpene concentrations. However, planting stock bred for higher monoterpene levels alone may not ensure that seedlings will be adequately protected from browse. Interactions between a plant's genetic background, developmental stage (ontogeny), nutritional quality, and silviculture, particularly appropriate deployment strategies that allow deer some level of choice, must all be considered to provide robust protection from browse. This paper outlines current knowledge, the ongoing breeding program at CLRS, and future research directions in western redcedar deer resistance.

Linking Deer Browse and Needle Monoterpene Concentrations in Western Redcedar

A subsample of individuals from a western redcedar population study with family structure at Holt Creek (HC) on southern Vancouver Island indicated that trees that were heavily browsed tended to have low monoterpene content (Vourc'h and others 2002b). A subsequent complete survey of all 2200 trial trees at HC confirmed this relationship (Russell nd). Both of these studies involved seedlings planted in an uncontrolled environment. To obtain a more precise measure of the relationship between deer browse and needle monoterpenes, a study was designed using multiple copies of clones planted in deer enclosures and in small openings in the forest with

heavy deer browsing pressure. Cuttings were rooted from 60 trees at HC that varied in monoterpene concentrations and planted at the USDA National Wildlife Research Center, Olympia Field Station, Olympia, Washington (Figure 1) and 2 reforestation sites on southern Vancouver Island. Consistent results were obtained at all 3 sites, demonstrating that browse preference is a function of total needle monoterpene content (Figure 2; Russell and Kimball nd).

Some Factors Affecting Needle Monoterpene Concentrations in Western Redcedar

Genetics

In the HC trial mentioned earlier, a complete sample of needles from all test trees indicated substantial genetic variation in monoterpenes. Narrow-sense heritabilities for 6 of the major needle monoterpenes varied from 0.25 to 0.44, and coefficients of additive genetic variation varied from 19% to 27% (Russell nd). There were significant seed source effects, that is, northern British Columbia populations had greater needle monoterpene concentrations than southern British Columbia sources. Thus, a base population with substantial genetic variation in needle monoterpenes for the initiation of a breeding program is available.

Ontogenetic and Age Effects

In a fertilization study with planted western redcedar in western Oregon, total needle monoterpenes increased linearly from age 1 to 3 (Kimball 2007). Similarly, 1-year-old rooted cuttings originating from 7-year-old western redcedar trees had greater total needle monoterpene concentrations and significantly reduced browse damage than genetically comparable 1-year-old seedlings (Figure 3; Russell nd). Utilizing either 2-year-old seedlings, or rooted cuttings from older hedged donors, can increase needle monoterpene levels and potentially minimize deer browsing.



Figure 1. Western redcedar feeding choice trial with Columbian black-tailed deer at the USDA National Wildlife Center, Olympia Field Station.




Figure 2. Relationship between total needle monoterpenes and mean failure by clone in western redcedar at the USDA National Wildlife Center, Olympia Field Station.




Figure 3. Relationship between deer browse and time since outplanting duration for western redcedar stocktypes.

Plant Nutrient Levels

Nutritional quality of seedlings at the time of planting can have a paradoxical effect. High nitrogen, for example, may make a seedling more palatable to deer, whereas it has been well documented for western redcedar that fertilization at planting stimulates early growth, and thus earlier free-to-grow status (Blevins and others 2006). A higher nitrogen level may also help seedlings recover following browse (see references in Jacobs 2005). Little research has explored the relationship between nutritional quality and monoterpene production. A current study is underway in the PNW examining the influence of mineral nutrition on susceptibility and recovery of planted western redcedar to animal browse (Jacobs 2005).

Breeding for Enhanced Needle Monoterpenes

A breeding program for enhanced needle monoterpene concentrations is currently underway at Cowichan Lake Research Station. Over 60 “not preferred” selections (minimal deer browse and enhanced needle monoterpenes) from a base population of approximately 2500 range-wide western redcedar trees have been selected for the



Figure 4. Breeding young western redcedar trees at Cowichan Lake Research Station.

breeding program. In addition, a number of “deer preferred” selections (high browse and low monoterpene concentrations) have also been included. Terpenes are expressed during a seedling’s first year, and breeding can be done within 2 years (Figure 4; Russell and Ferguson forthcoming). Additional selections for deer resistance can be quickly incorporated into the program.

Deployment of Deer-resistant Seedlots

Planting seedlings that are high in needle monoterpenes doesn't necessarily guarantee browse resistance. Plants contain beneficial and deleterious phytochemicals. If bitter, high monoterpene trees are the only available food source, deer will select based on the balance between energy input and cost of detoxification (Kimball and Provenza 2003). Results from a study deploying different mixtures of hybrid poplar clones in the Columbia Valley that were preferred and avoided by deer showed that a 2:1 mixture of "not preferred" to "preferred" trees was optimal to yield final free-to-grow stocking (Stanton 2004).

The next stage in developing deer-resistant western redcedar seedlots at CLRS is to outplant mixtures of trees with varying levels of needle monoterpenes in operational trials, including genetic selections, seedlings of varying age, and rooted cuttings. The ultimate goal is to develop custom-made seedlots with varying degrees of resistance tailored for specific sites.

References

- Baradat P, Bernard-Dagan C, Fillon C, Marpeau A, Pauly G. 1972. Les terpènes du Pin maritime: aspects biologiques et génétiques. II. Hérité de la teneur en mono-terpènes. *Annales des Sciences Forestières* 29:307-334.
- Blevins LL, Prescott CE, van Nienjenhuis A. 2006. The roles of nitrogen and phosphorous in increasing productivity of western hemlock and western redcedar plantations on northern Vancouver Island. *Forest Ecology and Management* 234:116-122.
- Coley PD, Barone JA. 1996. Herbivory and plant defenses in tropical forests. *Annual Review of Ecology and Systematics* 27:305-335.
- Duncan AJ, Hartley SE, Iason GR. 1994. The effect of monoterpene concentrations in Sitka spruce (*Picea sitchensis*) on the browsing behavior of red deer (*Cervus elaphus*). *Canadian Journal of Zoology* 72:1715-1720.
- Huber DPW, Ralph S, Bohlmann J. 2004. Genomic hardwiring and phenotypic plasticity of terpenoid-based defenses in conifers. *Journal of Chemical Ecology* 30:2399-2418.
- Jacobs DF. 2005. Influence of mineral nutrition on susceptibility and recovery of planted seedlings to animal browse [proposal submitted]. Salem (OR): Oregon Department of Forestry.
- Kimball B. 2007. Personal communication. Mesachie Lake (BC): British Columbia Ministry of Forests and Range, Cowichan Lake Research Station.
- Kimball BA, Provenza FD. 2003. Chemical defense and mammalian herbivores. In: Hardege JD, editor. *Encyclopedia of life support systems (EOLSS)*. URL: <http://www.eolss.net>. Oxford (United Kingdom): Developed under the auspices of the UNESCO, EOLSS Publishers.
- Powell JS, Raffa KF. 1999. Sources of variation in concentration and composition of foliar monoterpenes in tamarack (*Larix laricina*) seedlings: roles of nutrient availability, time of season and plant architecture. *Journal of Chemical Ecology* 25:1771-1797.
- Russell J. nd. [Unpublished data]. Located at: British Columbia Ministry of Forests and Range, Cowichan Lake Research Station, Mesachie Lake, British Columbia.
- Russell JH, Ferguson CR. Preliminary results from five generations of a western redcedar (*Thuja plicata*) selection study with self mating. *Tree Genetics and Genome*. Forthcoming.
- Russell J, Kimball B. nd. [Unpublished data]. Located at: British Columbia Ministry of Forests and Range, Cowichan Lake Research Station, Mesachie Lake, British Columbia.
- Schonwitz R, Lohwasser K, Kloos M, Ziegler H. 1990. Seasonal variation in the monoterpenes in needles of *Picea abies* (L.) Karst. *Trees: Structure and Function* 4:34-40.
- Stanton B. 2004. Hybrid poplar clonal deployment: A case study from the Northwest. In: IEG-40 meeting, advancing regeneration technologies for the deployment of elite southern pine germplasm; 2004 June 22-24; Jekyll Island, GA.
- van Nienjenhuis A. 2007. Personal communication. Saanich (BC): Saanich Forest Center, Western Forest Products Limited.
- Vourc'h G, de Garine-Wichatitsky M, Labbe A, Rosolowski D, Martin J-L, Fritz H. 2002a. Monoterpene effect on feeding choice by deer. *Journal of Chemical Ecology* 28:2405-2421.
- Vourc'h G, Russell JH, Martin J-L. 2002b. Linking deer browse, terpenes and genetic variation in Cupressaceae. *Journal of Heredity* 93(5):370-376.