

Is there a link between anthropogenic disturbance and the diversity and abundance of rodent flea communities ?

Megan M . Friggens

School of Forestry , Northern Arizona University , Flagstaff , AZ , USA 86011 . E-mail : mmf38@nau.edu

Key words : Flea assemblages , rodents , vector-borne disease , anthropogenic disturbance

Introduction Fleas are among the most common arthropod vectors of many of rodent borne diseases like plague (*Yersinia pestis*) . Human outbreaks of plague are commonly associated with the presence of peridomestic rodent species (Perry and Featherston, 1997) . Anthropogenic disturbance with respect to both habitat use and climate change can affect disease emergence and prevalence through its impact on hosts and host-parasite (vector) ecology (Daszak et al . , 2001) . Vector-borne disease transmission is often related to species diversity (Keesing et al . , 2006) . In addition , flea abundance and host spectrum (number of hosts infested) relate to the likelihood of flea-mediated disease transmission , particularly with respect to plague (Krasnov et al . , 2006) . Anthropogenic disturbance can influence rodent community diversity (Tikhonova et al . , 2006) , which in turn can affect flea diversity (Krasnov et al . , 2004) , and may lead to decreased host specificity among flea parasites (Gettlinger and Ernest , 1995) . The goal of this paper is to examine the influence of anthropogenic disturbance on flea communities from a variety of habitats across the world . In particular , this analysis focuses on the effect of disturbance on flea diversity and flea species abundance and specificity behaviors .

Methods Literature was compiled on studies conducted across the world that reported the entire flea assemblage from comprehensive rodent community surveys (Friggens , 2008 for comprehensive list) . Basic criteria for inclusion in this analysis were studies that demonstrated live capture trapping and active flea collection and provided habitat characterization and detailed data for flea and rodent species . Studies sites were categorized according to three impact levels : (1) High impact such as urban or densely populated areas ; (2) Intermediate , which included rural villages , and crop and rangelands ; and , (3) Low impact sites characterized as remote or wild habitat . For each study/habitat type , rodent and flea diversity (Shannon's H) and abundance (when available) , number of infested hosts and average flea burden per host species were calculated . Data was analyzed using t-tests or linear regression in SigmaPlot 9.0 .

Results Thirty seven studies were included in this analyses . These comprised 11 high impact , 16 intermediate , and 10 low impact sites . Desert , Grassland and Deciduous forest type habitats were represented in each impact category . Overall , rodent and flea diversity was lowest at high impact sites (Figure 1) . As rodent diversity increases , the average number of fleas/host decreased . Flea diversity and number of host species infested was highest in habitats which experienced intermediate levels of habitat disturbance (Figure 2) .




Figure 1 Rodent vs . Flea diversity (Shannon's H) for habitats divided according to 1 of 3 disturbance levels ; * $P < 0.05$ when compared to high impact sites .




Figure 2 Flea infestation parameters for rodents collected at 3 habitat types categorized according to disturbance level ; * $P < 0.05$ when compared to high impact sites .

Conclusions Transmission of zoonotic disease from wild reservoirs to humans most commonly occurs in rural environments . These sites provide the peridomestic rodent species essential to carrying disease from the wild reservoir hosts to commensal rodent living in proximity to humans . In addition , the results of this analysis suggest that the characteristics of flea assemblages within these communities appear to be conducive to plague transmission . Namely , flea diversity and host spectrum is greatest in areas that experience moderate amounts of disturbance .

Acknowledgements This research was funded by Paulette Ford (US Forest Service Rocky Mountain Research Station , Albuquerque , New Mexico , USA) .