

ETHANOL ACCUMULATION IN DROUGHT-STRESSED CONIFER SEEDLINGS

Daniel K. Manter¹ and Rick G. Kelsey

USDA Forest Service, Pacific Northwest Research Station, Corvallis, Oregon 97331, U.S.A.

In this study, we investigated the effect of drought stress on ethanol production and accumulation in tissues from seedlings of three conifers (Douglas-fir, lodgepole pine, and ponderosa pine) with increasing degrees of tolerance to drought stress, respectively. Significant ethanol accumulation was only observed in their aerial tissues when severely stressed (water potential < -3.0 MPa or water content < 0.4 g g⁻¹ dry mass), with needles accumulating greater quantities than the sapwood or phloem. All species had the same predawn water potentials when sampled, but they were experiencing different levels of stress based on their foliar water contents. Severely stressed Douglas-fir contained higher ethanol and lower water contents (except phloem) than the more drought-tolerant ponderosa pine seedlings. Lodgepole pine with intermediate drought tolerance tended to have intermediate quantities of both components. The mechanistic basis of ethanol accumulation associated with drought stress remains to be determined, especially in larger trees. We speculate that increased levels of in situ ethanol synthesis in seedlings may be associated with heat injury (e.g., membrane damage) due to a reduction in transpirational cooling after stomatal closure. Drought-induced hypoxia may also contribute to ethanol accumulation in the sapwood.

Keywords: Douglas-fir, fermentation, lodgepole pine, ponderosa pine, water stress.

Introduction

Water stress is often a prerequisite to outbreaks of phytophagous insects (Mattson and Haack 1987; Waring and Cobb 1992; Clancy et al. 1995) and pathogens (Houston 1987; Desprez-Loustau et al. 2006) in forest ecosystems. Water deficits can weaken chemical defenses of conifers by causing decreases in production and flow of terpene oleoresins (Waring and Cobb 1992; Kolb et al. 1998) and increases in the production of stress-induced volatiles (Kimmerer and Kozlowski 1982), such as ethanol, which functions as a kairomone, attracting ambrosia and some species of bark beetles (Moeck 1970; Klimetzek et al. 1986; Liu and McLean 1989; Kelsey and Joseph 2001). In addition, if ethanol is released in binary or tertiary combinations with host monoterpenes or pheromones, then the attraction of some bark beetles, wood borers, or root weevils can be greatly increased or synergized (Tilles et al. 1986; Phillips 1988; Schroeder and Lindelöw 1989; Rieske and Raffa 1991; Byers 1992; Joseph et al. 2001; Miller 2006). For example, heavy bark beetle attack (107 beetles m⁻²) has been recorded on severely water-stressed Douglas-fir branches with high amounts of ethanol, whereas adjacent branches with low ethanol were not attacked (Kelsey and Joseph 2001). These same bark beetles have been observed attacking drought-stressed Douglas-fir in the forest (Stoszek 1973).

Ethanol synthesis occurs whenever some form of stress impairs aerobic respiration, either by limiting the O₂ supply or

by damaging the mitochondria (Drew 1997; Tadege et al. 1999). The physiological role of ethanol fermentation is the regeneration of NAD⁺ that is needed in glycolysis for the continued production of pyruvate and ATP energy. This energy is critical for stabilizing cellular pH and avoiding lethal acidosis until aerobic respiration resumes, otherwise the cells die (Roberts et al. 1984, 1985; Fox et al. 1995; Gout et al. 2001; Felle 2005). Fermentation can also produce metabolic intermediates, but this may be more important in situations such as developing and germinating pollen where it occurs under aerobic conditions simultaneously with respiration (Tadege and Kühlemeier 1997; Mellema et al. 2002). The control of ethanol biosynthesis has been hypothesized to involve cellular pH regulating the activities of the key enzymes, with activation of pyruvate decarboxylase (PDC) initiating ethanol synthesis when cellular acidity increases in response to stress (pH-stat; Roberts et al. 1984, 1985; Fox et al. 1995; Gout et al. 2001; Felle 2005). An alternative hypothesis proposes that ethanol biosynthesis is regulated by the concentration of pyruvate and enzyme kinetics (PDH/PDC-stat; Tadege et al. 1999). Under aerobic conditions, pyruvate dehydrogenase (PDH) reacts with pyruvate for use in respiration, leaving minimal amounts available for ethanol synthesis. Upon impairment of respiration, pyruvate levels will increase until sufficient amounts of PDC are reached to initiate ethanol production.

In trees, a variety of stresses can cause the tissues to produce ethanol, including flooding, pathogens, freezing, crushing, heat, pollutants, and water deficits (Kimmerer and Kozlowski 1982; MacDonald and Kimmerer 1989, 1991; Kelsey and Joseph 1998, 2003). Severely water-stressed birch (*Betula papyrifera* Marsh.) leaves at -2.5 MPa contained elevated quantities of ethanol, whereas none was detected in leaves with a potential of -1.3 MPa (Kimmerer and Kozlowski 1982). The drier leaves

¹ Author for correspondence; current address: USDA–Agricultural Research Service–Northern Plains Area, Soil-Plant-Nutrient Research Unit, Fort Collins, Colorado 80526, U.S.A.; e-mail: daniel.manter@ars.usda.gov.

were wilted and slightly necrotic. Ebel et al. (1995) found that yellowing leaves from severely water-stressed (below -2.5 MPa) apple (*Malus domestica* Borhk.) seedlings released about half the quantity of ethanol emitted by leaves from well-watered seedlings, although the difference was not statistically significant. In this case, stomatal closure could have limited ethanol release from the stressed plants. Kelsey and Joseph (2001) induced cavitation and water stress in branches of Douglas-fir by freezing them with dry ice. After 3 wk, when water contents had declined by 60%, their ethanol concentrations were ca. 1000 and 100 times higher in the sapwood and needles, respectively, compared with controls. Aside from these observations, there is little known about the production, accumulation, or release of ethanol from drought-stressed trees.

The objectives of this study were (1) to determine to what extent ethanol accumulates in root, stem, and needle tissues of drought-stressed conifer seedlings; (2) to determine whether ethanol accumulation differs among conifer species of varying drought tolerance; (3) to determine how drought stress affects the ability of phloem tissues to synthesize ethanol; and (4) to determine whether stomatal closure and reduced photosynthesis from water stress affects carbohydrate contents of the phloem sufficiently to influence ethanol synthesis.

Material and Methods

Seedling Material

Two-year-old bare-root seedlings of ponderosa pine (*Pinus ponderosa* Dougl. ex Laws, zone 14), lodgepole pine (*Pinus contorta* Dougl. ex Loud. var. *latifolia*, zone 2), and interior Douglas-fir (*Pseudotsuga menziesii* [Mirb.] Franco var. *glauca*) were obtained from commercial sources, planted in 2-L pots in February or March 2003, and grown under ambient field conditions at the Oregon State University Botany Farm (Corvallis, OR). The potting mixture consisted of a 1 : 1 : 1 : 1 (v/v) mix of loam, peat, pumice, and fine Douglas-fir bark. Seedlings were fertilized in June 2003 with ca. 15 g of Osmocote 18 and 15 g of Scotts STEP Hi Mag Formula (Teufel Nursery, Portland, OR). Beginning on June 26, 2003, differential water treatments were applied to 20 seedlings of each species in a randomized complete block design. Water treatments were applied by two pressure-compensated drip emitters per pot designed to deliver a total of 2 L (high), 0.5 L (low), or 0 L (none) of water daily.

Sample Collection

The onset of drought stress was monitored by periodic measurements of predawn (0400 to 0600 hours) plant-water potentials on the treated seedling by excising a current-year lateral branch ($n \geq 16$) using a Scholander pressure bomb (PMS Instruments, Corvallis, OR). On the basis of these measurements, the ponderosa and lodgepole seedlings were sampled for ethanol, water content, and carbohydrates on July 22, 2003, and the Douglas-fir seedlings were sampled on August 4, 2003. On each sample date, the stem, root, and needle samples were removed from each seedling and stored on ice while being processed. Samples of 10 current-year needles from the uppermost whorl were used for constitutive ethanol analysis. A 5.0-cm stem segment was removed from 4 to 9 cm above the soil and cut into five 1.0-cm pieces. The phloem, with thin outer bark

attached on the outside and probably the cambium on the inside, was removed from four pieces; three were combined in a loosely capped vial and held on ice for measurement of anaerobic ethanol production rates and one for carbohydrate analysis was air-dried in an open vial. Phloem and sapwood from the fifth stem piece were separated and analyzed for constitutive ethanol. A random sample of ca. 1.5 g of live roots was washed in deionized water, blotted dry with paper towels, and used for constitutive ethanol analysis. All tissue samples, except the phloem samples for carbohydrate analysis or anaerobic treatment, were sealed in 4-mL glass vials with screw caps lined with Teflon (polytetrafluoroethylene; PTFE) and were immediately frozen with dry ice for storage at -36°C .

Constitutive Ethanol Analysis

In preparation for analysis, frozen samples were thawed on ice, weighed into headspace autosampler vials (22 mm $\times$ 75 mm, 22 mL volume), sealed with PTFE-lined butyl rubber septa, and heated for 30 min at 102°C to deactivate any remaining enzyme activity. Constitutive ethanol contents were analyzed by the headspace gas chromatography methods outlined by Kelsey and Manter (2004). External standard curves with three calibration levels were prepared for ethanol (100%, Apper Alcohol and Chemical, Shelbyville, KY) by diluting with deionized water. A syringe was used to deliver 5.0 μL of each solution into separate autosampler vials that were analyzed interspersed with the tissue samples. Replicates of these solutions were averaged for each calibration level in the curve.

Ethanol Production Rates in Anaerobic Phloem

All conifer tissues synthesize ethanol under anaerobic conditions, with the phloem and roots producing the greatest quantities (Kelsey et al. 1998). Enzymes for ethanol synthesis are maintained in tree tissues to accommodate brief periods of hypoxia or anoxia (Harry et al. 1988; Kimmerer and Stringer 1988). In maize root tips, the synthesis of additional fermentation enzymes occurs within 2 to 4 h after the onset of hypoxia (Sachs et al. 1980; Chang et al. 2000). In tobacco roots, the activity of PDC, the enzyme that initiates and regulates ethanol synthesis, increased by 25% after 8 h of hypoxic treatment (Tadege et al. 1998). Therefore, we used phloem ethanol production rates upon exposure to short periods of anoxia (i.e., ethanol production potential), but before the onset of carbohydrate substrate limitations, as a surrogate measure of the in situ content and activity of ethanol-related enzymes. Fresh phloem samples were sealed with septa in headspace vials (22 mL) that were purged with N_2 for 30 s and then incubated in the dark at 30°C for 2, 4, or 8 h (Kelsey et al. 1998). Immediately after incubation, the samples were heated for 30 min at 102°C to deactivate enzyme activity, and the ethanol contents were analyzed by the methods described earlier for constitutive ethanol. Ethanol production rates (g ethanol g^{-1} phloem [fresh mass] h^{-1}) were determined as the slope of the linear relationship between ethanol content and incubation time.

Carbohydrate Analysis

Air-dried phloem samples were ground with a Wiley mill (40 mesh) and stored at 4°C until extraction. Soluble carbohydrates

were extracted from samples of ca. 0.05 g with three 3-mL additions of methanol : chloroform : water (12 : 5 : 3; Dickson 1979). The aqueous layer from each extraction was combined in a new vial, followed by the addition of 1 mL H₂O and 1.4 mL chloroform for phase separation of pigments and lipids (Dickson 1979). Sugars in the aqueous layer were derivatized before gas chromatography analysis, using a procedure similar to that of Eklund and Little (1998). Briefly, 250 μ g of the internal standard phenyl- β -D-glucopyranoside was added to a 1-mL aliquot of the aqueous layer and evaporated to dryness. Samples were then redissolved in 100 μ L of pyridine containing 2.5 mg hydroxylamine HCL and heated for 20 min at 70°C, with occasional mixing to generate the oximes of fructose and glucose. Trimethylsilyl derivatives of the sugar oximes were formed by adding 100 μ L of trimethylsilylimidazole and incubating for 20 min at 70°C. Two microliters of the derivatized samples were analyzed with a Hewlett Packard 5890 Series II gas chromatograph equipped with a flame ionization detector and Phenomenex ZB-5 column (30 m $\times$ 0.25 mm inside diameter, 0.25 μ m film thickness). Helium was the carrier gas, with a column flow of 1.0 mL min⁻¹ and a split flow of 10 mL min⁻¹ (split ratio 1 : 10). The column temperature program started at 175°C. After an initial hold of 1 min, the temperature increased at 25°C min⁻¹ to a final temperature of 325°C, which was held for 6 min. Injector and detector temperatures were 350° and 375°C, respectively. Soluble carbohydrate contents (sum of sucrose, glucose, and fructose) were quantified using a four-level standard curve for each compound, with phenyl- β -D-glucopyranoside as an internal standard and derivatized by the process described earlier.

Statistical Analysis

An ANOVA was conducted using the General Linear Model procedure of SAS, version 9.1. The experiment was arranged as a randomized complete block with 20 replications for each treatment in a species. Fisher's least significant difference test was used for comparison of response variables between species and water treatments when the *F*-test for main effects was significant ($P = 0.05$). To meet the equal variance assumption of ANOVA, ethanol contents were log transformed for analysis. All reported values are the least squares means and pooled standard errors, except ethanol contents, which are back-transformed least squares means (median) and pooled standard errors. Tissue ethanol production potential was determined from the slope of the linear relationship between ethanol content and anaerobic incubation time (g ethanol g⁻¹ phloem [fresh mass] h⁻¹), using the line-fitting procedure in Microsoft Excel 2000.

Results

Water stress imposed by the differential drip-irrigation treatments was monitored by predawn water potential measurements. Measurements obtained from the morning of sampling (fig. 1) show significant differences among treatments (high: -0.48 ± 0.08 MPa, low: -0.85 ± 0.08 MPa, none: -3.48 ± 0.08 MPa). Although sampling was done on different days (July 22 for lodgepole and ponderosa pine, August 4




Fig. 1 Predawn xylem water potentials (MPa) in conifer seedlings exposed to various levels of drought stress at the time of sampling for ethanol, water, and carbohydrate contents. Treatments are high water (2.0 L d⁻¹), low water (0.5 L d⁻¹), and none (0.0 L d⁻¹).

for Douglas-fir), there were no significant differences in water potentials among species.

Constitutive ethanol content differed among species, watering treatment, and tissues. The ethanol in needles (fig. 2A), sapwood (fig. 2B), and phloem (fig. 2C) increased with increasing water stress between the low- and no-water treatments for all three species. In contrast, the ethanol content in root tissues either remained unchanged (ponderosa pine) or exhibited a small but statistically significant decline with increasing water stress (Douglas-fir and lodgepole pine; fig. 2D). Roots were the only tissue in which ethanol levels did not show an increase with stress and the only tissues rinsed with water while sampling (to remove debris). Ethanol in flooded Douglas-fir roots can diffuse into the surrounding water (Joseph and Kelsey 1997), but whether the rinse was sufficient to reduce root ethanol contents is uncertain. Given that the root water contents in each species showed the same pattern across treatments as the other three tissues (fig. 2E–2H), we suspect the rinse had minimal effect on ethanol levels.

Within each species, constitutive ethanol contents in the various tissues were highest in the needles, followed by the phloem (except for Douglas-fir with no water), sapwood, and root tissues (fig. 2A–2D). When each treatment was averaged across all three species, the greatest change in constitutive ethanol due to drought stress (low- to no-water treatment) was seen in the sapwood (about a fivefold increase), followed by the needles (about a fourfold increase), phloem (about a two-fold increase), and roots (ca. 0.7-fold change). The aerial tissues of Douglas-fir generally had the highest ethanol contents and those of ponderosa pine the least, for all treatments. In high- and low-water treatments, the ethanol in aerial tissues of lodgepole pine was intermediate between that of the other two species but usually significantly different from one or the other (fig. 2A–2C). At the most severe level of stress, ethanol in lodgepole pine sapwood and phloem was intermediate between that of the other two species (fig. 2B, 2C), whereas its needles had the highest concentrations (fig. 2A). Like constitutive ethanol, water content also differed among species, watering treatments, and tissues. However, the water contents of all




Fig. 2 Constitutive ethanol concentrations (A–D, back transformed) and water content (E–H) in tissues from conifer seedlings exposed to various levels of drought stress. DF = Douglas-fir, LP = lodgepole pine, PP = ponderosa pine. Treatments are high (2.0 L d^{-1}), low (0.5 L d^{-1}), and none (0.0 L d^{-1}). Bars with different letters are significantly different at $P < 0.05$; $n = 20$. DM = dry mass.

tissues (fig. 2E–2H) declined in each species with increasing water stress between the low- and no-water treatments.

Changes in phloem carbohydrate content in response to drought stress differed by species. They were unchanged for Douglas-fir, increased for ponderosa pine, and decreased for lodgepole pine between the low- and no-water treatments (fig. 3A); corresponding phloem water contents decreased between

these treatments (fig. 3B). Phloem ethanol production potentials (fig. 3C), estimated by incubating tissues in anaerobic conditions, declined with increasing drought stress for Douglas-fir and lodgepole and showed no significant differences for ponderosa pine.

Phloem ethanol production potentials showed a positive log-log relationship ($R^2 = 0.800$) with tissue water content (fig. 4).




Fig. 3 Carbohydrate content (A), water content (B), and ethanol production potential (C) from phloem tissues of conifer seedlings exposed to various levels of drought stress. DF = Douglas-fir, LP = lodgepole pine, PP = ponderosa pine. Treatments are high (2.0 L d^{-1}), low (0.5 L d^{-1}), and none (0.0 L d^{-1}). Bars with different letters are significantly different at $P < 0.05$; $n = 20$. DM = dry mass.

In ca. 10% of the samples, the 8-h time point was not used in calculating the production potential because ethanol contents had begun to increase exponentially, indicating the induction of additional enzyme synthesis. For all Douglas-fir tissues except roots, ethanol accumulation occurred once water contents fell below ca. 0.4 g g^{-1} dry mass (fig. 5). Similar results were observed for ponderosa and lodgepole pine (data not shown).

Discussion

In this study, we examined ethanol accumulation in various tissues from seedlings of three conifer species after exposure to high-, low-, and no-water treatments. Those in the low-water treatment received 25% of the water supplied to the high-water seedlings. Although this caused a statistically significant de-

crease in predawn water potentials, it was not limiting enough to create a mild level of water stress (-1 to -2 MPa). The no-water treatment, however, did cause severe levels of stress (-3 to -4 MPa) similar to that reported by Anekonda et al. (2002). The proportion of severely stressed seedlings that might have recovered had they been watered at the end of our treatments was not determined. However, Pharis (1966) reports that the dry-mass water content of needles on 2-yr-old seedlings, like ours, is indicative of the whole-plant lethal point for predicting probability of mortality from water stress. Comparing the needle water contents of our seedlings (fig. 2E) with his values suggests that, had they been rewatered, 50%–100% of our severely stressed ponderosa pine would have lived, whereas few, if any, rewatered Douglas-fir seedlings from the same treatment group would have recovered. Pharis (1966) did not test lodgepole pine, but its hydraulic properties are similar to those of ponderosa pine (Pinol and Sala 2000). Assuming their lethal foliar water contents are similar, ca. 25%–50% of the severely stressed lodgepole seedlings might have recovered with watering. These projected levels of survival among the three species are the same as their relative ranking of drought tolerance (Minore 1979).

The amounts of ethanol accumulated varied among treatments, species, and tissue types. Within species, similar tissues from seedlings without stress in the high- and low-water treatments contained the same quantities of ethanol, except for roots of lodgepole pine (fig. 2A–2D). All aerial tissues from severely stressed seedlings contained greater ethanol contents than the corresponding tissues from seedlings without stress. When severely stressed, the least drought-tolerant Douglas-fir seedlings contained higher ethanol and lower water levels (except in phloem) than the most drought-tolerant ponderosa pine seedlings. Lodgepole pine, considered intermediate in drought tolerance, tended to have intermediate quantities of ethanol and water, with the exceptions of ethanol in the needles and water in the phloem.




Fig. 4 Relationship between ethanol production potential and water content in phloem tissues from conifer seedlings exposed to various levels of drought stress. DF = Douglas-fir, LP = lodgepole pine, PP = ponderosa pine. Vertical reference lines correspond to water contents of 0.4 and 2.8 g g^{-1} dry mass (DM; back transformed).




Fig. 5 Relationship between ethanol accumulation and water content in Douglas-fir needles (A), sapwood (B), and phloem (C). Values were not transformed; therefore, the median point of each treatment is comparable to the bars shown in fig. 2. DM = dry mass.

Ethanol accumulation was most pronounced at very low water contents (ca. 0.4 g g^{-1} dry mass) as shown for Douglas-fir when cells were either near death or dying based on needle water contents discussed earlier. We monitored ethanol production potential and carbohydrate contents of the phloem tissue (because of its capacity to produce ethanol in woody plants; Kelsey et al. 1998) in drought-stressed seedlings, which we hypothesized a priori may potentially explain in situ rates of ethanol synthesis and subsequent accumulation. However, neither factor was significantly correlated with ethanol accumulation, and ethanol production potential declined dramatically with increasing drought stress. Therefore, some other physiological changes appear to be controlling ethanol accumulation in drought-stressed seedlings due to either an increased rate of synthesis or a decrease in dissipation or metabolism.

In regard to the latter, reduced sap flux after stomatal closure and subsequent tissue drying may somehow interrupt aerobic respiration by reducing oxygen supplies or damaging the mito-

chondria, since one of these is required for ethanol synthesis (Tadege et al. 1999). Reduced sap flow might have contributed to sapwood hypoxia because there is increasing evidence that oxygen supplied by the transpirational stream is critical for adequate aeration of its live parenchyma cells (Hook and Brown 1972; Eklund and Lavigne 1995; Gansert et al. 2001; del Hierro et al. 2002; Mancuso and Marras 2003; Spicer and Holbrook 2005). Our seedlings had thin stems and bark, and it is unknown whether their lateral oxygen diffusion was sufficient to prevent sapwood hypoxia from reduce sap flow. Consequently, it is still necessary to determine whether drought-associated sapwood hypoxia can induce ethanol synthesis in seedlings and especially in mature trees with longer lateral gas diffusion paths as a consequence of thicker stems and bark.

Ethanol synthesis in the phloem was probably not initiated by hypoxia caused by reduced oxygen supplies in the transpirational stream, because most of its oxygen comes from lateral diffusion with the atmosphere (Hook and Brown 1972; Eklund and Lavigne 1995; Mancuso and Marras 2003). The dramatic ca. 93% (see fig. 4, difference between ethanol production at 0.4 and 2.8 g g^{-1} dry mass, denoted by the vertical lines) decline in phloem ethanol production potentials, an index of ethanol-related enzyme contents and activity, may be an indicator that hypoxia was minimal, since we have determined that carbohydrate availability was not restricting ethanol production (fig. 3A, 3C). Expression of PDC, the enzyme controlling ethanol synthesis, is upregulated under hypoxia (Tadege et al. 1998; Kürsteiner et al. 2003). Thus, had the phloem in severely stressed seedlings been hypoxic, the ethanol production potential would have increased or remained unchanged instead of undergoing the sharp decline observed (fig. 3C). In the absence of hypoxia, ethanol production in the phloem would require some sort of damage to the mitochondria.

Loss of transpirational cooling after stomatal closure and subsequent increases in tissue temperatures sufficient to cause heat stress is another consequence of reduced sap flux that could stimulate ethanol synthesis in all aerial tissues, especially the foliage. Water potentials of our severely stressed seedlings (fig. 1) were well below the levels needed to close stomata: -1.6 , -1.8 , and -2.3 MPa for lodgepole, ponderosa, and Douglas-fir, respectively (Lopushinsky 1969). When water stress reduces maximal stomatal conductance of *Heteromeles arbutifolia* leaves in the range of 75% to 25% of well-watered plants, the leaf temperatures increase 2° – 3°C , with maximum temperature reaching 46° – 52°C (Valladares and Pearcy 1997). This level of heat can be lethal depending on its duration and the tissues' prior acclimation to elevated temperatures (Chen et al. 1988; Anderson et al. 1990). Heating pepper, corn, or wheat leaves from 45° to 55°C for 30 min results in electrolyte leakage and ethanol synthesis (Anderson 1994; Anderson and Peters 1994). Heat damage of the tonoplast (Chen et al. 1988) will allow acidic cell sap (Gout et al. 2001) to leak from the central vacuole into the cytoplasm. A declining cytoplasmic pH activates PDC and ethanol synthesis as postulated by the pH-stat mechanism (Roberts et al. 1984, 1985; Xia and Roberts 1994; Fox et al. 1995; Gout et al. 2001; Felle 2005). In addition, any heat damage to the mitochondria sufficient to reduce respiration would enhance the pool of pyruvate available for ethanol production as proposed by the PDH/PDC-stat mechanism (Tadege et al. 1999). This is the same mechanism proposed for ethanol synthesis and

accumulation in water-stressed Douglas-fir branches (Kelsey and Joseph 2001).

Animal and plant mitochondria can experience heat damage at ca. 45°C, given sufficient duration (Chen et al. 1988; Cole and Armour 1988). Stem tissues of mature ponderosa pine stressed by heat during a wildfire contain elevated ethanol concentrations 2 wk after the burn (Kelsey and Joseph 2003). Heating stems of well-watered Douglas-fir and ponderosa pine seedlings to 40°, 50°, or 60°C for 4–8 h results in ethanol accumulation (D. K. Manter and R. G. Kelsey, unpublished data). We suspect stems of severely water-stressed seedlings would give comparable results in a shorter period. Interestingly, roots were the only tissues that did not accumulate ethanol during drought stress, perhaps because of the lack of exposure to elevated temperatures (belowground) and heat stress.

In addition to increased in situ synthesis, ethanol accumulation is also affected by its dissipation. In the absence of water deficits, when ethanol is synthesized in roots, cambium, or phloem, it diffuses into the xylem sap and moves rapidly upward with the transpirational stream (MacDonald and Kimmerer 1991; Joseph and Kelsey 1997; Kreuzwieser et al. 1999). Loss of sap flow with stomatal closure restricts ethanol movement to diffusion from sites of synthesis, thus contributing to higher tissue concentrations. All aerobic cells can metabolize ethanol back into primary cellular constituents, allowing the tree to capture its carbon and energy, which is more beneficial on a whole-tree basis than releasing it into the atmosphere (MacDonald and Kimmerer 1993). Leaves are particularly effective, metabolizing 99.5% of the ethanol supplied in the transpirational stream (MacDonald and Kimmerer 1993; Kreuzwieser et al. 1999, 2001). We did not measure ethanol metabolism, but it is reasonable to believe that heat-stress damage to membranes and changes in cytoplasmic pH greatly limit this mechanism for dissipating ethanol in severely drought-stressed tissues.

Release of ethanol from leaves to the atmosphere is minimal, with only 0.5% or less supplied by the transpirational stream escaping (MacDonald and Kimmerer 1993; Kreuzwieser

et al. 1999, 2001), primarily through the stomata (Schade and Goldstein 2001, 2002). Some ethanol may also volatilize from the stems, as demonstrated for lodgepole pine (Gara et al. 1993), but, again, the amounts are small, with only 0.2% or less of the total produced in flooded lodgepole roots escaping from lenticels (Crawford and Finegan 1989). Loss of ethanol by volatilization may change with the severity of water stress. Since ethanol is extremely soluble in water, the highest amounts may escape during initial tissue drying while stomata are open. After the stomata close and stem shrinkage occurs (Ogigirigi et al. 1970), the rates of water and ethanol losses to the atmosphere probably decline.

In conclusion, all aerial tissues of our drought-stressed conifer seedlings accumulated ethanol, but only when the tissues were near death or dying. At the time of sampling, all three species exhibited similar predawn water potentials; however, foliar water contents differed significantly among species. We observed an inverse relationship between ethanol accumulation and tissue water contents. Douglas-fir, the least drought-tolerant species was the most stressed, and its tissues contained the highest levels of ethanol. Ponderosa pine, the most drought-tolerant species, was the least stressed and had the lowest tissue ethanol concentrations. Further work is needed to clarify all factors contributing to in situ ethanol synthesis and dissipation in water-stressed seedlings. Whether these same responses for ethanol can be detected in larger mature trees under natural drought conditions remains to be determined and is important ecologically because of ethanol's potential influence on selection of stressed trees as hosts by various types of beetles.

Acknowledgments

We thank Marion Brodhagen and Jane L. Hayes for their review of an early version of this manuscript. Manufacturer and product brand names are given for the reader's convenience and do not reflect endorsement by the U.S. government. This article was the work of U.S. government employees engaged in their official duties and is exempt from copyright.

Literature Cited

- Anderson J, G McCollum, W Roberts 1990 High temperature acclimation in pepper leaves. *Hortscience* 25:1272–1274.
- Anderson JA 1994 Production of methanol from heat-stressed pepper and corn leaf disks. *J Am Soc Hortic Sci* 119:468–472.
- Anderson JA, DC Peters 1994 Ethylene production from wheat seedlings infested with biotypes of *Schizaphis graminum* (Homoptera: Aphididae). *Environ Entomol* 23:992–998.
- Anekonda TS, MC Lomas, WT Adams, KL Kavanagh, SN Aitken 2002 Genetic variation in drought hardiness of coastal Douglas-fir seedlings from British Columbia. *Can J For Res* 32:1701–1716.
- Byers JA 1992 Attraction of bark beetles, *Tomicus piniperda*, *Hylurpogs palliatus*, and *Trypodendron domesticum* and other insects to short-chain alcohols and monoterpenes. *J Chem Ecol* 18: 2385–2402.
- Chang WWP, L Huang, M Shen, C Webster, AL Burlingame, JKM Roberts 2000 Patterns of protein synthesis and tolerance of anoxia in root tips of maize seedlings acclimated to a low-oxygen environment, and identification of proteins by mass spectrometry. *Plant Physiol* 122:295–317.
- Chen Y-R, M Chou, S-S Ren, Y-M Chen, C-Y Lin 1988 Observations of soybean root meristematic cells in response to heat shock. *Protoplasma* 144:1–9.
- Clancy KM, MR Wagner, PB Reich 1995 Ecophysiology and insect herbivory. Pages 118–180 in WK Smith, TM Hinkle, eds. *Ecophysiology of coniferous forests*. Academic Press, San Diego, CA.
- Cole A, EP Armour 1988 Ultrastructural study of mitochondrial damage in CHO cells exposed to hyperthermia. *Radiat Res* 115:421–435.
- Crawford RMM, DM Finegan 1989 Removal of ethanol from lodgepole pine roots. *Tree Physiol* 5:53–61.
- del Hierro AM, W Kronberger, P Hietz, I Offenthaler, H Richter 2002 A new method to determine the oxygen concentration inside the sapwood of trees. *J Exp Bot* 53:559–563.

- Desprez-Loustau M, B Marçais, L Nageleisen, D Piou, A Vannini 2006 Interactive effects of drought and pathogens in forest trees. *Ann For Sci* 63:597–612.
- Dickson RE 1979 Analytical procedures for the sequential extraction of ^{14}C -labeled constituents from leaves, bark, and wood of cottonwood plants. *Physiol Plant* 45:480–488.
- Drew MC 1997 Oxygen deficiency and root metabolism: injury and acclimation under hypoxia and anoxia. *Annu Rev Plant Physiol Plant Mol Biol* 48:223–250.
- Ebel RC, JP Mattheis, DA Buchanan 1995 Drought stress of apple trees alters leaf emissions of volatile compounds. *Physiol Plant* 93:709–712.
- Eklund L, MB Lavigne 1995 Restricted lateral gas movement in *Pinus strobus* branches. *Trees* 10:83–85.
- Eklund L, HA Little 1998 Ethylene evolution, radial growth and carbohydrate concentrations in *Abies balsamea* shoots ringed with Ethrel. *Tree Physiol* 18:383–391.
- Felle HH 2005 pH regulation in anoxic plants. *Ann Bot* 96:519–532.
- Fox GG, NR McCallan, RG Ratcliffe 1995 Manipulating cytoplasmic pH under anoxia: a critical test of the role of pH in the switch from aerobic to anaerobic metabolism. *Planta* 195:324–330.
- Gansert D, M Burgdorf, R Lösch 2001 A novel approach to the in situ measurement of oxygen concentrations in the sapwood of woody plants. *Plant Cell Environ* 24:1055–1064.
- Gara RI, WR Littke, DF Rhoades 1993 Emission of ethanol and monoterpenes by fungal infected lodgepole pine trees. *Phytochemistry* 34:987–990.
- Gout E, A-M Boisson, S Aubert, R Douce, R Bligny 2001 Origin of the cytoplasmic pH changes during anaerobic stress in higher plant cells carbon-13 and phosphorous-31 nuclear magnetic resonance studies. *Plant Physiol* 125:912–925.
- Harry DE, CS Kinlaw, RR Sederoff 1988 The anaerobic stress response and its use for studying gene expression in conifers. Pages 275–290 in JW Hanover, DE Keathley, eds. *Genetic manipulation of woody plants*. Plenum, New York.
- Hook DD, CL Brown 1972 Permeability of the cambium to air in trees adapted to wet habitats. *Bot Gaz* 133:304–310.
- Houston DR 1987 Forest tree declines of past and present: current understanding. *Can J Plant Pathol* 9:349–360.
- Joseph G, RG Kelsey 1997 Ethanol synthesis and water relations of flooded *Pseudotsuga menziesii* (Mirb.) Franco (Douglas-fir) seedlings under controlled conditions. *Int J Plant Sci* 158:844–850.
- Joseph G, RG Kelsey, RW Peck, CG Niwa 2001 Response of some scolytids and their predators to ethanol and 4-allylanisole in pine forests of central Oregon. *J Chem Ecol* 27:697–715.
- Kelsey RG, G Joseph 1998 Ethanol in Douglas-fir with black-stain root disease (*Leptographium wageneri*). *Can J For Res* 28:1207–1212.
- 2001 Attraction of *Scolytus unispinosus* bark beetles to ethanol in water-stressed Douglas-fir branches. *For Ecol Manag* 144:229–238.
- 2003 Ethanol in ponderosa pine as an indicator of physiological injury from fire and its relationship to secondary beetles. *Can J For Res* 33:870–884.
- Kelsey RG, G Joseph, EA Gerson 1998 Ethanol synthesis, nitrogen, carbohydrates, and growth in tissues from nitrogen fertilized *Pseudotsuga menziesii* (Mirb.) Franco and *Pinus ponderosa* Dougl. ex Laws seedlings. *Trees* 13:103–111.
- Kelsey RG, DK Manter 2004 Effects of Swiss needle cast on Douglas-fir physiological parameters that contribute to bark beetle host selection and susceptibility. *For Ecol Manag* 190:241–253.
- Kimmerer TW, TT Kozlowski 1982 Ethylene, ethane, acetaldehyde, and ethanol production by plants under stress. *Plant Physiol* 69:840–847.
- Kimmerer TW, MA Stringer 1988 Alcohol dehydrogenase and ethanol in the stem of trees. *Plant Physiol* 87:693–697.
- Klimetzek E, J Köhler, JP Vité, U Kohnle 1986 Dosage response to ethanol mediates host selection by “secondary” bark beetles. *Naturwissenschaften* 73:270–272.
- Kolb TE, KM Holmberg, MR Wagner, JE Stone 1998 Regulation of ponderosa pine foliar physiology and insect resistance mechanisms by basal area treatments. *Tree Physiol* 18:375–381.
- Kreuzwieser J, FJM Harren, LJJ Laarhoven, I Boamfa, S te Lintel-Hekkert, U Scheerer, C Hüglin, R Rennenberg 2001 Acetaldehyde emission by the leaves of trees: correlation with physiological and environmental parameters. *Physiol Plant* 112:41–49.
- Kreuzwieser J, U Scheerer, H Rennenberg 1999 Metabolic origin of acetaldehyde emitted by poplar (*Populus tremula* $\times$ *P. alba*) trees. *J Exp Bot* 50:757–765.
- Kürsteiner O, I Dupuis, C Kuhlemeier 2003 The *pyruvate decarboxylase1* gene of *Arabidopsis* is required during anoxia but not other environmental stresses. *Plant Physiol* 132:968–978.
- Liu Y-B, JA Mclean 1989 Field evaluation of responses of *Gnathotrichus sulcatus* and *G. retusus* (Coleoptera: Scolytidae) to semiochemicals. *J Econ Entomol* 82:1687–1690.
- Lopushinsky W 1969 Stomatal closure in conifer seedlings in response to leaf moisture stress. *Bot Gaz* 130:258–263.
- MacDonald RC, TW Kimmerer 1989 Aerobic ethanol production by leaves: evidence for air pollution stress in trees of the Ohio River Valley, USA. *Environ Pollut* 62:337–351.
- 1991 Ethanol in the stems of trees. *Physiol Plant* 82:582–588.
- 1993 Metabolism of transpired ethanol by eastern cottonwood (*Populus deltoids* Bartr.). *Plant Physiol* 102:173–179.
- Mancuso S, AM Marras 2003 Different pathways of the oxygen supply in the sapwood of young *Olea europaea* trees. *Planta* 216:1028–1033.
- Mattson WJ, RA Haack 1987 The role of drought stress in provoking outbreaks of phytophagous insects. Pages 365–407 in P Barbosa, JC Schultz, eds. *Insect outbreaks*. Academic Press, San Diego, CA.
- Mellema S, W Eichenberger, A Rawyler, M Suter, M Tadege, C Kuhlemeier 2002 The ethanolic fermentation pathway supports respiration and lipid biosynthesis in tobacco pollen. *Plant J* 30:329–336.
- Miller DR 2006 Ethanol and (–)- α -pinene: attractant kairomones for some large wood-boring beetles in southeastern USA. *J Chem Ecol* 32:779–794.
- Minore D 1979 Comparative autecological characteristics of northwestern tree species: a literature review. USDA Forest Service General Technical Report PNW-87. Pacific Northwest Forest and Range Experiment Station, Portland, OR.
- Moeck H 1970 Ethanol as the primary attractant for the ambrosia beetle *Trypodendron lineatum* (Coleoptera: Scolytidae). *Can Entomol* 102:985–995.
- Ogigirigi MA, TT Kozlowski, S Sasaki 1970 Effect of soil moisture depletion on stem shrinkage and photosynthesis of tree seedlings. *Plant Soil* 32:33–49.
- Pharis RP 1966 Comparative drought resistance of five conifers and foliage moisture content as a viability index. *Ecology* 47:211–221.
- Phillips TW, AJ Wilkening, TH Atkinson, JL Nation, RC Wilkinson, JL Foltz 1988 Synergism of turpentine and ethanol as attractants for certain pine-infesting beetles (Coleoptera). *Environ Entomol* 17:456–462.
- Pinol J, A Sala 2000 Ecological implications of xylem cavitation for several pinaceae in the Pacific northern USA. *Funct Ecol* 17:538–545.
- Rieske LK, KF Raffa 1991 Effects of varying ethanol and turpentine levels on attraction of two pine root weevil species, *Hylobius pales* and *Pachylobius picivorus* (Coleoptera: Curculionidae). *Environ Entomol* 20:48–52.
- Roberts JKM, FH Andrade, IC Anderson 1985 Further evidence that cytoplasmic acidosis is a determinant of flooding intolerance in plants. *Plant Physiol* 77:492–494.

- Roberts JKM, J Callis, D Wemmer, V Walbot, O Jardetzky 1984 Mechanism of cytoplasmic pH regulation in hypoxic maize root tips and its role in survival under hypoxia. *Proc Natl Acad Sci USA* 81: 3379–3383.
- Sachs MM, M Freeling, R Okimoto 1980 The anaerobic proteins of maize. *Cell* 20:761–767.
- Schade GW, AH Goldstein 2001 Fluxes of oxygenated volatile organic compounds from a ponderosa pine plantation. *J Geophys Res* 106:3111–3123.
- 2002 Plant physiological influences on the fluxes of oxygenated volatile organic compounds from ponderosa pine. *J Geophys Res* 107:ACH 2, doi:10.1029/2001JD000532.
- Schroeder LM, Å Lindelöw 1989 Attraction of scolytids and associated beetles by different absolute amounts and proportions of α -pinene and ethanol. *J Chem Ecol* 15:807–817.
- Spicer R, NM Holbrook 2005 Within-stem oxygen concentration and sap flow in four temperate tree species: does long-lived xylem parenchyma experience hypoxia? *Plant Cell Environ* 28:192–201.
- Stoszek KJ 1973 A contribution to the biology of the *Pseudohylesinus nebulosus* (Leconte) (Coleoptera: Scolytidae), especially in relation to the moisture stress of its host, Douglas-fir. PhD diss. Oregon State University, Corvallis.
- Tadege M, R Brändle, C Kuhlemeier 1998 Anoxia tolerance in tobacco roots: effect of overexpression of pyruvate decarboxylase. *Plant J* 14:327–335.
- Tadege M, I Dupuis, C Kuhlemeier 1999 Ethanol fermentation: new functions for an old pathway. *Trends Plant Sci* 4:320–325.
- Tadege M, C Kuhlemeier 1997 Aerobic fermentation during tobacco pollen development. *Plant Mol Biol* 35:343–354.
- Tilles DA, K Sjödin, G Nordlander, HH Eidmann 1986 Synergism between ethanol and conifer host volatiles as attractants for the pine weevil, *Hylobius abietis* (L.) (Coleoptera: Curculionidae). *J Econ Entomol* 79:970–973.
- Valladares F, RW Pearcy 1997 Interactions between water stress, sun-shade acclimation, heat tolerance, and photoinhibition in the sclerophyll *Heteromeles arbutifolia*. *Plant Cell Environ* 20:25–36.
- Waring CL, NS Cobb 1992 The impact of plant stress on herbivore population dynamics. Pages 167–226 in E Bernays, ed. *Insect-plant interactions*. Vol 4. CRC, Boca Raton, FL.
- Xia J-H, JKM Roberts 1994 Improved cytoplasmic pH regulation, increased lactate efflux, and reduced cytoplasmic lactate levels are biochemical traits expressed in root tips of whole maize seedlings acclimated to a low-oxygen environment. *Plant Physiol* 105:651–657.

