

Interacting elevated CO₂ and tropospheric O₃ predisposes aspen (*Populus tremuloides* Michx.) to infection by rust (*Melampsora medusae* f. sp. *tremuloidae*)

D. F. KARNOSKY*, KEVIN E. PERCY†, BIXIA XIANG*, BRENDA CALLAN‡, ASKO NOORMETS*, BLANKA MANKOVSKA§, ANTHONY HOPKIN¶, JAAK SOBER*, WENDY JONES*, R. E. DICKSON** and J. G. ISEBRANDS**

*Michigan Technological University, 1400 Townsend Drive, Houghton, Michigan 49931 USA, †Natural Resources Canada, Canadian Forest Service-Atlantic Forestry Centre, PO Box 4000, Fredericton, NB E3B 5P7, Canada, ‡Natural Resources Canada, Canadian Forest Service-Pacific Forestry Centre, 506 West Burnside Road, Victoria, BC V8Z 1M5, Canada, §Forest Research Institute, T.G. Masarykova Street 2195, 960 92 Zvolen, Slovakia, ¶Natural Resources Canada, Canadian Forest Service-Great Lakes Forestry Centre, 1219 Queen Street, PO Box 490, Sault Ste. Marie, ON P6A 5M7, Canada, **USDA Forest Service, North Central Research Station, Forestry Sciences Laboratory, 5985 Highway K, Rhinelander, Wisconsin 54501 USA

Abstract

We investigated the interaction of elevated CO₂ and/or (Ozone) O₃ on the occurrence and severity of aspen leaf rust (*Melampsora medusae* Thuem. f. sp. *tremuloidae*) on trembling aspen (*Populus tremuloides* Michx.). Furthermore, we examined the role of changes in leaf surface properties induced by elevated CO₂ and/or O₃ in this host–pathogen interaction. Three- to five-fold increases in levels of rust infection index were found in 2 consecutive years following growing-season-long exposures with either O₃ alone or CO₂ + O₃ depending on aspen clone. Examination of leaf surface properties (wax appearance, wax amount, wax chemical composition, leaf surface and wettability) suggested significant effects by O₃ and CO₂ + O₃. We conclude that elevated O₃ is altering aspen leaf surfaces in such a way that it is likely predisposing the plants to increased infection by aspen leaf rust.

Keywords: epicuticular wax, FACE experiment, leaf rust, leaf surface properties

Received 15 February 2001; revised version received 1 October 2001 and accepted 3 November 2001

Introduction

Global atmospheric CO₂ concentration has risen by nearly 30% since preindustrial times (Barnola *et al.* 1995), largely because of the industrial emissions (Keeling *et al.* 1995). Similarly, background concentrations of tropospheric ozone (O₃) related to emissions of nitrogen oxides (NO_x) and volatile organic compounds (VOC) from fossil fuel, such as thermal generation and transportation, have increased from *circa*. 10 ppb to over 40 ppb (Finlayson-Pitts & Pitts 1997; Fowler *et al.* 1998; Stevenson *et al.* 1998). Fowler *et al.* (1999) suggest that nearly one-quarter of the earth's forests are currently at risk from tropospheric O₃ when July peak concentrations exceed 60 ppb.

Elevated CO₂ and O₃ affect trees through different mechanisms. With trembling aspen (*Populus tremuloides*

Michx.) elevated CO₂ stimulates photosynthesis (Tjoelker *et al.* 1998; Noormets *et al.* 2001), delays foliar senescence in autumn (Isebrands, personal communication), and stimulates above-ground (Isebrands *et al.* 2001) and below-ground (King *et al.* 2001) growth. Trees grown with elevated CO₂ generally have lower nitrogen concentrations in their foliage, lower Rubisco concentrations, altered defense compounds (Lindroth *et al.* 1993, 1997) and decreased concentration of antioxidants (Wustman *et al.* 2001).

In contrast to the largely beneficial effects of CO₂ on aspen, O₃ is generally detrimental to aspen growth and productivity. Ozone has been shown to induce foliar injury (Karnosky 1976), decrease foliar chlorophyll content (Gagnon *et al.* 1992), accelerate leaf senescence (Karnosky *et al.* 1996), decrease photosynthesis (Coleman *et al.* 1995a), alter carbon allocation (Coleman *et al.* 1995b), alter epicuticular wax structure and composition

Correspondence: D. F. Karnosky, e-mail Karnosky@mtu.edu

(Mankovska *et al.* 1998; Karnosky *et al.* 1999), and decrease growth (Wang *et al.* 1986; Karnosky *et al.* 1992, 1996, 1998; Coleman *et al.* 1996). Extrapolation of data on the impacts of O₃ on aspen in open-top chambers to aspen in nature suggests that 14–33% biomass loss may be occurring over 50% of aspen's natural range in the eastern US (Hogsett *et al.* 1997).

Current climate change scenarios predict further increases in global CO₂ concentrations (Stott *et al.* 2000) and area of global forests exposed to damaging levels of O₃ (increases to 49% or 17×10^6 km²) (Stevenson *et al.* 1998; Fowler *et al.* 1999). Predicting the potential impacts of plant diseases in this background of climate change is difficult as little is known about the occurrence and severity of disease under interacting CO₂ and O₃ (Chakraborty *et al.* 2000; Sandermann 2000). Thus, it is not possible at present to predict whether increased levels of these gases will lead to a lower or higher disease likelihood in particular plant-pathogen systems (Sandermann 2000). For example, very little is known about how foliar pathogens of forest trees will respond to elevated CO₂ and/or O₃ (Kickert & Krupa 1990).

Ozone predisposes some plants to infection by facultative leaf pathogens (Wukasch & Hofstra 1977; Rist & Lorbeer 1984). However, there is uncertainty about the role of O₃ in affecting infection by obligate pathogens such as rusts. Previous studies with poplar response to O₃ and foliar pathogens have shown both decreased (Coleman *et al.* 1987, 1988) and increased (Beare *et al.* 1999) disease levels with elevated O₃. In a short-term (5 h) greenhouse chamber study, Coleman *et al.* (1987) showed that an acute O₃ dose decreased the susceptibility of two eastern cottonwood (*Populus deltoides* Bartr.) clones to *Melampsora medusae*. However, Beare *et al.* (1999) found opposite response when a longer O₃ exposure period (17 d) increased severity of *Melampsora* rust on 'Balsam spire' hybrid poplar (*Populus trichocarpa* Torr. & Gray $\times$ *Populus balsamifera* L.). To the best of our knowledge, no one has examined the impact of O₃ on *Melampsora* rust occurrence on trembling aspen and furthermore, no one has examined the role of interacting elevated CO₂ on the host $\times$ O₃ $\times$ rust interaction in trees.

In this paper, we present observations on the occurrence and severity of *Melampsora medusae* Thuem. f. sp. *tremuloidae* Shain infection on three trembling aspen genotypes differing in O₃ tolerance and grown for multiple years under interacting elevated CO₂ and/or O₃ in a free-air enrichment (FACE) system. The objectives of the study were to determine (a) if O₃ increases or decreases the occurrence and severity of *M. medusae* infection and if this response varies by genotype and its O₃ susceptibility (b) if CO₂ ameliorates, aggravates or has no effect on O₃-induced responses, and (c) if O₃-induced changes to leaf surface physiochemical

characteristics and properties might predispose the trees to rust attack.

Materials and methods

The impacts of rapidly changing gaseous (CO₂, O₃) composition of the atmosphere on forest trees is being investigated at the Aspen FACE site located on the USDA Forest Service Harshaw Research Farm near Rhinelander, Wisconsin (Karnosky *et al.* 1999; Dickson *et al.* 2000). The experiment comprises a full factorial with twelve 30 m diameter treatment rings composed of 3 control rings, 3 rings with elevated O₃, 3 rings with elevated CO₂, and 3 rings with elevated O₃ + elevated CO₂. All rings are a minimum of 100 m apart. The rings were planted in mid summer 1997 and treatments ran from aspen budbreak to budset in 1998, 1999 and 2000. The eastern one-half of each ring was randomly planted with 3- to 6-month-old potted trees (20–40 cm in height) at 1 m $\times$ 1 m spacing in two tree plots of five aspen clones differing in O₃ tolerance (8 L, 216 and 271 = relatively tolerant and 42E and 259 = relatively sensitive). For this study of rust occurrence and foliar surface properties, we focused on the three clones (216, 259, and 271) for which we had extensive previous information on O₃ tolerance and pest susceptibility (Karnosky *et al.* 1992, 1996, 1998, 1999).

CO₂ and O₃ were released during the daylight hours. The targeted elevated CO₂ treatment concentration was 560 ppm, 200 ppm above ambient and O₃ exposure was at 1.5 $\times$ ambient. O₃ was not administered during periods of cold weather, or when leaf surfaces were wetted from fog, dew, or rain events. Complete details on the experimental design and pollutant generation and monitoring can be found in Karnosky *et al.* (1999) and Dickson *et al.* (2000).

Rust assessment

Melampsora medusae Thuem. f. sp. *tremuloidae* was identified by observations of urediniospores with scanning electron microscopy and/or light microscopy. The inoculum was natural with the primary source likely being adjacent stands of larch (*Larix* spp. and hybrids). Voucher specimens of rusted leaves, representing each of the five severity classes, were placed in the Pacific Forestry Centre's Forest Pathology Herbarium (DAVFP 25858) in Victoria, BC, Canada. *Melampsora* rust incidence was assessed on September 27–28, 1999 and on August 29–30, 2000. The percentage of infected leaves per tree for every tree in the core of each ring ($n = 1839$ trees total) was estimated and the severity of rust occurrence was scored on an average of 20 leaves per tree from 1 to 5 with 1 = 1–20% of the leaf area covered with urediniospores, 2 = 21–40%, 3 = 41–60%, 4 = 61–80%, and 5 = 81–100%. An index of mean

severity of infection was then calculated as percent leaves injured $\times$ severity of rust occurrence.

Leaf surface properties

Epicuticular waxes provide a physical barrier to protect leaves from pathogen infection (Mendgen 1996). It is well known that the appearance of these waxes can be seriously altered by air pollutants (Kerstiens 1996) and that these changes are dramatically manifested in leaves of aspen growing under elevated O₃ (Mankovska *et al.* 1998; Karnosky *et al.* 1999).

To characterize the impacts of O₃ on wax structure, we collected leaves (recently mature, LPI = 10–15 (Larson & Isebrands 1971)) in mid August of 1999 and 2000. Leaf segments (5 mm $\times$ 10 mm) were air-dried, gold-coated on a cold stage, and examined under a JSM 6400 (JEOL Ltd., Peabody, Massachusetts) scanning electron microscope (SEM). Five leaves per clone were examined for each treatment.

Epicuticular waxes were recovered by a chloroform rinse of five fully expanded (LPI 8–12) leaves pooled per wax sample. Five samples were collected randomly across the two-tree plantings per clone, per ring ($n = 15$ samples per treatment). The solvent/wax solution was filtered, solvent evaporated, and epicuticular wax weighed to ± 10 μ g and expressed as μ g cm⁻² leaf area. Quantitative wax chemical composition ($\pm 0.001\%$) was determined with a high-temperature capillary Varian 3410 (Varian Inc., Walnut Creek, California) gas chromatograph (Percy *et al.* 1994). Homologue assignments and peak integration were determined with Varian version 6.0 WorkStar software and injection of reference homologues and wax mixtures. Final confirmation of homologue assignments was determined with GC-MS.

Leaf surface wettability was assessed by measuring leaf surface-droplet contact angle (DCA) (Percy & Baker 1988) on five leaves per clone per treatment. Small volume (0.2 μ L; *circa.* 700 μ m dia.) distilled water droplets were placed on leaf surfaces as described by Percy & Baker (1988). Equilibrium angles were measured ($\pm 1^\circ$) using an NRL Contact Angle Goniometer (Ramé-Hart Inc., Mountain Lakes, New Jersey).

Data analysis

The experiment was designed as a randomized complete block design with four treatment rings in three replicate blocks. An experimental ring was regarded as an experimental unit for all statistical analyses and the mean parameter values for individual rings were used as input values.

The main and interactive effects of clone, year, and CO₂ and O₃ levels on the percent of infected leaves,

severity of infection and infection index were calculated with analysis of variance (ANOVA) using general linear models procedure (SAS, SAS Institute, Cary, NC, USA). The significance of clonal differences within a given treatment was calculated with Duncan's Multiple Range Test.

Epicuticular wax amount and alkane ratios [Σ C 29–C35(long)/ Σ C23–C28(short) homologues; homologue value expressed per unit leaf area (μ g cm⁻²)] were analysed using ANOVA (SAS 1993). Main effects tested were treatment, block, clone, and interactions. Differences between means were analysed using Duncan's multiple range test. The DCA data were analysed using ANOVA with an additional main effect leaf surface included.

Results and discussion

Treatment summary

Treatment rings with elevated CO₂ had an average concentration of 543 ± 75 ppm CO₂ in 1999 and 531 ± 86 ppm CO₂ in 2000 during daylight hours. Background daylight ambient CO₂ averaged 350 ± 10 ppm in 1999 and 347 ± 13 ppm in 2000. Daytime ambient O₃, O₃ treatment rings, and O₃ + CO₂ treatment rings averaged 36.9, 51.1 and 52.3 ppb O₃, in 1999, and 36.0, 47.4, and 48.9 ppb in 2000, respectively (Table 1, Fig. 1). Actual hourly average O₃ data for the 1999 and 2000 growing season are presented in Fig. 2.

The CO₂ concentrations we used (ambient plus 200 ppm) are predicted to occur globally within the next 40–60 years (Stott *et al.* 2000). The O₃ concentrations are similar to those characterized for several forested areas of the eastern United States (Pinkerton & Lefohn 1987). The O₃ values are similar in terms of the $1 \times$ O₃ seasonal dose for the $1 \times$ treatments (SUM 0 of 50–70 ppmh) described by Karnosky *et al.* (1996) which decreased stem biomass for the O₃ sensitive clone 259 but not for the more O₃ tolerant clones 216 and 271.

Table 1 Daytime (7 a.m.–7 p.m) ozone (O₃) average concentration (ppb) and SUM 0 (ppmh) and standard deviation for ambient air plots and for O₃ and O₃ + CO₂ treatments in the Aspen FACE project in 1999 and 2000

Year	Treatment	Daytime average (ppb)	SUM 0 (ppmh)
1999	Ambient	36.9 ± 6.5	62.8 ± 11.1
	O ₃ treatments	51.1 ± 17.1	87.9 ± 29.4
	CO ₂ + O ₃ treatments	52.3 ± 18.0	90.0 ± 31.9
2000	Ambient	36.0 ± 11.0	58.2 ± 17.7
	O ₃ treatments	47.4 ± 21.6	78.8 ± 35.9
	CO ₂ + O ₃ treatments	48.9 ± 22.2	80.9 ± 36.7




Fig. 1 The hourly profiles (mean hourly values over the growing season) for O₃ (grey) concentrations in O₃ (top) and CO₂ + O₃ treatments (bottom) and ambient rings (black) are shown from 1999 data. The 2000 profiles were similar.

Rust assessment

Significant differences in rust occurrence and severity occurred owing to clone and treatment (Table 2, Fig. 3). Seasonal exposures to O₃ and CO₂ + O₃ both resulted in significantly increased rust occurrence and severity (three- to five-fold) for all the three clones. Clonal differences in rust susceptibility were greater in 1999 than 2000, although some of these differences may be in part because of the earlier sampling time of 2000. Some of the more severely rusted leaves might have already fallen from the trees by late September 1999. Weather conditions varied considerably for the 2 years as 2000 was on average cooler and wetter than 1999 (Heilman, unpublished). The O₃ sensitive clone (259) was consistently the most susceptible to rust occurrence though it was not always significantly different from the other two clones.

Our results support the findings of Beare *et al.* (1999) that O₃ exacerbates the extent of *Melampsora* rust infection in *Populus* and is counter to the results of Coleman *et al.* (1987) who found that *Melampsora* rust infection was decreased in the presence of O₃. The desperate findings could be because of the different *Populus* species used in the three experiments or to the differences in O₃ exposure durations and techniques. Ozone is known to predispose some host plants to pathogens (Manning & Tiedemann 1995; Sandermann 1996). Coleman *et al.* (1987) noted the impact of environment on genotypic reactions and the potential impact on disease resistance. Our data suggest that the more rust-resistant aspen clones show increased

Table 2 Analysis of variance for the effects of clone, year, CO₂, O₃ and CO₂ + O₃ on the occurrence and severity of *Melampsora medusae* rust on *Populus tremuloides*

Source of variation	df	%L ¹	Severity ²	Index ³
Clone	2	0.0001	0.0026	0.0003
Year	1	0.0056	n.s.	n.s.
Clone × Year	2	0.0002	0.0049	0.0008
CO ₂	1	0.0035	n.s.	0.0153
Clone × CO ₂	2	0.0131	n.s.	0.0453
Year × CO ₂	1	n.s.	n.s.	n.s.
Clone × Year × CO ₂	2	n.s.	n.s.	n.s.
O ₃	1	0.0001	0.0001	0.0001
Clone × O ₃	2	0.0094	n.s.	0.0001
Year × O ₃	1	0.0194	n.s.	n.s.
Clone × Year × O ₃	2	0.0037	n.s.	0.0055
CO ₂ × O ₃	1	0.0007	0.0500	0.0022
Clone × CO ₂ × O ₃	2	n.s.	n.s.	n.s.
Year × CO ₂ × O ₃	1	0.0032	n.s.	0.0334
Clone × Year × CO ₂ × O ₃	2	n.s.	n.s.	n.s.

¹Percent leaves with *Melampsora medusae* uridinospores

²Severity of *Melampsora medusae* infection (see Fig. 3 for details)

³Index comprised of average percent leaves infected × average severity

levels of infection after extended exposure to O₃, suggesting suppression of host resistance after prolonged exposure to O₃. Coleman *et al.* (1987) only tested very short-term O₃ exposures (5 h) whereas the Beare *et al.* (1999) O₃




Fig. 2 The hourly average O₃ concentrations in the O₃ treatments (grey) and ambient rings (dark) are shown for 1999 and 2000.

exposures were over multiple weeks and our study involved seasonal exposures. In two of the few CO₂/O₃/pathogen interaction studies reported, Tiedemann & Firsching (1998, 2000) found that O₃ decreased the occurrence of *Puccinia recondita* rust on spring wheat while elevated CO₂ had no effect on rust occurrence. Because of the conflicting results, predicting interactions among trees and foliar pathogens under rapidly increasing

atmospheric CO₂ and/or O₃ concentrations will be difficult at best (Kickert & Krupa 1990; Chakraborty *et al.* 2000; Seem *et al.* 2000).

Leaf surface physicochemical characteristics

Ozone had a large effect on the structure of wax deposits formed on aspen leaves in our study. Leaves from trees




Fig. 3 *Melampsora medusae* occurrence (percent of leaves infected), severity of infection (1 = 1–20% of the leaf area covered with urediniospores, 2 = 21–40%, 3 = 41–60%, 4 = 61–80%, and 5 = 81–100%), and infection index (percent leaves infected × severity) for aspen clones 216, 259 and 271 exposed to growing season – long conditions of elevated CO₂ (ambient + 200 ppm), O₃ (1.5 × ambient), or CO₂ (ambient + 200 ppm) + O₃ (1.5 × ambient) at the Aspen FACE site in Rhinelander, Wisconsin during 1999 (top) and 2000 (bottom panels). All values are means ± SE.

growing with elevated O₃ had an increased proportion of amorphous; flat deposits (Fig. 4) crystallized over a dense array of plate crystallites (Fig. 4) normally observed distributed between pronounced cuticular ridges on control leaves.

Allen (1991) demonstrated the key role of topographical signals from the leaf surface in the development of appressoria from urediniospores of *M. medusae* (unspecified f. sp. but probably *deltoidae*) collected from *P. deltoides*. Optimal ridge heights preferred for




Fig. 4 Ozone impacts on the adaxial leaf surface waxes are shown here in a comparison of scanning electron micrographs of leaves from aspen clone 259 grown under control (left) and elevated O₃ (right) in the FACTS II (Aspen FACE) project. The visible changes of wax structure from crystallite (left) to amorphous forms (right) have been confirmed by extensive SEM characterization of hundreds of leaves (Mankovska *et al.* 1998; Karnosky *et al.* 1999) and by subsequent wax chemical composition analysis data (see Fig. 6).




Fig. 5 Epicuticular wax amounts recovered from three aspen clones exposed to control, elevated CO₂, elevated O₃ or elevated CO₂ + O₃ over the 1999 growing season in the Aspen FACE project near Rhinelander, Wisconsin. Data are means $\pm$ SE.

appressorium formation (a prerequisite for leaf penetration by the pathogen) are quite low (0.4–0.8 μm). This morphology would likely be similar to the flattened amorphous epicuticular wax deposits we see on aspen leaves exposed to O₃ (Fig. 4b) and as described by Mankovska *et al.* (1998) and Karnosky *et al.* (1999). The O₃-induced topography is in marked contrast to the large ridges seen in waxes of control aspen leaves (Fig. 4a).

To be sure that the structural changes we saw were not induced during specimen preparation or SEM examination, we examined the production and chemical composition of aspen leaf surface waxes following seasonal O₃ and/or CO₂ exposures. Averaged across all the four treatments in our experiment, the O₃ sensitive clone (259) had significantly greater ($P < 0.0001$) adaxial/abaxial wax deposits (45.7 $\mu\text{g cm}^{-2}$) compared to the O₃ tolerant

clones 216 (42.8 $\mu\text{g cm}^{-2}$) and 271 (36.6 $\mu\text{g cm}^{-2}$). Ozone and CO₂ + O₃ treatments resulted in changes in wax production in two of the three clones (Fig. 5). Wax amount on O₃ treated (42.8 $\mu\text{g cm}^{-2}$) and CO₂ + O₃-treated (34.9 $\mu\text{g cm}^{-2}$) leaves of clone 271 were significantly greater ($P < 0.001$) than on control (29.1 $\mu\text{g cm}^{-2}$) leaves. In the O₃ sensitive clone 259, wax amount on O₃-treated (47.1 $\mu\text{g cm}^{-2}$) and CO₂ + O₃-treated (47.6 $\mu\text{g cm}^{-2}$) leaves was significantly greater ($P < 0.001$) than on control (38.4 $\mu\text{g cm}^{-2}$) leaves (Fig. 5). Increased wax amount implies an O₃-mediated stimulation of *de novo* synthesis, which may be a protective adaptation similar to the generalized wounding mechanism of plant response to O₃ (Heath 1999).

Significant ($P < 0.01$) treatment effects on epicuticular wax chemical composition were detected for two (216 and 259) of the three clones. Along with increases in




Fig. 6 Alkane carbon chain ratio [$(\Sigma C29-C35/\Sigma C23-C28)$ homologues; expressed per unit leaf area ($\mu\text{g cm}^{-2}$)] averaged across three aspen clones exposed to control, elevated CO_2 , elevated O_3 , elevated $\text{CO}_2 + \text{O}_3$ during the 1999 growing season in the Aspen FACE project near Rhinelander, Wisconsin. Data are means ± 1 SE.

proportions of wax classes crystallizing largely into amorphous forms (alkyl esters, fatty acids), there were proportional decreases in classes crystallizing as plates (alkanols, alkanes). Early work by Holloway (1969) showed that alkanes were most hydrophobic, marginally more so than esters and considerably more so than fatty acids. Changes in wax class composition therefore would be expected to result in changes to leaf wettability, possibly leading to a biological effect.

Ozone additionally induced significant changes in wax biosynthesis in the homologue carbon chains length. For instance, the ratio of long to short alkane carbon chains was significantly increased ($P < 0.05$) because of the O_3 in clone 259 (Fig. 6). More of the carbon allocated to the elongase-decarboxylase pathway was metabolized into the longer ($> C28$) chains, possibly contributing to the observed subtle alteration in leaf surface properties.

Leaf surface wettability

The key role of cuticle wettability in fungal pathogen spore survival has been pointed out by Kerstiens (1996). It is also known that wettability of tree foliar cuticles can be altered by O_3 (Percy *et al.* 1994). Contact angles on both adaxial and abaxial surfaces were affected by O_3 . With particular reference to *Melampsora medusae* urediniospores which germinate on aspen abaxial leaf surfaces, exposure to O_3 decreased ($P < 0.0001$) the contact angle


(leaves were more wettable) in the two O_3 tolerant clones but not in the O_3 sensitive clone 259 (Fig. 7). Interestingly, cofumigation with CO_2 ameliorated the negative effects of O_3 to different degrees on contact angle in the two tolerant clones.

Conclusions

Our results clearly show that exposure to O_3 alters leaf surface integrity. Leaf surface topography, microroughness and physicochemical characteristics such as chemical composition and epicuticular wax structure (conferred by its chemical composition) all combine to determine leaf surface properties such as wettability, retention of solutes, and foliar uptake among others. The microenvironment thus produced at the phylloplane is a critical determinant in the process of fungal attachment and infection (Kerstiens 1996; Mendgen 1996).

We believe that the O_3 -induced changes to leaf surface physicochemical characteristics and wettability will significantly increase incidence of *Melampsora medusae* as documented here and these changes may have deleterious short-term impacts on carbon fixation in aspen and influence long-term competitive ability of the different clones. Further experiments, such as the artificial inoculation of aspen leaf surfaces having altered physicochemical characteristics produced under controlled O_3 exposure are suggested to test this hypothesis.

Fig. 7 Droplet contact angle (DCA) of the abaxial leaves of three aspen clones exposed to control, elevated CO₂, elevated O₃, elevated CO₂ + O₃ over the 1999 growing season in the Aspen FACE project near Rhinelander, Wisconsin. Data are means $\pm$ SE.



Acknowledgements

This research was partially supported by the US Department of Energy, Office of Biological Research (DE-FG02-95ER6215), the USDA Forest Service Northern Global Change Program, the National Science Foundation (DBI-9601942; IBN-9652675), the National Council of the Paper Industry for Air and Stream Improvement (NCASI), Michigan Technological University, the Slovakian Forest Research Institute, the Proxair Foundation, the USDA NRI program, the Canadian Forest Service, and the USDA Forest Service.

References

- Allen EA (1991) Appressorium formation in response to topographical signals by 27 rust species. *Phytopathology*, **81**, 323–331.
- Barnola JM, Anklin M, Porheron J, Raynaud D, Schwander J, Stauffer BTI (1995) CO₂ evolution during the last millennium as recorded by Antarctic and Greenland ice. *Tellus B*, **47**, 264–272.
- Beare JAA, Archer SA, Bell JNB (1999) Effects of *Melampsora* leaf rust disease and chronic ozone exposure on poplar. *Environmental Pollution*, **105**, 419–426.
- Chakraborty S, Tiedemann AV, Teng PS (2000) Climate change: potential impact on plant diseases. *Environmental Pollution*, **108**, 317–326.
- Coleman MD, Dickson RE, Isebrands JG, Karnosky DF (1995a) Photosynthetic productivity of aspen clones varying in sensitivity to tropospheric ozone. *Tree Physiology*, **15**, 585–592.
- Coleman MD, Dickson RE, Isebrands JG, Karnosky DF (1995b) Carbon allocation and partitioning in aspen clones varying in sensitivity to tropospheric ozone. *Tree Physiology*, **15**, 593–604.
- Coleman MD, Dickson RE, Isebrands JG, Karnosky DF (1996) Root growth and physiology of potted and field-grown trembling aspen exposed to tropospheric ozone. *Tree Physiology*, **16**, 145–152.
- Coleman JS, Jones CG, Smith WH (1987) The effect of ozone on cottonwood–leaf rust interactions: independence of abiotic stress, genotype and leaf ontogeny. *Canadian Journal of Botany*, **65**, 949–953.
- Coleman JS, Jones CG, Smith WH (1988) Interactions between an acute ozone dose, eastern cottonwood and Marssonina leaf spot: implications for pathogen community dynamics. *Canadian Journal of Forest Research*, **66**, 863–868.
- Dickson RE, Lewin KF, Isebrands JG *et al.* (2000) Forest atmosphere carbon transfer and storage-II (FACTS-II). The aspen free-air CO₂ and O₃ enrichment (FACE) project: An Overview. USDA. Forest Service, North Central Research Station General Technical Report, NC-214, St. Paul, MN, 68 pp.
- Finlayson-Pitts BJ, Pitts JN Jr (1997) Tropospheric air pollution: ozone, airborne toxics, polycyclic aromatic hydrocarbons, and particulates. *Science*, **276**, 1045–1051.
- Fowler D, Cape JN, Coyle M *et al.* (1999) The global exposure of forests to air pollutants. *Water, Air, and Soil Pollution*, **116**, 5–32.
- Fowler D, Flechard C, Skiba U, Coyle M, Cape JN (1998) The atmospheric budget of oxidized nitrogen and its role in ozone formation and deposition. *New Phytologist*, **139**, 11–23.
- Gagnon ZE, Karnosky DF, Dickson RE, Isebrands JG (1992) Effects of ozone on chlorophyll content in *Populus tremuloides*. *American Journal of Botany*, **79** (Suppl.), 107.
- Heath RL (1999) Biochemical processes in an ecosystem: How should they be measured? *Water, Air and Soil Pollution*, **116**, 279–298.
- Hogsett WE, Weber JE, Tingey D, Herstrom A, Lee EH, Laurence JA (1997) An approach for characterizing tropospheric ozone risk to forests. *Environmental Management*, **21**, 105–120.
- Holloway PJ (1969) Chemistry of leaf waxes in relation to wetting. *Journal of the Science of Food and Agriculture*, **20**, 124–128.
- Isebrands JG, McDonald EP, Kruger E, Hendrey G, Pregitzer K, Percy K, Sober J, Karnosky DF (2001) Growth responses of *Populus tremuloides* clones to interacting carbon dioxide and tropospheric ozone. *Environmental Pollution*, **115**, 359–371.
- Karnosky DF (1976) Threshold levels for foliar injury to *Populus tremuloides* Michx. by sulfur dioxide and ozone. *Canadian Journal of Forest Research*, **6**, 166–169.

- Karnosky DF, Gagnon ZE, Dickson RE, Coleman MD, Lee EH, Isebrands JG (1996) Changes in growth, leaf abscission, and biomass associated with seasonal tropospheric ozone exposures of *Populus tremuloides* clones and seedlings. *Canadian Journal of Forest Research*, **16**, 23–27.
- Karnosky DF, Gagnon ZE, Reed DD, Witter JA (1992) Growth and biomass allocation of symptomatic and asymptomatic *Populus tremuloides* clones in response to seasonal ozone exposures. *Canadian Journal of Forest Research*, **22**, 1785–1788.
- Karnosky DF, Percy KE, Mankovska B, Dickson RE, Isebrands JG (1999) Influence of ambient O₃ on the growth, competitive ability, leaf surface properties, and pest tolerance of trembling aspen genotypes. *Water, Air and Soil Pollution*, **116**, 311–322.
- Karnosky DF, Podila GK, Gagnon Z *et al.* (1998) Genetic control of responses to interacting tropospheric ozone and CO₂ in *Populus tremuloides*. *Chemosphere*, **36**, 807–812.
- Keeling CM, Whort TP, Wahlen M, Vander Pliet J (1995) International extremes in the rate of rise of atmospheric carbon dioxide since 1980. *Nature*, **375**, 666–670.
- Kerstiens G (1996) Barrier properties of the cuticle to water, solutes and pest and pathogen penetration in leaves of plants grown in polluted atmospheres. In: *Plant Response to Air Pollution* (Yunus M, Iqbal M eds). John Wiley and Sons Ltd, New York, pp. 167–178.
- Kickert RN, Krupa SV (1990) Forest responses to tropospheric ozone and global climate change: An analysis. *Environmental Pollution*, **68**, 29–65.
- King JS, Pregitzer KS, Zak DR *et al.* (2001) Fine root biomass and fluxes of soil carbon in young stands of paper birch and trembling aspen is affected by elevated CO₂ and tropospheric O₃. *Oecologia*, **128**, 237–250.
- Larson PR, Isebrands JG (1971) The plastochron index as applied to developmental studies of cottonwood. *Canadian Journal of Forest Research*, **1**, 1–11.
- Lindroth RL, Kinney KK, Platz CL (1993) Responses of deciduous trees to elevated atmospheric CO₂: Productivity, phytochemistry, and insect performance. *Ecology*, **74**, 763–777.
- Lindroth RL, Roth S, Kruger EL, Volin JC, Koss PA (1997) CO₂-mediated changes in aspen chemistry: effects on gypsy moth performance and susceptibility to virus. *Global Change Biology*, **3**, 279–289.
- Mankovska B, Percy K, Karnosky DF (1998) Impact of ambient tropospheric O₃, CO₂, and particulates on the epicuticular waxes of aspen clones differing in O₃ tolerance. *Ekológia*, **18**, 200–210.
- Manning WJ, Tiedemann AV (1995) Climate change: potential effects of increased atmospheric carbon dioxide (CO₂), ozone (O₃), and ultraviolet-B (UV-B) radiation on plant diseases. *Environmental Pollution*, **88**, 219–245.
- Mendgen K (1996) Fungal attachment and penetration. In: *Plant Cuticles* (Kerstiens G ed.). Bios. Scientific Publishers Ltd, Oxford, pp. 175–188.
- Noormets A, Sober A, Pell EJ *et al.* (2000) Stomatal and non-stomatal control of photosynthesis in trembling aspen (*Populus tremuloides* Michx.) exposed to elevated CO₂ and/or O₃. *Plant, Cell and Environment*, **24**, 327–336.
- Percy KE, Baker EA (1988) Effect of simulated acid rain on leaf wettability, rain retention and uptake of some inorganic ions. *New Phytologist*, **108**, 75–82.
- Percy KE, McQuattie CJ, Rebbeck JA (1994) Effects of air pollutants on epicuticular wax chemical composition. In: *Air Pollutants and the Leaf Cuticle* (Percy, KE, Cape, JN, Jagels, R, Simpson, CM, eds) NATO ASI Series, Vol. G. 36. Springer-Verlag, Heidelberg, pp. 67–79.
- Pinkerton JE, Lefohn AE (1987) The characterization of ozone data for sites located in forested areas of the eastern United States. *Journal of the Air Pollution Control Association*, **37**, 1005–1010.
- Rist DL, Lorbeer JW (1984) Ozone-enhanced leaching of onion leaves in relation to lesion production by *Botrytis cinerea*. *Phytopathology*, **74**, 1217–1220.
- Sandermann H Jr. (1996) Ozone and plant health. *Annual Review of Phytopathology*, **34**, 347–366.
- Sandermann H Jr. (2000) Ozone/biotic disease interactions: molecular biomarkers as a new experimental tool. *Environmental Pollution*, **108**, 327–332.
- Seem RC, Margarey RD, Zack JW, Russo JM (2000) Estimating disease risk at the whole plant level with General Circulation Models. *Environmental Pollution*, **108**, 389–395.
- Stevenson DS, Johnson CE, Collins WJ, Derwent RG, Shine KP, Edwards JM (1998) Evolution of tropospheric ozone radiative forcing. *Geophysical Research Letters*, **25**, 3819–3822.
- Stott PA, Tett SFB, Jones GS, Allen MR, Mitchell JFB, Jenkins GJ (2000) External control of 20th century temperature by natural and anthropogenic forcing. *Science*, **290**, 2133–2137.
- Tiedemann AV, Firsching KH (1998) Combined whole-season effects of elevated ozone and carbon dioxide concentrations on a simulated wheat leaf rust (*Puccinia recondita* f. sp. *tritici*) epidemic. *Journal of Plant Disease Protection*, **105**, 555–566.
- Tiedemann AV, Firsching KH (2000) Interactive effects of elevated ozone and carbon dioxide on growth and yield of leaf rust-infected versus non-infected wheat. *Environmental Pollution*, **108**, 357–363.
- Tjoelker MG, Oleksyn J, Reich PB (1998) Seedlings of five boreal tree species differ in acclimation of net photosynthesis to elevated CO₂ and temperature. *Tree Physiology*, **18**, 715–726.
- Wang D, Karnosky DF, Bormann FH (1986) Effects of ambient ozone on the productivity of *Populus tremuloides* Michx. grown under field conditions. *Canadian Journal of Forest Research*, **16**, 47–55.
- Wukasch RT, Hofstra G (1977) Ozone and *Botrytis* spp. interactions in onion-leaf dieback: open-top chambers. *Phytopathology*, **67**, 1080–1084.
- Wustman BA, Oksanen E, Karnosky DF, Sober J, Isebrands GR, Hendrey GR, Pregitzer KS, Padila GK (2001) Effects of elevated CO₂ and O₃ on aspen clones vary in O₃ sensitivity: can CO₂ ameliorate the harmful effects of O₃? *Environmental Pollution*, **115**, 473–481.