

United States
Department of
Agriculture

Forest Service



Volume 30 • Issue 1



Forest Nursery Notes

Winter 2010







**Please send address changes to
Rae Watson. You may use the
Literature Order Form at the end of the
New Nursery Literature section.**

*This international technology transfer service
is produced by the
US. Department of Agriculture, Forest Service,
Natural Resources Conservation Service,
National Agroforestry Center (Lincoln, Nebraska),
with funding from the Forest Service,
State and Private Forestry,
through the Center for Reforestation,
Nurseries, and Genetics Resources.*

Forest Nursery Notes Team

R. Kasten Dumroese, Editor-In-Chief
USDA Forest Service
Rocky Mountain Research Station
1221 S. Main Street
Moscow, ID 83843-4211
TEL: 208.883.2324
FAX: 208.883.2318
[E-Mail: kdumroese@fs.fed.us](mailto:kdumroese@fs.fed.us)

Tom D. Landis, Lead Author & Editor
Forest Nursery Consultant
3248 Sycamore Way
Medford, OR 97504-9005
TEL: 541.210.8108
FAX: 541.858.6110
[E-Mail: nurseries@aol.com](mailto:nurseries@aol.com)

Diane L. Haase, Author
USDA Forest Service
PO Box 3623
Portland, OR 97208
TEL: 503.808.2349
FAX: 503.808.2339
[E-Mail: dlhaase@fs.fed.us](mailto:dlhaase@fs.fed.us)

Rae Watson, Layout and Author
USDA Forest Service
2606 Old Stage Road
Central Point, OR 97502
TEL: 541.858.6131
FAX: 541.858.6110
[E-Mail: rewatson@fs.fed.us](mailto:rewatson@fs.fed.us)

Laura Hutchinson, Library Services
USDA Forest Service
North Central Research Station
1992 Folwell Avenue
St. Paul, MN 55108
TEL: 651.649.5272
[E-Mail: lhutchinson@fs.fed.us](mailto:lhutchinson@fs.fed.us)

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202)720-6382 (TDD). USDA is an equal opportunity provider and employer.

Nursery Meetings

16th Wildland Shrub Symposium

This symposium will be held May 26 to 27, 2010 at Utah State University in Logan, Utah. Papers on Climate Change, Wildlife, Energy Extraction, Invasive Species, Restoration, Wildfire, Recreation, Livestock Grazing, Social and Economic Aspects, and Shrub Biology are encouraged.

For more information contact:

Tom Monaco

[E-Mail: tom.monaco@ars.usda.gov](mailto:tom.monaco@ars.usda.gov)

Or

Eugene Schupp

[E-Mail: Eugene.schupp@usu.edu](mailto:Eugene.schupp@usu.edu)

Forest Nurseries in the Natural State: Biennial meeting of the Southern Forest Nursery Association

This meeting will be held on July 26-29, 2010 in Little Rock, Arkansas.

Early registration will be \$300 and late registration \$400. Lodging is available at The Peabody Little Rock with single rooms at \$88 and doubles at \$128 plus tax. A full registration packet will be sent out in April.

Program and exhibitor information is available by contacting Allan Murray at 501-907-2486 or allan.murray@arkansas.gov

Registration services will be provided by Western Forestry and Conservation Association, 503-226-4562 or www.westernforestry.org



Target Seedling Symposium 2010

August 24-26, 2010

Sheraton Airport Hotel, Portland, Oregon

Joint meeting of:

**Forest Nursery Association of British Columbia
and Western Forest and Conservation Nursery Association**



August 24-25

Target Plant Characteristics

August 24

Choice of two field trips

Blooming Nursery: wholesale nursery with Oregon's largest solar thermal energy installation to circulate warm water and heat a 54,000 sq. ft greenhouse.

OR

Conifer Seedling Production: Visit both a container nursery producing a wide array of conifer seedlings primarily for reforestation and a bareroot nursery providing transplants for reforestation projects.

August 25

Hands-on sessions

August 26, 2010

Current and Emerging Technologies

Please contact:

Richard Zabel

Western Forestry and Conservation Association
503-226-4562

or mail to:

Richard@westernforestry.org

*Registration information, lodging information,
and the full event schedule will be available in April at*

www.westernforestry.org

Western Region of the International Plant Propagators Society

This year's meeting will be held in Bellingham, Washington on September 8 - 11, 2010

The agenda is still being developed but will include a wide variety of presentations on all aspects of plant propagation.

For more information go to the following website:

<http://www.ipps.org>

The Fifth Western Native Plants Conference

This conference will be held on December 7-9, 2010 in Portland, Oregon. This event happens every 3 years and covers challenges and strategies for propagation and restoration of native plants.

A variety of experts will speak about many topics such as invasives, genetics, climate change, monitoring, etc. The first day will consist of an optional all-day field tour of nurseries and restoration projects in the area.

The agenda and more information and registration will be posted at www.westernforestry.org or contact Diane for further information:

Diane L. Haase

USDA Forest Service, Cooperative Programs

PO Box 3623 - 333 SW First Avenue

Portland, OR 97208-3623

Phone: 503.808.2349

Fax: 503.808.2339

[E-Mail: dlhaase@fs.fed.us](mailto:dlhaase@fs.fed.us)



Fertigation - Injecting soluble fertilizers into the irrigation system: Part 2

by Thomas D. Landis, Jeremy R. Pinto, and Anthony S. Davis

The first part of this article in the Summer 2009 issue covered basic mineral nutrition, the 3 components of a fertigation system, and the chemical calculations for formulating your own custom fertigation solutions. In this second and final part, we'll discuss types of fertilizer injectors, fertigation scheduling, and how to check injector function and determine exactly how much liquid fertilizer is going on your crop.

The simplest way to fertigate is to mix a large batch of applied strength solution and just spray it directly on your crops. Some bareroot nurseries fertigate their beds by spraying an applied strength fertilizer solution through a tractor-drawn sprayer (Triebwasser and Altsuler 1995).

Because of the sheer volume of fertigation solution and higher labor costs for mixing and application, this method is only practical in smaller container facilities.

One obvious benefit of applying a diluted fertilizer solution is that there is no risk of fertilizer burn.

Types of Injectors

Most fertigation systems use some type of mechanical injector to mix small volumes of concentrated fertilizer solutions into the irrigation water; a wide variety of injectors are available (Table 1). The best and most current information on fertilizer injectors can be found on-line (Kessler and Pennisi 2004; Pennisi and Kessler 2003), and be sure to check the manufacturers websites for the latest information. For example, recent publications mention one injector, the Gewa, but that company has recently gone out of business and only replacement parts are now available.


The simplest and least expensive fertilizer injectors are the venturi types, such as the Hozon (Figure 1A),

Table 1 – Technical specifications for common fertilizer injectors

Brand name	Multiple heads	Injection ratios	Water Flow rate (gpm)	Water supply pipe diameter (in.)	Approx. cost (2010)	Acid injection possible	Remarks
Venturi							
Hozon	No	1:12 to 1:16	Any	0.75	\$30	No	Requires 35 psi water pressure
E-Z Flo	No	Varies with model: 1:400 to 1:1,000	Any	Varies with model: 0.75 to 3.00	\$50 to \$750	No	www.ezfloinjection.com/
Positive displacement hydraulic pump							
Smith MeasureMix	Yes	Varies with model: 1:100 or 1:200	Varies with model: 3 to 200	Varies with model: 0.75 to 6.00	\$1,600 to \$4,600	Yes	www.smithpumps.com
DosMatic	No	1:50 to 1:200	Varies with Model: 5 to 20	0.75	\$500 to \$700	Yes	www.DosMatic.com
Dosatron	No	1:50 to 1:500	Varies with model: 0.5 to 100	Varies with model: 0.75 to 2.00	\$500 to \$2,000	Yes	www.dosatronicsusa.com
Flow-metered hydraulic or electric pump							
Anderson	Yes	Adjustable: 1:100 to 1:20,000	Varies with model: 0.75 to 300	Varies with model: 0.75 to 10.00	\$800 to \$5,000	Yes	www.heanderson.com

which continuously injects stock solution at an approximate 1:16 ratio. As water passes through the Hozon, it creates a negative pressure that sucks the fertilizer solution from the stock tank (Figure 1B). One limitation is that a water pressure of at least 35 psi (pounds per square inch) is needed to create sufficient suction. Siphon injectors can be used to apply other water-soluble chemicals, such as insecticides and fungicides, but cannot be used to inject acids (Pennisi and Kessler 2003). More sophisticated injectors, such as the Smith

Measuremix* (Figure 1C), feature a water motor that injects stock solution at a specified ratio. For example, an injector with a 1:100 ratio injects one part fertilizer stock solution for every 100 parts of irrigation water (Figure 1D). Many of these injectors have separate heads to inject two or more solutions and some models have plastic parts that are compatible with acid injection. Again, check the web publications and manufacturer website for specifications (Table 1).



Several things should be considered before purchasing an injector (Kessler and Pennisi 2004; Pennisi and Kessler 2003; Weiler and Sailus 1996):

Size and complexity of your nursery - Small nurseries growing a few species with hand watering or with an irrigation system with only a couple of zones can get by with a simple and inexpensive injector such as the Hozon or E-Z Flo. However, as the number of crops and the area to be fertigated increases, more sophisticated injectors are required. If you haven't done so already, it's a good idea to separate your different crops into nutrient requirement zones such as low, medium, and high. Native plant crops vary considerably in their response to fertilization, especially nitrogen, so grouping species by fertility zones makes fertigation much easier and more efficient.

Water flow rate - Because injectors supply a proportionate amount of liquid fertilizer to a given amount of water, you must know how much water your irrigation system can supply per unit of time. Flow rates can be divided into three categories based on gallons per minute (gpm): low (0.05 to 12 gpm), medium (12 to 40 gpm) or high (> 40 gpm). If you don't know your water flow rate, there are a couple of ways to find out. The simplest is to turn your irrigation on full, and measure how long it takes to fill a container or tank of known volume. Dividing the volume in gallons by the time in minutes gives you gpm.

If your nursery has a permanent irrigation system, then you hopefully have an in-line water meter that measures total volume usage; if not, we'd recommend getting one installed. Knowing your irrigation flow rates is essential to effective fertigation; writing down the starting and ending water usage along with the amount of stock solution consumed in a daily log book is an easy and effective way to confirm the actual injection ratio and check if the injector is working properly.

Injection ratio - This is simply the ratio of the amount of fertilizer injected per volume of irrigation water and most fertilizer injectors can be ordered with a wide variety of injection ratios (Table 1). Most injectors have a fixed injection ratio and the most common are 1:100 or 1:200, but some brands feature adjustable injection ratios. In-line venturi injectors used in hand watering have relatively low injector ratios. For automated irrigation systems, injectors with ratios less than 1:100 aren't practical because a very large fertigation tank would be required. On the other hand, using injectors with ratios >1:200 means that the fertilizer solution

must be very concentrated, which leads to insolubility problems.

Multiple injector heads - Simpler injectors such as the Hozon can handle only one fertilizer stock solution at a time, but many fertilizer injectors can be ordered with two or more injection ports, or heads (Table 1). Commercial brands of soluble fertilizer can be mixed in a single stock solution tank so an injector with one head is adequate. However, when injecting acids to correct high water pH or when formulating custom fertigation solutions from stock chemicals, separate injector heads are necessary (Figure 2A). For example, calcium and sulfate cannot be mixed in the same stock solution tank because they form an insoluble precipitate (gypsum) that can plug up the injector or irrigation nozzles (Figure 2B). A list of incompatible fertilizer chemicals can be found in Landis and others (1989).

Water quality - The amount of dissolved chemicals or particulate matter suspended in your water supply must also be considered before purchasing an injector. With




Figure 2 - Fertilizer injectors with multiple heads, such as the Anderson Ratio:Feeders® (A), are necessary when using more than one fertilizer stock solution. In stock solutions, high fertilizer concentrations cause chemical reactions that form insoluble precipitates (B) that can plug up the injector or irrigation nozzles.

the simpler venturi-type injectors water quality isn't as much of an issue but, with more sophisticated injectors, high amounts of sediment or very hard water can cause excessive wear of the pump mechanism.

Mobility - Fertilizer injectors are typically installed in a permanent protected location, such as a headhouse, where the fertilizer solutions can be mixed and the stock solutions stored. These injectors are plumbed directly into the main irrigation line with valves and a bypass to allow normal irrigation. Some injector models can be mounted on a dolly or cart with quick-connections so that they can be used at several different locations.

Installation of Fertilizer Injectors


When permanently installing a fertilizer injector, we recommend the following. First, install injectors in the headhouse or other insulated building to prevent freezing damage and wear from exposure to the elements. Second, install a filter in the water supply line before the injector to filter debris and reduce wear. Third, install water pressure gauges before and after the filter — a large difference in the pressure readings means the filter is plugging up (Pennisi and Kessler 2003). Plumbing codes require that all potable water systems be protected with a backflow prevention device to insure that contaminated water is not accidentally mixed with water that is used for human consumption. Injecting any chemical without backflow prevention is against the law. Backsiphoning occurs when negative water pressure causes contaminated water to be sucked back into the water supply line. The most commonly used backflow preventer is the vacuum breaker (Figure 3). Under normal water pressure, the valve remains closed (Figure 3A); however, if the

pressure in the supply line drops below a predetermined level, the check valve will close and shut off the water supply (Figure 3B). Backflow devices should be installed between the last control valve of the supply system and the fertilizer injector (Koths and others 1976).

Scheduling Fertigation

Two basic schedules for applying liquid fertilizers are constant and periodic. The application of a dilute fertilizer solution each time the crop is irrigated is known as constant fertilization (Landis and others 1989), and the concentration of this applied fertilizer solution is exactly the nutrient concentration desired in the growing medium solution. Periodic fertilization consists of applying a more concentrated fertilizer solution according to some fixed schedule, such as once a week or every other irrigation. The applied fertilizer solution during periodic fertilization may therefore be several times more concentrated to allow for the dilution that occurs during subsequent irrigations. Because periodic application applies a more concentrated solution, growers should rinse crop foliage with irrigation water following each fertigation, as well as carefully monitor to avoid fertilizer salt build-up in the growing medium. An example of a periodic fertilization schedule is given in Table 2. One option is to use continuous fertigation early in the growing season to force growth and build-up plant nutrient reserves, and then change to periodic fertilizer applications to finish the crop. In one study, early season continuous fertigation followed by late season weekly fertigation reduced fertilizer costs by approximately 50% without any growth loss (Struve and Rose 1998). An alternative method is to fertigate using the exponential fertilization method whereby plants receive proportional amounts of fertilizer relative to their growth rate and size (Dumroese and others 2005).

Figure 3 - Vacuum breakers are one type of backflow device that prevents water that has been mixed with fertilizer from being sucked back into the water supply line (modified from Koths and others 1976).



Plant Growth Phase	Timing (weeks)	Type of Fertilizer	Frequency
Germination	0 to 4	H ₃ PO ₄	As needed
Establishment	4 to 8	7-40-17	Every irrigation
Rapid Growth	8 to 12	20-7-19	Every other irrigation
Hardening	12 to 14	4-25-35	Every third irrigation
Pre-Shipping	Prior to harvest	20-7-19	Once




Figure 4 - Checking the electrical conductivity at each stage of the fertigation process (1-5) ensures that the injector is working properly and regular foliar analysis (6) proves that your crop is receiving the proper levels of all mineral nutrients (A). The most critical check is the applied fertilizer solution that goes on your crop (B).

The best way to determine when to fertigate is to carefully monitor plant growth and foliar nutrient levels. Accumulating test results in a spreadsheet program along with seedling growth data allows easy analysis and creates a permanent database that only improves as you gain experience with each crop. As growth versus nutrient curves are developed, it is easy to identify the critical point in the curve when growth begins to flatten out. When this happens, applying more fertilizer will only lead to luxury consumption and, in the case of nitrogen and phosphorus, may cause environmental pollution. Inexperienced growers have the tendency to overfertilize "just to make sure!" and because fertilizer is relatively inexpensive (Landis and others 2005).

How to Monitor Fertigation

Fertigation is a powerful cultural tool but must be carefully monitored. There are two ways to check your fertigation program: electrical conductivity (EC) and foliar nutrient levels. The best way to determine if your fertilizer injector is working properly is to monitor the EC of the various fertilizer solutions. EC is a measure of the salinity (total salt level) of a solution and therefore gives an indication of the dissolved fertilizer salts. An EC meter measures the electrical charge carried by the ions that are dissolved in a solution — the more concentrated the ions, the higher the reading. By checking the EC at each step in the process (Figure 4A), you can be sure that your injector is functioning properly.

1. **Irrigation water** - The base EC of the irrigation water should be monitored monthly, or until you are certain that it does not vary significantly during the season.

2. **Fertilizer stock solutions** - The efficiency of the fertilizer injector can be checked by making an "applied strength" dilution of the fertilizer stock solution and measuring the EC level. For a 1:100 injector, add one part of stock solution to 100 parts of irrigation water. The EC reading of the diluted fertilizer solution should be approximately the same (within 10%) as the EC of the fertigation solution that is applied to your crop.

3. **Applied fertilizer solution** - The applied fertilizer solution is by far the most important of the fertilization checks because this solution actually contacts the seedling foliage and enters the root zone. Even if you don't check anything else, be sure to do this test regularly. The applied solution is collected directly from the irrigation nozzle (Figure 4B) and the EC reading should be approximately the sum of the base salinity of the irrigation water plus the salts added by the fertilizer stock solution. Send a sample of this solution to a testing laboratory and check the levels of the mineral nutrients against your calculated values.

4. **Growing medium extract** - Samples of the irrigation water and the applied fertilizer solution can be collected directly, but liquid samples must be extracted from the growing medium. The amount of growing medium solution is relatively small and is strongly absorbed, and so special sampling techniques must be used to collect enough solution to measure. The amount of growing medium solution is relatively small and is strongly

absorbed, and so special sampling techniques must be used to collect enough solution to measure. See Landis and Dumroese (2006) for details on the various options.

5. Leachate - The final check involves taking EC readings on the "leachate" solution that drains from the bottom of the containers. Leachate can be obtained by taping a test tube or other container to the drain hole of the container or by placing a tray under a block of containers during fertigation. If the EC of the leachate significantly exceeds the EC of the applied fertilizer solution, then excess salinity is building up in the growing medium and proper leaching is not occurring.

6. Foliar nutrient levels - While EC readings can reveal when overall problems with your fertigation system, the only comprehensive test is to chemically analyze the foliage of your crop and determine its nutrient status. The mineral nutrient concentration of the seedling foliage reflects the actual uptake of all the mineral nutrients. Several commercial suppliers of horticultural products are offering chemical testing of irrigation water, fertilizer solutions, growing media, and seedling tissue at very attractive prices. These labs are equipped with the latest analytical equipment such as the ICAP (Inductively Coupled Argon Plasma) spectrometer and so the tests are done quickly and accurately. They will even telephone, FAX, or email the results back to the nursery so that cultural corrections can be made within a matter of days. Interpretation of foliar tests can be intimidating, but general standards and helpful hints can be found in Landis and others (2005).

Summary

Fertigation is one of the most efficient ways for growers to fertilize their crops because all the essential mineral nutrients are applied at their ideal concentration and in the proper balance. In addition, fertigation does not suffer from the delayed response of solid fertilizers because the nutrients are already dissolved in water and can be quickly absorbed by the roots. Most growers use some type of injector to mix concentrated fertilizer solution into the irrigation system and a wide variety of injectors are available to meet the needs of any size of nursery. Injectors must be properly installed with a backflow device to prevent siphoning of liquid fertilizer back into the water source. Fertigation can be applied with each irrigation or at scheduled intervals; the choice will depend on crop response and the risk of excessive nutrient runoff. A well-designed fertigation system can be monitored at several stages in the process to ensure that the injector is working properly and that the plants are receiving the proper amount of fertilizer.

References

- Dumroese RK, Page-Dumroese DS, Salifu KF, Jacobs DF. 2005. Exponential fertilization of *Pinus monticola* seedlings: nutrient uptake efficiency, leaching fractions, and early outplanting performance. *Canadian Journal of Forest Research* 35(12): 2961-2967.
- Kessler R, Pennisi B. 2004. Greenhouse fertilizer injectors: selection, maintenance and calibration. Normal (AL): Alabama Cooperative Extension. Publication ANR-1243. 24 p. URL: <http://www.aces.edu/pubs/docs/A/ANR-1243> (accessed 12 Mar 2010).
- Koths JS, Judd RW Jr, Maisano JJ Jr, Griffin GF, Bartok JW, Ashley RA. 1976. Nutrition of greenhouse crops. Storrs (CT): University of Connecticut, Cooperative Extension Service. 20 p.
- Landis TD, Dumroese RK. 2006. Monitoring electrical conductivity in soils and growing media. Central Point (OR): USDA Forest Service, Pacific Northwest Region. Publication No. R6-CP-TP-04-2006: 6-10.
- Landis TD, Tinus RW, McDonald SE, Barnett JP. 1989. Seedling nutrition and irrigation, vol 4. The Container Tree Nursery Manual. Washington (DC): USDA Forest Service. Agriculture Handbook 674. 119 p.
- Landis TD, Haase DL, Dumroese RK. 2005. Plant nutrient testing and analysis in forest and conservation nurseries. In: Dumroese RK, Riley LE, Landis TD, technical coordinators. National proceedings, forest and conservation nursery associations-2004. Ft Collins (CO): USDA Forest Service, Rocky Mountain Research Station. Proceedings RMRS-P-35: 76-84.
- Pennisi B, Kessler R. 2003. Fertilizer Injectors: Selection, Maintenance and Calibration. Athens (GA): University of Georgia, College of Agricultural and Environmental Sciences, Cooperative Extension Service. Bulletin 1237. 16 p. URL: <http://www.aces.edu/pubs/docs/A/ANR-1243/ANR-1243.pdf> (access ed 12 Mar 2010).
- Triebwasser MT, Altsuler SL. 1995. Fertilization practices and application procedures at Weyerhaeuser. In: Landis TD, Clegg B, technical coordinators. National proceedings: forest and conservation nursery associations-1995. Portland (OR): USDA Forest Service, Pacific Northwest Research Station. General Technical Report GTR-365: 84-88.
- Weiler TC, Sailus M, editors. 1996. Water and nutrient management for greenhouses. Ithaca (NY): Northeast Regional Agricultural Engineering Service. Publication NRAES-56. 102 p.

Managing surplus or holdover nursery stock

By Thomas D. Landis

So, you've survived another hectic lift-and-pack season but you are left with some surplus or "holdover" plants. Maybe your sowing factors were a little too generous or you overestimated the market for a particular species. Sometimes, poor outplanting site conditions or operational problems means that some stock must be held over for another season. Surplus plants can also show up during grading. Sowing problems, poor weather or cultural shortcomings can result in perfectly good plants that don't meet grading specifications. Overly dense seedbeds produce stunted plants that lack the desired stem diameter, or excessive fertilization produces excessive height growth. What about the large plants that exceed both height and stem diameter specifications? These are often the genetically superior plants that you just hate to throw away. Even if you were aware of the surplus and didn't harvest the stock, you can't leave them in the seedbed or in the containers for too long.

One of the most difficult concepts for novice nursery managers and their customers is that, unlike many products, plants have a shelf-life. Nursery stock is at its peak quality when the plants are harvested and graded and, ideally, they can be shipped and outplanted soon afterwards. Of course, that often isn't possible so the plants must be placed in some sort of storage that maintains that quality. In the days before refrigerated storage, bareroot nurseries "heeled-in" their stock but this mainly protected the roots from desiccation and eventually plants would break bud. Early container nurseries tried to maintain plant quality by placing their stock in lathhouses or other shaded storage, but again, this was a short-term solution. Eventually, the plants would begin to grow and quality would suffer. Refrigerated storage will maintain plant quality for months but eventually, if you can't sell or ship them, something must be done.



Figure 1 – The best option for surplus or holdover bareroot stock is to root prune and transplant them (A); transplanting increases in-row growing space, which produces greater stem diameter (B).

One of the greatest challenges in nursery management is what to do with plants that have reached target size but haven't been sold or shipped for outplanting. Surplus or holdover stock happens in both bareroot and container nurseries but, as we will see, is a much greater problem in containers.

Bareroot stock — If the surplus plants are still in the ground, you need to evaluate the seedbed density. Seedbeds that are not too dense and still have lateral room to grow can be root pruned and/or wrenched to slow shoot growth, develop a more fibrous root system, and increase stem diameter. In most cases, however, plants are already too dense and would become stunted if left in place for another season. Overly dense crops are also an ideal breeding ground for fungal pathogens such as *Botrytis cinerea*. So, the best option is to harvest the plants, grade, and transplant them (Figure 1A). These plants and stock that is already in storage will have to be root pruned because the target root length for outplanting is much too long for transplanting. Transplanting increases growing space with the row, which produces plants with greater stem diameter (Figure 1B). The greatest challenge will be to keep the

shoot-to-root ratio in balance, so these transplants will have to be root pruned or wrenched, which will also increase root fibrosity.

Container stock — Container nursery culture has led to increased growth rates because of the greater control over most potentially limiting environmental factors. The challenge comes when we want to stop that growth, especially in roots because they don't go dormant. Shoots can be coaxed into dormancy by cultural manipulations of daylength (blackout), mineral nutrition, and water supply, but how do you stop roots from growing? The only way to do this is with cold temperatures, which is why refrigerated storage has become so popular (Landis and others 2010). Nurseries in milder climates that use open or sheltered storage can have a serious problem because roots continue to grow even after shoot growth has stopped. Tropical nurseries suffer the greatest risk of plants becoming "rootbound" because their stock never goes dormant.

Rootbound nursery stock can be defined as plants that have grown too large for their containers, resulting in severe matting and tangling of the root system (Figure 2A).




Figure 2 – When container plants have been held too long, they become “rootbound” (A). Transplanting them to bareroot beds or larger containers after making vertical cuts along the rootplug (B) is the recommended practice to hold them over for another season.

Observations have related rootbinding to the length of time that the plant has been in the container (Balisky and others 1995). Logically, the larger the container, the longer it takes for the plant to become rootbound. But time alone is not the only controlling factor, because root growth is also affected by cultural conditions at the nursery. A species growing rapidly in one nursery will become rootbound faster than the same species growing more slowly in another nursery. Similarly, a species in a large container given large amounts of fertilizer may become rootbound as fast as the same species in a smaller container given smaller amounts.

The fact that excessive root growth can be a quality issue in container plants has been known for decades. During the early 1980s there was considerable concern about "toppling" of lodgepole pine (*Pinus contorta*) container stock which was proven to be caused by poor root egress after outplanting (Burdett and others 1986). This led to the development of copper-coated containers and then to sideslit containers which encourage roots to egress all along the length of the plug instead of just at the bottom. Trials into extending the growing period to produce larger stocktypes also resulted in plants with excessive root biomass for their respective container volumes. Whatever the cause, container plants that have plugs with high root densities suffered poor survival and growth for several years after outplanting (Salonius and others 2002).

How to characterize rootbound plants has been a challenge. South and Mitchell (2006) propose a "root-bound index" based on root-collar diameter divided by container diameter or volume, but this index must be calculated for each container type. From an operational standpoint, establishing a maximum stem diameter along with a visual assessment of root binding might be the most practical system (Landis and others 2010).

Okay, what do you do if your container plants have become rootbound? The best option is to transplant into larger volume containers or into bareroot beds. In fact, the relatively new plug+one stocktype was originally developed as a way to hold over container seedlings (Hahn 1984). If you want to keep your plants as container stock, then you can just transplant into another container that is large enough to support new root growth. It's a good idea to cut the root plug from top to bottom at a couple of places as well as trim the roots at the drainage hole (Figure 2B). This process is time consuming but encourages new root growth all along the length of the original root plug.

Summary

Surplus or holdover stock is sometimes inevitable but careful planning and good communication with customers can reduce the instances. Bareroot stock should have their roots trimmed and then be transplanted; the plants may need to be pruned or wrenched to maintain a good shoot-to-root ratio. Holdover container stock is more of a challenge because the plugs can become severely rootbound. Make vertical cuts along the root plug before transplanting them to bareroot beds or containers large enough to promote new root growth while retarding excessive shoot growth.

References

- Balisky AC, Salonius P, Walli C, Brinkman D. 1995. Seedling roots and the forest floor: misplaced and neglected aspects of British Columbia's reforestation effort? *Forestry Chronicle* 71: 59-65.
- Burdett AN, Coates H, Eremko R, Martin PAF. 1986. Toppling in British Columbia's lodgepole pine plantations: significance, cause and prevention. *Forestry Chronicle* 62(5): 433-439.
- Hahn PF. 1984. Plug + 1 seedling production. In: Duryea ML, Landis TD, editors. *Forest nursery manual: production of bareroot seedlings*. The Hague/Boston/Lancaster: Martinus Nijhoff/DrW. Junk Publishers, for Forest Research Laboratory, Oregon State University, Corvallis. p 165-181.
- Landis TD, Dumroese RK, Haase DL. 2010. Seedling processing, storage, and outplanting, vol 7. *The Container Tree Nursery Manual*. Washington (DC): USDA Forest Service. *Agriculture Handbook* 674. 200 p.
- Salonius P, Hallett R, Beaton K, French C. 2002. Extended nursery rearing compromises field performance of container-reared conifer seedlings. Fredericton (NB): Canadian Forest Service, Atlantic Forestry Centre, Information Report M-X-214E. 21 p.
- South DB, Mitchell RG. 2006. A root-bound index for evaluating planting stock quality of container-grown pines. *Southern African Forestry Journal* 207: 47-54.

Late-Season Fertilization of Nursery Stock

By Diane L. Haase and Thomas D. Landis

Regulating seedling nutrient content through fertilization is a key component of nursery culture for both bareroot and container stock. Typically, fertilizers are applied early in the growing season to fuel active shoot growth. Then, fertilization (especially nitrogen) is reduced or stopped to induce budset and promote development of cold hardiness, usually during July through September depending on species, seed source, and stocktype specifications. However, a significant amount of root and stem growth can still occur late in the growing season as long as temperatures remain within favorable ranges. This increase in biomass late in the growing season can lead to nutrient dilution within the plant unless more nutrients are supplied through fertilization. If nutrient concentrations drop below the adequate range, there may be inadequate reserves for vigorous growth following outplanting. However, many growers are concerned about the traditional belief that fertilizing too late in the season will cause budbreak, stimulate additional shoot growth, or delay or reduce cold hardiness.

To prevent nutrient dilution, some nurseries apply late-season fertilizers after shoot growth has ceased (Figure 1A). Because seedlings are actively growing roots well into the fall, there is great potential to increase seedling nutrient content with fertilization. Nutrient loading is a relatively recent cultural practice in which late-season fertilization is used to increase seedling nutrient reserves with the objective of promoting additional growth after outplanting. Nutrient concentration in nursery plants follows a classic uptake curve and nutrient loading is the uptake of nutrients beyond the adequate range, but not so much that toxicity is reached (Figure 1B). This enhanced internal nutrient reserve is thought to increase root egress and promote




Figure 1 – Late season fertilization is applied after shoot growth has ceased (A) with the objective of “loading” the plants with extra mineral nutrients (B) to promote better growth after outplanting. (B, modified from Chapman 1967).

faster shoot growth immediately following outplanting. Nutrient loading is promoted to improve seedling field performance, especially on nutrient-poor sites or sites with heavy competition (Timmer 1997).

Numerous research studies on late season fertilization have been conducted, and results have been variable to say the least (Table 1). One explanation for this variation is that studies have been done on different crops,

Table 1 - Summary of research studies on with late-season fertilization of conifer seedlings

Source	Stocktype	Type and Rate of fertilizer	Application timing	Effects on nursery stock
Benzian and others (1974)	Bareroot	NH ₄ NO ₃ +CaCO ₃ or K ₂ SO ₄ at 62 or 125 lb/ac	Early September	Increased foliar nitrogen (N) concentrations; earlier budbreak; increased growth of spruce but reduced survival of fir
Birchler and others (2001)	Bareroot	NH ₄ NO ₃ +K ₂ SO ₄ or (NH ₄) ₂ SO ₄ +KCL at 0 to 285 lb N and K/ac	September to November	Increased foliar N; root growth potential and cold hardiness unaffected; earlier budbreak; no effect on outplanting performance
Boivin and others (2002)	Container	Soluble 20-20-20 NPK at 6 or 12 mg N per seedling	During hardening period	N, P and K uptake increased up to 164, 70 and 32%, respectively; greater biomass; lower shoot-to-root ratio
Boivin and others (2004)	Container	Soluble 20-20-20 NPK at 0 to 48 mg N / seedling	9 weeks after bud set	Increased N uptake and growth after outplanting; reduced survival at the highest rate
Hinesley and Maki (1980)	Bareroot	Macronutrient fertilizers at 150 N lb/ac	Late October	Increased seedling size and foliar nutrient concentrations
Irwin and others (1998)	Bareroot	NH ₄ NO ₃ at 51 lb N/ac applied 0 to 3 times	November to December	Increased N concentration with no change in morphology; increased first-year field height and survival
Islam and others (2009)	Bareroot	NH ₄ NO ₃ at 0 to 80 lb N/ac	Mid-September	Increased shoot height, bud size, number of needle primordia, N concentration, and cold hardiness
Montville and others (1996)	Container	foliar fertilization (27-15-12) at 324 to 972 ppm	Twice weekly during budset	Increased stem diameter, bud length, shoot biomass, and N concentration
South and Donald (2002)	Bareroot	0, 134 lb/ac N, 134 lb/ac of N + 134 lb/ac of P, or 134 lb/ac of K	Early October	Increased N concentration with no effect on morphology; variable effects after outplanting
Sung and others (1997)	Bareroot	NH ₄ NO ₃ at 0 to 36 lb/ac	Mid-September	Fewer culls at high rate; increased first order lateral roots and dry weights
van den Driessche (1985)	Bareroot	Macronutrient fertilizers at 0 to 71 lb N/ac	July to October	Increased N concentration and new roots; higher relative growth rate in sand culture but not in artificial soil; earlier budbreak
VanderSchaaf and McNabb (2004)	Bareroot	NH ₄ NO ₃ at 0 to 178 lb N /ac	January	Increased N concentration; no effect on morphology; greater growth after outplanting
Williams and South (1992)	Container	(NH ₄) ₂ HPO ₄ at 0 to 180 lb N/ac	September to November	Fertilization did not delay the progression of the bud dormancy cycle; temporary effects on cell division




Figure 2 – When black spruce (*Picea mariana*) seedlings were loaded with NPK fertilizer, nitrogen showed the greatest increase. For each nutrient, bars with the same letter are not statistically different ($\alpha \leq 0.05$). Adapted from Boivin and others (2004).

with different fertilizers, at different rates, and applied at different times. Therefore, it is impossible to generalize about the effects of late season fertilization on your particular crop. However, one consistent finding is that late season fertilization greatly increases seedling nutrient concentrations, especially nitrogen (Figure 2). The effects of late season fertilization on seedling morphology, physiology, and subsequent growth after outplanting are also variable, but two out of three studies report some type of positive response. In some cases, seedling size increased in the nursery (Boivin and Timmer 2002; Hinesly and Maki 1980; Islam and others 2009; Montville and others 1996; Sung and others 1997) but no additional growth was measured in other studies (Boivin and others 2004; Irwin and others 1988; South and Donald 2002; VanderSchaaf and McNabb 2004). Late season fertilization can affect budbreak the following year; several studies report that budbreak was earlier in the following season for nutrient-loaded seedling compared to control seedlings (Benzian and Freeman 1974; Birchler and others 2001; van den Driessche 1985). After outplanting, growth of seedlings with increased nutrient reserves was found to be greater in some cases (Benzian and Freeman 1974; Boivin and others 2004; Hinesly and Maki 1980; Irwin and others 1988; South and Donald 2002; van den Driessche 1985; VanderSchaaf and McNabb 2004) but was unaffected in others (Benzian and Freeman 1974; Birchler and others 2001; South and Donald 2002; van den Driessche 1985). In a few cases, very high fertilization rates resulted in reduced survival after outplanting (Benzian and Freeman 1974; Boivin and others 2004; South and Donald 2002) indicating the need to avoid excessive nutrient loading into the toxic range (Figure 1B).

Late season fertilizer applications must be carefully scheduled to avoid negative effect on phenology. By applying fertilizers after plants have been exposed to cold nights, the chances of stimulating Lamm growth will be lessened. One of the biggest concerns about late season fertilization is that it would decrease cold hardiness. One study found that cold hardiness is unaffected by properly applied, late-season fertilization (Birchler and others 2001) and, in fact, the development of cold hardiness can actually be impaired if plant nutrient concentrations are too low.

Summary

Late-season fertilization has potential for improving seedling quality and outplanting success but nurseries should conduct trials on fertilizer formulations and rates in order to develop an optimum treatment for a specific crop. Monitor mineral nutrient uptake through seedling nutrient analysis to ensure that fertilization treatments are effective, and prevent overfertilization which can reduce plant quality and accelerate nutrient runoff.

References

- Benzian B, Brown RM, Freeman SCR. 1974. Effect of late-season top-dressings of N (and K) applied to conifer transplants in the nursery on their survival and growth on British forest sites. *Forestry* 47: 153-184
- Birchler TM, Rose R, Haase DL. 2001. Fall fertilization with N and K: Effects on Douglas-fir seedling quality and performance. *Western Journal of Applied Forestry* 16: 71-79.
- Boivin JR, Miller BD, Timmer VR. 2002. Late-season fertilization of *Picea mariana* seedlings under greenhouse culture: biomass and nutrient dynamics. *Annals of Forest Science* 59: 255-264.
- Boivin JR, Salifu KF, Timmer VR. 2004. Late-season fertilization of *Picea mariana* seedlings: intensive loading and outplanting response on greenhouse bioassays. *Annals of Forest Science* 61: 737-745.
- Chapman HD. 1967. Plant analysis values suggestive of nutrient status of selected crops. In: *Soil Testing and Plant Analysis II*. Madison, WI: Soil Science Society of America. Special Publication Series No. 2: 77-92.
- Hinesley LH, Maki TE. 1980. Fall fertilization helps longleaf pine nursery stock. *Southern Journal of Applied Forestry* 4: 132-135.
- Irwin KM, Duryea ML, Stone EL. 1998. Fall-applied nitrogen improves performance of 1-0 slash pine nursery seedlings after outplanting. *Southern Journal of Applied Forestry* 22: 111-116.
- Islam MA, Apostol KG, Jacobs DE, Dumroese RK. 2009. Fall fertilization of *Pinus resinosa* seedlings: nutrient uptake, cold hardiness, and morphological development. *Annals of Forest Science*. 66: 704.
- Montville ME, Wenny DL, Dumroese RK. 1996. Foliar fertilization during bud initiation improves container-grown ponderosa pine seedling viability. *Western Journal of Applied Forestry* 11:114-119.
- South DB, Donald DGM. 2002. Effect of nursery conditioning treatments and fall fertilization on survival and early growth of *Pinus taeda* seedlings in Alabama, U.S.A. *Canadian Journal of Forest Research* 32: 1171 - 1179.
- Sung SS, Black CC, Kormanik TL, Zarnoch SJ, Kormanik PP, Counce PA. 1997. Fall nitrogen fertilization and the biology of *Pinus taeda* seedling development. *Canadian Journal of Forest Research* 27: 1406 - 1412.
- Timmer VR. 1997. Exponential nutrient loading: a new fertilization technique to improved seedling performance on competitive sites. *New Forests* 13:279-299.
- van den Driessche R. 1985. Late-season fertilization, mineral nutrient reserves, and retranslocation in planted Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) seedlings. *Forest Science* 31: 485-496.
- VanderSchaaf C, McNabb K. 2004. Winter nitrogen fertilization of loblolly pine seedlings. *Plant and Soil* 265:295-299,
- Williams H, South DB. 1992. Effects of fall fertilizer applications on mitotic index and bud dormancy of loblolly pine seedlings. *Forest Science* 38:336-349.

Bird Damage to Sown Seeds or Emerging Seedlings

By Thomas D. Landis

It's spring so most growers are sowing their crops or will be doing so in the next couple of months. Many things can go wrong during sowing (Figure 1), and all gardeners know the anxiety of waiting for those germinating plants to "poke their heads" out of the soil or growing medium. Seeds are small packets of high-energy food and therefore are especially attractive to birds. Trying to keep these critters from eating all your crop seeds is one of those concerns that man has faced since the beginning of recorded time.

It is difficult to quantify the overall impact of animal damage because the incidents are generally episodic. Bird predation is often due to migrating flocks, which can do severe damage in a short time. One nursery that participated in the Container Nursery Survey reported that from 25 to 50% of their sown seeds were eaten by goldfinches (Landis and others 1989). Bird damage is often localized -- losses varied from 0 to 75% of ponderosa pine (*Pinus ponderosa*) seeds in one bareroot nursery (Landis 1976).

Current information on bird predation is hard to locate but the best sources are websites. For instance, up-to-date information on all types of animal damage can be found at the Internet Center for Wildlife Damage Management, which is a non-profit, grant-funded site that provides research-based information on how to responsibly handle wildlife damage problems (Vantassel 2010).

On that site, you can download Prevention and Control of Wildlife Damage, which contains information on diagnosing and controlling all types of wildlife pests including birds (Hygnstrom and others 1994). Not much has been published on bird predation in forest, conservation, and native plant nurseries so much of this article was based on Volume Five: Nursery Pests and Mycorrhizae of the Container Tree Nursery Manual series (Landis and others 1989).

Hosts - Birds will eat seeds of all conifer and native plant species but prefer the large-seeded pines such as white pine, sugar pine, and pinyon. Crows and ravens can damage seed beds of large hardwood seeds such as oaks.

Symptoms/damage - If seeds cannot be located in the container or seedbed, but the seed covering has been scattered around, then bird predation is a possibility (Figure 1A). However, it can be difficult to distinguish between bird and rodent damage. Birds generally eat seeds immediately leaving spent seedcoats, whereas rodents often cache uneaten seeds. Rodents feed mostly at night, whereas birds usually feed during the day. Rodent predation can occur in open and closed growing areas, but bird predation is more common in open compounds. Birds also cause clipping injury to emerging seedlings by feeding on the seed coat that clings to the cotyledons (Figure 1 B). Larger seedlings can sometimes recover from clipping injury, although severely damaged




Figure 1 - A damage key is helpful to separate bird predation from the many things that can happen to newly-sown seeds (A). Besides consuming seeds, birds can clip the tops of germinants, which usually results in their death (B) (A, modified from Landis and others 1989).




Figure 2 – Netting has been effective in preventing bird damage in both bare-root (A) and container nurseries. Noise repellents, such as the propane cannon (B) work initially, but birds soon become tolerant.

germinants are weakened and prone to other pests, such as damping-off fungi.

Management - Bird predation is best controlled through prevention: exclusion through proper growing area design and use of screens, and elimination of suitable habitat around the growing area. Control measures include (Fuller and others 1984):

1. Netting - Plastic bird netting is available in several lengths and widths. It is applied over sown containers or seedbeds (Figure 2A) and physically protects seeds and germinants until they are established. Wire

hoops support the netting and the mesh size is large enough to allow rain or irrigation to reach the plants. A system for mechanically applying bird netting to bare-root seedbeds has been developed (Skakel and Washburn 1989).

2. Trapping and baiting - Birds are opportunists and can sometimes be lured away from crop seeds if you supply another more desirable food source away from the nursery. Surplus seeds can be scattered around to attract birds and keep them from reaching the crop seeds.

Table 1 - Characteristics of chemical bird repellents used in forest and conservation nurseries *

Product	Active Ingredient	Remarks
Dithianon	Anthraquinone	A quinone fungicide that is registered by the EPA for bird control. It also occurs naturally in some plants, such as aloe, senna, rhubarb, and Cascara buckthorn
Mesuro [®] 75 W	Methiocarb	Registered by EPA as an insecticide and molluscicide. Commonly used by homeowners to kill snail and slugs
Thiram [®] 50 WP, Arasan, Tersan 74	Dithiocarbamate	Registered by EPA as a fungicide that is applied as a seed treatment
Rejex-It, Bird Shield [™]	Methyl anthranilate	Several formulations are available that make birds sick without killing them. Active ingredient is found in concord grapes, and is used as a flavoring in grape soda
Hot Sauce [®] Animal Repellent, Deer Away	Capsaicin	Available commercially in concentrations from 2.5 to 6.2% Homemade formulations are made by grinding dried, ripe <i>Capsicum frutescens</i> chili peppers into a fine powder and mixing with a solvent.

* Modified from Colorado State University Extension, 2007. All pesticide information is subject to change so be sure to check the labels or on-line sources

3. Chemical repellents - You might think that the simplest and most effective way to prevent bird predation of seeds is to apply a repellent before sowing. Several chemicals have been used to repel birds in nurseries (Table 1).

Unfortunately, birds and mammals apparently have large differences in tolerances to various repellents. For instance, pen studies have shown that capsaicin products have effectively repelled deer and elk but other observations have shown that birds will readily consume seeds treated with capsaicin concentrations as high as 2% (Colorado State University Extension 2007).

4. Visual repellents - Altering the appearance of seeds may help delay or prevent predation. Aluminum powder has traditionally been used in forest nurseries to keep conifer seeds from sticking together during mechanical sowing. More recently, some nurseries coat their seeds with DayGlo® paint pigment to make sown seeds easier to see in the furrow or container. No published research exists, but changing the color of seeds may be an effective bird repellent.

5. Noise repellents - Many devices have been used for frightening birds including portable propane cannons (Figure 2B). While initially effective, most birds eventually become accustomed to the noise.

A good source for all types of animal control chemicals and equipment can be found on-line (Hygnstrom and others 1994).

Final Thoughts

Considering the cost of seed and the amount of time and energy expended during the sowing process, it only makes sense to try and protect newly-sown crops from bird predation:

1. Effectiveness is a factor of motivation and habituation. Repellents are less effective when birds are hungry or other sources of food are unavailable. All animals are creatures of habit, and habituation can complicate repellent efforts.

2. Phytotoxicity, worker safety, and environment hazards. Any chemical applied to seeds has the potential to adversely affect seed germination or young seedlings. Pesticides registered as safe for other crops may not have been tested on trees and other native plants. The repellents listed in Table 1

vary widely in their potential toxicity to nursery workers or other animals.

3. Availability & cost. Most commercial repellents are readily available through garden centers or reforestation suppliers, and their cost is minimal compared to the potential crop damage. Others, such as pepper sprays, can be homemade.

References

- Colorado State University Extension. 2007. Chemical, visual and auditory repellents for reducing problems with urban wildlife. URL: <http://www.colostate.edu/Depts/CoopExt/wildlife/repellents.htm> (accessed 31 Mar 2010).
- Fuller R, Landis T, Cummings J, Guarino J. 1984. Mesurol 75% seed treater as a bird repellent seed coat treatment. *Tree Planters' Notes* 35(1):12-17.
- Hygnstrom SE, Timm RM, Larson GE, editors. 1994. Prevention and control of wildlife damage. Lincoln (NE): University of Nebraska-Lincoln. 2 vols. URL: <http://icwdm.org/handbook/index.asp> (1 Apr 2010).
- Landis TD, Tinus RW, McDonald SE, Barnett JP. 1989. The biological component: nursery pests and mycorrhizae, vol 5. *The Container Tree Nursery Manual*. Washington (DC): USDA Forest Service. Agriculture Handbook 674. 171 p.
- Landis TD. 1976. An analysis of seed and seedling losses at Mt. Sopris tree nursery (CO). Denver (CO): USDA Forest Service, Forest Insect and Disease Management. Biological Evaluation R2-76-18. 7 p.
- Skakel S, Washburn J. 1989. Mechanized application of bird netting to protect germinating seedlings. *Tree Planters' Notes* 40(4):14-17.
- Vantassel S. 2010. Internet center for wildlife damage management. Lincoln (NE): University of Nebraska-Lincoln. URL: <http://icwdm.org/> (accessed 1 Apr 2010).



A compact disk with all the following journal articles or publications in Adobe PDF format can be ordered using the Literature Order Form on the last page of this section. Note that there are a 2 restrictions:

1. **Copyrighted Material.** Items with © are copyrighted and require a fee for each copy, so we will only send you the title page and abstract. If you want the entire article, you can order copies on-line or from a library service.
2. **Special Orders (SO).** Special orders are books or other publications that, because of their size or cost, require special handling. For some, the Forest Service has procured copies for free distribution, but others will have to be purchased. Prices and ordering instructions are given following each listing in the New Nursery Literature section.

Ordering information for:


Volume Seven of the Container Tree Nursery Manual series

"Yes, I know that it's been almost 2 years since I first announced its publication, but the government moves at a snail's pace. We sent the final draft to the Forest Service editors in Washington, DC a month ago, and they have contracted with the US Government Printing Office to print the book. The Forest Service has purchased a limited quantity of books for free distribution, and since we expect demand to be high, I would recommend ordering a copy right away"

Ask for a free copy of **Agriculture Handbook 674:**
The Container Tree Nursery Manual - Volume Seven.

Order from:

Richard D. Schneider, Publications Distribution
 USDA-FS Rocky Mountain Research Station
 240 W. Prospect Road
 Ft. Collins, CO 80526-2098 USA
 TEL: 970.498.1392 • FAX: 970.498.1396
[E-mail: rschneider@fs.fed.us](mailto:rschneider@fs.fed.us)



1. © Producing liminal space: gender, age and class in northern Ontario's tree planting industry. Sweeney, B. Gender, Place and Culture 16(5):569-586. 2009.



Container Production

2. © Cavity size and copper root pruning affect production and establishment of container-grown longleaf pine seedlings. Sayer, M. A. W., Haywood, J. D., and Sung, S.-J. S. *Forest Science* 55(5):377-389. 2009.
3. Degradable pots could help resolve disposal issues. Hayes, B. *Greenhouse Management and Production* 30(1):25-29. 2010.
4. Improving root growth and morphology of containerized Oregon white oak seedlings. Devine, W. D., Harrington, C. A., and Southworth, D. *Tree Planters' Notes* 53(2):29-34. 2009.
5. New tree techniques at Lancaster Farms. Jones, S. T. *International Plant Propagators' Society, combined proceedings*, 2008, 58:552-554. 2009.
6. Retractable roof greenhouse (RRG) versus bare-root tree liner post harvest survival and growth in nursery production. Mathers, H. M., Case, L. T., and Rivera, D. *International Plant Propagators' Society, combined proceedings*, 2008, 58:260-267. 2009.



Diverse Species

7. © Assisted revegetation in a subarctic environment: effects of fertilization on the performance of three indigenous plant species. Deshaies, A., Boudreau, S., and Harper, K. A. *Arctic, Antarctic, and Alpine Research* 41(4):434-441. 2009.
8. Bringing alpines down to earth. Fieseler, K. *International Plant Propagators' Society, combined proceedings*, 2008, 58:297-298. 2009.
9. © Can pine forest restoration promote a diverse and abundant understory and simultaneously resist nonnative invasion? McGlone, C. M., Springer, J. D., and Laughlin, D. C. *Forest Ecology and Management* 258:2638-2646. 2009.
10. Combined definition of seed transfer guidelines for ecological restoration in the French Pyrenees. Malaval, S., Luga, B., Regnault-Roger, C., and Largier, G. *Applied Vegetation Science* 13:113-124. 2010.
11. © Comparing direct abiotic amelioration and facilitation as tools for restoration of semiarid grasslands. Pueyo, Y., Alados, C. L., Garcia-Avila, B., and Kefi, S. *Restoration Ecology* 17(6):908-916. 2009.
12. © Controls on the sexual and asexual regeneration of Salicaceae along a highly dynamic, braided river system. Moggridge, H. L. and Gurnell, A. M. *Aquatic Science* 71:305-317. 2009.
13. © The disconnect between restoration goals and practices: a case study of watershed restoration in the Russian River Basin, California. Christian Smith, J. and Merenlender, A. M. *Restoration Ecology* 18(1):95-102. 2009.
14. The easy track to fern spore propagation. Edgren, M. *International Plant Propagators' Society, combined proceedings*, 2008, 58:348. 2009.
15. © The effects of seeding sterile triticale on a native plant community after wildfire in a pinyon pine - mountain mahogany woodland. Waitman, B. A., Draper, T. M., and Esque, T. C. *International Journal of Wildland Fire* 18:659-664. 2009.
16. © Effects of shade on growth and nodulation of three native legumes. Houx, J. H. III, McGraw, R. L., Fritsch, F. B., and Navarrete-Tindall, N. E. *Native Plants Journal* 10(3):233-238. 2009.
17. © Establishment and survival of native legumes on upland sites in Louisiana. Pitman, W. D. *Native Plants Journal* 10(3):240-250. 2009.
18. Evaluation of native warm season grass cultivars for riparian zones. Skinner, R. H., Zobel, R. W., van der Grinten, M., and Skaradek, W. *Journal of Soil and Water Conservation* 64(6):413-422. 2009.
19. Genetic diversity in populations of *Lupinus elegans* Kunth, implications for ecological restoration. Lara-Cabrera, S., Alejandre-Melena, N., Medina-Sanchez, E. I., and Lindig-Cisneros, R. *Rev. Fitotec. Mex.* 32(2):79-86. 2009.
20. © Germination, survival, and growth of grass and forb seedlings: effects of soil moisture variability. Fay, P. A. and Schultz, M. J. *Acta Oecologica* 35:679-684. 2009.
21. Integrating plains natives into captivity. Ackerman, R. R. *International Plant Propagators' Society, combined proceedings*, 2008, 58:228-232. 2009.

22. Micropropagation of upland native species for landscape restoration. Wright, N. International Plant Propagators' Society, combined proceedings, 2008, 58:452-455. 2009.
23. © Moving from model to application: cultural keystone species and reclamation in Fort McKay, Alberta. Garibaldi, A. Journal of Ethnobiology 29(2):323-338. 2009.
24. Multiple causes of seedling rarity in scrub plus, *Prunus geniculata* (Rosaceae), an endangered shrub of the Florida scrub. Weekley, C. W., Zaya, D. N., Menges, E. S., and Faivre, A. E. American Journal of Botany 97(1):144-155. 2010.
25. © Multiple drivers of apparent competition reduce re-establishment of a native plant in invaded habitats. Orrock, J. L. and Witter, M. S. Oikos 119:101 - 108. 2010.
26. © Native bamboo [*Arundinaria gigantea* (Walter) Muhl., Poaceae] establishment and growth after the removal of an invasive non-native shrub (*Ligustrum sinense* Lour., Oleaceae): implications for restoration. Osland, M. J., Pahl, J. W., and Richardson, C. J. Castanea 74(3):247-258. 2009.
27. © Notice of release of Dilley germplasm slender grama. Smith, F. S., Ocumpaugh, W. R., Maywald, P. D., and Lloyd-Reilly, J. Native Plants Journal 10(3):295-298. 2009.
28. © Pocket gophers and the invasion and restoration of native bunchgrass communities. Watts, S. M. Restoration Ecology 18(1):34-40. 2010.
29. Propagating *Arctostaphylos* and *Ceanothus*. Smith, M. N. International Plant Propagators' Society, combined proceedings, 2008, 58:291-294. 2009.
30. Propagating grasses, rushes, and sedges. Low, J. International Plant Propagators' Society, combined proceedings, 2008, 58:287-290. 2009.
31. Propagating Northwest natives. Buzzo, R. J. International Plant Propagators' Society, combined proceedings, 2008, 58:270-275. 2009.
32. Propagation of Colorado natives at Little Valley. Core, B. International Plant Propagators' Society, combined proceedings, 2008, 58:224-227. 2009.
33. Propagation of thirteen Alaska native plants by summer stem cuttings. Holloway, P. S. International Plant Propagators' Society, combined proceedings, 2008, 58:417. 2009.
34. Propagation of *Trichostema lanatum*. Navarez, K. International Plant Propagators' Society, combined proceedings, 2008, 58:374-375. 2009.
35. Propagation of wildflowers from wild collected seeds or cuttings. Thetford, M., Heather, A. E., Perez, H. E., and Wilson, S. B. International Plant Propagators' Society, combined proceedings, 2008, 58:555-560. 2009.
36. Propagation protocol for several high elevation California natives. Funston, N. International Plant Propagators' Society, combined proceedings, 2008, 58:276-278. 2009.
37. © Regional genetic variation in three native grasses in northern California. Kitzmiller, J. H. Native Plants Journal 10(3):263-280. 2009.
38. © Response of two perennial herbaceous Texas legumes to shade. Muir, J. P., Bow, J. R., and Boggs, L. L. Native Plants Journal 10(3):252-261. 2009.
39. © Rethinking species selection for restoration of arid shrublands. Padilla, F. M., Ortega, R., Sanchez, J., and Pugnaire, F. I. Basic and Applied Ecology 10:640-647. 2009.
40. © A role for assisted evolution in designing native plant materials for domesticated landscapes. Jones, T. A. and Monaco, T. A. Frontiers in Ecology and Environment 7(1):541-547. 2009.
41. Seed propagation techniques for selected California natives at Suncrest Nurseries. Smith, M. N. International Plant Propagators' Society, combined proceedings, 2008, 58:403-407. 2009.
42. Seed storage potential for dwarf birch (*Betula glandulosa* Michx.). Forbes, K. and Beardmore, T. Propagation of Ornamental Plants 9(3):143-150. 2009.
43. © Seedling competition between native cottonwood and exotic saltcedar: implications for restoration. Bhattacharjee, J., Taylor, J. P., Jr., Smith, L. M., and Haukos, D. A. Biol. Invasions 11:1777-1787. 2009.
44. © Sulfuric acid scarification of *Callicarpa americana* L. (Lamiaceae) seeds improves germination. Contreras, R. N. and Ruter, J. M. Native Plants Journal 10(3):283-286. 2009.
45. © Survival and genet growth and development of field-planted giant cane (*Arundinaria gigantea*) over time in southern Illinois. Zaczek, J. J., Baer, S. G., Hartleb, J. L., Brendecke, W. W., and Schoonover, J. E. Castanea 74(3):286-299. 2009.
46. Variation in development and response to root-zone pH among seedlings of *Dirca palustris* (Thymelaeaceae) from three provenances. Peterson, B. J. and Graves, W. R. HortScience 44(5):1319 - 1322. 2009.



Fertilization and Nutrition

47. An advisor's experience of the use of compost teas in nursery stock production. Hutchinson, D. International Plant Propagators' Society, combined proceedings, 2008, 58:440-442. 2009.

48. Chlorophyll fluorescence response of *Pinus radiata* clones to nitrogen and phosphorus supply. Bown, H. E., Mason, E. G., Clinton, P. W., and Watt, M. S. Ciencia e Investigacion Agraria 36(3):451-464. 2009.

49. The difference between nutrient solubility and mobility. Mattson, N. and Leatherwood, W. R. Greenhouse Management and Production 29(12):22-26. 2009.

50. Effect of N and NPK fertilizers on early field performance of narrow-leaved ash, *Fraxinus angustifolia*. Cicek, E., Yilmaz, F., and Yilmaz, M. Journal of Environmental Biology 31:109-114. 2010.

51. © Fall fertilization of *Pinus resinosa* seedlings: nutrient uptake, cold hardiness, and morphological development. Islam, M. A., Apostol, K. G., Jacombs, D. F., and Dumroese, R. K. Annals of Forest Science 66:704-712. 2009.

52. © Foliar iron-fertilisation of fruit trees: present knowledge and future perspectives - a review. Fernandez, V., Orera, I., Abadia, J., and Abadia, A. Journal of Horticultural Science & Biotechnology 84(1):1-6. 2009.

53. © Low soil temperature reduces the positive effects of high nutrient supply on the growth and biomass of white birch seedlings in ambient and elevated carbon dioxide concentrations. Amebebe, T. F., Dang, Q.-L., and Marfo, J. Botany 87:905-912. 2009.

54. Understanding plant nutrition: common high media-EC problems. Fisher, P. and Argo, B. Greenhouse Grower July 2009. 2009.

General and Miscellaneous



55. © Modelling the time course of shade, temperature, and wood recovery in streams with riparian forest restoration. Davies-Colley, R. J., Meleason, M. A., Hall, G. M. J., and Rutherford, J. C. New Zealand Journal of Marine and Freshwater Research 43:673-688. 2009.

56. Nursery worker turnover and language proficiency. Mathers, H. M., Acuna, A. A., Behe, B. K., and Hodges, A. W. HortScience 45(1):71-77. 2010.

57. © An outbreak of Pontiac fever due to *Legionella longbeachae* serogroup 2 found in potting mix in a horticultural nursery in New Zealand. Cramp, G. J., Harte, D., Douglas, N. M., and Graham, F. Epidemiology and Infection 138:15-20. 2010.

58. A picture is worth... Johnson, J. International Plant Propagators' Society, combined proceedings, 2008, 58:143-152. 2009.

59. Reforestation: challenges and themes in reforestation research (Chapter 1). Southworth, J. and Nagendra, H. IN: Reforesting landscapes: Linking pattern and process, p. 1-14. Nagendra, H. and Southworth, J., eds. Springer Science. 2010. This book integrates research findings from scientists working in a range of contexts and continents to examine reforestation. It is targeted to research community working on biophysical, geographic, socioeconomic, and institutional processes associated with reforestation.

60. Reforestation: conclusions and implications (Chapter 16). Nagendra, H. and Southworth, J. IN: Reforesting landscapes: Linking pattern and process, p. 357-367. Nagendra, H. and Southworth, J., eds. Springer Science. 2010.

61. © Reforestation planning using Bayesian networks. Galan, C. O., Matias, J. M., Rivas, T., and Bastante, F. G. Environmental Modelling and Software 24:1285-1292. 2009.

62. © Reforesting "Bare Hills" in Vietnam: social and environmental consequences of the 5 million hectare reforestation program. McElwee, P. Ambio 38(6):325-333. 2009.

Genetics and Tree Improvement



63. © Benefits of using genetically improved white spruce in Quebec: the forest landowner's viewpoint. Petrinovic, J. F., Gelinis, N., and Beaulieu, J. *Forestry Chronicle* 85(4):571-582. 2009.

64. © Genetic variation in seed size and germination patterns and their effect on white spruce seedling characteristics. Carles, S., Lamhamedi, M. S., Beaulieu, J., and Stowe, D. C. *Silvae Genetica* 58(4):152-161. 2009.

Mycorrhizae & Beneficial Microorganisms



65. © Molecular genetic tools to infer the origin of forest plants and wood. Finkeldey, R., Leinemann, L., and Gailing, O. *Applied Microbiology and Biotechnology* 85:1251-1258. 2010.

66. Infectivity and effectiveness of arbuscular mycorrhizal fungi in horticultural practices. Corkidi, L., Evans, M., and Bohn, J. *International Plant Propagators' Society, combined proceedings, 2008*, 58:241-244. 2009.

67. © Low *Frankia* inoculation potentials in primary successional sites at Mount St. Helens, Washington, USA. Seeds, J. D. and Bishop, J. G. *Plant and Soil* 323:225-233. 2009.

68. Opportunities from Down Under: How mycorrhizal fungi can benefit nursery propagation and production systems. Davies, F. T., Jr. *International Plant Propagators' Society, combined proceedings, 2008*, 58:539-548. 2009.

69. Truffles and oak selection. Fitzpatrick, N. *International Plant Propagators' Society, combined proceedings, 2008*, 58:91-93. 2009.

Nursery Structures & Equipment



Choosing the best biomass heating option. Sanford, S. *Greenhouse Management and Production* 29(11):20, 22-25. 2009.

Keratin biodegradable plastics. Tefteau, M. *American Nurseryman* 209(9):44-45. 2009.

Save fuel and electricity with energy/shade screens. Bartok, J. W., Jr. *Greenhouse Management and Production* 29(9):38-39. 2009.

Selecting a greenhouse curtain for better environmental control. Parhst, K. *Greenhouse Management and Production* 29(122):16, 18-20. 2009.

74. Upgrading the greenhouse water supply system. Bartok, J. W., Jr. *Greenhouse Management and Production* 29(12):30-31. 2009.

Outplanting Performance



75. Competitive interaction between *Microstegium vimineum* and first-year seedlings of three central hardwoods. Marshall, J. M., Buckley, D. S., and Franklin, J. A. *Journal of the Torrey Botanical Society* 136(3):342-349. 2009.

76. The effect of mining and vegetation scarification on the survival and establishment of *Pinus rotundata* Link. and *P. sylvestris* L. in contrasting peat bog habitats. Mach, J., Lanta, V., and Bastl, M. *Polish Journal of Ecology* 57(2):239-250. 2009.

77. © The effect of root and shoot pruning on early growth of hybrid poplars. DesRochers, A. and Tremblay, F. *Forest Ecology and Management* 258:2062-2067. 2009.

78. © The effect of soil compaction on germination and early growth of *Eucalyptus albens* and an exotic annual grass. Skinner, A. K., Lunt, I. D., Spooner, P., and McIntyre, S. *Austral Ecology* 34:698-704. 2009.

79. © Effects of nursery shading on seedling quality and post-planting performance in two Mediterranean species with contrasting shade tolerance. Puertolas, J., Benito, L. F., and Penuelas, J. L. *New Forests* 38:295-308. 2009.
80. © Fall planting and tree shelters for reforestation in the east Washington Cascades. Taylor, M., Haase, D. L., and Rose, R. L. *Western Journal of Applied Forestry* 24(4):173-179. 2009.
81. © Growth responses of *Salix gracilistyla* cuttings to a range of substrate moisture and oxygen availability. Nakai, A., Yurugi, Y., and Kisanuki, H. *Ecological Research* 24:1057-1065. 2009.
82. © Influence of ash substrate proximity on growth and survival of planted mixed-conifer seedlings. York, R. A., Thomas, Z., and Restaino, J. *Western Journal of Applied Forestry* 24(3):117-123. 2009.
83. © Integration of soil moisture, xylem water potential, and fall-spring herbicide treatments to achieve the maximum growth response in newly planted Douglas-fir seedlings. Dinger, E. J. and Rose, R. *Canadian Journal of Forest Research* 39:1401-1414. 2009.
84. The interaction of fertilization in nursery and field on survival, growth and the frost heaving of birch and spruce. Oskarsson, H. and Brynleyfsdottir, S. J. *Icelandic Agricultural Science* 22:59-68. 2009.
85. © Long-term biomass production and nutrient uptake of birch, alder and willow plantations on cut-away peatland. Hytonen, J. and Saarsalmi, A. *Biomass and Bioenergy* 33:1197-1211. 2009.
86. © Long-term effects of site preparation treatments, complete competition control, and repeated fertilization on growth of slash pine plantations in the flatwoods of the southeastern United States. Zhao, D., Kane, M., Borders, B., and Harrison, M. *Forest Science* 55(5):403-410. 2009.
87. © Long-term response of planted conifers, natural regeneration, and vegetation to harvesting, scalping, and weeding on a boreal mixedwood site. Man, R., Rice, J. A., and MacDonald, G. B. *Forest Ecology and Management* 258:1225-1234. 2009.
88. Nursery quality and first-year response of American chestnut (*Castanea dentata*) seedlings planted in the southeastern United States. Clark, S. L., Schweitzer, C. J., Schlarbaum, S. E., Dimov, L. D., and Hebard, F. V. *Tree Planters' Notes* 53(2):13-21. 2009.
89. Planting depth during container production and landscape establishment affects growth of *Ulmus parvifolia*. Bryan, D. L., Arnold, M. A., Voider, A., and Watson, W. T. *HortScience* 45(1):54-60. 2010.
90. © Reforestation after the Fountain fire in northern California: an untold success story. Zhang, J., Webseter, J., Powers, R. F., and Mills, J. *Journal of Forestry* 106(8):425-430. 2008.
91. © Stock type, subsoiling, and density impact productivity and land value of a droughty site. Blazier, M. A. and Dunn, M. *Southern Journal of Applied Forestry* 32(4):154-162. 2008.
92. © Survival and growth of drought hardened *Eucalyptus pilularis* Sm. seedlings and vegetative cuttings. Thomas, D. S. *New Forests* 38:245-259. 2009.
93. © Survival and growth of planted yellow-cedar seedlings and rooted cuttings (stecklings) near Ketchikan, Alaska. Hennon, P. E., McClellan, M. H., Spores, S. R., and Orlikowska, E. H. *Western Journal of Applied Forestry* 24(3):144-150. 2009.
94. A techno-economic evaluation of Bracke and M-Planter tree planting devices. Rantala, J., Harstela, P., Saarinen, V.-M., and Tervo, L. *Silva Fennica* 43(4):659-667. 2009.
95. Tree seedling survivorship, growth, and allocation in the Cross Timbers ecotone of Oklahoma, USA. Myster, R. W. *Plant Ecology* 205:193-199. 2009.
96. Using trees as a restoration tool in Tunisian arid areas: effects on understorey vegetation and soil nutrients. Jeddi, K. and Chaieb, M. *The Rangeland Journal* 31:377-384. 2009.
97. © Whitebark pine planting guidelines. McCaughey, W., Scott, G. L., and Izlar, K. L. *Western Journal of Applied Forestry* 24(3):163-166. 2009.



98. Effectiveness of *Trichoderma* spp. obtained from re-used soilless substrates against *Pythium ultimum* on cucumber seedlings. Liu, J. B., Gilardi, G., Gullino, M. L., and Garibaldi, A. *Journal of Plant Diseases and Protection* 116(4):156-163. 2009.
99. © Evaluation of several commercial biocontrol products on European and North American populations of *Phytophthora ramorum*. Elliott, M., Shamoun, S. E., Sumampong, G., and James, D. *Biocontrol Science and Technology* 19(1):1007-1021. 2009.

100. Experiments in liverwort management for nursery crops. Brennan, K. International Plant Propagators' Society, combined proceedings, 2008, 58:347. 2009.

101. Mode of action group 11: strobilurins. Chase, A. R. Greenhouse Management and Production 30(1):20, 22-24. 2010.

102. Mode of action group ml: copper. Chase, A. R. Greenhouse Management and Production 30(2):26, 28-29. 2010.

103. Moss control in a Proteaceae crop. Parlby, D. International Plant Propagators' Society, combined proceedings, 2008, 58:68-74. 2009.

104. The natural choice. Francis, J. American Nurseryman 210(1):36-39. 2010.

105. Nematodes: the good and the bad. Clark, B. American Nurseryman 209(12):8-9. 2009.

106. New chemical tools for control of plant diseases. Chase, A. R. International Plant Propagators' Society, combined proceedings, 2008, 58:245-247. 2009.


107. New control products for Botrytis blight. Hausbeck, M. Greenhouse Management and Production 29(8):42-43. 2009.

108. Resistance management of arthropod pests. Cloyd, R. A. and Cowles, R. S. American Nurseryman 209(10):36-39, 41-45. 2009.

109. Seed diseases and seedborne pathogens of North America. Cram, M. M. and Fraedrich, S. W. Tree Planters' Notes 53(2):35-44. 2009.

110. Vermicompost suppresses *Rhizoctonia solani* Kuhn in cucumber seedlings. Ersahin, Y. S., Haktanfi, K., and Yanar, Y. Journal of Plant Diseases and Protection 116(4):182-188. 2009.

111. Vertebrate control in container production. Coker, C. E. H. International Plant Propagators' Society, combined proceedings, 2008, 58:549-551. 2009.



112. Not all pesticide formulations are created equal. Capuzzi, G. Greenhouse Management and Production 30(2):30-31. 2010.

Seedling Harvesting and Storage



113. Late termination of freezer storage increases the risk of autumn frost damage to Norway spruce seedlings. Hanninen, H., Luoranen, J., Rikala, R., and Smolander, H. Silva Fennica 43(5):817-830. 2009.

Seedling Physiology and Morphology



114. © A comparison of root architecture and shoot morphology between naturally regenerated and container-grown seedlings of *Quercus ilex*. Tsakalidimi, M., Tsitsoni, T., Ganatsas, P., and Zagas, T. Plant and Soil 324:103-113. 2009.

115. © Effect of frost nights and day and night temperature during dormancy induction on frost hardiness, tolerance to cold storage and bud burst in seedlings of Norway spruce. Sogaard, G., Granhus, A., and Johnsen, O. Trees 23:1295-1307. 2009.

116. © The effect of mechanical stimulation on root and shoot development of young containerised *Quercus robur* and *Robinia pseudoacacia* trees. Reubens, R., Pannemans, B., Danjon, F., and De Proft, M. Trees 23:1213-1228. 2009.

117. © The effects of water, nutrient availability and their interaction on the growth, morphology and physiology of two poplar species. Yin, C. Pang X. and Chen, K. Environmental and Experimental Botany 67:196-203. 2009.

118. Root hydraulic conductance, gas exchange and leaf water potential in seedlings of *Pistacia lentiscus* L. and *Quercus suber* L. grown under different fertilization and light regimes. Hernandez, E. I., Vilagrosa, A., Luis, V. C., and Llorca, M. Environmental and Experimental Botany 67:269-276. 2009.


119. Root morphology and growth of bare-root seedlings of Oregon white oak. Gould, P. J. and Harington, C. A. Tree Planters' Notes 53(2):22-28. 2009.

120. © Seedling quality and field performance of commercial stocklots of containerized holm oak (*Quercus ilex*) in Mediterranean Spain: an approach for establishing a quality standard. del Campo, A. D., Navarro, R. M., and Ceacero, C. J. *New Forests* 39:19-37. 2010.

121. © Soil nitrogen limitation does not impact nighttime water loss in *Populus*. Howard, A. R. and Donovan, L. A. *Tree Physiology* 30:23-31. 2009.

122. © Soil salinity and drought alter wood density and vulnerability to xylem cavitation of baldcypress (*Taxodium distichum* (L.) Rich.) seedlings. Stiller, V. *Environmental and Experimental Botany* 67:164-171. 2009.

123. © Warm temperature accelerates short photoperiod-induced growth cessation and dormancy induction in hybrid poplar (*Populus* x spp.). Kalcsits, L. A., Silim, S., and Tanino, K. *Trees* 23:971-979. 2009.



124. A closer look at seed germination and dormancy. Geneve, R. L. and Dutt, M. *International Plant Propagators' Society, combined proceedings*, 2008, 58:561-565. 2009.

125. Creative germination techniques for difficult seeds. Krautman, M. *International Plant Propagators' Society, combined proceedings*, 2008, 58:280-283. 2009.

126. Environmental variation in seed and seedling characteristics of *Pinus roxburghii* Sarg. from Uttarakhnad, India. Ghildiyal, S. K., Sharma, C. M., and Gairola, S. *Applied Ecology and Environmental Research* 7(2):121-129. 2009.

127. © Identification of alkyl substituted 2H-Furo [2,3-clpyran-2-ones as germination stimulants present in smoke. Flematti, G. R., Ghisalberti, E. L., Dixon, K. W., and Trengove, R. D. *Journal of Agricultural and Food Chemistry* 57:9475-9480. 2009.

128. © Seed germination and seedling vigor of weevil-damaged acorns of red oak. Lombardo, J. A. and McCarthy, B. C. *Canadian Journal of Forest Research* 39:1600-1605. 2009.

129. Understanding lodgepole pine seed germination for improved utilization. El-Kassaby, Y. A., Kolotelo, D., and Reid, D. *Seed Science and Technology* 37:316-328. 2009.

130. © Variation in seed germination between populations of five sub-alpine woody species from eastern Qinghai-Tibet Plateau following dry storage at low temperatures. Want, J. H., Baskin, C. C., Chen, W., and Du, G. Z. *Ecological Research* 25:195-203. 2010.

**Soil Management
& Growing Media**

Peat Moss
Selected Canadian Sphagnum

131. Changes in physical properties of a pine tree substrate in containers over time. Jackson, B. E. and Wright, R. D. *International Plant Propagators' Society, combined proceedings*, 2008, 58:587-592. 2009.

132. © Compost and vermicompost as nursery pot components: effects on tomato plant growth and morphology. Lazcano, C., Arnold, J., Tato, A., Zaller, J. G., and Dominguez, J. *Spanish Journal of Agricultural Research* 7(4):944-951. 2009.

133. Composted poultry litter as an amendment for substrates with high wood content. Marble, S. C., Gilliam, C. H., Sibley, J. L., and Fain, G. B. *International Plant Propagators' Society, combined proceedings*, 2008, 58:524-532. 2009.

134. Container medium pH in a pine tree substrate amended with peatmoss and dolomitic limestone affects plant growth. Jackson, B. E., Wright, R. D., and Gruda, N. *HortScience* 44(7):1983-1987. 2009.

135. Developments in growing media policy and use in the U.K. Dawson, C. *International Plant Propagators' Society, combined proceedings*, 2008, 58:435-439. 2009.

136. © Effects of peat moss substitution with arboretum and greenhouse waste compost for use in container media. Olszewski, M. W., Trego, T. A., and Kuper, R. *Compost Science and Utilization* 17(3):151-157. 2009.

137. Field estimation of soil water content: a practical guide to methods, instrumentation and sensor technology. Heng, L. K. *International Atomic Energy Agency, Training Course Series No. 30*. 131 p. 2008.

138. A field method for soil erosion measurements in agricultural and natural lands. Hsieh, Y. P., Grant, K. T., and Bugna, G. C. *Journal of Soil and Water Conservation* 64(6):374-382. 2009.

139. Measuring air filled porosity for container substrates. Bilderback, T. *International Plant Propagators' Society, combined proceedings, 2008*, 58:129-131. 2009.

140. Methods of constructing a pine tree substrate from various wood particle sizes, organic amendments, and sand for desired physical properties and plant growth. Jackson, B. E., Wright, R. D., and Barnes, M. C. *HortScience* 45(1):103-112. 2010.

141. © Migration of heavy metals in soil as influenced by compost amendments. Farrell, M., Perkins, W. T., Hobbs, P. J., Griffith, G. W., and Jones, D. L. *Environmental Pollution* 158:55-64. 2010.

142. Physico-chemical changes in bauxite residue following application of spent mushroom compost and gypsum. Courtney, R. G., Jordan, S. N., and Harrington, T. *Land Degradation and Development* 20:572-581. 2009.

143. Research and development of potting media in Australia. Nichols, D. *International Plant Propagators' Society, combined proceedings, 2008*, 58:96-100. 2009.

144. A suitable substrate alternative. Jackson, B. E. *American Nurseryman* 209(9):28-30, 32-33. 2009.

145. Use of switchgrass as a nursery container substrate. Altland, J. E. and Krause, C. *HortScience* 44(7):1861-1865. 2009.



146. © Amending soils with hydrogels increases the biomass of nine tree species under non-water stress conditions. Orikiriza, L. J. B., Agaba, H., Tweheyo, M., and Ellu, G. *Clean* 37(8):615-620. 2009.

147. © Love of nurse plants is not enough for restoring oak forests in a seasonally dry tropical environment. Badano, E. I., Perez, D., and Vergara, C. H. *Restoration Ecology* 17(5):571-576. 2009.

148. © Physiological and morphological responses of young mahogany (*Swietenia macrophylla* King) plants to drought. Cordeiro, Y. E. M., Pinheiro, H. A.,

dos Santos Filho, B. G., and Correa, S. S. *Forest Ecology and Management* 258:1449-1455. 2009.

149. Potential of the seedling community of a forest fragment for tropical forest restoration. Viani, R. A. G. and Rodrigues, R. R. *Scientia Agricola (Piracicaba, Braz.)* 66(6):772-779. 2009.

150. © Predicting field performance of five irrigated tree species using seedling quality assessment in Burkina Faso, West Africa. Bayala, J., Dianda, M., Wison, J., Ouedraogo, S. J., and Sanon, K. *New Forests* 38:309-322. 2009.



151. © Evaluation of fall versus spring dormant planting of hardwood willow cuttings with and without soaking treatment. Tilley, D. J. and Hoag, J. C. *Native Plants Journal* 10(3):288-294. 2009.

152. Hybrid pines: opportunities for life on the edge. Hargreaves, C., Sibley, M., Menzies, M., Dungey, H., and van Ballekom, S. *International Plant Propagators' Society, combined proceedings, 2008*, 58:162-167. 2009.

153. Lightweight aggregate HydRocks: a novel approach to rooting and bare root cuttings. Pickens, J. M., Sibley, J. L., Gilliam, C. H., and Fain, G. B. *International Plant Propagators' Society, combined proceedings, 2008*, 58:573-577. 2009.

154. Propagate plants from cuttings using dry-dip rooting powders and water-based rooting solutions. Kroin, J. *International Plant Propagators' Society, combined proceedings, 2008*, 58:360-372. 11682.

155. © Seasonal fluctuation of root carbohydrates in hybrid aspen clones and its relationship to the sprouting efficiency of root cuttings. Stenvall, N., Piisila, M., and Pulkkinen, P. *Canadian Journal of Forest Research* 39:1531-1537. 2009.

156. Vegetative propagation of old growth conifers. Werner, B. *International Plant Propagators' Society, combined proceedings, 2008*, 58:317-322. 2009.

Water Management



- 157. Container-grown ornamental plant growth and water runoff nutrient content and volume under four irrigation treatments.** Warsaw, A. L., Fernandez, R. T., Cregg, B. M., and Andresen, J. A. HortScience 44(6):1573-1580. 2009.
- 158. Control of pathogens in irrigation water using chlorine without injury to plants.** Zheng, Y., Cayanan, D. F., and Dixon, M. International Plant Propagators' Society, combined proceedings, 2008, 58:248-249. 2009.
- 159. Is it time to water? Wireless soil moisture monitors provide the answer.** Davies, M. A. and Etter, T. R. USDA Forest Service, Technology and Development Program, Tech Tips, 0924-2316. 6 p. 2009.
- 160. Nurseries surveyed in southern California adopt best practices for water quality.** Mangiafico, S. S., Newman, J., Mochizuki, M., and Zurawski, J. California Agriculture 64(1):26-30. 2010.
- 161. © Prospective use of collected fog water in the restoration of degraded burned areas under dry Mediterranean conditions.** Estrela, M. J., Valienta, J. A., Core11, D., Fuentes, D., and Valdecantos, A. Agricultural and Forest Meteorology 149:1896-1906. 2009.
- 162. Reclaimed water start to finish.** Yeager, T. and Larsen, C. International Plant Propagators' Society, combined proceedings, 2008, 58:503-507. 2009.
- 163. Recycled irrigation water chlorination and pathogen prevention.** Black, R. International Plant Propagators' Society, combined proceedings, 2008, 58:508-511. 2009.
- 164. Soil erosion and runoff prevention by plant covers: a review.** Zuazo, V. H. D. and Pieguezuelo, C. R. R. IN: Sustainable Agriculture, p. 785-811. E. Eichtfous, et al, eds. 2009.
- 165. Using wireless sensor technology to schedule irrigations and minimize water use in nursery and greenhouse production systems.** Lea-Cox, J. D., Ristvey, A. G., and Kantor, G. F. International Plant Propagators' Society, combined proceedings, 2008, 58:512-518. 2009.
- 166. Water: what is on the horizon and knowing what to grow.** Adams, B. International Plant Propagators' Society, combined proceedings, 2008, 58:498-500. 2009.

Weed Control



- 167. Can Roundup be safely used over the top of nursery crops?** Van Hoogmoed, A. J., Gilliam, C. H., Wehtje, G. R., and Olive, J. W. International Plant Propagators' Society, combined proceedings, 2008, 58:568-572. 2009.
- 168. Efficacy comparison of some new natural-product herbicides for weed control at two growth stages.** Abouziena, H. F. H., Omar, A. A. M., Sharma, S. D., and Singh, M. Weed Technology 23:431-437. 2009.
- 169. Herbicide-resistant weeds in the United States and their impact on Extension.** Scott, B. A., Vangessel, M. J., and White-Hansen, S. Weed Technology 23:599-603. 2009.
- 170. Managing weeds in the greenhouse.** Smith, T. Greenhouse Management and Production 29(12):32-33. 2009.
- 171. © Results from six *Pinus taeda* nursery trials with the herbicide pendimethalin in the USA.** South, D. B. and Hill, T. E. Southern Forests 71(3):179-185. 2009.
- 172. © Twenty-year response of ponderosa pine (*Pinus ponderosa* Laws.) to treatment with hexazinone in northeastern Oregon.** Lindsay, A., Oester, P., and Cole, E. Western Journal of Applied Forestry 24(3):151-156. 2009.
- 173. Using preemergence herbicides in containerized rootstock during grafting.** LeBude, A. V., Upchurch, B. L., and Neal, J. C. International Plant Propagators' Society, combined proceedings, 2008, 58:208-212. 2009.
- 174. Weed response to flame weeding at different developmental stages.** Sivesind, E. C., Leblanc, M. L., Cloutier, D. C., Seguin, P., and Stewart, K. A. Weed Technology 23:438-443. 2009.

Literature Order and Mailing List Update Form Winter 2010

Please fill out a separate order form for each person receiving FNN.
For items that require a copyright fee, you will receive the title page with abstract and ordering instructions if you want the entire article. Fax or mail this form to:

Forest Nursery Notes
J.H. Stone Nursery
2606 Old Stage Rd.
Central Point, OR 97502

TEL: 541.858.6166

FAX: 541.858.61 10

[E-mail: rewatson@fs.fed.us](mailto:rewatson@fs.fed.us)

Name: Position:

Department: Nursery or Company:

Mailing address:

Street Address:

City: State or Province:

Country: Zip or Postal Code:

Telephone: FAX:

E-mail: Website:

☐

= Yes, please send me a CD with all the aarticles in the New Nursery Literature Section

☐

= Yes, please keep me listed on the FNN mailing list.

Contact Information for Reforestation, Nurseries, and Genetic Resources (RNGR) Team http://www.rngr.net		
Technology Transfer Services	Area of Responsibility	Who to Contact
• National Nursery Specialist • Forest Nursery Notes • Container Tree Nursery Manual • Proceedings of Nursery Meetings • Native Plants Journal	US and International	Kas Dumroese USDA Forest Service 1221 S. Main Street Moscow, ID 83843 TEL: 208.883.2324 • FAX: 208.883.2318 E-Mail: kdumroese@fs.fed.us
• Technical Assistance about Forest, Conservation, and Native Plant Nurseries	Western US	Diane L. Haase USDA Forest Service PO Box 3623 Portland, OR 97208 TEL: 503.808.2349 • FAX: 503.808.2339 E-Mail: dlhaase@fs.fed.us
• Technical Assistance about Tree Improvement and Genetic Resources • Technical Assistance about Forest and Conservation Nurseries	Southeastern US and International	George Hernandez USDA Forest Service 1720 Peachtree Road NW Atlanta, GA 30367 TEL: 404.347.3554 • FAX: 404.347.2776 E-Mail: ghernandez@fs.fed.us
• Technical Assistance about Tree and Shrub Seed	US and International	Bob Karrfalt National Seed Laboratory 5675 Riggins Mill Road Dry Branch, GA 31020 TEL: 478.751.4134 • FAX: 478.751.4135 E-Mail: rkarrfalt@fs.fed.us
• Technical Assistance about Tree Improvement and Genetic Resources • Technical Assistance about Forest, Conservation, and Native Plant Nurseries • Tree Planters' Notes	Northeastern US and International	Ron Overton Regeneration Specialist USDA Forest Service Purdue University 715 West State Street West Lafayette, IN 47907 TEL: 765.496.6417 • FAX: 765.494.9461 E-Mail: roverton@fs.fed.us
• Technical Assistance to Native Americans regarding Nurseries, Reforestation, and Restoration	US and International	Jeremy Pinto USDA Forest Service 1221 S. Main Street Moscow, ID 83843 TEL: 208.883.2352 • FAX: 208.883.2318 E-Mail: jpinto@fs.fed.us

