

THE 3 Ps OF OAK REGENERATION: PLANNING, PERSISTENCE, AND PATIENCE

Dale R. Weigel, Daniel C. Dey, and John Kabrick

Oak regeneration research in the United States has been ongoing in earnest since the late 1950s. Most research has focused on specific silvicultural practices, regeneration processes, site characteristics, and local limiting factors such as deer browsing or interfering species. Research has evaluated the effects of thinning on regeneration development, methods for oak planting, post-harvest treatments to control competing vegetation, and many other aspects of oak silviculture. All of these have provided solutions to individual problems in oak regeneration for local to regional areas.

However, with all this research we still have difficulty regenerating oak forests. One question remains “How do we insure that oaks are present at desired levels in the next stand following harvest?” We believe the answer is more a managerial problem than biological. The long-term and more universal solution is based on the 3Ps of oak regeneration: planning, persistence, and patience. Because these three steps are not consistently followed nor their importance recognized, oak regeneration often fails.

Research and operational silviculture have been focused on the application of one or several treatments over a short period of years. Oak regeneration is a long-term ecological process requiring long-term planning. Two important questions that must be answered in the planning process are: when do you want to regenerate, and where or which stands do you want to regenerate to oak? It is necessary for oak advanced regeneration (OAR) to be present before harvest for oaks to have a chance of developing in the next stand (Sander and others 1976). OAR is increased through acorn germination. Unfortunately acorn crops are sporadic and unpredictable (Beck 1977, Dey 1995, Godman and Mattson 1976). Planting can supplement OAR in order to decrease the time necessary to develop sufficient regeneration. Planting research has been completed across the entire eastern hardwood region (Dey and Parker 1997, Johnson and others 1986, Spetich and others 2009, Weigel and Johnson 2000).

Persistence in treatments is required both pre-harvest to enable oak regeneration to develop and post-harvest to

keep oak regeneration competitive (Carvell and Tryon 1961). Repeated treatments may be required to maintain increased light levels in the lower canopy and shrub layer. These treatments can include herbicide, mechanical, and prescribed fire. Fire has been present on the landscape dating back to at least the 1600s (Guyette and Dey 2006). The use of prescribed fire has been shown to benefit oak regeneration (Brose and others 2006). Post-harvest thinning and crop tree release are necessary to keep oak competitive (Perky and Wilkins 1993, Schuler and Miller 1999).

Because oak is a species physiologically adapted to repeated disturbances over decades, patience in the regeneration process is necessary. Oak’s growth habit of favoring early root growth over shoot growth helps oaks persist through repeated disturbances better than competitors (Johnson and others 2009, fig. 10.1). But the limited shoot elongation puts it at a competitive disadvantage with other species in the absence of disturbances such as fire and drought.

By completing these three steps; planning, persistence, and patience, oak regeneration can be accomplished.

LITERATURE CITED

- Beck, D.E.** 1977. Twelve-year acorn yield in southern Appalachian oaks. Res. Note SE-244. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 8 p.
- Brose, P.H.,** Schuler, T.M., and Ward, J.S. 2006. Responses of oak and other hardwood regeneration to prescribed fire: what we know as of 2005. In: Dickinson, M.B., ed. Fire in Eastern oak forests: delivering science to land managers. Gen. Tech. Rep. NRS-P-1. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station: 123-135.
- Carvell, K.L.** and Tryon, E.H. 1961. The effect of environmental factors on the abundance of oak regeneration beneath mature oak stands. Forest Science 7: 98-105.
- Dey, D.C.** 1995. Acorn production in red oak. For. Res. Info. Pap. No. 127. Sault Ste. Marie, ON: Ontario Ministry of Natural Resources, Ontario Forest Research Institute. 22 p.

-
- Dey, D.C.** and Parker, W.C. 1996. Regeneration of red oak (*Quercus rubra* L.) using shelterwood systems: ecophysiology, silviculture and management recommendations. For. Res. Info. Pap. No. 126. Sault Ste. Marie, ON: Ontario Ministry of Natural Resources, Ontario Forest Research Institute. 59 p.
- Godman, R.M.** and Mattson, G.A. 1976. Seed crops and regeneration problems of 19 species in northeastern Wisconsin. Res. Pap. NC-123. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 5 p.
- Guyette, R.P.**, Dey, D.C., Stambaugh, M.C., and Muzika, R.M. 2006. Fire scars reveal variability of Eastern fire regimes. In: Dickinson, M.B., ed. Fire in Eastern oak forests: delivering science to land managers. Gen. Tech. Rep. NRS-P-1. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station: 20-39.
- Johnson, P.S.**, Dale, C.D., Davidson, K.R., and Law, J.R. 1986. Planting northern red oak in the Missouri Ozarks: a prescription. Northern Journal of Applied Forestry 3: 66-68.
- Johnson, P.S.**, Shifley, S.R., and Rogers, R. 2009. The ecology and silviculture of oaks 2nd ed. Cambridge, MA: CABI.580 p.
- Perkey, A.W.** and Wilkins, B.L. 1993. Crop tree management in eastern hardwoods. Tech. Pap. NA-TP. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry. 19-23.
- Sander, I.L.**, Johnson, P.S., and Watt, R.F. 1976. A guide for evaluating the adequacy of oak advance reproduction. Gen. Tech. Rep. NC-23. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 7 p.
- Schuler, T.M.** and Miller, G.W. 1999. Releasing sheltered northern red oak during the early stem exclusion stage. In: Stringer, J.W. and Loftis, D.L. eds. Proceedings 12th central hardwood forest conference. Gen. Tech. Rep. SRS-24. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station: 195-201.
- Spetich, M.A.**, Dey, D.C., and Johnson, P.S. 2009. Shelterwood-planted northern red oaks: integrated costs and options. Southern Journal of Applied Forestry 33: 182-186.
- Weigel, D.R.** and Johnson, P.S. 2000. Planting red oak under oak/yellow-poplar shelterwoods: a provisional prescription. Gen. Tech. Rep. NC-210. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Research Station. 16 p.