

# Economic analysis of prescribed burning for wildfire management in Western Australia<sup>1</sup>

Veronique Florec<sup>2</sup>, David Pannell<sup>3</sup>, Michael Burton<sup>4</sup>, Joel Kelso<sup>5</sup>, Drew Mellor<sup>6</sup>, and George Milne<sup>7</sup>

## Abstract

Wildfires can cause significant damage to ecosystems, life and property, and wildfire events that do not involve people and property are becoming rare. With the expansion of the rural–urban interface in Western Australia and elsewhere, objectives of life and property protection become more difficult to achieve. We applied the cost plus net value change (C+NVC) model to a synthetic landscape, representative of the northern jarrah forest of the south west of Western Australia. The most economically efficient level of prescribed burning corresponds to a strategy where 5% of the simulated landscape is prescribed-burned per year. Our results are sensitive to changes in the average cost per hectare of prescribed burning, the probabilities of fire occurrence, urban area values (in average dollars per hectare) and suppression costs.

*Keywords: wildfire, fire management, economic analysis, cost plus net value change.*

## Introduction

Wildfires can cause significant damage to ecosystems, life and property and in many parts of the world the frequency of large, disastrous fires, often referred to as mega-fires, appears to have increased (Morgan 2009, Williams *and others* 2011). Catastrophic wildfires have occurred in Australia, the United States, Canada, Russia, China, South Africa, and Spain, among others. The Black Saturday fires in Victoria,

---

<sup>1</sup> An abbreviated version of this paper was presented at the IV International Symposium on Fire Economics, Planning, and Policy: Climate Change and Wildfires, November 5-11, 2012, Mexico City, Mexico.

<sup>2</sup> PhD candidate, School of Agricultural and Resource Economics, The University of Western Australia, Crawley, Australia (email: veronique.florec@uwa.edu.au)

<sup>3</sup> Winthrop Professor, School of Agricultural and Resource Economics, Director, Centre for Environmental Economics and Policy, The University of Western Australia, Crawley, Australia.

<sup>4</sup> Professor, School of Agricultural and Resource Economics, The University of Western Australia, Crawley, Australia.

<sup>5</sup> Post-doctoral researcher, School of Computer Science and Software Engineering, The University of Western Australia, Crawley, Australia.

<sup>6</sup> Post-doctoral researcher, School of Computer Science and Software Engineering, The University of Western Australia, Crawley, Australia.

<sup>7</sup> Winthrop Professor, School of Computer Science and Software Engineering, The University of Western Australia, Crawley, Australia.

Australia on 7 February 2009 resulted in the highest loss of life and property from a wildfire in Australian history (Teague and others 2010), causing the death of 173 people and widespread destruction of assets and infrastructure. Since 1998, nine states in the US have suffered their worst wildfires in history (Williams and others 2011). Although these fires are usually rare, they cause profound and long-lasting social, economic and environmental impacts where they occur (Handmer and Proudley 2008).

Urban development in fire prone areas has amplified the complexity of the problem. Wildfire events that do not involve people and property are becoming rare (Mutch and others 2011). The number of houses and infrastructure located within or close to areas of high fire risk continues to increase in Australia (Morgan and others 2007) and elsewhere (USA, Canada, Europe, among others, see Smalley 2003, Mozumder and others 2009, Mell and others 2010, Stockmann and others 2010, Mutch and others 2011), increasing the wildfire risk to life and property.

In light of the increasing wildfire threat, fire agencies have often responded with greater suppression capacity, involving increasing suppression costs. However, this has not solved the problem of catastrophic wildfires (Williams and others 2011). In Australia suppression expenditures have followed an escalating trend and the country “currently runs the risk of spending ever-greater amounts of money on wildfire suppression, while becoming even less successful in its management of fire in the landscape than is currently the case.” (Morgan and others 2007, p. 1). Other countries with fire-prone landscapes appear to face similar issues. In the US annual suppression expenditures have increased remarkably over the past several years while the western part of the country has been severely affected by large and intense wildfires since the 1980s (Calkin and others 2005).

Economics can provide improved understanding and comprehensive appraisals of wildfire costs and benefits in order to devise wildfire mitigation and management programs that optimally allocate resources and express informed, evidence-based judgements about trade-offs between available options (Handmer and Proudley 2008). However, the use of economics in the wildfire literature is still relatively limited. Despite the abundance of theoretical studies on the subject, empirical economic analyses of wildfire management are scarce (Mercer and others 2007).

In this paper we apply the cost plus net value change (C+NVC) model to a synthetic landscape, representative of the northern jarrah forest of the south west of Western Australia (WA). The purpose of the study is to determine the most economically efficient pre-suppression strategy for the synthetic landscape and evaluate which parameters significantly affect the results. We focus on prescribed burning as the main pre-suppression strategy. The primary objective of this model is to provide preliminary results which may inform the development of a more

complete model based on actual areas of WA.

## Methods

We simulated wildfires in a synthetic landscape under varying climatic conditions and different prescribed burning (pre-suppression) strategies using the AUSTRALIS Wildfire Simulator, which was developed at the School of Computer Science and Software Engineering, The University of Western Australia (see Johnston and others 2008 for a description of the fire simulator). The synthetic landscape generated for the simulations is a square landscape of 100,000 ha, containing a flat terrain with homogenous northern jarrah forest fuel.

We tested three prescribed burning strategies, each with three patch sizes. The strategies involved prescribed-burning 5, 10 or 20 % of the total area per year, corresponding to rotation cycles of 20, 10 and 5 years respectively. The landscape was partitioned into square-shaped patches of 50, 500 and 4000 ha, and each strategy could be carried out in burning patches of these sizes. The age of the fuel in each patch was a random integer value from  $[0, n]$ , where  $n = 5, 10$  or  $20$ , depending on the burning strategy. In addition to these 9 combinations ( $3 \text{ strategies} \times 3 \text{ patch sizes} = 9$ ), we defined a baseline strategy for comparison where the fuel age was uniformly set at 15 years across the entire treatment area and used this baseline as the 0% prescribed burning or no-strategy case. Each strategy-patch size combination was simulated under high, very high, extreme and catastrophic forest fire danger conditions (FFDI), giving  $(9+1) \times 4 = 40$  scenarios. Finally, each scenario was tested under 30 random ignitions, making a total of  $30 \times 40 = 1200$  simulations.

We used the “McArthur Mk V” forest fire meter (Noble *and others* 1980; Sirakoff 1985) to determine the rate of spread. Fuel load was determined using the fuel accumulation table for Jarrah forest in Sneeuwjagt and Peet (1998), which gives fuel load as a function of fuel age. Fire ignition points were generated according to a random uniform distribution. The weather was constant during simulation. Spotting effects were not modelled. **Error! Reference source not found.** summarises the simulator settings. It is assumed that if fire intensity is below a threshold of 2,000 kW/m, then the fire is suppressed.

We used the C+NVC model, which is currently the most commonly accepted model for economic evaluations of wildfire management programs (Ganewatta 2008, Gebert and others 2008). From Donovan and Rideout (2003), the C+NVC model can be expressed as:

$$\text{Min } C + NVC = W^P P + W^S S(P) + NVC(P, S(P)) \quad (1)$$

in which  $W^P$  is the price of pre-suppression;  $P$  is the pre-suppression effort;  $W^S$  is

the price of suppression;  $S$  is the suppression effort, which is dependent on pre-suppression; and  $NVC$  is the net fire damage (fire damage less fire benefit).

**Table 1—** Summary of simulator settings for prescribed burning experiments

|                                    | Value                                                 |                          |                   |                      |                   |
|------------------------------------|-------------------------------------------------------|--------------------------|-------------------|----------------------|-------------------|
| Scenario parameters                |                                                       |                          |                   |                      |                   |
| Patch sizes                        | 50, 500, 4000 ha                                      |                          |                   |                      |                   |
| Rotation cycles                    | 5, 10, 20 y                                           |                          |                   |                      |                   |
| Weather conditions                 | High, Very High, Extreme, and Catastrophic            |                          |                   |                      |                   |
| Ignition points                    | Uniformly randomly placed across the entire landscape |                          |                   |                      |                   |
|                                    |                                                       |                          |                   |                      |                   |
| Weather conditions                 | Temperature<br>(°Celsius)                             | Relative<br>Humidity (%) | Wind<br>direction | Wind speed<br>(km/h) | Drought<br>factor |
| High                               | 30                                                    | 25                       | North             | 30                   | 5                 |
| Very High                          | 35                                                    | 20                       | North             | 30                   | 7                 |
| Extreme                            | 35                                                    | 10                       | North             | 30                   | 9                 |
| Catastrophic                       | 40                                                    | 10                       | North             | 50                   | 10                |
|                                    |                                                       |                          |                   |                      |                   |
| Fuel                               |                                                       |                          |                   |                      |                   |
| Rate of spread meter               | McArthur Mk V, Forest (Sneeuwjagt and Peet 1998)      |                          |                   |                      |                   |
| Fuel accumulation rules            | Northern Jarrah Fuel                                  |                          |                   |                      |                   |
| Canopy cover                       | 60%                                                   |                          |                   |                      |                   |
|                                    |                                                       |                          |                   |                      |                   |
| Topography                         | Flat                                                  |                          |                   |                      |                   |
|                                    |                                                       |                          |                   |                      |                   |
| Cell grid                          |                                                       |                          |                   |                      |                   |
| Cell spacing                       | 50 m                                                  |                          |                   |                      |                   |
| Cell neighbourhood                 | Cells up to 6 links away are considered adjacent      |                          |                   |                      |                   |
|                                    |                                                       |                          |                   |                      |                   |
| Simulator configuration            |                                                       |                          |                   |                      |                   |
| Lateral rate of spread             | Rate of spread at zero wind speed                     |                          |                   |                      |                   |
| Maximum duration of simulated time | 36 h                                                  |                          |                   |                      |                   |

In our model, we assumed a negative relationship between prescribed burning effort (annual prescribed-burned area as a proportion of the entire landscape) and suppression costs. Prescribed burning is generally expected to improve directly the probability of successful suppression (Fernandes and Botelho 2003) because prescribed burning decreases the intensity with which wildfires burn. Preliminary results of both empirical and modelling studies suggest that the relationship between the percentage of landscape prescribed-burned and the probability of unplanned high intensity fire at a point may be represented by a complex multiplicative model with a convex shape (Cary and others 2003). Hence, as the intensity of fires increases, suppression becomes more difficult and more costly (Chatto and Tolhurst 2004), since more expensive resources such as water bombers are needed when direct attack methods can no longer be used.

The functional relationship between prescribed burning effort and suppression

costs is expressed as  $s = ke^{-ax}$ , where  $s$  is suppression expenditure;  $x$  is the proportion of area prescribed-burned;  $k$  represents the maximum suppression expenditure and  $a$  is a coefficient of prescribed burning effectiveness that affects the marginal benefit of each extra prescribed-burned hectare in the landscape. *Ceteris paribus*, the higher coefficient  $a$  is, the lower the expenditure on suppression for a given proportion of prescribed-burned area.

Since we simulated fires under different weather conditions, which have different probabilities of occurrence, we multiplied the outcomes of the fires by their probabilities of occurrence and assumed that under low and moderate weather conditions the fires would be suppressed relatively quickly and have only a minimal effect on the results. **Error! Reference source not found.** shows the probabilities that we used for fire occurrence.

**Table 2—Probabilities of fire occurrence**

| <i>Fire category</i> | <i>Probability of incident occurrence per year</i> |
|----------------------|----------------------------------------------------|
| Catastrophic         | 0.0001770                                          |
| Extreme              | 0.0002360                                          |
| Very High            | 0.0007080                                          |
| High                 | 0.0475789                                          |

We assumed in our analysis that a small town of 1,500 ha is located within the synthetic landscape in order to approach our evaluation of a rural-urban interface scenario. Thus, we included values of urban structures and public infrastructure and smoke and fire-related (prescribed or wildfire) health costs. Although the economic literature on human health impacts from exposure to wildfire or prescribed burning smoke is scarce<sup>8</sup> and the differences in estimates between available studies is large, omitting health impacts in economic evaluations of fire management programs could result in underinvestment in pre-suppression activities (Richardson and others 2012).

In our analysis, we have not accounted for number of days of exposure to the smoke caused by wildfires but linked intensity and area burned to health costs. We assumed an exponential relationship between area burned by wildfire and health costs, which changes with the level of intensity. As intensity and area burned by wildfires increase, fuel combustion and biomass burning emissions (such as particulate matter with an aerodynamic diameter less than 2.5  $\mu\text{m}$ , PM<sub>2.5</sub>) escalate, and ultimately, as they become greater, wildfires may cause serious injuries and

<sup>8</sup> Some studies that have estimated the adverse health effects from prescribed burning or bushfires smoke exposure include Richardson *and others* (2012), Butry and others (2001), Martin and others (2007), Rittmaster and others (2006). For a comprehensive review of the literature analysing economic cost of smoke-related health effects see Kochi and others (2010).

casualties. Hence we use an exponential relationship that increases as area burned and wildfire intensity become extreme. The relationship can be expressed as  $h = be^{c\omega}$ , where  $h$  is health costs per person at risk;  $\omega$  is the size of the simulated fires (area burned);  $b$  is a coefficient that integrates the effects of intensity on health costs and  $a$  is a coefficient that reflects the impact of area burned on health costs per capita, both coefficients determine the marginal cost of each extra hectare burned by wildfires at a particular level of intensity in the landscape.

Some fires however, can be relatively small and still cause an increase in  $PM_{2.5}$  concentration levels in a neighbouring town. We have then to assume that a smoke plume of a rectangular shape is released by the fire and has a probability of reaching the town  $P(m, A) = m(\omega)/A$ , with  $m$  the size of the smoke plume, which is dependent on the size of the fires; and  $A$  the size of the whole landscape.

## Results and Discussion

Our results indicate that the most efficient level of prescribed burning corresponds to a strategy where 5% of the simulated landscape is prescribed-burned per year. With our assumed costs of prescribed burning (\$/ha), this is equivalent to an annual investment in prescribed burning of approximately AU\$405,000 over an area of 100,000 ha. The minimum of the C+NVC curve equals about AU\$785,000. **Error! Reference source not found.** show the curves obtained from the C+NVC model.





Figure 2—The cost plus net value change (C+NVC) curve


We conducted a sensitivity analysis to evaluate the robustness of the results and determine which parameter values most affect the results. **Error! Reference source not found.** shows the change in the minimum of the costs plus net value change curve in dollars when some parameter values are reduced by 50% or increased by 50%. The parameters shown are those to which the results are most sensitive. As shown in **Error! Reference source not found.**, a change in prescribed burning costs (in average dollars per hectare) greatly affects the results.

As prescribed burning costs increase, they quickly become a large proportion of the C+NVC curve. If the slope of prescribed burning costs changes, the point where marginal costs equal marginal benefits shifts accordingly. If the average cost per hectare for prescribed burning is very high, then the minimum of the C+NVC curve corresponds to a strategy of 0% prescribed burned area. This case is illustrated in **Figure 2**—Change in the minimum of the costs plus net value change curve with an increase and a decrease of 50% in the value of selected parameters

. The minimum value of the C+NVC curve is then the sum of suppression costs and damages for the 0% strategy.




**Figure 2**—Change in the minimum of the costs plus net value change curve with an increase and a decrease of 50% in the value of selected parameters



**Figure 3—50% increase in average prescribed burning costs per hectare**

When the average prescribed burning cost per hectare is reduced by 50%, the minimum value of the C+NVC curve decreases by 34% (**Error! Reference source not found.**) and the most efficient strategy is between 5 and 10% area prescribed burned per year (**Error! Reference source not found.**). Comparing the three figures of the C+NVC curve, it can be seen that in our initial estimation () there is a wide range of prescribed burning levels that is near-optimal, but this is not the case when the cost of prescribed burning is modified (**Error! Reference source not found.** and **Error! Reference source not found.**).





**Figure 4—50% decrease in average prescribed burning costs per hectare**

Our results are also sensitive to changes in the probabilities of fire occurrence, urban area values (in average dollars per hectare) and suppression costs. **Error! Reference source not found.** shows how the most efficient prescribed burning strategy changes when we increase or reduce the value of the selected parameters described above.

**Table 3—Change in the most efficient prescribed burning strategy**

|                                       | <i>Most efficient prescribed burning strategy (% of landscape prescribed burned)</i> |               |              |
|---------------------------------------|--------------------------------------------------------------------------------------|---------------|--------------|
|                                       | Initial estimation                                                                   | 50% reduction | 50% increase |
| Prescribed burning costs (\$/ha)      | 5%                                                                                   | >5% and <10%  | 0%           |
| Probabilities type of fire occurrence | 5%                                                                                   | 0%            | >5% and <10% |
| Urban area value (\$/ha)              | 5%                                                                                   | 0%            | >5% and <10% |
| Suppression costs                     | 5%                                                                                   | 0%            | >5% and <10% |

This encapsulates the main challenges faced by fire agencies in developing sustainable fire management practices: climate change, the rural-urban interface and the effectiveness of suppression. Numerous uncertainties still exist regarding the behaviour of wildfires under severe weather conditions and the sustainability of different fire management practices in the context of climate change (Thornton 2010). With the expansion of the rural-urban interface, objectives of life and property

protection become more difficult to achieve. Smalley (2003) identifies the expansion of the rural-urban interface as “one of the three major factors that will propagate the pressures of the interface on communities. The other two are unusually severe weather events (from prolonged drought to severe heating periods and floods that erode soils and vegetation) and inadequate infrastructure due to the rapidity of growth or aging.” (p. 5)

The patch size of the prescribed burns had no significant effect on the results of the analysis. The severity measures obtained from the simulated fires were not significantly affected by changes in patch size (which is consistent with previous studies that use simulation to examine the efficacy of prescribed burning, e.g. Finney and others 2007, King and others 2008) and hence they had no significant impact on the economic analysis.

Although the C+NVC model has been extensively used to assess annual investments in fire prevention and protection, the model has been through some reformulation in the past two decades. Rideout and colleagues (Rideout and Omi 1990, Donovan and others 1999, Hesseln and Rideout 1999, Donovan and Rideout 2003, Rideout and Ziesler 2008) identified three inherent errors in the model that have been perpetuated from the earlier least cost plus loss model formulated by Sparhawk (1925). First, suppression expenditure is determined solely as a function of fire occurrence. Donovan and Rideout (2003) argued that both pre-suppression and suppression should be modelled as endogenous decision variables, with the benefits of suppression depending on the level of pre-suppression. As commonly applied, the C+NVC minimum may differ from the one obtained when pre-suppression and suppression are modelled correctly.

Second, suppression and pre-suppression expenditures are incorrectly modelled as negatively correlated (Donovan and Rideout 2003, Rideout and Ziesler 2008). And third, by analysing a single fire season, the long term effects of natural fuel accumulation processes, unplanned fires, fuel treatments and land management strategies on wildfire risk are overlooked (Hesseln and Rideout 1999). Fire scars, whether caused by fuel reduction treatments or unplanned fires, affect fire behaviour and intensity for longer than one year. Likewise, natural fuels accumulate over time and if left undisturbed, they gradually increase the risk of catastrophic fires year by year. Despite the limitations mentioned above, the application of the C+NVC model in its current formulation can help fire managers identify potential benefits and costs of different fire management options for a given year, even if a global minimum is not obtained (Rodriguez y Silva and Gonzalez-Caban 2010). We recognise the limitations of the model and their implications, and hope to address them in future work. The C+NVC framework has been used here as a first step towards a more comprehensive analysis.

## Summary

We applied the cost plus net value change (C+NVC) model to a synthetic landscape of 100,000 ha, representative of the northern jarrah forest of the south west of Western Australia. We used the AUSTRALIS Wildfire Simulator to simulate wildfires in this landscape under varying climatic conditions. We tested three different prescribed burning (pre-suppression) strategies and a no-strategy option. We found that the most economically efficient level of prescribed burning corresponds to a strategy where 5% of the simulated landscape is prescribed-burned per year over an area of 100,000 ha. Our results are sensitive to changes in the average cost per hectare of prescribed burning, the probabilities of fire occurrence, urban area values (in average dollars per hectare) and suppression costs.

## Acknowledgments

We thank Neil Burrows from the Department of Environment and Conservation Western Australia for his valuable contributions. We also thank the Bushfire Cooperative Research Centre, Australia for funding.

## References

- Butry, D., Mercer, D.E., Prestemon, J.P., Pye, J.M., Holmes, T.P. 2001. What is the price of catastrophic wildfire? *Journal of Forestry* **99**, 9.
- Calkin, D.E., Gebert, K.M., Jones, G., Neilson, R.P. 2005. Forest service large fire area burned and suppression expenditure trends, 1970-2002. *Journal of Forestry* **103**, 179-183.
- Cary, G., Lindenmayer, D., Dovers, S. (Eds). 2003. 'Australia Burning: Fire Ecology, Policy and Management Issues.' (CSIRO Publishing: Collingwood, VIC)
- Chatto, K., Tolhurst, K.G. 2004. A Review of the Relationship between Fireline Intensity and the Ecological and Economic Effects of Fire, and Methods Currently Used to Collect Fire Data. Report commissioned by Fire Management, Department of Sustainability and Environment, Victoria Available at <http://www.dse.vic.gov.au/fire-and-other-emergencies/living-with-fire-victorias-bushfire-strategy/fire-research-reports/research-report-67>.
- Donovan, G.H., Rideout, D.B. 2003. A reformulation of the cost plus net value change (C+NVC) model of wildfire economics. *Forest Science* **49**, 318-323.
- Donovan, GH, Rideout, DB, Omi, PN, Usda. 1999. 'The economic efficiency of the National Fire Management Analysis System and FIREPRO, Proceedings of Symposium on Fire Economics, Planning and Policy: Bottom Lines.' Available at <Go to ISI>://WOS:000167781700012
- Fernandes, PM, Botelho, HS .2003. A review of prescribed burning effectiveness in fire hazard reduction. *International Journal of Wildland Fire* **12**, 117-128.
- Finney, MA, Seli, RC, McHugh, CW, Ager, AA, Bahro, B, Agee, JK. 2007. Simulation of long-term landscape-level fuel treatment effects on large wildfires. *International Journal of Wildland Fire* **16**, 712-727.

- Ganewatta, G. 2008.** The economics of bushfire management. In 'Community Bushfire Safety.' (Eds J Handmer, K Haynes.) pp. 151-159. (CSIRO Publishing: Collingwood, VIC)
- Gebert, K, Calkin, DE, Huggett, RJ, Abt, K. 2008.** Toward a unified economic theory of fire program analysis with strategies for empirical modelling. In 'The Economics of Forest Disturbances: Wildfires, Storms and Invasive Species.' (Eds TP Holmes, JP Prestemon, K Abt.) pp. 295-322. (Springer: Dordrecht, The Netherlands)
- Handmer, J., Proudley, B. 2008.** The economics of interface wildfires. In 'Gonzalez-Caban, A. (technical coordinator) 'Proceedings of the second international symposium on fire economics, planning, and policy: a global view'. Cordoba, Spain', 19-22 April 2004. (General Technical Report PSW-GTR-208, U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. Available at [http://www.fs.fed.us/psw/publications/documents/psw\\_gtr208en/psw\\_gtr208en\\_fm.pdf](http://www.fs.fed.us/psw/publications/documents/psw_gtr208en/psw_gtr208en_fm.pdf))
- Hesseln, H, Rideout, DB. 1999.** 'Using control theory to model the long-term economic effects of wildfire, Proceedings of Symposium on Fire Economics, Planning and Policy: Bottom Lines.' Available at <Go to ISI>://WOS:000167781700013
- Johnston, P, Kelso, J, Milne, GJ. 2008.** Efficient simulation of wildfire spread on an irregular grid. *International Journal of Wildland Fire* **17**, 614-627.
- King, KJ, Bradstock, RA, Cary, GJ, Chapman, J, Marsden-Smedley, JB. 2008.** The relative importance of fine-scale fuel mosaics on reducing fire risk in south-west Tasmania, Australia. *International Journal of Wildland Fire* **17**, 421-430.
- Kochi, I, Donovan, GH, Champ, PA, Loomis, JB. 2010.** The economic cost of adverse health effects from wildfire-smoke exposure: a review. *International Journal of Wildland Fire* **19**, 803-817.
- Martin, WE, Brajer, V, Zeller, K. 2007.** Valuing the health effects of a prescribed fire. In 'Wildfire Risk: Human Perceptions and Management Implications.' (Eds WE Martin, C Raish, BJ Kenny.) pp. 244-261. (Resources for the Future:: Washington, DC, USA)
- Mell, WE, Manzello, SL, Maranghides, A, Butry, D, Rehm, RG. 2010.** The wildland–urban interface fire problem – current approaches and research needs. *International Journal of Wildland Fire* **19**, 238–251.
- Mercer, DE, Prestemon, JP, Butry, DT, Pye, JM. 2007.** Evaluating alternative prescribed burning policies to reduce net economic damages from wildfire. *American Journal of Agricultural Economics* **89**, 63-77.
- Morgan, G. 2009.** Asia and Australasia wildfire management: a regional perspective. In 'Gonzalez-Caban, A. (Technical Coordinator) Proceedings of the third international symposium on fire economics, planning, and policy: common problems and approaches. Carolina, Puerto Rico', 29 April - 2 May 2008. pp. 8-23. (General Technical Report PSW-GTR-227, U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. Available at [http://www.fs.fed.us/psw/publications/documents/psw\\_gtr227en/psw\\_gtr227\\_en.pdf](http://www.fs.fed.us/psw/publications/documents/psw_gtr227en/psw_gtr227_en.pdf))
- Morgan, G, Bartlett, T, Leonard, M. 2007.** Fire management and the bush – from an ad hoc seasonal pre-occupation to 'policy centre-stage'. In 'Tassie Fire Conference Proceedings. Hobart, Tasmania', 18 - 20 July 2007. Available at [http://proceedings.com.au/tassiefire/papers\\_pdf/thurs\\_gmorgan.pdf](http://proceedings.com.au/tassiefire/papers_pdf/thurs_gmorgan.pdf)
- Mozumder, P, Helton, R, Berrens, RP. 2009.** Provision of a wildfire risk map: informing residents in the wildland–urban interface. *Risk Analysis* **29**, 1588-1600.

- Mutch, RW, Rogers, MJ, Stephens, SL, Gill, MA. 2011.** Protecting lives and property in the wildland–urban interface: communities in Montana and southern California adopt Australian paradigm. *Fire Technology* **47**, 357-377.
- Noble, IR, Gill, AM, Bary, GAV. 1980.** McArthur's fire-danger meters expressed as equations. *Austral Ecology* **5**, 201-203.
- Richardson, LA, Champ, PA, Loomis, JB. 2012.** The hidden cost of wildfires: Economic valuation of health effects of wildfire smoke exposure in southern California. *Journal of Forest Economics* **18**, 14-35.
- Rideout, DB, Omi, PN. 1990.** Alternate expressions for the economic theory of forest fire management. *Forest Science* **36**, 614-624.
- Rideout, DB, Ziesler, PS. 2008.** 'Three Great Myths of Wildland Fire Management. In Gonzalez-Caban, A. (Coordinator) Proceedings of the second international symposium on fire economics, planning, and policy: a global view.' Albany, CA, USA, 19-22 April 2004. (U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, General Technical Report PSW-GTR-208. Available at [http://www.fs.fed.us/psw/publications/documents/psw\\_gtr208en/psw\\_gtr208en\\_319-326\\_rideout.pdf](http://www.fs.fed.us/psw/publications/documents/psw_gtr208en/psw_gtr208en_319-326_rideout.pdf)
- Rittmaster, R, Adamowicz, WL, Amiro, B, Pelletier, RT. 2006.** Economic analysis of health effects from forest fires. *Canadian Journal of Forest Research* **36**, 868-877.
- Rodriguez y Silva, F, Gonzalez-Caban, A. 2010.** 'SINAMI': a tool for the economic evaluation of forest fire management programs in Mediterranean ecosystems. *International Journal of Wildland Fire* **19**, 927-936.
- Sirakoff, C. 1985.** A correction to the equations describing the McArthur forest fire danger meter. *Austral Ecology* **10**, 481-481.
- Smalley, J. 2003.** Firewise communities: Recasting the future of the wildland-urban interface. In 'Paper presented at the 3rd International Wildland Fire Conference and Exhibition, Sydney. Available at <http://www.fire.uni-freiburg.de/summit-2003/3-IWFC/Papers/3-IWFC-017-Smalley.pdf>
- Sneeuwjagt, RJ, Peet, GB. 1998.** 'Forest Fire Behaviour Tables for Western Australia.' (Department of Conservation and Land Management, Perth, Western Australia:
- Sparhawk, WN. 1925.** The use of liability ratings in planning forest fire protection. *Journal of Agricultural Research* **30**, 693-762.
- Stockmann, KD, Burchfield, J, Calkin, DE, Venn, TJ. 2010.** Guiding preventative wildland fire mitigation policy and decisions with an economic modeling system. *Forest Policy and Economics* **12**, 147-154.
- Teague, B, Mcleod, R, Pascoe, S. 2010.** The 2009 Victoria Bushfires Royal Commission Final Report. Government Printer for the State of Victoria, Melbourne, VIC. Available at <http://www.royalcommission.vic.gov.au/Commission-Reports/Final-Report>.
- Thornton, R. 2010.** Research Programs: An Overview. Bushfire CRC, Melbourne, VIC. Available at [http://www.bushfirecrc.com/sites/default/files/1009\\_researchlist\\_lowres.pdf](http://www.bushfirecrc.com/sites/default/files/1009_researchlist_lowres.pdf).
- Williams, J, Albright, D, Hoffmann, AA, Eritsov, A, Moore, PF, Mendes De Morais, JC, Leonard, M, San Miguel-Ayaz, J, Xanthopoulos, G, van Lierop, P. 2011.** Findings and implications from a coarse-scale global assessment of recent selected mega-fires. In 'FAO at the Vth International Wildland Fire Conference. Sun City, South Africa', 9-13 May 2011. pp. 27-40. (Forestry Department, Fire Management Division Working Paper FM/27/E, Food and Agriculture Organization of the United Nations. Available at <http://www.fao.org/docrep/014/am663e/am663e00.pdf>