

Contributions of ammonia-oxidizing archaea and bacteria to nitrification in Oregon forest soils



Xinda Lu^{a,1}, Peter J. Bottomley^{a,b}, David D. Myrold^{a,*}

^a Department of Crop and Soil Science, Oregon State University, USA

^b Department of Microbiology, Oregon State University, USA

ARTICLE INFO

Article history:

Received 10 December 2014

Received in revised form

25 February 2015

Accepted 28 February 2015

Available online 14 March 2015

Keywords:

Nitrification

Ammonia oxidation

Archaea

Bacteria

Forest soil

ABSTRACT

Ammonia oxidation, the first step of nitrification, is mediated by both ammonia-oxidizing archaea (AOA) and bacteria (AOB); however, the relative contributions of AOA and AOB to soil nitrification are not well understood. In this study we used 1-octyne to discriminate between AOA- and AOB-supported nitrification determined both in soil-water slurries and in unsaturated whole soil at field moisture. Soils were collected from stands of red alder (*Alnus rubra* Bong.) and Douglas-fir (*Pseudotsuga menziesii* Mirb. Franco) at three sites (Cascade Head, the H.J. Andrews, and McDonald Forest) on acidic soils (pH 3.9–5.7) in Oregon, USA. The abundances of AOA and AOB were measured using quantitative PCR by targeting the *amoA* gene, which encodes subunit A of ammonia monooxygenase. Total and AOA-specific (octyne-resistant) nitrification activities in soil slurries were significantly higher at Cascade Head (the most acidic soils, pH < 5) than at either the H.J. Andrews or McDonald Forest, and greater in red alder compared with Douglas-fir soils. The fraction of octyne-resistant nitrification varied among sites (21–74%) and was highest at Cascade Head than at the other two locations. Net nitrification rates of whole soil without NH_4^+ amendment ranged from 0.4 to 3.3 mg N kg⁻¹ soil d⁻¹. Overall, net nitrification rates of whole soil were stimulated 2- to 8-fold by addition of 140 mg $\text{NH}_4^+\text{-N kg}^{-1}$ soil; this was significant for red alder at Cascade Head and the H.J. Andrews. Red alder at Cascade Head was unique in that the majority of NH_4^+ -stimulated nitrifying activity was octyne-resistant (73%). At all other sites, NH_4^+ -stimulated nitrification was octyne-sensitive (68–90%). The octyne-sensitive activity—presumably AOB—was affected more by soil pH whereas the octyne-resistant (AOA) activity was more strongly related to N availability.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Nitrification, the oxidation of ammonia (NH_3) to nitrate (NO_3^-), is mediated by microorganisms and is a key component of the nitrogen (N) cycle. The importance of nitrification in forest soils has been debated for a long time because N turnover and availability limit net primary productivity. In the conifer-dominated forests of the Pacific Northwest, N turnover varies between red alder (*Alnus rubra* Bong.) sites and conifer sites (Binckley et al., 1992; Perakis et al., 2012) and a several-fold increase in the rates of N turnover by the inclusion of red alder has been reported (Hart et al., 1997; Boyle et al., 2008). As a result of red alder having the potential to

fix 50–300 kg N ha⁻¹ y⁻¹ (Bottomley and Myrold, 2015), there is an increase in the N capital and N availability of the soil. A negative consequence is that red alder can reduce soil pH, to as low as 3.6–3.9 (Yarwood et al., 2010). The pH decline under red alder probably occurs as a result of increased nitrification (Binkley and Sollins, 1990). Although ammonia-oxidizing bacteria (AOB) have been isolated from soils of pH < 5 (De Boer and Kowalchuk, 2001), it remains equivocal whether or not they can effectively oxidize NH_3 at that pH. Heterotrophic nitrification, mediated by heterotrophic bacteria and fungi, has been suggested to be an alternate option (Hynes and Knowles, 1982; Stein, 2011; Zhang et al., 2011). Another possibility is the recent discovery of ammonia-oxidizing *Thaumarchaeota* (AOA) that have the potential for NH_3 oxidation and growth on CO_2 . Recently, an acidophilic AOA has been isolated (Lehtovirta-Morley et al., 2011, 2013), and may represent a lineage of dominant AOA types in acidic soils (Nicol et al., 2008; Gubry-Rangin et al., 2010; Yao et al., 2011; Prosser and Nicol, 2012; Hu et al., 2014). Although no isolates of AOB have been found to grow at

* Corresponding author. Agriculture and Life Sciences Building 3017, Oregon State University, Corvallis, OR 97331-7306, USA. Tel.: +1 541 737 5737.

E-mail address: david.myrold@oregonstate.edu (D.D. Myrold).

¹ Currently University of Waterloo, Canada.

pH < 5 (De Boer and Kowalchuk, 2001), they exist in acidic soils and their potential for activity has been inferred by detection of AOB mRNA for *amoA* (Nicol et al., 2008). This might be due to residence in pH-neutral microsites (Hankinson and Schmidt, 1984), an ability to use urea and/or other organic N sources (Burton and Prosser, 2001), or through protective aggregation (De Boer et al., 1991; Spieck et al., 1992). In most soils, AOA outnumber AOB based on their *amoA* gene abundances, suggesting the potential for a greater role in nitrification (Leininger et al., 2006; Prosser and Nicol, 2008; Norman and Barrett, 2014). In some soils, however, AOB can be more abundant than AOA (Hu et al., 2014; Petersen et al., 2012; Yarwood et al., 2010). Thus, to date no clear consensus has emerged about the mechanism(s) controlling niche differentiation in acidic soils.

We designed a study using our newly developed assay, which is based upon AOB NH_3 -oxidizing activity being quickly and irreversibly inactivated by low concentrations of 1-octyne, whereas AOA activity is unaffected by the same range of concentrations (Taylor et al., 2013). The study involved a comparison of the relative contribution of AOA and AOB to soil nitrification in acidic forest soils under different tree types. Three sites spanning a range of soil types and containing replicated stands of Douglas-fir (*Pseudotsuga menziesii* Mirb. Franco) and red alder were selected. The soils varied in pH and organic C and N contents, in part because of the N inputs provided by the red alder-*Frankia* symbiosis. The aims of this study were to: (i) assess the relative contributions of AOA and AOB to nitrification; and (ii) determine what factors influence their distribution and contribution to nitrification. It was hypothesized that N input through N_2 -fixation by red alder would increase both ammonia oxidizer abundances and nitrification potential, and would affect the relative contributions of AOA and AOB to nitrification.

2. Materials and methods

2.1. Site description and soil properties

Soils were collected from three forest sites containing plots of either pure stands of Douglas-fir or red alder. At two of the sites the stands were planted approximately 30 years ago (Radosevich et al., 2006). One site lies within the Cascade Head Experimental Forest (45°01' N, 123°54'W, 330 m elevation), 1.6 km from the Pacific Ocean, and receives ~2400 mm y^{-1} of precipitation. Soils at the site are classified as Isomesic Fluvudands. A second site was located within the H.J. Andrews Experimental Forest (44°14' N, 122°11'W, elevation 800 m), along the western slopes of the Cascade Range, and receives ~2300 mm y^{-1} of precipitation. Soils in the area are classified as Typic Haplumbrepts (Dyrness, 2001). The third site was located within the McDonald-Dunn Forest (44°42' N, 123°20'W, 350 m elevation), on the eastern foothills of the Coast Range. The soils are classified as Xeric Palehumults. Annual precipitation averages ~1070 mm y^{-1} . At McDonald Forest, natural stands of red alder are found along streams within a matrix of Douglas-fir; both about 30 years old. All soil samples were collected during the late spring of 2013. Soil was collected from each of three field replicate plots of each tree type at each location, kept in separate bags, and stored at 4 °C and used within six months. Samples (10 g) were removed from each bag and frozen (−20 °C) for DNA extraction at a later date.

Soils spanned a relatively wide range of pH (3.9–5.5), a three-fold range in total C, a four-fold range of total N, and an eight-fold range of NH_4^+ and NO_3^- (Table 1). Soil pH was measured in deionized water; total C and N were measured by combustion; concentrations of NH_4^+ and NO_3^- plus NO_2^- (referred to simply as

NO_3^- afterwards) in 2 M KCl extracts were determined colorimetrically (Hood-Nowotny et al., 2010; Kandeler and Gerber, 1988).

2.2. Nitrification potential assay

Nitrification potentials of soils were measured using a shaken soil-slurry method with supplemental NH_4^+ (1 mM NH_4Cl ; Taylor et al., 2010). Field moist soil (2.5 g) was added to 150-ml serum bottles loosely capped by phenolic caps fitted with butyl stoppers, and pre-incubated at room temperature (22 ± 2 °C) for 2 d. Deionized water (15 ml) supplemented with 1 mM NH_4Cl was added to each bottle (final concentration ~140 mg $\text{NH}_4^+\text{-N kg}^{-1}$ soil). Bottles were shaken at 200 rpm at 30 °C. Samples of soil slurries (1 ml) were taken after 15 min of shaking and at 24 and 48 h, centrifuged for 3 min at 13,000× g, and NO_3^- determined. At the beginning of the slurry assays, a set of bottles were amended with 2 kPa of either acetylene (6 μM aqueous) or octyne gas (4 μM aqueous) as described by Taylor et al. (2013). Acetylene was used to block NH_3 oxidation by AOA and AOB and to evaluate the possibility of acetylene-insensitive heterotrophic nitrification. Octyne was used to block AOB activity while allowing AOA activity to proceed (Taylor et al., 2013). For Cascade Head and McDonald Forest soils, nitrification potentials were calculated from NO_3^- accumulation during the first 24 h. For H.J. Andrews soils, nitrification potentials were calculated from NO_3^- accumulation occurring between 24 and 48 h due to lack of significant accumulation during 0–24 h. Nitrification potentials were calculated by subtracting the NO_3^- accumulation in the acetylene treatment from values measured in the treatments without acetylene to focus primarily on acetylene-sensitive AOA and AOB activities. The pH of the soil slurries was measured at the end of the incubation.

2.3. Whole soil nitrification assay

A whole soil assay was used to measure the potential contributions of AOA and AOB to soil nitrification. Soil samples (5 g) were incubated at three different NH_3 levels (equivalent to 0, 14, and 140 mg $\text{NH}_4^+ \text{kg}^{-1}$ soil) achieved by adding sufficient anhydrous NH_3 gas to the headspace of 150-ml serum bottles sealed by phenolic caps fitted with butyl stoppers. Prior to initiation of the assay, soils were pre-incubated at room temperature (22 ± 2 °C) for 2 d. Three treatments were imposed at each NH_4^+ level: (i) positive control (no octyne or acetylene amendment), (ii) acetylene amendment (2 kPa = 6 μM aqueous), and (iii) octyne amendment (2 kPa = 4 μM aqueous). Soil samples were incubated at 25 °C, NH_4^+ and NO_3^- concentrations were determined at 2 and 7 d, and nitrification rates were calculated after subtracting NO_3^- accumulation in the acetylene treatment. The pH at the beginning of whole soil incubation was measured 12 h after NH_3 gas was added, when NH_3 gas had fully defused into soil samples and pH came to equilibrium. The pH of soil samples was measured at the end of the incubation.

2.4. Copy numbers of *amoA* genes

DNA was extracted from soil (0.25 g dry weight equivalent) using a MoBio PowerSoil™ DNA isolation kit (MoBio Laboratories Inc, Carlsbad, CA) according to the manufacturer's instructions. DNA extracts were quantified with a NanoDrop™ ND-1000 UV–Vis Spectrophotometer (NanoDrop Technologies, Wilmington, DE) and diluted to 1 ng μl^{-1} . DNA extracts were stored at −20 °C for future use.

The quantitative polymerase chain reaction (qPCR) was used to determine AOA and AOB abundance by quantifying the number of *amoA* gene copies for each group. All qPCR reactions were performed in triplicate by using an ABI PRISM 7500 FAST (Carlsbad, CA)

Table 1
Properties of Douglas-fir and red alder soils at each experimental site. Data are mean $\pm$ standard error ($n = 3$); superscripts denote differences based on two-way ANOVA ($p < 0.05$, Fisher's least significant difference (LSD) procedure).

Site	Tree species	Total C	Total N	C:N	NH ₄ ⁺	NO ₃ ⁻	pH	Water content
		(g kg ⁻¹ soil) ^a			(mg N kg ⁻¹ soil)			(kg kg ⁻¹ soil)
Cascade Head	Red alder	144 $\pm$ 10.4 ^a	9.2 $\pm$ 0.9 ^a	15.7 $\pm$ 0.5 ^a	1.5 $\pm$ 0.6 ^{ab}	25.3 $\pm$ 2.6 ^a	3.9 $\pm$ 0.0 ^a	1.24 $\pm$ 0.07 ^a
	Douglas-fir	128 $\pm$ 2.3 ^a	6.7 $\pm$ 0.1 ^b	19.1 $\pm$ 0.0 ^a	0.5 $\pm$ 0.2 ^a	9.9 $\pm$ 0.9 ^b	4.7 $\pm$ 0.2 ^b	0.97 $\pm$ 0.09 ^{abc}
H.J. Andrews	Red alder	82 $\pm$ 12.1 ^b	3.4 $\pm$ 0.2 ^c	23.8 $\pm$ 2.4 ^b	1.2 $\pm$ 0.6 ^a	7.5 $\pm$ 1.0 ^{bc}	4.8 $\pm$ 0.1 ^b	0.55 $\pm$ 0.03 ^{bc}
	Douglas-fir	90 $\pm$ 7.5 ^b	2.7 $\pm$ 0.1 ^c	33.2 $\pm$ 2.1 ^c	2.3 $\pm$ 1.3 ^{ab}	3.3 $\pm$ 0.5 ^c	5.0 $\pm$ 0.1 ^{bc}	0.60 $\pm$ 0.04 ^{bc}
McDonald Forest	Red alder	52 $\pm$ 9.4 ^c	2.8 $\pm$ 0.5 ^c	18.6 $\pm$ 0.2 ^a	1.8 $\pm$ 0.5 ^{ab}	10.2 $\pm$ 2.4 ^b	5.2 $\pm$ 0.1 ^{cd}	1.08 $\pm$ 0.36 ^{ab}
	Douglas-fir	53 $\pm$ 1.5 ^c	2.7 $\pm$ 0.1 ^c	19.6 $\pm$ 0.8 ^a	4.1 $\pm$ 1.4 ^b	3.6 $\pm$ 0.6 ^c	5.5 $\pm$ 0.2 ^d	0.44 $\pm$ 0.23 ^c

^a Total C and total N data are modified from Boyle et al. (2008) and Zeglin et al. (2011).

sequence detection system set to read SYBR green fluorescence. Standard curves were constructed using *Nitrosomonas europaea* genomic DNA (AOB *amoA*) or *Nitrososphaera viennensis* genomic DNA (AOA *amoA*). AOB *amoA* genes were amplified using primers *amoA*-1F (5'-GGG GGT TTC TAC TGG TGG T; Stephen et al., 1998) and *amoA*-2R (5'-CCC CTC KGS AAA GCC TTC TTC; Rothauwe et al., 1997). AOA *amoA* genes were amplified using primers Cren-*amoA*23f (5'-ATG GTC TGG CTW AGA CG; Tourna et al., 2008) and Cren-*amoA*616r (5'-GCC ATC CAT CTG TAT GTC CA; Tourna et al., 2008). Copy numbers are reported as *amoA* gene copies g⁻¹ dry soil. Primers sets, thermal protocols master mix recipes, standard curve r^2 , and reaction efficiency are summarized in Supplemental Material (Table S1).

2.5. Statistical analysis

Treatment differences were evaluated using analysis of variance (ANOVA), followed by the Fisher's least significant differences (LSD) test. Unless otherwise noted, only significant ($p < 0.05$) interactions are discussed. To meet the assumptions of normality and homogeneity of variance, whole soil nitrification rates were square-root transformed and *amoA* gene copy numbers were log transformed. Relationships among variables were examined by correlation analysis. Analyses were done using StatGraphics Centurion Version 16.1.03 (Statpoint Technologies, Inc., Warrenton, VA) or SigmaPlot 12.3 (Systat Software, Inc., San Jose, CA).

3. Results

3.1. Soil properties

McDonald Forest soils contained the lowest amount of total C and Cascade Head soils the most (Table 1). There was no significant difference in total C between soils under red alder and Douglas-fir at any of the sites. Total N of Cascade Head soils was significantly higher than soils from other two sites, which were not significantly different. Total N was consistently higher in red alder compared to Douglas-fir, but this was statistically significant only at Cascade Head. Soil pH ranged from 3.9 to 5.5 with highly significant difference among soils. Cascade Head soils had the lowest pH; McDonald Forest soils had the highest pH values. At Cascade Head, soils under red alder had significantly lower pH than soils under Douglas-fir. This trend also held for soils from the other two sites but differences were not significant. There was little difference in extractable NH₄⁺ among sites or between tree species. Cascade Head soils had more extractable NO₃⁻ than the other two sites, but at all sites NO₃⁻ was higher under red alder than under Douglas-fir. Most soil properties, except for C:N ratio, were highly correlated with each other; for example, pH was inversely correlated with total N ($r = -0.83$) but not to C:N ratio ($r = 0.26$). Thus, relationships between nitrification rates and soil properties will focus on C:N ratios (an index of N availability) and pH.

3.2. Soil slurry assay

When acetylene was added to soil slurries, no significant NO₃⁻ accumulation was measured in any of the soils during the first 24 h of incubation; however, from 24 to 48 h there was some acetylene-resistant activity in red alder soils at the H.J. Andrews (2.8 $\pm$ 0.4 mg N kg⁻¹ soil d⁻¹), which may indicate the presence of a low population of heterotrophic nitrifiers in that soil. To eliminate any contribution from heterotrophic nitrification, all nitrification potentials were adjusted by subtracting the NO₃⁻ values of the acetylene treatment prior to statistical analysis of the nitrification potential data.

3.2.1. Nitrification potentials of soil slurries

Two-way ANOVA showed that the nitrification potentials varied significantly among sites and between tree types (Fig. 1). Cascade Head soils averaged 11.0 mg N kg⁻¹ soil d⁻¹, which was significantly higher than McDonald Forest (2.4 mg N kg⁻¹ soil d⁻¹) and H.J. Andrews (1.2 mg N kg⁻¹ soil d⁻¹) soils. Across all sites, nitrification potential was two-fold greater in red alder soils compared with Douglas-fir soils. No significant NO₃⁻ accumulation was detected in H.J. Andrews Douglas-fir soils.

The initial pH of soil slurries varied significantly among sites ($p < 0.01$) and tree types ($p < 0.01$): 4.4–5.4 at Cascade Head, 5.1–5.6 at the H.J. Andrews, and 5.6–6.2 at McDonald Forest. At the end of the nitrification potential incubations, pH values of the H.J. Andrews and McDonald Forest soil slurries remained similar to those measured initially, but the pH of Cascade Head soil slurries




Fig. 1. Potential nitrification activities of octyne-resistant (archaeal) and octyne-sensitive (bacterial) ammonia oxidizers in short-term soil slurry assays supplemented with 1 mM NH₄⁺ (~140 mg NH₄⁺ N kg⁻¹ soil). Data are means $\pm$ standard errors ($n = 3$). No potential nitrification activity was found for the Douglas-fir soil at the H.J. Andrews. Two-way ANOVA table shows the statistical significance (p -value) of site, tree type, and their interaction.

had decreased significantly to pH 3.9–4.6 ($p = 0.02$), presumably due to greater H^+ associated with higher nitrification activities and differential buffering among the soils.

3.2.2. Effect of octyne on nitrification potentials of soil slurries

Two-way ANOVA showed that the octyne-resistant nitrification potential varied significantly among sites and tree type, although there was a barely significant interaction (Fig. 1). The octyne-resistant nitrification potential at Cascade Head averaged $7.1 \text{ mg N kg}^{-1} \text{ soil d}^{-1}$, 12-fold greater than at McDonald Forest and 20-fold greater than at the H.J. Andrews. The fraction of the nitrification potential that was octyne-resistant ranged from 21 to 74%, indicating that the relative contribution of AOA varied among soils. At Cascade Head, nitrification potentials with or without octyne were not significantly different, suggesting that the high potential nitrification activity was dominated by AOA. A decrease in nitrification potential was detected at McDonald Forest for Douglas-fir ($p = 0.05$) and red alder ($p = 0.08$) soils when exposed to octyne, indicating an increased AOB contribution (octyne sensitive). The effect of octyne on nitrification potentials was compared to the effect of $100 \mu\text{M}$ ATU (1-allyl-2-thiourea; known to inhibit AOB but not AOA) at two randomly selected soils (Cascade Head and H.J. Andrews red alder soils). The sensitivities of nitrification potential to octyne and ATU were strongly correlated ($r^2 = 0.74$, $p < 0.01$), adding further evidence to suggest that AOA were responsible for the octyne-resistant potential nitrification activity.

3.2.3. Environmental factors

Nitrification potentials decreased significantly in a curvilinear manner with both soil pH ($r^2 = 0.60$) and C:N ratio ($r^2 = 0.51$; Fig. 2a,b). Similar trends were found for the fraction of octyne-resistant activity but with a weaker relationship for pH ($r^2 = 0.26$) than C:N ratio ($r^2 = 0.59$; Fig. 2c,d).

3.3. Whole soil nitrification assay

When acetylene was added to whole soil, no significant NO_3^- accumulation was detected in soils from McDonald Forest; some accumulation of NO_3^- was measured in soils from Cascade Head ($<3.2 \text{ mg N kg}^{-1} \text{ soil}$) and the H.J. Andrews ($<5.4 \text{ mg N kg}^{-1} \text{ soil}$) during the 0–2 d interval, but no further significant NO_3^- accumulation occurred between 2 and 7 d. As with nitrification potentials, whole soil nitrification rates were adjusted by subtracting the NO_3^- values of the acetylene-treated samples.

3.3.1. Nitrification rate

Measurable accumulation of NO_3^- was detected within all soils at 7 d. Three-way ANOVA showed that site, tree type, and NH_3 addition level (Fig. 3) affected nitrification rates. Whole soil nitrification rates of all soils were greater when NH_3 was added, however, only the high NH_3 concentration treatment showed significantly greater nitrification rates compared with the no and low NH_3 addition treatments, which were not statistically different.

Focusing on the nitrification rates in whole soil without NH_3 addition, which is similar to *in situ* conditions, two-way ANOVA showed that net NO_3^- accumulation was significantly influenced by site ($p = 0.04$). Cascade Head, had the highest nitrification rate of $2.8 \text{ mg N kg}^{-1} \text{ soil d}^{-1}$, with McDonald Forest ($1.3 \text{ mg N kg}^{-1} \text{ soil d}^{-1}$), and the H.J. Andrews ($0.9 \text{ mg N kg}^{-1} \text{ soil d}^{-1}$) being lower (Fig. 3). Nitrification rates tended to be higher in red alder soils than in Douglas-fir soils across all sites ($p = 0.09$).

Nitrification rates with high NH_3 addition were about three-fold higher than that of the no NH_3 control (Fig. 3). No significant differences in rates were found among sites ($p = 0.13$) but it is interesting that the order of nitrification rates changed relative to the no

NH_3 control, with McDonald Forest ($7.2 \text{ mg N kg}^{-1} \text{ soil d}^{-1}$) highest, followed by Cascade Head ($5.8 \text{ mg N kg}^{-1} \text{ soil d}^{-1}$) and H.J. Andrews soils ($3.2 \text{ mg N kg}^{-1} \text{ soil d}^{-1}$). Ammonia-amended nitrification rates were significantly higher in red alder compared to Douglas-fir stands ($p = 0.03$).

After 7 d of incubation, the pH of whole soil was lower than the original pH in all treatments, and lowest at Cascade Head. Without NH_3 addition, the final pH was 3.5–4.1 at Cascade Head, 0.6 units lower than original pH; 5.1–5.8 at the H.J. Andrews, 0.5 units lower than original pH, and 5.4–5.8 at McDonald Forest, 0.5 units lower than original pH. With high NH_3 concentration treatment, final pH was 3.6–4.5 at Cascade Head, 0.9 units lower than original pH; 5.1–5.8 at H.J. Andrews, 0.7 units lower than original pH and 5.3–5.6 at McDonald Forest, 0.8 units lower than original pH.

3.3.2. Effect of octyne

Three-way ANOVA showed that octyne-resistant activity varied significantly among sites, tree types, and NH_3 addition (Fig. 3). Nitrate accumulation rate at Cascade Head was $2.72 \text{ mg N kg}^{-1} \text{ soil d}^{-1}$, which was significantly higher than the rate at the H.J. Andrews and McDonald Forest. No significant difference was observed between the latter two sites. Soils under red alder had a three-fold higher NO_3^- accumulation rate compared to soils under Douglas-fir across all sites ($p < 0.01$). Octyne-resistant activity at the high concentration of NH_3 addition was significantly higher than the non-amended control, but that of the low NH_3 concentration did not differ from the other two treatments.

The fraction of octyne-resistant activity was significantly influenced by site ($p < 0.01$) and tree type ($p < 0.01$) but not by NH_3 additions ($p = 0.58$). Octyne-resistant activity at Cascade Head accounted for 62% of the total nitrifying activity, significantly higher than the fraction at the H.J. Andrews (13%) and McDonald Forest (19%). Octyne-resistant activity was significantly higher under red alder (45%) than under Douglas-fir (17%).

3.3.3. Environmental factors

In non-amended whole soils, the nitrification rate was positively, linearly correlated with total N ($r^2 = 0.32$) but not with pH or C:N ratio (Suppl. Fig. 1a,b). The fraction of octyne-resistant activity showed a linear positive correlation with total N ($r^2 = 0.38$), and negative curvilinear correlations with soil pH ($r^2 = 0.52$) and C:N ratio ($r^2 = 0.47$; Suppl. Fig. 1c,d). In soils amended with a high concentration of NH_3 , no significant correlations were found between nitrification rate and soil properties, but significant correlations were found for the octyne-resistant fraction. This fraction showed positive linear correlation with total N ($r^2 = 0.64$), negative curvilinear correlations with soil pH ($r^2 = 0.75$) and C:N ratio ($r^2 = 0.49$; Fig. 4).

3.4. AOA and AOB *amoA* gene abundance

Copies of AOA and AOB *amoA* genes were detected in all soil samples, but copy numbers varied widely among samples. Across all soils, AOA *amoA* gene abundance ranged from 1×10^4 to $2 \times 10^8 \text{ amoA genes g}^{-1} \text{ soil}$ and AOB *amoA* gene abundance ranged from 4×10^4 to $3 \times 10^7 \text{ amoA genes g}^{-1} \text{ soil}$ (Fig. 5). The site-by-tree interaction was significant for AOA ($p = 0.02$) and AOB ($p < 0.01$) *amoA* gene abundances. Abundances of both AOA and AOB *amoA* genes were higher in red alder than Douglas-fir soils, but this difference was statistically significant only at the H.J. Andrews. Ammonia-oxidizing archaeal *amoA* gene copy numbers were significantly higher under both tree types at Cascade Head than the other two sites, and higher at McDonald Forest than at the H.J. Andrews under Douglas-fir. Abundances of AOB *amoA* genes were significantly higher under both tree types at McDonald Forest than




Fig. 2. Relationship between total nitrification potential (a, b) or octyne-resistant (AOA) fraction of the nitrification potential (c, d) and soil slurry pH (a, c) and C:N ratio (b, d). Soils are identified by site (CH—Cascade Head, black; HJA—H.J. Andrews, white; MF—McDonald Forest, gray) and tree type (DF—Douglas-fir, triangle; RA—red alder, circle). Lines represent best fit, exponential regressions; all are significant at $p < 0.05$.

the other two sites; and higher at Cascade Head than at the H.J. Andrews under Douglas-fir. The AOA:AOB ratio varied significantly across sites ($p < 0.01$; Fig. 5). Cascade Head soils were strongly dominated by AOA, whereas McDonald Forest soils were dominated by AOB; H.J. Andrews soils contained approximately equal

numbers of AOA and AOB. Tree species also influenced the AOA:AOB ratio, which tended to be higher in soils under red alder than soils under Douglas-fir ($p = 0.08$).

Ammonia-oxidizing archaeal *amoA* gene copy numbers decreased significantly as soil pH increased ($r^2 = 0.32$), but there was no correlation between AOB *amoA* gene copy numbers and pH (Fig. 6a). Both AOA ($r^2 = 0.68$) and AOB ($r^2 = 0.63$) *amoA* gene copy numbers significantly decreased as soil C:N ratio increased (Fig. 6b).

3.5. Linking abundance with activity

Regression analysis showed that nitrification potential was linearly and positively correlated with AOA ($r^2 = 0.53$) but not AOB ($r^2 < 0.01$) *amoA* gene copy numbers, and was not improved if gene copy numbers of both groups of ammonia oxidizers were included in a multiple regression analysis. The octyne-resistant nitrification potential was even more highly correlated with AOA *amoA* gene copy numbers (Fig. 7; $r^2 = 0.80$) whereas octyne-sensitive nitrification potential was not correlated with AOB *amoA* gene copy numbers ($r^2 = 0.05$).

Nitrification rates of either non-amended or NH_4^+ -amended whole soils were not significantly correlated with AOA or AOB *amoA* gene copy numbers, either individually or in a multiple regression approach; however, the octyne-resistant nitrification rate of whole soil was significantly correlated with AOA *amoA* copy numbers in both NH_4^+ -amended ($r^2 = 0.88$) or non-amended ($r^2 = 0.73$) whole soils. No correlation between octyne-sensitive nitrification and




Fig. 3. Nitrification activities of octyne-resistant (archaeal) and octyne-sensitive (bacterial) ammonia oxidizers in whole soil assays supplemented with NH_3 equivalent to 0 or 140 $\text{mg NH}_4\text{-N kg}^{-1}$ soil. Data are means $\pm$ standard errors ($n = 3$). Three-way ANOVA table shows the statistical significance (p -value) of site, tree type, and NH_3 addition; interactions were not significant ($p > 0.05$).




Fig. 4. Relationship between the octyne-resistant (AOA) fraction of nitrification in whole soil supplemented with NH_3 (equivalent to $140 \text{ mg NH}_4^+\text{-N kg}^{-1}$ soil) and soil pH (a) and C:N ratio (b). Soils are identified by site (CH—Cascade Head, black; HJA—H.J. Andrews, white; MF—McDonald Forest, gray) and tree type (DF—Douglas-fir, triangle; RA—red alder, circle). Lines represent best fit, exponential regressions; all are significant at $p < 0.05$.




Fig. 5. Abundance of ammonia-oxidizing archaea (AOA) and bacteria (AOB) based on copy numbers of their *amoA* genes. Data are means $\pm$ standard error ($n = 3$). Number above a set of bars is the AOA:AOB ratio.

AOB *amoA* gene copy numbers was found in either non-amended or NH_4^+ -amended whole soils.

4. Discussion

4.1. Activity and abundance of ammonia oxidizers

The nitrification rates measured across the three sites fall within the range of about $0.1\text{--}14 \text{ mg N kg}^{-1} \text{ soil d}^{-1}$ that has been reported for forest soils worldwide (Vitousek et al., 1982; Wertz et al., 2012). The inclusion of red alder in conifer-dominated forests of the Pacific Northwest, with its large N input from N_2 fixation, has been found to increase both gross and net nitrification in soils from both low- and high-productivity sites with a concomitant decrease soil pH (Binkley and Sollins, 1990; Hart et al., 1997; Boyle et al., 2008). This combination of shifts in soil pH and N availability raises questions about what kinds of ammonia oxidizers might contribute to soil nitrification, such as AOA versus AOB, and the possibility that different phylotypes of each might contribute depending on the severity of the acidity. Both nitrification potential and whole soil nitrification assays showed a strong inverse relationship between




Fig. 6. Relationship between ammonia-oxidizing archaea (AOA) or bacteria (AOB), based on copy numbers of their *amoA* genes, and soil pH (a) and C:N ratio (b). Copy numbers of AOA *amoA* genes (black circles, solid lines) were significantly correlated with soil pH ($R^2 = 0.32$, $p = 0.01$) and C:N ratio ($R^2 = 0.68$, $p < 0.01$), whereas AOB *amoA* gene copy numbers (white circles, dashed lines) were correlated with only soil C:N ratio ($R^2 = 0.63$, $p < 0.01$) but not pH ($p = 0.81$).




Fig. 7. The rate of octyne-resistant (AOA) nitrification potential is strongly correlated with AOA *amoA* gene copy numbers. Soils are identified by site (CH—Cascade Head, black; HJA—H.J. Andrews, white; MF—McDonald Forest, gray) and tree type (DF—Douglas-fir, triangle; RA—red alder, circle). Line represents exponential regression ($R^2 = 0.80$, $p < 0.01$).

nitrification rate and soil C:N ratio, with nitrification rate sharply increasing below a threshold of C:N around 20. This finding is consistent with earlier findings (Aber et al., 2003; Bengtsson et al., 2003; Ross et al., 2004), and a similar trend has been reported in several European sites with a threshold of C:N ratio 24–27 (Emmett et al., 1998; Gundersen et al., 1998).

Regression analysis showed soil pH to have less of an influence on the abundance of ammonia oxidizers than soil C:N ratio, which suggests that production of NH_4^+ may be a greater determinant of the carrying capacity of ammonia oxidizers than soil acidity. This is clearest for AOA, which were present in highest abundance in the soil with the lowest pH but also lowest C:N ratio (Cascade Head), consistent with the idea that AOA activity may be tightly linked to organic N mineralization (Levičnik-Höfferle et al., 2012). Ammonia-oxidizing bacteria, on the other hand, were most dominant in the soils of the highest pH but intermediate C:N ratio (McDonald Forest), consistent with the generalization that AOB are less tolerant of acidity than AOA (Gubry-Rangin et al., 2011; Hu et al., 2014; Zhang et al., 2012).

4.2. Relative contribution of AOA and AOB to nitrification

Although we have evidence to suggest that 1-octyne is a potentially selective inhibitor of AOB activity in agricultural soils (Taylor et al., 2013), that concept was not tested on acidic forest soils. In this current study, we showed that a positive relationship exists between AOA *amoA* gene copy abundance and octyne-resistant nitrifying activity in both slurries (Fig. 7) and whole soil assays (data not shown). Based on the slope of this relationship, an NH_3 oxidation rate of approximately $2 \times 10^{-16} \text{ mol cell}^{-1} \text{ h}^{-1}$ could be calculated. To place this value into context, the per-cell rate of NH_3 oxidation measured in the marine AOA, *Nitrosopumilus maritimus*, ranges between 2.5 and $3.5 \times 10^{-16} \text{ mol cell}^{-1} \text{ h}^{-1}$ (Könneke et al., 2005). From published data we calculated a per-cell rate of $1\text{--}2 \times 10^{-16} \text{ mol cell}^{-1} \text{ h}^{-1}$ for the soil AOA *Nitrososphaera viennensis* grown at circumneutral pH (Tourna et al., 2011); however, per-cell rates calculated from two different data sets for the acidophilic soil AOA, *Nitrosotalea devanattera* (grown at pH 4.5), provided somewhat lower estimates ($\sim 0.2\text{--}1 \times 10^{-16} \text{ mol cell}^{-1} \text{ h}^{-1}$; Lehtovirta-Morley et al., 2011, 2013). Clearly, the qPCR based estimates of AOA populations found in the Oregon acidic forest soils could account for

the octyne-resistant activity measured, suggesting that most of the AOA in these acidic forest soils were active. By contrast, no significant correlation was found between AOB abundance and octyne-sensitive activity, perhaps because not all measured AOB were physiologically active at the low pH of these soils, that the proportion of those that were active varied among the soils, or that some other type of octyne-sensitive microorganisms may contribute to the measured activity. Despite these caveats, the per-cell rate for AOB obtained from averaging across all soils is about $6 \times 10^{-15} \text{ mol cell}^{-1} \text{ h}^{-1}$, which is similar to that calculated for soil AOB (Jiang and Bakken, 1999; Taylor and Bottomley, 2006).

Measuring *amoA* gene expression might be a complementary, or even alternative, approach to the selective inhibitor method that we used. For example, both potential NH_3 -oxidizing activity and *amoA* mRNA increased following the resuscitation of starved cultures of an AOB, but the authors cautioned against making direct correlations because *amoA* mRNA persisted during starvation (Bollmann et al., 2005). In soil, AOA and AOB *amoA* transcripts were observed to increase following wetting of dried soils, however, NH_3 -oxidizing activities of the two groups were not measured (Placella and Firestone, 2013). In the future, a comparison of AOA and AOB NH_3 -oxidizing activity and their expression of *amoA* genes would be worthwhile as it is important to determine whether there is a direct relationship between transcript numbers and activity under a range of soil and environmental conditions.

4.3. Niche differentiation

Although considerable evidence now suggests that AOA oxidize NH_3 in acidic soils (Nicol et al., 2008; Gubry-Rangin et al., 2010; Stopnisek et al., 2010; Yao et al., 2011; Zhang et al., 2011), it remains a fact that in all of these studies, except one (Stopnisek et al., 2010), the soils also contained populations of AOB. In our study we detected AOB in all soils, and found that NH_4^+ -stimulated activity was octyne sensitive in all soils except red alder at Cascade Head. As mentioned in the introduction, various hypotheses have been put forward through the years to invoke why generally acid-sensitive AOB might persist and be locally active in acidic forest soils, including protective microsites. In this context, octyne-sensitive nitrification rates in soils from McDonald Forest and the H.J. Andrews were lower in disaggregated soil slurries than in whole soils amended with the equivalent amount of NH_4^+ .

By contrast, Cascade Head soils, particularly under red alder, were of lower pH and where dominated by AOA activity. There is precedent in the literature for small pH differences among acidic soils affecting AOA composition. Yao et al. (2011) in a survey of about 20 acidic tea soils from China reported that some types of AOA *amoA* sequences were restricted to soils of pH < 4.4, whereas others were found only in soils with pH > 5. In a survey of AOA phylotypes among a large number of soils Gubry-Rangin et al. (2011) separated the soils above and below pH 5, and found some AOA types restricted to soils of pH < 5. Furthermore, Lehtovirta-Morley et al. (2011) showed that NH_3 oxidation by the acidophilic AOA *Nitrosotalea devanattera* was optimum between pH 4–5, and dropped off sharply above pH 5, implying that this type of AOA might prefer to occupy soil of pH < 5. Boyle-Yarwood et al. (2008) found that AOA communities were different in soils under red alder and Douglas-fir at Cascade Head also supporting the hypothesis that small difference in the level of soil acidity might be sufficient to invoke these shifts. Nonetheless, a tree type effect that is independent of pH cannot be ruled out in that study. It is possible that AOA phylotypes exist in the acidic forest soils that are more sensitive to inhibition by octyne, and whose contribution to soil nitrifying activity is affected by pH.

Interestingly, although AOA are known to have a high affinity for NH_3 , addition of high concentrations of NH_3 stimulated both octyne-resistant and octyne-sensitive nitrification, suggesting that both AOA and AOB were substrate limited in these forest soils. The response of AOB to added NH_4^+ is common, but not all studies have found AOA to be responsive (Verhamme et al., 2011; Levcik-Höfferle et al., 2012; Taylor et al., 2013). The positive response to NH_3 additions in these forest soils is likely related to their low pH and accompanying lower soil solution NH_3 concentration, particularly at Cascade Head. In such soils, AOA activity may be tightly linked to organic N mineralization, as suggested by a study showing AOA activity being stimulated by addition of mineralizable N: urea, glutamate, or yeast extract (Levcik-Höfferle et al., 2012).

5. Summary

The distribution and relative contribution of AOA and AOB to nitrification in forest soils remains a topic of debate. This study demonstrated that soil pH, substrate concentration, and C:N ratio are all key factors that differentiate niches occupied by AOA and AOB. The activity of both AOA and AOB responded the addition of substrate but AOA activity was more closely linked to N availability, as indicated by soil C:N ratio, whereas AOB activity was affected more by soil pH.

Acknowledgements

This research was partially funded by USDA NIFA Award No. 2012-67019-3028 and the Oregon Agricultural Research Foundation. We thank Dr. Anne Taylor and Andrew Giguere for assistance with the nitrification assays, and Dr. Kristin Kasschau and Rachel Danielson for help with the qPCR assays.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.soilbio.2015.02.034>.

References

- Aber, J.D., Goodale, C.L., Olinger, S.V., Smith, M.-L., Magill, A.H., Martin, M.E., Hallett, R.A., Stoddard, J.L., 2003. Is nitrogen deposition altering the nitrogen status of Northeastern forests? *Bioscience* 53, 375–390.
- Bengtsson, G., Bengtson, P., Månsson, K.F., 2003. Gross nitrogen mineralization-, immobilization-, and nitrification rates as a function of soil C/N ratio and microbial activity. *Soil Biology and Biochemistry* 35, 143–154.
- Binkley, D., Sollins, P., 1990. Factors determining differences in soil pH in adjacent conifer and alder-conifer stands. *Soil Science Society of America Journal* 54, 1417–1433.
- Binkley, D., Sollins, P., Bell, R., Sachs, D., Myrold, D., 1992. Biogeochemistry of adjacent conifer and alder-conifer stands. *Ecology* 73, 2022–2033.
- Bollmann, A., Schmidt, I., Saunders, A.M., Nicolaisen, M.H., 2005. Influence of starvation on potential ammonia-oxidizing activity and *amoA* mRNA levels of *Nitrosopirra briensis*. *Applied and Environmental Microbiology* 71, 1276–1282.
- Bottomley, P.J., Myrold, D.D., 2015. Biological and anthropogenic N inputs and their role in ecosystem services. In: Paul, E.A. (Ed.), *Soil Microbiology, Ecology and Biochemistry*. Academic Press, San Diego, CA, pp. 447–470.
- Boyle, S.A., Yarwood, R.R., Bottomley, P.J., Myrold, D.D., 2008. Bacterial and fungal contributions to soil nitrogen cycling under Douglas fir and red alder at two sites in Oregon. *Soil Biology and Biochemistry* 40, 443–451.
- Boyle-Yarwood, S., Bottomley, P.J., Myrold, D.D., 2008. Community composition of ammonia-oxidizing bacteria and archaea in soils under stands of red alder and Douglas fir in Oregon. *Environmental Microbiology* 10, 2956–2965.
- Burton, S.A.Q., Prosser, J.I., 2001. Autotrophic ammonia oxidation at low pH through urea hydrolysis. *Applied and Environmental Microbiology* 67, 2952–2957.
- De Boer, W., Gunnewiek, P.J., Veenhuis, M., Bock, E., Laanbroek, H.J., 1991. Nitrification at low pH by aggregated chemolithotrophic bacteria. *Applied and Environmental Microbiology* 57, 3600–3604.
- De Boer, W., Kowalchuk, G.A., 2001. Nitrification in acid soils: micro-organisms and mechanisms. *Soil Biology and Biochemistry* 33, 853–866.
- Dyrness, C., 2001. *Soil Descriptions and Data for Soil Profiles in the Andrews Experimental Forest, Selected Reference Stands, Research Natural Areas, and National Parks, 1962 & 1996*. Long-Term Ecological Research.
- Emmett, B.A., Boxman, D., Bredemeier, M., Gundersen, P., Kjonaas, O.J., Moldan, F., Schleppi, P., Tietema, A., Wright, R., 1998. Predicting the effects of atmospheric nitrogen deposition in conifer stands: evidence from the NITREX ecosystem-scale experiments. *Ecosystems* 1, 352–360.
- Gubry-Rangin, C., Hai, B., Quince, C., Engel, M., Thomson, B.C., James, P., Schlöter, M., Griffiths, R.I., Prosser, J.I., Nicol, G.W., 2011. Niche specialization of terrestrial archaeal ammonia oxidizers. *Proceedings of the National Academy of Sciences of the United States of America* 108, 21206–21211.
- Gubry-Rangin, C., Nicol, G.W., Prosser, J.I., 2010. Archaea rather than bacteria control nitrification in two agricultural acidic soils. *FEMS Microbiology Ecology* 74, 566–574.
- Gundersen, P., Callesen, I., de Vries, W., 1998. Nitrate leaching in forest ecosystems is related to floor forest C: N ratios. *Environmental Pollution* 102, 403–407.
- Hankinson, T.R., Schmidt, E.L., 1984. Examination of an acid forest soil for ammonia- and nitrite-oxidizing autotrophic bacteria. *Canadian Journal of Microbiology* 30, 1125–1132.
- Hart, S.C., Binkley, D., Perry, D.A., 1997. Influence of red alder on soil nitrogen transformations in two conifer forests of contrasting productivity. *Soil Biology and Biochemistry* 29, 1111–1123.
- Hood-Nowotny, R., Hink-Najera Uman, N., Inselbacher, E., Oswald-Lachoni, P., Wanek, W., 2010. Alternative methods for measuring inorganic, organic and total dissolved nitrogen in soil. *Soil Science Society of America Journal* 74, 1018–1027.
- Hu, H.W., Xu, Z.H., He, J.Z., 2014. Ammonia-oxidizing archaea play a predominant role in acid soil nitrification. In: Sparks, D.L. (Ed.), *Advances in Agronomy*, vol. 125, pp. 261–302.
- Hynes, R.K., Knowles, R., 1982. Effect of acetylene on autotrophic and heterotrophic nitrification. *Canadian Journal of Microbiology* 28, 334–340.
- Jiang, Q.Q., Bakken, L.R., 1999. Comparison of *Nitrosospora* strains isolated from terrestrial environments. *FEMS Microbiology Ecology* 30, 171–186.
- Kandeler, E., Gerber, H., 1988. Short-term assay of soil urease activity using colorimetric determination of ammonium. *Biology and Fertility of Soils* 6, 68–72.
- Könneke, M., Bernhard, A.E., de la Torre, J.R., Walker, C.B., Waterbury, J.B., Stahl, D.A., 2005. Isolation of an autotrophic ammonia-oxidizing marine archaeon. *Nature* 437, 543–566.
- Lehtovirta-Morley, L.E., Verhamme, D.T., Nicol, G.W., Prosser, J.I., 2013. Effect of nitrification inhibitors on the growth and activity of *Nitrosotalea devanterra* in culture and soil. *Soil Biology and Biochemistry* 62, 129–133.
- Lehtovirta-Morley, L.E., Stoecker, K., Vilcinskas, A., Prosser, J.I., Nicol, G.W., 2011. Cultivation of an obligate acidophilic ammonia oxidizer from a nitrifying acid soil. *Proceedings of the National Academy of Sciences of the United States of America* 108, 15892–15897.
- Leininger, S., Urich, T., Schlöter, M., Schwark, L., Qi, J., Nicol, G.W., Prosser, J.I., Schuster, S.C., Schleper, C., 2006. Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature* 442, 806–809.
- Levcik-Höfferle, S., Nicol, G.W., Ausec, L., Mandić-Mulec, I., Prosser, J.I., 2012. Stimulation of thaumarchaeal ammonia oxidation by ammonia derived from organic nitrogen but not added inorganic nitrogen. *FEMS Microbiology Ecology* 80, 114–123.
- Nicol, G.W., Leininger, S., Schleper, C., Prosser, J.I., 2008. The influence of soil pH on the diversity, abundance and transcriptional activity of ammonia oxidizing archaea and bacteria. *Environmental Microbiology* 10, 2966–2978.
- Norman, J.S., Barrett, J.E., 2014. Substrate and nutrient limitation of ammonia-oxidizing bacteria and archaea in temperate forest soil. *Soil Biology and Biochemistry* 69, 141–146.
- Perakis, S.S., Matkins, J.J., Hibbs, D.E., 2012. N_2 -fixing red alder indirectly accelerates ecosystem nitrogen cycling. *Ecosystems* 15, 1182–1193.
- Petersen, D.G., Blazewicz, S.J., Firestone, M., Herman, D.J., Turetsky, M., Waldrop, M., 2012. Abundance of microbial genes associated with nitrogen cycling as indices of biogeochemical process rates across a vegetation gradient in Alaska. *Environmental Microbiology* 14, 993–1008.
- Placella, S.A., Firestone, M.K., 2013. Transcriptional response of nitrifying communities to wetting of dry soil. *Applied and Environmental Microbiology* 79, 3294–3302.
- Prosser, J.I., Nicol, G.W., 2008. Relative contributions of archaea and bacteria to aerobic ammonia oxidation in the environment. *Environmental Microbiology* 10, 2931–2941.
- Prosser, J.I., Nicol, G.W., 2012. Archaeal and bacterial ammonia oxidisers in soil: the quest for niche specialisation and differentiation. *Trends in Microbiology* 20, 523–531.
- Radoosevich, S.R., Hibbs, D.E., Ghersa, C.M., 2006. Effects of species mixtures on growth and stand development of Douglas-fir and red alder. *Canadian Journal of Forest Research* 36, 768–782.
- Ross, D.S., Lawrence, G.B., Fredriksen, G., 2004. Mineralization and nitrification patterns at eight northeastern USA forested research sites. *Forest Ecology and Management* 188, 317–335.
- Rothauwe, J.H., Witzel, K.P., Liesack, W., 1997. The ammonia monooxygenase structural gene *amoA* as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing populations. *Applied and Environmental Microbiology* 63, 4704–4712.
- Spieck, E., Meincke, M., Bock, E., 1992. Taxonomic diversity of *Nitrosospora* strains isolated from building sandstones. *FEMS Microbiology Letters* 102, 21–26.

- Stephen, J.R., Kowalchuk, G.A., Bruns, M.-A.V., McCaig, A.E., Phillips, C.J., Embley, T.M., Prosser, J.I., 1998. Analysis of β -subgroup proteobacterial ammonia oxidizer populations in soil by denaturing gradient gel electrophoresis analysis and hierarchical phylogenetic probing. *Applied and Environmental Microbiology* 64, 2958–2965.
- Stein, L.Y., 2011. Heterotrophic nitrification and nitrifier denitrification. In: Ward, B.B., Arp, D.J., Klotz, M.G. (Eds.), *Nitrification*. ASM Press, Washington, DC, pp. 95–114.
- Stopnisek, N., Gubry-Rangin, C., Hofferle, S., Nicol, G.W., Mandic-Mulec, I., Prosser, J.I., 2010. Thaumarchaeal ammonia oxidation in an acidic forest peat soil is not influenced by ammonium amendment. *Applied and Environmental Microbiology* 76, 7626–7634.
- Taylor, A.E., Bottomley, P.J., 2006. Nitrite production by *Nitrosomonas europaea* and *Nitrosospora* sp. AV in soils at different solution concentrations of ammonium. *Soil Biology and Biochemistry* 38, 828–836.
- Taylor, A.E., Vajjala, N., Giguere, A.T., Gitelman, A.I., Arp, D.J., Myrold, D.D., Sayavedra-Soto, L., Bottomley, P.J., 2013. Use of aliphatic n-alkynes to discriminate soil nitrification activities of ammonia-oxidizing thaumarchaea and bacteria. *Applied and Environmental Microbiology* 79, 6544–6551.
- Taylor, A.E., Zeglin, L.H., Dooley, S., Myrold, D.D., Bottomley, P.J., 2010. Evidence for different contributions of archaea and bacteria to the ammonia-oxidizing potential of diverse Oregon soils. *Applied and Environmental Microbiology* 76, 7691–7698.
- Tourna, M., Freitag, T.E., Nicol, G.W., Prosser, J.I., 2008. Growth, activity and temperature responses of ammonia-oxidizing archaea and bacteria in soil microcosms. *Environmental Microbiology* 10, 1357–1364.
- Tourna, M., Stieglmeier, M., Spang, A., Könneke, M., Schintlmeister, A., Urich, T., Engel, M., Schlöter, M., Wagner, M., Richter, A., Schleper, C., 2011. *Nitrososphaera viennensis*, an ammonia oxidizing archaeon from soil. *Proceedings of the National Academy of Sciences of the United States of America* 108, 8420–8425.
- Verhamme, D.T., Prosser, J.I., Nicol, G.W., 2011. Ammonia concentration determines differential growth of ammonia-oxidising archaea and bacteria in soil microcosms. *ISME Journal* 5, 1067–1071.
- Vitousek, P.M., Gosz, J.R., Grier, C.C., Melillo, J.M., Reiners, W.A., 1982. A comparative analysis of potential nitrification and nitrate mobility in forest ecosystems. *Ecology Monographs* 52, 155–177. <http://dx.doi.org/10.2307/1942609>.
- Wertz, S., Leigh, A.K.K., Grayston, S.J., 2012. Effects of long-term fertilization of forest soils on potential nitrification and on the abundance and community structure of ammonia oxidizers and nitrite oxidizers. *FEMS Microbiology Ecology* 79, 142–154.
- Yao, H., Gao, Y., Nicol, G.W., Campbell, C.D., Prosser, J.I., Zhang, L., Han, W., Singh, B.K., 2011. Links between ammonia oxidizer community structure, abundance, and nitrification potential in acidic soils. *Applied and Environmental Microbiology* 77, 4618–4625.
- Yarwood, S.A., Bottomley, P.J., Myrold, D.D., 2010. Soil microbial communities associated with Douglas-fir and red alder stands at high- and low-productivity forest sites in Oregon, USA. *Microbial Ecology* 60, 606–617.
- Zeglin, L.H., Taylor, A.E., Myrold, D.D., Bottomley, P.J., 2011. Bacterial and archaeal *amoA* gene distribution covaries with soil nitrification properties across a range of land uses. *Environmental Microbiology Reports* 3, 717–726.
- Zhang, J.B., Muller, C., Zhu, T.B., Cheng, Y., Cai, Z.C., 2011. Heterotrophic nitrification is the predominant NO_3^- production mechanism in coniferous but not broad-leaf acid forest soil in subtropical China. *Biology and Fertility of Soils* 47, 533–542.
- Zhang, L.M., Hu, H.W., Shen, J.P., He, J.Z., 2012. Ammonia-oxidizing archaea have more important role than ammonia-oxidizing bacteria in ammonia oxidation of strongly acidic soils. *ISME Journal* 6, 1032–1045.