

Asymmetric hybridization between non-native winter moth, *Operophtera brumata* (Lepidoptera: Geometridae), and native Bruce spanworm, *Operophtera bruceata*, in the Northeastern United States, assessed with novel microsatellites and SNPs

N.P. Havill^{1*}, J. Elkinton², J.C. Andersen², S.B. Hagen³,
Hannah J. Broadley⁴, G.J. Boettner² and A. Caccone⁵

¹USDA Forest Service, Northern Research Station, Hamden, Connecticut, USA; ²Department of Environmental Conservation, University of Massachusetts, Amherst, Massachusetts, USA; ³NIBIO, Norwegian Institute for Bioeconomy Research, Svanvik, Norway; ⁴University of Massachusetts Amherst, Organismic and Evolutionary Biology, Amherst, Massachusetts, USA; ⁵Department of Ecology & Evolutionary Biology, Yale University, New Haven, Connecticut, USA

Abstract

The European winter moth, *Operophtera brumata*, is a non-native pest in the Northeastern USA causing defoliation of forest trees and crops such as apples and blueberries. This species is known to hybridize with *O. bruceata*, the Bruce spanworm, a native species across North America, although it is not known if there are hybrid generations beyond F1. To study winter moth population genetics and hybridization with Bruce spanworm, we developed two sets of genetic markers, single nucleotide polymorphisms (SNPs) and microsatellites, using genomic approaches. Both types of markers were validated using samples from the two species and their hybrids. We identified 1216 SNPs and 24 variable microsatellite loci. From them we developed a subset of 95 species-diagnostic SNPs and ten microsatellite loci that could be used for hybrid identification. We further validated the ten microsatellite loci by screening field collected samples of both species and putative hybrids. In addition to confirming the presence of F1 hybrids reported in previous studies, we found evidence for multi-generation asymmetric hybridization, as suggested by the occurrence of hybrid backcrosses with the winter moth, but not with the Bruce spanworm. Laboratory crosses between winter moth females and Bruce spanworm males resulted in a higher proportion of viable eggs than the reciprocal cross, supporting this pattern. We discuss the possible roles of population demographics, sex chromosome genetic incompatibility, and bacterial symbionts as causes of this asymmetrical hybridization and the utility of the developed markers for future studies.

*Author for correspondence
Phone: 203-230-4320
Fax: 203-230-4315
E-mail: nphavill@fs.fed.us

Keywords: hybridization, invasive species, microsatellite development, next generation sequencing, RAD-seq

(Accepted 24 August 2016; First published online 23 November 2016)

Introduction

The European winter moth, *Operophtera brumata* L. (Lepidoptera: Geometridae), is a non-native invasive pest in the Northeastern USA that is currently spreading and causing defoliation and tree mortality in the Northeastern USA (Elkinton *et al.*, 2010; Simmons *et al.*, 2014). In its native range of Europe, winter moth undergoes 9–10-year cyclical outbreaks (Tenow *et al.*, 2013). North America has experienced three major winter moth epidemics. The first one occurred in Nova Scotia, Canada, where it was detected in the 1930s and caused widespread damage to oak forests and apple orchards in the 1950s (Cuming, 1961). Two parasitoids from Europe were released as biological controls: a tachinid fly, *Cyzenis albicans* (Fallén), and an ichneumonid wasp, *Agrypon flaveolatum* (Gravenhorst). By 1961, winter moth populations collapsed, as parasitoid populations increased. To this day, winter moth remains at low density in Nova Scotia (Roland & Embree, 1995). The second winter moth outbreak occurred during the 1970s in the Pacific Northwest of Canada and the USA, where it became a serious pest of apples, hazelnuts, cherries, and blueberries (Kimberling *et al.*, 1986). This outbreak subsided after release of the same parasitoids used in the Nova Scotia outbreak (Roland & Embree, 1995).

The most recent winter moth outbreak was first detected in Massachusetts in the late 1990s. It has since spread to adjacent states and is currently causing persistent and widespread defoliation of forest trees and crops such as apples, blueberries, and cranberries (Elkinton *et al.*, 2015). The expansion of its range in the Northeastern USA was unexpected, given successful control in other regions. The leading edge of spread is moving approximately 6 km year⁻¹, and defoliation caused by winter moth is expanding (Elkinton *et al.*, 2014). It is not known whether this outbreak originated from an existing North American population, or if it represents a new introduction from Europe.

Winter moth is known to hybridize with a closely related North American species, *O. bruceata* (Hulst), the Bruce spanworm (Elkinton *et al.*, 2010). Bruce spanworm is common across the Northern USA and Canada, but it rarely experiences outbreaks (Rose & Lindquist, 1982). Winter moth and Bruce spanworm are difficult to distinguish based on wing patterns, but the males have distinct genital morphology (Eidt *et al.*, 1966). Pivnick *et al.* (1988) observed males with intermediate genitalia at several locations in British Columbia, as did Troubridge & Fitzpatrick (1993), who speculated that 0.4% of the moths they examined from British Columbia with intermediate morphological characters may be hybrids.

Elkinton *et al.* (2010) confirmed the presence of field collected hybrids in the Northeastern USA using DNA sequence data from one mitochondrial gene (cytochrome oxidase subunit I (COI)) and one nuclear gene (glucose-6-phosphate dehydrogenase (G6PD)). Hybrids were heterozygous for G6PD alleles fixed in each species, or had a COI haplotype characteristic of one species, and a G6PD allele characteristic of the other. Since the discovery of field-collected hybrids using this method, yearly sampling using pheromone traps that

attract both species and their hybrids found 32 confirmed hybrids out of 667 samples along an East–West transect across central Massachusetts (4.8%; Elkinton *et al.*, 2014). However, since this assay relies on only one nuclear gene, it cannot be determined whether these hybrids represent generations beyond F1, and if they do, whether there is asymmetry in the direction of backcrosses.

If multi-generational transfer of genes and adaptive traits between the introduced and native species is occurring (reviewed in Harrison & Larson, 2014), this could impact both the invasiveness of winter moth and the pest status of Bruce spanworm. Determining whether winter moth/Bruce spanworm F1 hybrids can reproduce will therefore help us decide whether more detailed studies of hybridization are warranted in this system.

In this study, we develop single nucleotide polymorphism (SNP) loci using Restriction-Associated DNA Sequencing (RAD-seq; Baird *et al.*, 2008) and microsatellite loci from shotgun genomic reads. Microsatellite markers are historically difficult to develop for Lepidoptera because microsatellites and their flanking regions tend to be duplicated in the genome (Zhang, 2004). We resolve this by filtering thousands of microsatellites from genomic reads to identify likely single copy markers. Microsatellite loci were then evaluated for utility in intra- and inter-specific population genetic studies of winter moth and Bruce spanworm and used to determine whether hybrid generations beyond F1 occur in the field using a group of moths collected in the Northeastern USA that had been previously classified as hybrids using G6PD (Elkinton *et al.*, 2014). We also performed laboratory crosses within and between the species to compare egg viability for the pure species versus their hybrids.

Materials and methods

RAD-seq SNP development

A RAD-seq library was prepared using DNA extracted from the head and thorax of 48 male moth samples using the DNeasy Blood & Tissue Kit (Qiagen) following manufacturer instructions. Individually barcoded samples included 23 Bruce spanworms, 19 winter moths, four putative hybrids, and two *Operophtera fagata* Scharfenberg samples included as an outgroup (Table S1). Library preparation and sequencing were performed by Floragenex Inc. (Eugene, OR) using the methods of Baird *et al.* (2008). Sequence reads for each sample were aligned with the software BWA (Li & Durbin, 2009) to the recently published winter moth genome (Derks *et al.*, 2015) accessed from NCBI BioProject (Accession Number PRJNA263715). The resulting alignments were used to identify SNPs with the software Stacks (Catchen *et al.*, 2013). Summary statistics were generated using a minimum read coverage of 4× and including only loci present in ≥75% of individuals. One random SNP was exported from each ‘stack’ for subsequent analysis. To evaluate whether individuals putatively identified as Bruce spanworm, winter moth, or hybrids could be identified with these SNPs, the program

fastStructure (Raj *et al.*, 2014) was used to calculate the probability of assignment (q) of individuals to $k = 2$ clusters.

To identify a subset of SNPs diagnostic for distinguishing winter moth and Bruce spanworm and their hybrids, the output from Stacks was filtered to exclude polymorphic sites that were within 50 bp of each other and sorted by allele frequencies to identify those that were fixed in Bruce spanworm and winter moth. For each candidate SNP, 300 bp of flanking sequence was exported and a single primer pair was designed for each locus using the Primer3 v. 2.3.4 plug-in (Untergasser *et al.*, 2012) in Geneious v. 8.1.6 (Kearse *et al.*, 2012), with the following specifications: primers at least 25 base pairs from the SNP, minimum product size of 100 bp, maximum product size of 300 bp, minimum T_m of 52°C, and minimum 20% GC content.

Microsatellite development

Genomic reads for microsatellite discovery were generated using an Ion Torrent Personal Genome Machine Sequencer (ThermoFisher). DNA was extracted from the head and thorax of a single male winter moth collected in a pheromone trap (see Elkinton, *et al.*, 2010 for trapping methods) in Lugo, Spain in December 2008 using the DNeasy Blood & Tissue Kit. 100 ng of DNA was sheared to approximately 300–500 bp fragments using Ion Shear enzymatic fragmentation and a library was prepared using the Ion Xpress Plus Library Kit (Life Technologies). Size selection was performed using a Pippin-Prep Blue (Sage Science), targeting 475 bp fragments. The library was prepared using the Ion PGM Sequencing 400 kit (Life Technologies). Sequencing was performed using 850 flows on an Ion 318 sequencing chip. Base-calling was performed with TorrentSuite 2.2 (Life Technologies).

Microsatellite discovery and primer design were performed using QDD 3.1.2 (Meglecz *et al.*, 2014). The default parameters were used except that the minimum sequence length was set to 200 bp. Finally, sequences with successful primer design were compared with the RepBase database of known transposable elements (Jurka, 2000) to identify potentially duplicated genomic regions. The 'best' primer pair per target sequence was selected as described in Meglecz *et al.*, (2014), based on primer alignment score, the distance from the microsatellite, and length of the PCR product. In addition, loci were selected that contained only pure microsatellites (not compound) with at least eight repeats, and fewer than ten reads in consensus sequences (to avoid duplicated loci).

The resulting primer pairs were tested for amplification success in four male winter moth samples collected in the Czech Republic, Spain, Scotland, and Italy, and four male Bruce spanworm samples from the USA collected in Washington, New Jersey, New York, and Pennsylvania. Reverse primers were modified with the addition of a 5' pig-tail sequence (GTTT) to reduce stutter (Brownstein *et al.*, 1996), and forward primers were modified with the addition of an M13 tail (TCCCAGTCACGACGT) to allow incorporation of a 6-FAM labeled M13 primer during PCR (Schuelke, 2000). PCR reactions (10 μ l) contained 1 $\times$ PCR buffer, 1.0 μ l dNTPs (10 mM each), 0.8 μ l $MgCl_2$ (25 mM), 0.1 μ l BSA (10 mg ml^{-1}), 0.025 μ l of forward primer (10 mM), 0.25 μ l of reverse primer (10 mM), 0.05 μ l of fluorescently labeled M13 primer (100 mM), 0.10 μ l Go Taq DNA polymerase (Promega), and 1.0 μ l sample DNA template. PCRs were amplified using a touch-down program with an annealing temperature that started at 61°C and decreased 2°C for each of the first five cycles,

then continued at 51°C for 30 cycles. PCR products were run on an ABI 3730 sequencer (Life Technologies) at the DNA Analysis Facility on Science Hill at Yale University with a Liz 500 internal size standard (Gel Company). Alleles were scored using the software Geneious.

Loci that had detectable peaks for at least three of four winter moth samples were further characterized using 24 male moths each from two winter moth populations collected with pheromone traps in Marcillac, France and Reinhardshagen, Germany. Primer pairs that also had readable peak patterns in Bruce spanworm were also characterized using 24 male moths each from two Bruce spanworm populations collected with pheromone traps in Harbor Creek, Pennsylvania and Oconto, Wisconsin. These populations were located approximately 600 and 1300 km, respectively, from the nearest known winter moth population (Elkinton *et al.*, 2010, 2014).

ARLEQUIN 3.5 (Excoffier *et al.*, 2005) was used to calculate allelic diversity, observed and expected heterozygosities, to test for Hardy–Weinberg equilibrium, to test for linkage disequilibrium, and to estimate differentiation between populations and species. Multiple comparisons were controlled using the method of Benjamini & Hochberg (1995) with a false discovery rate of 0.05. MICRO-CHECKER 2.2.3 (van Oosterhout *et al.*, 2004) was used to test for null alleles.

Finally, the samples from the four winter moth and Bruce spanworm populations were used to assess multiplexing combinations of 24 loci that amplified most consistently and were in Hardy–Weinberg equilibrium. The ten of these loci that amplified for both species were kept together for multiplexed combinations to allow for efficient genotyping of populations with potential hybrids.

Analysis of field-collected putative hybrids

To evaluate the ability to classify hybrids using the ten microsatellite loci that consistently co-amplified for winter moth and Bruce spanworm, hybrid genotypes were simulated with Hybridlab 1.0 (Nielsen *et al.*, 2006). Parental genotypes from 43 known Bruce spanworms from Maine, New York, and Pennsylvania, and 43 known winter moths from Connecticut, Massachusetts, Rhode Island, New Hampshire, and New York were used to generate 100 genotypes each of pure winter moth, pure Bruce spanworm, F1 hybrids, F2 hybrids, winter moth backcrosses, and Bruce spanworm backcrosses. A data set that included the 86 known samples, plus the 600 simulated samples was analyzed with Structure 2.3.4 (Pritchard *et al.*, 2000) and NewHybrids 1.1 (Anderson & Thompson, 2002). Structure was run with $k = 2$ clusters, known individuals coded as learning samples for updating allele frequencies, and 20,000 burn-in iterations followed by 100,000 sample iterations. NewHybrids was run with 10,000 burn-in iterations followed by 100,000 sample iterations, with known individuals identified. The probability of assignment was estimated for four different genotype frequency categories: each pure species, F1 hybrids, F2 hybrids, and backcross to each species.

Fifty-four putatively hybrid male moths collected with pheromone traps in Massachusetts, Rhode Island, Connecticut, and New Hampshire that were previously classified using the G6PD gene (Elkinton *et al.*, 2014) were genotyped using the ten microsatellite loci. Genetic ancestry and hybrid class of the putatively hybrid moths were analyzed using Structure and NewHybrids with the same run




Fig. 1. Bar plot showing individual assignment to Bruce spanworm or winter moth for the known pure species and putative hybrids analyzed with 1216 SNPs using fastStructure (Raj *et al.*, 2014). The height of each colored bar represents the proportion of an individual's genotype assigned to each species. The individual sample number is below each bar corresponding to Table S1.

conditions as above. The data set included the putative hybrids plus the 43 reference moths for each parental species.

For each putative hybrid, the 658 base pair 'DNA Barcoding' portion of the mitochondrial COI gene was amplified and sequenced using the primers LepF1 and LepR1 (Hebert *et al.*, 2004). Sequences were compared with known samples of each species (Gwiazdowski *et al.*, 2013) to determine the maternal lineage of each sample.

Laboratory crosses

Laboratory crosses were performed to produce pure winter moth, pure Bruce spanworm, and hybrid offspring with reciprocal female and male parents from each species. In 2014 and 2015, female moths were reared from field collected larvae, while males were either reared from larvae or collected as adults using pheromone traps. The larvae and adults were collected in Massachusetts on either side of the documented hybrid zone: winter moth in the east and Bruce spanworm in the west (Elkinton *et al.*, 2010). Larvae were reared in ventilated 5-gallon buckets with the tree foliage from which they were found. When they were in their last instar, sifted peat moss was added to the bottom of the buckets for pupation. The resulting pupae were sifted, sorted by sex, and stored at 10–12°C until adult emergence at the end of October to beginning of November.

Matings were set up in clear, 12-ounce cups lined with paper, in an incubator (Percival) set at 10–12°C with a 13.5 h dark to 10.5 h light schedule. One or two females were combined with 3–30 males, as available, as they emerged (median ratio of 2:5). Females laid eggs on the paper lining of the mating cups. The temperature in the incubator with the eggs was reduced to 9.5°C on 4 December, to 5°C on 12 December, and to 2°C on 18 January.

Egg condition was measured 4–5 weeks following mating. Eggs were scored according to color (orange or green) as an indication of viability (Peterson, 1962; du Merle & Brunet, 1991). Orange eggs were considered to have been fertilized and were classified as viable while green eggs were considered unfertilized and were classified as non-viable. This proved true as green eggs inevitably shriveled and died. Egg viability data were analyzed using a Chi-square (χ^2) test to determine

differences between observed and expected proportions of viable eggs among the four cross types using RStudio 0.98.501 (RStudio Team, 2015).

Results

RAD-seq SNP development

De-multiplexed reads from 46 of the 48 moth samples (sequencing failed for one winter moth and one putative hybrid sample) were aligned to the reference winter moth genome. A total of 92,852,319 sequence reads (mean of 2,018,528 reads per sample with mean coverage of 258×; Table S1) were analyzed with Stacks, which identified 11,779 stacks with 104,033 polymorphic SNPs. Filtering these loci to include one SNP per stack, and those with at least 4X coverage in at least 75% of individuals resulted in 1216 polymorphic loci. Analysis of these loci with fastStructure correctly identified all pure species individuals and identified two of the three putative hybrids as having mixed ancestry (fig. 1). The third putative hybrid was identified as pure Bruce spanworm.

Further filtering of the 1216 SNPs to those that were separated by >50 bp and fixed in each species identified 95 SNPs suitable for species-diagnostic marker development. A file with 300 bp of leading and trailing sequence centered on each SNP, and a list of the primer sequences with expected fragment size are available in the Supplementary Materials.

Microsatellite development

Genomic sequencing for microsatellite discovery resulted in 4,461,215 sequences after filtering for polyclonal and low-quality reads. Median sequence length was 373 bp. Raw sequence reads have been deposited in NCBI BioProject Accession Number PRJNA330716. A total of 3,551,925 sequences were at least 200 bp long, and 207,331 contained microsatellites. The sequence comparison and alignment step of analysis with QDD took 18.6 days to complete on a High Performance Computing Cluster at Yale University. This resulted in 14,491 singletons and 14,148 unique consensus sequences, for a total of 28,639 from which to design primer pairs. Primers were successfully designed for 17,141 of these. Stringent filtering resulted in 104 loci. Of these, 17 matched an

Table 1. Characterization of 32 microsatellite loci tested with 24 individuals each from winter moth populations from Reinhardshagen, Germany and Marcillac, France.

Locus	Motif	Size range	No. of alleles	Hardy–Weinberg equilibrium					
				Reinhardshagen			Marcillac		
				Ho	He	P	Ho	He	P
WM00070	AACAT	190–215	6	0.2500	0.5488	0.0009*	0.2857	0.5691	0.0013*
WM00485	ATC	302–323	8	0.4583	0.7305	0.0092*	0.4500	0.7628	0.0011*
WM00672	AC	318–360	14	0.6667	0.7943	0.3820	0.6667	0.8688	0.0011*
WM01184	ATC	243–267	7	0.4583	0.5089	0.1682	0.5417	0.5931	0.8647
WM01585	ACAT	174–234	12	0.7917	0.8245	0.6078	0.5833	0.7997	0.0624
WM01762	ATC	158–209	13	0.6364	0.8552	0.2259	0.5455	0.8235	0.0114
WM01835	ATC	239–266	6	0.2273	0.3594	0.0574	0.1667	0.3626	0.0002*
WM02191	AT	184–212	10	0.3684	0.7425	0.0021*	0.5556	0.7238	0.1295
WM02392	AG	172–178	4	0.5652	0.6541	0.8612	0.4546	0.5740	0.0691
WM02534	AC	272–308	15	0.4583	0.8236	0.0007*	0.5000	0.8014	0.0005*
WM02544	ATC	249–264	5	0.1739	0.2425	0.1995	0.3044	0.2773	1.0000
WM02565	ATC	188–209	8	0.4783	0.7845	0.0046*	0.7500	0.7385	0.5733
WM02953	AG	186–206	10	0.7500	0.6844	0.6758	0.9546	0.7981	0.0054*
WM03270	ATC	142–307	8	0.7083	0.7287	0.5558	0.4167	0.5319	0.0186
WM03280	ATC	278–317	10	0.6087	0.7092	0.5218	0.6522	0.7314	0.0597
WM03296	AT	150–156	4	0.2083	0.4149	0.0054*	0.2500	0.2988	0.0648
WM03463	AAG	199–214	5	0.4762	0.5563	0.6168	0.4546	0.5127	0.7954
WM03667	AC	300–342	14	0.7826	0.8899	0.3013	0.5000	0.8425	0.0031*
WM05159	ATC	289–325	12	0.7500	0.8564	0.4206	0.7500	0.7713	0.3714
WM12042	AC	248–366	30	0.9583	0.9140	0.6105	0.8261	0.8976	0.4007
WM13672	AC	124–130	4	0.2381	0.3752	0.0556	0.5217	0.7101	0.0366
WM16696	AC	246–264	6	0.3636	0.6607	0.0007*	0.5000	0.6507	0.0616
WM17017	AATG	240–276	6	0.1177	0.7825	0.0000*	0.0000	0.7196	0.0000*
WM17094	AC	248–264	7	0.2727	0.5539	0.0005*	0.2174	0.1981	1.0000
WM18760	AC	194–208	8	0.6667	0.7651	0.5057	0.7500	0.7598	0.1608
WM24011	AG	186–226	19	0.9546	0.9049	0.0673	0.9130	0.9034	0.9112
WM24441	ATC	136–148	4	0.5417	0.4619	0.8098	0.3636	0.4577	0.5315
WM24979	ACT	164–200	8	0.7083	0.7172	0.1632	0.5833	0.6551	0.2577
WM31399	AC	160–186	14	0.8696	0.8522	0.2842	0.7826	0.8908	0.1638
WM32817	AC	250–264	7	0.6667	0.7145	0.2703	0.6364	0.7495	0.0856
WM32864	AAG	304–316	5	0.2917	0.2677	1.0000	0.2500	0.2695	0.1488
WM33617	ATC	221–236	4	0.2917	0.2642	1.0000	0.1250	0.1215	1.0000

Asterisks indicate significant deviation from Hardy–Weinberg equilibrium after correcting for false discovery rate.

arthropod transposable element (12 from Lepidoptera) and were discarded. An additional primer pair was also discarded because the reverse primer had a TTTCTTT sequence on the 5' end that might have interfered with the pig-tail modification, leaving 86 remaining primer pairs to test. Forty-six of these had decipherable peak patterns, with one or two alleles in at least three of the four winter moth individuals tested, and were further characterized in two winter moth populations (tables 1 and 2). These loci included 22 dinucleotides, 20 trinucleotides, two tetranucleotides, and two pentanucleotides. Fourteen primer pairs also had decipherable peak patterns for at least three of the four Bruce spanworm individuals and were further characterized with two Bruce spanworm populations (table 2). Primer sequences are shown in Table S2.

The number of alleles per locus ranged from 4 to 30 in winter moth and 3 to 14 in Bruce spanworm (tables 1 and 2). Four loci deviated significantly from Hardy–Weinberg equilibrium in both of the winter moth populations, and ten loci deviated in one of the populations (tables 1 and 2), after controlling for false discovery rate. Two loci deviated from Hardy–Weinberg equilibrium in both of the Bruce spanworm populations, and two deviated in one of the populations (table 2). For winter moth, significant linkage was found for 19 out of 1035 pairwise

comparisons of loci in the Marcillac winter moth population, and three in the Reinhardshagen population after controlling for false discovery rate. WM03296 and WM12042 was the only pair that was linked in both populations. No loci were linked in either Bruce spanworm population. Seven loci showed evidence for null alleles in both winter moth populations, five in Reinhardshagen only, and four in Marcillac only. One locus showed evidence of null alleles in both Bruce spanworm populations, one in Harborcreek only, and one in Oconto only (Table S3).

The 24 loci that amplified most consistently for winter moth and were in Hardy–Weinberg equilibrium in at least one of the populations are indicated in Table S2. Using these 24 loci, genetic differentiation (F_{st}) between the two winter moth populations was 0.0067. Using the ten loci that amplified consistently in both species, F_{st} between the two winter moth populations was 0.0007, and between the two Bruce spanworm populations was 0.0130. Pairwise F_{st} values between populations of different species were: 0.3824 (Marcillac versus Harborcreek), 0.4030 (Marcillac versus Oconto), 0.3917 (Reinhardshagen versus Harborcreek), and 0.4112 (Reinhardshagen versus Oconto). When populations were combined ($N = 48$), F_{st} between species was 0.3952. Suitable PCR multiplex combinations for the 24 tested loci found are shown in Table S4.

Table 2. Characterization of 14 microsatellite loci tested with 24 individuals each from two winter moth populations from Reinhardshagen, Germany and Marcillac, France, and two Bruce spanworm populations from Harbortcreek, Pennsylvania and Oconto, Wisconsin.

Locus	Motif	Size range	No. of alleles	Hardy–Weinberg equilibrium					
				Winter moth			Bruce spanworm		
				Reinhardshagen			Harbortcreek		
				WM	BSw		Ho	He	P
							Ho	He	P
WM00863	ATC	145–187	9	3	0.4167	0.5656	0.0519	0.6812	0.0003*
WM00952	AG	161–375	28	14	0.7917	0.7846	0.7274	0.8118	0.0650
WM01619	ATC	182–215	5	5	0.6667	0.6215	0.6153	0.5652	0.3768
WM02232	AC	152–180	5	5	0.8750	0.5895	0.0075	0.3044	0.0357
WM02339	AC	110–440	10	5	0.6667	0.7385	0.4086	0.8333	0.3745
WM03475	AC	292–312	5	8	0.2500	0.2668	0.3949	0.3224	0.2290
WM07650	AC	203–243	10	10	0.7083	0.7952	0.2423	0.7391	0.3543
WM12853	ATC	242–290	8	7	0.6667	0.6853	0.2110	0.7500	0.3127
WM22363	ATC	303–321	4	5	0.2500	0.2278	1.0000	0.1818	1.0000
WM28247	AG	204–286	23	6	0.8750	0.8511	0.8969	0.9583	0.8747
WM29309	AC	162–188	13	5	0.9167	0.8360	0.0791	0.9167	0.9267
WM30125	ATC	296–314	7	6	0.4583	0.5993	0.0132	0.5000	0.1828
WM32985	ATC	263–290	5	4	0.3750	0.4229	0.0351	0.2609	0.1106
WM34463	AAATG	233–358	18	4	0.7500	0.8404	0.1303	0.7083	0.0524

Asterisks indicate significant deviation from Hardy–Weinberg equilibrium using after correcting for false discovery rate.

Analysis of field-collected putative hybrids

Structure analysis of simulated pure parental and hybrid genotypes (fig. 2) resulted in a mean probability of assignment (q) to Bruce spanworm of 0.97 for simulated Bruce spanworm genotypes, and 0.03 for simulated winter moth winter moth genotypes. Mean q values for simulated F1, F2, Bruce spanworm backcross, and winter moth backcross genotypes were 0.74, 0.51, 0.50, and 0.25, respectively. NewHybrids analysis resulted in simulated genotypes being assigned with the highest probability to the correct hybrid class for 100% of simulated F1s (mean probability = 0.92), 62% of simulated F2s (mean probability = 0.64), for 91% of simulated Bruce spanworm backcrosses (mean probability = 0.84), and 95% of simulated winter moth backcrosses (mean probability = 0.84).

Structure analysis of field-collected putative hybrids resulted in q values ranging from 0.009 to 0.991, indicating a mix of pure Bruce spanworm, hybrids, and pure winter moth (fig. 3; Table S5). Assignment of putative hybrids to pure species or hybrid classes using NewHybrids resulted in 6 moths assigned with highest probability as Bruce spanworm, three as winter moth, 24 as F1 hybrids, three as F2 hybrids, 18 as winter moth backcross, and none as Bruce spanworm backcrosses (Table S5).

Of the 45 moths classified with highest probability to a hybrid class, 30 had COI mitochondrial haplotypes matching winter moth and 15 had haplotypes matching Bruce spanworm (Table S5). Sequences have been deposited in GenBank (Accessions KX591003–KX591056). Broken down by hybrid class, 18 out of 24 F1 hybrids (75%), one out of three F2 hybrids (33%), and 11 out of 18 (61%) winter moth backcrosses had winter moth COI haplotypes.

Laboratory crosses

There was no significant difference in the proportion of viable eggs between years, so the data were combined for analysis. Sixty-two females laid 2262 eggs in the pure winter moth crosses (mean = 42.9 eggs/female), 39 females laid 1869 eggs in the pure Bruce spanworm crosses (mean = 47.9 eggs/female), 358 females laid 14,023 eggs in the winter moth female × Bruce spanworm male crosses (mean = 39.2 eggs/female), and six females laid 344 eggs in the Bruce spanworm female × winter moth male crosses (mean = 57.3 eggs/female). The numbers of crosses that include Bruce spanworm females were limited by the difficulty in collecting Bruce spanworm larvae, which are rare. There was a strong effect of cross type on egg viability ($\chi^2 = 2044.9$, $df = 3$, $P < 0.0001$). Pure winter moth and pure Bruce spanworm crosses had 93.5 and 94.1% viable eggs, respectively, while crosses between winter moth females and Bruce spanworm males had 60.8% viable eggs and crosses between Bruce spanworm females and winter moth males had 22.1% viable eggs (fig. 4). The latter two proportions were significantly different ($\chi^2 = 209.56$, $df = 1$, $P < 0.0001$).

Discussion

The SNP and microsatellite loci that we developed will be useful for studying population dynamics of winter moth in Europe, the introduction history of winter moth in North America, and spatial patterns of hybridization between winter moth and Bruce spanworm. The suite of SNP loci, together with the available draft winter moth genome (Derks *et al.*,




Fig. 2. Histograms of the probability of species assignment (q) using Structure (Pritchard *et al.*, 2000) for simulated pure species and select hybrid classes using ten microsatellite loci.




Fig. 3. Bar plot showing individual assignment to Bruce spanworm or winter moth for the known pure species and putative hybrids analyzed with ten microsatellite loci using Structure (Pritchard *et al.*, 2000). The height of each colored bar represents the proportion of an individual's genotype assigned to each species.




Fig. 4. Proportion of viable eggs by laboratory cross type between pure winter moths, pure Bruce spanworms, and their reciprocal female to male hybrids.

2015), could be especially useful for tracking selective introgression of particular regions of the genome.

Isolation of microsatellites by stringent filtering of genomic DNA reads has successfully identified a suite of 24 loci useful for winter moth population genetics. Ten of these loci were also shown to be suitable for characterizing hybrids between winter moth and Bruce spanworm through simulation of hybrid genotypes. Additional suitable loci could undoubtedly be developed by using less stringent filtering parameters than we used in this study.

Our study confirms the presence of field collected F1 hybrid moths and is the first to report later hybrid generations (F2 and winter moth backcross individuals), which opens up the possibility that selective introgression between the species could be impacting invasion. For example, hybridizing with a ubiquitous native species could relax Allee effects (the difficulty of finding mates) that limit population growth and establishment (Elkinton *et al.*, 2014). In addition, since Bruce spanworm can persist in colder regions than winter moth

(Roelofs *et al.*, 1982; Elkinton *et al.*, 2010), the transfer of alleles that confer cold tolerance could allow winter moth to invade colder areas than if they were not hybridizing.

Finally, the microsatellite data indicate a strong asymmetry among winter moth/Bruce spanworm hybrids in the Northeastern USA. Forty percent of the hybrids were classified as winter moth backcrosses and none were Bruce spanworm backcrosses. In addition, 75% of the moths classified as F1 hybrids had winter moth COI haplotypes (Table S5) indicating that it is more likely for winter moth females to successfully reproduce with Bruce spanworm males than the reciprocal cross. This result is consistent with Smith & Ring (unpublished, cited in Underhill *et al.*, 1987), who found that winter moth females and Bruce spanworm males from British Columbia could successfully hybridize in cages to at least the F2 generation, but that the reciprocal cross between Bruce spanworm females and winter moth males did not occur. Our data did not indicate a similarly complete directional incompatibility. This discrepancy could be explained by the Bruce spanworms in British Columbia observed by Smith & Ring being a different subspecies (*O. brumata occidentalis*) than those found on the East Coast of North America (Gwiazdowski *et al.*, 2013). The discrepancy could also be an artifact of the imperfect ability to assign individuals to a hybrid class. Among our simulated hybrid genotypes, 8% of the F2 hybrids were incorrectly classified as F1 hybrids (fig. 2), so some of the field collected moths that were classified as F1 hybrids could actually be F2. Laboratory crosses between winter moth females and Bruce spanworm males resulted in a higher proportion of viable eggs than the reciprocal cross (fig. 4), supporting this pattern, and implicating asymmetrical egg viability as an important source of mortality contributing to this pattern.

There are several possible causes for the observed asymmetrical, sex-biased hybridization between winter moth and Bruce spanworm. It could simply be a result of population demographics in the sampled region, where winter moths greatly outnumber Bruce spanworms during an outbreak (Elkinton *et al.*, 2015), making it less likely to sample Bruce spanworm backcrosses. Alternatively, there could be a sex-linked genetic incompatibility. Females are the heterogametic sex in Lepidoptera and there could be genes on the Bruce spanworm sex chromosome that are incompatible with genes on winter moth autosomes (Jiggins *et al.*, 2001). Finally, a symbiotic micro-organism, such as *Wolbachia*, could be causing sex-biased directional incompatibility. Bruce spanworms could harbor a

maternally transmitted symbiont that is incompatible with the winter moth genome or with a winter moth symbiont (Werren *et al.*, 2008). Sequencing the genome of a winter moth collected in the Netherlands detected *Wolbachia* DNA (Derks *et al.*, 2015), but it is not yet known whether it is present in North American populations of winter moth or Bruce spanworm.

The observation of multi-generation, asymmetrical hybridization between non-native winter moths and native Bruce spanworms, highlights the need to further study the causes and consequences of this pattern. For example, samples from across the ranges of both species and through hybrid zones can be used to test whether hybridization could be contributing to the unexpected new outbreak of winter moth in the Northeastern USA.

Acknowledgements

The authors gratefully acknowledge funding from the USDA Forest Service, Forest Health Technology Enterprise Team, and The Norwegian Institute for Bioeconomy Research (NIBIO). We are grateful to Jason Boone and Jenna Gribbon (Floragenex) for RAD-seq support and Ben Evans (Yale University) who helped with preliminary data analysis. We thank Frank Krueger (Nordwestdeutsche Forstliche Versuchsanstalt), Sandy Gardosik (Pennsylvania Department of Agriculture), Linda Williams (Wisconsin Department of Natural Resources), Alain Roques (Institut National de la Recherche Agronomique), Marek Turcáni (Czech University of Life Sciences), Maria Lombardero (University of Santiago de Compostela), Adam Vanbergen (NERC Centre for Ecology & Hydrology), Chad Phillips (Washington State Department of Agriculture), Saul Vaiciunas (New Jersey Department of Agriculture), Kenneth Carnes (New York State Department of Agriculture & Markets), and Andrea Battisti (University of Padova) for collecting the samples used in this study. We thank Ida Fløystad, Siv Grethe Aarnes, and DeAdra Newman for technical assistance.

Supplementary material

The supplementary material for this article can be found at <https://dx.doi.org/10.1017/S0007485316000857>.

References

- Anderson, E.C. & Thompson, E.A. (2002) A model-based method for identifying species hybrids using multilocus genetic data. *Genetics* **160**, 1217–1229.
- Baird, N.A., Etter, P.D., Atwood, T.S., Currey, M.C., Shiver, A.L., Lewis, Z.A., Selker, E.U., Cresko, W.A. & Johnson, E.A. (2008) Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PLoS ONE* **3**, e3376.
- Benjamini, Y. & Hochberg, Y. (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B, Methodological* **5**, 289–300.
- Brownstein, M.J., Carpten, J.D. & Smith, J.R. (1996) Modulation of non-templated nucleotide addition by tag DNA polymerase: primer modifications that facilitate genotyping. *Biotechniques* **20**, 1004–1010.
- Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A. & Cresko, W.A. (2013) Stacks: an analysis tool set for population genomics. *Molecular Ecology* **22**, 3124–3140.
- Cuming, F.G. (1961) The distribution, life history, and economic importance of the winter moth, *Operophtera brumata* (L.) (Lepidoptera, Geometridae), in Nova Scotia. *Canadian Entomologist* **93**, 135–142.
- Derks, M.F.L., Smit, S., Salis, L., Schijlen, E., Bossers, A., Mateman, C., Pijl, A.S., de Ridder, D., Groenen, M.A.M., Visser, M.E. & Megens, H.J. (2015) The genome of winter moth (*Operophtera brumata*) provides a genomic perspective on sexual dimorphism and phenology. *Genome Biology and Evolution* **7**, 2321–2332.
- du Merle, P. & Brunet, S. (1991) From green to yellow or yellowish white: egg-color changes in relation to oviposition rank in the fir budworm *Choristoneura murinana* (Hb) (Lep., Tortricidae). *Journal of Applied Entomology* **111**, 342–348.
- Eidt, D.C., Embree, D.G. & Smith, C.C. (1966) Distinguishing adults of winter moth *Operophtera brumata* (L.) and Bruce spanworm *O. bruceata* (Hulst) (Lepidoptera: Geometridae). *Canadian Entomologist* **98**, 258–261.
- Elkinton, J., Boettner, G., Liebhold, A. & Gwiazdowski, R. (2015) *Biology, Spread, and Biological Control of the Winter Moth in the Eastern United States*. Morgantown, WV, USDA Forest Service, Forest Health Technology Enterprise Team.
- Elkinton, J.S., Boettner, G.H., Sremac, M., Gwiazdowski, R., Hunkins, R.R., Callahan, J., Scheufele, S.B., Donahue, C.P., Porter, A.H., Khrimian, A., Whited, B.M. & Campbell, N.K. (2010) Survey for winter moth (Lepidoptera: Geometridae) in northeastern North America with pheromone-baited traps and hybridization with the native Bruce spanworm (Lepidoptera: Geometridae). *Annals of the Entomological Society of America* **103**, 135–145.
- Elkinton, J.S., Liebhold, A., Boettner, G.H. & Sremac, M. (2014) Invasion spread of *Operophtera brumata* in northeastern United States and hybridization with *O. bruceata*. *Biological Invasions* **16**, 2263–2272.
- Excoffier, L., Laval, G. & Schneider, S. (2005) Arlequin ver. 3.0: an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* **1**, 47–50.
- Gwiazdowski, R.A., Elkinton, J.S., deWaard, J.R. & Sremac, M. (2013) Phylogeographic diversity of the winter moths (Lepidoptera: Geometridae: Operophtera) *O. brumata*, and *O. bruceata* in Europe and North America. *Annals of the Entomological Society of America* **106**, 143–151.
- Harrison, R.G. & Larson, E.L. (2014) Hybridization, introgression, and the nature of species boundaries. *Journal of Heredity* **105**, 795–809.
- Hebert, P.D.N., Penton, E.H., Burns, J.M., Janzen, D.H. & Hallwachs, W. (2004) Ten species in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astrartes fulgerator*. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 14812–14817.
- Jiggins, C.D., Linares, M., Naisbit, R.E., Salazar, C., Yang, Z.H. & Mallet, J. (2001) Sex-linked hybrid sterility in a butterfly. *Evolution* **55**, 1631–1638.
- Jurka, J. (2000) Repbase Update: a database and an electronic journal of repetitive elements. *Trends in Genetics* **9**, 418–420.
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Mentjies, P. & Drummond, A. (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–1649.
- Kimberling, D.N., Miller, J.C. & Penrose, R.L. (1986) Distribution and parasitism of winter moth, *Operophtera brumata* (Lepidoptera, Geometridae), in western Oregon. *Environmental Entomology* **15**, 1042–1046.

- Li, H. & Durbin, R. (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760.
- Meglec, E., Pech, N., Gilles, A., Dubut, V., Hingamp, P., Trilles, A., Grenier, R. & Martin, J.F. (2014) QDD version 3.1: a user-friendly computer program for microsatellite selection and primer design revisited: experimental validation of variables determining genotyping success rate. *Molecular Ecology Resources* **14**, 1302–1313.
- Nielsen, E.E., Bach, L.A. & Kotlicki, P. (2006) HYBRIDLAB (version 1.0): a program for generating simulated hybrids from population samples. *Molecular Ecology Notes* **6**, 971–973.
- Peterson, A. (1962) Some eggs of insects that change color during incubation. *The Florida Entomologist* **45**, 81–87.
- Pivnick, K.A., Barton, D.L., Millar, J.G. & Underhill, E.W. (1988) Improved pheromone trap exclusion of the Bruce spanworm *Operophtera bruceata* (Hulst) (Lepidoptera: Geometridae) when monitoring winter moth *Operophtera brumata* (L.) populations. *Canadian Entomologist* **120**, 389–396.
- Pritchard, J.K., Stephens, M. & Donnelly, P. (2000) Inference of population structure using multilocus genotype data. *Genetics* **155**, 945–959.
- Raj, A., Stephens, M. & Pritchard, J.K. (2014) fastSTRUCTURE: variational inference of population structure in large SNP data sets. *Genetics* **197**, 573–589.
- Roelofs, W.L., Hill, A.S., Linn, C.E., Meinwald, J., Jain, S.C., Herbert, H.J. & Smith, R.F. (1982) Sex pheromone of the winter moth, a geometrid with unusually low-temperature pre-copulatory responses. *Science* **217**, 657–659.
- Roland, J. & Embree, D.G. (1995) Biological control of the winter moth. *Annual Review of Entomology* **40**, 475–492.
- Rose, A.H. & Lindquist, O.H. (1982) *Insects of Eastern Hardwood Trees*. Canadian Forest Service Publication, Forestry Technical Report 29. Ottawa, Natural Resources Canada.
- RStudio Team (2015) *RStudio: Integrated Development for R*. Boston, MA, RStudio, Inc. Available online at <http://www.rstudio.com/>.
- Schuelke, M. (2000) An economic method for the fluorescent labeling of PCR fragments. *Nature Biotechnology* **18**, 233–234.
- Simmons, M.J., Lee, T.D., Ducey, M.J. & Dodds, K.J. (2014) Invasion of winter moth in New England: effects of defoliation and site quality on tree mortality. *Forests* **5**, 2440–2463.
- Tenow, O., Nilssen, A.C., Bylund, H., Pettersson, R., Battisti, A., Bohn, U., Carouille, F., Ciornei, C., Cs Ka, G., Delb, H., De Prins, W., Glavendekic, M., Gninenko, Y.I., Hrasovec, B., Matosevic, D., Meshkova, V., Moraal, L., Netoiu, C., Pajares, J., Rubtsov, V., Tomescu, R., Utkina, I. & Gurney, W. (2013) Geometrid outbreak waves travel across Europe. *Journal of Animal Ecology* **82**, 84–95.
- Troubridge, J.T. & Fitzpatrick, S.M. (1993) A revision of the North American *Operophtera* (Lepidoptera, Geometridae). *Canadian Entomologist* **125**, 379–397.
- Underhill, E.W., Millar, J.G., Ring, R.A., Wong, J.W., Barton, D. & Giblin, M. (1987) Use of a sex attractant and an inhibitor for monitoring winter moth and Bruce spanworm populations. *Journal of Chemical Ecology* **13**, 1319–1330.
- Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B. C., Remm, M. & Rozen, S.G. (2012) Primer3-new capabilities and interfaces. *Nucleic Acids Research* **40**, e115.
- van Oosterhout, C., Hutchinson, W.F., Wills, D.P.M. & Shipley, P. (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**, 535–538.
- Werren, J.H., Baldo, L. & Clark, M.E. (2008) *Wolbachia*: master manipulators of invertebrate biology. *Nature Reviews Microbiology* **6**, 741–751.
- Zhang, D.X. (2004) Lepidopteran microsatellite DNA: redundant but promising. *Trends in Ecology & Evolution* **19**, 507–509.