

A remotely sensed pigment index reveals photosynthetic phenology in evergreen conifers

John A. Gamon^{a,b,c,1}, K. Fred Huemmrich^d, Christopher Y. S. Wong^{e,f}, Ingo Ensminger^{e,f,g}, Steven Garrity^h, David Y. Hollingerⁱ, Asko Noormets^j, and Josep Peñuelas^{k,l}

^aCenter for Advanced Land Management Information Technologies, School of Natural Resources, University of Nebraska–Lincoln, Lincoln, NE 68583; ^bDepartment of Earth and Atmospheric Sciences, University of Alberta, Edmonton, AB, Canada T6G 2E3; ^cDepartment of Biological Sciences, University of Alberta, Edmonton, AB, Canada T6G 2E9; ^dNASA Goddard Space Flight Center, University of Maryland, Baltimore County, Greenbelt, MD 20771; ^eDepartment of Biology, University of Toronto, Mississauga, ON, Canada L5L1C6; ^fGraduate Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada M5S 1A1; ^gGraduate Department of Cell and Systems Biology, University of Toronto, Toronto, ON, Canada M5S 3G5; ^hDecagon Devices, Inc., Pullman, WA 99163; ⁱUS Forest Service, Northern Research Station, Durham, NH 03824; ^jDepartment of Forestry and Environmental Resources, North Carolina State University, Raleigh, NC 27695; ^kConsejo Superior de Investigaciones Científicas (CSIC), Global Ecology Unit, Center for Ecological Research and Forestry Applications (CREAF)-CSIC-Campus de Bellaterra, Bellaterra 08193, Catalonia, Spain; and ^lCREAF, Cerdanyola del Vallès 08193, Catalonia, Spain

Edited by Christopher B. Field, Carnegie Institution of Washington, Stanford, CA, and approved September 29, 2016 (received for review April 17, 2016)

In evergreen conifers, where the foliage amount changes little with season, accurate detection of the underlying “photosynthetic phenology” from satellite remote sensing has been difficult, presenting challenges for global models of ecosystem carbon uptake. Here, we report a close correspondence between seasonally changing foliar pigment levels, expressed as chlorophyll/carotenoid ratios, and evergreen photosynthetic activity, leading to a “chlorophyll/carotenoid index” (CCI) that tracks evergreen photosynthesis at multiple spatial scales. When calculated from NASA’s Moderate Resolution Imaging Spectroradiometer satellite sensor, the CCI closely follows the seasonal patterns of daily gross primary productivity of evergreen conifer stands measured by eddy covariance. This discovery provides a way of monitoring evergreen photosynthetic activity from optical remote sensing, and indicates an important regulatory role for carotenoid pigments in evergreen photosynthesis. Improved methods of monitoring photosynthesis from space can improve our understanding of the global carbon budget in a warming world of changing vegetation phenology.

carotenoid pigments | evergreen conifers | gross primary productivity | chlorophyll/carotenoid index | CCI

The biosphere helps regulate atmospheric composition and climate, in part through the exchange of radiatively active gases, primarily carbon dioxide. About half of the “extra” carbon added to the atmosphere by human activity is rapidly absorbed by the biosphere, effectively slowing climate change relative to what would occur without this uptake (1, 2). The exact mechanism and spatiotemporal distribution of the terrestrial component of this carbon sink have been ongoing research topics for many years. In a warming world, the timing of photosynthetic activity is also changing, with unknown impacts on ecosystem productivity. These shifting patterns of seasonal photosynthetic activity, or “photosynthetic phenology,” affect the biospheric-atmospheric gas exchange, further influencing atmospheric composition and climate (3–5). Faced with these uncertainties, quantifying the spatial and temporal patterns of biosphere/atmosphere carbon fluxes for different biomes and understanding their proximal controls remain central goals of global carbon cycle science.

Northern forests make a large contribution to global photosynthetic carbon fixation and are an important component of the global carbon budget. However, northern evergreen conifers, including evergreen conifers of the vast boreal regions, present particular challenges to global carbon cycle monitoring (6). Their seasonal activity may be changing with an earlier growing season, with important implications for the biospheric carbon budget. A simple hypothesis has been that a longer growing season results in greater carbon uptake, particularly for northern ecosystems where

photosynthetic activity has been temperature-limited (3, 7). By contrast, warmer growing seasons are also more likely to cause drought, restricting ecosystem carbon uptake and enhancing ecosystem respiration, resulting in accelerated carbon losses to the atmosphere (8). The actual outcome of changing seasonality on the biospheric carbon budget remains an open question, with multiple factors likely to be important.

A primary tool for assessing terrestrial carbon uptake has been eddy covariance, which provides near-direct assessment of surface/atmosphere carbon fluxes. Although it provides an excellent means of sampling the gas exchange of representative ecosystems for limited regions (9), it must be supplemented by other less costly and more spatially extensive methods for global carbon flux assessments. Remote sensing provides an ideal means of extrapolating flux measurements beyond the sampling footprint of individual flux towers to larger regions. Accurate methods of tracking photosynthetic phenology using remote sensing are critical to a full understanding of the impact of climate variation on terrestrial gross primary productivity (GPP) and to a proper assessment of the global carbon budget. With global, daily satellite coverage, we now have the means of generating wall-to-wall images of photosynthetic carbon uptake and net primary productivity for virtually the entire planet. For example, the MOD17 algorithm based on the light-use efficiency (LUE) model (10) states that daily GPP, or gross primary production, is a product of absorbed radiation (*APAR*) and the

Significance

Evergreen photosynthetic activity has been difficult to determine from remote sensing, causing errors in terrestrial photosynthetic carbon uptake models. Using a reflectance chlorophyll/carotenoid index (CCI) sensitive to seasonally changing chlorophyll/carotenoid pigment ratios, we demonstrate a method of tracking photosynthetic phenology in evergreen conifers. The CCI reveals seasonally changing photosynthetic rates and detects the onset of the growing season in evergreen foliage. This method could improve our understanding of changing photosynthetic activity in a warming climate, and could improve assessment of the evergreen component of the terrestrial carbon budget, which has been elusive.

Author contributions: J.A.G., K.F.H., and C.Y.S.W. designed research; J.A.G., K.F.H., and C.Y.S.W. performed research; J.A.G., K.F.H., C.Y.S.W., D.Y.H., and A.N. analyzed data; and J.A.G., K.F.H., C.Y.S.W., I.E., S.G., D.Y.H., A.N., and J.P. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. Email: jgamon@gmail.com.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1606162113/-DCSupplemental.

efficiency (ε) with which this absorbed energy is converted to fixed carbon:

$$GPP = APAR \times \varepsilon \quad [1]$$

$APAR$, in turn, is a product of irradiance in the photosynthetically active radiation (PAR) region of the spectrum (i.e., 400–700 nm) and the fraction of that PAR irradiance that is absorbed by green vegetation (f_{APAR}):

$$APAR = f_{APAR} \times PAR \quad [2]$$

The f_{APAR} term is closely related to the normalized difference vegetation index (NDVI), a measure of vegetation greenness, and the PAR term is typically obtained from meteorological data (10, 11).

For much of the world's deciduous or annual vegetation, where the seasonal expression of photosynthetic activity closely follows green canopy display, the $APAR$ term largely captures the seasonal photosynthetic dynamics (11). However, evergreen vegetation poses particular challenges for the LUE model. For evergreens that retain their foliage through the seasons, the f_{PAR} component of the $APAR$ term is relatively stable compared with f_{PAR} for annual or deciduous vegetation, providing insufficient information on the actual seasonal dynamics of photosynthetic activity. Assessing ε has been more challenging than measuring $APAR$ from remote sensing, in part because ε is highly variable in time and space (12). Particularly for northern evergreen conifers that experience periods of photosynthetic down-regulation during winter dormancy or chronic stress, ε emerges as an important determinant of seasonal photosynthetic activity (13). Conventional vegetation indices (e.g., NDVI) that are responsive to changes in green leaf area typically fail to detect these seasonal photosynthetic dynamics (12, 14), leading to errors in satellite-driven ecosystem productivity models. Not surprisingly, although the MOD17 algorithm depicts broad-scale seasonal dynamics of photosynthetic activity for different biomes, it sometimes fails to reproduce faithfully annual sums or seasonal patterns of ecosystem photosynthetic activity measured by flux towers (15), and it does not capture local heterogeneity in ε (16).

Due to these challenges, many studies have sought to improve the LUE model by assessing ε directly from remote sensing (12). One approach involves remote detection of the xanthophyll cycle, which adjusts the distribution of absorbed light energy through the interconversion of a group of three carotenoid pigments (17). This biochemical response can be detected with proximal remote sensing using the photochemical reflectance index (PRI), which provides a direct method of assessing changing LUE over short time spans (18–20). However, over annual time periods, the primary driver of the PRI at leaf and canopy scales is the changing leaf carotenoid pigment pool, typically expressed as the changing ratio of chlorophyll to carotenoid pigments (or its inverse), and not the xanthophyll cycle per se (21–25). This observation of changing pigment content is particularly relevant for northern evergreen conifers that undergo large seasonal swings in photosynthetic activity and carotenoid pigment levels associated with temperature extremes (24–28). These findings have led us to reconsider the function of carotenoid pigments in the context of detecting photosynthetic phenology in evergreen conifers.

Carotenoids serve many important roles in plants. They function as light capture and photoprotective pigments; act as antioxidants; and are linked to the synthesis of isoprenoids, which are produced under high-temperature stress (29). Carotenoid levels increase when plants are exposed to a variety of environmental stressors. This finding leads to the hypothesis that enhanced carotenoid levels, typically expressed relative to chlorophyll content, provide optical indicators of reduced photosynthetic activity and LUE. A large body of literature indicates that chlorophyll/carotenoid ratios decrease in evergreen leaves during winter cold (27, 30–32). Field studies have confirmed that chlorophyll/

carotenoid ratios closely track the seasonal photosynthetic activity of Mediterranean climate evergreen leaves (21, 23). This conclusion is further supported by recent spectral and kinetic evidence that demonstrates a close coincidence between seasonal photosynthetic activity, chlorophyll/carotenoid ratios, and spectral reflectance in evergreen conifers (24–26). Together, these studies suggest that satellite indices of changing pigment levels might provide reliable indicators of evergreen photosynthetic phenology.

In this study, we consider whether newly available combinations of satellite bands from NASA's Moderate Resolution Imaging Spectroradiometer (MODIS) sensors can provide useful metrics of seasonal changing pigment levels and photosynthetic activity in evergreen conifers. Our analysis was made possible by the recent reprocessing of MODIS data (MODIS collection 6) that provides both ocean and land band surface reflectance products over terrestrial areas, affording new options for assessing evergreen photosynthetic activity. Because the original PRI bands (33, 34) are not available from the MODIS, we considered new band combinations (MODIS bands 1 and 11) indicative of chlorophyll/carotenoid ratios, and used leaf- and stand-level experiments to confirm the interpretation of these bands independently.

Results

In conifers, reflectance from evergreen leaves exhibits distinct seasonal changes, particularly in the green-red spectral region (550–650 nm) (Fig. 1). The wintertime increase in reflectance at these wavelengths indicates a decreased chlorophyll/carotenoid pigment ratio, often visible to a careful observer as a wintertime needle yellowing. MODIS bands 11 (531 nm) and 1 (645 nm) captured the contrasting behavior of evergreen leaf reflectance in these two spectral regions (Fig. 1). MODIS band 1 (645 nm, a terrestrial band) is clearly affected by the increased red reflectance, whereas band 11 (531 nm, an ocean band) changes little or undergoes a slight decline in reflectance during winter.

Parallel to the seasonal leaf reflectance changes (Fig. 1), the leaves exhibited substantial photosynthetic down-regulation during winter, detectable as changes in reflectance, pigment levels, and photosynthetic rates (Fig. 2). Leaf chlorophyll/carotenoid ratios declined gradually in fall and winter, and then recovered rapidly in spring (Fig. 2). Photosynthetic rates exhibited a similar rapid spring increase (Fig. 2). The chlorophyll/carotenoid index (CCI) responded near synchronously to these seasonal changes in pigment ratios and photosynthetic activity, both at the leaf and stand level (Fig. 2).

We then examined the CCI generated from satellite-derived surface reflectance for several evergreen-dominated North




Fig. 1. Needle reflectance spectra of *Pinus contorta* (lodgepole pine) seedlings in Edmonton, Alberta, Canada, exposed to a boreal winter (red line, March 19, 2013) and summer (black line, June 21, 2013). The positions of MODIS bands 1 and 11 are indicated in gray.

Although we describe these stands as predominantly “evergreen” conifer stands, they actually included varying deciduous components that likely contributed to variation in CCI and NDVI responses across stands (Table S1). This varying stand composition and levels of deciduousness helps explain the contrasting seasonal CCI and NDVI patterns for these sites (Fig. 3). These possibilities can be further investigated by independent ground studies characterizing physiological responses (as shown in Figs. 1 and 2), stand composition, and latitudinal (including angular) effects on the optical signals. Currently, there is a growing network of ground optical validation sites located at flux towers (38–40), and these sites can be equipped to test the relative merits of the NDVI, PRI, and CCI further for monitoring ecosystem photosynthesis. Such independent ground validation is needed because any two-band index can be affected by multiple factors, often causing misinterpretation of underlying biophysical traits or processes when satellite data are used alone. To clarify the contribution of these multiple factors, we propose that extended ground sampling networks using flux tower sites be used in future studies of changing photosynthesis and productivity from global satellite sensors. Despite these remaining questions, the strongly similar seasonal responses across the three sites and across spatial scales shown here, along with similar findings from Mediterranean evergreen vegetation (36), suggest that the CCI offers a widely applicable indicator of photosynthetic activity and GPP for evergreen-dominated ecosystems. We note, however, that our study does not consider all evergreen-dominated biomes (e.g., tropical moist broadleaved forests, where satellite and flux data observations are more limited).

More work is now needed to reconsider the parameterization of the LUE model in light of these findings, which could help develop new upscaling approaches. A current working hypothesis is that the NDVI, PRI, and CCI provide complementary information about photosynthetic phenology. The NDVI closely follows seasonal photosynthetic activity related to green leaf display in annual and deciduous vegetation, whereas the CCI and PRI add additional information about photosynthetic regulatory processes involving photoprotective pigments and appear to be particularly useful in evergreens (28). Like the CCI, the PRI responds to seasonal pigment shifts associated with photosynthetic down-regulation in evergreens (24, 25), but it is also sensitive to short periods of down-regulation mediated by the xanthophyll cycle (e.g., midday photosynthetic depression due to summer drought) (41). In the future, LUE model parameterizations should explicitly recognize that these two indices reveal contrasting effects on photosynthetic activity over different time scales (28).

Recently, a similar ability to track photosynthetic phenology from satellites has been reported using solar-induced fluorescence (SIF), which provides an alternate method of assessing photosynthetic activity from remote sensing (6, 42). Because the CCI index is based on reflectance, which has a well-established methodology and history, it can provide a strong foundation for interpreting the causes of changes in SIF. However, SIF is not currently available at a temporal or spatial scale comparable to the MODIS, making a direct comparison with a MODIS-derived CCI difficult without considerable data aggregation and associated loss of spatial or temporal resolution that would conceal the underlying mechanisms addressed here. Combined measurement of SIF and pigment indices has been proposed recently as part of the FLuorescence EXplorer (FLEX) satellite mission (43), and could lead to an improved assessment of photosynthetic phenology, particularly for evergreens that have been difficult to measure from satellites using more conventional vegetation indices (e.g., the NDVI). To fully understand the significance of plant pigments as functional indicators, a full spectral (“hyperspectral”) satellite imaging spectrometer would be ideal. Such sensors are available on airborne platforms (44) and as demonstration satellite missions (45), none of which provide frequent global coverage; and thus cannot properly assess diurnal or seasonal photosynthetic dynamics for large regions of the planet.

Proposed satellite-based imaging spectrometers (43, 46) could further our understanding of these functionally different pigment responses for different ecosystems, opening further unseen possibilities for monitoring photosynthesis from space.

This study extends our previous understanding of carotenoid pigments by relating their seasonal patterns to the seasonal patterns of reflectance and photosynthesis in evergreen conifers. The CCI derived from a new combination of MODIS bands 1 and 11 offers a promising tool for observing photosynthetic phenology of terrestrial ecosystems. If properly coupled with field validation, such measurements could greatly improve our understanding of the changing photosynthetic phenology of evergreen stands that have been difficult to assess with conventional satellite indices. Assessment of evergreen pigment activity can improve our ability to measure carbon cycle dynamics for this important component of the world’s vegetation.

Methods

Sites and Plant Materials. Leaf- and stand-scale measurements were conducted on potted 4-y-old lodgepole pine (*Pinus contorta*) seedlings grown outdoors under ambient conditions in full sun at the University of Alberta campus (Edmonton, Alberta, Canada; 53.5289°N, 113.5261°W). Plants were arranged in closed-canopy stands. Initially planted in 2010 in 2.3-L pots, they were repotted in subsequent years into 2.8-L and 6.2-L pots in a 1:2 soil mixture of sandy top soil and potting soil (Sunshine Mix 4; Sun Gro Horticulture). The seedlings were irrigated throughout the growing season to avoid drought stress. Stand-scale eddy covariance measurements (described below) were obtained from sites spanning a wide range of species, edaphic, climatic, geographic location, and canopy structural conditions (Table S1).

Leaf Pigment Measurements. Leaves were collected immediately after leaf reflectance measurements from the same six trees sampled every 1–2 wk. The sampled leaves were frozen in liquid nitrogen and later transferred into a –80 °C freezer for long-term storage. For each sampling date, six 1-cm-long segments from each seedling were pooled together for pigment analysis using HPLC (1260 Infinity; Agilent Technologies) following the method of Thayer and Björkman (47) to determine carotenoid and chlorophyll pigment quantities. Commercial pigment standards (DHI LAB Products) were used to calibrate the HPLC system for pigment quantification. Chlorophyll/carotenoid pigment ratios were determined from total chlorophyll (a and b) and total carotenoids (neoxanthin, violaxanthin, antheraxanthin, zeaxanthin, lutein, and beta-carotene concentrations).

Leaf Gas Exchange Measurements. Foliar gas exchange was measured using a portable photosynthesis system (LI-6400; LI-COR). Each measurement consisted of a light-response curve ranging from 0 to 1,500 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$. Measurements were taken at each light level when steady-state photosynthesis was obtained, typically within 3 min. The chamber CO_2 concentration was set to 395 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$, and temperature was set to match ambient conditions. Photosynthetic rate was expressed as light-saturated photosynthesis, which was estimated from the saturation point of the light response curves. The same six plants from the reflectance measurements were monitored on a weekly basis during spring recovery of photosynthetic activity.

Flux Data. Daily GPP values were retrieved from the FLUXNET “LaThuile” dataset (www.fluxdata.org/default.aspx) for three sites (Table S1). These values were calculated from eddy covariance measurements that were filtered, despiked, and gap-filled, and the net ecosystem exchange was partitioned into daily GPP and ecosystem respiration following standard algorithms (48, 49).

Leaf Spectral Measurement. Leaf-scale reflectance was measured with a portable spectrometer (UniSpec SC; PP Systems) equipped with a bifurcated fiber optic (UNI410; PP Systems) and needle leaf clip (UNI501; PP Systems). Leaves of six plants (five randomly selected, sunlit leaves per plant) were monitored through the course of the study. Leaves were sampled at around 1300 hours under sunlight every 1–2 wk. Each set of leaf measurements for a given plant consisted of an average of 10 samples and was preceded by a dark scan and a reference scan from a Spectralon white standard (Lab-sphere) to obtain reflectance values.

Whole-Stand Spectral Measurements. Reflectance was also measured on experimental stands of the same seedlings used for leaf reflectance with a portable spectrometer (UniSpec-DC; PP Systems). An upward-facing fiber

(UNI686; PP Systems) attached to a cosine receptor (UNI435; PP Systems) measured incoming irradiance. A downward-facing fiber (UNI684; PP Systems) fitted with a field-of-view (FOV) restrictor (UNI688; PP Systems) limited the FOV to 15° for measuring stand radiance. For each measurement date, 12 measurements were taken from a position ~0.5 m above the canopy, each covering a different location of the stand at around 1330 hours every week during spring recovery. Each set of canopy measurements was preceded by a dark scan and a reference scan from a white standard panel (Spectralon; Labsphere) to obtain reflectance values.

MODIS Satellite Data. Optical data from MODIS onboard the *Aqua* satellite were processed to surface reflectance using the multi-angle implementation of atmospheric correction (MAIAC) algorithm (50, 51). The MAIAC algorithm makes use of time-series and spatial analyses for cloud screening and aerosol retrievals (52, 53). MAIAC terrestrial surface reflectances were gridded at a resolution of 1 km, and both MODIS land and ocean bands were processed (MODIS bands 1–12), providing reflectance in spectral bands that have previously been used on a limited basis for terrestrial applications (16, 37, 54). In this analysis, a time series of surface reflectance values was extracted for the 1-km grid that included the location of the flux tower. All cloud-free observations with high data quality and view zenith angles less than 45° were used.

CCI Calculations. From the MODIS satellite sensor, the CCI was calculated from reflectance using MODIS bands 1 and 11 reflectance (ρ) as $(\rho_{B11} - \rho_{B1})/(\rho_{B11} + \rho_{B1})$. From leaf- and stand-level reflectance, the reflectance spectra were first interpolated to 1-nm intervals. MODIS band reflectance values were then simulated by convolving the 1-nm reflectance values against the MODIS

bandpass responses for bands 1 and 11. The CCI was then calculated using the simulated MODIS bands 1 and 11, respectively.

Statistical Analysis. Regression statistics and analyses of covariance were conducted in R (R Studio).

ACKNOWLEDGMENTS. We thank Dave Landis (GST, Inc.) for MODIS data processing. J.A.G. was supported by the Alberta Informatics Circle of Research Excellence/Alberta Innovates Technology Futures (Grants G224150012 and 200700172), Natural Sciences and Engineering Research Council (NSERC) (Grant RGPIN-2015-05129), Canada Foundation for Innovation (CFI) (Grant 26793), and NASA Arctic-Boreal Vulnerability Experiment (Grant NNX15AT78A). K.F.H. was supported by the NASA Science of Aqua and Terra program (Grant NNX14AJ65G). I.E. was supported by the NSERC (Grant RGPIN-2015-06514), CFI (Grant 27330), and Ontario Ministry of Research and Innovation (Grant ER10-07-015). J.P. was supported by European Research Council Synergy Grant SyG-2013-610028 IMBALANCE-P. Data and logistical support for the Wind River site were provided by the US Forest Service (FS) Pacific Northwest Research Station and the University of Washington (Grant 11-JV-11261952-030). Support for the Howland Forest AmeriFlux core site was provided by the US Department of Energy (DOE)'s Office of Science and the US Department of Agriculture (USDA) Forest Service Northern Research Station. The work at the Parker Tract, NC (US-NC2) site was supported by USDA FS Eastern Forest Environmental Threat Assessment Center Cooperative Agreements 03-CA-11330147-073 and 04-CA-11330147-238, DOE-National Institute for Climate Change Research (NICCR) Award 08-SC-NICCR-1072, Department of the Interior (DOI) Southeast Climate Science Center Award G10AC00624, DOE Terrestrial Ecosystem Science Award 11-DE-SC-0006700, and USDA NIFA Awards 2011-68002-30185 and 2014-67003-22068.

- Canadell JG, et al. (2007) Contributions to accelerating atmospheric CO₂ growth from economic activity, carbon intensity, and efficiency of natural sinks. *Proc Natl Acad Sci USA* 104(47):18866–18870.
- Raupach MR, et al. (2014) The declining uptake rate of atmospheric CO₂ by land and ocean sinks. *Biogeosciences* 11:3453–3475.
- Randerson JT, Field CB, Fung IY, Tans PP (1999) Increases in early season ecosystem uptake explain recent changes in the seasonal cycle of atmospheric CO₂ at high northern latitudes. *Geophys Res Lett* 26:2765–2768.
- Peñuelas J, Rutishauser T, Filella I (2009) Ecology. Phenology feedbacks on climate change. *Science* 324(5929):887–888.
- Richardson AD, et al. (2013) Climate change, phenology, and phenological control of vegetation feedbacks to the climate system. *Agriculture and Forest Meteorology* 169:156–173.
- Walther S, et al. (2016) Satellite chlorophyll fluorescence measurements reveal large-scale decoupling of photosynthesis and greenness dynamics in boreal evergreen forests. *Glob Change Biol* 22(9):2979–2996.
- Myneni RB, Keeling CD, Tucker CJ, Asrar G, Nemani RR (1997) Increased plant growth in the northern high latitudes from 1981 to 1991. *Nature* 386:698–702.
- Trahan MW, Schubert BA (2016) Temperature-induced water stress in high-latitude forests in response to natural and anthropogenic warming. *Glob Change Biol* 22(2):782–791.
- Baldocchi DD, Hicks BB, Meyers TP (1988) Measuring biosphere-atmosphere exchanges of biologically related gases with micrometeorological methods. *Ecology* 69:1331–1340.
- Heinsch FA, et al. (2006) Evaluation of remote sensing based terrestrial productivity from MODIS using regional tower eddy flux network observations. *IEEE Trans Geosci Remote Sens* 44:1908–1925.
- Running SW, et al. (2004) A continuous satellite-derived measure of global terrestrial primary production. *Bioscience* 54:547–560.
- Garbulsky MF, et al. (2010) Patterns and controls of the variability of radiation use efficiency and primary productivity across terrestrial ecosystems. *Glob Ecol Biogeogr* 19:253–267.
- Garbulsky MF, Peñuelas J, Gamon J, Inoue Y, Filella I (2011) The photochemical reflectance index (PRI) and the remote sensing of leaf, canopy and ecosystem radiation use efficiencies: A review and meta-analysis. *Remote Sens Environ* 115:281–297.
- Gamon JA, et al. (1995) Relationships between NDVI, canopy structure, and photosynthesis in three Californian vegetation types. *Ecol Appl* 5:28–41.
- Turner DP, et al. (2005) Site-level evaluation of satellite-based global terrestrial gross primary production and net primary production monitoring. *Glob Change Biol* 11:666–684.
- Drolet GG, et al. (2008) Regional mapping of gross light-use efficiency using MODIS spectral indices. *Remote Sens Environ* 112:3064–3078.
- Demmig-Adams B, Adams WW (1996) Chlorophyll and carotenoid composition in leaves of *Euonymus kiautschovicus* acclimated to different degrees of light stress in the field. *Aust J Plant Physiol* 23:649–659.
- Gamon JA, Peñuelas J, Field CB (1992) A narrow-waveband spectral index that tracks diurnal changes in photosynthetic efficiency. *Remote Sens Environ* 41:35–44.
- Gamon JA, Serrano L, Surfus JS (1997) The photochemical reflectance index: An optical indicator of photosynthetic radiation use efficiency across species, functional types, and nutrient levels. *Oecologia* 112:492–501.
- Peñuelas J, Garbulsky MF, Filella I (2011) Photochemical reflectance index (PRI) and remote sensing of plant CO₂ uptake. *New Phytol* 191(3):596–599.
- Stylinski CD, Gamon JA, Oechel WC (2002) Seasonal patterns of reflectance indices, carotenoid pigments and photosynthesis of evergreen chaparral species. *Oecologia* 131:366–374.
- Nakaji T, Oguma H, Fujinuma Y (2006) Seasonal changes in the relationship between photochemical reflectance index and photosynthetic light use efficiency of Japanese larch needles. *Int J Remote Sens* 27:493–509.
- Filella I, et al. (2009) PRI assessment of long-term changes in carotenoids/chlorophyll ratio and short-term changes in de-epoxidation state of the xanthophyll cycle. *Int J Remote Sens* 30:4443–4455.
- Wong CYS, Gamon JA (2015) The photochemical reflectance index provides an optical indicator of spring photosynthetic activation in evergreen conifers. *New Phytol* 206(1):196–208.
- Wong CYS, Gamon JA (2015) Three causes of variation in the photochemical reflectance index (PRI) in evergreen conifers. *New Phytol* 206(1):187–195.
- Fréchette E, Wong CYS, Junker LV, Chang CY-Y, Ensminger I (2015) Zeaxanthin-independent energy quenching and alternative electron sinks cause a decoupling of the relationship between the photochemical reflectance index (PRI) and photosynthesis in an evergreen conifer during spring. *J Exp Bot* 66(22):7309–7323.
- Ensminger I, et al. (2004) Intermittent low temperatures constrain spring recovery of photosynthesis in boreal Scots pine forests. *Glob Change Biol* 10:995–1008.
- Gamon JA (2015) Reviews and syntheses: Optical sampling of the flux tower footprint. *Biogeosciences* 12:4509–4523.
- Peñuelas J, et al. (2013) Photochemical reflectance index as an indirect estimator of foliar isoprenoid emissions at the ecosystem level. *Nat Commun* 4:2604.
- Adams WW, III, Demmig-Adams B (1994) Carotenoid composition and down regulation of photosystem II in three conifer species during the winter. *Physiol Plant* 92:451–458.
- Ottander C, Campbell D, Oquist G (1995) Seasonal changes in photosystem II organization and pigment composition in *Pinus sylvestris*. *Planta* 197:176–183.
- Oquist G, Huner NPA (2003) Photosynthesis of overwintering evergreen plants. *Annu Rev Plant Biol* 54:329–355.
- Gamon JA, Filella I, Peñuelas J (1993) The dynamic 531-nanometer Δ reflectance signal: A survey of twenty angiosperm species. *Photosynthetic Responses to the Environment*, eds Yamamoto HY, Smith CM (American Society of Plant Physiologists, Rockville, MD), pp 172–177.
- Peñuelas J, Filella I, Gamon JA (1995) Assessment of photosynthetic radiation-use efficiency with spectral reflectance. *New Phytol* 131:291–296.
- Jahns P, Holzwarth AR (2012) The role of the xanthophyll cycle and of lutein in photoprotection of photosystem II. *Biochim Biophys Acta* 1817(1):182–193.
- Garbulsky MF, Peñuelas J, Ogaya R, Filella I (2013) Leaf and stand-level carbon uptake of a Mediterranean forest estimated using the satellite-derived reflectance indices EVI and PRI. *Int J Remote Sens* 34:1282–1296.
- Goerner A, Reichstein M, Rambal S (2009) Tracking seasonal drought effects on ecosystem light use efficiency with satellite-based PRI in a Mediterranean forest. *Remote Sens Environ* 113:1101–1111.
- Gamon JA, Rahman AF, Dungan JL, Schildhauer M, Huemmrich KF (2006) Spectral Network (SpecNet)—What is it and why do we need it? *Remote Sens Environ* 103:227–235.
- Gamon JA, et al. (2010) SpecNet revisited: Bridging flux and remote sensing communities. *Can J Remote Sens* 36:5376–5390.
- Porcar-Castell A, et al. (2015) EUROSPEC: At the interface between remote-sensing and ecosystem CO₂ flux measurements in Europe. *Biogeosciences* 12:6103–6124.
- Ripullone F, et al. (2011) Effectiveness of the photochemical reflectance index to track photosynthetic activity over a range of forest tree species and plant water statuses. *Funct Plant Biol* 38:177–186.

42. Joiner J, et al. (2014) The seasonal cycle of satellite chlorophyll fluorescence observations and its relationship to vegetation phenology and ecosystem atmosphere carbon exchange. *Remote Sens Environ* 152:375–391.
43. Kraft S, et al. (2013) FLORIS: Phase A status of the Fluorescence Imaging Spectrometer of the Earth Explorer mission candidate FLEX. *Proc SPIE* 8889(88890T):1–12.
44. Green RO, et al. (1998) Imaging spectroscopy and the Airborne Visible Infrared Imaging Spectrometer (AVIRIS). *Remote Sens Environ* 65:227–248.
45. Middleton EM, et al. (2013) The Earth Observing One (EO-1) satellite mission: Over a decade in space. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* 6:243–256.
46. Lee CM, et al. (2015) An introduction to the NASA Hyperspectral InfraRed Imager (HyspIRI) mission and preparatory activities. *Remote Sens Environ* 167:6–19.
47. Thayer SS, Björkman O (1990) Leaf Xanthophyll content and composition in sun and shade determined by HPLC. *Photosynth Res* 23(3):331–343.
48. Papale D, et al. (2006) Towards a standardized processing of Net Ecosystem Exchange measured with eddy covariance technique: Algorithms and uncertainty estimation. *Biogeosciences* 3:571–583.
49. Reichstein M, et al. (2005) On the separation of net ecosystem exchange into assimilation and ecosystem respiration: Review and improved algorithm. *Glob Change Biol* 11:1424–1439.
50. Lyapustin A, Martonchik J, Wang Y, Laszlo I, Korkin S (2011) Multiangle implementation of atmospheric correction (MAIAC): 1. Radiative transfer basis and look-up tables. *J Geophys Res Atmos* 116, 10.1029/2010jd014985.
51. Lyapustin A, et al. (2011) Multiangle implementation of atmospheric correction (MAIAC): 2. Aerosol algorithm. *J Geophys Res Atmos* 116, 10.1029/2010jd014986.
52. Lyapustin A, Wang Y, Frey R (2008) An automatic cloud mask algorithm based on time series of MODIS measurements. *J Geophys Res Atmos* 113, 10.1029/2007jd00964.
53. Lyapustin A, Wang Y, Laszlo I, Korkin S (2012) Improved cloud and snow screening in MAIAC aerosol retrievals using spectral and spatial analysis. *Atmos Meas Tech* 5: 843–850.
54. Rahman AF, Cordova VD, Gamon JA, Schmid HP, Sims DA (2004) Potential of MODIS ocean bands for estimating CO₂ flux from terrestrial vegetation: A novel approach. *Geophys Res Lett* 31:L10503.
55. Hollinger DY, et al. (1999) Seasonal patterns and environmental control of carbon dioxide and water vapour exchange in an ecotonal boreal forest. *Glob Change Biol* 5:891–902.
56. Noormets A, et al. (2010) Response of carbon fluxes to drought in a coastal plain loblolly pine forest. *Glob Change Biol* 16:272–287.
57. Wharton S, Falk M, Bible K, Schroeder M, Paw UKT (2012) Old-growth CO₂ flux measurements reveal high sensitivity to climate anomalies across seasonal, annual and decadal time scales. *Agric For Meteorol* 161:1–14.