

Insect outbreak shifts the direction of selection from fast to slow growth rates in the long-lived conifer *Pinus ponderosa*

Raul de la Mata^{a,b,1,2}, Sharon Hood^{a,c}, and Anna Sala^a

^aDivision of Biological Sciences, University of Montana, Missoula, MT 59812; ^bDepartment of Renewable Resources, University of Alberta, Edmonton, AB, Canada T6G 2H1; and ^cRocky Mountain Research Station, Fire, Fuel, and Smoke Science Program, US Department of Agriculture Forest Service, Missoula, MT 59808

Edited by Ian T. Baldwin, Max Planck Institute for Chemical Ecology, Jena, Germany, and approved May 19, 2017 (received for review January 2, 2017)

Long generation times limit species' rapid evolution to changing environments. Trees provide critical global ecosystem services, but are under increasing risk of mortality because of climate change-mediated disturbances, such as insect outbreaks. The extent to which disturbance changes the dynamics and strength of selection is unknown, but has important implications on the evolutionary potential of tree populations. Using a 40-y-old *Pinus ponderosa* genetic experiment, we provide rare evidence of context-dependent fluctuating selection on growth rates over time in a long-lived species. Fast growth was selected at juvenile stages, whereas slow growth was selected at mature stages under strong herbivory caused by a mountain pine beetle (*Dendroctonus ponderosae*) outbreak. Such opposing forces led to no net evolutionary response over time, thus providing a mechanism for the maintenance of genetic diversity on growth rates. Greater survival to mountain pine beetle attack in slow-growing families reflected, in part, a host-based life-history trade-off. Contrary to expectations, genetic effects on tree survival were greatest at the peak of the outbreak and pointed to complex defense responses. Our results suggest that selection forces in tree populations may be more relevant than previously thought, and have implications for tree population responses to future environments and for tree breeding programs.

fluctuating selection | growth-survival trade-offs | selection response | *Pinus ponderosa* | *Dendroctonus ponderosae*

Understanding the dynamics of selection over time is fundamental for understanding life-history evolution (1) and predicting evolutionary change under climate change (2, 3). To date, such understanding is based almost exclusively on data for relatively short-lived species (4, 5), but virtually nonexistent for long-lived organisms, such as trees (ref. 6; but see ref. 7). Trees and forests provide critical ecological and commercial services, including impacts on global carbon cycles, species diversity, water quality, and climate regulation (8). Recent forest mortality (9, 10) highlights the importance of understanding how climate change and climate change-driven disturbances may impact forests (11, 12). Trees may live for hundreds of years and experience varying selection pressures associated with fluctuating climate (13), disturbance regimes (14), and biotic interactions (15), all of which may be magnified under climate change. The extent to which these events may change the strength and direction of selection and contribute to the maintenance of genetic diversity and evolutionary potential is unknown. Of special relevance are insect outbreaks, a biotic interaction expected to increase with climate change (16) but unaccounted for in models to predict the evolutionary potential of tree populations (17, 18).

Mountain pine beetle (MPB; *Dendroctonus ponderosae* Hopkins) is a native, irruptive forest insect in western North America that uses numerous *Pinus* species as hosts. Via pheromone-mediated mass attacks that overcome host defenses, the MPB kills the host in the process of reproducing and completing its life cycle (19). Attacks have the binary outcome of either a successful mass

attack/dead tree or unsuccessful attack/surviving tree, and mechanisms of tolerance as common in other plant-herbivore systems (20) are not involved. MPB outbreaks have recently caused unprecedented mortality in pine forests (21) and have been linked to warming effects on insect development and increased drought stress of hosts (19). Evidence that tree-killing insect outbreaks exert strong selective pressures, potentially changing selection patterns in tree populations and their underlying basis, is rare. Such information is critical to evaluate the evolutionary potential of tree populations under climate change and increasing incidence of insect outbreaks (17).

How insect outbreaks affect selection is important because relative fitness contribution of phenotypic traits changes over time in response to environmental change (2) and ontogenetic development (22). Conflicting selection pressures during ontogeny are common and often arise because of trade-offs among traits (23, 24). For example, growth rate has clear fitness benefits but also incurs intrinsic and extrinsic costs (23, 25). In trees, fast growth rates and large size may provide fitness benefits via increased competitive ability, faster time to reproduction, and increased chances of early survival (26). Fast growth rates are particularly important during early, vulnerable ontogenetic stages when rapid growth minimizes exposure time to mortality agents and suppression by neighbors, and allows a size advantage to monopolize resources (27). However, consistent with results in animals (28), fast early growth in trees has been associated

Significance

Understanding selection dynamics over time is fundamental for predicting evolutionary potential. This is critical in long-lived organisms, particularly trees, which are at increased risk of mortality because of climate change-related drought and associated stresses. For ponderosa pine, we show rare direct empirical evidence of strong context-dependent fluctuating selection on growth rates over time: fast growth was selected for before a mountain pine beetle outbreak but against after the outbreak, when a growth-survival trade-off emerged. Our results provide insight into the mechanisms contributing to the maintenance of genetic diversity in a long-lived organism, and show that strong and fluctuating selection can contribute to the evolutionary potential of trees under climate change.

Author contributions: R.d.l.M., S.H., and A.S. designed research; R.d.l.M., S.H., and A.S. performed research; R.d.l.M. analyzed data; and A.S., S.H., and R.d.l.M. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: The data reported in this paper have been deposited in the US Forest Service Research Data Archive (<https://doi.org/10.2737/RDS-2017-0026>).

¹Present address: Research Institute of Food Technology and Agriculture-IRTA, Fruit Production Program, Torre Marimon, 08140 Caldes de Montbui, Spain.

²To whom correspondence should be addressed. Email: delamatapombo@yahoo.es.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1700032114/-DCSupplemental.

with shorter lifespan (29), with long-term consequences on lifetime reproductive output. Thus, selection pressures for fast growth at early life stages may conflict with fitness costs at later stages (29, 30). These conflicts may explain the substantial intraspecific genetic variation in growth rates observed both within and among natural populations of organisms (23).

Another potential cost of fast growth is decreased defense against consumers (31, 32). Growth-defense trade-offs could contribute to changes in magnitude or even reversals of directional selection over a tree's lifespan, depending on the intensity and timing of herbivory (6). For example, tree-killing pests can exert strong selective pressures favoring better-defended (33), slower-growing trees (30), and dampen or reverse directional selection for fast growth. Therefore, earlier selection for fast growth but later context-dependent selection for slow growth could contribute to the maintenance of genetic diversity in growth rates (1, 23). Understanding such pressures and their consequences on selection requires following growth dynamics over the entire lifespan of the organisms in question and its consequences on fitness; such data are rarely available for long-lived woody plants (34).

We took advantage of an existing common garden genetic experiment of ponderosa pine (*Pinus ponderosa* Lawson and C. Lawson var. *ponderosa* C. Lawson) that was impacted by a MPB outbreak when the trees were approximately 40 y old. We asked the following main question: Does strong herbivory pressure caused by an insect outbreak at the mature stage change the strength and direction of selection on growth rates relative to that in juvenile stages? We also asked: Does this change arise because of a growth-survival trade-off? Defenses against MPB are critical when beetle populations are low to moderate, but their role is thought to decrease at the epidemic stage when mass attacks can overwhelm even well-defended individuals (35). Although resistance to bark beetles is under genetic control (33, 36), it is not known whether genetic effects on tree resistance change during the course of the outbreak. Therefore, we also tested if these effects change over the outbreak. We hypothesized positive selection for fast growth at juvenile stages, but negative selection under strong herbivory pressure at mature stages as a result of a growth-survival trade-off. We also hypothesized stronger genetic effects on tree resistance earlier during the outbreak relative to the peak of the outbreak, when the overwhelming effect of mass attacks at high MPB pressure may render defenses irrelevant.

Results

Survival Over Time. Overall survival 38 y after planting was 41%. Sixty-three percent of trees planted survived before the outbreak, with 64% of those surviving the outbreak (Fig. S1). Before the outbreak, survival was lowest up to age 2 y (78%) and progressively increased to $\geq 97\%$ for each interval after age 11 y. During the outbreak, survival was lowest at ages 35 and 36 y (85% of the living trees each year) (Fig. S1). At age 36 y, 106 elite trees (identified based on superior growth at age 27 y) were sprayed with insecticide for protection against the MPB. Sprayed trees exhibited much higher survival (93%) compared with background survival (64%; $P < 0.05$) (SI Text). However, spraying effects were only significant on target sprayed trees and not surrounding trees (SI Text and Figs. S2–S4), and patterns of heritability of survival (see below) did not change after deleting these trees from the analyses (SI Text and Fig. S5).

The MPB Outbreak Reversed Patterns of Selection. The MPB outbreak strongly changed the survival ranking of families relative to before the outbreak [significant genotype by environment (G×E) interaction; χ^2 for log-likelihood comparison between full and restricted model: $P < 0.001$]. On average, family survival rank changed 76 positions (of the 204 families in total) from before to after the outbreak (Fig. 1), with two extreme cases where family rank changed 192 and 189 positions (Fig. 1). Before the outbreak, genetic correlations between survival at a given sampling age and survival at the previous sampling age were almost equal to one (Fig. S6). Therefore, families most likely to survive in a specific year were also most likely to survive the following year.




Fig. 1. Family survival rank before and after beetle outbreak for 204 families (lower rank means higher survival). Family ranks were estimated based on survival 27 y after planting (before) and by accounting mortality during the outbreak (after). In bold are drastic rank changes of two families.

In contrast, at the onset of the outbreak (from age 34 y) genetic correlations decreased significantly, particularly from ages 34–36 y, indicating a change in the genetic patterns of mortality (the MPB started to kill trees from families that did not necessarily exhibit low survival before the outbreak). Note that we used cumulative survival, so that potential changes in the genetic correlations between two successive sampling ages are only because of the killed trees between both ages. Given that after age 2 y survival rates were above 92% between every two successive ages (Fig. S1), small reductions in the genetic correlations indicate significant changes in the genetic patterns of survival.

The MPB outbreak caused a switch in selection patterns on intrinsic growth potential [measured as diameter at breast height (DBH) at age 27 to prevent confounding growth responses to reduced competition because of MPB-caused mortality]. Although a positive genetic response to selection in growth rate occurred before the outbreak from age 2–4 y, the genetic response was negative from age 35–36 y, the period of highest mortality during the outbreak (Fig. 2A and Fig. S1). The negative selection during the outbreak was one-third of the positive selection at early stages, but it became stronger when the entire outbreak was considered a single selection event (Fig. 2A). Similarly, cumulative responses were positive during most of the MPB-free period but became nonsignificant after age 34 y, indicating no evolutionary response in the long term (Fig. 2B). Despite the positive genetic response before the outbreak, phenotypic selection differentials were not statistically different from zero, regardless of whether they were calculated on a period-by-period basis or on a cumulative basis (Fig. 2). However, from the onset of the outbreak yearly selection differentials were significantly negative (i.e., preferential selection of slow-growing phenotypes), causing cumulative differentials to also become negative from age 35 y and later. When the outbreak was considered a single selection event, the selection differential was significant and was larger than that at any period before the outbreak (Fig. 2A). This finding indicates a strong environmental effect on strength and direction of selection during the outbreak.

A Growth-Survival Trade-Off Emerged During the MPB Outbreak. Before the outbreak, surviving trees were larger than trees that died, whereas the reverse was true after the outbreak (Fig. 3).




Fig. 2. Responses to selection and selection differentials over time for intrinsic growth potential (diameter at age 27 y) based on (A) fitness (survival) at each sampling time (black lines, closed and open circles, respectively) and considering the outbreak as a single period (gray lines, closed and open triangles, respectively) and (B) cumulative fitness. Bars are 95% credibility intervals (for clarity, credibility intervals before the outbreak are only shown for responses to selection). An asterisk (*) denotes significant selection responses; an "x" denotes significant selection differentials.

However, sprayed trees (a subset of faster growing trees) exhibited faster growth both before and after the outbreak, indicating that when MPB was excluded, fast growth was not selected against during the outbreak (actual selection estimates within sprayed trees were not possible because of small sample size). Consistently, faster growing families exhibited greater survival before the outbreak (significant positive genetic correlation between survival at the end of the MPB-free period and growth potential) (Fig. 4A), but lower survival after the outbreak (significant negative genetic correlation) (Fig. 4B), indicating a growth–survival trade-off. The pattern before the outbreak was driven by positive genetic correlations from ages 0–2 and 2–4 y (Fig. 4C and Fig. S7), whereas that after the onset of the outbreak was driven by a negative genetic correlation from ages 35–36 y (Fig. 4D and Fig. S7), the period with highest survival heritability (Fig. 5D).

Genetic Effects Were Strongest at the Peak of the Outbreak but Were Delayed. Heritability estimates for growth since planting were low to moderate ($h_i^2 = 0.19$ – 0.47) and tended to decrease with time, with the largest decrease during the first years of life (between ages 4 and 11 y) and stabilizing thereafter (Fig. 5A). Heritability for survival since planting was consistently lower than for growth ($h_i^2 = 0.11$ – 0.14) (Fig. 5B). The genetic effect on survival to MPB attack (i.e., resistance) measured at age 34–38 y was much stronger than that on survival over the entire period. Individual heritabilities for cumulative survival during the outbreak were always significant after age 34 y, with estimates starting at low values at age 34 y ($h_i^2 = 0.07$) and reaching a maximum at age 37 y ($h_i^2 = 0.34$) (Fig. 5C). On an annual basis during the outbreak, the genetic effect on beetle resistance varied over time, and became significant at ages 35 and 36 y, coinciding with maximum estimated heritability of 0.35 at age 36 y (Fig. 5D).

Discussion

High genetic diversity and strong selection are critical for the evolutionary potential of populations of long-lived species,

where long generation times limit rapid evolutionary change. We provide direct evidence of fluctuating selection during ontogeny in *P. ponderosa*, a long-lived species. Fast growth was selected during juvenile stages, but slow growth was selected under intense herbivory pressure by a MPB outbreak at mature stages (34 to 38 y old; i.e., growth–survival trade-off). Such opposing selection led to no net evolutionary response of growth rates, thus contributing to the maintenance of genetic diversity of this trait. Our data suggest that the growth–survival trade-off during the outbreak was not solely driven by MPB's preference for larger, resource-rich phenotypes at the epidemic stage (even when well-defended), but it likely reflected an intrinsic host-based life-history trade-off. In contrast to our prediction, strong genetic differentiation in survival to the MPB was delayed and occurred the second year of highest mortality, pointing to complex resistance responses under genetic control. The evolutionary consequences of selective events driven by biotic interactions are relevant under climate change, given the expected increase of insect pests and pathogens (16).

Three lines of evidence support that fluctuating selection was caused by novel environmental conditions imposed by intense herbivory pressure as a result of the MPB outbreak. First, family survival rankings changed significantly from before to the end of the outbreak (Fig. 1). Second, genetic correlations between survival rates at successive sampling periods indicate that at the onset of the outbreak the MPB started to target families that were more likely to survive in the absence of intense herbivory (Fig. S6). Third, when MPB was excluded by insecticide spraying, fast growth was not selected against (Fig. 3). These results support classic accounts of selection in natural populations as a result of biotic interactions, often modulated by abiotic factors (2, 37).

Growth rates have fitness consequences and selection is expected (23). Indeed, we found significant genetic responses to selection on growth rates, but importantly, these responses changed in direction and strength over time (Fig. 2A). Fast growth was positively selected before the outbreak, but negatively selected during the outbreak, clearly showing that intense herbivory shifted selection patterns. Fast growth in trees under competitive environments is critical for light acquisition and resource capture, and slow growth is selected against and underrepresented at mature stages (27). Consistently, selection for fast growth was strongest during early seedling establishment




Fig. 3. Mean diameter of surviving and dead trees over time and of a subset of surviving fast growing (elite) trees treated by insecticide. Preoutbreak surviving trees grew faster than trees that died. The reverse was true post-outbreak. When beetles were excluded in elite trees, fast growth was not selected against and 93% survived. Thus, selection against larger trees in the overall population was beetle-driven. Bars are SEs (often smaller than symbols). Mean diameters postoutbreak are from age 27 y after accounting for mortality. Dashed lines between ages 27 and 34 y illustrate that different sets of trees survive and die pre- and postoutbreak.




Fig. 4. Genetic correlations between intrinsic growth potential (DBH, age 27 y) and (A) cumulative survival preoutbreak (age 27 y), (B) at the end of the outbreak (age 38 y; only the live-tree pool at age 27 y was considered), and survival from ages (C) 2–4 (preoutbreak) and (D) 35–36 y (during the beetle outbreak). Intervals from ages 2–4 y and 35–36 y were chosen based on most significant genetic correlations (Fig. S7). Data points represent best linear unbiased predictors for 204 open-pollinated pine families. Regression lines, bivariate genetic correlation coefficients (r_G), and 95% credibility intervals (CI) shown.

when density-dependent mortality in trees typically occurs (38) and when the proportion of seeds that attain maturity is usually very small (6). The MPB outbreak, however, caused significant selection differentials in the opposite direction (positive selection for slow-growing phenotypes), which triggered a negative genetic response after the outbreak (Fig. 24). Our results are consistent with studies showing that fast early growth within tree species correlates with decreased longevity (29) and increased herbivory at maturity (30), and provide strong empirical evidence of the conflicting effects of growth rates on fitness during ontogeny.

Importantly, growth potential did not experience a net evolutionary response at the end of the sampling period (Fig. 2B), and even though the negative genetic response caused by the outbreak was noticeably smaller than the positive response at juvenile stages, it was strong enough to counteract the expected positive selection for fast growth in trees. Thus, positive and negative selection responses (1) reduced the cumulative effects of selection, clearly illustrating the importance of assessing selection over time (4).

Directional selection is expected to reduce genetic variability. The decrease in heritability for growth over time (Fig. 5A) suggests this occurred to some degree. However, the reduction was stronger at earlier stages when genetic response to selection for fast growth was strongest, but dampened thereafter. This and the negligible net response to selection clearly indicate that in natural populations subjected to fluctuating and often unpredictable selection pressures, consequences of growth rates are highly context-dependent, thus leading to fluctuating selection and the maintenance of genetic variation (4, 22, 39).

Trees show high intrapopulation genetic diversity relative to other plant species (40), which contributes to their evolutionary potential (18). However, increased potential for microevolution in trees contrasts with slow macroevolutionary rates (6). The current most-favored explanation for this paradox is that selection is not a major driver of genetic diversity (41). In nonsessile organisms, selection has been shown to contribute very little to phenotypic changes over time and changing climate (42). Our results, in contrast, suggest that selection in trees (long-lived sessile organisms) may be more relevant than previously thought and that the accumulation of context-dependent fluctuating, episodic, and opposing selection forces over time contribute to the maintenance of genetic diversity within populations (43). Such diversity is critical to buffer changes in selective pressures as a result of environmental variation over the extended lifespan of trees (18), and likely contributes to their evolutionary success (6).

As the MPB population increases to the epidemic stage, host defenses are more easily overwhelmed and allow beetles to behaviorally switch from weakened, stressed trees toward larger, resource-rich hosts, even if such hosts are better defended (35). This behavioral switch is consistent with the plant-vigor hypothesis, posing that more vigorous, highly nutritious hosts are preferred by insects (44). Consistently, we found that the MPB clearly preferred larger trees (Fig. 3), despite the fact that larger trees at our site have higher resin duct defenses and slightly higher resin flow (45). Negative genetic correlations are often interpreted as evidence of intrinsic life-history trade-offs (46). However, because faster growing families have larger hosts on average, and at the epidemic stage beetles switch preference for larger hosts regardless of defenses, the growth-survival trade-off we found could be entirely driven by the MPB behavioral switch rather than reflect a host-based life-history trade-off (31, 32). However, two lines of evidence suggest that the growth-survival trade-off we found reflects in part a host-based life-history trade-off. First, despite the significant environmental effect caused by the MPB behavior switch (Fig. S7), a significant negative genetic correlation between growth and survival was superimposed (and somewhat stronger) at age 36 y during the peak of the outbreak. Therefore, MPB not only targeted larger phenotypes, but trees from faster growing families. Second, such effect was not driven by a depletion of large hosts in slower-growing families during the first year of high mortality (age 35 y) (Fig. S8). Note also that survival to the MBP outbreak at our site was quite high (64%), likely leaving suitable hosts in all families where variability in growth rates (size) is high. Furthermore, our results are conservative because original selected mothers were vigorous trees, thus likely limiting the range of growth rates. To the extent that mortality during the outbreak was exclusively caused by the beetle, the host-based growth-survival trade-off most likely reflects a growth-resistance trade-off. Unfortunately, however, we lack direct data on defenses during the outbreak (which cannot be obtained in this retrospective study; see below) and we cannot unambiguously demonstrate a growth-resistance trade-off.

Consistent with previous results in *Pinus* species (33, 36), we found high genetic differentiation in survival (resistance) during the outbreak (Fig. 5C). However, genetic effects varied strongly and unexpectedly over time: they were insignificant early and late in the outbreak (Fig. 5D), but very strong at the peak of the outbreak (at age 36 y, the second consecutive year of highest mortality; approximately 15% for both years). Such a delayed




Fig. 5. Individual heritability of: (A) growth rates measured as diameter and height over entire period, (B) cumulative survival over entire period, and (C) cumulative survival and (D) survival at each age postoutbreak (live-tree pool only). Bars are 95% credibility intervals. Heritabilities are all significant except when “ns” is noted (“0” value not included in the credibility interval). Heritability for growth traits after the beetle outbreak (A) were estimated based on age 27 y and accounting for mortality thereafter.

of animal models, above) between intrinsic growth potential (DBH at age 27 y) and survival at each sampling period, and also at ages 27 y (overall correlation before the outbreak) and at the end of the outbreak, excluding trees killed before age 27 y (overall correlation during the outbreak). Because mortality after age 27 y was exclusively because of MPB, a negative growth–survival genetic correlation indicates a growth–survival trade-off.

Because individuals from a given family were planted in four-tree family-row plots, insecticide effects on neighbors could have confounded genetic effects. We tested whether the insecticide application on 106 trees in 2010 affected our results by including this as a fixed factor in the quantitative genetic model (see discussion of animal models, above). We considered five groups of trees potentially affected by the insecticide application: targeted sprayed trees and trees one, two, three, and more than three trees away from sprayed trees, respectively. Post hoc comparisons indicated that the

insecticide affected survival of the target trees only (*SI Text*). We then deleted the trees affected by the insecticide from the analysis and re-estimated genetic variances and heritabilities. The number of trees sprayed was too small to allow meaningful estimates of selection (only 106 trees were sprayed).

ACKNOWLEDGMENTS. We thank Alexandra Ginter, Gilia Patterson, Katie Arledge, and Jordi Martínez-Vilalta for field support; Marc Rust for Inland Empire Tree Improvement Cooperative data through 2001; and Rafael Zas, David Coltman, Thomas Martin, Lila Fishman, and two anonymous reviewers for thoughtful comments. The Barrié Foundation provided postdoctoral support to R.d.I.M. Research funds were provided by McIntire-Stennis Cooperative Forestry Research Grant MONZ-1206 (College of Forestry and Conservation, University of Montana).

- Schluter D, Price TD, Rowe L (1991) Conflicting selection pressures and life history trade-offs. *Philos T R Soc B* 246:11–17.
- Grant PR, Grant BR (2002) Unpredictable evolution in a 30-year study of Darwin's finches. *Science* 296:707–711.
- Hoffmann AA, Sgrò CM (2011) Climate change and evolutionary adaptation. *Nature* 470:479–485.
- Siepielski AM, DiBattista JD, Carlson SM (2009) It's about time: The temporal dynamics of phenotypic selection in the wild. *Ecol Lett* 12:1261–1276.
- Kingsolver JG, Diamond SE (2011) Phenotypic selection in natural populations: what limits directional selection? *Am Nat* 177:346–357.
- Petit RJ, Hampe A (2006) Some evolutionary consequences of being a tree. *Annu Rev Ecol Syst* 37:187–214.
- Alía R, Chambel R, Notivol E, Climent J, González-Martínez SC (2014) Environment-dependent microevolution in a Mediterranean pine (*Pinus pinaster* Aiton). *BMC Evol Biol* 14:200.
- Franklin J, Serra-Díaz JM, Syphard AD, Regan HM (2016) Global change and terrestrial plant community dynamics. *Proc Natl Acad Sci USA* 113:3725–3734.
- Hansen MC, et al. (2013) High-resolution global maps of 21st-century forest cover change. *Science* 342:850–853.
- Allen CD, Breshears DD, McDowell NG (2015) On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* 6:art129.
- Seidl R, Spies TA, Peterson DL, Stephens SL, Hicke JA (2016) Searching for resilience: Addressing the impacts of changing disturbance regimes on forest ecosystem services. *J Appl Ecol* 53:120–129.
- Speer JH, Swetnam TW, Wickman BE, Youngblood A (2001) Changes in pandora moth outbreak dynamics during the past 622 years. *Ecology* 82:679–697.
- Gutschick VP, BassiriRad H (2003) Extreme events as shaping physiology, ecology, and evolution of plants: Toward a unified definition and evaluation of their consequences. *New Phytol* 160:21–42.
- Sousa WP (1984) The role of disturbance in natural communities. *Annu Rev Ecol Syst* 15:353–391.
- Boege K, Marquis RJ (2005) Facing herbivory as you grow up: The ontogeny of resistance in plants. *Trends Ecol Evol* 20:441–448.
- Anderegg WRL, et al. (2015) Tree mortality from drought, insects, and their interactions in a changing climate. *New Phytol* 208:674–683.
- Aitken SN, Yeaman S, Holliday JA, Wang T, Curtis-McLane S (2008) Adaptation, migration or extirpation: Climate change outcomes for tree populations. *Evol Appl* 1: 95–111.
- Alberto FJ, et al. (2013) Potential for evolutionary responses to climate change—Evidence from tree populations. *Glob Change Biol* 19:1645–1661.
- Raffa KF, et al. (2008) Cross-scale drivers of natural disturbances prone to anthropogenic amplification: The dynamics of bark beetle eruptions. *Bioscience* 58:501–517.
- Stowe KA, Marquis RJ, Hochwender CG, Simms EL (2000) The evolutionary ecology of tolerance to consumer damage. *Annu Rev Ecol Syst* 31:565–595.
- Meddens AJH, Hicke JA, Ferguson CA (2012) Spatiotemporal patterns of observed bark beetle-caused tree mortality in British Columbia and the western United States. *Ecol Appl* 22:1876–1891.
- Monro K, Marshall DJ (2014) Faster is not always better: Selection on growth rate fluctuates across life history and environments. *Am Nat* 183:798–809.
- Arendt JD (1997) Adaptive intrinsic growth rates: An integration across taxa. *Q Rev Biol* 72:149–177.
- Boege K, Dirzo R, Siemens D, Brown P (2007) Ontogenetic switches from plant resistance to tolerance: Minimizing costs with age? *Ecol Lett* 10:177–187.
- Mangel M, Stamps J (2001) Trade-offs between growth and mortality and the maintenance of individual variation in growth. *Evol Ecol Res* 3:583–593.
- Lanner RM (2002) Why do trees live so long? *Ageing Res Rev* 1:653–671.
- Landis RM, Peart DR (2005) Early performance predicts canopy attainment across life histories in subalpine forest trees. *Ecology* 86:63–72.
- Lee W-S, Monaghan P, Metcalfe NB (2013) Experimental demonstration of the growth rate–lifespan trade-off. *Philos T R Soc B* 280:20122370.
- Bigler C, Veblen TT (2009) Increased early growth rates decrease longevities of conifers in subalpine forests. *Oikos* 118:1130–1138.
- Ruel J, Whitham TG (2002) Fast-growing juvenile pinyons suffer greater herbivory when mature. *Ecology* 83:2691–2699.
- Coley PD, Bryant JP, Chapin FS, 3rd (1985) Resource availability and plant anti-herbivore defense. *Science* 230:895–899.
- Herm DA, Mattson WJ (1992) The dilemma of plants: To grow or defend. *Q Rev Biol* 67:283–335.
- Strom BL, Goyer RA, Ingram LL, Boyd GDL, Lott LH (2002) Oleoresin characteristics of progeny of loblolly pines that escaped attack by the southern pine beetle. *For Ecol Manage* 158:169–178.
- Doak DF (1992) Lifetime impacts of herbivory for a perennial plant. *Ecology* 73: 2086–2099.
- Boone CK, Aukema BH, Bohlmann J, Carroll AL, Raffa KF (2011) Efficacy of tree defense physiology varies with bark beetle population density: A basis for positive feedback in eruptive species. *Can J Res* 41:1174–1188.
- Yanchuk AD, Murphy JC, Wallin KF (2008) Evaluation of genetic variation of attack and resistance in lodgepole pine in the early stages of a mountain pine beetle outbreak. *Tree Genet Genomes* 4:171–180.
- Bell G (2010) Fluctuating selection: The perpetual renewal of adaptation in variable environments. *Philos Trans R Soc Lond B Biol Sci* 365:87–97.
- Dodd ME, Silvertown J (2000) Size-specific fecundity and the influence of lifetime size variation upon effective population size in *Abies balsamea*. *Heredity (Edinb)* 85: 604–609.
- Dmitriev CM (2011) The evolution of growth trajectories: What limits growth rate? *Biol Rev Camb Philos Soc* 86:97–116.
- Hamrick JL, Godt M (1996) Effects of life history traits on genetic diversity in plant species. *Philos T R Soc B* 351:1291–1298.
- Scotti I, González-Martínez SC, Budde KB, Lalagüe H (2016) Fifty years of genetic studies: What to make of the large amounts of variation found within populations? *Ann Sci* 73:69–75.
- Ozgul A, et al. (2009) The dynamics of phenotypic change and the shrinking sheep of St. Kilda. *Science* 325:464–467.
- Takahata N, Ishii K, Matsuda H (1975) Effect of temporal fluctuation of selection coefficient on gene frequency in a population. *Proc Natl Acad Sci USA* 72:4541–4545.
- Price PW (1991) The plant vigor hypothesis and herbivore attack. *Oikos* 62:244–251.
- Hood S, Sala A (2015) Ponderosa pine resin defenses and growth: Metrics matter. *Tree Physiol* 35:1223–1235.
- Agrawal AA, Conner JK, Rasmann S (2010) Tradeoffs and negative correlations in evolutionary ecology. *Evolution Since Darwin: The First 150 Years*, eds Bell M, Eanes W, Futuyma D, Levinton J (Sinauer Associates, Sunderland, MA), pp 243–268.
- Haukioja E, Suomela J, Neuvonen S (1985) Long-term inducible resistance in birch foliage: Triggering cues and efficacy on a defoliator. *Oecologia* 65:363–369.
- Franceschi VR, Krokene P, Christiansen E, Krekling T (2005) Anatomical and chemical defenses of conifer bark against bark beetles and other pests. *New Phytol* 167:353–375.
- Moreira X, et al. (2014) Trade-offs between constitutive and induced defenses drive geographical and climatic clines in pine chemical defences. *Ecol Lett* 17:537–546.
- Pinnell S (2016) Resin duct defenses in ponderosa pine during a mountain pine beetle outbreak: Genetic effects, mortality, and relationships with growth. Masters thesis (University of Montana, Missoula, MT).
- Kolb TE, et al. (2016) Observed and anticipated impacts of drought on forest insects and diseases in the United States. *For Ecol Manage* 380:321–334.
- White TL, Adams WT, Neale DB (2007) *Forest Genetics* (Cabi Publishing, Oxfordshire, UK).
- Snieszko RA (2006) Resistance breeding against nonnative pathogens in forest trees—Current successes in North America. *Can J Plant Pathol* 28:S270–S279.
- Franklin O, Palmroth S, Näsholm T (2014) How eco-evolutionary principles can guide tree breeding and tree biotechnology for enhanced productivity. *Tree Physiol* 34: 1149–1166.
- King KC, Lively CM (2012) Does genetic diversity limit disease spread in natural host populations? *Heredity (Edinb)* 109:199–203.
- Paquette A, Messier C (2010) The role of plantations in managing the world's forests in the Anthropocene. *Front Ecol Environ* 8:27–34.
- Preisler HK (1993) Modelling spatial patterns of trees attacked by bark-beetles. *Appl Stat* 42:501–514.
- Zas R (2006) Iterative kriging for removing spatial autocorrelation in analysis of forest genetic traits. *Tree Genet Genomes* 2:177–185.
- Zas R, Solla A, Sampedro L (2007) Variography and kriging allow screening *Pinus pinaster* resistant to *Armillaria ostoyae* in field conditions. *Forestry* 80:201–209.
- Hadfield JD (2010) MCMC methods for multi-response generalized linear mixed models: The MCMCglmm R package. *J Stat Softw* 33:1–22.
- Price GR (1970) Selection and covariance. *Nature* 227:520–521.
- Kruuk LEB (2004) Estimating genetic parameters in natural populations using the “animal model”. *Philos Trans R Soc Lond B Biol Sci* 359:873–890.