



United States Department of Agriculture



**NORTHWEST
FOREST PLAN**

Synthesis of Science to Inform Land Management Within the Northwest Forest Plan Area

Volume 2



**Forest
Service**

Pacific Northwest
Research Station

General Technical Report
PNW-GTR-966 Vol. 2

June
2018

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at http://www.ascr.usda.gov/complaint_filing_cust.html and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

USDA is an equal opportunity provider, employer, and lender.

Technical Coordinators

Thomas A. Spies is a senior scientist, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis, OR 97331; **Peter A. Stine** is a research program manager and biogeographer (retired), U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, and is a research associate, John Muir Institute for the Environment, University of California–Davis, 1 Shields Avenue, Davis, CA 95616; **Rebecca Gravenmier** is a science coordinator, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 1220 SW 3rd Avenue, Portland, OR 97204; **Jonathan W. Long** is an ecologist, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 1731 Research Park Drive, Davis, CA 95618; and **Matthew J. Reilly** is a postdoctoral researcher, Humboldt State University, Department of Biological Sciences, 1 Harpst Street, Arcata, CA 95521.

Cover photos: Upper left: coho spawning on the Salmon River; photo by Bureau of Land Management. Upper right: a typical stream in the Cispus River watershed, Washington; photo by Alanna Wong. Lower right: an adult captive red tree vole mechanically removing one of two unpalatable resin ducts in a Douglas-fir needle before eating the rest of the needle; photo by Michael Durham. Lower middle: orange coral fungus, Olympic National Forest; photo by USDA Forest Service. Lower left: centipede, one component of the wide array of soil invertebrates found in late-successional and old-growth forests of the Northwest Forest Plan region; photo by Bruce Marcot.

Synthesis of Science to Inform Land Management Within the Northwest Forest Plan Area

Volume 2

Thomas A. Spies, Peter A. Stine, Rebecca Gravenmier,
Jonathan W. Long, and Matthew J. Reilly, Technical Coordinators

U.S. Department of Agriculture
Forest Service
Pacific Northwest Research Station
Portland, Oregon
General Technical Report PNW-GTR-966 Vol. 2
June 2018

Abstract

Spies, T.A.; Stine, P.A.; Gravenmier, R.; Long, J.W.; Reilly, M.J., tech. coords. 2018.

Synthesis of science to inform land management within the Northwest Forest Plan area. Gen. Tech. Rep. PNW-GTR-966. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 1020 p. 3 vol.

The 1994 Northwest Forest Plan (NWFP) was developed to resolve debates over old-growth forests, endangered species, and timber production on federal forests in the range of the northern spotted owl. This three-volume science synthesis, which consists of 12 chapters that address various ecological and social concerns, is intended to inform forest plan revision and forest management within the NWFP area. Land managers with the U.S. Forest Service provided questions that helped guide preparation of the synthesis, which builds on the 10-, 15-, and 20-year NWFP monitoring reports and synthesizes the vast body of relevant scientific literature that has accumulated in the 24 years since the NWFP was initiated. It identifies scientific findings, lessons learned, and uncertainties and also evaluates competing science and provides considerations for management.

This synthesis finds that the NWFP has protected dense old-growth forests and maintained habitat for northern spotted owls, marbled murrelets, aquatic organisms, and other species despite losses from wildfire and low levels of timber harvest on federal lands. Even with reductions in the loss of older forests, northern spotted owl populations continue to decline. Moreover, a number of other goals have not been met, including producing a sustainable supply of timber, decommissioning roads, biodiversity monitoring, significant levels of restoration of riparian and dry forests, and adaptation and learning through adaptive management.

New conservation concerns have arisen, including a major threat to spotted owl populations from expanding populations of the nonnative barred owl, effects of fire suppression on forest succession, fire behavior in dry forests, and lack of development of diverse early-seral vegetation as a result of fire suppression in drier parts of moist forests. Climate change and invasive species have emerged as threats to native biodiversity, and expansion of the wildland-urban interface has limited the ability of managers to restore fire to fire-dependent ecosystems.

The policy, social, and ecological contexts for the NWFP have changed since it was implemented. The contribution of federal lands continues to be essential to the conservation and recovery of fish listed under the Endangered Species Act and northern spotted owl and marbled murrelet populations. Conservation on federal lands alone, however, is likely insufficient to reach the goals of the NWFP or the newer goals of the 2012 planning rule, which emphasizes managing for ecosystem goals (e.g. ecological resilience) and a few species of concern, rather than the population viability of hundreds of individual species.

The social and economic basis of many traditionally forest-dependent communities have changed in 24 years, and many are now focused on amenity values. The capacities of human communities and federal agencies, collaboration among stakeholders, the interdependence of restoration and the timber economy, and the role of amenity- or recreation-based communities and ecosystem services are important considerations in managing for ecological resilience, biodiversity conservation, and social and economic sustainability.

A growing body of scientific evidence supports the importance of active management or restoration inside and outside reserves to promote biodiversity and ecological resilience. Active management to promote heterogeneity of vegetation conditions is important to sustaining tribal ecocultural resources. Declines in agency capacity, lack of markets for small-diameter wood, lack of wood processing infrastructure in some areas, and lack of social agreement have limited the amount of active management for restoration on federal lands. All management choices involve social and ecological tradeoffs related to the goals of the NWFP. Collaboration, risk management, adaptive management, and monitoring are considered the best ways to deal with complex social and ecological systems with futures that are difficult to predict and affect through policy and land management actions.

Keywords: Northwest Forest Plan, science, management, restoration, northern spotted owl, marbled murrelet, climate change, socioeconomic, environmental justice.

Preface

In 2015, regional foresters in the Pacific Northwest and Pacific Southwest Regions of the USDA Forest Service requested that the Pacific Northwest and Pacific Southwest Research Stations prepare a science synthesis to inform revision of existing forest plans under the 2012 planning rule in the area of the Northwest Forest Plan (NWFP, or Plan). Managers provided an initial list of hundreds of questions to the science team, which reduced to them to 73 questions deemed most feasible for addressing through a study of current scientific literature. The stations assembled a team of 50 scientists with expertise in biological, ecological, and socioeconomic disciplines. At the suggestion of stakeholders, a literature reference database was placed online so the public could submit additional scientific literature for consideration. By spring 2016, writing was underway on 12 chapters that covered ecological and social sciences.

The draft synthesis, which was ready for peer and public review by fall 2016, went through a special review process because it was classified as “highly influential science” in accordance with the Office of Management and Budget’s 2004 “Final Information Quality Bulletin for Peer Review.” The synthesis was classified as such because it fit the category of a scientific assessment that is novel, controversial, or precedent-setting, or has significant interagency interest. Per the bulletin, the two research stations commissioned an independent entity, the Ecological Society of America (ESA), to manage the peer-review process, including the selection of peer reviewers.

The bulletin also stipulates that such an assessment be made available to the public through a public meeting to enable the public to bring scientific issues to the attention of peer reviewers. Accordingly, a public forum was held in Portland, Oregon, in December 2016. For those who could not travel to Portland, the forum was accessible via live Web stream, and multiple national forests within the NWFP area hosted remote viewing. Written comments on the draft synthesis were collected for 2 months. This generated 130 public comments, totaling 890 pages, which were given to the peer reviewers for consideration in their review, as they deemed appropriate. The OMB guidelines further direct that the peer-review process be transparent by making available to the public the ESA’s written guidance to the reviewers, the peer reviewer’s names, the peer review reports, and the responses of the authors to the peer reviewer comments—all of which are available at <https://www.fs.fed.us/pnw/research/science-synthesis/index.shtml>.

The peer reviewer comments, which were received in spring 2017 and informed by public input, resulted in substantive revisions to chapters of the synthesis. The result is this three-volume general technical report (an executive summary of the synthesis is available as a separate report). This document is intended to support upcoming management planning on all public lands in the Plan area, but is expected to serve primarily lands managed by the U.S. Forest Service. We hope it will be a valuable reference for managers and others who seek to understand the scientific basis and possible tradeoffs associated with forest plan revision and management decisions. The synthesis also provides an extensive list of published sources where readers can find further information.

We understand that the term “synthesis” can have many different meanings. For our purposes, it represents a compilation and interpretation of relevant scientific findings that pertain to key issues related to the NWFP that were identified by managers and by the authors of the document. Such a compilation not only summarizes science by topic areas but also interprets that science in light of management goals, characterizes competing science, and makes connections across scientific areas, addressing multilayered and interacting ecological and socioeconomic issues. In a few cases, simple analyses of existing data were conducted and methods were provided to reviewers.

The synthesis builds upon the 10-, 15-, and 20-year NWFP monitoring reports, and authors considered well over 4,000 peer-reviewed publications based on their knowledge as well as publications submitted by the public and others suggested by peer reviewers. For some of the questions posed by land managers, there was ample scientific research from the Plan area. For many of the questions, however, little research existed that was specific to the area. In such cases, studies from other regions or current scientific theory were used to address the questions to the extent possible. In many cases, major scientific uncertainties were found; these are highlighted by the authors.

The synthesis chapters characterize the state of the science but they do not develop management alternatives, analyze management tradeoffs, or offer recommendations as to what managers should do. The synthesis does identify ideas, facts, and relationships that managers may want to consider as they develop plans and make management decisions about particular issues. The final chapter attempts to integrate significant cross-cutting issues, e.g., ecological and socioeconomic interdependencies, compatibility of different management goals, and tradeoffs associated with different restoration actions. All the chapters identify where more research is needed to fill critical information gaps.

We would like to acknowledge the peer reviewers who considered hundreds of public comments as part of the process of reviewing our lengthy draft manuscripts. We also thank the many contributors to the development of the synthesis in draft and final form, including those who provided editing, layout, database, and other support services.

Contents

Volume 1

- 1 **Chapter 1: Introduction**
Peter A. Stine and Thomas A. Spies
- 29 **Chapter 2: Climate, Disturbance, and Vulnerability to Vegetation Change in the Northwest Forest Plan Area**
Matthew J. Reilly, Thomas A. Spies, Jeremy Littell, Ramona Butz, and John B. Kim
- 95 **Chapter 3: Old Growth, Disturbance, Forest Succession, and Management in the Area of the Northwest Forest Plan**
Thomas A. Spies, Paul F. Hessburg, Carl N. Skinner, Klaus J. Puettmann, Matthew J. Reilly, Raymond J. Davis, Jane A. Kertis, Jonathan W. Long, and David C. Shaw
- 245 **Chapter 4: Northern Spotted Owl Habitat and Populations: Status and Threats**
Damon B. Lesmeister, Raymond J. Davis, Peter H. Singleton, and J. David Wiens
- 301 **Chapter 5: Marbled Murrelet**
Martin G. Raphael, Gary A. Falxa, and Alan E. Burger

Volume 2

- 371 **Chapter 6: Other Species and Biodiversity of Older Forests**
Bruce G. Marcot, Karen L. Pope, Keith Slauson, Hartwell H. Welsh, Clara A. Wheeler, Matthew J. Reilly, and William J. Zielinski
- 371 **Introduction**
- 372 Setting and Background
- 373 New Learning and Recent Issues
- 373 **Guiding Questions**
- 374 **Agency Programs on Other Species and Biodiversity of Older Forests**
- 374 Survey and Manage Program
- 378 Interagency Special Status and Sensitive Species Program
- 378 Implications of Forest Service and BLM Planning Directions
- 379 **Key Findings**
- 379 New Information on Other Species and Biodiversity of Older Forests
- 408 Recent Research and Development on New Tools and Data
- 409 Recent and Ongoing Issues Adding Challenges and Opportunities
- 416 **Uncertainties and Research Needs**
- 416 Indicators and Surrogates
- 417 Climate Change
- 418 **Conclusions and Management Considerations**
- 418 Summary of Key Management Considerations
- 422 Issues of Species Conservation
- 425 Main Findings
- 426 **Acknowledgments**
- 427 **Metric Equivalents**
- 427 **Literature Cited**

461	Chapter 7: The Aquatic Conservation Strategy of the Northwest Forest Plan—A Review of the Relevant Science After 23 Years
	<i>Gordon H. Reeves, Deanna H. Olson, Steven M. Wondzell, Peter A. Bisson, Sean Gordon, Stephanie A. Miller, Jonathan W. Long, and Michael J. Furniss</i>
461	Introduction
463	Guiding Questions
463	Key Findings
463	Status of Species and Population Units Listed Under the Endangered Species Act on Federal Lands in the Northwest Forest Plan Area
468	Monitoring—Aquatic and Riparian Effectiveness Monitoring Program
483	Components of the Aquatic Conservation Strategy
483	Riparian Reserves
498	Key Watersheds
499	Watershed Analysis
499	New Perspectives on Conservation of Riparian and Aquatic Ecosystems
504	Non-Fish-Bearing Streams
506	Continuing and Emerging Topics of Concern
506	Water
506	Roads
510	Climate Change
531	Research Needs, Uncertainties, Information Gaps, and Limitations
533	Conclusions and Management Considerations
533	Guiding Questions
534	Science Foundation for the ACS (Question 1)
535	Monitoring of the ACS (Question 2)
536	Aquatic and Riparian Ecosystems (Questions 3–6)
537	Water Contributions From Federal Lands (Question 7)
537	Climate Change (Question 8)
537	Literature Cited
582	Appendix 1: Aquatic-Riparian Invasive Species of the Northwest Forest Plan Area
	<i>Deanna H. Olson and Rebecca L. Flitcroft</i>
582	Priority Aquatic Invasive Species
586	Taxonomic Summaries
586	Pathogens and Parasites
586	Aquatic Plants
587	Riparian-Terrestrial Plants
587	Aquatic Invertebrates
587	Aquatic Vertebrates
588	Riparian-Terrestrial Vertebrates
588	Research and Development, Monitoring, and Management
588	Research and Development

590	Monitoring and Management
593	Future Considerations
594	Other Nonnative Species on the Radar in the Northwest
594	Climate Change Projections
595	Research and Monitoring Priorities
596	Appendix 2: Influence of Climate Change on Life Stages of Pacific Salmon
	<i>Peter A. Bisson, Gordon H. Reeves, Nate Mantua, and Steven M. Wondzell</i>
596	Adults
598	Eggs and Alevins
600	Juveniles
603	Smolts
608	Glossary
624	Acknowledgments

Volume 3

625	Chapter 8: Socioeconomic Well-Being and Forest Management in Northwest Forest Plan-Area Communities
	<i>Susan Charnley, Jeffrey D. Kline, Eric M. White, Jesse Abrams, Rebecca J. McLain, Cassandra Moseley, and Heidi Huber-Stearns</i>
717	Chapter 9: Understanding Our Changing Public Values, Resource Uses, and Engagement Processes and Practices
	<i>Lee K. Cervený, Emily Jane Davis, Rebecca McLain, Clare M. Ryan, Debra R. Whitall, and Eric M. White</i>
809	Chapter 10—Environmental Justice, Low-Income and Minority Populations, and Forest Management in the Northwest Forest Plan Area
	<i>Susan Charnley, Delilah Jaworski, Heidi Huber-Stearns, Eric M. White, Elisabeth Grinspoon, Rebecca J. McLain, and Lee Cervený</i>
851	Chapter 11: Tribal Ecocultural Resources and Engagement
	<i>Jonathan Long, Frank K. Lake, Kathy Lynn, and Carson Viles</i>
919	Chapter 12: Integrating Ecological and Social Science to Inform Land Management in the Area of the Northwest Forest Plan
	<i>Thomas A. Spies, Jonathan W. Long, Peter Stine, Susan Charnley, Lee Cervený, Bruce G. Marcot, Gordon Reeves, Paul F. Hessburg, Damon Lesmeister, Matthew J. Reilly, Martin G. Raphael, and Raymond J. Davis</i>



Table Rock Wilderness, Oregon.
Photo by USDI Bureau of Land Management, Oregon-Washington State Office.

Chapter 6: Other Species and Biodiversity of Older Forests

Bruce G. Marcot, Karen L. Pope, Keith Slauson, Hartwell H. Welsh, Clara A. Wheeler, Matthew J. Reilly, and William J. Zielinski¹

Introduction

This chapter focuses mostly on terrestrial conditions of species and biodiversity associated with late-successional and old-growth forests in the area of the Northwest Forest Plan (NWFP). We do not address the northern spotted owl (*Strix occidentalis caurina*) or marbled murrelet (*Brachyramphus marmoratus*)—those species and their habitat needs are covered in chapters 4 and 5, respectively. Also, the NWFP's Aquatic and Riparian Conservation Strategy and associated fish species are addressed in chapter 7, and early-successional vegetation and other conditions are covered more in chapters 3 and 12.

We begin by summarizing a set of questions provided by management. We then review the state of knowledge of species, biodiversity, and ecosystem conditions gained from studies conducted since the 10-year synthesis of monitoring and research results (Haynes et al. 2006). We review agency programs on other species and biodiversity of older forests of the Pacific Northwest, including implementation of the NWFP Survey and Manage standards and guidelines, the Interagency Special Status and Sensitive Species Program (ISSSP), and other biodiversity consortia. We then review new findings on selected individual species and groups of species including fungi, lichens, bryophytes, and plants, as well as invertebrates. We also summarize

findings on amphibians, reptiles, and birds, and on selected carnivore species including fisher (*Pekania pennanti*), marten (*Martes americana*), and wolverine (*Gulo gulo*), and on red tree voles (*Arborimus longicaudus*) and bats. We close the section with a brief review of the value of early-seral vegetation environments. We next review recent advances in development of new tools and datasets for species and biodiversity conservation in late-successional and old-growth forests, and then review recent and ongoing challenges and opportunities for ameliorating threats and addressing dynamic system changes. We end with a set of management considerations drawn from research conducted since the 10-year science synthesis and suggest areas of further study.

The general themes reviewed in this chapter were guided by a set of questions provided by the U.S. Forest Service Pacific Northwest Region (Region 6) and Pacific Southwest Region (Region 5). The scientific publications we review were selected based on the specific subjects listed above, as pertinent to science findings on other species and biodiversity of late-successional and old-growth forest ecosystems in the area of the NWFP in the Pacific Northwest, United States. We include selected references on studies outside the NWFP and Pacific Northwest and references dating prior to the previous NWFP science synthesis, when such studies are nonetheless pertinent to understanding biological and ecological topics within the NWFP and Pacific Northwest. We also address selected topics such as early-successional forest ecosystems and effects of wildfire, fire suppression, and climate change, as guided by the availability of recent literature on NWFP species and biodiversity; these topics, raised by managers, are also covered more fully in other chapters of this science synthesis. The final chapter of this synthesis discusses the conceptual and practical implications of new science findings, remaining scientific uncertainties and research needs, and overall conclusions.

¹ **Bruce G. Marcot** is a research wildlife biologist, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 620 SW Main St., Suite 400, Portland, OR 97205; **Karen L. Pope** is a research wildlife biologist, **Keith Slauson** is a research fellow, **Hartwell H. Welsh** is a research wildlife biologist emeritus, **Clara A. Wheeler** is an ecologist, and **William J. Zielinski** is a research ecologist, U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, 1700 Bayview Drive, Arcata, CA 95521; **Matthew J. Reilly** is a postdoctoral researcher, Humboldt State University, Department of Biological Sciences, 1 Harpst Street, Arcata, CA 95521.

Setting and Background

Originally, the NWFP was developed as an ecosystem management plan to provide for the full suite of biodiversity at all taxonomic and functional levels, particularly in late-successional and old-growth environments, under an adaptive learning and management approach. The first decade of the NWFP, however, focused on the status of species; no biodiversity monitoring program per se was instituted under the NWFP (Marcot and Molina 2006). Since then, a broad assumption has been made that older-forest biodiversity in its full capacity would be provided by two complementary approaches of managing for “coarse filter” elements such as the dispersion and distribution of late-successional forest reserves and aquatic and riparian corridors, along with managing for “fine filter” elements of habitat needs of selected, individual late-successional and old-growth-associated species. The combined coarse- and fine-filter approach is intended to provide the same level of protection as would management and monitoring directed at specific biodiversity elements such as ecosystem processes of nutrient cycling, species’ ecological functions, and population genetics and viability (e.g., Noss 1990). A current challenge is to test this assumption within the changing tapestry of ecological processes and disturbance-influenced ecosystems of the Northwest.

The 2012 planning rule for guiding land and resource management plans on national forests differs from the 1982 rule that was in place when the NWFP was first instituted and that guided the NWFP. The 2012 planning rule puts more weight on coarse-filter approaches and on ecological integrity (based in part on natural range of variation) but still calls for both coarse- and fine-filter approaches. In the Forest Service’s evaluations of the alternatives to the planning rule (USDA FS 2012), the terms coarse filter and fine filter are referred to extensively as “well-developed concept[s] in the scientific literature [with] broad support from the scientific community and many stakeholders.” However, the debate continues on an appropriate balance between coarse-filter (ecosystem and biodiversity) and

fine-filter (species-specific) planning direction (Hayward et al. 2016,² Schultz et al. 2013).

Also, although not part of the NWFP per se, some previous elements of the U.S. Forest Service’ Forest Inventory and Analysis (FIA) program³—which have since been drastically reduced or are no longer being carried out—provided much-needed information for monitoring biodiversity of trees, vegetation, and lichens.⁴ The FIA program has become a de facto biodiversity monitoring program, at least for selected vegetation and floral elements.

In the previous science synthesis, Marcot and Molina (2006) concluded that NWFP directions for establishing effectiveness monitoring of forest biodiversity elements for other than selected species remained mostly unmet (beyond the FIA-identified biodiversity indicators). This remains true today, but much information has been provided by research studies and gathered by agency programs on basic occurrence, distribution, and ecology of rare and poorly known late-successional and old-growth-associated species. The 2006 synthesis also provided the following suggestions:

- Engage research partnerships to fill key information gaps on rare and little-known late-successional and old-growth species.
- Clarify objectives and expectations of implementing a coarse- and fine-filter conservation approach to managing for viable and persistent species populations.
- Validate the use of surrogates (e.g., indicator and focal species) for species and conservation objectives.
- Develop and maintain databases from ongoing inventory, survey, and any monitoring programs.

² Hayward, G.D.; Flather, C.H.; Rowland, M.M.; Terney, R.; Mellen-Mclean, K.; Malcolm, K.D.; McCarthy, C.; Boyce, D.A. 2016. Applying the 2012 planning rule to conserve species: a practitioner’s reference. Unpublished paper. On file with: Bruce Marcot, Forestry Sciences Laboratory, 620 SW Main, Portland, OR 97205. 78 p.

³ <http://www.fia.fs.fed.us/>.

⁴ McCune, B. Personal communication. Professor, Department of Botany and Plant Pathology, Oregon State University, 2082 Cordley Hall Corvallis, OR 97331-2902. <http://bmccune.weebly.com/>.

- Develop, test, and implement species survey designs.
- Explore habitat modeling and decision-support tools to meet some conservation objectives.
- Develop and implement an effectiveness monitoring framework.

The current synthesis determines the degree to which these suggestions have been met.

New Learning and Recent Issues

Much has been learned since the 2006 synthesis (Haynes et al. 2006) about conditions and dynamics of forest ecosystems and their organisms in the Pacific Northwest and throughout the West. The issue of climate change and its known and projected impacts on systems has become a foremost research topic (Bagne et al. 2011, Vose et al. 2012). Occurrence and effects of large-scale wildfire have become major issues of research and management focus (Sheehan et al. 2015, Wimberly and Liu 2014). Studies on the effect of fire suppression on vegetation succession are needed, however, as are studies examining how suppression activities affect subsequent fire behavior but also how it changes vegetation conditions as habitat for many species (see chapter 3). Concern over invasive species also has elevated (Jones et al. 2010, Wilson et al. 2009; also see chapter 3). We address these and other issues in sections that follow.

Additionally, of increased focus is how early-successional vegetation provides habitats for many species (Hagar 2007, Swanson et al. 2011). Another topic of continued interest is the importance of conditions in the managed-forest matrix and connectivity of late-successional and old-growth forests and late-successional reserves (LSRs) in the face of fire, climate change, and increased pressure on matrix-land resources (Suzuki and Olson 2008, Wilson and Puettmann 2007). As well, the roles and conditions of rare and little-known species have been addressed (Raphael and Molina 2007). In general, much more detailed information is now available on vertebrates than on most other species groups.

Guiding Questions

This chapter reviews the scientific understanding of the ecology and conservation of species associated with late-successional and old-growth forests. We summarize science findings on the conservation strategy of the NWFP and its provision for these species; scientific progress since the previous NWFP evaluations (Diaz and Haynes 2002, Haynes and Perez 2001, Haynes et al. 2006); and the outcome of the NWFP Survey and Manage program.

We review advancements on science and conservation through the following questions:

- What is the current scientific understanding of the rarity of late-successional and old-growth-associated species?
- Is forest management under the NWFP providing habitat for rare and uncommon species as planned? Are rare and uncommon species maintaining populations under NWFP management? How effective are the management recommendations for habitat conservation in retaining these species across treated landscapes?
- Have we accumulated enough information to change the management status of these species? Are there species originally ranked as having low potential for persistence that are now of less concern, particularly with the reduction in harvest levels of late-successional and old-growth forest that has occurred under the NWFP? Are there late-successional and old-growth species originally ranked as high persistence or not initially identified as conservation concerns that have been added to lists of species of concern?
- What are results of research on the effects of prescribed fire and wildfire on rare and uncommon late-successional and old-growth species?
- What are results of research on the effectiveness of site buffers as compared with landscape-scale habitat management for ensuring late-successional and old-growth species persistence, dispersal, and habitat connectivity?
- How has the ISSSSP served to provide information on late-successional and old-growth-associated species under the NWFP?

- Does the current list of special status and sensitive species adequately represent rare late-successional and old-growth species with risks to population persistence?
- What are new issues related to conservation of biodiversity in the NWFP area?

Agency Programs on Other Species and Biodiversity of Older Forests

Survey and Manage Program

Following the 1993 report of the Forest Ecosystem Management Assessment Team or FEMAT (1993), and as part of the initial creation of the NWFP, the NWFP Survey and Manage program was instituted in 1993 as part of a final environmental impact statement and record of decision for amendments to U.S. Department of Agriculture (USDA) Forest Service and U.S. Department of the Interior (USDI) Bureau of Land Management (BLM) planning documents for federal public lands within the range of the northern spotted owl. The Survey and Manage program was then amended by a 2001 record of decision for amendment to the Survey and Manage, protection buffer, and other mitigation measures standards and guidelines. The amendment established (1) an annual species review panel process to evaluate monitoring and research findings and to recommend to the regional forester of Region 6 appropriate conservation categories for all late-successional and old-growth-associated species not otherwise provided for by the NWFP guidelines, and (2) a set of site survey protocols⁵ and management recommendations for detecting and conserving sites with rare and little-known species under the NWFP. The annual species review sessions were designed as rigorous, 10-person panels consisting of 5 biologists and 5 managers and used a Bayesian network decision modeling construct to help evaluate knowledge and explicitly represent uncertainty of each species in documented, repeatable procedures (Marcot et al. 2006). Mostly because of high costs and administrative complexities, no formal annual species review has been conducted since 2003.

The Survey and Manage program was established under the NWFP as a means of collecting information on, and providing appropriate conservation direction for, rare and poorly known late-successional and old-growth-associated species under the precautionary principle (resisting implementation of untested or disputed activities that may have adverse effects) and an adaptive management process (Marcot et al. 2006, Molina et al. 2003, USDA and USDI 2001). From the initial list of 1,120 late-successional and old-growth-associated species evaluated by FEMAT (1993), various mitigation means under the NWFP Survey and Manage program narrowed the list in 2001 to 296 individual species and 4 arthropod species groups. The Survey and Manage program was then abolished, and, under a management policy decision of the agencies, 152 of the 296 species were moved to the USDA Forest Service Sensitive Species program and the USDI BLM Special Status Species program, but the court then mandated that the Survey and Manage record of decision be reinstated. Eventually, the two agencies' species programs were merged into the ISSSSP, discussed more fully below, which has since held the responsibility for evaluation of late-successional and old-growth species in the region. Also in the interim, a set of new national forest planning regulations have been instituted that provide impetus for considering other species, biodiversity, ecosystems, dynamics, and functions of both older and early-seral forests (Schultz et al. 2013). We discuss these updates to Forest Service and BLM planning guidelines and regulations further below.

The Survey and Manage program has had an unstable existence, having been established in 1994 (USDA and USDI 1994) with corrections to its standards and guidelines published in 2001 (USDA and USDI 2001), abolished by the agencies in 2004 (USDA and USDI 2004), reinstated by the court in 2006, again abolished by the agencies in 2007, challenged in 2008, and with a court ruling in 2009 that the 2007 Forest Service environmental impact statement was flawed and the court subsequently approving a settlement agreement in 2011. The timber industry then challenged the settlement agreement in 2011 (and subsequently dropped their appeal in 2015), the Ninth Circuit Court of Appeals reversed and remanded the approval of the settlement

⁵ <http://www.blm.gov/or/plans/surveyandmanage/protocols/>.

agreement in 2013, and then in 2014, the 2007 records of decision were vacated.

Vacatur of the two 2007 records of decision (in 2007, BLM and the Forest Service each issued separate records of decision) has had the effect of returning the agencies to the status quo in existence prior to the 2007 records of decision. The status quo existing before the 2007 records of decision was defined by three previous court rulings, as follows. First is the 2006 court order reinstating the 2001 record of decision, including any amendments or modifications that were in effect as of March 21, 2004. This ruling incorporated the 2001, 2002, and 2003 annual species review changes. Second was the 2006 court-ordered categories of activities that could proceed without conducting predisturbance surveys or site management for species: (1) thinning in forests less than 80 years old; (2) replacement or removal of water culverts; (3) activities for improvement of riparian and stream areas; and (4) treatment of hazardous fuels, including use of prescribed fire; these reinstatements were retained in the later court rulings mentioned above. Third was the 2006 court ruling that vacated the 2001 and 2003 annual species review category change and subsequent removal of reference to the red tree vole in a portion of its range, returning the species to its prior monitoring status throughout its range.

At present, oversight of the Survey and Manage standards and guidelines implementation is consigned to staff members within the ISSSSP (Region 6 and Oregon BLM) and the Region 5 regional wildlife program manager within the Ecosystem Conservation staff. These individuals coordinate revision of management recommendations and survey protocols, assist field specialists in implementing the standards and guidelines, resolve issues between Survey and Manage species management and meeting other resource objectives, coordinate data management between the agencies preparing for an Annual Species Review, stay abreast of taxonomic updates, and coordinate methods for filling information gaps. To clarify, Forest Service Region 5 is not formally a part of the ISSSSP, which is unique to Forest Service Region 6 and Oregon-Washington BLM.

The list of late-successional and old-growth-associated species as provided by FEMAT (1993) had been evaluated by the Forest Service and BLM under the Survey and

Manage program's annual species reviews, using a set of published guidelines (table 6-1) to determine species' potential need for more specific and additional conservation. Based on an evaluation of the occurrence of, and scientific knowledge on, the species, about 400 species of amphibians, bryophytes, fungi, lichens, mollusks, vascular plants, arthropod functional groups, and one mammal were deemed to be potentially at-risk, and the rest of the species were deemed to be adequately provided under the NWFP guidelines; the genealogy through 2006 of the many species lists are covered by Marcot and Molina (2006) and Molina et al. (2006). The annual species reviews developed and adopted use of a Bayesian network decision modeling approach to help wade through the complex evaluation guidelines (table 6-1) and to document results on each species (Marcot et al. 2006).

Under the Survey and Manage program, about 68,000 sites with presence of Survey and Manage species were identified by surveys, and new ecological knowledge was gained on about 100 species leading to their being removed from the protection list (Molina et al. 2006). Additionally, a set of field and management guides were produced on aquatic and terrestrial mollusks and fungi,⁶ and guidelines were published on assessing rare species of lichens (Edwards et al. 2004), fungi (Castellano et al. 2003, Molina 2008), and other taxa. Eventually, the high cost of maintaining the Survey and Manage program, running into several tens of millions of dollars, with its annual species reviews and all other activities associated with compiling scientific and monitoring information on late-successional and old-growth-associated species, was a factor considered by managers in their decision to abolish the program and enfold it into the ISSSSP.

Understanding the distributions and disturbance responses of rare species is a perennial problem in ecology, the main issues of which include securing adequate sample sizes for statistical analyses (Cunningham and Lindenmayer 2005). Methods for increasing confidence in such studies include stratifying samples, such as demonstrated by Edwards et al. (2005) with five rare epiphytic macrolichens in the Pacific Northwest United States.

⁶ <http://www.blm.gov/or/plans/surveyandmanage/field.php>.

Table 6-1—Guidelines for determining whether late-successional and old-growth forest (late-successional and old-growth)-associated species under the Northwest Forest Plan (NWFP) may need additional conservation consideration, as required under the Survey and Manage program 2001 record of decision (USDA and USDI 2001)

Evaluation category^a	Description in record of decision (USDA and USDI 2001)
1. Geographic range	The species must occur within the NWFP area or near the NWFP area and have potentially suitable habitat within the NWFP area.
2. Late-successional and old-growth association	A species is considered to be closely associated with late-successional and old-growth forests if it meets at least one of the following criteria: • The species is significantly more abundant in late-successional and old-growth forest than in young forest, in any part of its range. • The species shows association with late-successional and old-growth forest and may reach highest abundance there, and the species requires habitat components that are contributed by late-successional and old-growth forest. • The species is associated with late-successional and old-growth forest, based on field study, and is on a federal U.S. Fish and Wildlife Service (USFWS) list or state threatened or endangered list; the USFWS candidate species list; a Bureau of Land Management or Forest Service special status species list in California, Oregon, or Washington; or is listed by the states of California, Oregon, or Washington as a species of special concern or as a sensitive species. • Field data are inadequate to measure strength of association with late-successional and old-growth forest; the species is listed as a federal USFWS threatened and endangered species; and the Forest Ecosystem Management Assessment Team suspected, or the panel doing the final placement in Species Review Process suspects, that it is associated with late-successional and old-growth forest.
3. Plan provides for persistence	The reserve system and other standards and guidelines of the NWFP do not appear to provide for a reasonable assurance of species persistence. Criteria indicating a concern for persistence, i.e., one or more of the following criteria must apply: • Low to moderate number of likely extant known sites/records in all or part of a species range. • Low to moderate number of individuals. • Low to moderate number of individuals at most sites or in most populations. • Very limited to somewhat limited range. • Distribution within habitat is spotty or unpredictable in at least part of its range. • Very limited to somewhat limited habitat. Criteria indicating little or no concern for persistence, usually, most of the following criteria must apply: • Moderate to high number of likely extant sites/records. • Sites are relatively well distributed within the species range. • High proportion of sites and habitat in reserve land allocations; or limited number of sites within reserves, but the proportion or amount of potential habitat within reserves is high and there is a high probability that the habitat is occupied. • Matrix standards and guidelines or other elements of the NWFP provide a reasonable assurance of species persistence.
4. Data sufficiency	Information is insufficient to determine whether Survey and Manage basic criteria are met, or to determine what management is needed for a reasonable assurance of species persistence.

Table 6-1—Guidelines for determining whether late-successional and old-growth forest (late-successional and old-growth)-associated species under the Northwest Forest Plan (NWFP) may need additional conservation consideration, as required under the Survey and Manage program 2001 record of decision (USDA and USDI 2001) (continued)

Evaluation category ^a	Description in record of decision (USDA and USDI 2001)
5. Practicality of survey	Surveys are considered “practical” if all of the following criteria apply: • The taxon appears annually or predictably, producing identifying structures that are visible for a predictable and reasonably long time. • The taxon is not so minuscule or cryptic as to be barely visible. • The taxon can authoritatively be identified by more than a few experts, or the number of available experts is not so limited that it would be impossible to accomplish all surveys or identifications for all proposed habitat-disturbing activities in the NWFP area needing identification within the normal planning period for the activity. • The taxon can be readily distinguished in the field and needs no more than simple laboratory or office examination to confirm its identification. • Surveys do not require unacceptable safety (5a) or species risks. • Surveys can be completed in two field seasons (about 7 to 18 months). • Credible survey methods for the taxon are known or can be developed within a reasonable time period, i.e., about 1 year.
6a. Relative rarity	The species is relatively rare and all known sites or population areas are likely to be necessary to provide reasonable assurance of species persistence, as indicated by one or more of the following: • The species is poorly distributed within its range or habitat. • Limited dispersal capability on federal lands. • Reproduction or survival not sufficient. • Low number of likely extant sites/records on federal lands indicates rarity. • Limited number of individuals per site. • Declining population trends. • Low number of sites in reserves or low likelihood of sites or habitat in reserves. • Highly specialized habitat requirements (narrow ecological amplitude). • Declining habitat trend. • Dispersal capability limited relative to federal habitat. • Habitat fragmentation that causes genetic isolation. • Microsite habitat limited. • Factors beyond management under the NWFP affect persistence, but special management under the NWFP will help persistence.
6b. Relative uncommonness	The species is relatively uncommon rather than rare, and not all known sites or population areas are likely to be necessary for reasonable assurance of persistence, as indicated by one or more of the following: • A higher number of likely extant sites/records does not indicate rarity of the species. • Low to high number of individuals/site. • Less restricted distribution pattern relative to range or potential habitat. • Moderate to broad ecological amplitude. • Moderate to high likelihood of sites in reserves. • Populations or habitats are stable.

^a If criteria for any evaluation category were met, then the species may be further considered for needing additional conservation beyond what the NWFP generally provides; such further consideration was addressed during annual species reviews under the NWFP Survey and Manage program.

Interagency Special Status and Sensitive Species Program

The ISSSSP⁷ was formed in 2005 as an interagency Forest Service Region 6 and BLM Oregon/Washington program for regional-level approaches for conservation and management of rare (but neither federally listed threatened nor endangered) species that would meet criteria for the two agencies' lists of special status species and sensitive species. Its geographic and ecological scope includes and exceeds that of the NWFP and late-successional and old-growth forests in Washington and Oregon. The ISSSSP is not a reformulation of the NWFP Survey and Manage program, although it has taken on some of those functions pertaining to evaluation of the conservation status of species, development of some species survey and monitoring protocols, and other items. The ISSSSP addresses species across Forest Service and BLM lands in Oregon and Washington (but not California), implementing the Forest Service sensitive species policy (FSM 2670) and BLM special status species policy (BLM 6840) and providing oversight of the Survey and Manage standards and guidelines. Criteria for determining Forest Service sensitive species are quite different from the Survey and Manage species criteria discussed above (also see table 1).

California, particularly northwest California within the NWFP area, does not have an organization equivalent to the ISSSSP, which is a collaboration unique to Washington and Oregon. In California, instead, the Forest Service Region 5 implements the national Forest Service sensitive species policy (FSM 2670) and results are overseen by various Forest Service regional office staff for the entire state, not just for the NWFP area and the six national forests therein. California BLM includes lands within the NWFP area, and those are overseen by the BLM Redding Resource Area, Arcata Resource Area, and the Kings Range National Conservation Area, all within the BLM Ukiah District.

The ISSSSP has produced a wide array of products related to conservation of rare, nonlisted species. Products

include species fact sheets, conservation assessments, conservation strategies, inventory reports, inventory and survey protocols and methods workshops, and results of studies. The most recent program update⁸ (June 2015) mentions reorganization of the program's conservation and inventory information on bats and fungi (covered below). The ISSSSP partners with and supports a variety of research and academic institutions to provide key information on rare species of conservation concern within its geographic venue.

Unique among federal land management agencies, the ISSSSP has developed criteria used in common with Forest Service and BLM for including species on sensitive and special status lists. The ISSSSP considers species for such listing by using independent information from the Oregon Biodiversity Information Center⁹ Washington Natural Heritage Program¹⁰ and NatureServe.¹¹

The current list of Survey and Manage species¹² dates to December 2003 and includes 298 species: 189 fungi, 15 bryophytes, 40 lichens, 12 vascular plants, 36 snails and slugs (mollusks), 4 amphibians, 1 mammal (red tree vole, treated below), and 1 bird (great gray owl, *Strix nebulosa*).

Implications of Forest Service and BLM Planning Directions

The current planning rule for the U.S. Forest Service (2012: 21174)¹³ states that its intent is

... to provide for the diversity of plant and animal communities, and keep common native species common, contribute to the recovery of threatened and endangered species, conserve proposed and candidate species, and maintain species of conservation concern within the plan area, within Agency authority and the inherent capability of the land.

⁷ <http://www.fs.fed.us/r6/sfpnw/issssp/>.

⁸ <http://www.fs.fed.us/r6/sfpnw/issssp/documents3/update-2015-06.pdf>.

⁹ <http://inr.oregonstate.edu/orbic>.

¹⁰ <http://www.dnr.wa.gov/natural-heritage-program>.

¹¹ <http://www.natureserve.org/>.

¹² <http://www.blm.gov/or/plans/surveyandmanage/files/sm-fs-enc3-table1-1-dec2003wrtv.pdf>.

¹³ <http://www.fs.usda.gov/detail/planningrule/home/?cid=stel-prdb5359471>.

The rule establishes guidelines and mandates for ecological sustainability, particularly for ecosystem integrity defined as the maintenance or restoration of terrestrial and aquatic ecosystems and watersheds, and their structure, function, composition, and connectivity. The rule also explicitly adopts a coarse- and fine-filter approach (further discussed below) to managing for diversity of plant and animal communities beginning with maintaining or restoring the diversity of ecosystem and habitat types including rare aquatic and terrestrial plant and animal communities, as well as identifying species of conservation concern to be designated by the responsible official, in coordination with the regional forester, based on scientific information. In the Pacific Northwest, the Forest Service is currently producing a draft list of potential species of conservation concern to facilitate efficiencies when the region undergoes plan revision under the 2012 planning rule.

Additional parts of the 2012 planning rule for the Forest Service provide guidance on monitoring, which it defines as “a systematic process of collecting information to evaluate effects of actions or changes in conditions or relationships” (USDA FS 2012:21271). The planning rule also provides guidance on managing ecological systems at the broad scale and for specific ecosystem elements, such as individual species, at finer scales. As such, specifically, the planning rule refers to coarse-filter management as “designing ecosystem ... connectivity based on landscape patterns of forests, grasslands, rangelands, streams, and wetlands that were created under ecological processes and landscape disturbance regimes that occurred before extensive human alteration” (section 23.11b: Ecosystem Integrity), and fine-filter management as “species-specific plan components, including standards and guidelines, for each of those species” (section 23.13: Species-Specific Plan Components for At-Risk Species).

Schultz et al. (2013) recommended directly monitoring selected species of conservation concern and focal species because of inconsistencies in the 2012 planning rule between its operational requirements and its generous discretionary allowances. They suggested that monitoring should evaluate viability of such species and that management should do no harm to species for which viability

cannot be provided solely on Forest Service lands; and that monitoring should specify trigger points to spark reviews of management activities affecting species conservation.

As a point of history, BLM proposed a new rule for resource management planning nationally on BLM lands (USDI BLM 2016), but in February 2017, Congressional action nullified the regulations. In the NWFP area, BLM Resource Management Plans (RMPs) are in effect from records of decision signed in August 2016.¹⁴ The new RMPs are intended to provide protection for northern spotted owls, listed fish species, and water resources, and provide for jobs, recreation, and timber harvest. At this point, it is unclear if the RMPs will provide additional guidelines for conservation of other old-forest species and biodiversity under the NWFP in addition to guidelines provided by ISSSP.

Key Findings

New Information on Other Species and Biodiversity of Older Forests

In this section, we review new research information on individual species and species groups under the NWFP and in old-forest environments, conducted mostly since the previous science synthesis (Haynes et al. 2006).

Fungi—

Fungi are an important part of forest ecosystems. Fungi have always been a conservation challenge in terms of species identification, taxonomic designation, inventory and monitoring of furtive and seldom-appearing species, and understanding of their key ecological roles in late-successional and old-growth forest ecosystems. Many fungi are rare or little known, but much has been learned about some aspects of species occurrence and distribution since the previous science synthesis. Some of this work is presented in peer-reviewed publications, and other work is available through an agency peer review process. Recent regional work provides information on California fungal species. Other work provides a better understanding of the status of fungal species in the NWFP area.

¹⁴ <https://www.blm.gov/or/plans/rmpswesternoregon/>.

Occurrence of fungal species is influenced, at least in part, by the type and intensity of disturbances, time since disturbance, and vegetation development, and by forest stand management and forest age class (Heithecker and Halpern 2006, Trofymow et al. 2003). Studies by Hebel et al. (2009) suggested that high-severity wildfire can reduce or prevent colonization of, and can kill, beneficial arbuscular mycorrhizal fungi. As noted in chapter 3 and in Reilly et al. (2017), current rates of high-severity fire in the NWFP area are very low even in the moist forests where fire was historically infrequent. In the dry forests, where fires were historically frequent, recent fire rotations have been well below the historical levels; however, in forests that historically had low-severity fire and little high-severity fire, the recent amounts of high-severity fire appear to be higher. It is unclear how current fire regimes might affect forest structure, age class, and disturbance intensity.

Presumably, native fungi species were able to persist across landscapes with frequent to very frequent fire (<50 years) and in landscapes with occasional, moderately frequent, mixed-severity fire (50 to 200 years). Fire suppression has affected the various forest ecosystems of the NWFP area in different ways (chapter 3), although little is known about effects of fire suppression activities on fungi. Luoma and Eberhard (2005) urged the conservation of rare truffle and mushroom species "... in a manner that recognizes their different responses to forest disturbance." They also hypothesized that fire suppression activities may have favored mushroom production over truffle production, and that presence of fire is a factor in the reproductive evolution of ectomycorrhizal fungi. Thus, they concluded that providing for ectomycorrhizal fungi would include restoring forest health from the adverse effects of decades of fire suppression.

Many fungi species are soil dwellers and mostly subterranean, and, with intermittent fruiting cycles, they are not easy to detect and collect. Determining presence can be difficult, but they can play major roles in nutrient uptake by trees and other plants and in aiding coarse-wood decomposition, contributing to soil organic matter, and maintaining overall forest health.

Luoma (2001)¹⁵ found that retention of green trees in late-successional forests helped retain *Arcangeliella camphorata*, a rare truffle fungus, as compared to the species' loss in clearcuts. Trappe et al. (2009) provided an extensive study of the distribution, ecology, and conservation of truffle species in the Pacific Northwest. In southwest Oregon, Clarkson and Mills (1994) and Amaranthus et al. (1994) also had previously demonstrated that clearcuts ranging from 4 to 27 years since harvest nearly eliminated truffle production and that retention of mature trees and coarse woody debris promote their diversity and abundance. Marcot (2017) reviewed the role of fungi in wood decay of forest ecosystems of the Pacific Northwest.

Some fungi are dispersed in unusual ways, such as on the beaks of foraging and cavity-excavating woodpeckers (Jusino et al. 2016). Fungi such as truffles and their ectomycorrhizal sporocarps are key food resources for northern flying squirrels (*Glaucomys sabrinus*) (Lehmkuhl et al. 2004); in turn, flying squirrels are a key prey species of northern spotted owls in parts of the owl's range. Deliberate introduction of fungi in live trees is a management tactic sometimes used to induce wood decay and create partially dead trees and snags for wildlife use such as in western Washington (Bednarz et al. 2013).

In 1994, the Survey and Manage program listed some 234 rare fungi species associated with late-successional and old-growth forests (Molina 2008). Molina's (2008) review of mycology herbaria eventually yielded 14,400 records of these species, with 55 percent of the species found at 20 or fewer sites and 42 percent found at 10 or fewer sites; it is unclear which of the 42 percent are rare or undersurveyed. Some 90 percent of the species had some fraction of their locations occurring within reserves, but only a third of the species had all their locations within reserves. This led Molina (2008) to conclude that fine-filter conservation of rare species outside reserves was needed to help ensure conservation of the entire late-successional and old-growth-associated fungal biota.

¹⁵ Luoma, D.L. 2001. Monitoring of fungal diversity at the Siskiyou integrated research site with special reference to the survey and manage species *Arcangeliella camphorata* (Singer & Smith) Pegler & Young. Unpublished report. On file with: Chetco Ranger District, Siskiyou National Forest, Brookings, OR 97415.

Overall, although this work has provided much information on presence and distribution of rare fungi, in general their species-specific status and trends are still not well known.

The ISSSSP Fungi Work Group has compiled an annotated bibliography on rare fungi species on the Special Status Species list for California, Oregon, and Washington.¹⁶ This work includes 174 references, each with annotated findings and indexed to keywords pertaining to ecological and management topics.

Mushroom collection for individual use is a popular recreational activity in forests of the Pacific Northwest (Trappe et al. 2009). Individual national forests in the NWFP area may have specific regulations on commercial mushroom harvests. For example, Siuslaw National Forest (2007) sells individual commercial collection permits, up to 1,000 permits per year, with permits being unlimited by weight or amount of mushrooms collected. Alexander et al. (2002) compared the economics of timber harvest with

mushroom harvest in the Pacific Northwest and found that some mushrooms (e.g., chanterelles, morels) have lower value and some (matsutake) have about the same value as commercial timber, as measured by soil expectation value analysis. At present, though, we have encountered no studies on the impacts of mushroom collection on these species' populations in the NWFP area.

Lichens—

Much has been learned over the past decade about the occurrence and rarity of forest lichens (table 6-2) and the effects of forest management in the Pacific Northwest. Lichens play important ecological roles in late-successional and old-growth forests of the Pacific Northwest (chapter 3). Epiphytic macrolichens, because of their complete reliance on atmospheric sources of water and nutrients, are useful for monitoring air quality and climate (fig. 6-1) (Geiser and Neitlich 2007; similar work by Root et al. 2015 followed

¹⁶ <http://www.fs.fed.us/r6/sfpnw/issssp/documents3/cpt-fu-effects-guidelines-att3-annotated-bibliography-2013-10.docx>.

Table 6-2—Selected recent findings on lichen species associated with late-successional and old-growth forest conditions within the Northwest Forest Plan area

Lichen taxa	Topic	Source
<i>Bryoria subcana</i> to <i>B. fuscescens</i>	Taxonomy of <i>Bryoria</i> section Implexae	Velmala et al. 2014
<i>Dermatocarpon</i>	<i>Dermatocarpon luridum</i> now found to not occur in the United States. The species in the Pacific Northwest (PNW) is <i>D. meiophyllizum</i> .	Glavich 2009, Glavich and Geiser 2004
<i>Fuscopannaria</i>	<i>Fuscopannaria saubinetii</i> (misidentification) does not occur in the PNW. Previous records were misidentifications of <i>F. pacifica</i> , a common species in the PNW.	Jørgensen 2000
<i>Leptogium</i>	<i>Leptogium burnetiae</i> var. <i>hirsutum</i> is now a synonym of <i>L. hirsutum</i> , which does not occur in the PNW. The taxon of conservation concern is <i>L. burnetiae</i> .	Esslinger 2015
<i>Leptogium</i>	<i>Leptogium rivale</i> is changed to <i>Scytinium rivale</i> . <i>L. teretiusculum</i> is changed to <i>S. teretiusculum</i> .	Otálora et al. 2014
<i>Usnea</i>	<i>Usnea hesperina</i> is changed to <i>U. subgracilis</i> . Previously, <i>U. hesperina</i> and <i>U. subgracilis</i> were both considered synonyms of <i>U. schadenbergiana</i> ; <i>U. schadenbergiana</i> is distinct from <i>U. hesperina</i> and <i>U. subgracilis</i> .	Truong et al. 2013



Figure 6-1—Some arboreal, epiphytic lichens, such as this *Alectoria* in Wind River Experimental Forest in the south Washington Cascade Range, are fully dependent on atmospheric sources of water and nutrients, and thus can serve as sensitive indicators for monitoring of air pollution and climate change.

mostly east of the NWFP area).¹⁷ Trofymow et al. (2003) found that arboreal lichen abundance and species richness differed between mature (75 to 95 years old) and old-growth (>240 years old) conifer forest stands on Vancouver Island, and thus can serve as indicators of late-successional and old-growth forest conditions. Arsenault and Goward (2016) reported that, in inland wet conifer forests of British Columbia, some macrolichens, such as *Lobaria pulmonaria*, are good indicators of old-growth forests in some ecological conditions but not in others, highlighting the

need for caution when determining species' association with, and dependence on, old-growth forest conditions. Sillett et al. (2000) showed that dispersal can limit the development of some lichens (e.g., *Lobaria oregana*) in late-successional and old-growth forests, as well as in young forest plantations.

Bokhorst et al. (2015) reported that lichen species in southern Norway have a large impact on associated invertebrate communities, varying with lichens that differ in nitrogen fixation and nutrient concentration, thallus structure, and terricolous versus arboreal or epiphytic habitat. Whether such contributions of lichen diversity to overall lichen-invertebrate communities occur within

¹⁷ Also see: <http://people.oregonstate.edu/~mccuneb/epiphytes.htm> and <http://people.oregonstate.edu/~mccuneb/biblio.htm>.

late-successional and old-growth forests of the Pacific Northwest is apparently unstudied.

Effects of wildfire and fire suppression activities on late-successional and old-growth-associated lichens have been poorly studied. Large-scale, stand-replacing fires will likely reduce distribution and biomass of late-successional and old-growth-associated lichens over decades to centuries, as a function of the species' dispersal mechanisms and limitations (Sillett et al. 2000).

Recent studies suggest that variable-density thinning treatments in young forests of Douglas-fir (*Pseudotsuga menzeisii*) and western hemlock (*Tsuga heterophylla*) can provide habitat for a variety of lichen species (Root et al. 2010) that could complement lichen assemblages found in unthinned late-successional and old-growth forest environments. Earlier studies suggested that uniform thinning of dense forest stands provides only minor improvement of habitat of lichens and bryophytes, whereas some variable-density thinning can help promote development of tree branches lower in the canopy profile as substrates for lichens and bryophytes. Earlier studies (e.g., Neitlich and McCune 1997) also revealed that forest structure may play a greater role in determining lichen diversity and biomass than forest age per se from observations that lichen biomass can be greater in structurally diverse young forests with gaps and older remnant trees than in some old-growth forests. Neitlich and McCune (1997) suggested, for conserving epiphytic macrolichens, to protect forest gaps, wolf trees (large trees with wide-spreading crowns), and old remnant and legacy trees. Retention of propagule sources, such as old legacy trees in younger stands, and older-forest patches in cutting units, can be critical to providing for old-growth-associated epiphytes (Sillett et al. 2000) and cryptogams (Hofmeister et al. 2015).

Much work has also been done over the past decade on riparian lichens. Three groups of lichens depend on riparian areas:¹⁸ (1) nonepiphytic, instream species (e.g., *Dermatocarpon meiohyllizum*, *Leptogium rivale*, and *Peltigera hydrothyria*); (2) epiphytic species occurring in

streamside and lakeshore forest and woodland environments (e.g., *Leptogium polycarpum*, *Ramalina thrausta*, *Sticta fuliginosa*, *Usnea wirthii*, and many cyanolichen species); and (3) epiphytic species in forested wetlands, particularly in Oregon ash (*Fraxinus latifolia*) swamps (e.g., *Fusco-pannaria mediterranea*, *Hypotrachyna riparia*, and many cyanolichens). A number of these riparian lichens are likely afforded habitat under the NWFP's Aquatic and Riparian Conservation Strategy. For further information on riparian lichens in the Pacific Northwest, see McCune et al. (2002) and Ruchty (2000).

As with fungi, many species of lichens and other rare and poorly known species in the Pacific Northwest are difficult to detect, inventory, monitor, and study because they require specialized expertise in the field and the laboratory. To address such problems, the Survey and Manage program spurred the development of methods for statistically determining the occurrence and frequency of rare species of lichens (Edwards et al. 2004, 2005). A guide was produced under the Survey and Manage program on natural history and management considerations of selected lichen species under the NWFP (Leshner et al. 2003).

Additionally, Miller et al. (2017) tested techniques for surveying *Lobaria oregana*, a rare canopy lichen, at the southern edge of its range in northwest California within the NWFP area. They found the species on branches of large trees in the mid-crown, and on boles of small trees near ground level, suggesting that the species benefits from cool, wet microclimates, and that maintaining such microhabitats is key to its long-term viability. They also concluded that ground surveys are useful for detecting abundant lichen species, but that tree-climbing to conduct canopy surveys also may be needed to detect low-abundance species associated with mid- or upper-canopy conditions.

Bryophytes—

Bryophytes—including hornworts, liverworts, and mosses (in general, nonvascular plants)—are a conspicuous component of many late-successional and old-growth forests of the Pacific Northwest, and many species have been part of the Survey and Manage program. Many late-successional and old-growth-associated bryophytes may be sensitive to disturbance and may require old-forest stands larger than

¹⁸ McCune, B. Personal communication. Professor, Oregon State University, Corvallis, OR 97331.

2.5 ac (1 ha) (Halpern et al. 2012) or with more than 15 percent retention of older green trees in dispersed retention harvests (Heithecker and Halpern 2006).

Mölder et al. (2015) reported that 31 bryophyte species in north Germany can serve as ancient woodland indicators; a number of bryophyte species of the Pacific Northwest can serve similar functions (Rambo and Muir 1998, 2002). In an older study, Ryan et al. (1998) found that cryptogams in mixed-conifer forests of southern Vancouver Island were largely associated with humus, and especially coarse woody debris and rock substrates, and that old-growth forests variously excluded shade-intolerant species and selected for shade-tolerant species.

A recent taxonomic update has changed *Diplophyllum plicatum* to *Douinia plicata* (Konstantinova et al. 2013).

Pacific Northwest forests are the main source of commercially harvested moss in North America with dozens of species likely affected by the practice (Peck 2006, Peck and Christy 2006). The main mosses collected are epiphytes that require 15 to 25 years to recover, and which are most abundant in riparian and low-elevation forests and absent or much less abundant in young (<70 years) Douglas-fir plantations (Peck 2006). Commercial harvest methods of stripping entire moss mats can greatly impede recovery of the species (Peck and Frelich 2008) and may be a major, local threat to this group.¹⁹ Peck and Moldenke (2011) reported a wide array of invertebrates (205 morphospecies) associated with subcanopy epiphytic bryophyte mats subject to commercial harvest in the Coast and Cascade Ranges of the Pacific Northwest, where more than 8.2 million lbs (3.7 million kg) of bryophytes are harvested per year. However, the impact of moss harvest on populations of these invertebrate species is undetermined.

Overall, few studies are available on the status and trends of the full suite of bryophyte species (and associated other taxa) considered under the NWFP and ISSSSP. In the NWFP area, some individual national forests in Region 6 (Washington and Oregon) have their own direction regarding commercial moss collection, although there is no

overall regional direction. For example, Siuslaw National Forest and Willamette National Forest may have instituted local direction of moss collection under their special forest products or nontimber forest products regulations (e.g., Siuslaw National Forest 2007). Siuslaw National Forest's direction, for instance, permits commercial moss harvests to be limited to 16,000 lb (7258 kg) per year, from harvest areas that would be open for 12 years, with only one harvest area open at a time and rotated every 12 years, and harvest areas consisting of forest stands <110 years old (Siuslaw National Forest 2007).

Vascular plants—

Vascular plants are a conspicuous and important component of late-successional and old-growth forests in the NWFP area. The diversity of vascular plants tends to increase during structural development following logging in western hemlock-Douglas-fir, and to peak in old-growth forest conditions (Halpern and Spies 1995), but less is known in other vegetation zones of the area. Late-successional and old-growth-related vascular plants in the Pacific Northwest are generally considered to be protected under the NWFP guidelines and within LSRs, and have constituted only a small fraction of all Survey and Manage species thought to be closely associated with late-successional and old-growth forests (Marcot and Molina 2006).

Most studies of changes in plant communities over seral development have used a chronosequence (space-for-time) approach and compared composition or abundance in young logged stands with naturally regenerated stands in middle or later stages of development. These studies generally agree that understory plant species diversity increases following disturbance then decreases during canopy closure, and subsequently increases again and sometimes peaks in old-growth conditions (Halpern and Spies 1995, Jules et al. 2008). Few perennial understory plant species are either absent from, or restricted to, any specific stage of successional development, although the abundance of individual species changes over time (Halpern and Spies 1995). Some species are most abundant in old growth or are closely associated with old growth (table 6-3), but none of these species is on the current Survey and Manage species list and might be candidates

¹⁹ For a compendium of literature on moss harvesting, see: <http://bryophytes.science.oregonstate.edu/MossHarvest.htm>.

Table 6-3—Vascular plant species found to be associated with late-successional and old-growth forests in the Northwest Forest Plan area^a

Species	Halpern and Spies 1995 ^b	Lindh and Muir 2004 ^c	Jules et al. 2008 ^d
Trees/shrubs:			
<i>Abies amabilis</i>	✓		
<i>Acer circinatum</i>	✓		
<i>Taxus brevifolia</i>	✓		
<i>Tsuga heterophylla</i>	✓		
<i>Rhododendron macrophyllum</i>		✓	
<i>Vaccinium alaskaense</i>	✓		
<i>Vaccinium membranaceum</i>	✓	✓	
<i>Vaccinium parvifolium</i>	✓		
<i>Ribes lacustre</i>			✓
Low/subshrubs:			
<i>Berberis nervosa</i>	✓		
<i>Chimaphila menziesii</i>			
<i>Chimaphila umbellata</i>	✓	✓	
<i>Cornus canadensis</i>	✓		
<i>Gaultheria ovatifolia</i>	✓		
<i>Goodyera oblongifolia</i>		✓	✓
<i>Linnaea borealis</i>	✓	✓	
Herbs, forbs, and fern allies:			
<i>Achlys triphylla</i>	✓	✓	✓
<i>Adenocaulon bicolor</i>	✓		✓
<i>Anemone oregana</i>			✓
<i>Coptis laciniata</i>	✓	✓	
<i>Clintonia uniflora</i>	✓		✓
<i>Disporum hookeri</i>	✓		
<i>Galium</i> spp.			✓
<i>Lycopodium clavatum</i>	✓		
<i>Osmorhiza chilensis</i>			✓
<i>Pyrola asarifolia</i>	✓	✓	
<i>Senecio bolanderii</i>			✓
<i>Smilacena stellata</i>			✓
<i>Synthyris reniformis</i>	✓		
<i>Tiarella trifoliata</i>	✓	✓	
<i>Trillium ovatum</i>			✓
<i>Vancouveria hexandra</i>	✓		

Table 6-3—Vascular plant species found to be associated with late-successional and old-growth forests in the Northwest Forest Plan area^a (continued)

Species	Halpern and Spies 1995 ^b	Lindh and Muir 2004 ^c	Jules et al. 2008 ^d
Saprophytes/root parasites:			
<i>Corallorhiza maculata</i>			✓
<i>Corallorhiza mertensiana</i>	✓		
<i>Hemitomes congestum</i>			
<i>Hypopitys monotropa</i>			
<i>Monotropa uniflora</i>			
<i>Pterospora andromeda</i>			

^a None of these species is on the current Survey and Manage species list.

^b Oregon Coast Range and Oregon and Washington Cascade Range.

^c H.J. Andrews Experimental Forest, western Cascade Range, Oregon.

^d California Klamath Province.

for such consideration. Bailey et al. (1998) suggested that understory communities are relatively resilient to past cutting as few species differed in frequency of occurrence between young- and old-growth forests in the Oregon Coast and western Cascade Ranges. These results contrast with those of Halpern and Spies (1995) but are based on a much smaller sample size (9 vs. 196 plots) and are only from the drier part of the moist forest region.

The affinity of some species of vascular plants to old-growth forest conditions has been attributed to multiple mechanisms including the presence of canopy gaps and unique microclimatic conditions, greater heterogeneity in resources, and sensitivity to disturbance (e.g., logging and fire) coupled with low rates of growth and reestablishment (Halpern and Spies 1995). Jules (1998) found that ~97 percent of *Trillium ovatum* were eliminated following clear-cutting and planting of conifers. Furthermore, fragmentation of habitat had negative demographic consequences including decreased recruitment and seed production in this species (Jules and Rathcke 1999). Slow recovery rates of dispersal-limited, perennial herbs (e.g., *Trillium* spp., *Cornus canadensis*, *Clintonia uniflora*, *Disporum hookeri* var. *hookeri*) may take centuries for populations to recover following clearcutting (Kahmen and Jules 2005). Halpern and Spies (1995) suggested that long-term rotations (150 to 300 years) may be needed to maintain understory plant spe-

cies that require long recovery times following disturbance. Observations are lacking on how these dynamics differ from historical dynamics following natural disturbance (e.g., fire).

The response of understory vascular plants to management is known primarily from western hemlock-Douglas-fir forests (e.g., Halpern et al. 1999, Puettmann et al. 2013) and generally suggest that certain management activities can increase later seral species in previously managed stands (i.e., clearcuts). Precommercial thinning can increase compositional similarity of shrubs and herbs between young- and old-growth stands and increase the abundance of late-seral herbs (Lindh and Muir 2004). Conversely, thinning in 60- to 80-year-old stands can increase diversity and the abundance of early-seral species but have little effect on late-seral species (Puettmann et al. 2013). North et al. (1995) found that richness of herb and shrub species was greater in a green tree retention harvest than in an adjacent clearcut and in an intact 65-year stand (also see Halpern et al. 2005).

Retention patches of late-successional and old-growth forest scattered between late-successional forest reserves might serve as conservation centers for some understory vascular plants. Nelson and Halpern (2005) studied the short-term (1 and 2 years) response of understory plants to patterns of aggregated retention harvest and found that

old-forest patches larger than 2.5 ac (1 ha) helped retain populations of late-seral plant species, although disturbance within 33 ft (10 m) of the stand edge (“depth of edge” influence) was evident by the incursion of early-seral plant species. Late-seral herbs were more frequently extirpated in harvested portions of aggregated treatments as opposed to dispersed treatments

Fire exclusion may have affected understory communities in late-successional and old-growth forests in vegetation zones where fire was frequent historically (Loya and Jules 2008), but these effects are poorly understood and more research is needed. Some evidence suggests increases in shrub cover may be associated with fire exclusion (Loya and Jules 2008). Loss of bear grass (*Xerophyllum tenax*) in anthropogenically maintained savannahs has also been attributed to fire exclusion (Peter and Shebitz 2006). Studies from mixed-conifer forests in the eastern Washington Cascades and from ponderosa pine (*Pinus ponderosa*) forests suggest that a single application of prescribed fire can reduce shrub cover, but effects on understory response were minor and limited to slight increases in vegetation cover (Dodson et al. 2008) and diversity (Busse et al. 2000). Donato et al. (2009) found that almost all species in mature and old-growth stands were present following high-severity fire in the Biscuit Fire. There is some evidence that reintroduction of wildfire to dry forests of the Sierra Nevada had little effect on diversity of vascular plant species but may increase the distribution of species that may have been negatively affected by fire exclusion (Webster and Halpern 2010).

Invertebrates—

The ISSSSP provides a series of products on selected invertebrate species.²⁰ Range maps are provided on a number of mollusks species²¹ and conservation assessments and species fact sheets are provided for all sensitive invertebrates including bumblebees, beetles, true bugs, butterflies and

moths, damselflies and dragonflies, amphipods, and other taxa. The following sections summarize other recent studies on various invertebrate taxa including soil invertebrates, mollusks, and other insects and arthropods.

Soil invertebrates—Soil invertebrates constitute a wide array of taxa, including microorganisms, nematodes, mites, springtails, microspiders, centipedes, millipedes, earthworms, and others, with an equally diverse set of ecological functional roles as comminutors (chewers), detritivores and saprophages (detritus-feeders), fungivores (fungi-eaters), prey, predators, parasites, and more (Berg and Laskowski 2005). In addition, there are four functional groups of arthropods—litter- and soil-dwelling species, coarse wood chewers, understory and forest gap herbivores, and canopy herbivores—that are listed as Survey and Manage species in the Oregon and California Klamath, California Cascade, and California Coast Range physiographic provinces.

Some functional roles of soil invertebrates are yet to be discovered and may be surprising. As an example, Duhamel et al. (2013) reported that fungivorous springtails may play the ecological role of transferring secondary metabolites (catalpol) from host plants to arbuscular mycorrhizal fungi that then helps prevent the mycorrhizae from being grazed, thus protecting the host plant that uses the symbiotic fungi as a source of soil nutrient update.

Soil microorganisms include protozoa, rotifers, bacteria, and others that also play key ecological functions contributing to forest health, resilience, and productivity (Luo et al. 2016). Overall, soil invertebrates, including microorganisms, of late-successional and old-growth forests in the Pacific Northwest have been little studied but likely include species that are undescribed and some that closely associate with closed-canopy, late-successional, and old-growth forest conditions (see chapter 3 for a discussion of closed-canopy and open-canopy late-successional and old-growth forests and associated degree of shade-tolerant understory plants). Several older identification guides and species lists for the Pacific Northwest or selected sites therein are available such as on oribatid mites (Moldenke and Fichter 1988), arthropods (Parsons et al. 1991), and spiders (Moldenke et al. 1987).

²⁰ <http://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-invertebrates.shtml>.

²¹ <http://www.fs.fed.us/r6/sfpnw/issssp/planning-tools/species-distribution-maps.shtml#invertebrates>.

Forest invertebrate species vary in their sensitivity to disturbance and to forest canopy closure and tree density conditions. For example, Brand (2002) found that species richness, frequency, and density of epigeic (litter-dwelling) springtails in the Midwest are sensitive to fire; findings included the springtail *Tomocerus flavescens*, which is also a component of Pacific Northwest forests. Brand suggested using scattered refuges to maintain survival of fire-sensitive invertebrates in the face of fires; whether this is practical in the face of increasing rates of fire in the drier forests of the NWFP area is untested. Within the NWFP area, Moldenke and Fichter (1988) documented that oribatid mites, which are diverse and numerically dominant in ecosystems worldwide, occurred in very different species groups in open- and closed-canopy forests. The closed-canopy species were found throughout all stages of closed-canopy succession but differed markedly in relative abundance between young and old forests.

Milcu et al. (2006) found various responses of earthworms, springtails, and soil microorganisms—all decomposers—to grass and herb species diversity, functional group diversity, and growth form, but that the response of the decomposers did not correlate with plant productivity. Niwa and Peck (2002) found variable responses to time since prescribed fire burning in the occurrence and abundance of species of spiders (Araneae) and beetles (Carabidae) in southwest Oregon.

Some late-successional and old-growth-associated invertebrates, particularly dispersal-limited species, can serve as indicators of biodiversity and climate change risk (Ellis 2015, Homburg et al. 2014), and indicators of response of species groups to thinning (Yi and Moldenke 2005). Such potential indicator species include flightless terrestrial carabid beetles (e.g., Brumwell et al. 1998, Driscoll and Weir 2005, Eggers et al. 2010), which are a component of the entomofauna of LSRs in the Pacific Northwest, and flightless saproxylic weevils (Coleoptera: Curculionidae) (Buse 2012), which are also found in late-successional and old-growth reserves in the Pacific Northwest.

Yi and Moldenke (2005) studied the effects of thinning in 40- to 60-year-old Douglas-fir forest and associated

changes in forest floor environmental conditions in Willamette National Forest in the Oregon Cascade Range. They reported that Formicidae ants preferred heavy thinning intensities; Araneae spiders, Carabidae ground beetles, and Polydesmida millipedes positively correlated with litter moisture, which in turn was sensitive to season and negatively correlated with thinning intensity; and Gryllacrididae camel-cricket were negatively associated with litter moisture.

Rykken et al. (2007) found that invertebrates associated with headwater streams on the Willamette National Forest of Oregon, within the NWFP area, constitute spatially constrained but extremely species-rich communities. This occurred particularly within 3 ft (1 m) of the stream edge in mature forests and in 100-ft (30-m) riparian buffers within managed landscapes, such as may be provided as part of the Aquatic and Riparian Conservation Strategy of the NWFP (see chapter 7).

Caesar et al. (2005) assessed the genetic structure of a soil-inhabiting beetle (*Acrotrichis xanthocera*: Ptiliidae) in LSRs in the Klamath-Siskiyou ecoregion of northern California within the NWFP area. They concluded that the reserve system currently maintains high genetic variation for the species and suggested that intervening habitat gaps among the reserves should be reduced to maintain connectivity among the beetle populations within the reserves, apparently under the tacit assumption that the LSR system would continue in its current form under its current fire and fuels management approach (e.g., Fire suppression and limited use of prescribed fire) (see chapter 3).

Of concern for the conservation of invertebrates in LSRs is the incursion by invasive species. Of high concern are exotic earthworms (Ewing et al. 2015), e.g., *Lumbricus terrestris*, and about two dozen related species of Lumbricidae, introduced from Europe and currently found throughout North America including the Pacific Northwest although not yet in natural conifer stands. Exotic earthworms have been shown to reduce germination of tree seeds and survival of seedlings in southern Quebec, Canada (Drouin et al. 2014). In greenhouse and laboratory experiments, Eisenhauer and Scheu (2008) and Gundale (2002) found that exotic earthworms aid establishment of invader plants. In northern hardwood forests of Michigan

and New York, Gundale et al. (2005) and Eusenhauer and Sheu (2008), respectively, found that exotic earthworms reduce nitrogen retention. Hendirx and Bohlen (2002) reported that exotic earthworms, in general, can adversely affect soil processes and can introduce pathogens. In northern temperate forests of North America, Bohlen et al. (2004a, 2004b) reported that invasive exotic earthworms can adversely alter soil nutrient content (total carbon and phosphorus, and carbon-nitrogen ratios) and soil food webs. In the Northeastern United States, increased abundance of white-tailed deer (*Odocoileus virginianus*) can facilitate the spread of invasive earthworms; the deer's herbivory promotes the spread of nonnative understory vegetation that, in turn, fosters less acidic soils that are more favorable to the invasive earthworm species (Dávalos et al. 2015). Ziemba et al. (2015) reported that invasive *Amyntas* earthworms are altering detrital soil communities in North America, adversely affecting eastern red-backed salamanders (*Plethodon cinereus*) (also see Ransom 2012). Other invasive invertebrates occur throughout the Western United States (e.g., Chen and Seybold 2014) but are largely unstudied within the NWFP reserve system.

Few such studies of the specific impacts of exotic earthworms have been conducted in the NWFP region. Bailey et al. (2002) compared the influence of native and exotic earthworm species on soil and vegetation in remnant forests in the Willamette Valley of western Oregon. They discovered

two genera of native earthworms and the exotic earthworms were present in all five forest study sites, and reported that although there was no direct evidence that exotic species were affecting native species, they did not detect numerous native species presumed present in the region. They further suggested experimental studies to more clearly identify impacts of exotic species on native fauna.

Mollusks—Mollusks of the Pacific Northwest—particularly slugs and snails—have been the earlier focus of inventory, monitoring, and taxonomic study under the NWFP. Field guides have been produced under the Survey and Manage program for freshwater mollusks (Frest and Johannes 1999) and terrestrial mollusks (Kelley et al. 1999). Recent findings on mollusk species taxonomy are presented in table 6-4.

In a study in northern California, Dunk et al. (2004) sampled five species of terrestrial mollusks (*Ancotrema voyanum*, *Helminthoglypta talmadgei*, *Monadenia churchi*, *M. fidelis klamathica*, and *M. f. ochromphalus*). In a comparison across randomly selected sample plots of various forest conditions, they found that *A. voyanum* was associated with late-successional and old-growth forests, *M. churchi* was a habitat generalist, and data were insufficient to determine habitat associations of the remaining three species. As a result of the information on *M. churchi* being a habitat generalist, the species was removed from the Survey and Manage species list owing to not being specifically associated with late-successional and old-growth forests.

Table 6-4—Selected recent findings on taxonomy of mollusk species associated with late-successional and old-growth forest conditions within the Northwest Forest Plan area

Mollusk taxa	Topic	Source
<i>Fluminicola</i> (aquatic snail)	Updated taxonomy in northern California: <i>F. n. sp. 14 = F. potemicus</i> <i>F. n. sp. 15, 16, & 17 = F. multifarious</i> <i>F. n. sp. 18 = F. anserinus</i> <i>F. n. sp. 19 & 20 = F. umbilicatus</i>	Hershler et al. 2007
<i>Lyogyryus</i> (aquatic snail)	<i>Lyogyryus n. sp. 1</i> is now described as <i>Colligyryus greggi</i> . The genus was transferred and then the species fully described.	Hershler 1999, Liu et al. 2015
<i>Deroceras</i>	<i>Deroceras hesperium</i> is now considered <i>D. leave</i> , a common and widespread slug in North America	Roth et al. 2013
<i>Pristiloma</i>	<i>P. arcticum crateris</i> is now changed to <i>P. crateris</i>	Roth 2015

In a study of species in old-forest leaf islands (up to 1 ac or 0.4 ha) in western Oregon, Wessell (2005) found that leaf island size positively correlated with overall slug and snail density, and density of three mollusk species groups. Studies conducted beyond the Pacific Northwest also indicate that mollusk species richness is related to habitat area (Horsák et al. 2012) or hardwood forest stand age (Moning and Müller 2009). Other studies elsewhere (Moss and Hermanutz 2010) suggest the importance of monitoring for nonnative gastropods, particularly slugs, and especially along the margins of disturbed (burned) areas. Invasive slugs can negatively affect native slug diversity and plant regeneration and should be included in the monitoring of protected areas (Moss and Hermanutz 2010).

Jordan and Black (2015) provided a conservation assessment and finding of a potential threat for the mollusk *Cryptomastix devia*, otherwise known as the Puget Oregonian (Jordan and Black 2015). This terrestrial Gastropoda snail is strongly associated with large, old bigleaf maples (*Acer macrophyllum*) (fig. 6-2) growing among conifers, typically Douglas-fir, western hemlock, and western redcedar (*Thuja plicata*), or among other hardwoods such as black cottonwood (*Populus trichocarpa* ssp. *trichocarpa* and red alder (*Alnus rubra*). A primary threat identified in the conservation assessment for this species is reduction or loss of large, old bigleaf maples through suppression by Douglas-fir and other conifers, and through selective commercial thinning of hardwoods. Further, over the past 10 to 15 years,

Bruce G. Marcot



Figure 6-2—Large, old bigleaf maples are important hosts for the terrestrial gastropod snail known as the Puget Oregonian. Loss of maples in the southern Washington Cascade Range from commercial thinning of hardwoods, suppression by conifers, and disease may be threatening this snail.

there has been widespread mortality of bigleaf maples in the Upper Cowlitz and Cispus River drainages of the south Washington Cascades,²² the area with the vast majority of known locations of this mollusk. Other threats may include vertebrate and invertebrate predators (predatory snails, and beetles), and the occurrence of invasive slugs. As with nearly all Survey and Manage species, there is no ongoing research and monitoring program for this mollusk.

Other insects and arthropods—Studies of macromoths in H.J. Andrews Experimental Forest in Oregon (and in companion study sites in Mount Jirisan National Park in South Korea) revealed that most species consisted of the families Noctuidae and Geometridae, and that more than 3 years of sampling are needed to determine 90 percent of species richness (Choi and Miller 2013). Miller et al. (2003) studied 15 species of uncommon to rare moths (Noctuidae: Plusiinae) in the Cascade Range and identified three guilds of conifer, hardwood tree and shrub, and herbaceous-feeding species. They concluded that uncommon and rare species with special or restricted habitat requirements add to overall biodiversity, and that diverse environments such as meadows and early-successional vegetation contribute to the species' diversity.

A number of regional species identification guides have been recently produced on insects (Acorn and Sheldon 2001, Haggard and Haggard 2006), lepidoptera (Miller and Hammond 2003), macromoths (Miller and Hammond 2000), butterflies (Pyle 2002), bumblebees (Koch et al. 2012), dragonflies and damselflies (Kerst and Gordon 2011), and others, adding to previous identification keys such as on arboreal spiders (Moldenke et al. 1987).

Additional arthropods of late-successional and old-growth forests of the Pacific Northwest include millipedes and centipedes. Taxonomy, ecology, and habitat elements of many of these species remain poorly known. A study of riparian-associated millipedes in southwest Washington by Foster and Claeson (2011) revealed 15 species among 10 families, and that millipede species assemblages varied

between spring and fall and among sites. The authors urged further study on taxonomy and habitats and emphasized the importance of the ecological role of millipedes as detritivores in the forest ecosystem.

We did not find any recent autecology studies on centipedes in late-successional and old-growth forests of the Pacific Northwest. Insights into habitat associations of centipedes from studies elsewhere only suggest that some native species may be more closely associated with interior forest habitat conditions, and exotic species more associated with edge and disturbed habitat conditions (Hickerson et al. 2005).

A fairly recent management concern is the reduction in populations of pollinators on federal lands. In the Pacific Northwest, native bee populations are declining because of parasites, pathogens, pesticides, and invasive species (Spivak et al. 2011). Recent research has aimed at providing methods for restoring and conserving habitat for honeybee (*Apis* spp., nonnative species), bumblebees (*Bombus* spp.), and other pollinators (Decourtye et al. 2010, Wratten et al. 2012). The work highlights the economic benefit and ecosystem services provided by pollinating insects (Losey and Vaughan 2006).

Amphibians and reptiles—

Amphibians and reptiles of forest systems—Recent studies on the distribution, movement, and habitat relationships of forest amphibians and reptiles help assess how well the NWFP has served to provide habitat for late-successional and old-growth forest-associated amphibians and reptiles.

Old-growth forest-associated amphibians and reptiles are often described as secretive because most are fossorial and only seasonally encountered on the ground surface under down wood, rocks, leaves, and moss. Nonetheless, creative study approaches and modern landscape modeling and genetics techniques have furthered our understanding of the distribution, life history, and status of these species in the Pacific Northwest. For example, genetics were used to identify two new species: Scott Bar salamander (*Plethodon asupak*) and forest sharp-tailed snake (*Contia longicauda*). The Scott Bar salamander is associated with cool, moist forests and talus slopes

²² Kogut, T. Personal communication. Wildlife biologist (retired), P.O. Box 258, Packwood, WA 98361.

of an extremely restricted range in northern California; and, similar to other plethodontid salamanders (lungless salamanders of the family Plethodontidae), it is likely sensitive to changes in its habitat and in microclimate conditions (DeGross and Bury 2007, Mead et al. 2005). The forest sharp-tailed snake is associated with mesic and dense canopied forests of northern California and southwestern Oregon (Feldman and Hoyer 2010). Although little is known about this new species, the more wide-ranging species, *C. tenuis*, displays life history characteristics such as slow growth, late maturity, and low fecundity that are consistent with functional rarity and intrinsic vulnerability to population declines (Govindarajulu et al. 2011).

However, in an analysis of conservation risk of wildlife, including amphibians and reptiles, of Washington and Oregon, Lehmkuhl et al. (2001) determined that although life history traits can be used to qualitatively determine potential vulnerability, alone they are not an adequate quantitative predictor of vulnerability without additional information on species' habitat selection, habitat breadth, demography, and other factors. Clearly, additional survey and monitoring efforts and research are needed to better delineate species ranges and to improve our understanding of species' distributions, status, habitat requirements, and potential vulnerability (Gibbs 1998).

Another example of a furtive species yielding surprises upon study is the Shasta salamander (*Hydromantes shastae*) that, prior to 2012, was thought to be endemic to limestone outcroppings in the Shasta Lake watershed (Mooney 2010). Management decisions and land use project planning were based on this assumed association. Yet, Shasta salamanders subsequently have been found at sites lacking limestone outcrops and in an adjacent watershed (Lindstrand et al. 2012, Nauman and Olson 2004).

The Survey and Manage program and the ISSSSP have greatly increased scientific knowledge of forest-dwelling salamanders and their relationships with environments in the Pacific Northwest. Conservation Assessments for seven terrestrial salamanders (*Aneides flavipunctatus*, *Batrachoseps attenuatus*, *B. wrighti*, *Plethodon larselli*, *P. stormi*, *P. van-dykei*, and *P. asupak*) have been completed since 2005.²³ The

Density Management and Riparian Buffer Study of western Oregon (Anderson et al. 2007, Burton et al. 2016, Olson and Burton 2014) was initiated in 1994 when the NWFP implemented riparian buffers along non-fish-bearing streams (Cissel et al. 2006). This study assessed the long-term effects of thinning and differing riparian buffer widths on the distribution, abundance, and movement of terrestrial and aquatic amphibians. It showed that terrestrial salamanders are more abundant and more mobile in unthinned, densely canopied (with visible sky ranging 5 to 7 percent) (Anderson et al. 2007) riparian buffers as compared to nearby upland thinned forest (with visible sky ranging 9 to 12 percent) (Anderson et al. 2007; Kluber et al. 2008; Olson et al. 2014a, 2014b). This work highlights the importance of maintaining areas of dense canopies within at least 50 ft (15.2 m) of perennial streams in managed forests to serve as habitat for terrestrial species and as corridors for their dispersal (Olson et al. 2014b).

Recent studies have determined that the assumed dependence of terrestrial salamanders to late-successional and old-growth forests is more nuanced than previously thought (Bosakowski 1999). Some species have been found to be more abundant or occur more frequently in late-successional and old-growth forests (*B. attenuatus*, *Ensatina eschscholtzii*, Welsh and Hodgson 2013; *P. elongatus*, Welsh et al. 2008), but studies are inconsistent (e.g., *P. stormi*, Bull et al. 2006, Suzuki et al. 2008, Welsh et al. 2008). Relationships between plethodontid salamanders and cool, moist microhabitat conditions characteristic of late-successional and old-growth forests have been identified, but these conditions can be provided by other environmental features such as proximity to streams, talus slopes, and shading by vegetation, aspect, or topography (*P. stormi*, Suzuki et al. 2008; Welsh et al. 2007; *P. larselli*, Crisafulli et al. 2008). For species associated with forest floor attributes such as amount of down wood, a high volume of down wood may mediate the otherwise adverse effects of changes to the microhabitat conditions after thinning (*E. eschscholtzii*, Welsh et al. 2015; *E. eschscholtzii* and *P. vehiculum*, Kluber et al. 2008, Rundio and Olson 2007; *B. wrighti*, Clayton and Olson 2009).

²³ <http://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-amphibians.shtml>.

Substrate type and availability are important correlates of some species (*P. vehiculum*, Kluber et al. 2008; *P. larselli*, Crisafulli et al. 2008; *H. shastae*, Mooney 2010; *P. vandykei*, McIntyre et al. 2006, Olson and Crisafulli 2014), and down logs may provide thermal refugia for salamanders in thinned upslope zones (Kluber et al. 2009). In areas deficient in preferred substrates that provide cool refugia, salamanders may be associated with old-growth forest and related microhabitat conditions (e.g., *P. larselli*, Crisafulli et al. 2008). Maintaining cool and humid refugia may be central in protecting populations in areas that may experience disturbances.

Recent studies have started to unravel the complex trophic relationships of amphibians in forest and woodland food webs. Best and Welsh (2014) noted that plethodontid salamanders serve as predators of invertebrates, and as prey for carnivores. In an experiment, the authors found that *E. eschscholtzii* predation on diverse invertebrate prey suppressed some invertebrate taxa and released others resulting in increased retention of leaf litter and increased carbon sequestration, although such functions varied by the timing and amount of precipitation. We expect many other species of reptiles and amphibians play ecological roles influencing the trophic and nutrient dynamics of forest ecosystems to varying degrees, although quantitative studies are few and wanting.

Amphibians of stream systems—

The Aquatic Conservation Strategy under the NWFP provides for conservation and restoration of selected stream and riparian systems of the Pacific Northwest (Reeves et al. 2006). Although most protections are for fish-bearing streams, the importance of protecting non-fish bearing headwater streams and associated riparian environments has been increasingly recognized (Sedell and Froggatt 1984, Wilkins and Peterson 2000). First- and second-order headwater channels comprise the majority of stream miles (kilometers) within the Pacific Northwest and serve as cold water refugia in a warming climate (Isaak et al. 2016). In addition, they serve as headwater linkages between watersheds that provide important dispersal corridors for aquatic and terrestrial species, alike (Olson and Burnett 2009, Richardson and Neill 1998). Within stream networks,

amphibians occur from the headwaters to the alluvial flood plains, and their community structure is closely tied to channel types and within-channel attributes (Welsh and Hodgson 2011). For this reason, combined with their sensitivity to environmental perturbations, stream amphibians have been recognized as biometrics of stream health (Welsh and Hodgson 2008). In a study predating the NWFP, in the Oregon Cascades and Coast Range, Corn and Bury (1989) found that four species of aquatic amphibians—coastal giant salamander (*Dicamptodon tenebrosus*), Olympic salamander (*R. olympicus*), coastal tailed frog (*Ascaphus truei*), and Dunn's salamander (*P. dunni*)—had greater occurrence and abundance in streams flowing through uncut forests than in forests logged 14 to 40 years ago, and that tailed frogs and Olympic salamanders may be extirpated from headwaters in clearcuts. More recently, Olson et al. (2014a, 2014b) found that streambank amphibian species, including Dunn's salamander and western red-backed salamander, were sensitive to forest thinning within 20 to 49 ft (6 to 15 m) of small streams including headwaters.

Research has greatly advanced in the Pacific Northwest to incorporate amphibians to answer questions about the role of riparian buffers in the retention of aquatic ecosystem services. Studies have addressed how wide no-entry buffer zones should be to retain sensitive amphibian species or their critical habitat conditions and whether no-entry buffers are needed under differing upslope ecological forestry approaches. They are also asking whether or not it is appropriate to thin riparian zones to accelerate riparian restoration to achieve biodiversity goals in some cases. Using a meta-analysis to test the effectiveness of riparian buffers for conserving terrestrial fauna, Marczak et al. (2010) reported that riparian buffers, in general, maintain fewer amphibians compared to riparian forests in unharvested areas. Stream-associated amphibians such as coastal tailed frog and torrent salamanders are highly sensitive to timber harvest and the associated instream effects of increases in water temperature and sedimentation rates (Ashton et al. 2006, Bury 2008, Emel and Storfer 2015, Olson et al. 2007a, Pollet et al. 2010, Welsh and Hodgson 2011). Coastal giant salamanders appear to be more resilient to timber harvest practices than are other stream-associated

amphibians of the Pacific Northwest (Leuthold et al. 2012, Pollet et al. 2010). While the general findings that certain species are more sensitive to timber harvest than others holds relatively consistent in the literature, alternative findings in some areas (e.g., Raphael et al. 2002) highlight the importance of recognizing that abiotic and biotic factors interact with management actions in complex ways to influence the distribution and density of stream amphibians (Kroll et al. 2009).

The preponderance of evidence suggests that streams with riparian buffers within harvested landscapes do help to ameliorate some of the adverse effects of timber removal on instream conditions (Olson et al. 2014a, Pollet et al. 2010, Stoddard and Hayes 2005). In a replicated field experiment, Olson et al. (2014a) concluded that the NWFP Aquatic Conservation Strategy riparian buffers of ~230 to 476 ft (~70 to 145 m) seem to protect the headwater amphibian biota, and in a multiscaled survey, Stoddard and Hayes (2005) found that presence of a 150-ft (46-m)-wide riparian buffer predicted increased occurrence of tailed frogs and two species of torrent salamanders in a managed landscape. In the Oregon Coast Range, Kluber et al. (2008) found that ground surface attributes such as amount of rock or fine substrate determined the response of riparian and upland amphibian species to forest thinning along headwater streams, and that variable-width riparian buffer retention can provide for such microhabitat conditions. Olson and Burton (2014) examined the effects of alternative riparian management approaches including three no-harvest buffer treatments (site-potential tree-height (~230 ft [~70 m]), variable width with a 49-ft (15-m) minimum buffer on each side of the stream, and streamside retention (~20 ft [~6 m]) and a thin-through treatment whereby overstory tree density in a ~476-ft (~145-m) buffer treatment was reduced from 430 to 600 trees per acre (hectare) to ~150 trees per acre. They found that densities of torrent salamanders decreased along streams with the narrowest buffer, ~20 ft (~6 m), and that densities of Dunn's salamanders and coastal giant salamanders decreased in thin-through buffers (Olson and Burton 2014). The authors recommend the use of a 49-ft (15-m) or wider buffer to retain headwater stream amphibians (Olson and Burton 2014).

Riparian buffers can also maintain habitat connectivity and headwater linkage areas for dispersing amphibians. Using landscape genetics, Emel and Storfer (2015) found that gene flow among populations of the southern torrent salamander (*R. variegatus*) becomes restricted in streams with low canopy cover and high heat loads. The authors suggest that maintaining stream corridors with continuous riparian buffers may increase connectivity in managed landscapes. Coastal tailed frogs disperse more readily through intact forests than clearcuts (Wahbe et al. 2004) and, thus, may benefit from headwater management for connectivity across ridgelines, or the creation of "headwater linkage areas" (Olson and Burnett 2009). Olson and Burnett (2009) recommend linking headwater drainages across 7th-code hydrologic units to maintain landscape connectivity for headwater species. They propose extending buffers or alternative forest management practices that maintain canopy structure and shading to link neighboring watersheds over ridgelines (Olson and Burnett 2009, Olson et al. 2007).

Amphibians of still-water systems—In the Pacific Northwest, still-water (lentic) environments such as lakes, ponds, and wet meadows, and their adjacent riparian and terrestrial environments, also provide for species diversity. Many such sites do not provide habitat for fish, however, if they are impermanent. Lentic breeding amphibians such as red-legged frogs (*Rana aurora*), Oregon spotted frogs (*R. pretiosa*), Columbia spotted frogs (*R. luteiventris*), Cascades frogs (*R. cascadae*), northwestern salamanders (*Ambystoma gracile*), western toads (*Anaxyrus boreas*), and Pacific chorus frogs (*Pseudacris regilla*) flourish in these environments. Yet modifications and disturbances to these species' habitats (e.g., damming to increase water storage, ditching to increase drainage, siltation owing to timber harvest in the watershed, and overgrazing), combined with diseases and the spread of invasive species such as American bullfrogs (*Lithobates catesbeianus*) and brook trout (*Salvelinus fontinalis*), have resulted in some population declines (Adams 1999, Fisher and Shaffer 1996, Pearl et al. 2007).

The Oregon spotted frog is one of the most threatened amphibians in the Pacific Northwest; much information on this species can be found from the U.S. Fish and Wildlife Service Federal Register of notice of threatened status for

this species (USDI FWS 2014). Blouin et al. (2010) found that low connectivity between populations has resulted in decreased genetic diversity, and they recommended maintaining habitat connectivity and expanding the availability of appropriate wetlands.

Whereas palustrine wetlands and the amphibians that depend on them are not directly protected under the NWFP, riparian buffers likely benefit these species. In a literature survey of the use of wetland buffers by reptiles and amphibians, Semlitsch and Bodie (2003) found that core habitat of wetlands, as used by breeding populations, ranged from 522 to 951 ft (159 to 290 m) for amphibians and 417 to 948 ft (127 to 289 m) for reptiles from the edge of the aquatic site, and thus that adjacent terrestrial environments also were essential for population persistence.

Recent and ongoing issues for amphibians: fire and timber harvest—For most of the species discussed, effects of large disturbances such as wildfire and broad-scale timber harvest on populations, have not been adequately studied, and monitoring programs are not in place to assess the status of species and effectiveness of the NWFP in protecting these species. With recent decades of fire suppression in the Pacific Northwest, most of the forests have experienced much less fire than they did historically (see chapters 2 and 3). The effect of this change on habitats and populations of wildlife, including amphibians that require shaded, cool, moist microsites, is not known.

For the Larch Mountain salamander (*P. larselli*), Van Dyke's salamander (*P. vandykei*), California slender salamander (*B. attenuatus*), and Shasta salamander (*H. shastae*), a high proportion of known occurrences are within federal reserves associated with the NWFP where timber harvest is minimal compared to on nonreserve lands and thus these species are more protected, at least from timber harvest, under the NWFP. Species with a significant portion of their range on nonreserve land allocations (Siskiyou Mountains salamander (*P. stormi*) and Scott Bar salamander (*P. asupak*), Clayton et al. 2005, Nauman and Olson 2008; Oregon slender salamander (*B. wrighti*), Clayton and Olson 2009; black salamander (*A. flavipunctatus*), Olson 2008), however, might still be negatively affected by future land management activities if they deviate from the LSR

standards, and management of these lands may be crucial to protect these species. Furthermore, much of the range of many plethodontids is on nonfederal forest land (Suzuki and Olson 2008). Adaptive management areas (AMAs) and matrix lands were expected to function as experimental areas to address unresolved questions and to conduct long-term studies on timber harvesting and fire effects on forest salamanders and their response to these disturbances (Stankey et al. 2006, Suzuki and Olson 2008). However, the AMA program was never fully instituted under the NWFP, although AMAs were designated.

Recent and ongoing issues for amphibians: climate change—Climate change models for the Pacific Northwest region predict a warming of 0.2 to 1.1 °F (0.1 to 0.6 °C) per decade, with wetter autumns and winters, but drier summers (Mote and Salathe 2010). Significant warming is projected for the Pacific Northwest by the end of the 21st century, with simulation results projecting an increase in warming of 5.9 to 17.5 °F (3.3 to 9.7 °C) in Washington and Oregon and 2.7 to 8.1 °F (1.5 to 4.5 °C) in northern California, with generally increasing aridity but with higher uncertainty about specific precipitation levels (see chapter 2).

Changes in the climate can affect amphibians by altering habitat features such as vegetation, soil, and hydrology directly or via interactions with other threats such as timber harvest, wildfire, and disease (Blaustein et al. 2010). Plethodontid salamanders are particularly vulnerable to climate warming because they are specialized to cool microclimates, have limited dispersal capabilities, and may lack physiological tolerance to warm-induced stresses (Bernardo and Spotila 2006, Velo-Antón et al. 2013). Climate change may influence the distribution of suitable environments, which can result in fragmenting populations and shifting or retracting species ranges (Blaustein et al. 2010). More precipitation during the autumn and winter may lengthen the period of surface activity, but warmer and drier summers could result in fewer available surface and subsurface refugia that prevent desiccation. Forest management practices that retain canopy cover, and maintain or supplement surface refugia such as down wood and logs may help to ameliorate the effects of climate change (Shoo et al. 2011).

Recent and ongoing issues for amphibians: pests and pathogens—Minimal research has been conducted on the effects of nonnative animals or plants on amphibians in the Pacific Northwest, and the degree to which the NWFP influences establishment and spread of such invasives. Introduced American bullfrogs (*L. catesbeianus*) have become widely established in lentic environments in the Pacific Northwest and are considered an important predator of native pond-breeding amphibians. Most documented effects are from lowland areas (<0.2 mi [<240 m]) such as the Willamette Valley where Pearl et al. (2004) found greater effects of bullfrogs on Oregon spotted frogs compared to northern red-legged frogs. Breeding bullfrog populations have been documented from the Oregon Cascades (Garcia et al. 2009), but environmental conditions such as high UV-B levels (Garcia et al. 2015) may preclude extensive colonization.

As noted previously, the invasive Asian earthworm *Amyntas* that has invaded much of North America has been found to alter the behavior of eastern red-backed salamanders, but similar studies have not been conducted with salamander species found in the Pacific Northwest.

The threat of novel diseases entering populations of amphibians of the Pacific Northwest appears increasingly pressing. The deadly amphibian disease, chytridiomycosis, emerged in the 1970s and has become a prominent threat to amphibian biodiversity worldwide (Olson et al. 2013, Skerratt et al. 2007). Chytridiomycosis is caused by the fungus *Batrachochytrium dendrobatidis* (*Bd*) and has been identified as causing decline or extinction of over 200 amphibian species globally (Skerratt et al. 2007). It affects a broad host-range among amphibians, including salamanders, and has been reported in 516 (42 percent) of 1,240 amphibian species evaluated (Olson et al. 2013). *Bd* is widespread in the Pacific Northwest and has been found in northern red-legged frog, Columbia spotted frog, Oregon spotted frog, and Cascades frog (Pearl et al. 2007, 2009). Although the risk posed by *Bd* to these species is mostly unclear, evidence suggests that Cascades frogs at the southern extent of their range in northern California have experienced severe declines owing to chytridiomycosis (De León et al. 2016, Piovia-Scott et al. 2014, Pope et al. 2014).

More recently, in 2010, a second infectious chytrid pathogen, *B. salamandrivorans* (*Bsal*), emerged in Europe where it has decimated fire salamanders (*Salamandra salamandra*) (Martel et al. 2014). Unlike *Bd*, *Bsal* appears to affect only salamanders, not anurans (Van Rooij et al. 2015). As *Bsal* is not yet known to occur in North America, amphibian disease specialists are bracing for its arrival, which will probably occur through international pet trade routes (Grant et al. 2016, Gray et al. 2015, Martel et al. 2014). Models of amphibian habitat suitability predict that the Pacific Northwest is among the highest risk areas in North America (Richgels et al. 2016, Yap et al. 2015). The Pacific Northwest represents a hotspot for salamander biodiversity and contains numerous species from the two most *Bsal*-susceptible families, Plethodontidae and Salamandridae (Martel et al. 2014). In January of 2016, the U.S. Fish and Wildlife Service responded to the threat of *Bsal* by enacting a temporary rule restricting the importation of 201 species of salamanders for the pet trade (USFWS 2016). Although a critical step, the risk of spread of *Bsal* to North America, home to the world's richest salamander fauna, is still great and surveillance, research, and management actions are mandatory for successful mitigation of spread and response to this emerging infectious disease (Grant et al. 2016, Gray et al. 2015, Yap et al. 2015).

Recent and ongoing issues for amphibians: connectivity—The movement capabilities of most Pacific Northwest amphibians, especially plethodontid salamanders, are presumed to be limited, and fragmentation of their habitats may inhibit gene flow, and thus could isolate subpopulations. In forest watersheds with harvested upland areas, amphibians may disperse along riparian corridors that provide cooler, humid conditions (Olson et al. 2007a, 2014b). Stream buffers, riparian corridors, and down wood “chains” may provide connectivity for terrestrial species between their riparian and upland habitats, their adjacent stream and terrestrial habitats, and over ridges (Olson and Burnett 2009, 2013; Olson et al. 2014b). Providing linkage areas between adjacent watersheds and using various combinations of alternative management approaches (riparian buffers, thinning, down wood, leave islands, and uncut blocks) to retain forested areas along headwater ridgelines may facilitate upland dispersal and connectivity between subpopulations (Olson and Burnett 2009, 2013).

New methods for amphibians—Recently, environmental DNA has been applied to identifying the presence of endangered freshwater biodiversity including amphibians in lakes, ponds, and streams (Thomsen et al. 2012). Environmental DNA originates from cellular material shed by organisms (via skin, excrement, etc.) into aquatic environments that can be sampled, sequenced, and assigned back to the species of origin. Such methodology is important for the early detection of invasive species as well as the detection of rare and cryptic species. Environmental DNA approaches have been applied to imperiled aquatic amphibians in the Southeastern United States (McKee et al. 2015); an invasive salamander in Australia (Smart et al. 2015); and the trematode *Ribeiroia ondatrae*, a pathogenic parasite on North American amphibians (Huver et al. 2015). They have recently been applied to amphibians of aquatic systems in the NWFP area (e.g., Welsh and Cummings).²⁴

Birds—

Bird species pertinent to the development of the LSRs and late-successional and old-growth forests of the NWFP are covered elsewhere (see chapters 4 and 5). Many other bird species live in or migrate through the NWFP area, contributing to the region's biodiversity; here we address new science since the 10-year synthesis.

Recent studies have addressed various aspects of avian ecology in late-successional and old-growth forests, for example on pileated woodpecker (*Dryocopus pileatus*) (Aubry and Raley 2002a, Raley and Aubry 2006) and its role as a keystone species providing such ecosystem engineering functions as cavity excavation (Aubry and Raley 2002b). Other studies have provided information on forest habitat selection by white-headed woodpeckers (*Picoides albolarvatus*) (Lorenz et al. 2015), black-backed woodpeckers (*P. arcticus*) (Bonnot et al. 2009), and others. These studies suggest the role of disturbance (fire and forest pathogens) in providing habitat for these species.

For example, white-headed woodpeckers were found to establish home ranges within forest patches that had undergone recent disturbance including small patches

of prescribed burning and incidence of disease. Other studies have suggested that habitat associations of white-headed woodpeckers may differ by location and condition. Latif et al. (2015) found that the species in dry conifer forests on the east side of the Cascade Range in Oregon was closely associated with canopy openings adjacent to closed-canopy forests. In unburned, dry conifer forests of central Oregon, Hollenbeck et al. (2011) found that nest sites of white-headed woodpeckers were associated with low elevation, high density of large trees, low slope, and interspersed-juxtaposition of low- and high-canopy cover ponderosa pine patches. In postfire ponderosa pine forests of south-central Oregon, Wightman et al. (2010) found that low nest survival of white-headed woodpeckers was due to predation and was not correlated with habitat and abiotic features, and that survival in recently burned forests is likely enhanced with mosaic burn patterns with retention of larger, decayed snags.

Fontaine et al. (2009) studied response of bird communities to single fires (2 to 3 and 17 to 18 years following fire), repeated high-severity fire (2 to 3 years after fires repeated at 15-year intervals), and stand-replacement fire (>100 years following fire) in mature and old-growth mixed-evergreen forests of the Klamath-Siskiyou region of southwest Oregon. They reported that bird species richness (number of bird species) did not differ significantly among the various postfire conditions; that bird density was lowest 2 years after fire and highest 17 to 18 years after fire; and that repeat-fire conditions favored shrub-nesting and ground-foraging species, and unburned mature forests favored conifer-nesting and foliage-gleaning species. In that ecoregion, the researchers reported that bird communities were structured mostly by conditions of broadleaf hardwoods and shrubs, by extended periods of early-seral broadleaf dominance and short-interval high-severity fires.

Natural disturbance patterns, including fire, wind-throw, and other forces, can leave remnant and legacy elements of older forests such as old-forest patches, large-diameter green trees and snags, and large-diameter down wood, as key habitat elements for a variety of species. Subsequent regrowth of the vegetation through secondary succession then creates early-seral structures

²⁴ Unpublished data.

more complex than those resulting from final timber harvests such as clearcuts. The complexity of naturally regenerating vegetation following natural disturbances can often contain habitat for a wide variety of birds and other wildlife associated with early-successional environments (Marcot 1983, 1985; Swanson et al. 2011).

Vogeler et al. (2013) adopted the use of light detection and ranging to evaluate forest structure conditions for brown creepers (*Certhia americana*), an old-forest associate. In the Eastern United States, Poulin and Villard (2011) determined that brown creeper nest survival was greater in forest interiors at least 328 ft (100 m) from the forest edge, and that young forest conditions—plantations from previous management—in the managed matrix can adversely influence productivity of the species.

In a study of the influence of forest land ownership on focal species, McComb et al. (2007) projected habitat decreases for warbling vireo (*Vireo gilvus*) and increases for pileated woodpecker, as early-successional vegetation is projected to decline in area and older forests to increase. They also noted that land ownership patterns affected dispersion of habitat for both species, and that, for wide-ranging species, public lands provided habitat less available or not present on adjacent private lands.

On southeastern Vancouver Island, British Columbia, Canada, just outside the NWFP area, Hartwig et al. (2003) noted that pileated woodpeckers selected for largest diameter trees and for patches of older forests for nest cavities in coastal forests of Douglas-fir and western hemlock. They recommend retaining large live and dead grand fir, Douglas-fir, and red alder for pileated woodpecker nest sites, and to provide mature climax stands with greater proportions of bigleaf maple and grand fir.

On the Washington Pacific coast, Aubry and Raley (2002a) found that pileated woodpeckers selected Pacific silver fir (*Abies amabilis*) for nesting and western redcedar for roosting, and selected against western hemlock for both activities; that trees used for just nesting ranged from 26 to 61 inches (65 to 154 cm) diameter at breast height (d.b.h.), and trees used for just roosting ranged from 61 to 122 inches (155 to 309 cm) d.b.h.; that trees used for both

nesting and roosting were at least 90 ft (27.5 m) tall; and that trees less than 49 inches (125 cm) d.b.h. or less than 57 ft (17.5 m) tall were generally selected against. The authors suggested that habitat management for the species may need to also consider roost trees by providing snags and decadent live trees with heart-rot decay fungi. Raley and Aubry (2006) further reported that pileated woodpeckers in coastal forests foraged almost exclusively in closed-canopy stands, that they selected for relatively tall, large-diameter snags in early to moderate decay stages, and that they did not appear to use down logs for foraging because down logs did not support their primary prey of carpenter ants.

Great gray owls are considered late-successional and old-growth forest associated within the NWFP area and are still a Survey and Manage species. A conservation assessment was developed for the species in 2012, and the most recent survey protocol is from 2004, as posted on the ISSSSP site.

Carnivores—

This section summarizes new science associated with the taxa of mammalian carnivores that were either ranked as “not likely to remain well distributed” by the FEMAT or that had state or federal listing status. One species and one subspecies met both criteria (the fisher and the coastal population of the Pacific marten), and two species met the latter criterion only (the lynx [*Felis lynx*] and wolverine).

Taxonomic and listing updates of carnivores—Recent phylogenetic studies have changed the relationships of several of the species considered here. Within the Mustelidae, the fisher—formerly recognized as *Martes pennanti*—has been shown to be more closely related to the wolverine and tayra (*Eira barbara*) than to martens (*Martes* sp.) (Koepfli et al. 2008, Sato et al. 2012). Consequently, the fisher is now recognized as being the sole extant member of its own genus—*Pekania*—and is now recognized as *P. pennanti*. In April 2016, the U.S. Fish and Wildlife Service determined that the West Coast Distinct Population Segment of fisher does not warrant federal listing status. No other federal listing pertains to the species in the NWFP area.

Dawson and Cook (2012) evaluated phylogenetic and morphological evidence across the distribution of the American marten (*M. americana*) in North America and agreed with previous work (Carr and Hicks 1997) that there are two species of martens in North America. The American marten occurs east of the crest of the Rocky Mountains and across northern Alaska, and the Pacific marten (*M. caurina*) occurs west of the Rocky Mountains and includes the NWFP region. Dawson and Cook (2012) found significant genetic substructuring consistent with existing subspecific designations in the Pacific marten, and Slauson et al. (2009) agreed with these designations except in coastal California and coastal Oregon. Populations of the Pacific marten in coastal Oregon and coastal California appear to be a single evolutionary unit and have been proposed to be referred to collectively as the Humboldt marten (*M. c. humboldtensis*), a taxon formerly described as occurring only along the coast in northern California (Grinnell and Dixon 1926, Slauson et al. 2009). As of this writing, the U.S. Fish and Wildlife Service has determined that there is insufficient data to warrant federal listing of the Humboldt marten, but this decision is being appealed. The Humboldt marten is a state candidate species in California and is under a 1-year review for listing there.

The contiguous U.S. Distinct Population Segment of Canada lynx is currently listed as threatened in Washington, Oregon, and beyond²⁵ (USFWS 2000). On August 13, 2014, the Fish and Wildlife Service withdrew a proposal to list the North American wolverine in the contiguous United States as a threatened species under the Endangered Species Act; but then on April 4, 2016, the U.S. District Court for the District of Montana ordered U.S. Fish and Wildlife Service to reconsider whether to list the wolverine as a threatened species, and it currently sits as proposed threatened status.²⁶ A proposed rule for listing of wolverine will be finalized in the future (Further, recent information on life history of each species is available through Web sites noted above).

Current distribution of carnivores—In the NWFP area, the fisher continues to be restricted to a single native population in northwestern California and southwestern Oregon (Lofroth et al. 2010), a single small introduced population in the southern Oregon Cascades from source populations in Canada and Minnesota (Aubry and Lewis 2003), and recently reintroduced populations on the Olympic peninsula (Lewis 2014) and in the Washington Cascades (Lewis 2013).

The Pacific marten appears to remain fairly well distributed in the high-elevation forests of the inland mountain ranges but occupies less than 5 percent of its historical range in the coastal forests of California and less than 10 percent of its historical range in Oregon (Slauson et al., in press), and it is known from a single recent verifiable record from the Olympic peninsula.²⁷

In the NWFP area, lynx are known only from high-elevation (>4,600 ft [1400 m]) spruce (*Picea* spp.) and lodgepole pine (*Pinus contorta*) forest habitats of Washington in the northern Cascades, and from disjunct mountain ranges in the counties bordering Canada.

Verifiable wolverine occurrences in the NWFP area continue to be limited to high-elevation alpine and sub-alpine habitats in the central and northern Cascades of Washington (Aubry et al. 2007). However, recent detections in northeastern Oregon²⁸ and of a single male in the northern Sierra Nevada (Moriarty et al. 2009) suggest that wolverines have the potential to occur in the Cascades of Oregon but have yet to be detected there.

Habitat and feeding ecology of carnivores—We summarize here the new insights on habitat relationships of each species, and how these relate to each species' key life history needs for feeding, resting, and reproducing. Where possible, we link these new insights to management actions that may benefit each species. Our emphasis is primarily on martens and fishers, which occur across a larger proportion of the NWFP area than do lynx and wolverine.

²⁵ <http://ecos.fws.gov/ecp0/profile/speciesProfile?slId=3652>.

²⁶ <http://ecos.fws.gov/ecp0/profile/speciesProfile?slId=5123>.

²⁷ K. Aubry, personal communication, Emeritus scientist, Pacific Northwest Research Station, 3625 93rd Ave., Olympia, WA 98512.

²⁸ A. Magoun, personal communication, Wildland Research and Management, 3680 Non Road, Fairbanks, AK 99709.

Fisher—In the NWFP area, fisher home ranges vary from 2.9 to 24.5 mi² (7.4 to 63.5 km²) for males and 0.7 to 49.5 mi² (1.7 to 128.3 km²) for females (Lewis 2014, Lofroth et al. 2010), with home ranges decreasing in size from north to south. Overall, home ranges tend to be composed of mosaics of forest stand types and seral stages, but still include high proportions of mid- to late-seral forest (Raley et al. 2012). Notable exceptions exist in the coastal forests in northwestern California, where fishers occur both in areas with mosaics of young regenerating forest and residual mature forest (Matthews et al. 2013), as well as in areas with high proportions of young regenerating forest (Thompson et al. 2008). Higher use of young regenerating forest in these areas is related to the abundance of dusky-footed woodrats (*Neotoma fuscipes*), which are an important prey species (Lofroth et al. 2010) and which reach peak densities in young even-aged stands 5 to 20 years postharvest (Hamm and Diller 2009). Fishers typically avoid entering open areas devoid of or with significantly reduced overhead cover and escape cover, and exhibit avoidance by positioning home ranges to minimize overlap with large open areas (Raley et al. 2012) or by moving within home ranges to avoid such large open areas.

Fishers are dependent on large-diameter live and dead trees and downed logs with features such as cavities, platforms, and chambers as daily resting sites and seasonal den sites for females raising young (Raley et al. 2012). Female fishers are obligate cavity users for reproduction, with cavities in trees and snags providing secure environments for kits by regulating temperature extremes and providing protection from predators (Raley et al. 2012). These structures take hundreds of years to develop (Raley et al. 2012), emphasizing the critical importance of retaining these features for conservation of the species.

Rest structures for fishers primarily include large deformed or deteriorating live trees and secondarily include snags and logs (Raley et al. 2012, Weir et al. 2012). The species of trees and logs used for resting appear less important than the presence of a suitable microstructure, such as a cavity or platform. A meta-analysis by Aubry et al. (2013) of resting site selection by fishers across the NWFP area and adjacent areas found that rest site characteristics selected by

fishers were consistent across the Pacific States and British Columbia, and occurred on steeper slopes, in cooler micro-environments, with denser overhead cover, greater volumes of logs, and a greater prevalence of large-diameter trees and snags than were generally available. This meta-analysis provides managers with empirical support for managing for these conditions where maintenance or restoration of resting habitat conditions is an objective, even in locations where fishers have not been studied. Cavities used by female fishers for birthing and preweaning periods of kits were most often created by heart rot in older trees, and most often in the largest diameter trees available (Raley et al. 2012). During the postweaning period, female fishers continue to use tree cavities but also make use of downed logs.

Truex and Zielinski (2013) studied the influence of fuel treatments in forests of the Sierra Nevada, California, and found that mechanical treatments and fire treatments together, but not separately, reduced resting sites for fishers. Also, late-season burns adversely affected resting sites, but early-season burns did not. The authors suggested evaluating the effects of fuel treatment activities at various scales—fisher resting sites, home range, and landscape—to best balance fuel reduction with restoration and maintenance of fisher habitat.

Marten—In the NWFP area, martens occur in two distinct forest regions: (1) high-elevation forests of the inland mountain ranges, such as the Cascades of California, Oregon, and Washington and the Marble-Salmon-Trinity Mountains of California, that receive significant snowfall and (2) low- to mid-elevation coastal forests in Oregon and California that receive little snowfall. In mesic west-slope forest types in inland mountain ranges, martens select riparian areas and avoid landscapes with large patches of young forest; in contrast, in xeric east-side forests in inland mountain ranges, martens select sites with higher tree canopy cover and avoid fragmented forests (Shirk et al. 2014).

Pacific martens have recently been located in forests of the central and south coast regions of coastal Oregon (Moriarty et al. 2016a). In coastal California and Oregon, they occur in three environmental conditions: productive soil types where Douglas-fir, hemlock, and redwood forest associations occur; and two restricted environments on

low-productivity soils, including near-coast serpentine forest environments restricted to south coast Oregon and north coast California, and shore pine (*Pinus contorta* var. *contorta*) forest associations found in stabilized dunes in central coastal Oregon (Slauson et al. 2007, in press). Although the tree composition and vegetation structure vary among these three environmental conditions, they have one feature in common: a dense (more than 60 percent cover) shrub understory dominated by ericaceous species such as salal (*Gaultheria shallon*) or evergreen huckleberry (*Vaccinium ovatum*) (Slauson et al. 2007). In coastal forests at the home-range scale, martens select for large patches (more than 250 ac or 100 ha) of old-growth and late-mature forest or serpentine areas (Slauson et al. 2007).

Complex vertical and horizontal forest structure is an important characteristic of marten habitat, providing features used to meet daily resting and seasonal denning needs as well as providing food and cover for prey species. Just outside the NWFP area in north-central California, Moriarty et al. (2016b) found that martens strongly selected complex-structured forest stands over simple stands and openings. Stand structure influenced martens' movement patterns, being quicker and more erratic and linear in simple stands and openings, likely related to predator behavior; and more sinuous and slower in complex stands, likely related to foraging under cover from predators.

In coastal California, the summer and fall diet of martens is dominated by sciurid and cricetid rodents (chipmunks [*Tamias* sp.], Douglas squirrels [*Tamiasciurus douglasii*], and red-backed voles [*Myodes californicus*]) and medium-sized birds. During winter, martens shift to larger bodied prey species such as medium-sized birds and northern flying squirrels (*Glaucomys sabrinus*) (Slauson et al., in press). Many of these key prey species of the Humboldt marten reach their highest densities in forest stands with mature and late-successional structural features (e.g., Carey 1991, Carey and Johnson 1995, Hayes and Cross 1987, Rosenberg et al. 1994, Waters and Zabel 1995) where their key food resources—conifer seed crops, truffles (the fruiting bodies of ectomycorrhizal fungi), and late-successional and old-growth-associated pendant lichens—typically reach their greatest abundances (Luoma et al. 2003, Smith et al.

2002). Dense ericaceous shrub layers are also positively associated with key prey species including chipmunks (Hayes et al. 1995), northern flying squirrels (Carey 1995), and small mammals in general (Carey and Johnson 1995). Coarse woody debris also has been shown to positively influence foraging efficiency for martens by making prey more vulnerable to capture; where it has been reduced from timber harvest, marten foraging efficiency declines (Andruskiw et al. 2008).

Like fishers, martens also depend on large decadent woody structures including live and dead trees and downed logs with features such as cavities, platforms, and chambers for daily use as resting sites and as seasonal den sites for females raising young (Raphael and Jones 1997, Slauson and Zielinski 2009, Thompson et al. 2012). In summer and fall, martens in coastal forests rest predominantly in large-diameter snags, downed logs, and live trees, and less commonly in slash piles, boulder piles, and shrub clumps (Slauson and Zielinski 2009). Use of the sizes of trees, snags, and downed logs is similar between coastal and inland mesic forests, but tree diameters are larger than reported for inland xeric forest (Raphael and Jones 1997). Female martens are also obligate cavity users for reproduction, with cavities in trees and snags providing secure environments for kits by regulating temperature extremes and providing protection from predators (Raphael and Jones 1997, Slauson and Zielinski 2009, Thompson et al. 2012). Reproductive den structure types are more restricted than are rest sites (Ruggiero et al. 1998), and all natal dens have occurred in aerial cavities in large-diameter hardwoods, and maternal dens have included cavities and platforms in both large-diameter hardwoods and broken top live and dead conifers (Slauson and Zielinski 2009; Slauson, in press). In contrast to inland forest types, the dens found to date in coastal forests have been predominantly in hardwoods.

Lynx—In northern Washington, Koehler et al. (2008) found that lynx exhibited the strongest selection for Engelmann spruce (*Picea engelmannii*) and subalpine fir (*Abies lasiocarpa*) forest on moderate slopes, and to a lesser degree, forests with moderate canopy cover, at elevations of 5,000 to 6,000 ft (1525 to 1829 m), and either avoided, or used proportional to their availability, the following cover types:

lower elevation Douglas-fir, ponderosa pine forests, forest openings, and recently burned areas with sparse canopy and understory cover. Lynx in Washington spend more time hunting in patches of Engelmann spruce and subalpine fir where their predominant prey, snowshoe hare (*Lepus americanus*), occur at their highest densities (Maletzke et al. 2008). Lynx den sites typically occur in sheltered spaces most often created by downed logs, root wads of fallen trees and less often by boulder fields, slash piles, and live trees (Moen et al. 2008, Squires et al. 2008). In Montana, lynx den sites typically occurred in mature and mid-seral conifer stands with abundant woody debris, located in drainages or drainage-like basins (Squires et al. 2008).

Wolverine—The combination of historical records (Aubry et al. 2007) and contemporary studies of wolverines (Copeland et al. 2007, 2010) reveals that, during the denning and nondenning periods of the year, the species is closely associated with persistent spring (April to May) snow cover, and that this represents an obligate association with this bioclimatic feature. Wolverine habitat selection is strongly influenced by the distribution of seasonal prey resources, which tend to be dominated by large ungulates (e.g., moose, deer, elk, and caribou) (Copeland 1996, Lofroth et al. 2007) and large rodents (e.g., beaver, hoary marmots, and porcupine) (Lofroth et al. 2007).

Female wolverines have been shown to be more sensitive than males to predation risk and human disturbance. Females make greater use of rugged terrain to avoid predators and avoid areas with winter ski recreation and higher road densities within their home ranges (Krebs et al. 2007). Wolverine den sites tend to occur in areas with persistent spring snow cover and with large boulders and downed trees where females find secure sites for giving birth to and raising kits (Dawson et al. 2010, Magoun and Copeland 1998).

Landscape habitat suitability and connectivity for carnivores—Studies and habitat modeling of carnivores conducted at the landscape scale primarily address issues related to population processes, such as dispersal, gene flow, habitat connectivity, and population persistence. Regional landscape habitat models have been developed for the fisher

in various portions of the NWFP area (Carroll et al. 1999, Davis et al. 2007, Halsey et al. 2015, Lewis 2013, Zielinski et al. 2010). Collectively, the most important predictors of suitable fisher habitat measured at various scales include areas with higher annual precipitation, moderate to dense tree canopy cover, higher basal area and diameter classes of hardwoods and conifers, and topographically complex areas at mid to low elevations.

The southern portion of the NWFP area supports the largest extant fisher population in the Pacific Northwest. Modeled suitable habitat for fishers in the southern portion is projected to be well distributed and well connected, and the habitat models provide managers with a tool for planning activities and for monitoring habitat changes.

In the coastal forests of California and Oregon, Slauson et al. (n.d.)²⁹ modeled landscape habitat suitability for the Humboldt marten where the most important predictors, at various scales, were old-growth forest structure, amount of serpentine environment, and areas of higher precipitation, all positively correlated with marten detection locations versus nondetection locations. This model predicted that less than 15 percent of the historical habitat of this subspecies in the NWFP area is currently suitable. In the southeastern portion of the NWFP area, Kirk and Zielinski (2009) modeled landscape habitat suitability for martens in inland forest areas and found that marten occurrence was best predicted by a large amount of the biggest size classes of mesic forest types, by landscape configuration (more habitat patches), and by more habitat in public land management.

Moriarty et al. (2015) used new methods for evaluating functional connectivity of Pacific martens across fragmented forest landscapes: use of food incentive experiments, and nonincentive locations of martens collected with global positioning system telemetry. They found that Pacific martens selected complex stands and crossed open areas in the winter when enticed by bait, and without bait they avoided openings and simple-structured forest

²⁹ Slauson, K.M.; Zielinski, W.J.; Laplante, D.; Kirk, T. [N.d.]. Landscape habitat suitability model for the Humboldt marten (*Martes caurina humboldtensis*). Manuscript in preparation. On file with: Keith Slauson, U.S. Forest Service, Pacific Southwest Research Station, 1700 Bayview Drive, Arcata, CA 95521. kslauson@fs.fed.us.

stands. Pacific martens also selected complex stands during summer and winter alike. Baiting can induce movements and introduce bias in habitat selection analysis intended to represent bait-free conditions. Dispersal and connectivity of Pacific martens thus seemed to be afforded partially by snow cover across open areas, and more fully by networks of complexly structured forest stands during summer.

Koehler et al. (2008) developed a model of lynx habitat suitability from snow tracking and estimated that 1,467 mi² (3800 km²) of habitat suitable for lynx is present in the NWFP region, far less than prior estimates based on the correlation of historical records to vegetation types (McKelvey et al. 2000). Most lynx habitat identified by Koehler et al. (2008) occurs in western Okanogan County, Washington, and it was estimated that this amount of habitat could support a population of 87 lynx. Murray et al. (2008) assessed the state of lynx research and conservation in the species' southern range, including the NWFP area, and concluded based on their review of the existing science that successful lynx conservation will require protection and management of large tracts of lynx and snowshoe hare habitat, and ensuring connectivity between lynx populations at the core and periphery of the species' range.

The global distribution of wolverines is closely associated with areas of persistent spring snow cover (Copeland et al. 2010). In the NWFP area, the most significant areas with persistent spring snow cover are high elevations of the north-central Washington Cascades, where most records of wolverines occur. Although smaller areas with these conditions occur in the Oregon and California Cascades, limited connectivity to areas known to be occupied by wolverines may impede the ability for wolverines to reoccupy these areas. Other than models of areas with persistent spring snow cover across the NWFP area (Copeland et al. 2010), no additional habitat suitability models have been developed for wolverine. Similar to the lynx, ensuring connectivity between the largest areas of suitable habitat for the wolverine in the NWFP area with areas occupied by wolverines in British Columbia will be one of the most critical requirements to help ensure wolverine conservation in the NWFP area.

Landscape-scale habitat connectivity is a significant requirement for the carnivores considered here. For lynx and wolverine, connectivity to population source areas outside the NWFP is essential for population persistence (Aubry et al. 2007, Murray et al. 2008, Schwartz et al. 2009). For the fisher in the Washington Cascades (Lewis and Hayes 2004), and for lynx (Koehler et al. 2008), a current lack of connectivity likely is a main factor inhibiting the recolonization of suitable habitat in the northern portions of the NWFP area. For species that occupy high-elevation areas, particularly in the Cascades (i.e., lynx, wolverine, marten), climate change will likely reduce the amount and connectivity of habitat (Lawler and Safford 2012, McKelvey et al. 2011). In the coastal forests of northwestern California, the lack of connectivity of habitat for the Pacific marten is likely inhibiting the recolonization of suitable habitat patches within their dispersal capability (Slauson, in press).

In general, current and future problems of habitat connectivity for carnivores include wide distances among patches of suitable habitat, low quality of intervening habitat, and adverse effects of natural and anthropogenic movement filters and barriers such as rivers and roads on dispersal behavior and survival (Crooks et al. 2011). Owing to the high cost of creating habitat corridors and enhancing connectivity such as by constructing wildlife crossings on highways, or the long time necessary for regenerating late-successional forests where they have been lost, connectivity enhancements that can benefit multiple species (e.g., Singleton et al. 2002) could be most efficient at achieving objectives for carnivore conservation.

New threats to carnivores: predation—Studies in the NWFP area, as well as studies from adjacent areas in the Pacific States (Bull et al. 2001), have identified the bobcat (*Lynx rufus*) as a major predator of fishers (Wengert et al. 2014) and martens (Slauson et al., in press). Whereas predation is a natural mortality factor, its magnitude may have increased from historical times owing to the increase in early- to mid-seral forests in the NWFP area, which provide conditions suitable for predators such as bobcats and coyotes (*Canis latrans*) that have general habitat needs. In the NWFP area, studies of bobcat habitat use and diet in

coastal forests indicate that they select for young regenerating stands (Slausonet al., in press; Wengert 2013) where they find key prey species such as lagomorphs, woodrats, and mountain beavers (*Apodonta rufa*) (Knick et al. 1984; Slauson et al., in press). Halsey et al. (2015) modeled bobcat habitat to account for predation risk in evaluating reintroduction locations for fishers in the southern Washington Cascades, and found that reintroduction sites at middle to higher elevations rather than at lower elevation would likely reduce risk of predation by bobcats.

New threats to carnivores: rodenticides—The widespread use of rodenticides and other toxic chemicals found at illegal marijuana-growing operations has emerged as a major new threat. In California, most [dead] fishers tested for toxicant exposure tested positive for one or more anticoagulant rodenticides, which were ingested when the fishers consumed contaminated rodent prey. Several fishers recently have been confirmed to have died from acute rodenticide poisoning on the Hoopa Valley Indian Reservation located in the southern portion of the NWFP area (Gabriel et al. 2012, 2015). Exposure rates of Humboldt martens to rodenticides is less well understood, but in California, rodenticide exposure has been confirmed in one of six (17 percent) [dead] martens tested (Slauson et al., in press). This threat may be limited to the southern portion of the NWFP area where cultivation of marijuana is illegal on federal land but common, and where most suspected use of rodenticides is at such illegal growing sites on public, tribal, and private land alike.

New threats to carnivores: wildfire and fuels treatments—In 2007, the Fisher Conservation Strategy Biology Team held an expert panel workshop to conduct a threats analysis on fishers within the West Coast Distinct Population Segment and British Columbia portions of the species' range in the Pacific Northwest. The team evaluated 20 types of potential threats and ranked four equally as the greatest threats: uncharacteristically severe wildfire, forest canopy and overstory reduction, reduction of forest structural elements, and forest habitat fragmentation (Marcot et al. 2012a).

Wildfire is a natural process that can alter environments in various ways, including leaving behind remnants

of older forest stands and large wood components that can serve as legacy structures as the vegetation subsequently regenerates. However, major loss or degradation of forests, especially from extensive high-intensity wildfire, can significantly affect habitat for the carnivores considered here. Effects of wildfire will be the most severe where fire suppression has resulted in the unnatural buildup of fuels. Severe wildfire occurrence is primarily in the southern portion of the NWFP region and at lowest elevations, putting the fisher at greater risk to loss of habitat than the other carnivores considered here. These are also the areas where fuels treatments such as mechanical thinning, which can potentially negatively affect fisher habitat, are most commonly prescribed. High-severity wildfire can have positive effects on some prey species, particularly herbivorous species such as Leporids (*Lepus americanus* and *Sylvilagus bachmani*) and dusky-footed woodrats (*Neotoma fuscipes*), by creating early-seral conditions that favor these populations. The degree to which high-severity wildfire can benefit the carnivores considered here depends on their reliance on these species as important components of their diets and the degree to which fires result in landscape patterns (e.g., mixed severity in the range of the fisher) compatible with supporting the rest of their life history needs.

The effects of fire suppression on carnivore forest habitat in the Pacific Northwest have received little study. Fire suppression activities often result in denser forests with more canopy layers and more dead wood from shade-tolerant trees. Whether these outcomes increase habitat quality or amount for forest carnivores is unstudied.

Initial research on the effects of fuels treatments on fishers outside the NWFP area suggests that treatments have significant indirect benefits by protecting fisher habitat from wildfire (Scheller et al. 2011) and that some types of fuels treatments may be compatible with fisher occupancy as long as they do not exceed a particular rate and extent (Sweitzer et al. 2016a, Zielinski et al. 2013) and they do not affect the density of large structures (logs, snags, live trees) used by fishers for foraging, resting, and denning.

Loss of habitat from wildfire was ranked as one of the top threats to the Humboldt marten in coastal California (Slauson et al., in press). In the last two decades, several

fires have burned in marten-occupied areas with unknown effects on the population. Fuels treatments are typically used to minimize the severity of wildfire in these situations, but to avoid reducing marten habitat, they could be carefully planned and monitored. Fuels treatments that reduce surface and ladder fuels produce stand structures that martens either avoid or travel through quickly (Moriarty et al. 2016), similar to how marten respond to open linear features such as ski runs (Slauson et al 2017) and seismic lines (Tigner et al. 2015). Therefore to minimize impacts to martens from fuels treatments, they could be carefully designed to minimize their overlap with habitats supporting important aspects of marten life history such as denning and have a spatial arrangement that most likely represents a tradeoff between affecting fire behavior and minimizing negative effects on connectivity and foraging for martens (Moriarty et al. 2015, 2016b). Since 1985, fires have burned more than 386 mi² (1000 km²) of forest habitat in Okanogan County, the only region in Washington where lynx has been verified recently; this loss of forest habitat from fire, coupled with the naturally fragmented distribution of suitable habitat, represent a present, significant threat for lynx conservation where they persist in the NWFP area (Koehler et al. 2008).

The effect of wildfire and fuels treatments on wolverines has not been directly studied.

New monitoring methods for carnivores—New developments in methods for monitoring carnivores include methods of detection, improvement of statistical survey designs, and advances in statistical analyses of carnivore survey data. Once the objectives of monitoring are identified, a suite of survey designs can be considered (Long and Zielinski 2008, O’Connell et al. 2011). Statistical advances in occupancy modeling (MacKenzie et al. 2006, Royle et al. 2008, Slauson et al. 2012) and spatially explicit capture-recapture analysis (Royle et al. 2014) provide two new statistical frameworks to conduct carnivore monitoring. Because of the importance of designing surveys capable of reliably detecting a species, detecting magnitudes of change as specified in the monitoring objective, and optimizing the costs of such designs, Ellis et al. (2014) developed a power-analysis tool to evaluate the effects of alternative design elements for monitoring wolverines that has also been evaluated for fishers (Tucker 2013).

Monitoring population status via sampling DNA from non-invasively collected hair samples is an increasingly realistic goal for carnivores (Kendall and McKelvey 2008, Pauli et al. 2011, Schwartz and Monfort 2008). Recent advances also have been made in the use of camera-trap sampling of carnivores (Mcfadden-Hiller and Hiller 2015), including fishers (Sweitzer et al. 2016b), and mountain lions (*Puma concolor*) (Caruso et al. 2015), and carnivores elsewhere (e.g., Herrera et al. 2016, LaPoint et al. 2013) including arboreal situations (Cotsell and Vernes 2016).

Red tree vole and *Arborimus* allies—

Since the 2006 synthesis (Haynes et al. 2006), much work has taken place on the morphology, genetics, and taxonomy of red tree voles and related species (Bellinger et al. 2005; Miller et al. 2006, 2010). Three species of *Arborimus* forest voles are now recognized to occur within the NWFP area. The white-footed vole (*A. albipes*) occurs in the Coast Range, Willamette Valley, and a portion of the western Cascades, from northwestern California north through Oregon to the Columbia River border with Washington. The red tree vole (*A. longicaudus*) occupies a subset of the range of the white-footed vole but does not extend as far into California, is absent in extreme northwest Oregon, and extends into the Cascades east of Portland, still constrained by the Columbia River to the north. The Sonoma tree vole (or California red tree mouse; *A. pomio*) is found only in coastal northern California, extending northward to where the red tree vole’s range ends. Red tree voles and Sonoma tree voles are arboreal, making nests and using movement runways in canopies of late-successional and old-growth forests; white-footed voles, however, are only partially arboreal.

Red tree voles and Sonoma tree voles are particularly ecologically interesting for their arboreal life history and their reliance on needles and twigs of conifers as food sources. Chinnici et al. (2012) documented association of Sonoma tree voles with unharvested and partially harvested old-growth Douglas-fir forest in California. Genetics analyses by Blois and Arbogast (2006) revealed that, although Sonoma tree voles consist of one panmictic (interbreeding) population, some genetic evidence suggests that southern Sonoma tree voles may occur as a distinct management unit within the species.

Swingle and Forsman (2009) found that red tree voles occasionally moved short distances across small forest openings or across small logging roads less than 82 ft (25 m) wide. However, they saw no evidence of tree voles moving across large nonforest areas and thereby speculated that such areas are barriers to dispersal, and that long-term persistence of red tree voles depends on size and connectivity of forest cover and structure of the forest canopy.

Dunk and Hawley (2009) recognized threats to red tree vole populations, including fire and recent historical conversion of forests to nonforest uses. They developed habitat association models, and their most statistically significant model identified the best habitat correlates as site percentage of slope, basal area of trees 18 to 35 inches (45 to 90 cm) d.b.h., maximum tree height, and variation in diameters of conifers. They applied the model to the distribution of the species and determined that reserves have significantly higher quality habitat than do nonreserved lands.

Studies by Wilson and Forsman (2013) noted that silvicultural thinning of young forest stands to promote late-seral forest conditions can provide habitat for a number of vertebrate species, but thinning can delay the development of a mid-story layer of trees, which is an important habitat component for red tree voles (and also of northern flying squirrels, *Glaucomys sabrinus*). The authors suggested that long-term forest and habitat management goals should include developing more structurally and biologically complex forests across the landscape at scales appropriate to the species' ecologies.

Other recent research on red tree voles includes determining their distribution and abundance based on occurrence in pellets regurgitated by a principal predator, northern spotted owls (Forsman et al. 2004). Working in Oregon, the authors found that the incidence of red tree voles in the owl pellets occurred most commonly in the central and south coastal regions, were relatively sparse in the northern Coast Range and Cascades areas of the state, and were absent from pellets on the east slope of the Cascade Mountains and most of the dry forest area in south central Oregon. Incidence also declined by elevation in the Cascades, being most common in owl pellets below 3,200 ft (975 m) and rare above 4,000 ft (1220 m) elevation. The authors also speculated that red tree vole populations have declined in landscapes dominated

by young forests and with increased occurrence of logging, fire, and human development. Their finding of low incidence of red tree voles in spotted owl pellets along the north coast area of Oregon was also supported by direct surveys there in Tillamook and Clatsop State Forests by Price et al. (2015) who found 33 tree vole nests at only 4 out of 86 randomly selected sites. They speculated that the dearth of nests there is because of logging or burning in the early 1900s and subsequent intensive forest management, and suggested that remnant old-forest stands on BLM and state lands serve as source populations.

Rosenberg et al. (2016) presented models of red tree vole habitat, comparing and combining their best performing model with previously published models. They concluded that the ensemble combination of models provided the most accurate predictions of red tree vole occurrence and habitat suitability, and that it could be used to reduce survey costs and to guide management decisions and habitat conservation strategies for the species.

Swingle et al. (2010) discovered that a key predator of red tree voles in western Oregon is weasels (*Mustela* spp.). Activity behavior and dispersal timing of red tree voles were studied by Forsman et al. (2009) using video cameras at nest sites, and additional studies of red tree vole home ranges and activity patterns were conducted by Swingle and Forsman (2009). At present, a major review has been completed on the distribution, habitat, and diet of red tree voles and Sonoma tree voles (Forsman et al. 2016), and an annotated bibliography has been compiled on all three *Arborimus* species discussed here (Swingle and Forsman 2016).

The ISSSSP's 2016 Red Tree Vole High Priority Site Management Recommendations³⁰ provides guidance to field personnel for identifying land use allocations consistent with red tree vole conservation, high-priority sites outside of those allocations, and connectivity areas for linking sites, as well as nonpriority sites and key information gaps.

On the International Union for Conservation of Nature and Natural Resources Red List, although relatively rare, white-footed voles are listed as of "least concern," whereas red tree voles and Sonoma tree voles are both listed as "near

³⁰ <http://www.blm.gov/or/plans/surveyandmanage/files/RTV-HPS-MR-201604-final.pdf>.

threatened” because of their restriction to mature forests—old-growth forests for the red tree vole—and from threats of logging and fragmentation of their forest habitats. In 2009, the U.S. Fish and Wildlife Service evaluated whether red tree voles in the North Coast Range of Oregon—sometimes referred to as “dusky tree voles” that seem to differ in diet and other aspects from other red tree voles—deserve designation as a distinct population segment under the Endangered Species Act. The Fish and Wildlife Service convened a panel of species experts and managers to evaluate the evidence (Marcot and Livingston 2009), and ultimately announced that the north Oregon Coast Range population warrants listing as a distinct population segment but is precluded from development of a proposed listing rule because of higher priority actions (USFWS 2011). Morphological study of the dusky tree vole (putatively *A. longicaudus silvicola*) did not find evidence to support the proposed, separate subspecies designation (Miller et al. 2010). The current listing status of red tree vole is Continuing Candidate; the Fish and Wildlife Service’s species assessment³¹ also provides a summary of recent biological and ecological information on the species.

Bats—

At least 14 species of bats are found within the NWFP area. Some of these species are late-successional and old-growth associated, and their forest habitats are likely provided by NWFP reserve guidelines, although studies are incomplete.

Kroll et al. (2012) reviewed the state of knowledge on birds and bats dependent on cavities and snags (dead standing trees) in the Pacific Northwest. They concluded that relatively little is known about how the viability of populations of these species is affected by distributions of snags and by the proximity and amount of mature and late-successional forests across the landscape. Rodhouse et al. (2012) provided models of the distributions of little brown bats (*Myotis lucifugus*) in the Northwestern United States, finding a link of the species with forest cover and productivity.

Lacki et al. (2012) studied use of snags for roosting by long-legged myotis (*M. volans*) in the Pacific Northwest,

and found that fir snags (*Abies* spp.) had the lowest persistence rates (highest falling rates) over the course of their 5-year study (2001–2006), and that roost snags that persist standing the longest are larger in diameter, shorter in height, and with fewer branches. The authors recommend reserving sufficient, large-diameter leave-trees to provide for future use by roosting bats, given the persistence and fall rates of the snags they observed. Luszcz et al. (2016) studied three bat foraging guilds in southwest British Columbia, Canada, and reported that foraging activity of long-eared myotis increased with increasing forest age, and in general, *Myotis* spp. feeding along edges and in forest gaps was greater in black cottonwood stands and in the interior of old Douglas-fir stands than in other types of forests.

On a far broader scale, Duchamp et al. (2007) suggested managing for bats at a landscape scale to accommodate their nocturnal foraging movements, alternative forest roost sites for some species, and reduce adverse effects of forest habitat fragmentation. The authors cite evidence that some bat species seem sensitive to local habitat fragmentation but also require mixes of conditions across broad geographic areas to meet all resource needs.

Few studies are available on response of bats to fire. Buchalski et al. (2013) studied the response of bats to wildfire severity in a mixed-conifer forest in the Sierra Nevada of California and reported use in unburned, moderate-severity burn, and high-severity burn locations, with large-bodied bats showing no preference. They concluded there was no evidence that burn severity affected site selection and that bats in their region are resilient to landscape-scale fire that served to open up the stands and increase the availability of prey and roost structures.

One issue of growing concern is the recent discovery (March 2016) of white-nose syndrome in a little brown bat (Lorch et al. 2016) 30 mi (48 km) east of Seattle, Washington.³² White-nose syndrome is caused by the fungus *Pseudogymnoascus* (prev. *Geomyces*) *destructans* (Verant et al. 2014) and is a major cause of mortality in a variety of bat species in the Eastern United States, having killed over 6 million bats since 2006. So far, white-nose syndrome

³¹ http://ecos.fws.gov/docs/candidate/assessments/2015/r1/A0J3_V02.pdf.

³² <http://wdfw.wa.gov/conservation/health/wns/>.

seems to exist mainly in species of bats that roost or hibernate in cave or cave-like environments; at least four species of bats in the Pacific Northwest are known to use such locations: little brown bat, big brown bat (*Eptesicus fuscus*), Townsend's big-eared bat (*Corynorhinus [Plecotus] townsendii*), and the poorly known long-eared myotis (*M. evotis*). Whether white-nose syndrome will spread geographically or among species in the Pacific Northwest, or to bat species using other roost situations such as large trees with sloughing bark and dense canopy foliage, is unknown.

The 2001 amended standards and guidelines for NWFP implementation provided specific direction on bat conservation and management, including safety considerations for conducting bat surveys, survey protocols for bat surveys in buildings, environmental education on bat ecology and presence, and plans for building bat boxes. Other issues regarding bat conservation in the NWFP area include gating of caves and mines to prevent disturbance of bat colonies, bat mortality from wind energy installations (Rodhouse et al. 2015, Weller and Baldwin 2012), and concerns for human health and safety from bat colonies occupying buildings.

The ISSSSP has developed a set of management and safety guidelines, and survey and monitoring protocols, for bat conservation in the Pacific Northwest.³³ Forest Service Region 6 has also conducted bat surveys using a Bat Grid Monitoring design and protocol.³⁴ The Bat Grid Monitoring study was conducted from 2003 to 2010 to provide a baseline of knowledge on bat distribution for future change detection, and produced predictive distribution maps of 11 bat species in the Pacific Northwest.

Recent Research and Development on New Tools and Data

Along with scientific studies specifically conducted on species, their habitats, and biodiversity of the NWFP area since the 2006 synthesis (Haynes et al. 2006), several advancements have been made in the development of new scientific tools, datasets, study methods, and assessments.

Much work has focused on developing geographic models and map projections, particularly related to projected effects of climate change. The Nature Conservancy has assessed and mapped the degree of resilience of ecosystems through a broad area of western North America including portions of the Pacific Northwest and NWFP area (east Cascades, west Cascades, north Cascades, Klamath Mountains, California north coast, Pacific Northwest coast, Willamette Valley, and Puget Trough).³⁵ They have also developed a mapping tool for assessing connectivity of “resilient terrestrial landscapes” in the Pacific Northwest (McRae et al. 2016). The U.S. Geological Survey has evaluated vulnerability of species and ecosystems to future climate in the Pacific Northwest.³⁶ A number of other institutions, academic studies, and agencies also have addressed climate change impacts in the NWFP area. Other sources of information on biodiversity and species in the NWFP area include the Washington Biodiversity Council³⁷ and the Oregon Biodiversity Information Center.³⁸

Species and natural-area mapping studies and evaluations, specifically in the Klamath province, have been provided also by Sarr et al. (2015) and Olson et al. (2012). Sarr et al. compared three ecoregional classification systems and found that the top one (Omernik) served to map the distribution of tree species best, but the distribution of mammal species poorest. Olson et al. (2012) projected effects of climate change and determined that some old-growth forest conditions (“microrefugia”) were not captured in the existing reserve designs. These microrefugia conditions would lend to the reserve system being more resilient under climate change; they include old-growth and intact forests on north-facing slopes and in canyon bottoms, lower and middle elevations, wetter coastal mountains, and along elevational gradients. However, the degree to which these findings for the Klamath province can be applied beyond is unevaluated, and we anticipate that other patterns might

³³ <http://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-mammals-bats.shtml>.

³⁴ <http://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-mammals-bats-grid-monitoring.shtml>.

³⁵ <https://www.conservationgateway.org/ConservationByGeography/NorthAmerica/UnitedStates/oregon/science/Pages/Resilient-Landscapes.aspx>.

³⁶ <http://gec.cr.usgs.gov/projects/effects/vulnerability/index.html>.

³⁷ http://www.rco.wa.gov/biodiversity/about_the_council.shtml.

³⁸ <http://inr.oregonstate.edu/book/export/html/549>.

occur in more moist forest environments farther north in the NWFP area (see chapter 2)

A number of tools for analyzing species' habitat connectivity are available (e.g., Brost and Beier 2012). Zielinski et al. (2006) used MARXAN to analyze habitat of spotted owls and fisher in northern California under the NWFP, and concluded that spotted owls require a greater percentage of planning units with habitat than do fisher, and that the current location of the late-successional reserves may not be best to maintain habitat connections for the two species. The individual movement modeling shell HexSim (Schumaker 2013) has been used in the NWFP area to assess habitat patch size and dispersion patterns for northern spotted owls (Marcot et al. 2013, Schumaker et al. 2014). Trumbo et al. (2013) used the modeling tool Circuitscape (McRae and Shah 2009) to evaluate landscape genetics relationships of Cope's giant salamander (*Dicamptodon copei*). Using an example of American pika (*Ochotona princeps*), Schwalm et al. (2015) championed the use of species distribution models for evaluating effects of climate change and habitat conditions on landscape genetics. They found that wide local variation in pika response, ranging from complete extirpation in some areas to stable occupancy patterns in others, as well as habitat composition and connectivity, were important to include in the pika model. Many other simulation and modeling tools are available.

Another recent advancement in mapping species distributions is the development of maps of probability of suitable habitat coupled with maps of uncertainty, such as the one produced by Rodhouse et al. (2012, 2015) and hosted by the ISSSSP, for bat species of the Pacific Northwest (see footnote 34), thereby providing managers with information on locations of habitat as well as confidence in the projections shown spatially. Other approaches to evaluating the effects of forest management on rare species can address the need for balanced sampling designs, such as using the application EstimateS (Colwell 2013) to statistically compare samples among various land condition categories (treatments).³⁹

The modeling framework of MaxEnt (maximum entropy) has recently become popular, as used to map potential habitat in the Pacific Northwest for northern spotted owls (e.g., Carroll et al. 2010, Loehle et al.; also see chapter 4), white-headed woodpeckers (Latif et al. 2015), historical range of California condors (*Gymnogyps californianus*, D'Elia et al. 2015), fisher (Zielinski et al. 2012), and other species. Cautions in the use of the MaxEnt approach, however, have been offered by a number of researchers, such as Yackulic et al. (2013).

Olson et al. (2007b) summarized the strategic surveys conducted under the NWFP guidelines. Strategic surveys (Molina et al. 2003) are designed to fill key information gaps on species distributions and ecologies, the basic criteria for Survey and Manage, to help answer three questions: (1) Does the species occur in or occur close to the NWFP area? (2) Is the species associated with late-successional and old-growth forest? (3) Does the reserve system and other standards and guidelines of the NWFP provide for a reasonable assurance of species persistence? Olson et al. (2007b) reported that all strategic surveys were conducted under the Survey and Manage program from 1994 to 2004 and from 2006 to 2007 (other types of surveys are ongoing on a district-by-district basis). Survey results in the form of projects, publications, and reports on 10 taxa (fungi, lichens, bryophytes, vascular plants, arthropods, mollusks, amphibians, red tree voles, great gray owl, and bats) are now available in a regional repository of the ISSSSP⁴⁰ that is currently managed for a list of sensitive species, some of which are Survey and Manage species.

Recent and Ongoing Issues Adding Challenges and Opportunities

We have raised a number of issues related to other species under the NWFP, in the above text. Here we summarize these and additional challenges and opportunities, regarding conservation of other species, that have emerged since the 2006 synthesis (Haynes et al. 2006).

³⁹ As suggested by D. Luoma. Personal communication. Assistant professor, Department of Forest Ecosystems & Society, Oregon State University, 239A Richardson Hall, Corvallis, OR 97331.

⁴⁰ <http://www.fs.fed.us/r6/sfpnw/issssp/>.

Disturbance ecology and system dynamics—

Wildfire and fire suppression—In this chapter, we have reviewed the mostly adverse direct effects of severe wildfire on fungi, soil arthropods, forest salamanders, forest carnivores, and tree voles. Chapter 3 reviews historical fire regimes of moist and dry forests, effects of recent historical fire suppression on increasing fuel loads and current fire risk, fire suppression effects on reducing early-successional environments, use of fire for restoration, and how the role of fire in late-successional reserves varies across the NWFP area.

Healey et al. (2008) reported that, since the initiation of the NWFP in 1994, loss of large-diameter forests to fire has exceeded loss from harvest across large areas of the region including federal and nonfederal lands (also see Davis et al. 2015). They also found that increased fire incidence is correlated with climate change and that retention of dry-site forests will hinge on coordinated, landscape-scale fire management (also see Halofsky et al. 2014, Podur and Wotton 2010, Sheehan et al. 2015). Healey et al. (2008: 1117) cited several sources for types of strategic fuel management approaches that “may have the potential to reduce the [future] impact of fire on the landscape...”

Although severe wildfire in the NWFP area is an increasing threat to conservation of existing older forest species and ecosystems, it is likely that some dense late-successional and old-growth forests of the Western United States resulted or persisted from decades of fire suppression (Sollman et al. 2016), and others likely were maintained because of topographic positions protected from fire. Fire suppression may have contributed to the development and continuation of some old-forest conditions on the east side of the Cascade Range, such as on the Deschutes National Forest (also see chapter 3). Fuels reduction activities also have changed habitat for many NWFP species in the managed forest matrix, particularly in the dry forest types in the southern part of the NWFP region, by removing fuel loads that in turn reduces presence and amounts of fine litter fuels, and in some cases, large down wood used as den sites by fishers (Sweitzer et al. 2016a).

Historically, frequent fires in dry forest types of the Pacific Northwest resulted in a diversity of vegetation ages and structures among multiple seral stages, and natural legacies of old-forest elements such as large trees and large down wood often remained postfire. But, acting as a double-edged sword, fires of various intensities and frequencies can either create standing and dead wood structures useful to wildlife, or burn them up, unless they are protected or unless fires are of generally low intensity.

As fire regimes in the region shift location and increase in intensity (see chapter 3), there may arise difficult management and social choices between fire suppression and natural fire restoration. Wildfire in the Pacific Northwest is inevitable, and massive “megafires” have the potential of posing threats to old-forest species, such as reported on spotted owls in California by Jones et al. (2016).

Prescribed fire, on the other hand, can be a valuable management tool for maintaining open old-growth forests, forest openings, and early-successional vegetation. Fuels management can help reduce risk of spread of large, high-severity fire. Management options include the use of fuelbreaks, managing for low fuel loading on the forest floor, thinning heavily stocked “doghair” and stagnant stands, pruning to eliminate fire ladders, and other methods. However, there is disagreement over the need to use such treatments, even in fire-prone systems (see chapter 3). Whereas fuel treatments and restoration management can affect fire behavior, these actions expose bare soil that, in turn, could promote invasion of exotic plant species that could spread into adjacent lands, and where nonnative plant cover is twice as high on fuel breaks as in adjacent wildlands (Merriam et al. 2006; also see chapter 3).

Modeling by Ager et al. (2007) suggested that treating 20 percent of the landscape as fuel breaks and for fuel reduction can result in a 44 percent increase in the probability of conserving late-successional and old-growth spotted owl habitat (see chapter 4). Moriarty et al. (2016) suggested that fuel treatments that simplify stand structure can adversely affect movement and habitat connectivity for martens. The authors noted that martens avoided openings

at landscape- and home-range scales, but thinned stands were avoided mostly at only the home range scale because such stands were uncommon in their study area of Lassen National Forest, in northeastern California, just outside the NWFP area. The authors also suggested that fuel treatments that would create landscape- or home range-scale openings be done at elevations lower than where martens typically occur. However, the effectiveness of such fuel-treatment approaches in providing for marten habitat likely varies with other considerations, in particular, habitat considerations for other species. Such treatments also may be more effective on south slopes and ridgetops, and less effective on north aspects and in drainage bottoms.

Further, fuels treatments as mentioned above could have adverse effects on the many species, within late-successional and old-growth forest stands and other forest age and structure classes, that occupy litter and duff (the organic matter soil horizon) and down wood of various decay, size, and cover classes and tree species, as well as standing partially dead trees and snags. The flip side of this, however, is that species associated with open stands of late-successional and old-growth forest or from burn conditions might benefit from lower severity wildfire. As mentioned below, balancing reduction of fire risk with providing such microhabitat elements for species and biodiversity conservation can be the subject of a decision analysis and decision science process (see chapter 12).

Early-seral vegetation environments—Early-successional vegetation conditions provide habitats for young-seral species and thus contribute to overall biodiversity of the region (Swanson et al. 2011, 2014). For example, in the Klamath-Siskiyou province, early-successional vegetation, including grass- and shrub-dominated areas and areas of young trees and shrubs, helps support abundant populations of dusky-footed woodrats, which are a primary prey of northern spotted owls in that area. Lists of special status and sensitive species of Forest Service and BLM in Washington and Oregon include a few species associated with early-successional vegetation conditions, such as green-tailed towhee (*Pipilo chlorurus*).

However, natural early-successional vegetation conditions have been reduced throughout the Western United States, including the Pacific Northwest, from plantation forestry operations of tree planting, control of competing vegetation (typically broad-leaf or hardwood species), fire suppression, and some thinning (Bormann et al. 2015). The result is reduction in habitat for some species closely associated with complex, natural early-successional conditions and with landscape mosaics of early-successional and old-growth forests. For example, Betts et al. (2010) reported declines in bird species associated with early-seral broadleaf-dominated vegetation following stand-replacing disturbance, including timber harvest and fire, in the Pacific Northwest. However, variable-density thinning and heavy thins can promote subsequent growth of hardwood shrubs and trees and provide habitat for some species of lichens, invertebrates, and other organisms (Schowalter et al. 2003, Wilson and Puettmann 2007), but more study is needed on specific conditions and effects on late-successional and old-growth species (Wilson and Forsman 2013). In general, though, early-successional vegetation created naturally from fire, windstorm, or other natural disturbances usually differs substantially in structure and composition of plant and animal species from that created by timber harvest subsequently subject to tree planting and control of competing vegetation.

Early-successional vegetation conditions created artificially also can become sources of plants, invertebrates, and other species that can compromise adjacent forest-interior conditions in late-successional and old-growth forest patches. We have noted that recent historical increases in artificially created early-successional vegetation have likely provided increased habitat for predators of mustelids, such as bobcats and coyotes.

Depending on amounts and dispersion patterns, early-successional vegetation can serve to fragment otherwise extensive and connected late-successional and old-growth forest cover and impede movement, population connectivity, and population size of canopy-associated late-successional and old-growth species such as red tree voles, martens,

fishers, northern flying squirrels, and brown creepers. Sollmann et al. (2016) found that forest thinning reduced local densities of northern flying squirrels which moved into adjacent unthinned stands; that there was no difference in effect from even thinning and variable thinning; and that their results underscore the need to consider effects of stand management in a broader landscape context, particularly regarding movement and dispersal (functional response) of the squirrels.

Full range of conditions—The full suite of native species and biodiversity of the Pacific Northwest occurs only with a mix of the full range of environmental conditions, including all forest successional stages that support the full array of terrestrial vertebrates (e.g., Raphael 1988, Raphael and Marcot 1986) and geophysical conditions such as montane wet meadows that support associated invertebrates. The NWFP was designed to provide aquatic and late-successional and riparian forest environments; the most appropriate balance with other conditions (e.g., mixed of early- and late-successional stages) is beyond the scope of the NWFP’s original mandate and its guidelines under the 1982 planning rule, and is a matter of public policy informed by the best science. The 2012 planning rule increases emphasis on ecosystem approaches to conserving species and ecological integrity, based in part on the coarse-filter approach of providing structures and functions of ecosystems. Thomas et al. (2006) called for the NWFP to be recast as an integrative conservation strategy with federal forests managed as dynamic ecosystems.

Climate change: indicators, tipping points, and thresholds—

Complicating management objectives to provide the full array of environmental conditions are the dynamic and potentially sudden influences of climate change on vegetation, fire, species distributions, and development of so-called novel ecosystems, that is, specific groups of species in assemblages and communities that have not previously occurred in those combinations (Collier 2015, Pace et al. 2015). The current locations of late-successional

forest reserves under the NWFP may not be fully resilient to expected effects of climate change. We know little about how management of vegetation, fuels, and fire will be affected by changing climate and thus the future of late-successional and old-growth forests and associated species and biodiversity (Pfeifer-Meister et al. 2013, Wimberly and Liu 2014), although some indications are that, under the NWFP, large reserves in moist forests have been relatively stable compared to smaller reserves, and that moist forests may be at less risk of disturbance from climate change than are dry forests (see chapter 3).

Changes in climate, particularly increased temperatures and aridity, can adversely affect amphibians by altering regimes of local hydrology (see chapter 7 for more discussion of climate change effects on aquatic ecosystems), fire, and disease occurrence and transmission. More directly, it may cause physiological stress on species adapted to cool and moist microclimates. If climate change reduces availability and connectivity of higher elevation older forests, populations of associated carnivores such as lynx, wolverine, and marten might suffer, as summarized above (“Carnivores” section). In a study of 34 species of small mammals in montane environments in the Sierra Nevada Mountains and Lassen Volcanic National Park in California, Rowe et al. (2015) reported that 25 species shifted their ranges since the early 20th century in response to climate change, with high-elevation species contracting their lower range limits downslope and low-elevation species showing variable responses.

Some lichens and bryophytes found in late-successional and old-growth forests could serve as useful indicators of changes in, and thresholds of, forest conditions and climate and air quality (Ellis 2015, Hofmeister et al. 2015, Mölder et al. 2015). Some late-successional and old-growth-associated invertebrates also can serve as indicators of biodiversity and climate change risk (Yi and Moldenke 2005). Halpern et al. (2012) suggested that disturbance has passed a threshold level for bryophyte conservation in Pacific Northwest forests, and that 15 percent retention of late-successional conifer forests—which the authors cited as the minimum

standard on federal forest lands in the Pacific Northwest⁴¹—is inadequate to maintain the abundance or diversity of late-successional and old-growth species.

Turner et al. (2015) developed models to simulate potential impacts of climate change and disturbances on vegetation in the Willamette River basin of Oregon. Their projections under climate-warming scenarios suggest that potential forest types will transition from evergreen needleleaf to a mix of broadleaf and needleleaf forms, with as much as a nine-fold increase in area burned by the end of the 21st century. Implications for lowland plants and animals suggest a shift to more warm-temperate and fire-tolerant species mixes. Research by Creutzburg et al. (2017) suggests a more modest change, as climate change will increase forest productivity and carbon storage capacity that will not be offset by wildfire under the legacy of management and fire suppression activities.

Carroll et al. (2010) modeled the NWFP reserve network under climate change stressors. They concluded that using spotted owls as an umbrella species for reserve design, as under the NWFP, included habitat for other dispersal-limited (“localized”) species but will not include much of their core habitats, as defined in the Zonation model they used, under projections of climate change. Thus, the current array of fixed late-successional and old-growth reserves designed for the owls may not suffice for other species that may change locations as their habitats shift under climate change. The authors suggested that a fixed-reserve system should be designed for resilience under anticipated climate change by including additional areas, particularly by adding reserves in higher elevation sites (Carroll 2010). However, not all species will respond to climate change by migrating their distributions to higher elevations. Crimmins

et al. (2011) reported anomalous downhill shifts of some plant species in California, following regional changes in climatic water balance rather than responding to changes in temperature. Tingley et al. (2012) projected effects of climate change on altitudinal distributions of birds in the Sierra Nevada of California, and found that species will likely respond differently and individually at their range margins to increasing temperature and increasingly variable precipitation. Similarly, variable responses by plants, birds, and other taxa in the NWFP area are likely. The lesson is that the response to changing climate and associated weather conditions will vary, sometimes greatly, among species, and such variability will likely occur in all taxa and species groups.

Bagne et al. (2014) evaluated the efficacy of managing for special status species under climate change and its secondary effects on providing for overall biodiversity in the Southeastern United States. They concluded that 74 percent of terrestrial vertebrates potentially vulnerable to climate change were not included in current lists of special status species, and omissions were greatest for birds and reptiles. Current lists of special status species—potentially including those of the ISSSSP in the Pacific Northwest under the NWFP—could be evaluated for potentially adding species that might be vulnerable to climate threats, particularly with regard to those environmental conditions that would exceed physiological thresholds, if such species were lacking. Such analyses have not been conducted across all taxa, although this chapter and others in this compendium (e.g., chapter 2) provide insights on some species that might be vulnerable to future climate change effects, such as marten, pika, and others.

Aside from climate change effects are questions of balancing conservation between biodiversity and individual species. For example, Arthur et al. (2004) modeled tradeoffs in western Oregon between maximizing overall species richness and providing for individual endangered species. They found that the tradeoff was nonlinear, whereby increasing endangered species protection from

⁴¹ The reference here to 15 percent retention might more clearly pertain to the 1994 Standards and Guidelines (USDA and USDI 1994, page C-41) that state “Landscape areas where little late-successional forest persists should be managed to retain late-successional patches. This standard and guideline will be applied in the fifth field watersheds (20 to 200 mi² [52 to 518 km²]) in which federal forest lands are currently comprised of 15 percent or less late-successional forest.”

90 to 99 percent caused a decline in all-species protection 2 to 14 percent. Their risk tradeoff approach could be applied elsewhere throughout the Pacific Northwest and NWFP area to evaluate variations in proportion of landscapes in late-successional and old-growth reserves and managed matrix lands (Arthur et al. 2002). However, their model did not quantify species persistence or viability, and it did not evaluate endemic, rare, or specialized species considerations that could be added to their analysis.

Matrix lands and habitat connections—

Many studies suggest that habitat connectivity, fragmentation, and isolation are among the most dire stressors on populations, and that, under environmental changes, movement barriers for many species are likely to shift over space and time (Caplat et al. 2016). Much has been written in the conservation biology literature on wildlife habitat corridors and connectivity among habitat patches and reserve areas (e.g., Beier and Brost 2010; Beier et al. 2008, 2011). The literature has addressed the influence of conditions in matrix lands on arthropods (Shields et al. 2008), butterflies (Ross et al. 2005), birds (Neuschulz et al. 2013), and mammals (Kurek et al. 2014). Beier et al. (2008) suggested designing habitat linkages between wildlands to provide for multiple species rather than single focal species. Brady et al. (2009) used measures of disturbance, structure, and floristics in habitat core areas, habitat patch edges, and matrix landscapes to assess small mammal populations in northeast Australia. They found that small mammal conservation entails controlling disturbances and providing for habitat preservation and restoration within matrix lands. Wessell's (2005) study of abundance and diversity of vascular plants, arthropods, amphibians, and mollusks in western Oregon suggested the value of managing forest matrix lands for habitat heterogeneity.

Several studies in the Pacific Northwest have suggested a need to view the LSR system under the NWFP as a habitat network (Molina et al. 2006), and to provide further connectivity of forest reserves to account for anticipated climate-change shifts in habitat quality for sundry species (Beier and Brost 2010; Carroll et al. 2010; Rudnick et al. 2012; see also Spies et al. 2010). Proulx and

Santos-Reis (2012) determined the value of conserving late-successional and old-growth forests and structures in the Pacific Northwest as habitat for Humboldt and Pacific martens. Habitat connectivity for organisms such as northern flying squirrels and red tree voles that depend on contiguous tree canopies can be severed with heavy thinning of forests designed to accelerate diameter growth of trees (Wilson and Forsman 2013).

In general, these studies suggest also that the degree to which protection of species' habitats in the matrix may contribute to population conservation is species-specific, and depends in part on the size and configuration of habitat in core areas, patches, and habitat linkages, and in the amount of habitat occurring in land already conserved (e.g., NWFP LSRs).

Coarse and fine filters in conservation—

Since the last science synthesis, much has been written on sundry approaches to management of species, biodiversity, and ecosystems. These approaches include coarse- and fine-filter management; use of surrogate species such as indicators, umbrellas, keystones, and flagships; management of ecosystem services; management for the range of natural or historical variation; and much more. Much of this literature poses approaches with scant empirical testing and validation of their efficacy, however. In this section, we review recent literature on such topics as pertains to the Pacific Northwest.

A popular method for conservation assessment and management uses both coarse filters and fine filters (Hunter 1991, Noss 1987). This approach first addresses coarse-level, spatially broad environmental conditions to meet general habitat associations assumedly of a large number of species, and next adds specific environmental components and locations to meet the needs of additional species not sufficiently provided by the coarse-filter level. Such an approach was used to craft the late-successional forest and the riparian and aquatic reserve system under the NWFP. Further, the Survey and Manage standards and guidelines and its annual species reviews were specifically designed to determine which late-successional and old-growth-dependent species would not be sufficiently provided by the coarse-level reserves and may need additional fine-filter management. Determinations

were made based on a rigorous, structured approach to evaluating new information (Marcot et al. 2006).

In the Pacific Northwest, Lehmkuhl et al. (2007) applied a coarse-filter approach to define the historical range of variability of the composition and pattern of forest conditions under historical fire regimes, and then a fine-filter approach to establish fuel reduction treatments at the landscape scale to provide conditions for the food web involving northern spotted owls and their prey of northern flying squirrels and bushy-tailed woodrats (*Neotoma cinerea*). Overton et al. (2006) used a coarse-filter regression approach with Gap Analysis Program information to evaluate use by band-tailed pigeons (*Patagioenas fasciata*) of abandoned mineral sites in western Oregon. In a more complex variant to the approach, Higgins et al. (2005) devised a four-tier spatial approach to identify areas critical to conservation of freshwater biodiversity in the Columbia River basin of the Pacific Northwest.

Thompson et al. (2009) used a coarse-filter approach to define the historical range of variability of forests in the Oregon Coast Range as a target for biodiversity conservation, coupled with what they termed the social range of variability to account for resource and land use patterns. They found that land development, shifts in anthropogenic fire regimes, and climate change will likely impede the use of historical range of variability as a management objective, and that more complex planning is needed in a continuous process of negotiation.

A coarse-filter approach to ecosystem management can also assume that emulating natural processes and disturbance dynamics, such as natural fire regimes, will provide for forest biota at natural population levels (Armstrong et al. 2003). Other variants of the coarse-filter approach include identifying and protecting biodiversity “hot spots” under the assumption that most or all biota will be provided. Some applications of the “hot spot” approach focus not on overall species richness but on identifying concentrations of locally or regionally endemic species, which tend to be range restricted and perhaps extinction prone. “Endemism hot spots” could signal climate refugia, and protecting them could be a useful tool for avoiding extinction under climate change (e.g., Harrison and Noss 2017).

The strength of the coarse- and fine-filter approach is that it can account for a wide array of species generally associated with an ecosystem condition such as late-successional or old-growth forests. The weakness of this approach is that it may require site surveys, habitat and ecological studies, and intensive assessments of individual species to determine which require fine-filter attention. Fine-filter management is typically viewed as management and conservation of species-specific habitats. This means that understanding species taxonomy is key to determining biological entities of potential conservation concern. In this chapter, we cite several instances of recent taxonomic findings describing newly identified species. Recent work on mitochondrial DNA (mtDNA) of flying squirrels now suggests that a geographically limited Pacific coastal form of northern flying squirrel ranges from southern British Columbia to southern California (Arbogast et al. 2017); the authors suggest the new species be designated Humboldt’s flying squirrel, using the previously suggested epithet *Glaucomys oregonensis*. Nearly all of the occurrence of flying squirrels in the NWFP area now pertains to this newly described species, although there seems to be overlap with the northern flying squirrel (*G. sabrinus*) in western Washington. Thus, two species of flying squirrels may occupy the NWFP area, and their individual conservation status has yet to be determined.

Hunter (2005) suggested use of an additional “mesofilter” level to complement the coarse-filter strategy for conservation of entire ecosystems in reserves and the fine-filter strategy for conservation of selected individual species. The mesofilter (“middle” filter) level would focus on conserving critical ecosystem components, specifically microhabitat attributes such as logs, snags, and pools used by invertebrates, fungi, nonvascular plants, and other species groups that are often overlooked in conservation planning. Hunter suggested that the mesofilter approach is particularly useful with seminatural conditions, for sustaining biodiversity and commodity production. In a sense, management for snags, down wood, riparian conditions, and other ecosystem components, as already part of agency activities in the NWFP area, constitutes a mesofilter approach.

Assumptions of the coarse- and fine-filter approach, however, are seldom empirically tested in a controlled and rigorous manner, and their blind application may lead to inappropriate expectations for species conservation (Cushman et al. 2008, Hunter 2005). In one test in Oregon, Fagan and Kareiva (1997) found that hot spots of biodiversity (species richness) did not coincide with locations of rare and endangered butterflies. In another test in Oregon, Cushman et al. (2008) found that only 4 percent of the variation in bird species abundance was explained by using a general characterization of habitat as a proxy to bird species abundance and vegetation cover type as a proxy to habitat. Further, Molina (2008) found that 90 percent of fungal species has some fraction of their known sites occurring within LSRs in the NWFP area, but some 66 percent of all sites of Survey and Manage fungal species occurred outside the reserves, so that the coarse-filter level alone to identifying late-successional and old-growth forest reserves would not provide for many rare fungi. There may be some conservation salvation for these sites outside late-successional and old-growth forest reserves, insofar as the reserve strategy has changed, at least de facto, with little to no harvest of federal-land late-successional and old-growth forests outside the original NWFP LSRs (see chapter 12).

Related to the coarse- and fine-filter approach to conservation is the application of species- and system-level approaches to biodiversity conservation. Species-level approaches include use of population viability analyses; surrogate species approaches with the use of umbrella species, focal species, guild surrogates, habitat assemblage surrogates, management indicator species, biodiversity indicator species, apex predators, and flagship species; multiple-species approaches with use of entire guilds and entire habitat assemblages; and geographically based approaches with identifying locations of target species at risk and species hot spots or concentration of biodiversity (Marcot and Flather 2007). System-level approaches include the use of range of natural variability, key habitat conditions, keystone species, ecosystem engineers, and approaches to emulate or provide for regimes of fire, herbivory, key ecological functions of species assemblages, and food webs (Marcot and Sieg 2007). Much has been written about most of these species-

and system-level approaches (e.g., Branton and Richardson 2014, Breckheimer et al. 2014, Hunter et al. 2016), including due caution in using them without clarity of definitions (Barua 2011) and empirical verification of their assumptions (e.g., Bifulchi and Lodé 2005).

Each of these species- and system-level approaches carries strengths and weaknesses depending on management objectives. No one approach is fully effective in providing for species diversity, genetic diversity, and ecosystem diversity conservation objectives, which is why the 2012 planning rule specifies use of both coarse- and fine-filter management. Raphael et al. (2007) and Raphael and Marcot (2007) suggested that a combination of approaches may be efficacious for meeting multiple conservation objectives. They suggested a series of steps to identify the suite of species- and system-level approaches for conservation of rare or little-known species and their habitats, and they emphasized the critical importance of setting clear goals, identifying measurable short- and long-term objectives, and including learning objectives to increase knowledge.

Uncertainties and Research Needs

Indicators and Surrogates

The debate over use of surrogate species in its many forms—indicators, sentinels, flagships, umbrellas, keystones, and others—bears closer scrutiny. The literature seems divided over cautioning against its use for more holistic ecosystem management, and using it with caution for more focused needs with clearly stated objectives. The Forest Service guidance on implementing the 2012 planning rule calls for the use of a surrogate-species approach to evaluating conditions contributing to viability of groups of species of conservation concern. In this approach, the viability of the surrogates is assumed to represent viability of the broader subset of species of conservation concern species with similar ecological requirements and similar responses to environmental conditions and changes. Surrogates are chosen also on the basis of their having more stringent ecological requirements than others of the group. Most important, surrogates are meant to represent ecological conditions that contribute to viability of the broader species group and are not meant to directly represent their

population dynamics *per se*; this qualification may add some uncertainty over the response of the fuller species groups (Wiens et al. 2008) but is truer to the concerns expressed in the literature over use of indicators of population viability.

In many ways, the evolving planning and management framework under the NWFP has largely met earlier calls for conservation of forest biodiversity and other species. This includes viewing forests as dynamic ecosystems, explicitly providing for ecosystem processes, addressing little-known species, and attending to conditions in the managed forest matrix (Franklin 1993). Recent scientific publications have continued to suggest using management indicator species and umbrella species only with caution and empirical evaluation (Branton and Richardson 2010, 2014; Wiens et al. 2008), or not at all, because the terms are typically poorly defined and the concepts often untested. Simberloff (1998) argued that moving toward an ecological functional basis for species and biodiversity management, rather than using simpler proxies of indicator species, is more likely to succeed, particularly with application of ecological forestry approaches. Also, much recent scientific literature has moved beyond the contentious era of arguing whether species or systems should be the focus for conservation; the approaches are, as Lindenmayer et al. (2007) and Sergio et al. (2003) noted, complementary, including with the debate over conserving species versus ecosystem services (Hunter et al. 2014, Kline et al. 2016; also see chapter 8). Ultimately, successfully combining species and system approaches depends on clearly articulating management objectives and determining the efficacy of management through monitoring and research.

Climate Change

Climate change is at the heart of many conservation concerns in the Pacific Northwest, but much empirical study remains to better understand its effects on species, their habitats, and their ecological functions, and to avoid the easy but potentially erroneous assumption that it is the proximate cause of most or all observed perils, such as with the decline of amphibians (Davidson et al. 2001, 2002). Climate change might act indirectly, such as by reducing snowfall (Corn 2003, Forister et al. 2010) and increasing evaporation, leading to loss of wetlands (Corn 2005); by increasing fire

frequency and extent; and by improving environmental conditions for diseases or pathogens that in turn could affect other organisms, again such as with the decline of amphibians (Carey and Alexander 2003).

Further, there may be higher order effects whereby climate change in turn alters biotic interactions among species, such as Preston et al. (2008) reported on an endangered butterfly and a threatened bird species in southern California, which led the authors to suggest that considering species interactions is important when designing reserve systems for conservation of sensitive species. How this can influence any redesign of reserves under the NWFP would require study.

Microbiota and ecosystem functioning—

There is a dearth of knowledge and understanding of Wilson's (1987) "little things that run the world," referring to the invertebrates, particularly soil microorganisms and mesoarthropods, along with the suite of poorly known fungi and allies, that collectively play crucial roles in organic matter breakdown, nutrient cycling, and maintenance of forest health (Berg and Laskowski 2005, Ulyshen 2016, Tolkkinen et al. 2015). There is also much to be learned about how biodiversity affects ecosystem functioning (Hooper et al. 2005, Marcot 2007). Here is where an interplay and complementarity of species-focused and ecological-process research and management can usefully provide for forest ecosystem conservation and restoration to meet early visions of achieving sustainable forestry for conservation of late-successional and old-growth forests and their biodiversity (Beebe 1991, Crow 1990).

The future of other species is not written, but is progressively being tilted by fluctuating dynamics of fire regimes, shifts of forest composition and structure from changing climates, increasing stressors of invasive species, and changes in societal values with increasing needs for progressively scarce forest resources and their ecosystem services. Successfully addressing these and related issues for restoration and conservation of other species of late-successional and old-growth forests necessitates continuing on the path to whole-ecosystem research and adaptive management in a socioecological scope (Ban et al. 2013, Schmiedel et al. 2016; also see chapters 8 and 11).

Conclusions and Management Considerations

Summary of Key Management Considerations

Here we apply our key findings to the set of guiding questions posed after the introduction of this chapter. We also address this additional question: If all late-successional and old-growth forests are now protected under the NWFP (and under designated critical habitat for the northern spotted owl), is there a need for additional fine-filter approaches, beyond those for the northern spotted owl and marbled murrelet, for species assumed to be associated with said forests? We will conclude the affirmative, that sufficient uncertainty exists, particularly with future changes in old-forest environments and species-specific habitats, particularly for species that are rare or poorly known or that are expected to be increasingly adversely affected by disturbances and changing climate. As to which species may need to be so studied and provided is at present unclear, although advances in identifying species of conservation concern under the 2012 planning rule will be a major step forward.

Current scientific understanding of late-successional and old-growth-associated species rarity—

Although no specific analysis has been conducted to quantify the degree of rarity of late-successional and old-growth-associated species in the latest Survey and Manage species list, the general trends since the 2006 synthesis (Haynes et al. 2006) are:

1. Scientific understanding has greatly expanded on the occurrence, distribution, ecology, and potential threats of many of these species, as noted by the rich array of inventory reports and conservation planning documents conducted by the ISSSSP.
2. Numerous Survey and Manage species had been removed from that list, during the last rounds conducted on the annual species reviews, largely because predisturbance site surveys suggested that the species are more common or frequent than previously thought and because they no longer met the concern for the basic criteria for persistence of Survey and Manage species.
3. The best scientific information to date on many of the remaining Survey and Manage species, as summarized by the ISSSSP, suggests continued rarity

or potential vulnerability to habitat disturbance. It is true that some old-forest associated species are naturally rare and perhaps more vulnerable than are more abundant species, but neither the NWFP guidelines nor the 2012 planning rule state that naturally rare species are to be discounted. Although it may not be possible or feasible to increase the distribution or abundance of naturally rare species, their preservation or conservation is very much in the spirit of the NWFP. The specific lists of species currently considered under the ISSSSP are found on the web site cited above.

Planned provision of habitat for rare and uncommon species under the NWFP—

The NWFP was initially envisioned as providing habitats for late-successional and old-growth-associated species through a combination of LSRs, AMAs, aquatic and riparian reserves and buffers, and selected conservation of late-successional and old-growth forest stands in the managed forest matrix. At present, however, there is little to no harvest of late-successional and old-growth forest stands on BLM and Forest Service lands within the NWFP area, including within and outside reserves (see chapter 3). In this way, the current implementation of the NWFP is not the same as initially envisioned.

Further, there has been no official effectiveness monitoring program under the NWFP, so there is limited information for determining the degree to which all old-forest associated species are being provided, and determining their habitat and population trends. Without a formal effectiveness monitoring or research program for other species and biodiversity under the NWFP, much has to be inferred from vegetation and late-successional and old-growth conditions and trends, although some research is available on selected species and taxa, as reviewed in this chapter.

Rare and uncommon species populations under NWFP management—

The degree to which remaining late-successional and old-growth forest is adequately providing for persistent and sustainable populations of rare and uncommon species is known for only those few species for which demographic and monitoring studies have been conducted, such as with northern spotted owls and marbled murrelets. Lichen

sampling, if reinstituted in the Forest Service's FIA program, could provide answers to this question on late-successional and old-growth-associated species. Otherwise, as under the 2012 planning rule, assumptions need to be made on the efficacy of coarse-filter approaches to providing general ecosystem conditions and their adequacy to in turn provide habitat and resource conditions for individual species. To gain greater knowledge and understanding of the adequacy of these assumptions may entail further species-specific research and monitoring, perhaps including species previously deemed secure but now perhaps with uncertain futures from climate change and other disturbances unforeseen in the initial NWFP guidelines.

Efficacy of habitat conservation management recommendations—

There has been no effectiveness monitoring program for the bulk of Survey and Manage species. Various studies suggest some degree of success for specific species groups, such as headwater amphibians. For most species, however, no monitoring data are available.

Sufficient information to change species management status—

This would best be determined through an evaluation of recent scientific data, survey information, and other sources, through a structured species review process (e.g., the previously instituted annual species reviews under the Survey and Manage program). The ISSSSP has compiled and organized much survey and scientific information on the Survey and Manage species but has not been charged with conducting such species reviews per se. It is likely that significant information has been gathered on some species for making this determination, but at least some of the determination will not be made until or unless an adaptive learning and management procedure, along the lines of the annual species reviews (last conducted in 2003), is implemented again.

Species persistence under NWFP reduced late-successional and old-growth harvest—

No analysis has been conducted on threats to late-successional and old-growth-associated species, comparing threat levels perceived under the NWFP as initially envisioned with the NWFP as currently implemented. The NWFP as initially envisioned included late-successional and old-growth reserves, aquatic and riparian reserves and buffers,

AMAs, and site-specific conservation of late-successional and old-growth forest stands. The NWFP as currently implemented includes little to no harvest of late-successional and old-growth forests on Forest Service and BLM lands within the NWFP area (see chapter 3); and provides for some management in LSRs in dry forests. Such a comparison between initial and current management conditions could recognize the greater roles and potential changes from shifts in local climate and especially in fire suppression, fuels management, and alternative ecological forestry approaches providing for the development of natural, complex early-seral and young-age forest vegetation conditions.

Under this suite of considerations, it is likely that some species originally ranked as having low potential for persistence might be viewed as having higher potential, particularly if their late-successional and old-growth forest habitats and specific habitat elements are better provided throughout the matrix as anticipated over time. Other species tied to specific locations that are vulnerable to fire and disturbances might not fare as well, but no evaluation is currently available. It is likely that location data accumulated from predisturbance and strategic surveys might suggest adequate numbers by which to reduce concern for the persistence of some late-successional and old-growth-associated species, but predisturbance surveys alone do not provide information on whether the species survived the disturbance, only that they were present before the disturbance.

Late-successional and old-growth species added to species-of-concern lists—

Several "species lists" cover the NWFP area, including the lists of Survey and Manage species; Forest Service and BLM special status and sensitive species; U.S. Fish and Wildlife Service's candidate, threatened, and endangered species under the Endangered Species Act; and various lists of species of conservation or vulnerability concern from International Union for Conservation of Nature Red Data Lists, state lists, Washington Natural Heritage Program, Oregon Biodiversity Information Center, and others. The Forest Service also is currently compiling a regional list of potential species of conservation concern that will also intersect with the NWFP area, its late-successional and old-growth forest-associated species, and many of the other lists noted here. There has been no evaluation of the composite set of species among all these lists, comparing

to the list of late-successional and old-growth species originally ranked as high persistence or not identified as conservation concern under the NWFP.

This report has summarized new threats or new scientific information revealed over the past decade, suggesting potentially growing conservation concern for late-successional and old-growth forest-associated bats, should white-nosed syndrome spread in the Pacific Northwest; highly locally endemic mollusks and other invertebrates, under threat of invasive invertebrate species or reduction in key habitat elements (e.g., large, old bigleaf maples for the Puget Oregonian snail); new species recently split taxonomically and that occupy a smaller portion of the former species' range and that might associate with late-successional and old-growth forest environments that could be at risk from increasing wildfire (e.g., the new species of sharp-tail snake recently identified from northwest California and southwest Oregon); and other examples. For some species, a more extensive evaluation is in progress; for others, it awaits.

In the past annual species reviews, no species was added to the initial Survey and Manage species list.⁴²

⁴² New information to support adding a species to the Survey and Manage list (USDA and USDI 2001:15-16) must address the following three basic criteria including the specific factors used as a basis for determining concern for persistence, in addition to criteria for late-successional and old-growth association:

- The species must occur within the NWFP area, or occur close to the NWFP area and have potentially suitable habitat within the NWFP area.
- The species must be closely associated with late-successional or old-growth forest.
- The reserve system and other standards and guidelines of the NWFP do not appear to provide for a reasonable assurance of species persistence.

The specific factors must apply to at least an identified portion of the species range, on federal lands, within the NWFP area. One or more of the following factors may indicate that persistence is a concern. These factors must be considered in the context of other standards and guidelines (other than those related to Survey and Manage) in the NWFP:

- Low to moderate number of likely extant known sites/records in all or part of species range.
- Low to moderate number of individuals.
- Low to moderate number of individuals at most sites or in most populations.
- Very limited to somewhat-limited range.
- Very limited to somewhat-limited habitat.
- The distribution of the species within habitat is spotty or unpredictable in at least part of its range.

Effects of fire on rare and uncommon late-successional and old-growth species—

Species and ecosystems of the NWFP experience diverse forest types and disturbance regimes (Bunnell 1995).

Regarding effects of fire on rare and uncommon species associated with older forests, research results are limited and mixed. For example, for some species such as white-headed woodpecker, small patches of prescribed burning can be used to create or elevate the quality of their habitat. For other species, prescribed fire (particularly associated with sanitation and safety harvests) and associated fuels reduction activities can diminish key late-successional and old-growth habitat components such as large hollow trees, large snags, and large down logs, in turn reducing habitat for a wide variety of denning and cavity-using wildlife species, including fishers, Humboldt marten, and other mammalian carnivores (fig. 6-3).

Effects of high-severity, stand-replacing wildfire are perceived as largely adverse to species that use dense, multilayered older forests, although much of the literature has not clarified levels of wildfire intensity, extent, and location when speculating or concluding on wildfire effects. Research suggests that (high-severity) fire can reduce beneficial mycorrhizal and other desirable fungi, but for the most part, the effects of wildfire on individual, rare and uncommon late-successional and old-growth species are largely unstudied. Insofar as atypically severe wildfire can reduce canopy closure and fragment dense, contiguous-canopy stands, it adversely affects habitat for red tree voles.

Little has been studied on the direct effects of suppression of wildfire on wildlife in the Pacific Northwest.

Efficacy of site buffers compared with landscape-scale habitat management for species persistence, dispersal, and habitat connectivity—

No studies have specifically compared site buffers with landscape-scale management of late-successional and old-growth forests, as influencing persistence, dispersal, and habitat connectivity of late-successional and old-growth-associated species. Similarly, little research has been done in the effectiveness of landscape-scale management in assuming a reasonable assurance of persistence



Bruce G. Marcot

Figure 6-3—Naturally hollow Douglas-fir log in a lowland old-growth conifer forest, west Cascade Mountains of northern Oregon. Such logs are prime denning sites for fishers, Humboldt marten, and other mammalian carnivores. Some prescribed fire, timber salvage operations, and fuels reduction activities can reduce such habitat elements.

of late-successional and old-growth-associated species. However, some studies have noted that small patches of late-successional and old-growth forests, and legacy elements of late-successional and old-growth forest such as large green trees, large snags, and large down wood, can provide limited habitat for old-forest-associated species in the forest matrix. Such patches can also be used as points of restoration of older forest. Studies of old-forest remnants and fragments suggest that, by themselves, they provide only a fraction of the original forest biodiversity. Although they may be valuable to conserve for that purpose, if they are the only remaining elements of late-successional and old-growth forests in an area, they cannot be counted on to

provide for persistence, dispersal, and habitat connectivity of late-successional and old-growth-associated species.

In general, riparian stream buffers, including unthinned riparian vegetation and adjacent upland forests, have been shown to have immense conservation values, particularly for amphibians. Buffers of sufficient width along streams and wetlands can serve as movement corridors and landscape linkages for aquatic frogs, salamanders, reptiles, and birds (Perry et al. 2011, Vergara 2011). Streams with riparian buffers can provide for some birds (Nimmo et al. 2016) and arthropods. Studies suggest that buffer widths differ to accommodate different species or species groups, as noted in summaries above.

The ISSSSP under the NWFP—

The ISSSSP has provided a wealth of information to agency researchers and managers on late-successional and old-growth-associated species, such as conservation assessments, summaries of surveys, threats evaluations, range maps, survey protocols, and ecological studies. It should be remembered that the ISSSSP pertains only to Washington and Oregon; in California, the Forest Service and BLM each have their own species programs and lists.

The current special status and sensitive species list—

Some Survey and Manage species also have sensitive or strategic species status, but the ISSSSP list does not include Survey and Manage species, which are maintained as separate lists based on different criteria. The current special status and sensitive species list includes those Survey and Manage species that qualify based on the ISSSSP criteria. The current ISSSSP list is broader in application covering all of Oregon and Washington with updates as recent as July 2015. The ISSSSP list includes a broad array of fungi, lichens, bryophytes, vascular plants, vertebrates, and invertebrates. Late-successional and old-growth species, or any species for that matter, with risks to persistence would likely have been included in the list update. No changes have been made to the Survey and Manage species list since 2003 owing to capacity and funding limitations to conduct an annual species review (the adaptive management process that removes or adds species or changes Survey and Manage category based on new information).

Additional fine-filter approaches for species assumed to be associated with late-successional and old-growth forest—

Not all late-successional and old-growth forests are being protected, but surveys for Survey and Manage species are being conducted in stands over 80 years of age. Much remains to be learned about demography, persistence, viability, dispersal, and habitat connectivity of most late-successional and old-growth-associated species. The coarse-filter approach assumes that providing for general habitat conditions at a broad scale, such as through LSRs and aquatic and riparian reserves, will provide for many

specific elements, but research informs on situations to be wary of taking that assumption at face value without additional information and tests. Additional knowledge is needed especially on many late-successional and old-growth-associated species still known as rare or uncommon, or that are little-known—such as most soil and canopy invertebrates, fungi, and many amphibians—and on highly locally endemic species such as some aquatic mollusks, before fine-filter (species-specific) conservation measures can be confidently replaced by coarse-filter guidelines.

Issues of Species Conservation

As a program to replace or complement the Survey and Manage program under the NWFP, the ISSSSP—spanning the Forest Service Region 6 and BLM Oregon and Washington—has produced much material since the 2006 NWFP science synthesis on a wide variety of little-known or rare late-successional and old-growth-associated species. Their products pertain directly to contributing information on conservation and management of Survey and Manage species under the NWFP and sensitive species across Oregon and Washington. The products include basic ecological information, field inventories and monitoring results, protocols for surveys, and conservation guidelines. Ongoing local monitoring efforts at district levels will continue to inform on the status of little-known and rare late-successional and old-growth-associated species, including if and when individual species should be of additional conservation concern or are doing well.

The Survey and Manage program produced guidelines and approaches on natural history and management considerations for hidden, rare, or little-known species, including soil invertebrates and some lichens. Still, research is needed on their habitat associations, ecology, and degree of tolerance to disturbances of many rare and little-known, late-successional and old-growth-associated species of fungi, lichens, bryophytes, and vascular plant species.

Beyond the biological sciences, successfully conserving, restoring, and managing for other species under the NWFP will take additional attention to social needs and

interests (see chapters 8, 9, and 11). Fostering or retaining public support for conservation and restoration of rare and little-known species may require addressing social beliefs and values, and clarifying impacts and rationale of management policies, context, and actions (Stankey and Shindler 2006; see chapters 9, 10, and 11). Such was the case in the formulation of the initial NWFP guidelines, and such may be required again if the Plan is to be expanded to include other species, their habitats, or objectives beyond late-successional and old-growth forest, aquatic, and riparian environments.

Accounting for dynamic systems and uncertainties—

Recent science findings suggest that successfully managing for the future of other species under the NWFP needs to explicitly account for disturbance dynamics (Odion and Sarr 2007) in forest ecosystems of the Pacific Northwest, principally fire and anthropogenic changes to species' habitats in the managed forest matrix. Collectively, our findings suggest that such changes to forest age classes, structures, and dispersion patterns may have had greater impact on species metapopulation viability and on late-successional and old-growth forest resilience than initially thought by FEMAT, in large part because of increasing occurrence of high-severity fire in recent years.

Recent science findings also have highlighted the need to consider how continuing climate change will affect such disturbance dynamics and landscape-level connectivity of habitats for late-successional and old-growth-associated species. Researchers have suggested developing adaptation planning for climate change effects, in large part for (1) flexing and migrating boundaries of LSRs (as originally intended by FEMAT (1993), (2) conserving habitat in higher elevations, and (3) providing for site-specific conservation of late-successional and old-growth forest conditions within the forest matrix to serve as connections between the reserves. The federal matrix lands are not treated as they were when the NWFP was implemented; rather, late-successional and old-growth forests in the matrix lands are generally not being harvested at the rate they had been previously (see chapter 3), which may contribute to conservation of

old-forest habitat for some species, although the specific degree of contribution is unstudied. Also, the Aquatic Conservation Strategy, with its riparian buffers and headwater conservation approach, seems to be providing for nonfish aquatic species, particularly stream-associated amphibians and arthropods.

There will always be uncertainty on species' ecologies and conservation effectiveness. Several researchers have suggested use of decision science and risk analysis approaches to evaluating and managing for NWFP late-successional and old-growth-associated species under uncertainty (e.g., Kerns and Ager 2007). Such approaches might prove increasingly useful as we further enter into an era of climate change-induced disturbances, by presenting evaluations of potential changes and expected effects of alternative management actions as probabilities of outcomes in a risk analysis and risk management framework. Borrmann (2004) suggested a new approach to managing forests under uncertainty, which they termed "options forestry" that would entail diversifying management operations for the purpose of learning under an adaptive management framework. Franklin and Johnson (2012) suggested a restoration strategy ("ecological forestry;" see also Franklin et al. 2007, Hanson et al. 2012) for federal public forests in Washington and Oregon that would produce ecological and economic benefits by reserving older forest stands, thinning plantations to encourage complex vegetation structures, and conducting variable-retention harvests in younger forests to provide diverse early-seral environments.

Continuing the conservation of other species and biodiversity under the NWFP may benefit from use of decision science approaches including decision support models (Staus et al. 2010) and structured decisionmaking frameworks (Marcot et al. 2012b) that clearly articulate objectives; involve managers, decisionmakers, stakeholders, and analysts in all planning stages; and provide for monitoring and adaptive management to revisit, reaffirm, or revise management objectives and guidelines (Thompson et al. 2013), particularly under dynamics of changing social values, climates, and disturbance regimes (e.g., Jactel et al. 2012). The topic is explored more fully in chapter 12.

Coarse-filter, fine-filter, and landscape-scale management—

We have discussed recent findings on the use and testing of the coarse- and fine-filter approach to management, which is still the prevailing framework under the NWFP, the Survey and Manage standards and guidelines, and the ISSSSP.

The approach has been conflated with the use of a variety of species- and system-level approaches, including use of umbrella species, flagship species, management indicator species, focal species, surrogate species, guild indicators, indicator groups, and others (e.g., Lawler et al. 2003). The coarse- and fine-filter approach is a simple two-tier method of (1) providing general conditions, such as late-successional and old-growth forests in LSRs, and aquatic and riparian environments in the NWFP's Aquatic Conservation Strategy, and (2) then testing the efficacy for conservation of all species and biodiversity attributes of interest, and devising and implementing additional management activities and requirements as needed to provide for the full suite of species' habitats and biodiversity elements. What recent science has found is that the results—the combination of environmental conditions—will undoubtedly shift under climate change and disturbance dynamics, and thus might need to be at least intermittently reevaluated and readjusted.

Further, the coarse- and fine-filter approach focuses on environmental conditions and species' habitats except with the northern spotted owl and marbled murrelet, which include monitoring of population-level demographic status and trends. Whether the approach assures, to a degree acceptable to management, the long-term viability of all populations for the remaining 80 years of the NWFP is not known.

Recent studies and evaluations of how well the NWFP provides for other species and biodiversity also suggest that the role and efficacy of late-successional (and aquatic and riparian) reserves needs to be put into the broader context of the managed forest matrix and other systems of the Pacific Northwest found mainly on federal public lands, specifically subalpine forests, rock and ice substrates, headwaters, complex early-successional vegetation, and unharvested postfire conditions. However, maintaining late-successional and old-growth forest ecosystems and their biota may not be fully achievable only with old-forest

reserves, and only on federal forest lands, given effects of forest fragmentation and fire suppression and dynamics (e.g., Perault and Lomolino 2000, Sheehan et al. 2015). For these and other reasons, McAlpine et al. (2007) concluded that conservation of regional biodiversity under broad forest plans including the NWFP are better achieved with sustainable forest management practices implemented on—or at least coordinated across—all ownerships and by all stakeholders, not just for focused forest reserves. Social and economic considerations for any such forest management are discussed in chapters 8 and 12.

Other considerations—

Several other considerations may be pertinent and useful for managing for biodiversity and other species other than the northern spotted owl, marbled murrelet, and salmonids under the NWFP. One idea from recent science findings is the value of providing older-forest substrates and elements outside of LSRs that can contribute to species' habitats and habitat connections in the managed forest matrix (e.g., Dunk and Hawley 2009). An example pertains to dead wood. Approximately two-thirds of Survey and Manage species under the NWFP are associated in some way with, and benefit from mitigation for, partially dead trees, snags, and down wood (coarse woody debris). The Forest Service Decayed Wood Advisor (DecAID)⁴³ provides substantial information on wildlife use of snags, partially dead trees, and down wood, along with guidance on management of such elements at stand and landscape scales (Mellen et al. 2002).

Further, there may be an opportunity to bolster research, conservation, and restoration guidelines for providing such conditions within the managed forest matrix and across land ownerships. The objective would be to help connect LSRs, particularly now that late-successional and old-growth forests in the federal land matrix are currently being harvested at much less the rate they once were. In the past, thinning of plantations and young stands served to reduce dead wood, especially large down wood. Dead wood provides for a surprisingly vast array of life and its essential

⁴³ <http://www.fs.fed.us/r6/nr/wildlife/decaid/>; currently undergoing a major version update.

ecological processes (Brazee et al. 2014, Seibold et al. 2015). Providing more naturally regenerating, structurally and floristically complex early-successional stands, with the aim of encouraging development of dead wood and large down wood, may be part of restoration actions.

Protection of old-forest “legacies” such as large old green trees and snags and large-diameter down logs in the managed forest matrix has long been touted as a means of conserving biodiversity in Pacific Northwest old-growth forests following disturbances (Johnstone et al. 2016, North and Franklin 1990). Remnant, shade-tolerant old-growth trees (*Thuja* spp.) have been found to provide biological legacies and seed sources in postfire conditions in Pacific Northwest conifer forests (Keeton and Franklin 2005). However, retention of old-forest legacies may not fully substitute for retention of old-growth forest stands per se. For example, Price and Hochachka (2001) reported that epiphytic and alectoroid lichens were less abundant in mature (70- to 120-year-old) stands with structural retention of old-forest legacies, than in old-growth (at least 300-year-old) stands. The point here is that old-forest legacies retained in early-successional forests can complement but not fully supplant old-growth forests in providing habitat for old-forest species.

There may be considerations for tradeoffs among the kinds of species that can be provided under different silvicultural activities such as different intensities of forest thinning. For example, in the Pacific Northwest, Pollock and Beechie (2014) found that different sets of wildlife species associated with large-diameter live trees benefited more from heavy thinning in riparian areas than species associated with large-diameter deadwood benefited from light or no thinning. The authors suggested that because far more vertebrate species use large deadwood than large-diameter live trees, riparian areas may best be left to develop naturally in the absence of thinning for the benefit of terrestrial and aquatic species. Consideration for such tradeoffs of management objectives is discussed more fully in chapter 12.

Another consideration pertains to maintaining the full suite of ecological functions provided by late-successional and old-growth-associated species. This helps provide for “fully-functional” ecosystems, including full food webs and

ecosystem processes of all successional stages (Marcot 2002, Marcot and Sieg 2007, Marcot and Vander Heyden 2001).

Finally, there may be efficiencies and advantages to continue coordination of NWFP implementation and any amendments or updates, with other programs including climate science centers, disturbance research programs, and other protected area network programs.

Main Findings

Much has been learned since the 2006 synthesis (Haynes et al. 2006) about a wide variety of species and their occurrence, distribution, and rarity, and some on their ecology, reaffirming the role of LSRs and conservation of aquatic systems and riparian buffers in providing habitat for such species. Greater clarity also has been developed on the role of retaining old-forest components and substrates in the managed forest matrix to serve as connections among the reserves. Providing such connections is possible through the slowing of harvests of late-successional and old-growth forests in the federal matrix lands, the current critical habitat designation for northern spotted owls to protect all remaining owl habitat, and retention of old-forest structures in the forest matrix. Based on research on early-successional vegetation and associated biotic communities, how young stands in the forest matrix are managed will have a significant bearing on the future of older forest ecosystems throughout the NWFP area.

Recent work has also called for a far more dynamic approach to account for shifting movement barriers, fire disturbance to reserves and late-successional and old-growth forest stands, and other anthropogenic and climate-mediated changes in reserves and matrix lands alike. Habitats and population connections of other species of late-successional and old-growth forests are likely more vulnerable to the static placement of existing LSRs than previously envisioned. According to the research summarized in this chapter, a dynamic approach addressing the long-term scheduling of forest management activities and additions of reserves to further connect existing reserves and supplement them in higher elevations could better account for the influence of changing fire regimes, climate, and use of natural resources.

The Survey and Manage program, and more recently the ISSSSP, have provided much information on various rare and uncommon species associated with late-successional and old-growth environments. Their conservation evaluations provide evidence that the NWFP is generally providing habitat for late-successional and old-growth-associated species either under the late-successional, aquatic, and riparian buffer reserve system (coarse-filter management), or additionally through species-specific inventories, monitoring, and site protection (fine-filter management). Additional federal listing of late-successional and old-growth species as threatened or endangered, beyond northern spotted owl, marbled murrelet, and salmonids, has generally not proved necessary. Still, there may remain concerns for some species groups such as some late-successional and old-growth-associated bryophytes (Halpern et al. 2012) and other species groups because of the legacy of past forest harvesting patterns coupled with climate-change stressors (e.g., see section above on “Bryophytes”), but species-specific information is generally scant. For most of the other, rarer species addressed in this chapter, little to no information is available on population size and trend, especially their demographics across the managed forest matrix outside of reserves, although information is available on the general distribution or occurrence of some such species.

The Survey and Manage program, with its annual species reviews, provided the basis for reducing or removing concern for conservation of many uncommon to rare late-successional and old-growth-associated species that were originally ranked as having low potential for persistence. The lowering of conservation concern was due not so much to lower levels of harvest of older forests, but more to efforts locating such species during “pre-disturbance surveys” before local harvests and other management activities proceeded.

Since the 2006 synthesis (Haynes et al. 2006), no species have been added to the Survey and Manage species list; any additions would occur through a renewed annual species review process, and none was added during the reviews in 2001, 2002, and 2003. Those reviews resulted only in removing species from the list on the basis of new findings (viz., no concern for persistence, or the species

not being late-successional and old-growth associated) or changing their conservation and monitoring categories based on new information.

Other than the research summarized in this chapter, little effectiveness monitoring has been conducted on site buffers to confirm that they are indeed providing for species persistence at those sites. Limited research on riparian stream buffers suggest high value for protected associated species’ habitats.

Acknowledgments

We thank Rob Huff and Carol Hughes of the USDA Forest Service and USDI Bureau of Land Management’s ISSSSP for information on the program’s direction and content. We thank Daniel Luoma, Bruce McCune, and Tom Spies for helpful reviews of the manuscript. We also thank Jimmy Swingle and Eric Forsman for providing much information on tree voles. Dede Olson provided helpful comments on the reptiles and amphibians section. Diane Ikeda and Greg Schroer of Forest Service Region 5, Kim Mellen-McLean of Forest Service Region 6, and Carol Hughes provided very helpful management reviews. Anne Boeder provided information on the current status of resource management plans for BLM, and Mark Skinner provided information on regulations of harvests of special forest products. Coauthors K. Pope, H. Welsh, and C. Wheeler provided text on reptiles and amphibians; coauthor M. Reilly provided text on vascular plants; and coauthors K. Slauson and W. Zielinski provided text on mammalian carnivores.

Metric Equivalents

When you know:	Multiply by:	To find:
Inches	2.54	Centimeters
Feet (ft)	.305	Meters
Miles (mi)	1.609	Kilometers
Acres (ac)	.405	Hectares
Square miles (mi ²)	2.59	Square feet
Pounds (lbs)	.454	Kilograms
Degrees Fahrenheit (°F)	.56(°F – 32)	Degrees Celsius

Literature Cited

- Acorn, J.; Sheldon, I. 2001.** Bugs of Washington and Oregon. Edmonton, AB: Lone Pine Publishing. 160 p.
- Adams, M.J. 1999.** Correlated factors in amphibian decline: exotic species and habitat change in western Washington. *Journal of Wildlife Management*. 63(4): 1162–1171.
- Ager, A.A.; Finney, M.A.; Kerns, B.K.; Maffei, H. 2007.** Modeling wildfire risk to northern spotted owl (*Strix occidentalis caurina*) habitat in central Oregon, USA. *Forest Ecology and Management*. 246(1): 45–56. doi:10.1016/j.foreco.2007.03.070.
- Alexander, S.J.; Pilz, D.; Weber, N.S.; Brown, E.; Rockwell, V.A. 2002.** Mushrooms, trees, and money: value estimates of commercial mushrooms and timber in the Pacific Northwest. *Environmental Management*. 30(1): 129–141.
- Amaranthus, M.; Trappe, J.M.; Bednar, L.; Arthur, D. 1994.** Hypogeous fungal production in mature Douglas-fir forest fragments and surrounding plantations and its relation to coarse woody debris and animal mycophagy. *Canadian Journal Forest Research*. 24(11): 2157–2165.
- Anderson, P.D.; Larson, D.J.; Chan, S.S. 2007.** Riparian buffer and density management influences on microclimate of young headwater forests of western Oregon. *Forest Science*. 53(2): 254–269.
- Andruskiw, M.; Fryxell, J.M.; Thompson, I.D.; Baker, J.A. 2008.** Habitat-mediated variation in predation risk by the American marten. *Ecology*. 89(8): 2273–2280.
- Arbogast, B.S.; Schumacher, K.I.; Kerhoulas, N.J.; Bidlack, A.L.; Cook, J.A.; Kenagy, G.J. 2017.** Genetic data reveal a cryptic species of new world flying squirrel: *Glaucomys oregonensis*. *Journal of Mammalogy*. 98(4): 1027–1041. doi:10.1093/jmammal/gyx055.
- Armstrong, G.W.; Adamowicz, W.L.; Beck, J.A.; Cumming, S.G.; Schmiegelow, F.K.A. 2003.** Coarse filter ecosystem management in a nonequilibrating forest. *Forest Science*. 49(2): 209–223.
- Arsenault, A.; Goward, T. 2016.** Macrolichen diversity as an indicator of stand age and ecosystem resilience along a precipitation gradient in humid forests of inland British Columbia, Canada. *Ecological Indicators*. 69: 730–738.
- Arthur, J.L.; Camm, J.D.; Haight, R.G.; Montgomery, C.A.; Polasky, S. 2004.** Weighing conservation objectives: maximum expected coverage versus endangered species protection. *Ecological Applications*. 14(6): 1936–1945.
- Arthur, J.L.; Haight, R.G.; Montgomery, C.A.; Polasky, S. 2002.** Analysis of the threshold and expected coverage approaches to the probabilistic reserve site selection problem. *Environmental Modeling and Assessment*. 7(2): 81–89.
- Ashton, D.T.; Marks, S.B.; Welsh, H.H. 2006.** Evidence of continued effects from timber harvesting on lotic amphibians in redwood forests of northwestern California. *Forest Ecology and Management*. 221(1–3): 183–193.
- Aubry, K.B.; Lewis, J.C. 2003.** Extirpation and reintroduction of fishers (*Martes pennanti*) in Oregon: implications for their conservation in the Pacific states. *Biological Conservation*. 114: 79–90.
- Aubry, K.B.; McKelvey, K.S.; Copeland, J.P. 2007.** Distribution and broadscale habitat relations of the wolverine in the contiguous United States. *Journal of Wildlife Management*. 71(7): 2147. doi:10.2193/2006–548.

- Aubry, K.B.; Raley, C.M. 2002a.** The pileated woodpecker as a keystone habitat modifier in the Pacific Northwest. In: Laudenslayer, W.F., Jr.; Shea, P.J.; Valentine, B.E.; Weatherspoon, C.P.; Lisle, T.E., eds. Proceedings of the symposium on the ecology and management of dead wood in western forests. Gen. Tech. Rep. PSW-GTR-181. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 257–274.
- Aubry, K.B.; Raley, C.M. 2002b.** Selection of nest and roost trees by pileated woodpeckers in coastal forests of Washington. *Journal of Wildlife Management*. 66(2): 392–406.
- Aubry, K.B.; Raley, C.M.; Buskirk, S.W.; Zielinski, W.J.; Schwartz, M.K.; Golightly, R.T.; Purcell, K.L.; Weir, R.D.; Yaeger, J.S. 2013.** Meta-analyses of habitat selection by fishers at resting sites in the Pacific coastal region. *Journal of Wildlife Management*. 77(5): 965–974. doi:10.1002/jwmg.563.
- Bagne, K.E.; Friggens, M.M.; Coe, S.J.; Finch, D.M. 2014.** The importance of assessing climate change vulnerability to address species conservation. *Journal of Fish and Wildlife Management*. 5(2): 450–462.
- Bagne, K.E.; Friggens, M.M.; Finch, D.M. 2011.** A system for assessing vulnerability of species (SAVS) to climate change. Gen. Tech. Rep. RMRS-GTR-257. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 28 p.
- Bailey, J.D.; Mayrsohn, C.; Doescher, P.S.; St. Pierre, E.; Tappeiner, J.C. 1998.** Understory vegetation in old and young Douglas-fir forests of western Oregon. *Forest Ecology and Management*. 112(3): 289–302. doi:10.1016/s0378-1127(98)00408-3.
- Ban, N.C.; Mills, M.; Tam, J.; Hicks, C.C.; Klain, S.; Stoeckl, N.; Bottrill, M.C.; Levine, J.; Pressey, R.L.; Satterfield, T.; Chan, K.M.A. 2013.** A social-ecological approach to conservation planning: embedding social considerations. *Frontiers in Ecology and the Environment*. 11(4): 194–202.
- Barua, M. 2011.** Mobilizing metaphors: the popular use of keystone, flagship and umbrella species concepts. *Biodiversity and Conservation*. 20(7): 1427–1440.
- Bednarz, J.C.; Huss, M.J.; Benson, T.J.; Varland, D.E. 2013.** The efficacy of fungal inoculation of live trees to create wood decay and wildlife-use trees in managed forests of western Washington, USA. *Forest Ecology and Management*. 307: 186–195.
- Beebe, S.B. 1991.** Conservation in temperate and tropical rain forests: the search for an ecosystem approach to sustainability. *North American Wildlife and Natural Resources Conference*. 56: 595–603.
- Beier, P.; Brost, B. 2010.** Use of land facets to plan for climate change: conserving the arenas, not the actors. *Conservation Biology*. 24(3): 701–710.
- Beier, P.; Majka, D.R.; Spencer, W.D. 2008.** Forks in the road: choices in procedures for designing wildlife linkages. *Conservation Biology*. 22(4): 836–851.
- Beier, P.; Spencer, W.; Baldwin, R.; McRae, B. 2011.** Toward best practices for developing regional connectivity maps. *Conservation Biology*. 25(5): 879–892.
- Bellinger, M.R.; Haig, S.M.; Forsman, E.D.; Mullins, T.D. 2005.** Taxonomic relationships among *Phenacomys* voles as inferred by cytochrome b. *Journal of Mammalogy*. 86(1): 201–210. doi:10.1644/1545-1542(2005)086<0201:trapva>2.0.co;2.
- Berg, B.; Laskowski, R. 2005.** Decomposers: soil microorganisms and animals. *Advances in Ecological Research*. 38: 73–100.
- Bernardo, J.; Spotila, J.R. 2006.** Physiological constraints on organismal response to global warming: mechanistic insights from clinally varying populations and implications for assessing endangerment. *Biology Letters*. 2(1): 135–139. doi:10.1098/rsbl.2005.0417.
- Best, M.L.; Welsh, H.H., Jr. 2014.** The trophic role of a forest salamander: impacts on invertebrates, leaf litter retention, and the humification process. *Ecosphere*. 5(art16): <http://dx.doi.org/10.1890/ES13-00302.1>.

- Betts, M.G.; Hagar, J.C.; Rivers, J.W.; Alexander, J.D.; McGarigal, K.; McComb, B.C. 2010.** Thresholds in forest bird occurrence as a function of the amount of early-seral broadleaf forest at landscape scales. *Ecological Applications*. 20: 2116–2130.
- Bifulchi, A.; Lodé, T. 2005.** Efficiency of conservation shortcuts: an investigation with otters as umbrella species. *Biological Conservation*. 126(4): 523–527.
- Blaustein, A.R.; Walls, S.C.; Bancroft, B.A.; Lawler, J.J.; Searle, C.L.; Gervasi, S.S. 2010.** Direct and indirect effects of climate change on amphibian populations. *Diversity*. 2(2): 281–313. doi:10.3390/d2020281.
- Blois, J.L.; Arbogast, B.S. 2006.** Conservation genetics of the Sonoma tree vole (*Arborimus pomo*) based on mitochondrial and amplified fragment length polymorphism markers. *Journal of Mammalogy*. 87: 950–960.
- Blouin, M.S.; Phillipsen, I.C.; Monsen, K.J. 2010.** Population structure and conservation genetics of the Oregon spotted frog, *Rana pretiosa*. *Conservation Genetics*. 11(6): 2179–2194.
- Bohlen, P.J.; Groffman, P.M.; Fahey, T.J.; Fisk, M.C.; Suarez, E.; Pelletier, D.M.; Fahey, R.T. 2004a.** Ecosystem consequences of exotic earthworm invasion of north temperate forests. *Ecosystems*. 7: 1–12.
- Bohlen, P.J.; Scheu, S.; Hale, C.M.; McLean, M.A.; Migge, S.; Groffman, P.M.; Parkinson, D. 2004b.** Non-native invasive earthworms as agents of change in northern temperate forests. *Frontiers in Ecology and the Environment*. 2(8): 427–435.
- Bokhorst, S.; Asplund, J.; Kardol, P.; Wardle, D.A. 2015.** Lichen physiological traits and growth forms affect communities of associated invertebrates. *Ecology*. 96(9): 2394–2407.
- Bonnot, T.W.; Millsaugh, J.J.; Rumble, M.A. 2009.** Multi-scale nest-site selection by black-backed woodpeckers in outbreaks of mountain pine beetles. *Forest Ecology and Management*. 259(2): 220–228. doi:10.1016/j.foreco.2009.10.021.
- Bormann, B.T.; Darbyshire, R.L.; Homann, P.S.; Morrisette, B.A.; Little, S.N. 2015.** Managing early succession for biodiversity and long-term productivity of conifer forests in southwestern Oregon. *Forest Ecology and Management*. 340: 114–125. doi:10.1016/j.foreco.2014.12.016.
- Bormann, B.T.; Kiester, A.R. 2004.** Options forestry: acting on uncertainty. *Journal of Forestry*. 102: 22–27.
- Bosakowski, T. 1999.** Amphibian macrohabitat associations on a private industrial forest in western Washington. *Northwestern Naturalist*. 80: 61–69.
- Brady, M.J.; McAlpine, C.A.; Miller, C.J.; Possingham, H.P.; Baxter, G.S. 2009.** Habitat attributes of landscape mosaics along a gradient of matrix development intensity: matrix management. *Landscape Ecology*. 24(7): 879–891.
- Brand, R.H. 2002.** The effect of prescribed burning on epigeic springtails (Insecta: Collembola) of woodland. *American Midland Naturalist*. 148(2): 383–393.
- Branton, M.; Richardson, J.S. 2010.** Assessing the value of umbrella-species concept for conservation planning with meta-analysis. *Conservation Biology*. 25(1): 9–20.
- Branton, M.A.; Richardson, J.S. 2014.** A test of the umbrella species approach in restored floodplain ponds. *Journal of Applied Ecology*. 51(3): 776–785.
- Brazee, N.J.; Lindner, D.L.; D’Amato, A.W.; Fraver, S.; Forrester, J.A.; Mladenoff, D.J. 2014.** Disturbance and diversity of wood-inhabiting fungi: effects of canopy gaps and downed woody debris. *Biodiversity and Conservation*. 23(9): 2155–2172.

- Breckheimer, I.; Haddad, N.M.; Morris, W.F.; Trainor, A.M.; Fields, W.R.; Jobe, R.T.; Hudgens, B.R.; Moody, A.; Walters, J.R. 2014.** Defining and evaluating the umbrella species concept for conserving and restoring landscape connectivity. *Conservation Biology*. 28(6): 1584–1593.
- Brost, B.M.; Beier, P. 2012.** Use of land facets to design linkages for climate change. *Ecological Applications*. 22(1): 87–103.
- Brumwell, L.J.; Craig, K.G.; Scudder, G.G.E. 1998.** Litter spiders and carabid beetles in a successional Douglas-fir forest in British Columbia. *Northwest Science*. 72(Special issue): 94–95.
- Buchalski, M.R.; Fontaine, J.B.; Heady, P.A.I.; Hayes, J.P.; Frick, W.F. 2013.** Bat response to differing fire severity in mixed-conifer forest California, USA. *PLoS ONE*. 8(3): e57884.
- Bull, E.L.; Heather, T.W. 2001.** Survival, causes of mortality, and reproduction in the American marten in northeastern Oregon. *Northwestern Naturalist*. 82: 1–6.
- Bull, J.C.; Stopher, M.; Williams, D.R.; Morefield, K.; Croteau, J.M. 2006.** Report to the California Fish and Game Commission: status review of Siskiyou mountains salamander (*Plethodon stormi*) in California. Status Report 2006-01. [Place of publication unknown]: California Department of Fish and Game, Habitat Conservation Planning Branch. 54 p.
- Bunnell, F.L. 1995.** Forest-dwelling vertebrate faunas and natural fire regimes in British Columbia: patterns and implications for conservation. *Conservation Biology*. 9(3): 636–644.
- Burton, J.I.; Olson, D.H.; Puettmann, K.J. 2016.** Effects of riparian buffer width on wood loading in headwater streams after repeated forest thinning. *Forest Ecology and Management*. 372: 247–257.
- Bury, R.B. 2008.** Low thermal tolerances of stream amphibians in the Pacific Northwest: implications for riparian and forest management. *Applied Herpetology*. 5(1): 63–74. doi:10.1163/157075408783489211.
- Buse, J. 2012.** “Ghosts of the past”: flightless saproxylic weevils (Coleoptera: Curculionidae) are relict species in ancient woodlands. *Journal of Insect Conservation*. 16(1): 93–102.
- Busse, M.D.; Simon, S.A.; Riegel, G.M. 2000.** Tree-growth and understory responses to low-severity prescribed burning in thinned *Pinus ponderosa* forests of central Oregon. *Forest Science*. 46: 258–268.
- Caesar, R.M.; Gillette, N.; Cognato, A.I. 2005.** Population genetic structure of an edaphic beetle (Ptiliidae) among late-successional reserves within the Klamath-Siskiyou ecoregion, California. *Annals of the Entomological Society of America*. 98(6): 931–940.
- Caplat, P.; Edelaar, P.; Dudaniec, R.Y.; Green, A.J.; Okamura, B.; Cote, J.; Ekroos, J.; Jonsson, P.R.; Londahl, J.; Tesson, S.V.M.; Petit, E.J. 2016.** Looking beyond the mountain: dispersal barriers in a changing world. *Frontiers in Ecology and Evolution*. 14(5): 261–268.
- Carey, A.B. 1991.** The biology of arboreal rodents in Douglas-fir forests. Gen. Tech. Rep. PNW-GTR-276. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 46 p.
- Carey, A.B. 1995.** Sciurids in Pacific Northwest managed and old-growth forests. *Ecological Applications*. 5(3): 648–661.
- Carey, A.B.; Johnson, M.L. 1995.** Small mammals in managed, naturally young, and old-growth forests. *Ecological Applications*. 5(2): 336–352.
- Carey, C.; Alexander, M.A. 2003.** Climate change and amphibian declines: Is there a link? *Diversity and Distributions*. 9(2): 111–121. doi:10.1046/j.1472-4642.2003.00011.x.
- Carr, S.M.; Hicks, S.A. 1997.** *Martes*: taxonomy, ecology, techniques, and management. In: Bryant, H.N.; Proulx, G.; Woodard, P.M., eds. *Proceedings of the second international Martes symposium*. Edmonton, Alberta, Canada: Provincial Museum of Alberta: 15–28. <http://dx.doi.org/10.5962/bhl.title.102822>. (4 December 2017).

- Carroll, C. 2010.** Role of climatic niche models in focal-species-based conservation planning: assessing potential effects of climate change on northern spotted owl in the Pacific Northwest, USA. *Biological Conservation*. 143(6): 1432–1437. doi:10.1016/j.biocon.2010.03.018.
- Carroll, C.; Dunk, J.R.; Moilanen, A. 2010.** Optimizing resiliency of reserve networks to climate change: multispecies conservation planning in the Pacific Northwest, USA. *Global Change Biology*. 16(3): 891–904.
- Carroll, C.; Zielinski, W.J.; Noss, R.F. 1999.** Using presence-absence data to build and test spatial habitat models for the fisher in the Klamath region, U.S.A. *Conservation Biology*. 13(6): 1344–1359.
- Caruso, N.; Guerisoli, M.; Vidal, E.M.L.; Castillo, D.; Casanave, E.B.; Lucherini, M. 2015.** Modelling the ecological niche of an endangered population of puma concolor: first application of the GNESFA method to an elusive carnivore. *Ecological Modelling*. 297: 11–19.
- Castellano, M.A.; Cazares, E.; Fondrick, B.; Dreisbach, T. 2003.** Handbook to additional fungal species of special concern in the Northwest Forest Plan. Gen. Tech. Rep. PNW-GTR-572. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 144 p.
- Chen, Y.; Seybold, S.J. 2014.** Crepuscular flight activity of an invasive insect governed by interacting abiotic factors. *PLoS ONE*. 9(8) doi:10.1371/journal.pone.0105945.
- Chinnici, S.J.; Bigger, D.; Johnson, E. 2012.** Sonoma tree vole habitat on managed redwood and Douglas-fir forestlands in north coastal California. In: Standiford, R.B.; Weller, T.J.; Piirto, D.D.; Stuart, J.D., eds. *Proceedings of the coast redwood forests in a changing California: a symposium for scientists and managers*. Gen. Tech. Rep. PSW-GTR-238. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 389–397.
- Choi, S.W.; Miller, J.C. 2013.** Species richness and abundance among macromoths: a comparison of taxonomic, temporal and spatial patterns in Oregon and South Korea. *Entomological Research*. 43: 312–321.
- Cissel, J.H.; Anderson, P.D.; Olson, D.; Puettmann, K.J.; Berryman, S.; Chan, S.; Thompson, C. 2006.** BLM density management and riparian buffer study: establishment report and study plan. Scientific Investigations Report 2006-5087. Reston, VA: U.S. Department of the Interior, Geological Survey. 151 p.
- Clarkson, D.A.; Mills, L.S. 1994.** Hypogeous sporocarps in forest remnants and clearcuts in southwest Oregon. *Northwest Science*. 68(4): 259–265.
- Clayton, D.R.; Olson, D.H. 2009.** Conservation assessment for the Oregon slender salamander (*Batrachoseps wrighti*). USDA Forest Service Region 6 and USDI Bureau of Land Management. 76 p. <http://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-amphibians.shtml>. (4 December 2017).
- Clayton, D.R.; Olson, D.H.; Nauman, R.S. 2005.** Conservation assessment for the Siskiyou mountains salamander (*Plethodon stormi*). USDA Forest Service Region 6 and USDI Bureau of Land Management, Oregon and Washington. 32 p. <http://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-amphibians.shtml>. (4 December 2017).
- Collier, M.J. 2015.** Novel ecosystems and social-ecological resilience. *Landscape Ecology*. 30(8): 1363–1369.
- Colwell, R.K. 2013.** EstimateS: statistical estimation of species richness and shared species from samples. Version 9 user's guide and application. <http://purl.oclc.org/estimates>. (4 December 2017).
- Copeland, J.P. 1996.** Biology of the wolverine in central Idaho. Moscow, ID: University of Idaho. 152 p. Ph.D. dissertation.

- Copeland, J.P.; McKelvey, K.S.; Aubry, K.B.; Landa, A.; Persson, J.; Inman, R.M.; Krebs, J.; Lofroth, E.; Golden, H.; Squires, J.R.; Magoun, A.; Schwartz, M.K.; Wilmot, J.; Copeland, C.L.; Yates, R.E.; Kojola, I.; May, R. 2010.** The bioclimatic envelope of the wolverine (*Gulo gulo*): Do climatic constraints limit its geographic distribution? *Canadian Journal of Zoology*. 88(3): 233–246. doi:10.1139/z09-136.
- Copeland, J.P.; Peek, J.M.; Groves, C.R.; Melquist, W.E.; McKelvey, K.S.; McDaniel, G.W.; Long, C.D.; Harris, C.E. 2007.** Seasonal habitat associations of the wolverine in central Idaho. *Journal of Wildlife Management*. 71(7): 2201–2212.
- Corn, P.S. 2003.** Amphibian breeding and climate change: importance of snow in the mountains. *Conservation Biology*. 17(2): 622–625. doi:10.1046/j.1523-1739.2003.02111.x.
- Corn, P.S. 2005.** Climate change and amphibians. *Animal Biodiversity and Conservation*. 28(1): 59–67.
- Corn, P.S.; Bury, R.B. 1989.** Logging in western Oregon: responses of headwater habitats and stream amphibians. *Forest Ecology and Management*. 29(1–2): 39–57.
- Cotsell, N.; Vernes, K. 2016.** Camera traps in the canopy: surveying wildlife at tree hollow entrances. *Pacific Conservation Biology*. 22(1): 48–60.
- Creutzburg, M.K.; Scheller, R.M.; Lucash, M.S.; Leduc, S.D.; Johnson, M.G. 2017.** Forest management scenarios in a changing climate: trade-offs between carbon, timber, and old forest. *Ecological Applications*. 27(2): 503–518. doi:10.1002/eap.1460.
- Crimmins, S.M.; Dobrowski, S.Z.; Greenberg, J.A.; Abatzoglou, J.T.; Mynsberge, A.R. 2011.** Changes in climatic water balance drive downhill shifts in plant species' optimum elevations. *Science*. 331(6015): 324–327. doi:10.1126/science.1199040.
- Crisafulli, C.M.; Clayton, D.R.; Olson, D.H. 2008.** Conservation assessment for the Larch Mountain salamander (*Plethodon larselli*). USDA Forest Service Region 6 and USDI Bureau of Land Management Interagency Special Status and Sensitive Species Program. 36 p. <http://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-amphibians.shtml>. (22 February 2018).
- Crooks, K.R.; Burdett, C.L.; Theobald, D.M.; Rondinini, C.; Boitani, L. 2011.** Global patterns of fragmentation and connectivity of mammalian carnivore habitat. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 366(1578): 2642–2651.
- Crow, T. 1990.** Old growth and biological diversity: a basis for sustainable forestry. In: Perera, A.H.; Euler, D.L.; Thompson, I.D., eds. *Old growth forests: What are they? How do they work?* Proceedings from the conference on old-growth forests. Toronto, ON: Canadian Scholars' Press Inc.: 49–62.
- Cunningham, R.B.; Lindenmayer, D.B. 2005.** Modeling count data of rare species: some statistical issues. *Ecology*. 86(5): 1135–1142.
- Cushman, S.A.; McKelvey, K.S.; Flather, C.H.; McGarigal, K. 2008.** Do forest community types provide a sufficient basis to evaluate biological diversity? *Frontiers in Ecology and the Environment*. 6(1): 13–17. doi:10.1890/070039.
- Dávalos, A.; Simpson, E.H.; Nuzzo, V.; Blossey, B. 2015.** Non-consumptive effects of native deer on introduced earthworm abundance. *Ecosystems*. 18(6): 1029–1042.
- Davidson, C.; Bradley Shaffer, H.; Jennings, M.R. 2001.** Declines of the California red-legged frog: climate, UV-B, habitat, and pesticides hypotheses. *Ecological Applications*. 11(2): 464–479. doi:10.1890/1051-0761(2001)011[0464:dotcrl]2.0.co;2.
- Davidson, C.; Shaffer, H.B.; Jennings, M.R. 2002.** Spatial tests of the pesticide drift, habitat destruction, UV-B, and climate-change hypotheses for California amphibian declines. *Conservation Biology*. 16(6): 1588–1601. doi:10.1046/j.1523-1739.2002.01030.x.

- Davis, F.W.; Seo, C.; Zielinski, W.J. 2007.** Regional variation in home-range-scale habitat models for fisher (*Martes pennanti*) in California. *Ecological Applications*. 17(8): 2195–2213. doi:10.1890/06-1484.1.
- Davis, R.J.; Ohmann, J.L.; Kennedy, R.E.; Cohen, W.B.; Gregory, M.J.; Yang, Z.; Roberts, H.M.; Gray, A.N.; Spies, T.A. 2015.** Northwest Forest Plan—the first 20 years (1994–2013): status and trends of late-successional and old-growth forests. Gen. Tech. Rep. PNW-GTR-911. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 112 p. doi:10.2737/pnw-gtr-911.
- Dawson, F.N.; Magoun, A.J.; Bowman, J.; Ray, J.C. 2010.** Wolverine, *Gulo gulo*, home range size and denning habitat in lowland boreal forest in Ontario. *Canadian Field Naturalist*. 124(2): 139–144.
- Dawson, N.G.; Cook, J.A. 2012.** Behind the genes: diversification of north American martens (*Martes americana* and *M. caurina*). In: Aubry, K.B.; Zielinski, W.J.; Raphael, M.G.; Proulx, G.; Buskirk, S.W., eds. *Biology and conservation of martens, sables, and fishers: a new synthesis*. Ithaca, NY: Cornell University Press: 23–38.
- De León, M.E.; Vredenburg, V.T.; Piovia-Scott, J. 2017.** Recent emergence of a chytrid fungal pathogen in California Cascades frogs (*Rana cascadae*). *EcoHealth*. 14(1): 155–161. doi:10.1007/s10393-016-1201-1.
- Decourtye, A.; Mader, E.; Desneux, N. 2010.** Landscape enhancement of floral resources for honey bees in agro-ecosystems. *Apidologie*. 41: 264–277.
- Degross, D.J.; Bury, R.B. 2007.** Science review for the Scott Bar salamander (*Plethodon asupak*) and the Siskiyou mountains salamander (*P. stormi*): biology, taxonomy, habitat, and detection probabilities/occupancy. Open-File Report 2007-1352. Reston, VA: U.S. Department of the Interior, Geological Survey. 14 p.
- D’Elia, J.; Haig, S.M.; Johnson, M.; Marcot, B.G.; Young, R. 2015.** Activity-specific ecological niche models for planning reintroductions of California condors (*Gymnogyps californianus*). *Biological Conservation*. 184: 90–99.
- Diaz, N.M.; Haynes, R.W. 2002.** Highlights of science contributions to implementing the Northwest Forest Plan—1994 to 1998. Gen. Tech. Rep. PNW-GTR-540. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 22 p. doi:10.2737/pnw-gtr-540.
- Dodson, E.K.; Peterson, D.W.; Harrod, R.J. 2008.** Understory vegetation response to thinning and burning restoration treatments in dry conifer forests of the eastern Cascades, USA. *Forest Ecology and Management*. 255(8–9): 3130–3140. doi:10.1016/j.foreco.2008.01.026.
- Donato, D.C.; Fontaine, J.B.; Robinson, W.D.; Kauffman, J.B.; Law, B.E. 2009.** Vegetation response to a short interval between high-severity wildfires in a mixed-evergreen forest. *Journal of Ecology*. 97(1): 142–154. doi:10.1111/j.1365-2745.2008.01456.x.
- Driscoll, D.A.; Weir, T. 2005.** Beetle responses to habitat fragmentation depend on ecological traits, habitat condition, and remnant size. *Conservation Biology*. 19(1): 182–194.
- Drouin, M.; Bradley, R.; Lapointe, L.; Whalen, J. 2014.** Non-native anecic earthworms (*Lumbricus terrestris* L.) reduce seed germination and seedling survival of temperate and boreal trees species. *Applied Soil Ecology*. 75: 145–149.
- Duchamp, J.E.; Arnett, E.; Larson, M.; Swihart, R.K. 2007.** Ecological considerations for landscape-level management of bats. In: Lacki, M.J.; Hayes, J.P.; Kurta, A., eds. *Bats in forests: conservation and management*. Baltimore, MD: Johns Hopkins University Press: 237–261.

- Duhamel, M.; Pel, R.; Ooms, A.; Bücking, H.; Jansa, J.; Ellers, J.; Van Straalen, N.M.; Wouda, T.; Vandenkoornhuyse, P.; Kiers, E.T. 2013.** Do fungivores trigger the transfer of protective metabolites from host plants to arbuscular mycorrhizal hyphae? *Ecology*. 94(9): 2019–2029.
- Dunk, J.R.; Hawley, J.J.V.G. 2009.** Red-tree vole habitat suitability modeling: implications for conservation and management. *Forest Ecology and Management*. 258(5): 626–634. doi:10.1016/j.foreco.2009.04.041.
- Dunk, J.R.; Zielinski, W.J.; Preisler, H.K. 2004.** Predicting the occurrence of rare mollusks in northern California forests. *Ecological Applications*. 14(3): 713–729.
- Edwards, T.C.; Cutler, D.R.; Geiser, L.; Alegria, J.; McKenzie, D. 2004.** Assessing rarity of species with low detectability: lichens in Pacific Northwest forests. *Ecological Applications*. 14(2): 414–424. doi:10.1890/02-5236.
- Edwards, T.C.; Cutler, D.R.; Zimmermann, N.E.; Geiser, L.; Alegria, J. 2005.** Model-based stratifications for enhancing the detection of rare ecological events. *Ecology*. 86(5): 1081–1090. doi:10.1890/04-0608.
- Eggers, B.; Matern, A.; Drees, C.; Eggers, J.; Hardtle, W.; Assmann, T. 2010.** Value of semi-open corridors for simultaneously connecting open and wooded habitats: a case study with ground beetles. *Conservation Biology*. 24(1): 256–266.
- Eisenhauer, N.; Scheu, S. 2008.** Invasibility of experimental grassland communities: the role of earthworms, plant functional group identity and seed size. *Oikos*. 117(7): 1026–1036.
- Ellis, C.J. 2015.** Ancient woodland indicators signal the climate change risk for dispersal-limited species. *Ecological Indicators*. 53: 106–114.
- Ellis, M.M.; Ivan, J.S.; Schwartz, M.K. 2014.** Spatially explicit power analyses for occupancy-based monitoring of wolverine in the US Rocky Mountains. *Conservation Biology*. 28(1): 52–62.
- Emel, S.L.; Storfer, A. 2015.** Landscape genetics and genetic structure of the southern torrent salamander, *Rhyacotriton variegatus*. *Conservation Genetics*. 16: 209–221.
- Esslinger, T.L. 2015.** A cumulative checklist for the lichen-forming lichenicolous and allied fungi of the continental United States and Canada. Fargo, ND: North Dakota State University. <http://www.ndsu.edu/pubweb/~esslinge/chcklst/chcklst7.htm>. (19 April 2015).
- Ewing, H.A.; Tuininga, A.R.; Groffman, P.M.; Weathers, K.C.; Fahey, T.J.; Fisk, M.C.; Bohlen, P.J.; Suarez, E. 2015.** Earthworms reduce biotic 15-nitrogen retention in northern hardwood forests. *Ecosystems*. 18(2): 328–342.
- Fagan, W.F.; Kareiva, P.M. 1997.** Using compiled species lists to make biodiversity comparisons among regions: a test case using Oregon butterflies. *Biological Conservation*. 80: 249–259.
- Feldman, C.R.; Hoyer, R.F. 2010.** A new species of snake in the genus *Contia* (Squamata: Colubridae) from California and Oregon. *Copeia*. 2010(2): 254–267. doi:10.1643/ch-09-129.
- Fisher, R.N.; Shaffer, H.B. 1996.** The decline of amphibians in California's Great Central Valley. *Conservation Biology*. 10(5): 1387–1397.
- Fontaine, J.B.; Donato, D.C.; Robinson, W.D.; Law, B.E.; Kauffman, J.B. 2009.** Bird communities following high-severity fire: response to single and repeat fires in a mixed-evergreen forest, Oregon, USA. *Forest Ecology and Management*. 257(6): 1496–1504. doi:10.1016/j.foreco.2008.12.030.
- Forest Ecosystem Management Assessment Team. [FEMAT]. 1993.** Forest ecosystem management: an ecological, economic, and social assessment. Portland, OR: U.S. Department of Agriculture; U.S. Department of the Interior [and others]. [Irregular pagination].

- Forister, M.L.; McCall, A.C.; Sanders, N.J.; Fordyce, J.A.; Thorne, J.H.; O'Brien, J.; Waetjen, D.P.; Shapiro, A.M. 2010.** Compounded effects of climate change and habitat alteration shift patterns of butterfly diversity. *Proceedings of the National Academy of Sciences of the United States of America*. 107(5): 2088–2092. doi:10.1073/pnas.0909686107.
- Forsman, E.D.; Anthony, R.G.; Zabel, C.J. 2004.** Distribution and abundance of red tree voles in Oregon based on occurrence in pellets of northern spotted owls. *Northwest Science*. 78(4): 294–302.
- Forsman, E.D.; Swingle, J.K.; Davis, R.J.; Biswell, B.L.; Andrews, L.S. 2016.** Tree voles: an evaluation of their distribution and habitat relationships based on recent and historical studies, habitat models, and vegetation change. Gen. Tech. Rep. PNW-GTR-948. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 119 p.
- Forsman, E.D.; Swingle, J.K.; Hatch, N.R. 2009.** Behavior of red tree voles (*Arborimus longicaudus*) based on continuous video monitoring of nests. *Northwest Science*. 83(3): 262–272.
- Foster, A.D.; Claeson, S.M. 2011.** Habitats and seasonality of riparian-associated millipedes in southwest Washington, USA. *Terrestrial Arthropod Reviews*. 4(3): 203–220. doi:10.1163/187498311x591102.
- Franklin, J.F. 1993.** Preserving biodiversity: species, ecosystems, or landscapes? *Ecological Applications*. 3(2): 202–205.
- Franklin, J.F.; Johnson, K.N. 2012.** A restoration framework for federal forests in the Pacific Northwest. *Journal of Forestry*. 110(8): 429–439. doi:10.5849/jof.10-006.
- Franklin, J.F.; Mitchell, R.J.; Palik, B.J. 2007.** Natural disturbance and stand development principles for ecological forestry. Gen. Tech. Rep. NRS-GTR-19. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 44 p. doi:10.2737/nrs-gtr-19.
- Frest, T.J.; Johannes, E.J. 1999.** Field guide to survey and manage freshwater mollusk species. Portland, OR: U.S. Department of the Interior, Fish and Wildlife Service, Regional Ecosystem Office; and Bureau of Land Management Oregon State Office. 117 p.
- Gabriel, M.W.; Woods, L.W.; Poppenga, R.; Sweitzer, R.A.; Thompson, C.; Matthews, S.M.; Higley, J.M.; Keller, S.M.; Purcell, K.; Barrett, R.H.; Wengert, G.M. 2012.** Anticoagulant rodenticides on our public and community lands: spatial distribution of exposure and poisoning of a rare forest carnivore. *PLoS ONE*. 7(7): e40163. doi:10.1371/journal.pone.0040163.
- Gabriel, M.W.; Woods, L.W.; Wengert, G.M.; Stephenson, N.; Higley, J.M.; Thompson, C.; Matthews, S.M.; Sweitzer, R.A.; Purcell, K.; Barrett, R.H.; Keller, S.M.; Gaffney, P.; Jones, M.; Poppenga, R.; Foley, J.E.; Brown, R.N.; Clifford, D.L.; Sacks, B.N. 2015.** Patterns of natural and human-caused mortality factors of a rare forest carnivore, the fisher (*Pekania pennanti*) in California. *PLOS ONE*. 10(11): e0140640. doi:10.1371/journal.pone.0140640.
- Garcia, T.; Rowe, J.; Doyle, J. 2015.** A tad too high: sensitivity to UV-B radiation may limit invasion potential of American bullfrogs (*Lithobates catesbeianus*) in the Pacific Northwest invasion range. *Aquatic Invasions*. 10(2): 237–247. doi:10.3391/ai.2015.10.2.12.
- Garcia, T.S.; Paoletti, D.J.; Blaustein, A.R. 2009.** Correlated trait responses to multiple selection pressures in larval amphibians reveal conflict avoidance strategies. *Freshwater Biology*. 54(5): 1066–1077. doi:10.1111/j.1365-2427.2008.02154.x.
- Geiser, L.H.; Neitlich, P.N. 2007.** Air pollution and climate gradients in western Oregon and Washington indicated by epiphytic macrolichens. *Environmental Pollution*. 145: 203–218.
- Gibbs, J.P. 1998.** Amphibian movements in response to forest edges, roads, and streambeds in southern New England. *Journal of Wildlife Management*. 62(2): 584–589.

- Glavich, D.A. 2009.** Distribution, rarity and habitats of three aquatic lichens on federal land in the U.S. Pacific Northwest. *The Bryologist*. 112(1): 54–72. doi:10.1639/0007-2745-112.1.54.
- Glavich, D.A.; Geiser, L.H. 2004.** *Dermatocarpon meiophyllizum* Vainio in the US Pacific Northwest. *Evansia*. 21(3): 137–140.
- Govindarajulu, P.; Isaac, L.A.; Engelstoft, C.; Ovaska, K. 2011.** Relevance of life-history parameter estimation to conservation listing: case of the sharp-tailed snake (*Contia tenuis*). *Journal of Herpetology*. 45(3): 300–307.
- Grant, E.H.C.; Muths, E.L.; Katz, R.A.; Canessa, S.; Adams, M.J.; Ballard, J.R.; Berger, L.; Briggs, C.J.; Coleman, J.; Gray, M.J.; Harris, M.C.; Harris, R.N.; Hossack, B.R.; Huyvaert, K.P.; Kolby, J.E.; Lips, K.R.; Lovich, R.E.; McCallum, H.I.; Mendelson, J.R.; Nanjappa, P.; Olson, D.H.; Powers, J.G.; Richgels, K.L.D.; Russell, R.E.; Schmidt, B.R.; Spitzen-Van Der Sluijs, A.; Watry, M.K.; Woodhams, D.C.; White, C.L. 2016.** Salamander chytrid fungus (*Batrachochytrium salamandrivorans*) in the United States—developing research, monitoring, and management strategies. Open-File Report 2331-1258. Reston, VA: U.S. Department of the Interior, Geological Survey. doi:10.3133/ofr20151233.
- Gray, M.J.; Lewis, J.P.; Nanjappa, P.; Klocke, B.; Pasmans, F.; Martel, A.; Stephen, C.; Parra Olea, G.; Smith, S.A.; Sacerdote-Velat, A.; Christman, M.R.; Williams, J.M.; Olson, D.H. 2015.** *Batrachochytrium salamandrivorans*: the North American response and a call for action. *PLoS Pathogens*. 11(12): e1005251. doi:10.1371/journal.ppat.1005251.
- Grinnell, J.; Dixon, J.S. 1926.** Two new races of the pine marten from the Pacific coast of North America. *Zoology*. 21: 411–417.
- Gundale, M.J. 2002.** Influence of exotic earthworms on the soil organic horizon and the rare fern *Botrychium mormo*. *Conservation Biology*. 16(6): 1555–1561.
- Gundale, M.J.; Jolly, W.M.; Deluca, T.H. 2005.** Susceptibility of a northern hardwood forest to exotic earthworm invasion. *Conservation Biology*. 19(4): 1075–1083.
- Hagar, J.C. 2007.** Wildlife species associated with non-coniferous vegetation in Pacific Northwest conifer forests: a review. *Forest Ecology and Management*. 246(1): 108–122. doi:10.1016/j.foreco.2007.03.054.
- Haggard, P.; Haggard, J. 2006.** *Insects of the Pacific Northwest*. Portland, OR: Timber Press. 295 p.
- Halofsky, J.S.; Halofsky, J.E.; Burcsu, T.; Hemstrom, M.A. 2014.** Dry forest resilience varies under simulated climate-management scenarios in a central Oregon, USA landscape. *Ecological Applications*. 24(8): 1908–1925. doi:10.1890/13-1653.1.
- Halpern, C.B.; Evan, S.A.; Nelson, C.R.; McKenzie, D.; Liguori, D.A.; Hibbs, D.E.; Halaj, M.G. 1999.** Response of forest vegetation to varying levels and patterns of green-tree retention: an overview of a long-term experiment. *Northwest Science*. 73(Special issue): 27–44.
- Halpern, C.B.; Halaj, J.; Evans, S.A.; Dovciak, M. 2012.** Level and pattern of overstory retention interact to shape long-term responses of understories to timber harvest. *Ecological Applications*. 22: 2049–2064. doi:10.1890/12-0299.1.
- Halpern, C.B.; McKenzie, D.; Evans, S.A.; Maguire, D.A. 2005.** Initial responses of forest understories to varying levels and patterns of green-tree retention. *Ecological Applications*. 15(1): 175–195. doi:10.1890/03-6000.
- Halpern, C.B.; Spies, T.A. 1995.** Plant species diversity in natural and managed forests of the Pacific Northwest. *Ecological Applications*. 5(4): 913. doi:10.2307/2269343.
- Halsey, S.M.; Zielinski, W.J.; Scheller, R.M. 2015.** Modeling predator habitat to enhance reintroduction planning. *Landscape Ecology*. 30: 1257–1271.

- Hamm, K.A.; Diller, L.V. 2009.** Forest management effects on abundance of woodrats in northern California. *Northwestern Naturalist*. 90(2): 97–106.
- Hanson, J.J.; Lorimer, C.G.; Halpin, C.R.; Palik, B.J. 2012.** Ecological forestry in an uneven-aged, late-successional forest: simulated effects of contrasting treatments on structure and yield. *Forest Ecology and Management*. 270: 94–107.
- Harrison, S.; Noss, R. 2017.** Endemism hotspots are linked to stable climatic refugia. *Annals of Botany*. doi:10.1093/aob/mcw248.
- Hartwig, C.L.; Eastman, D.S.; Harestad, A.S. 2003.** Characteristics of pileated woodpecker (*Dryocopus pileatus*) cavity trees and their patches on southeastern Vancouver Island, British Columbia, Canada. *Forest Ecology and Management*. 187(2–3): 225–234.
- Hayes, J.P.; Cross, S.P. 1987.** Characteristics of logs used by western red-backed voles, *Clethrionomys californicus*, and deer mice, *Peromyscus maniculatus*. *Canadian Field-Naturalist*. 101: 543–546.
- Hayes, J.P.; Horvath, E.G.; Hounihan, P. 1995.** Townsend's chipmunk populations in Douglas-fir plantations and mature forests in the Oregon Coast Range. *Canadian Journal of Zoology*. 73: 67–73.
- Haynes, R.W.; Bormann, B.T.; Lee, D.C.; Martin, J.R. 2006.** Northwest Forest Plan—the first 10 years (1994–2003): synthesis of monitoring and research results. Gen. Tech. Rep. PNW-GTR-651. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 292 p. doi:10.2737/pnw-gtr-651.
- Haynes, R.W.; Perez, G.E. 2001.** Northwest Forest Plan research synthesis. Gen. Tech. Rep. PNW-GTR-498. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 130 p.
- Healey, S.P.; Cohen, W.B.; Spies, T.A.; Moeur, M.; Pflugmacher, D.; Whitley, M.G.; Lefsky, M. 2008.** The relative impact of harvest and fire upon landscape-level dynamics of older forests: lessons from the Northwest Forest Plan. *Ecosystems*. 11(7): 1106–1119. doi:10.1007/s10021-008-9182-8.
- Hebel, C.L.; Smith, J.E.; Cromack, K. 2009.** Invasive plant species and soil microbial response to wildfire burn severity in the Cascade Range of Oregon. *Applied Soil Ecology*. 42(2): 150–159. doi:10.1016/j.apsoil.2009.03.004.
- Heithecker, T.D.; Halpern, C.B. 2006.** Variation in microclimate associated with dispersed-retention harvests in coniferous forests of western Washington. *Forest Ecology and Management*. 226(1–3): 60–71. doi:10.1016/j.foreco.2006.01.024.
- Hendrix, P.F.; Bohlen, P.J. 2002.** Exotic earthworm invasions in North America: ecological and policy implications. *BioScience*. 52(9): 801–812.
- Herrera, J.M.; De Sá Teixeira, I.; Rodríguez-Pérez, J.; Mira, A. 2016.** Landscape structure shapes carnivore-mediated seed dispersal kernels. *Landscape Ecology*. 31(4): 731–743.
- Hershler, R. 1999.** A systematic review of the Hydrobiid snails (Gastropoda: Rissoidae) of the Great Basin, western United States. Part II: general *Colligyrus*, *Eremopyrgus*, *Fluminicola*, *Pristinicola*, and *Tryonia*. *Veliger*. 42(4): 306–337.
- Hershler, R.; Liu, H.-P.; Frest, T.J.; Johannes, E.J. 2007.** Extensive diversification of pebblesnails (Lithoglyphidae: *Fluminicola*) in the upper Sacramento River basin, northwestern USA. *Zoological Journal of the Linnean Society*. 149(3): 371–422. doi:10.1111/j.1096-3642.2007.00243.x.
- Hickerson, C.A.; Anthony, C.D.; Walton, B.M. 2005.** Edge effects and intraguild predation in native and introduced centipedes: evidence from the field and from laboratory microcosms. *Oecologia*. 146(1): 110–119.

- Higgins, J.V.; Bryer, M.T.; Khoury, M.L.; Fitzhugh, T.W. 2005.** A freshwater classification approach for biodiversity conservation planning. *Conservation Biology*. 19(2): 432–445. doi:10.1111/j.1523-1739.2005.00504.x.
- Hofmeister, J.; Hošek, J.; Brabec, M.; Dvůrák, D.; Beran, M.; Deckerová, H.; Burel, J.; Kríž, M.; Borovicka, J.; Běťák, J.; Vašutová, M.; Malíček, J.; Palice, Z.; Syrovátková, L.; Steinová, J.; Cernajová, I.; Holá, E.; Novozámská, E.; Cížek, L.; Iarema, V.; Baltaziuk, K.; Svoboda, T. 2015.** Value of old forest attributes related to cryptogam species richness in temperate forests: a quantitative assessment. *Ecological Indicators*. 57: 497–504
- Hollenbeck, J.P.; Saab, V.A.; Frenzel, R.W. 2011.** Habitat suitability and nest survival of white-headed woodpeckers in unburned forests of Oregon. *Journal of Wildlife Management*. 75(5): 1061–1071.
- Homburg, K.; Brandt, P.; Drees, C.; Assmann, T. 2014.** Evolutionarily significant units in a flightless ground beetle show different climate niches and high extinction risk due to climate change. *Journal of Insect Conservation*. 18(5): 781–790.
- Hooper, D.U.; Chapin, F.S.; Ewel, J.J.; Hector, A.; Inchausti, P.; Lavorel, S.; Lawton, J.H.; Lodge, D.M.; Loreau, M.; Naeem, S.; Schmid, B.; Setälä, H.; Symstad, A.J.; Vandermeer, J.; Wardle, D.A. 2005.** Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecological Monographs*. 75(1): 3–35. doi:10.1890/04-0922.
- Horsák, M.; Hájek, M.; Spitale, D.; Hájková, P.; Díte, D.; Nekola, J.C. 2012.** The age of island-like habitats impacts habitat specialist species richness. *Ecology*. 93(5): 1106–1114.
- Hunter, M., Jr.; Westgate, M.; Barton, P.; Calhoun, A.; Pierson, J.; Tulloch, A.; Beger, M.; Branquinho, C.; Caro, T.; Gross, J.; Heino, J.; Lane, P.; Longo, C.; Martin, K.; McDowell, W.H.; Mellin, C.; Salo, H.; Lindenmayer, D. 2016.** Two roles for ecological surrogacy: indicator surrogates and management surrogates. *Ecological Indicators*. 63: 121–125.
- Hunter, M.L. 1991.** Coping with ignorance: the coarse-filter strategy for maintaining biodiversity. In: Kohm, K.A., ed. *Balancing on the brink of extinction*. Covelo, CA: Island Press: 266–281.
- Hunter, M.L. 2005.** A mesofilter conservation strategy to complement fine and coarse filters. *Conservation Biology*. 19(4): 1025–1029.
- Hunter, M.L.; Redford, K.H.; Lindenmayer, D.B. 2014.** The complementary niches of anthropocentric and biocentric conservationists. *Conservation Biology*. 28(3): 641–645.
- Huwer, J.R.; Koprivnikar, J.; Johnson, P.T.J.; Whyard, S. 2015.** Development and application of an edna method to detect and quantify a pathogenic parasite in aquatic ecosystems. *Ecological Applications*. 25(4): 991–1002.
- Isaak, D.J.; Young, M.K.; Luce, C.H.; Hostetler, S.W.; Wenger, S.J.; Peterson, E.E.; Ver Hoef, J.M.; Groce, M.C.; Horan, D.L.; Nagel, D.E. 2016.** Slow climate velocities of mountain streams portend their role as refugia for cold-water biodiversity. *Proceedings of the National Academy of Sciences*. 113(16): 4374–4379. doi:10.1073/pnas.1522429113.
- Jactel, H.; Branco, M.; Duncker, P.; Gardiner, B.; Grodzki, W.; Langstrom, B.; Moreira, F.; Netherer, S.; Nicoll, B.; Orazio, C.; Piou, D.; Schelhaas, M.J.; Tojic, K. 2012.** A multicriteria risk analysis to evaluate impacts of forest management alternatives on forest health in Europe. *Ecology and Society*. 17(4): Article 52. <http://dx.doi.org/10.5751/ES-04897-170452>.

- Johnstone, J.F.; Allen, C.D.; Franklin, J.F.; Frelich, L.E.; Harvey, B.J.; Higuera, P.E.; Mack, M.C.; Meentemeyer, R.K.; Metz, M.R.; Perry, G.L.W.; Schoennagel, T.; Turner, M.G. 2016.** Changing disturbance regimes, ecological memory, and forest resilience. *Frontiers in Ecology and Evolution*. 14(7): 369–378.
- Jones, C.C.; Acker, S.A.; Halpern, C.B. 2010.** Combining local- and large-scale models to predict the distributions of invasive plant species. *Ecological Applications*. 20(2): 311–326.
- Jones, G.M.; Gutierrez, R.J.; Tempel, D.J.; Whitmore, S.A.; Berigan, W.J.; Peery, M.Z. 2016.** Megafires: an emerging threat to old-forest species. *Frontiers in Ecology and Evolution*. 14(6): 300–306.
- Jordan, S.F.; Black, S.H. 2015.** Conservation assessment for *Cryptomastix devia*, *Puget Oregonian* (originally issued 1999 as management recommendations by T. E. Burke, N. Duncan, and P. Jeske; revised 2005 by T. Kogut and N. Duncan). Portland, OR: Interagency Sensitive and Special Status Species Program, U.S. Department of Agriculture, Forest Service, Pacific Northwest Region and U.S. Department of the Interior, Bureau of Land Management, Oregon and Washington Office. 32 p.
- Jørgensen, P.M. 2000.** Survey of the lichen family Pannariaceae on the American continent, north of Mexico. *The Bryologist*. 103(4): 670–704. doi:10.1639/0007-2745(2000)103[0670:sotlfp]2.0.co;2.
- Jules, E.S. 1998.** Habitat fragmentation and demographic change for a common plant: *Trillium* in an old-growth forest. *Ecology*. 79: 1645–1656.
- Jules, E.S.; Rathcke, B.J. 1999.** Mechanisms of reduced *Trillium* recruitment along edges of old-growth forest fragments. *Conservation Biology*. 13(4): 784–793.
- Jules, M.J.; Sawyer, J.O.; Jules, E.S. 2008.** Assessing the relationships between stand development and understory vegetation using a 420-year chronosequence. *Forest Ecology and Management*. 255(7): 2384–2393. doi:10.1016/j.foreco.2007.12.042.
- Jusino, M.A.; Lindner, D.L.; Banik, M.T.; Rose, K.R.; Walters, J.R. 2016.** Experimental evidence of a symbiosis between red-cockaded woodpeckers and fungi. *Proceedings of the Royal Society B: Biological Sciences*. 283(1827): 20160106. doi:10.1098/rspb.2016.0106.
- Kahmen, A.; Jules, E.S. 2005.** Assessing the recovery of a long-lived herb following logging: *Trillium ovatum* across a 424-year chronosequence. *Forest Ecology and Management*. 210(1-3): 107–116.
- Keeton, W.S.; Franklin, J.F. 2005.** Do remnant old-growth trees accelerate rates of succession in mature Douglas-fir forests? *Ecological Monographs*. 75(1): 103–118.
- Kelley, R.; Dowlan, S.; Duncan, N.; Burke, T. 1999.** Field guide to survey and manage terrestrial mollusk species from the Northwest Forest Plan. Portland, OR: U.S. Department of the Interior, Bureau of Land Management Oregon State Office. 114 p.
- Kendall, K.C.; McKelvey, K.S. 2008.** Hair collection. In: Long, R.A.; Mackay, P.; Zielinski, W.J.; Ray, J.C., eds. *Noninvasive survey methods for carnivores*. Washington, DC: Island Press: 142–182.
- Kerns, B.K.; Ager, A. 2007.** Risk assessment for biodiversity conservation planning in Pacific Northwest forests. *Forest Ecology and Management*. 246(1): 38–44. doi:10.1016/j.foreco.2007.03.049.
- Keirst, C.; Gordon, S. 2011.** Dragonflies and damselflies of Oregon. Corvallis, OR: Oregon State University Press. 304 p.
- Kirk, T.A.; Zielinski, W.J. 2009.** Developing and testing a landscape habitat suitability model for the American marten (*Martes americana*) in the Cascades mountains of California. *Landscape Ecology*. 24(6): 759–773.

- Kline, J.D.; Harmon, M.E.; Spies, T.A.; Morzillo, A.T.; Pabst, R.J.; McComb, B.C.; Schnekenburger, F.; Olsen, K.A.; Csuti, B.; Vogeler, J.C. 2016.** Evaluating carbon storage, timber harvest, and habitat possibilities for a western Cascades (USA) forest landscape. *Ecological Applications*. 26(7): 2044–2059. doi:10.1002/eap.1358.
- Kluber, M.R.; Olson, D.H.; Puettmann, K.J. 2008.** Amphibian distributions in riparian and upslope areas and their habitat associations on managed forest landscapes in the Oregon Coast Range. *Forest Ecology and Management*. 256(4): 529–535. doi:10.1016/j.foreco.2008.04.043.
- Kluber, M.R.; Olson, D.H.; Puettmann, K.J. 2009.** Downed wood microclimates and their potential impact on Plethodontid salamander habitat in the Oregon Coast Range. *Northwest Science*. 83(1): 25–34. doi:10.3955/046.083.0103.
- Knick, S.T.; Sweeney, S.J.; Alldredge, J.R.; Brittell, J.D. 1984.** Autumn and winter food habits of bobcats in Washington state. *Great Basin Naturalist*. 44(1): 70–74.
- Koch, J.; Strange, J.; Williams, P. 2012.** Bumble bees of the western United States. Washington, DC: U.S. Department of Agriculture, Forest Service. 143 p.
- Koehler, G.M.; Maletzke, B.T.; Von Kienast, J.A.; Aubry, K.B.; Wielgus, R.B.; Naney, R.H. 2008.** Habitat fragmentation and the persistence of lynx populations in Washington state. *Journal of Wildlife Management*. 72(7): 1518–1524.
- Koepfli, K.P.; Deere, K.A.; Slater, G.J.; Begg, C.; Begg, K.; Grassman, L.; Lucherini, M.; Veron, G.; Wayne, R.K. 2008.** Multigene phylogeny of the Mustelidae: resolving relationships, tempo and biogeographic history of a mammalian adaptive radiation. *BMC Biology*. 6(10). doi:10.1186/1741-7007-6-10.
- Konstantinova, N.A.; Vilnet, A.A.; Söderström, L.; Hagborg, A.; Von Konrat, M. 2013.** Notes on early land plants today. 14. Transfer of two *Macrodiplophyllum* species to *Douinia* (Scapaniaceae, Marchantiophyta). *Phytotaxa*. 76(3): 31–32. doi:10.11646/phytotaxa.76.3.2.
- Krebs, J.; Lofroth, E.C.; Parfitt, I. 2007.** Multiscale habitat use by wolverines in British Columbia, Canada. *Journal of Wildlife Management*. 71(7): 2180. doi:10.2193/2007-099.
- Kroll, A.J. 2009.** Sources of uncertainty in stream-associated amphibian ecology and responses to forest management in the Pacific Northwest, USA: a review. *Forest Ecology and Management*. 257(4): 1188–1199. doi:10.1016/j.foreco.2008.12.008.
- Kroll, A.J.; Lacki, M.J.; Arnett, E.B. 2012.** Research needs to support management and conservation of cavity-dependent birds and bats on forested landscapes in the Pacific Northwest. *Western Journal of Applied Forestry*. 27(3): 128–136.
- Kurek, P.; Kapusta, P.; Holeksa, J. 2014.** Burrowing by badgers (*Meles meles*) and foxes (*Vulpes vulpes*) changes soil conditions and vegetation in a European temperate forest. *Ecological Research*. 29(1): 1–11.
- Lacki, M.J.; Baker, M.D.; Johnson, J.S. 2012.** Temporal dynamics of roost snags of long-legged myotis in the Pacific Northwest, USA. *Journal of Wildlife Management*. 76(6): 1310–1316.
- Lapoint, S.; Gallery, P.; Wikelski, M.; Kays, R. 2013.** Animal behavior, cost-based corridor models, and real corridors. *Landscape Ecology*. 28(8): 1615–1630.
- Latif, Q.; Saab, V.; Mellen-Mclean, K.; Dudley, J. 2015.** Evaluating habitat suitability models for nesting white-headed woodpeckers in unburned forest. *Journal of Wildlife Management*. 79: 263–273.

- Lawler, J.J.; Safford, H.D.; Girvetz, E.H. 2012.** Martens and fishers in a changing climate. In: Aubry, K.B.; Zielinski, W.J.; Raphael, M.G.; Proulx, G.; Buskirk, S.W., eds. Biology and conservation of martens, sables, and fishers: a new synthesis. Ithaca, NY: Cornell University Press: 371–397.
- Lawler, J.J.; White, D.; Sifneos, J.C.; Master, L.L. 2003.** Rare species and the use of indicator groups for conservation planning. *Conservation Biology*. 17(3): 875–882. doi:10.1046/j.1523-1739.2003.01638.x.
- Lehmkuhl, J.F.; Gould, L.E.; Cazares, E.; Hosford, D.R. 2004.** Truffle abundance and mycophagy by northern flying squirrels in eastern Washington forests. *Forest Ecology and Management*. 200(1–3): 49–65.
- Lehmkuhl, J.F.; Kennedy, M.; Ford, E.D.; Singleton, P.H.; Gaines, W.L.; Lind, R.L. 2007.** Seeing the forest for the fuel: integrating ecological values and fuels management. *Forest Ecology and Management*. 246(1): 73–80. doi:10.1016/j.foreco.2007.03.071.
- Lehmkuhl, J.F.; Marcot, B.G.; Quinn, T. 2001.** Characterizing species at risk. In: Johnson, D.H.; O'Neill, T.A., eds. Wildlife-habitat relationships in Oregon and Washington. Corvallis, OR: Oregon State University Press: 474–500.
- Leshner, R.D.; Derr, C.C.; Geiser, L.H. 2003.** Natural history and management considerations for Northwest Forest Plan Survey and Manage lichens based on information as of the year 2000. Natural Resources Tech. Paper R6-NR-S&M-TP-03-03. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region. 211 p.
- Leuthold, N.; Adams, M.J.; Hayes, J.P. 2012.** Short-term response of *Dicamptodon tenebrosus* larvae to timber management in southwestern Oregon. *Journal of Wildlife Management*. 76(1): 28–37. doi:10.1002/jwmg.269.
- Lewis, J.C. 2013.** Implementation plan for reintroducing fishers to the Cascade Mountain range in Washington. Olympia, WA: Washington Department of Fish and Wildlife. 29 p.
- Lewis, J.C. 2014.** Post-release movements, survival, and resource selection of fishers (*Pekania pennanti*) translocated to the Olympic Peninsula of Washington. Seattle, WA: University of Washington. 133 p Ph.D. dissertation.
- Lewis, J.C.; Hayes, G.E. 2004.** Feasibility assessment for reintroducing fishers to Washington. Olympia, WA: Washington Department of Fish and Wildlife, Wildlife Program. 70 p.
- Lindenmayer, D.B.; Fischer, J.; Felton, A.; Montague-Drake, R.; Manning, A.D.; Simberloff, D.; Youngentob, K.; Saunders, D.; Wilson, D.; Felton, A.M.; Blackmore, C.; Lowe, A.; Bond, S.; Munro, N.; Elliott, C.P. 2007.** The complementarity of single-species and ecosystem-oriented research in conservation research. *Oikos*. 116(7): 1220–1226.
- Lindh, B.C.; Muir, P.S. 2004.** Understory vegetation in young Douglas-fir forests: Does thinning help restore old-growth composition? *Forest Ecology and Management*. 192(2-3): 285–296. doi:10.1016/j.foreco.2004.01.018.
- Lindstrand, L.I.; Bainbridge, K.; Youngblood, B. 2012.** Habitat characteristics, a range extension, and an elevational record for Shasta salamanders. *California Fish and Game*. 98(4): 236–241.
- Liu, H.-P.; Hershler, R.; Rossel, C. 2015.** Taxonomic status of the Columbia dusksnail (*Truncatelloidea*, *Amnicolidae*, *Colligyrus*). *ZooKeys*. 514: 1–13. doi:10.3897/zookeys.514.9919.
- Loehle, C.; Irwin, L.; Manly, B.F.J.; Merrill, A. 2015.** Range-wide analysis of northern spotted owl nesting habitat relations. *Forest Ecology and Management*. 342: 8–20. doi:10.1016/j.foreco.2015.01.010.

- Lofroth, E.C.; Krebs, J.A.; Harrower, W.L.; Lewis, D. 2007.** Food habits of wolverine *Gulo gulo* in montane ecosystems of British Columbia, Canada. *Wildlife Biology*. 13(supp2): 31–37. doi:10.2981/0909-6396(2007)13[31:fhowgg]2.0.co;2.
- Lofroth, E.C.; Raley, C.M.; Higley, J.M.; Truex, R.L.; Yaeger, J.S.; Lewis, J.C.; Happe, P.J.; Finley, L.L.; Naney, R.H.; Hale, L.J.; Krause, A.L.; Livingston, S.A.; Myers, A.M.; Brown, R.N. 2010.** Conservation of fishers (*Martes pennanti*) in south-central British Columbia, western Washington, western Oregon, and California—volume I: Conservation assessment. Denver, CO: U.S. Department of the Interior, Bureau of Land Management. 163 p.
- Long, R.A.; Zielinski, W.J. 2008.** Designing effective noninvasive carnivore surveys. In: Long, R.A.; Mackay, P.; Zielinski, W.J.; Ray, J.C., eds. *Noninvasive survey methods for carnivores*. Island Press: 8–44.
- Lorch, J.M.; Palmer, J.M.; Lindner, D.L.; Ballmann, A.E.; George, K.G.; Griffin, K.; Knowles, S.; Huckabee, J.R.; Haman, K.H.; Anderson, C.D.; Becker, P.A.; Buchanan, J.B.; Foster, J.T.; Blehert, D.S. 2016.** First detection of bat white-nose syndrome in western North America. *mSphere*. 1(4):e00148. doi:10.1128/mSphere.00148-16.
- Lorenz, T.J.; Vierling, K.T.; Kozma, J.M.; Millard, J.E.; Raphael, M.G. 2015.** Space use by white-headed woodpeckers and selection for recent forest disturbances. *Journal of Wildlife Management*. 79(8): 1286–1297.
- Losey, J.E.; Vaughan, M. 2006.** The economic value of ecological services provided by insects. *BioScience*. 56(4): 311. doi:10.1641/0006-3568(2006)56[311:tevoes]2.0.co;2.
- Lowe, W.H.; Likens, G.E.; McPeck, M.A.; Buso, D.C. 2006.** Linking direct and indirect data on dispersal: isolation by slope in a headwater stream salamander. *Ecology*. 87(2): 334–339.
- Loya, D.T.; Jules, E.S. 2008.** Use of species richness estimators improves evaluation of understory plant response to logging: a study of redwood forests. *Plant Ecology*. 194(2): 179–194.
- Luo, W.; Callaway, R.M.; Atwater, D.Z. 2016.** Intraspecific diversity buffers the inhibitory effects of soil biota. *Ecology*. 97(8): 1913–1918.
- Luoma, D.L.; Eberhart, J.L. 2005.** Results from green-tree retention experiments: ectomycorrhizal fungi. In: Peterson, C.E.; Maguire, D., eds. *Balancing ecosystem values: innovative experiments for sustainable forestry—proceedings of a conference*. Gen. Tech. Rep. PNW-GTR-635. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 257–264. <http://dx.doi.org/10.2737/pnw-gtr-635>.
- Luoma, D.L.; Trappe, J.M.; Claridge, A.W.; Jacobs, K.M.; Cazares, E. 2003.** Relationships among fungi and small mammals in forested ecosystems. In: Zabel, C.J.; Anthony, R.G., eds. *Mammal community dynamics: management and conservation in the coniferous forests of western North America*. Cambridge, United Kingdom: Cambridge University Press: 343–373.
- Mackenzie, D.I.; Nichols, J.D.; Royle, J.A.; Pollock, K.H.; Bailey, L.L.; Hines, J.E. 2006.** Occupancy estimation and modeling: inferring patterns and dynamics of species occurrence. Burlington, MA: Academic Press. 324 p.
- Magoun, A.J.; Copeland, J.P. 1998.** Characteristics of wolverine reproductive den sites. *The Journal of Wildlife Management*. 62(4): 1313. doi:10.2307/3801996.
- Maletzke, B.T.; Koehler, G.M.; Wielgus, R.B.; Aubry, K.B.; Evans, M.A. 2008.** Habitat conditions associated with lynx hunting behavior during winter in northern Washington state. *Journal of Wildlife Management*. 72(7): 1473–1478.

- Marcot, B.G. 1983.** Snag use by birds in Douglas-fir clearcuts. In: Davis, J.W.; Goodwin, G.A.; Ockenfels, R.A., eds. Snag habitat management: proceedings of the symposium. Gen. Tech. Rep. RM-GTR-99. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 134–139.
- Marcot, B.G. 1985.** Habitat relationships of birds and young-growth Douglas-fir in northwestern California. Corvallis, OR: Oregon State University. 282 p. Ph.D. dissertation.
- Marcot, B.G. 2002.** An ecological functional basis for managing decaying wood for wildlife. In: Laudenslayer, W.F., Jr.; Shea, P.J.; Valentine, B.E.; Weatherspoon, C.P.; Lisle, T.E., eds. Proceedings of the symposium on the ecology and management of dead wood in western forests. Gen. Tech. Rep. PSW-GTR-181. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 895–910.
- Marcot, B.G. 2007.** Biodiversity and the lexicon zoo. *Forest Ecology and Management*. 246(1): 4–13.
- Marcot, B.G. 2017.** A review of the role of fungi in wood decay of forest ecosystems. Res. Note PNW-RN-575. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station and Pacific Northwest Region. 31 p.
- Marcot, B.G.; Flather, C.H. 2007.** Species-level strategies for conserving rare or little-known species. In: Raphael, M.G.; Molina, R., eds. Washington, DC: Island Press: 125–164.
- Marcot, B.G.; Hohenlohe, P.A.; Morey, S.; Holmes, R.; Molina, R.; Turley, M.; Huff, M.; Laurence, J. 2006.** Characterizing species at risk II: using Bayesian belief networks as decision support tools to determine species conservation categories under the Northwest Forest Plan. *Ecology and Society*. 11(2): 12. <http://www.ecologyandsociety.org/vol11/iss2/art12/>. (13 November 2017).
- Marcot, B.G.; Livingston, S. 2009.** Report on the red tree vole listing advisory panel, held August 25–26, 2009. Portland, OR: U.S. Department of the Interior, Fish and Wildlife Service, Oregon Office. 50 p.
- Marcot, B.G.; Molina, R. 2006.** Conservation of other species associated with older forest conditions. In: Haynes, R.; Bormann, B.T.; Lee, D.C.; Martin, J.R., eds. Northwest Forest Plan—the first 10 years (1994–2003): synthesis of monitoring and research results. Gen. Tech. Rep. PNW-GTR-651. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 145–179.
- Marcot, B.G.; Morey, S.; Naney, B.; Finley, L. 2012a.** Fisher threats workshop report. Appendix 1. In: Naney, R.H.; Finley, L.L.; Lofroth, E.C.; Happe, P.J.; Krause, A.L.; Raley, C.M.; Truex, R.L.; Hale, L.J.; Higley, J.M.; Kosc, A.D.; Lewis, J.C.; Livingston, S.A.; Macfarlane, D.C.; Myers, A.M.; Yaeger, J.S., eds. Conservation of fishers (*Martes pennanti*) in south-central British Columbia, western Washington, western Oregon, and California. Volume III: Threat assessment. Denver, CO: U.S. Department of the Interior, Bureau of Land Management: 47–55.
- Marcot, B.G.; Raphael, M.G.; Schumaker, N.H.; Galleher, B. 2013.** How big and how close? Habitat patch size and spacing to conserve a threatened species. *Natural Resource Modeling*. 26(2): 194–214. doi:10.1111/j.1939-7445.2012.00134.x.
- Marcot, B.G.; Sieg, C.H. 2007.** System-level strategies for conserving rare or little-known species. In: Raphael, M.G.; Molina, R., eds. Conservation of rare or little-known species: biological, social, and economic considerations. Washington, DC: Island Press: 165–186.
- Marcot, B.G.; Thompson, M.P.; Runge, M.C.; Thompson, F.R.; McNulty, S.; Cleaves, D.; Tomosy, M.; Fisher, L.; Bliss, A. 2012b.** Recent advances in applying decision science to managing national forests. *Forest Ecology and Management*. 285: 123–132.

- Marcot, B.G.; Vander Heyden, M. 2001.** Key ecological functions of wildlife species. In: Johnson, D.H.; O'Neill, T.A., eds. *Wildlife-habitat relationships in Oregon and Washington*. Corvallis, OR: Oregon State University Press: 168–186.
- Marczak, L.B.; Sakamaki, T.; Turvey, S.L.; Deguise, I.; Wood, S.L.R.; Richardson, J.S. 2010.** Are forested buffers an effective conservation strategy for riparian fauna? An assessment using meta-analysis. *Ecological Applications*. 20(1): 126–134.
- Martel, A.; Blooi, M.; Adriaensen, C.; Van Rooij, P.; Beukema, W.; Fisher, M.C.; Farrer, R.A.; Schmidt, B.R.; Tobler, U.; Goka, K.; Lips, K.R.; Muletz, C.; Zamudio, K.R.; Bosch, J.; Lotters, S.; Wombwell, E.; Garner, T.W.J.; Cunningham, A.A.; Spitzen-Van Der Sluijs, A.; Salvidio, S.; Ducatelle, R.; Nishikawa, K.; Nguyen, T.T.; Kolby, J.E.; Van Bocxlaer, I.; Bossuyt, F.; Pasmans, F. 2014.** Recent introduction of a chytrid fungus endangers western palearctic salamanders. *Science*. 346(6209): 630–631. doi:10.1126/science.1258268.
- Matthews, S.M.; Higley, J.M.; Rennie, K.M.; Green, R.E.; Goddard, C.A.; Wengert, G.M.; Gabriel, M.W.; Fuller, T.K. 2013.** Reproduction, recruitment, and dispersal of fishers (*Martes pennanti*) in a managed Douglas-fir forest in California. *Journal of Mammalogy*. 94(1): 100–108.
- McAlpine, C.; Spies, T.; Norman, P.; Peterson, A. 2007.** Conserving forest biodiversity across multiple land ownerships: lessons from the Northwest Forest Plan and the southeast Queensland regional forests agreement (Australia). *Biological Conservation*. 134(4): 580–592. doi:10.1016/j.biocon.2006.09.009.
- McComb, B.C.; Spies, T.A.; Olsen, K.A. 2007.** Sustaining biodiversity in the Oregon Coast Range: potential effects of forest policies in a multi-ownership province. *Ecology and Society*. 12(2): 29. <http://www.ecologyandsociety.org/vol12/iss2/art29/>. (15 January 2018).
- McCune, B.; Hutchinson, J.; Berryman, S. 2002.** Concentration of rare epiphytic lichens along large streams in a mountainous watershed in Oregon, U.S.A. *The Bryologist*. 105(3): 439–450. doi:10.1639/0007-2745(2002)105[0439:corela]2.0.co;2.
- McFadden-Hiller, J.E.; Hiller, T.L. 2015.** Non-invasive survey of forest carnivores in the northern Cascades of Oregon, USA. *Northwestern Naturalist*. 96(2): 107–117.
- McIntyre, A.P.; Schmitz, R.A.; Crisafulli, C.M. 2006.** Associations of the Van Dyke's salamander (*Plethodon vandykei*) with geomorphic conditions in headwall seeps of the Cascade range, Washington state. *Journal of Herpetology*. 40(3): 309–322. doi:10.1670/0022-1511(2006)40[309:aotvds]2.0.co;2.
- McKee, A.M.; Calhoun, D.L.; Barichivich, W.J.; Spear, S.F.; Goldberg, C.S.; Glenn, T.C. 2015.** Assessment of environmental DNA for detecting presence of imperiled aquatic amphibian species in isolated wetlands. *Journal of Fish and Wildlife Management*. 6(2): 498–510.
- McKelvey, K.S.; Aubry, K.B.; Ortega, Y.K. 2000.** History and distribution of lynx in the contiguous United States. In: Ruggiero, L.F.; Aubry, K.B.; Buskirk, S.W.; Koehler, G.M.; Krebs, C.J.; McKelvey, K.S.; Squires, J.R., tech. eds. *Ecology and conservation of lynx in the United States*. Boulder, CO: University Press of Colorado: 207–264. Chapter 8.
- McKelvey, K.S.; Copeland, J.P.; Schwartz, M.K.; Littell, J.S.; Aubry, K.B.; Squires, J.R.; Parks, S.A.; Elsner, M.M.; Mauger, G.S. 2011.** Climate change predicted to shift wolverine distributions, connectivity, and dispersal corridors. *Ecological Applications*. 21(8): 2882–2897. doi:10.1890/10-2206.1.
- McRae, B.H.; Popper, K.; Jones, A.; Schindel, M.; Buttrick, S.; Hall, K.; Unnasch, R.S.; Platt, J. 2016.** Conserving nature's stage: mapping omnidirectional connectivity for resilient terrestrial landscapes in the Pacific Northwest. Portland, OR: The Nature Conservancy. 47 p.

- McRae, B.H.; Shah, V.B. 2009.** Circuitscape user guide. Santa Barbara, CA: University of California, Santa Barbara. 12 p. <http://www.circuitscape.org>. (16 January 2018).
- Mead, L.S.; Clayton, D.R.; Nauman, R.S.; Olson, D.H.; Pfrender, M.E. 2005.** Newly discovered populations of salamanders from Siskiyou county California represent a species distinct from *Plethodon stormi*. *Herpetologica*. 61(2): 158–177.
- Mellen, K.; Ager, A. 2002.** A coarse wood dynamics model for the western Cascades. In: Laudenslayer, W.F., Jr.; Shea, P.J.; Valentine, B.E.; Weatherspoon, C.P.; Lisle, T.E., eds. Proceedings of the symposium on the ecology and management of dead wood in western forests. Gen. Tech. Rep. PSW-GTR-181. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 503–516.
- Milcu, A.; Partsch, S.; Langel, R.; Scheu, S. 2006.** The response of decomposers (earthworms, springtails and microorganisms) to variations in species and functional group diversity of plants. *Oikos*. 112(3): 513–524.
- Miller, J.C.; Hammond, P.C. 2000.** Macromoths of Northwest forests and woodland. FHTET-98-18. Morgantown, WV: U.S. Department of Agriculture, Forest Service, Forest Health Technology Enterprise Team. 133 p.
- Miller, J.C.; Hammond, P.C. 2003.** Lepidoptera of the Pacific Northwest: caterpillars and adults. FHTET-2003-03. Morgantown, WV: U.S. Department of Agriculture, Forest Service, Forest Health Technology Enterprise Team. 324 p.
- Miller, J.C.; Hammond, P.C.; Ross, D.N.R. 2003.** Distribution and functional roles of rare and uncommon moths (Lepidoptera: Noctuidae: Plusiinae) across a coniferous forest landscape. *Annals of the Entomological Society of America*. 96(6): 847–855.
- Miller, J.E.D.; Villella, J.; Carey, G.; Carlberg, T.; Root, H.T. 2017.** Canopy distribution and survey detectability of a rare old-growth forest lichen. *Forest Ecology and Management*. 392: 195–201.
- Miller, M.P.; Bellinger, M.R.; Forsman, E.D.; Haig, S.M. 2006.** Effects of historical climate change, habitat connectivity, and vicariance on genetic structure and diversity across the range of the red tree vole (*Phenacomys longicaudus*) in the Pacific northwestern United States. *Molecular Ecology*. 15(1): 145–159. doi:10.1111/j.1365-294x.2005.02765.x.
- Miller, M.P.; Forsman, E.D.; Swingle, J.K.; Miller, S.A.; Haig, S.M. 2010.** Size-associated morphological variation in the red tree vole (*Arborimus longicaudus*). *Northwestern Naturalist*. 91: 63–73.
- Moen, R.; Burdett, C.L.; Niemi, G.J. 2008.** Movement and habitat use of Canada lynx during denning in Minnesota. *Journal of Wildlife Management*. 72(7): 1507. doi:10.2193/2008-072.
- Moldenke, A.R.; Fichter, B.L. 1988.** Invertebrates of the H.J. Andrews Experimental Forest, western Cascade Mountains, Oregon: IV. The oribatid mites (Acari: Cryptostigmata). Gen. Tech. Rep. PNW-GTR-217. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 112 p.
- Moldenke, A.R.; Fichter, B.L.; Stephen, W.P.; Griswold, C.E. 1987.** A key to arboreal spiders of Douglas-fir and true fir forests of the Pacific Northwest. Gen. Tech. Rep. PNW-GTR-207. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 48 p.
- Mölder, A.; Schmidt, M.; Engel, F.; Schönfelder, E.; Schulz, F. 2015.** Bryophytes as indicators of ancient woodlands in Schleswig-Holstein (northern Germany). *Ecological Indicators*. 54: 12–30.
- Molina, R. 2008.** Protecting rare, little known, old-growth forest-associated fungi in the Pacific Northwest USA: a case study in fungal conservation. *Mycological Research*. 112(6): 613–638. doi:10.1016/j.mycres.2007.12.005.

- Molina, R.; Marcot, B.G.; Leshner, R. 2006.** Protecting rare, old-growth, forest-associated species under the Survey and Manage program guidelines of the Northwest Forest Plan. *Conservation Biology*. 20(2): 306–318. doi:10.1111/j.1523-1739.2006.00386.x.
- Molina, R.; McKenzie, D.; Leshner, R.; Ford, J.; Alegria, J.; Cutler, R. 2003.** Strategic survey framework for the Northwest Forest Plan Survey and Manage program. Gen. Tech. Rep. PNW-GTR-573. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 34 p.
- Moning, C.; Müller, J. 2009.** Critical forest age thresholds for the diversity of lichens, molluscs and birds in beech (*Fagus sylvatica* L.) dominated forests. *Ecological Indicators*. 9(5): 922–932.
- Mooney, T.L. 2010.** Predicting *Hydromantes shastae* occurrences in Shasta County, California. Los Angeles, CA: University of Southern California. 39 p. M.S. thesis.
- Moriarty, K.M.; Bailey, J.D.; Smythe, S.E.; Verschuyt, J. 2016a.** Distribution of Pacific marten in coastal Oregon. *Northwestern Naturalist*. 97(2): 71–81.
- Moriarty, K.M.; Epps, C.W.; Betts, M.G.; Hance, D.J.; Bailey, J.D.; Zielinski, W.J. 2015.** Experimental evidence that simplified forest structure interacts with snow cover to influence functional connectivity for Pacific martens. *Landscape Ecology*. 30(10): 1865–1877.
- Moriarty, K.M.; Epps, C.W.; Zielinski, W.J. 2016b.** Forest thinning changes movement patterns and habitat use by Pacific marten. *Journal of Wildlife Management*. 80(4): 621–633.
- Moriarty, K.M.; Zielinski, W.J.; Gonzales, A.G.; Dawson, T.E.; Boatner, K.M.; Wilson, C.A.; Schlexer, F.V.; Pilgrim, K.L.; Copeland, J.P.; Schwartz, M.K. 2009.** Wolverine confirmation in California after nearly a century: native or long-distance immigrant? *Northwest Science*. 83(2): 154–162. doi:10.3955/046.083.0207.
- Moss, M.; Hermanutz, L. 2010.** Monitoring the small and slimy—protected areas should be monitoring native and non-native slugs (Mollusca: Gastropoda). *Natural Areas Journal*. 30(3): 322–327.
- Mote, P.W.; Salathé, E.P. 2010.** Future climate in the Pacific Northwest. *Climatic Change*. 102(1–2): 29–50. doi:10.1007/s10584-010-9848-z.
- Murray, D.L.; Steury, T.D.; Roth, J.D. 2008.** Assessment of Canada lynx research and conservation needs in the southern range: another kick at the cat. *Journal of Wildlife Management*. 72(7): 1463. doi:10.2193/2007-389.
- Nauman, R.S.; Olson, D.H. 2004.** Surveys for terrestrial amphibians in Shasta County, California, with notes on the distribution of shasta salamanders (*Hydromantes shastae*). *Northwestern Naturalist*. 85(1): 35–38. doi:10.1898/1051-1733(2004)085<0035:sftais>2.0.co;2.
- Nauman, R.S.; Olson, D.H. 2008.** Distribution and conservation of *Plethodon* salamanders on federal lands in Siskiyou County, California. *Northwestern Naturalist*. 89(1): 1. doi:10.1898/1051-1733(2008)89[1:dacops]2.0.co;2.
- Neitlich, P.N.; McCune, B. 1997.** Hotspots of epiphytic lichen diversity in two young managed forests. *Conservation Biology*. 11(1): 172–182. doi:10.1046/j.1523-1739.1997.95492.x.
- Nelson, C.R.; Halpern, C.B. 2005.** Edge-related responses of understory plants to aggregated retention harvest in the Pacific Northwest. *Ecological Applications*. 15(1): 196–209. doi:10.1890/03-6002.
- Neuschulz, E.L.; Brown, M.; Farwig, N. 2013.** Frequent bird movements across a highly fragmented landscape: the role of species traits and forest matrix. *Animal Conservation*. 16(2): 170–179.
- Nimmo, D.G.; Haslem, A.; Radford, J.Q.; Hall, M.; Bennett, A.F. 2016.** Riparian tree cover enhances the resistance and stability of woodland bird communities during an extreme climatic event. *Journal of Applied Ecology*. 53(2): 449–458.

- Niwa, C.G.; Peck, R.W. 2002.** Influence of prescribed fire on carabid beetle (Carabidae) and spider (Araneae) assemblages in forest litter in southwestern Oregon. *Environmental Entomology*. 31(5): 785–796. doi:10.1603/0046-225x-31.5.785.
- North, M.; Chen, J.; Smith, G.; Krakowiak, L.; Franklin, J. 1995.** Initial response of understory plant diversity and overstory tree diameter growth to a green tree retention harvest. *Northwest Science*. 70: 24–35.
- North, M.; Franklin, J. 1990.** Post-disturbance legacies that enhance biological diversity in a Pacific Northwest old-growth forest. *The Northwest Environmental Journal*. 6: 427–429.
- Noss, R.F. 1987.** From plant communities to landscapes in conservation inventories: a look at The Nature Conservancy (USA). *Biological Conservation*. 41: 11–37.
- Noss, R.F. 1990.** Indicators for monitoring biodiversity: a hierarchical approach. *Conservation Biology*. 4(4): 355–364. doi:10.1111/j.1523-1739.1990.tb00309.x.
- O’Connell, A.F.; Nichols, J.D.; Karanth, K.U., eds. 2011.** Camera traps in animal ecology: methods and analyses. Springer Science & Business Media. doi:10.1007/978-4-431-99495-4_1.
- Odion, D.C.; Sarr, D.A. 2007.** Managing disturbance regimes to maintain biological diversity in forested ecosystems of the Pacific Northwest. *Forest Ecology and Management*. 246(1): 57–65. doi:10.1016/j.foreco.2007.03.050.
- Olson, D.; DellaSala, D.A.; Noss, R.F.; Strittholt, J.R.; Kass, J.; Koopman, M.E.; Allnutt, T.F. 2012.** Climate change refugia for biodiversity in the Klamath-Siskiyou ecoregion. *Natural Areas Journal*. 32(1): 65–74. doi:10.3375/043.032.0108.
- Olson, D.H. 2008.** Conservation assessment for the black salamander (*Aneides flavipunctatus*) in Oregon. Portland, OR: U.S. Department of Agriculture, Forest Service, Region 6 and U.S. Department of the Interior, Bureau of Land Management, Interagency Special Status and Sensitive Species Program. 23 p. <http://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-amphibians.shtml>. (5 December 2017).
- Olson, D.H.; Aanensen, D.M.; Ronnenberg, K.L.; Powell, C.I.; Walker, S.F.; Bielby, J.; Garner, T.W.J.; Weaver, G.; Fisher, M.C. 2013.** Mapping the global emergence of *Batrachochytrium dendrobatidis*, the amphibian chytrid fungus. *PLoS ONE*. 8(2): e56802. doi:10.1371/journal.pone.0056802.
- Olson, D.H.; Anderson, P.D.; Frissell, C.A.; Welsh, H.H.; Bradford, D.F. 2007a.** Biodiversity management approaches for stream-riparian areas: perspectives for Pacific Northwest headwater forests, microclimates, and amphibians. *Forest Ecology and Management*. 246(1): 81–107. doi:10.1016/j.foreco.2007.03.053.
- Olson, D.H.; Burnett, K.M. 2009.** Design and management of linkage areas across headwater drainages to conserve biodiversity in forest ecosystems. *Forest Ecology and Management*. 258: S117–S126. doi:10.1016/j.foreco.2009.04.018.
- Olson, D.H.; Burnett, K.M. 2013.** Geometry of forest landscape connectivity: pathways for persistence. In: Anderson, P.D.; Ronnenberg, K.L., eds. *Density management in the 21st century: west side story*. Gen. Tech. Rep. PNW-GTR-880. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 220–238. doi:10.2737/pnw-gtr-880.
- Olson, D.H.; Burton, J.I. 2014.** Near-term effects of repeated-thinning with riparian buffers on headwater stream vertebrates and habitats in Oregon, USA. *Forests*. 5: 2703–2729.

- Olson, D.H.; Crisafulli, C.M. 2014.** Conservation assessment for the Van Dyke's salamander (*Plethodon vandykei*). [Place of publication unknown]: U.S. Department of Agriculture, Forest Service Region 6 and U.S. Department of the Interior, Bureau of Land Management. 56 p. <http://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-amphibians.shtml>. (5 December 2017).
- Olson, D.H.; Kluber, M.R.; Kluber, M.R. 2014a.** Plethodontid salamander distributions in managed forest headwaters in western Oregon, USA. *Herpetological Conservation and Biology*. 9(1): 76–96.
- Olson, D.H.; Leirness, J.B.; Cunningham, P.G.; Ashley Steel, E. 2014b.** Riparian buffers and forest thinning: effects on headwater vertebrates 10 years after thinning. *Forest Ecology and Management*. 321: 81–93. doi:10.1016/j.foreco.2013.06.013.
- Olson, D.H.; Rugger, C. 2007.** Preliminary study of the effects of headwater riparian reserves with upslope thinning on stream habitats and amphibians in western Oregon. *Forest Science*. 53(2): 331–342.
- Olson, D.H.; Van Norman, K.J.; Huff, R.D. 2007b.** The utility of strategic surveys for rare and little-known species under the Northwest Forest Plan. Gen. Tech. Rep. PNW-GTR-708. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 48 p.
- Otálora, M.A.G.; Jørgensen, P.M.; Wedin, M. 2014.** A revised generic classification of the jelly lichens, Collemataceae. *Fungal Diversity*. 64(1): 275–293. doi:10.1007/s13225-013-0266-1.
- Overton, C.T.; Schmitz, R.A.; Casazza, M.L. 2006.** Linking landscape characteristics to mineral site use by band-tailed pigeons in western Oregon: coarse-filter conservation with fine-filter tuning. *Natural Areas Journal*. 26: 38–46.
- Pace, M.L.; Carpenter, S.R.; Cole, J.J. 2015.** With and without warning: managing ecosystems in a changing world. *Frontiers in Ecology and Evolution*. 13(9): 460–467.
- Parsons, G.L.; Cassis, G.; Moldenke, A.R.; Lattin, J.D.; Anderson, N.H.; Miller, J.C.; Hammond, P.; Schowalter, T.D. 1991.** Invertebrates of the H.J. Andrews Experimental Forest, western Cascade Range, Oregon. V: An annotated list of insects and other arthropods. Gen. Tech. Rep. PNW-GTR-290. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 168 p.
- Pauli, J.N.; Whiteman, J.P.; Marcot, B.G.; McClean, T.M.; Ben-David, M. 2011.** DNA-based approach to aging martens (*Martes americana* and *M. caurina*). *Journal of Mammalogy*. 92(3): 500–510.
- Pearl, C.A.; Adams, M.J.; Bury, R.B.; McCreary, B. 2004.** Asymmetrical effects of introduced bullfrogs (*Rana catesbeiana*) on native ranid frogs in Oregon. *Copeia*. 2004(1): 11–20. doi:10.1643/ce-03-010r2.
- Pearl, C.A.; Bowerman, J.; Adams, M.J.; Chelgren, N.D. 2009.** Widespread occurrence of the chytrid fungus *Batrachochytrium dendrobatidis* on Oregon spotted frogs (*Rana pretiosa*). *EcoHealth*. 6(2): 209–218. doi:10.1007/s10393-009-0237-x.
- Pearl, C.A.; Bull, E.L.; Green, D.E.; Bowerman, J.; Adams, M.J.; Hyatt, A.; Wente, W.H. 2007.** Occurrence of the amphibian pathogen *Batrachochytrium dendrobatidis* in the Pacific Northwest. *Journal of Herpetology*. 41: 145–149.
- Peck, J.E. 2006.** Towards sustainable commercial moss harvest in the Pacific Northwest of North America. *Biological Conservation*. 128(3): 289–297. doi:10.1016/j.biocon.2005.10.001.
- Peck, J.E.; Christy, J.A. 2006.** Putting the stewardship concept into practice: commercial moss harvest in northwestern Oregon, USA. *Forest Ecology and Management*. 225(1–3): 225–233. doi:10.1016/j.foreco.2005.12.054.
- Peck, J.E.; Frelich, L.E. 2008.** Moss harvest truncates the successional development of epiphytic bryophytes in the Pacific Northwest. *Ecological Applications*. 18(1): 146–158. doi:10.1890/07-0145.1.

- Peck, J.E.; Moldenke, A.R. 2011.** Invertebrate communities of subcanopy epiphyte mats subject to commercial moss harvest. *Journal of Insect Conservation*. 15(5): 733–742.
- Perault, D.R.; Lomolino, M.V. 2000.** Corridors and mammal community structure across a fragmented, old-growth forest landscape. *Ecological Monographs*. 70(3): 401–422.
- Perry, R.W.; Wigley, T.B.; Melchior, M.A.; Thill, R.E.; Tappe, P.A.; Miller, D.A. 2011.** Width of riparian buffer and structure of adjacent plantations influence occupancy of conservation priority birds. *Biodiversity and Conservation*. 20(3): 625–642.
- Peter, D.; Shebitz, D. 2006.** Historic anthropogenically maintained bear grass savannas of the southeastern Olympic Peninsula. *Restoration Ecology*. 14(4): 605–615. doi:10.1111/j.1526-100x.2006.00172.x.
- Pfeifer-Meister, L.; Bridgman, S.D.; Little, C.J.; Reynolds, L.L.; Goklany, M.E.; Johnson, B.R. 2013.** Pushing the limit: experimental evidence of climate effects on plant range distributions. *Ecology*. 94(10): 2131–2137. doi:10.1890/13-0284.1.
- Piovia-Scott, J.; Pope, K.; Joy Worth, S.; Rosenblum, E.B.; Poorten, T.; Refsnider, J.; Rollins-Smith, L.A.; Reinert, L.K.; Wells, H.L.; Rejmanek, D.; Lawler, S.; Foley, J. 2014.** Correlates of virulence in a frog-killing fungal pathogen: evidence from a California amphibian decline. *The ISME Journal*. 9(7): 1570–1578. doi:10.1038/ismej.2014.241.
- Pollett, K.L.; Maccracken, J.G.; Macmahon, J.A. 2010.** Stream buffers ameliorate the effects of timber harvest on amphibians in the Cascade Range of southern Washington, USA. *Forest Ecology and Management*. 260(6): 1083–1087. doi:10.1016/j.foreco.2010.06.035.
- Pollock, M.M.; Beechie, T.J. 2014.** Does riparian forest restoration thinning enhance biodiversity? The ecological importance of large wood. *Journal of the American Water Resources Association*. 50(3): 543–559. doi:10.1111/jawr.12206.
- Pope, K.; Brown, C.; Hayes, M.; Green, G.; Macfarlane, D. 2014.** Cascades frog conservation assessment. Gen. Tech. Rep. PSW-GTR-244. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 116 p. doi:10.2737/psw-gtr-244.
- Poulin, J.F.; Villard, M.A. 2011.** Edge effect and matrix influence on the nest survival of an old forest specialist, the brown creeper (*Certhia americana*). *Landscape Ecology*. 26(7): 911–922.
- Preston, K.L.; Rotenberry, J.T.; Redak, R.A.; Allen, M.F. 2008.** Habitat shifts of endangered species under altered climate conditions: importance of biotic interactions. *Global Change Biology*. 116 p. doi:10.1111/j.1365-2486.2008.01671.x.
- Price, A.L.; Mowdy, J.S.; Swingle, J.K.; Forsman, E.D. 2015.** Distribution and abundance of tree voles in the northern Coast Ranges of Oregon. *Northwestern Naturalist*. 96: 37–49.
- Price, K.; Hochachka, G. 2001.** Epiphytic lichen abundance: effects of stand age and composition in coastal British Columbia. *Ecological Applications*. 11(3): 904–913.
- Proulx, G.; Santos-Reis, M. 2012.** A century of change in research and management on the genus *Martes*. In: Aubry, K.B.; Zielinski, W.J.; Raphael, M.G.; Proulx, G.; Buskirk, S.W., eds. *Biology and conservation of martens, sables, and fishers: a new synthesis*. Ithaca, NY: Cornell University Press: 471–489.
- Puettmann, K.J.; Dodson, E.K.; Ares, A.; Berger, C. 2013.** Over- and understory responses to thinning treatments: Can we accelerate late successional stand structures? In: Anderson, P.D.; Ronneberg, K.L., eds. *Density management in the 21st century: west side story*. Gen. Tech. Rep. PNW-GTR-880. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 44–58.
- Pyle, R.M. 2002.** *The butterflies of Cascadia*. Seattle, WA: Seattle Audubon Society. 420 p.

- Raley, C.M.; Aubry, K.B. 2006.** Foraging ecology of pileated woodpeckers in coastal forests of Washington. *Journal of Wildlife Management*. 70(5): 1266–1275.
- Raley, C.M.; Lofroth, E.C.; Truex, R.L.; Scott Yaeger, J.; Mark Higley, J. 2012.** Habitat ecology of fishers in western North America: a new synthesis. In: Aubry, K.B.; Zielinski, W.J.; Raphael, M.G.; Proulx, G.; Buskirk, S.W., eds. *Biology and conservation of martens, sables, and fishers*. Ithaca, NY: Cornell University Press: 231–254.
- Rambo, T.R.; Muir, P.S. 1998.** Bryophyte species association with coarse woody debris and stand ages in Oregon. *The Bryologist*. 100(3): 366–376.
- Rambo, T.R.; Muir, P.S. 2002.** Forest floor bryophytes of *Pseudotsuga menziesii*-*Tsuga heterophylla* stands in Oregon: influences of substrate and overstory. *The Bryologist*. 101(1): 116–130.
- Ransom, T.S. 2012.** Behavioral responses of a native salamander to native and invasive earthworms. *Biological Invasions*. 14(12): 2601–2616. doi:10.1007/s10530-012-0255-4.
- Raphael, M. 1988.** Long-term trends in abundance of amphibians, reptiles, and mammals in Douglas-fir forests of northwest California. In: Szaro, R.C.; Severson, K.E.; Patton, D.R., eds. *Management of amphibians, reptiles, and small mammals in North America*. Gen. Tech. Rep. RM-GTR-166. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 23–31.
- Raphael, M.G.; Bisson, P.A.; Lawrence, C.C.; Foster, D. 2002.** Effects of stream-side forest management on the composition and density of stream and riparian fauna of the Olympic Peninsula. In: Johnson, A.C.; Haynes, R.W.; Monserud, R.A., eds. *Congruent management of multiple resources: proceedings from the Wood Compatibility Initiative workshop*. Gen. Tech. Rep. PNW-GTR-563. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 27–40. <http://dx.doi.org/10.2737/pnw-gtr-563>. (5 December 2017).
- Raphael, M.G.; Jones, L.C. 1997.** Characteristics of resting and denning sites of American martens in central Oregon and western Washington. In: Bryant, H.N.; Proulx, G.; Woodard, P.M., eds. *Martes: taxonomy, ecology, techniques, and management: proceedings of the second international Martes symposium*. Edmonton, Canada: Provincial Museum of Alberta. <http://dx.doi.org/10.5962/bhl.title.102822>. (5 December 2017).
- Raphael, M.G.; Marcot, B.G. 1986.** Validation of a wildlife-habitat-relationships model: vertebrates in a Douglas-fir sere. In: Verner, J.; Morrison, M.L.; Ralph, C.J., eds. *Wildlife 2000: modeling habitat relationships of terrestrial vertebrates*. Madison, WI: University of Wisconsin Press: 129–138.
- Raphael, M.G.; Marcot, B.G. 2007.** Introduction. In: Raphael, M.G.; Molina, R., eds. *Conservation of rare or little-known species: biological, social, and economic considerations*. Washington, DC: Island Press: 1–16.
- Raphael, M.G.; Molina, R.; Flather, C.H.; Holthausen, R.; Johnson, R.L.; Marcot, B.G.; Olson, D.H.; Peine, J.D.; Sieg, C.H.; Swanson, C.S. 2007.** A process for selection and implementation of conservation approaches. In: Raphael, M.G.; Molina, R., eds. *Conservation of rare or little-known species: biological, social, and economic considerations*. Washington, DC: Island Press: 334–362.
- Reeves, G.H.; Williams, J.E.; Burnett, K.M.; Gallo, K. 2006.** The aquatic conservation strategy of the Northwest Forest Plan. *Conservation Biology*. 20(2): 319–329. doi:10.1111/j.1523-1739.2006.00380.x.
- Richardson, J.S.; Neill, W.E. 1998.** Headwater amphibians and forestry in British Columbia: Pacific giant salamanders and tailed frogs. *Northwest Science*. 72(Special issue): 122–123.
- Richgels, K.L.D.; Russell, R.E.; Adams, M.J.; White, C.L.; Grant, E.H.C. 2016.** Spatial variation in risk and consequence of *Batrachochytrium salamandrivorans* introduction in the USA. *Royal Society Open Science*. 3(2): 150616. doi:10.1098/rsos.150616.

- Rodhouse, T.J.; Ormsbee, P.C.; Irvine, K.M.; Vierling, L.A.; Szewczak, J.M.; Vierling, K.T. 2012.** Assessing the status and trend of bat populations across broad geographic regions with dynamic distribution models. *Ecological Applications*. 22(4): 1098–1113.
- Rodhouse, T.J.; Ormsbee, P.C.; Irvine, K.M.; Vierling, L.A.; Szewczak, J.M.; Vierling, K.T. 2015.** Establishing conservation baselines with dynamic distribution models for bat populations facing imminent decline. *Diversity and Distributions*. 21(12): 1401–1413.
- Root, H.T.; Geiser, L.H.; Jovan, S.; Neitlich, P. 2015.** Epiphytic macrolichen indication of air quality and climate in interior forested mountains of the Pacific Northwest, USA. *Ecological Indicators*. 53: 95–105. doi:10.1016/j.ecolind.2015.01.029.
- Root, H.T.; McCune, B.; Neitlich, P. 2010.** Lichen habitat May be enhanced by thinning treatments in young *Tsuga heterophylla*–*Pseudotsuga menziesii* forests. *The Bryologist*. 113(2): 292–307. doi:10.1639/0007-2745-113.2.292.
- Rosenberg, D.K.; Davis, R.J.; Van Norman, K.J.; Dunk, J.R.; Forsman, E.D.; Huff, R.D. 2016.** Patterns of red tree vole distribution and habitat suitability: implications for surveys and conservation planning. *Ecosphere*. 7(12): 1–24. doi:10.1002/ecs2.1630.
- Rosenberg, D.K.; Swindle, K.A.; Anthony, R.G. 1994.** Habitat associations of California red-backed voles in young and old-growth forests in western Oregon. *Northwest Science*. 68(4): 266–272.
- Ross, J.A.; Matter, S.F.; Roland, J. 2005.** Edge avoidance and movement of the butterfly *Parnassius smintheus* in matrix and non-matrix habitat. *Landscape Ecology*. 20(2): 127–135.
- Roth, B. 2015.** Range of *Pristiloma crateris* Pilsbry, 1946 (Gastropoda: Pulmonata: Pristilomatidae) in the United States Pacific Northwest. *Check List*. 11(2): 1571. doi:10.15560/11.2.1571.
- Roth, B.; Jadin, R.; Jadin, R. 2013.** The taxonomic status of *Deroceras hesperium* Pilsbry, 1944 (Gastropoda: Pulmonata: Agriolimacidae), a species of conservation concern in Oregon, USA. *Zootaxa*. 3691(4): 453. doi:10.11646/zootaxa.3691.4.4.
- Rowe, K.C.; Rowe, K.M.C.; Tingley, M.W.; Koo, M.S.; Patton, J.L.; Conroy, C.J.; Perrine, J.D.; Beissinger, S.R.; Moritz, C. 2015.** Spatially heterogeneous impact of climate change on small mammals of montane California. *Proceedings of the Royal Society B: Biological Sciences*. 282(1799): 20141857–20141857. doi:10.1098/rspb.2014.1857.
- Royle, J.A.; Chandler, R.B.; Sollmann, R.; Gardner, B. 2014.** Spatial capture-recapture. Waltham, MA: Academic Press. 612 p.
- Royle, J.A.; Stanley, T.R.; Lukacs, P.M. 2008.** Statistical modeling and inference from carnivore survey data. In: Long, R.A.; Mackay, P.; Zielinski, W.J.; Ray, J.C., eds. *Noninvasive survey methods for carnivores*. Washington, DC: Island Press: 293–312.
- Ruchty, A.M. 2000.** The association of epiphytic macrolichens and bryophytes with riparian stand types along a valley continuum, Oregon Coast Range. Oregon State University. 109 p. M.S. thesis.
- Rudnick, D.A.; Ryan, S.J.; Beier, P.; Cushman, S.A.; Dieffenbach, F.; Epps, C.W.; Gerber, L.R.; Hartter, J.; Jenness, J.S.; Kintsch, J.; Merenlender, A.M.; Perkl, R.M.; Preziosi, D.V.; Trombulak, S.C. 2012.** The role of landscape connectivity in planning and implementing conservation and restoration practices. *Issues in Ecology*. 16: 21.
- Ruggiero, L.F.; Pearson, E.; Henry, S.E. 1998.** Characteristics of American marten den sites in Wyoming. *Journal of Wildlife Management*. 62(2): 663. doi:10.2307/3802342.
- Rundio, D.E.; Olson, D.H. 2007.** Influence of headwater site conditions and riparian buffers on terrestrial salamander response to forest thinning. *Forest Science*. 53(2): 320–330.

- Ryan, M.W.; Fraser, D.F.; Marshall, V.G.; Pollard, D.W.F. 1998.** Differences in the composition of vascular plants, bryophytes, and lichens among four successional stages on southern Vancouver Island. *Northwest Science*. 72(Special issue): 86–88.
- Rykken, J.J.; Moldenke, A.R.; Olson, D.H. 2007.** Headwater riparian forest-floor invertebrate communities associated with alternative forest management practices. *Ecological Applications*. 17(4): 1168–1183. doi:10.1890/06-0901.
- Sarr, D.A.; Duff, A.; Dinger, E.C.; Shafer, S.L.; Wing, M.; Seavy, N.E.; Alexander, J.D. 2015.** Comparing ecoregional classifications for natural areas management in the Klamath region, USA. *Natural Areas Journal*. 35(3): 360–377. doi:10.3375/043.035.0301.
- Sato, J.J.; Wolsan, M.; Prevosti, F.J.; D'Elia, G.; Begg, C.; Begg, K.; Hosoda, T.; Campbell, K.L.; Suzuki, H. 2012.** Evolutionary and biogeographic history of weasel-like carnivorans (Musteloidea). *Molecular Phylogenetics and Evolution*. 63(3): 745–757. doi:10.1016/j.ympev.2012.02.025.
- Scheller, R.M.; Spencer, W.D.; Rustigian-Romsos, H.; Syphard, A.D.; Ward, B.C.; Strittholt, J.R. 2011.** Using stochastic simulation to evaluate competing risks of wildfires and fuels management on an isolated forest carnivore. *Landscape Ecology*. 26(10): 1491–1504. doi:10.1007/s10980-011-9663-6.
- Schmiedel, D.; Wilhelm, E.G.; Roth, M.; Scheibner, C.; Nehring, S.; Winter, S. 2016.** Evaluation system for management measures of invasive alien species. *Biodiversity and Conservation*. 25(2): 357–374.
- Schowalter, T.D.; Zhang, Y.L.; Rykken, J.J. 2003.** Litter invertebrate responses to variable density thinning in western Washington forest. *Ecological Applications*. 13(2): 1204–1211.
- Schultz, C.A.; Sisk, T.D.; Noon, B.R.; Nie, M.A. 2013.** Wildlife conservation planning under the United States Forest Service's 2012 planning rule. *Journal of Wildlife Management*. 77: 428–444. doi:10.1002/jwmg.513.
- Schumaker, N.H. 2013.** HexSim. Version 2.4.5. Corvallis, OR: U.S. Environmental Protection Agency, Environmental Research Laboratory. <http://hexsim.net>. (13 November 2017).
- Schumaker, N.H.; Brookes, A.; Dunk, J.R.; Woodbridge, B.; Heinrichs, J.A.; Lawler, J.J.; Carroll, C.; Laplante, D. 2014.** Mapping sources, sinks, and connectivity using a simulation model of northern spotted owls. *Landscape Ecology*. 29(4): 579–592. doi:10.1007/s10980-014-0004-4.
- Schwalm, D.; Epps, C.W.; Rodhouse, T.J.; Monahan, W.B.; Vardaro, J.A.C.; Ray, C.; Jeffress, M.R. 2015.** Habitat availability and gene flow influence diverging local population trajectories under scenarios of climate change: a place-based approach. *Global Change Biology*. 22(4): 1572–1584. doi:10.1111/gcb.13189.
- Schwartz, M.K.; Copeland, J.P.; Anderson, N.J.; Squires, J.R.; Inman, R.M.; McKelvey, K.S.; Pilgrim, K.L.; Waits, L.P.; Cushman, S.A. 2009.** Wolverine gene flow across a narrow climatic niche. *Ecology*. 90(11): 3222–3232. doi:10.1890/08-1287.1.
- Schwartz, M.K.; Monfort, S.L. 2008.** Genetic and endocrine tools for carnivore surveys. In: Long, R.A.; Mackay, P.; Zielinski, W.J.; Ray, J.C., eds. *Noninvasive survey methods for carnivores*. Washington, DC: Island Press: 238–262.
- Sedell, J.R.; Froggatt, J.L. 1984.** Importance of streamside forests to large rivers: the isolation of the Willamette River, Oregon, USA, from its floodplain by snagging and streamside forest removal. *Verhandlungen des Internationalen Verein Limnologie*. 22: 1828–1834.

- Seibold, S.; Bässler, C.; Brandl, R.; Gossner, M.M.; Thorn, S.; Ulyshen, M.D.; Müller, J. 2015.** Experimental studies of dead-wood biodiversity—a review identifying global gaps in knowledge. *Biological Conservation*. 191: 139–149.
- Semlitsch, R.D.; Bodie, J.R. 2003.** Biological criteria for buffer zones around wetlands and riparian habitats for amphibians and reptiles. *Conservation Biology*. 17(5): 1219–1228.
- Sergio, F.; Pedrini, P.; Marchesi, L. 2003.** Reconciling the dichotomy between single species and ecosystem conservation: black kites (*Milvus migrans*) and eutrophication in pre-alpine lakes. *Biological Conservation*. 110(1): 101–111.
- Sheehan, T.; Bachelet, D.; Ferschweiler, K. 2015.** Projected major fire and vegetation changes in the Pacific Northwest of the conterminous United States under selected CMIP5 climate futures. *Ecological Modelling*. 317: 16–29.
- Shields, J.M.; Webster, C.R.; Storer, A.J. 2008.** Short-term community-level response of arthropods to group selection with seed-tree retention in a northern hardwood forest. *Forest Ecology and Management*. 255: 129–139.
- Shirk, A.J.; Raphael, M.G.; Cushman, S.A. 2014.** Spatiotemporal variation in resource selection: insights from the American marten (*Martes americana*). *Ecological Applications*. 24(6): 1434–1444.
- Shoo, L.P.; Olson, D.H.; McMenamin, S.K.; Murray, K.A.; Van Sluys, M.; Donnelly, M.A.; Stratford, D.; Terhivuo, J.; Merino-Viteri, A.; Herbert, S.M.; Bishop, P.J.; Corn, P.S.; Dovey, L.; Griffiths, R.A.; Lowe, K.; Mahony, M.; McCallum, H.; Shuker, J.D.; Simpkins, C.; Skerratt, L.F.; Williams, S.E.; Hero, J.-M. 2011.** Engineering a future for amphibians under climate change. *Journal of Applied Ecology*. 48(2): 487–492. doi:10.1111/j.1365-2664.2010.01942.x.
- Sillett, S.C.; McCune, B.; Peck, J.E.; Rambo, T.R.; Ruchty, A. 2000.** Dispersal limitations of epiphytic lichens result in species dependent on old-growth forests. *Ecological Applications*. 10(3): 789–799.
- Simberloff, D. 1998.** Flagships, umbrellas, and keystones: is single-species management passé in the landscape era? *Biological Conservation*. 83(3): 247–257. doi:10.1016/s0006-3207(97)00081-5.
- Singleton, P.H.; Gaines, W.L.; Lehmkuhl, J.F. 2002.** Landscape permeability for large carnivores in Washington: a geographic information system weighted-distance and least-cost corridor assessment. Res. Pap. PNW-RP-549. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 89 p.
- Siuslaw National Forest. 2007.** Decision notice—Siuslaw Commercial Special Forest Products Program. Corvallis, OR: U.S. Department of Agriculture, Forest Service. 12 p.
- Skerratt, L.F.; Berger, L.; Speare, R.; Cashins, S.; McDonald, K.R.; Phillott, A.D.; Hines, H.B.; Kenyon, N. 2007.** Spread of chytridiomycosis has caused the rapid global decline and extinction of frogs. *EcoHealth*. 4: 125–134.
- Slauson, K.; Baldwin, J.; Zielinski, W. 2012.** Occupancy estimation and modeling in *Martes* research and monitoring. In: Aubry, K.B.; Zielinski, W.J.; Raphael, M.G.; Proulx, G.; Buskirk, S.W., eds. *Biology and conservation of martens, sables, and fishers: a new synthesis*. Ithaca, NY: Cornell University Press: 343–368.
- Slauson, K.M.; Schmidt, G.A.; Zielinski, W.J.; Detrich, P.J.; Callas, R.L.; Thrailkill, J.; Devlin Craig, B.; Early, D.A.; Hamm, K.A.; Schmidt, K.N.; Transou, A.; West, C.J. [In press].** A conservation assessment and strategy for the Humboldt marten (*Martes caurina humboldtensis*) in California and Oregon. Gen. Tech. Rep. Arcata, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station.

- Slauson, K.M.; Zielinski, W.J. 2009.** Characteristics of summer and fall diurnal resting habitat used by American martens in coastal northwestern California. *Northwest Science*. 83(1): 35–45. doi:10.3955/046.083.0104.
- Slauson, K.M.; Zielinski, W.J.; Hayes, J.P. 2007.** Habitat selection by American martens in coastal California. *Journal of Wildlife Management*. 71(2): 458–468. doi:10.2193/2004-332.
- Slauson, K.M.; Zielinski, W.J.; Schwartz, M.K. 2017.** Ski areas affect Pacific marten movement, habitat use, and density. *Journal of Wildlife Management*. 81(5): 892–904. doi:10.1002/jwmg.21243.
- Slauson, K.M.; Zielinski, W.J.; Stone, K.D. 2009.** Characterizing the molecular variation among American marten (*Martes americana*) subspecies from Oregon and California. *Conservation Genetics*. 10(5): 1337–1341. doi:10.1007/s10592-008-9626-x.
- Smart, A.S.; Tingley, R.; Weeks, A.R.; Van Rooyen, A.R.; McCarthy, M.A. 2015.** Environmental DNA sampling is more sensitive than a traditional survey technique for detecting an aquatic invader. *Ecological Applications*. 25(7): 1944–1952.
- Smith, J.E.; Molina, R.; Huso, M.M.; Luoma, D.L.; McKay, D.; Castellano, M.A.; Lebel, T.; Valachovic, Y. 2002.** Species richness, abundance, and composition of hypogeous and epigeous ectomycorrhizal fungal sporocarps in young, rotation-age, and old-growth stands of Douglas-fir (*Pseudotsuga menziesii*) in the Cascade range of Oregon, U.S.A. *Canadian Journal of Botany*. 80(2): 186–204. doi:10.1139/b02-003.
- Sollmann, R.; White, A.M.; Tarbill, G.L.; Manley, P.N.; Knapp, E.E. 2016.** Landscape heterogeneity compensates for fuel reduction treatment effects on northern flying squirrel populations. *Forest Ecology and Management*. 373: 100–107. doi:10.1016/j.foreco.2016.04.041.
- Spies, T.A.; Giesen, T.W.; Swanson, F.J.; Franklin, J.F.; Lach, D.; Johnson, K.N. 2010.** Climate change adaptation strategies for federal forests of the Pacific Northwest, USA: ecological, policy, and socio-economic perspectives. *Landscape Ecology*. 25(8): 1185–1199. doi:10.1007/s10980-010-9483-0.
- Spivak, M.; Mader, E.; Vaughan, M.; Euliss, N.H., Jr. 2011.** The plight of bees. *Environmental Science and Technology*. 45: 34–38.
- Squires, J.R.; Decesare, N.J.; Kolbe, J.A.; Ruggiero, L.F. 2008.** Hierarchical den selection of Canada lynx in western Montana. *Journal of Wildlife Management*. 72(7): 1497. doi:10.2193/2007-396.
- Stankey, G.H.; Bormann, B.T.; Clark, R.N. 2006.** Learning to manage a complex ecosystem: adaptive management and the Northwest Forest Plan. Res. Pap. PNW-RP-567. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 194 p.
- Stankey, G.H.; Shindler, B. 2006.** Formation of social acceptability judgments and their implications for management of rare and little-known species. *Conservation Biology*. 20(1): 28–37.
- Staus, N.L.; Strittholt, J.R.; DellaSala, D.A. 2010.** Evaluating areas of high conservation value in western Oregon with a decision-support model. *Conservation Biology*. 24(3): 711–720. doi:10.1111/j.1523-1739.2010.01445.x.
- Stoddard, M.A.; Hayes, J.P. 2005.** The influence of forest management on headwater stream amphibians at multiple spatial scales. *Ecological Applications*. 15(3): 811–823.
- Suzuki, N.; Olson, D.H. 2008.** Options for biodiversity conservation in managed forest landscapes of multiple ownerships in Oregon and Washington, USA. *Biodiversity and Conservation*. 17(5): 1017–1039.

- Suzuki, N.; Olson, D.H.; Reilly, E.C. 2008.** Developing landscape habitat models for rare amphibians with small geographic ranges: a case study of Siskiyou mountains salamanders in the western USA. *Biodiversity and Conservation*. 17(9): 2197–2218. doi:10.1007/s10531-007-9281-4.
- Swanson, M.E.; Franklin, J.F.; Beschta, R.L.; Crisafulli, C.M.; DellaSala, D.A.; Hutto, R.L.; Lindenmayer, D.B.; Swanson, F.J. 2011.** The forgotten stage of forest succession: early-successional ecosystems on forest sites. *Frontiers in Ecology and the Environment*. 9(2): 117–125. doi:10.1890/090157.
- Swanson, M.E.; Studevant, N.M.; Campbell, J.L.; Donato, D.C. 2014.** Biological associates of early-seral pre-forest in the Pacific Northwest. *Forest Ecology and Management*. 324: 160–171. doi:10.1016/j.foreco.2014.03.046.
- Sweitzer, R.A.; Furnas, B.J.; Barrett, R.H.; Purcell, K.L.; Thompson, C.M. 2016a.** Landscape fuel reduction, forest fire, and biophysical linkages to local habitat use and local persistence of fishers (*Pekania pennanti*) in sierra Nevada mixed-conifer forests. *Forest Ecology and Management*. 361: 208–225.
- Sweitzer, R.A.; Thompson, C.M.; Green, R.E.; Barrett, R.H.; Purcell, K.L. 2016.** Survival of fishers in the southern Sierra Nevada region of California. *Journal of Mammalogy*. 97(1): 274–286.
- Swingle, J.K.; Forsman, E.D. 2009.** Home range areas and activity patterns of red tree voles (*Arborimus longicaudus*) in western Oregon. *Northwest Science*. 83(3): 273–286.
- Swingle, J.K.; Forsman, E.D. 2016.** Annotated bibliography of the red tree vole (*Arborimus longicaudus*), Sonoma tree vole (*A. pomio*), and white-footed vole (*A. albipes*). Gen. Tech. Rep. PNW-GTR-909. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 179 p.
- Swingle, J.K.; Forsman, E.D.; Anthony, R.G. 2010.** Survival, mortality, and predators of red tree voles (*Arborimus longicaudus*). *Northwest Science*. 84(3): 255–265. doi:10.3955/046.084.0305.
- Thomas, J.W.; Franklin, J.F.; Gordon, J.; Johnson, K.N. 2006.** The Northwest Forest Plan: origins, components, implementation experience, and suggestions for change. *Conservation Biology*. 20(2): 277–287. doi:10.1111/j.1523-1739.2006.00385.x.
- Thompson, I.D.; Fryxell, J.; Harrison, D.J. 2012.** Improved insights into use of habitat by American martens. In: Aubry, K.B.; Zielinski, W.J.; Raphael, M.G.; Proulx, G.; Buskirk, S.W., eds. *Biology and conservation of martens, sables, and fishers*. Ithaca, NY: Cornell University Press: 209–230. doi:10.7591/9780801466076-012.
- Thompson, J.L. 2008.** Density of fisher on managed timberlands in north coastal California. Arcata, CA: Humboldt State University. 40 p. M.S. thesis.
- Thompson, J.R.; Duncan, S.L.; Johnson, K.N. 2009.** Is there potential for the historical range of variability to guide conservation given the social range of variability? *Ecology and Society*. 14(1): 18. <http://www.ecologyandsociety.org/vol14/iss1/art18/>. (6 December 2017).
- Thompson, M.P.; Marcot, B.G.; Thompson, F.R.; McNulty, S.; Fisher, L.A.; Runge, M.C.; Cleaves, D.; Tomosy, M. 2013.** The science of decision making: applications for sustainable forest and grassland management in the National Forest System. Gen. Tech. Rep. WO-GTR-88. Washington, DC: U.S. Department of Agriculture, Forest Service. 54 p.
- Thomsen, P.F.; Kielgast, J.; Iversen, L.L.; Wiuf, C.; Rasmussen, M.; Gilbert, M.T.P.; Orlando, L.; Willerslev, E. 2012.** Monitoring endangered freshwater biodiversity using environmental DNA. *Molecular Ecology*. 21(11): 2565–2573.
- Tigner, J.; Bayne, E.M.; Boutin, S. 2015.** American marten respond to seismic lines in northern Canada at two spatial scales. *PLoS ONE*. 10(3): e0118720. doi:10.1371/journal.pone.0118720.

- Tingley, M.W.; Koo, M.S.; Moritz, C.; Rush, A.C.; Beissinger, S.R. 2012.** The push and pull of climate change causes heterogeneous shifts in avian elevational ranges. *Global Change Biology*. 18(11): 3279–3290. doi:10.1111/j.1365-2486.2012.02784.x.
- Tolkkinen, M.; Mykrä, H.; Annala, M.; Markkola, A.M.; Vuori, K.M.; Muotka, T. 2015.** Multi-stressor impacts on fungal diversity and ecosystem functions in streams: natural vs. anthropogenic stress. *Ecology*. 96(3): 672–683.
- Trappe, J.M.; Molina, R.; Luoma, D.L.; Cázares, E.; Pilz, D.; Smith, J.E.; Castellano, M.A.; Miller, S.L.; Trappe, M.J. 2009.** Diversity, ecology, and conservation of truffle fungi in forests of the Pacific Northwest. Gen. Tech. Rep. PNW-GTR-772. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 35 p.
- Trofymow, J.A.; Addison, J.; Blackwell, B.A.; He, F.; Preston, C.A.; Marshall, V.G. 2003.** Attributes and indicators of old-growth and successional Douglas-fir forests on Vancouver Island. *Environmental Reviews*. 11(S1): S187-S204. doi:10.1139/a03-007.
- Truex, R.L.; Zielinski, W.J. 2013.** Short-term effects of fuel treatments on fisher habitat in the Sierra Nevada, California. *Forest Ecology and Management*. 293: 85–91. doi:10.1016/j.foreco.2012.12.035.
- Trumbo, D.R.; Spear, S.F.; Baumsteiger, J.; Storfer, A. 2013.** Rangewide landscape genetics of an endemic Pacific northwestern salamander. *Molecular Ecology*. 22(5): 1250–1266. doi:10.1111/mec.12168.
- Truong, C.; Rodriguez, J.M.; Clerc, P. 2013.** *Pendulous usnea* species (Parmeliaceae, lichenized Ascomycota) in tropical South America and the Galapagos. *The Lichenologist*. 45(04): 505–543. doi:10.1017/s0024282913000133.
- Tucker, J.M. 2013.** Assessing changes in connectivity and abundance through time for fisher in the southern Sierra Nevada. Missoula, MT: University of Montana. 118 p. Ph.D. dissertation.
- Turner, D.P.; Conklin, D.R.; Bolte, J.P. 2015.** Projected climate change impacts on forest land cover and land use over the Willamette River basin, Oregon, USA. *Climatic Change*. doi:10.1007/s10584-015-1465-4.
- U.S. Department of Agriculture, Forest Service [USDA FS]. 2012.** National forest system land management planning. 36 C.F.R. part 219. Federal Register. 77: 21260–21276. <http://www.fs.usda.gov/planningrule>.
- U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management [USDA and USDI]. 1994.** Record of decision for amendments to Forest Service and Bureau of Land Management planning documents within the range of the northern spotted owl. [Place of publication unknown]. 74 p. [plus attachment A: standards and guidelines].
- U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management [USDA and USDI]. 2001.** Record of decision and standards and guidelines for amendments to the survey and manage, protection buffer, and other mitigation measures standards and guidelines in Forest Service and Bureau of Land Management planning documents within the range of the northern spotted owl. Portland, OR. 86 p.
- U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management [USDA and USDI]. 2004.** Record of decision to remove or modify the survey and manage mitigation measure standards and guidelines in Forest Service and Bureau of Land Management planning documents within the range of the northern spotted owl. BLM/OR/WA/PL-04/017-1792. Portland, OR. 44 p.
- U.S. Department of the Interior, Bureau of Land Management [USDI BLM]. 2016.** Resource management planning, Bureau of Land Management, proposed rule. 43 C.F.R. part 1600. Federal Register. 81: 9674–9734.

- U.S. Department of the Interior, Fish and Wildlife Service [USFWS]. 2000.** Endangered and threatened wildlife and plants; determination of threatened status for the contiguous U.S. distinct population segment of the Canada lynx and related rule. Federal Register. 65: 16052–16086.
- U.S. Department of the Interior, Fish and Wildlife Service [USFWS]. 2011.** Endangered and threatened wildlife and plants; 12-month finding on a petition to list a distinct population segment of the red tree vole as endangered or threatened. Federal Register. 76: 63720–63762.
- U.S. Department of the Interior, Fish and Wildlife Service [USFWS]. 2014.** Endangered and threatened wildlife and plants; threatened status for Oregon spotted frog. Federal Register. 79: 51658–51710.
- U.S. Department of the Interior, Fish and Wildlife Service [USFWS]. 2016.** Injurious wildlife species; listing salamanders due to risk of salamander chytrid fungus. Federal Register. 81: 1534–1556.
- Ulyshen, M.D. 2016.** Wood decomposition as influenced by invertebrates. Biological Reviews. 91: 70–85.
- Van Rooij, P.; Martel, A.; Haesebrouck, F.; Pasmans, F. 2015.** Amphibian chytridiomycosis: a review with focus on fungus-host interactions. Veterinary Research. 46(1): 137. doi:10.1186/s13567-015-0266-0.
- Velmala, S.; Myllys, L.; Goward, T.; Holien, H.; Halonen, P. 2014.** Taxonomy of *Bryoria* section *Implexae* (Parmeliaceae, Lecanoromycetes) in North America and Europe, based on chemical, morphological and molecular data. Annales Botanici Fennici. 51(6): 345–371. doi:10.5735/085.051.0601.
- Velo-Antón, G.; Parra, J.L.; Parra-Olea, G.; Zamudio, K.R. 2013.** Tracking climate change in a dispersal-limited species: reduced spatial and genetic connectivity in a montane salamander. Molecular Ecology. 22(12): 3261–3278. doi:10.1111/mec.12310.
- Verant, M.L.; Meteyer, C.U.; Speakman, J.R.; Cryan, P.M.; Lorch, J.M.; Blehert, D.S. 2014.** White-nose syndrome initiates a cascade of physiologic disturbances in the hibernating bat host. BMC Physiology. 14(10). doi:10.1186/s12899-014-0010-4.
- Vergara, P.M. 2011.** Matrix-dependent corridor effectiveness and the abundance of forest birds in fragmented landscapes. Landscape Ecology. 26(8): 1085–1096.
- Vogeler, J.C.; Hudak, A.T.; Vierling, L.A.; Vierling, K.T. 2013.** Lidar-derived canopy architecture predicts brown creeper occupancy of two western coniferous forests. Condor. 115(3): 614–622.
- Vose, J.M.; Peterson, D.L.; Patel-Weynand, T. 2012.** Effects of climatic variability and change on forest ecosystems: a comprehensive science synthesis for the U.S. Forest Service. Gen. Tech. Rep. PNW-GTR-870. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 265 p.
- Wahbe, T.R.; Bunnell, F.L.; Bury, R.B. 2004.** Terrestrial movements of juvenile and adult tailed frogs in relation to timber harvest in coastal British Columbia. Canadian Journal of Forest Research. 34(12): 2455–2466. doi:10.1139/x04-126.
- Waters, J.R.; Zabel, C.J. 1995.** Northern flying squirrel densities in fir forests of northeastern California. Journal of Wildlife Management. 59(4): 858–866.
- Webster, K.M.; Halpern, C.B. 2010.** Long-term vegetation responses to reintroduction and repeated use of fire in mixed-conifer forests of the Sierra Nevada. Ecosphere. 1(5): 9. doi:10.1890/es10-00018.1.
- Weir, R.D.; Phinney, M.; Lofroth, E.C. 2012.** Big, sick, and rotting: why tree size, damage, and decay are important to fisher reproductive habitat. Forest Ecology and Management. 265: 230–240. doi:10.1016/j.foreco.2011.10.043.

- Weller, T.J.; Baldwin, J.A. 2012.** Using echolocation monitoring to model bat occupancy and inform mitigations at wind energy facilities. *Journal of Wildlife Management*. 76(3): 619–631.
- Welsh, H.H., Jr.; Hodgson, G.R. 2008.** Amphibians as metrics of critical biological thresholds in forested headwater streams of the Pacific Northwest, U.S.A. *Freshwater Biology*. 53: 1470–1488.
- Welsh, H.H., Jr.; Hodgson, G.R. 2011.** Spatial relationships in a dendritic network: the herpetofaunal metacommunity of the Mattole River catchment of northwest California. *Ecography*. 34: 49–66.
- Welsh, H.H., Jr.; Hodgson, G.R. 2013.** Woodland salamanders as metrics of forest ecosystem recovery: a case study from California's redwoods. *Ecosphere*. 4(5): Article 59.
- Welsh, H.H.; Pope, K.L.; Wheeler, C.A. 2008.** Using multiple metrics to assess the effects of forest succession on population status: a comparative study of two terrestrial salamanders in the US Pacific Northwest. *Biological Conservation*. 141(4): 1149–1160. doi:10.1016/j.biocon.2008.02.014.
- Welsh, H.H.; Stauffer, H.; Clayton, D.R.; Ollivier, L.M. 2007.** Strategies for modeling habitat relationships of uncommon species: an example using the Siskiyou mountains salamander (*Plethodon stormi*). *Northwest Science*. 81(1): 15–36. doi:10.3955/0029-344x-81.1.15.
- Welsh, H.H., Jr.; Waters, J.R.; Hodgson, G.R.; Weller, T.J.; Zabel, C.J. 2015.** Responses of the woodland salamander *Ensatina eschscholtzii* to commercial thinning by helicopter in late-seral Douglas-fir forest in Northwest California. *Forest Ecology and Management*. 335: 156–165.
- Wengert, G.M. 2013.** Ecology of intraguild predation on fishers (*Martes pennanti*) in California. Davis, CA: University of California. 119 p. Ph.D. dissertation.
- Wengert, G.M.; Gabriel, M.W.; Matthews, S.M.; Higley, J.M.; Sweitzer, R.A.; Thompson, C.M.; Purcell, K.L.; Barrett, R.H.; Woods, L.W.; Green, R.E.; Keller, S.M.; Gaffney, P.M.; Jones, M.; Sacks, B.N. 2014.** Using DNA to describe and quantify interspecific killing of fishers in California. *Journal of Wildlife Management*. 78(4): 603–611. doi:10.1002/jwmg.698.
- Wessell, S.J. 2005.** Biodiversity in managed forests of western Oregon: species assemblages in leave islands, thinned, and unthinned forests. Corvallis, OR: Oregon State University. 74 p. M.S. thesis.
- Wiens, J.A.; Hayward, G.D.; Holthausen, R.S.; Wisdom, M.J. 2008.** Using surrogate species and groups for conservation planning and management. *BioScience*. 58(3): 241–252.
- Wightman, C.S.; Saab, V.A.; Forristal, C.; Mellen-Mclean, K.; Markus, A. 2010.** White-headed woodpecker nesting ecology after wildfire. *Journal of Wildlife Management*. 74(5): 1098–1106. doi:10.2193/2009-174.
- Wilkins, R.N.; Peterson, N.P. 2000.** Factors related to amphibian occurrence and abundance in headwater streams draining second-growth Douglas-fir forests in southwestern Washington. *Forest Ecology and Management*. 139(1–3): 79–91.
- Wilson, D.S.; Puettmann, K.J. 2007.** Density management and biodiversity in young Douglas-fir forests: challenges of managing across scales. *Forest Ecology and Management*. 246(1): 123–134. doi:10.1016/j.foreco.2007.03.052.
- Wilson, E.O. 1987.** The little things that run the world (the importance and conservation of invertebrates). *Conservation Biology*. 1(4): 344–346.

- Wilson, T.M.; Forsman, E.D. 2013.** Thinning effects on spotted owl prey and other forest-dwelling small mammals. In: Anderson, P.D.; Ronnenberg, K.L., eds. Density management for the 21st century: west side story. Gen. Tech. Rep. PNW-GTR-880. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 79–90.
- Wilson, T.M.; Schuller, R.; Holmes, R.; Pavola, C.; Fimbel, R.A.; McCain, C.N.; Gamon, J.G.; Speaks, P.; Seevers, J.I.; Demeo, T.E.; Gibbons, S. 2009.** Interagency strategy for the Pacific Northwest natural areas network. Gen. Tech. Rep. PNW-GTR-798. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 33 p.
- Wimberly, M.C.; Liu, Z. 2014.** Interactions of climate, fire, and management in future forests of the Pacific Northwest. *Forest Ecology and Management*. 327: 270–279. doi:10.1016/j.foreco.2013.09.043.
- Wratten, S.D.; Gillespie, M.; Decourtye, A.; Mader, E.; Desneux, N. 2012.** Pollinator habitat enhancement: benefits to other ecosystem services. *Agriculture, Ecosystems and Environment*. 159: 112–122.
- Yackulic, C.B.; Chandler, R.; Zipkin, E.F.; Royle, J.A.; Nichols, J.D.; Campbell Grant, E.H.; Veran, S. 2013.** Presence-only modelling using MAXENT: When can we trust the inferences? *Methods in Ecology and Evolution*. 4(3): 236–243. doi:10.1111/2041-210x.12004.
- Yap, T.A.; Koo, M.S.; Ambrose, R.F.; Wake, D.B.; Vredenburg, V.T. 2015.** Averting a North American biodiversity crisis. *Science*. 349(6247): 481–482. doi:10.1126/science.aab1052.
- Yi, H.; Moldenke, A. 2005.** Response of ground-dwelling arthropods to different thinning intensities in young Douglas-fir forests of western Oregon. *Environmental Entomology*. 34(5): 1071–1080.
- Zielinski, W.J.; Carroll, C.; Dunk, J.R. 2006.** Using landscape suitability models to reconcile conservation planning for two key forest predators. *Biological Conservation*. 133(4): 409–430. doi:10.1016/j.biocon.2006.07.003.
- Zielinski, W.J.; Dunk, J.R.; Gray, A.N. 2012.** Estimating habitat value using forest inventory data: the fisher (*Martes pennanti*) in northwestern California. *Forest Ecology and Management*. 275: 35–42.
- Zielinski, W.J.; Dunk, J.R.; Yaeger, J.S.; Laplante, D.W. 2010.** Developing and testing a landscape-scale habitat suitability model for fisher (*Martes pennanti*) in forests of interior northern California. *Forest Ecology and Management*. 260(9): 1579–1591. doi:10.1016/j.foreco.2010.08.006.
- Zielinski, W.J.; Thompson, C.M.; Purcell, K.L.; Garner, J.D. 2013.** An assessment of fisher (*Pekania pennanti*) tolerance to forest management intensity on the landscape. *Forest Ecology and Management*. 310: 821–826. doi:10.1016/j.foreco.2013.09.028.
- Ziemba, J.L.; Cameron, A.C.; Peterson, K.; Hickerson, C.A.M.; Anthony, C.D. 2015.** Invasive Asian earthworms of the genus *Amyntas* alter microhabitat use by terrestrial salamanders. *Canadian Journal of Zoology*. 93(10): 805–811.



Molalla River, Oregon.
Photo by Jeff Clark, USDI Bureau of Land Management, Oregon-Washington State Office.

Chapter 7: The Aquatic Conservation Strategy of the Northwest Forest Plan—A Review of the Relevant Science After 23 Years

Gordon H. Reeves, Deanna H. Olson, Steven M. Wondzell, Peter A. Bisson, Sean Gordon, Stephanie A. Miller, Jonathan W. Long, and Michael J. Furniss¹

Introduction

The Aquatic Conservation Strategy (ACS) is a regional strategy applied to aquatic and riparian ecosystems across the area covered by the Northwest Forest Plan (NWFP, or Plan), encompassing broad landscapes of public lands administered by the U.S. Department of Agriculture Forest Service and the U.S. Department of the Interior Bureau of Land Management (BLM) (USDA and USDI 1994a). The ACS was developed during the analysis (FEMAT 1993) that led to the NWFP, but its foundation was a refinement of earlier strategies: the Scientific Panel on Late-Successional Forest Ecosystems (“The Gang of Four”) (Johnson et al. 1991), PacFish (USDA and USDI 1994b), and the Scientific Analysis Team (Thomas et al. 1993).

The ACS uses an ecosystem approach to management of riparian and aquatic habitats (Everest and Reeves 2007) and was designed to (1) protect watersheds that had good-quality habitat and strong fish populations at the time the Plan was drafted, and (2) halt further declines in watershed condition and restore ecological processes that create and maintain favorable conditions in aquatic ecosystems in degraded ecosystems (FEMAT 1993). The long-term goal (100+ years) is

to develop a network of functioning watersheds that supports populations of fish and other aquatic and riparian-dependent organisms across the NWFP area (USDA and USDI 1994a). The ACS is based on preserving and restoring key ecological processes, including the natural disturbance regimes (USDA and USDI 1994a) that create and maintain habitat for native aquatic and riparian-dependent organisms, and it recognizes that periodic natural disturbances may be required to sustain ecological productivity. As a result, the ACS does not expect that all watersheds will be in favorable condition (highly productive for the same aquatic organisms) at any point in time, nor does it expect that any particular watershed will remain in a certain condition through time. If the ACS and the NWFP are effective, the proportion of watersheds in better condition (for native organisms) is expected to remain the same or increase over time (Reeves et al. 2004).

The primary objective of the ACS is to maintain and restore the distribution, diversity, and complexity of watershed-level features and processes to which aquatic and riparian species are uniquely adapted. Programs and actions under the ACS are to maintain, not prevent, attainment of this goal. The ACS designates watershed analysis as the tool for developing baseline conditions against which to assess maintenance and restoration conditions, and improvements in biological and physical processes are to be evaluated relative to the natural range of variability (USDA and USDI 1994a). ACS objectives address (1) diversity and complexity of watershed features; (2) spatial and temporal connectivity within and between watersheds; (3) physical integrity; (4) water quality; (5) sediment input, storage, and transport; (6) instream flows (e.g., both peak and low flows); (7) floodplain inundation; (8) riparian plant-species composition and structural diversity; and (9) habitat to support well-distributed populations of native, aquatic and riparian-dependent species of plants, invertebrates, and vertebrates.

The ACS sets out five components to meet its goals:

- **Riparian reserves:** Riparian reserves are specifically designated portions of the watershed most tightly coupled with streams and rivers that provide

¹ **Gordon H. Reeves** is a research fish ecologist, **Deanna H. Olson** is a research aquatic ecologist, and **Steven M. Wondzell** is a research geologist, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis, OR 97331; **Jonathan W. Long** is a research ecologist, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 1731 Research Park, Davis, CA 95618; **Peter A. Bisson** is a research fish ecologist (retired), U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 3625 93rd Avenue SW, Olympia, WA 98512; **Michael J. Furniss** is a hydrologist (retired), U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 1700 Bayview Drive, Arcata, CA 95521-6013; **Sean Gordon** is a research assistant professor, Institute for Sustainable Solutions, Portland State University, 1600 SW Fourth Avenue, Portland, OR 97204; **Stephanie A. Miller** is the Riparian Program lead, U.S. Department of the Interior, Bureau of Land Management, 20 M Street SE, Washington, DC 20001.

the ecological functions and processes necessary to create and maintain habitat for aquatic and riparian-dependent organisms over time, as well as habitat connectivity within and between watersheds. The reserve boundaries were considered interim until a watershed analysis was completed, at which time they could be modified based on suggestions made in the watershed analysis.

- **Key watersheds:** 5th-code (40,000 to 250,000 ac [16 187 to 101 171 ha]) to 6th-code (10,000 to 40,000 ac [4047 to 16 187 ha]) hydrologic units that were intended to serve as refugia for aquatic organisms, particularly in the short term for at-risk fish populations, and had the greatest potential for restoration, or to provide sources of high-quality water. At the time the NWFP was drafted, Tier 1 key watersheds had strong populations of fish, productive habitat that was in good condition, or high restoration potential. Tier 2 key watersheds provided sources of high-quality water.
- **Watershed analysis:** An analytical process that characterizes the features and processes of watersheds and identifies potential actions for addressing problems and concerns, along with possible management options. It assembles information necessary to determine the ecological characteristics and behavior of the watershed and contribute to the development of options to guide management in the watershed, including adjusting riparian-reserve boundaries.
- **Watershed restoration:** Includes actions deemed necessary to restore degraded ecological processes and habitat. Restoration activities focus on restoring the key ecological processes required to create and maintain favorable environmental conditions for aquatic and riparian-dependent organisms.
- **Standards and guidelines:** These directives impose specific requirements (standards) or recommended approaches (guidelines) for management activities in riparian reserves and key watersheds.

Note that a key philosophical shift occurred in the development of the ACS and NWFP as compared with

efforts prior to 1993. The ACS, along with PACFISH (USDA and USDI 1994b) and the riparian component of the Tongass Land and Resource Management Plan (USDA FS 1997), made two substantive changes in how riparian management was formulated (Everest and Reeves 2007). First, they addressed riparian management at the watershed scale (5th- to 6th-code hydrologic units), with specific emphasis on maintaining ecological functions over the long term. Second, they rejected the previous philosophy of trying to define and achieve the absolute minimum set of practices that would meet stated riparian-management goals, and the concept that goals could be met by implementing yet another set of best management practices. The new (at that time) management philosophy under the NWFP represented a paradigm shift in how managers viewed resource coordination. In previous riparian rule-sets, riparian and aquatic technical specialists shouldered the “burden of proof” to demonstrate resource damage from forestry activities and the need for more comprehensive forest-practices rules to meet riparian-management goals. Under the NWFP, the precautionary principle was invoked—the burden of proof shifted (Thomas et al. 2006, USDA and USDI 1994a). Forest managers who wanted to alter the comprehensive default prescriptions for riparian management under the NWFP (described above) to pursue other management goals were required to demonstrate through watershed analysis that changes would not compromise established riparian-management goals.

This chapter focuses on the scientific literature related to the management and conservation of aquatic ecosystems, particularly as it has developed since the 10-year NWFP review (Reeves 2006), with particular emphasis on the area of the NWFP. Among the key issues considered are:

- The ecological, physical, and biological importance of headwater and intermittent streams.
- The contribution of periodic disturbances to the resilience and productivity of aquatic ecosystems.
- The inherent variation of aquatic ecosystems in space and time.
- A better understanding of the variation in where key ecological processes occur within the stream network and the development of new tools to identify these locations.

- An understanding of the variation in the capacity of aquatic ecosystems to provide habitat for various fish species.
- Awareness of climate change and its potential effects.

We provide an update on the status of species listed under the Endangered Species Act (ESA) and the components and the associated monitoring program (Aquatic and Riparian Effectiveness Monitoring Program [AREMP]) of the ACS. We also assess the implications for the potential evolution of the ACS in the next round of forest plans. Reeves (2006) provided a thorough review of the literature in the first 10 years of the ACS, and readers should refer to that publication for a review of the relevant science during that time.

Guiding Questions

Federal land managers submitted many questions that they deemed necessary to consider in the NWFP science synthesis to help with any revisions of forest plans. Because there was substantial overlap among and duplication in the questions, we distilled them into categories represented by the following eight questions to guide our update and assessment:

1. Is the science foundation of the ACS still valid, or does science developed since 1993 suggest potential changes or adjustments that could be made to the ACS?
2. What is the basis of trends observed in the ACS monitoring program, and what are the limitations, uncertainties, and research needs related to monitoring?
3. What is known about the variation in characteristics of unmanaged streams and riparian ecosystems in relation to stream networks across the NWFP area?
4. What has been learned about the effects of riparian vegetation on stream habitat and environments?
5. What effects have human activities had on stream and riparian ecosystems?
6. What is the scientific basis for restoration management in riparian reserves, and how does restoration relate to the ecological goals of the ACS?
7. What is the capacity of federal lands in the NWFP area to contribute water for a suite of economic, recreational, and ecological uses?

8. What are the potential effects of climate change on aquatic ecosystems in the NWFP area, and are these adequately addressed by the ACS?

These eight questions are not answered specifically in sequence because of the overlap among them and the variety of topics they involve. They are, however, answered to the extent possible in different or multiple sections of the chapter, and are addressed in outline in the conclusions.

Key Findings

Status of Species and Population Units Listed Under the Endangered Species Act on Federal Lands in the Northwest Forest Plan Area

In 1993, only the Sacramento winter Chinook salmon (*Oncorhynchus tshawytscha*), and the shortnose sucker (*Chasmistes brevirostris*), and Lost River sucker (*Deltistes luxatus*, both native to the Klamath River system) were listed as threatened or endangered under the ESA in the area covered by the NWFP. Within a few years of the development of the ACS, 23 evolutionarily significant units of Pacific salmon and 3 distinct population segments of bull trout (*Salvelinus confluentus*) were listed under the ESA (table 7-1). There have been three additions since the 10-year review (Reeves 2006): the Oregon Coast coho salmon evolutionarily significant unit (*O. kisutch*), and two other fish species, the Oregon chub (*Oregonichthys crameri*) and the Pacific eulachon (*Thaleichthys pacificus*). No population units of Pacific salmon or bull trout have warranted delisting since the ACS was developed.² However, the Oregon chub was delisted in 2015 (USFWS 2015), becoming the first fish to be delisted because of increases in numbers. Habitat on the Willamette National Forest contributed to its recovery.

The developers of the ACS anticipated the ESA listing of distinct population segments of various species of Pacific salmon, evolutionarily significant units, and other fish species. The ACS was not expected to prevent the listing of any species or distinct population segment because factors outside the responsibility and control of federal

² http://www.westcoast.fisheries.noaa.gov/publications/status_reviews/salmon_steelhead/2016_status_review.html.

Table 7-1—Evolutionarily significant units (ESUs) of Pacific salmon and trout (*Oncorhynchus* spp.), distinct population segments (DPSs) of bull trout (*Salvelinus confluentus*), and fish and amphibian species listed under the Endangered Species Act that occur in the area covered by the Northwest Forest Plan

Species ^a	ESU/DPS/species	National forests (NFs) and Bureau of Land Management (BLM) districts in which ESU, DPS, or species occur
1. Fish		
Coho salmon	Lower Columbia/ southwest Washington	Gifford Pinchot and Mount Hood NFs
	Oregon coast	Siskiyou, Siuslaw, and Umpqua NFs; Coos Bay, Eugene, Roseburg, and Salem BLM districts
	Southern Oregon/northern California	Klamath, Mendocino, Rogue River–Siskiyou, Shasta-Trinity, and Six Rivers NFs; Arcata, Medford, and Redding BLM districts; Kings Range National Conservation Area (NCA)
	Central California coast	Ukiah BLM district
Chinook salmon	Puget Sound	Gifford Pinchot, Mount Baker–Snoqualmie, and Olympic NFs
	Lower Columbia	Gifford Pinchot and Mount Hood NFs; Salem BLM district
	Upper Columbia	Okanogan-Wenatchee NF
	Upper Willamette	Mount Hood and Willamette NFs; Eugene and Salem BLM districts
	California coastal	Mendocino and Six Rivers NFs; Arcata and Ukiah BLM districts; Kings Range NCA
	Sacramento River winter run	Mendocino and Shasta-Trinity NFs; Mendocino BLM district
	Central Valley spring run	Shasta-Trinity NF; Mendocino and Redding BLM districts
Chum salmon	Hood Canal summer	Olympic NF
	Columbia River	Salem BLM district
Steelhead	Puget Sound	Gifford Pinchot, Mount Baker–Snoqualmie, and Olympic NFs
	Lower Columbia	Gifford Pinchot and Mount Hood NFs; Salem BLM district
	Mid-Columbia	Gifford Pinchot, Mount Hood, and Wenatchee NFs
	Upper Columbia	Okanogan-Wenatchee NF
	Upper Willamette	Willamette NF; Eugene and Salem BLM districts
	Northern California	Mendocino and Six Rivers NFs; Arcata, Mendocino, and Ukiah BLM districts; Kings Range NCA
	Central California coast	Arcata BLM district; Kings Range NCA
Bull trout	Central Valley, California	Mendocino and Shasta-Trinity NFs Mendocino BLM
	Klamath River	Fremont-Winema NF
	Columbia River	Deschutes, Gifford Pinchot, Mount Hood, Okanogan-Wenatchee, and Willamette NFs; Eugene BLM district
	Coastal—Puget Sound	Mount Baker–Snoqualmie and Olympic NFs
Lost River sucker		Fremont-Winema NF
Shortnose sucker		Fremont-Winema NF
Pacific eulachon		Siuslaw and Six Rivers NFs

Table 7-1—Evolutionarily significant units (ESUs) of Pacific salmon and trout (*Oncorhynchus* spp.), distinct population segments (DPSs) of bull trout (*Salvelinus confluentus*), and fish and amphibian species listed under the Endangered Species Act that occur in the area covered by the Northwest Forest Plan (continued)

Species ^a	ESU/DPS/species	National forests (NFs) and Bureau of Land Management (BLM) districts in which ESU, DPS, or species occur
2. Amphibians		
Oregon spotted frog (T)		Deschutes, Fremont-Winema, Gifford Pinchot, Mount Hood, and Willamette NFs; Klamath Falls and Medford BLM districts; Columbia River Gorge National Scenic Area (NSA) (S)
Cascades frog (petitioned)		Deschutes, Gifford Pinchot, Mount Baker–Snoqualmie, Mount Hood, Okanagan-Wenatchee, Olympic, Rogue River–Siskiyou, Umpqua, and Willamette NFs; Medford (S), Roseburg, and Salem BLM districts
Oregon slender salamander (petitioned)		Mount Hood and Willamette NFs; Columbia River Gorge NSA
Cascade torrent salamander (petitioned)		Gifford Pinchot, Mount Hood, and Willamette NFs; Eugene and Salem BLM districts; Columbia River Gorge NSA
Columbia torrent salamander (petitioned)		Siuslaw NF; Salem (S) BLM district
Western pond turtle (petitioned)		Fremont Winema, Mount Hood, Rogue River–Siskiyou, Siuslaw, Umpqua, and Willamette NFs; Columbia River Gorge NSA; Coos Bay, Eugene, Klamath Falls, Medford, Roseburg, and Salem (S) BLM districts

^a Petitioned = under review for Endangered Species Act listing; T = threatened; S = suspected occurrence.

land managers contribute to the decline and recovery of fish populations and will strongly influence their recovery.

These factors include:

- Degradation and loss of freshwater and estuarine habitats on nonfederal lands (McConnaha et al. 2006, NRC 1996).
- Excessive harvest in commercial and recreational fisheries (NRC 1996).
- Migratory impediments, such as dams (McConnaha et al. 2006, NRC 1996).
- Loss of genetic integrity from the effects of hatchery practices and introductions, combined with undesirable interactions (e.g., competition and predation) involving hatchery and naturally produced fish (Araki and Schmid 2010, NRC 1996).

Thus, the ACS was an attempt to develop a strategy to guide management of aquatic ecosystems on federal lands in the NWFP area that would meet potential ESA requirements. The ACS was expected to make significant contributions to the recovery of the ESA-listed fish by increasing the quantity and quality of freshwater habitat for Pacific salmon and protecting and enhancing habitats of other species (FEMAT 1993). Although the condition of habitat in aquatic ecosystems on federal lands appears to have improved at least slightly over the NWFP area, this has not been sufficient to change the status of most listed fish.

The potential of federal lands to contribute to the recovery of listed fish, particularly Pacific salmon, in many parts of the NWFP area is likely more limited than was recognized when the ACS was developed. The primary reason for this difference




Figure 7-1—Proportion of the length of fish-bearing streams categorized as most ecologically sensitive in the six study watersheds of Reeves et al. (2016a) in western Oregon, by ownership. Ecologically sensitive areas are portions of the stream network that (1) have moderate- to high-quality habitat potential for coho and Chinook salmon and steelhead, (2) have moderate to high potential to warm if the riparian ecosystem is modified, and (3) have moderate to high potential for erosion or debris flows. From Reeves et al. 2016a.

is that, in many situations, federal lands (figs. 7-1 and 7-2) have a limited capacity to provide high-quality habitat for some of the listed fish. Federally managed lands are generally located in the middle to upper portions of watersheds, which tend to have steeper gradients and more confined valleys and floodplains, making them inherently less productive for some fish (Burnett et al. 2007, Lunetta et al. 1997, Reeves et al. 2016a). Federal lands may, however, be important sources of wood, sediment (Reeves et al. 2016a), and water (Brown and Froemke 2010, 2012) for downstream nonfederal lands, and will be important for the potential recovery of most populations. Nevertheless, their contribution to recovery may in many cases be insufficient without parallel contributions from nonfederal land ownerships elsewhere in the basin (Grantham et al. 2017).

The numbers of Pacific salmon and other anadromous fish returning to freshwater in the NWFP area are strongly influenced by ocean conditions, which are highly variable over time. Favorable conditions (cold water) tend to occur in the negative phase of the Pacific Decadal Oscillation (PDO) and the La Niña phase of the El Niño-Southern Oscillation (ENSO), when fish growth is strong and survival is high, resulting in strong returns of adults to freshwater (Mantua et al. 1997). Survival is low and numbers decline during warmer

periods, especially during the positive phase of the PDO and the El Niño phase of the ENSO. Winters are cold and wet in the negative PDO–La Niña phase, which also creates more favorable conditions in freshwater (Mantua et al. 1997). A positive PDO–El Niño produces dry, warm winters, reducing streamflows, increasing water temperatures, and increasing the occurrence of fire (see chapter 3). The last extended period of high productivity was from the late 1940s to 1976 (Mantua et al. 1997), with brief periods of favorable conditions in 1984–1988, 1999–2002,³ and 2010–2011 (Bond et al. 2015). However, beginning in 2013, abnormally warm conditions in the Pacific Ocean (“the Blob”) developed because of lower-than-normal heat loss from the ocean to the atmosphere, combined with a relatively weak mixing of the upper ocean layer owing to an unusually high atmospheric pressure (Bond et al. 2015). Initial effects were most notable in the North Pacific Ocean off Alaska. Ocean conditions changed noticeably along the NWFP area in 2014 as a result, and fish returns are expected to decline over the next few years.

³ Mantua, N. 2017. Personal communication. Leader Landscape Ecology Team, National Marine Fisheries Service–Southwest Fisheries Science Center, 8901 LaJolla Shores Drive, Santa Cruz, CA 92037. nate.mantua@noaa.gov.




Figure 7-2—Distribution of ecologically sensitive stream reaches on federal and nonfederal lands in the McKenzie River watershed, Oregon. From Reeves et al. 2016a.

We are unable to separate the influence of ocean conditions over the last 10 years from the influence of changes in the condition of freshwater ecosystems on federal lands that may have occurred under the NWFP and ACS. The actual contribution of freshwater habitats to the persistence and recovery of anadromous salmon and trout will be relatively more important when ocean conditions move into a less-productive phase (Lawson 1993). Improvements in the quantity and quality of freshwater habitat resulting from the ACS could result in relatively greater numbers of fish entering the ocean, thus increasing the likelihood of persistence of many populations during periods of low ocean productivity. However, as noted previously, the contribution of federal lands may be more limited than expected because their potential to provide high-quality habitat is less than originally recognized when the ACS was developed.

The status of other aquatic-riparian species in the NWFP area is not as well monitored as that of Pacific salmon. The Oregon spotted frog (*Rana pretiosa*) was listed as threatened under the ESA in 2014. It is a pond-breeding amphibian now restricted to isolated populations that overlap the NWFP area in western Washington and Oregon.⁴ Five other aquatic-riparian amphibian and reptile species are petitioned for ESA listing and are under status review: (1) Columbia torrent salamander, *Rhyacotriton kezeri*; (2) Cascade torrent salamander, *R. cascadae*; (3) Cascades frog, *Rana cascadae*; (4) Oregon slender salamander, *Batrachoseps wrighti*; and (5) western pond turtle, *Actinemys marmorata*. The two torrent salamanders are headwater forest species, occurring predominantly in and along the banks of small streams, with significant portions of their ranges on nonfederal lands. Nevertheless, federal riparian reserves contribute habitat for localized populations of Columbia torrent salamanders and more extensive areas for Cascade torrent salamanders. The Oregon slender salamander is found in proximity to down wood on the forest floor in riparian and upland forests, and has associations with older forest conditions (Clayton and Olson 2007). Cascades frogs are pond breeders at higher elevations in the Cascade Range, where they may be affected by multiple stressors (Pope et al.

2014). Similarly, multiple threats appear to affect western pond turtles, which may occur in stream and pond systems in the NWFP area (Rosenberg et al. 2009).

Monitoring—Aquatic and Riparian Effectiveness Monitoring Program

Watershed conditions—

The Aquatic and Riparian Effectiveness Monitoring Program is responsible for monitoring, assessing, and reporting on watershed conditions on lands governed by the NWFP. Although NWFP implementation began in 1994, AREMP implementation was delayed to accommodate the time needed for its design. The scope of the AREMP sampling design includes field-data collection across 250 watersheds in the Plan area, with a rotation of sampling among watersheds conducted each year so that the entire population of watersheds selected for monitoring would be sampled over an 8-year period. In addition, using geographic information system (GIS) and remotely sensed data are used to quantify roads and vegetation in 1,974 watersheds with federal lands in the Plan area and assess the condition of upslope and riparian areas.

Pilot monitoring of watershed conditions began in 2000, and the monitoring plan was finalized in 2003 (Reeves et al. 2004). The first full rotation of watershed visits was conducted in years 2002–2009, assessing initial status, and the second full rotation is scheduled to occur in 2010–2018 where paired assessments of most watersheds were possible owing to watersheds being resampled a second time. Reporting is on a 5-year cycle, in synchrony with NWFP establishment, with the first report covering up to year 10 of Plan implementation (Gallo et al. 2005), the second report covering up to year 15 (Lanigan et al. 2012), and the third to year 20 (Miller et al. 2017). The 20-year report includes assessment of data from the first rotation of watershed visits (2002–2009) and the first 4 years of the second rotation (2010–2013), and hence includes trend assessments based on a subset of sampled watersheds.

Changes in data collection and aggregation procedures, and in application of analytical methods, were anticipated from the onset of the development of AREMP (Hohler et al. 2001). In the late 1990s, our understanding of watershed

⁴ <http://www.fs.fed.us/r6/sfpnw/issssp/agency-policy/>.

ecology and watershed-condition assessment approaches was acknowledged to be limited, and advances in both ecological and statistical disciplines were expected to contribute to further development of AREMP assessments. Indeed, both data sources and analyses have changed over time, with the consequence being that the results from each of the reports are not directly comparable. For example, relative to data sources, some data-collection procedures changed as attribute variability became apparent. Relative to analytical approaches, the 10- and 15-year analyses used decision support models (Reeves et al. 2004, Reynolds et al. 2014) that depended on empirical relations and expert judgment to evaluate data. The 20-year report employed a more statistical focus, with expert opinion and independent analysis of upland, riparian, and in-channel metrics. Additional discussion of adaptive processes through AREMP implementation, including anticipated next steps for research, is presented following the key 20-year findings. Although data analysis and assessment methods changed, each report reanalyzes the entire spatial and temporal dataset available at the time, and is intended to represent the most current understanding of status and trends since the beginning of the Plan.

Key 20-year findings—

The 20-year AREMP report (Miller et al. 2017) examined upslope-riparian and in-channel datasets separately. This segregation acknowledged that the source data differed significantly between these two components. Upslope-riparian data were derived from remote sensing and GIS landscape data covering all NWFP watersheds (watersheds containing more than 5 percent federal ownership, a total of 1,974 watersheds). In contrast, in-channel data were derived from annual field measurements, and therefore were limited to 213 sampled watersheds. Upslope-riparian assessments integrated five data types reflecting watershed processes: sedimentation, wood recruitment, riparian condition and processes, hydrology, and fish passage. In-channel analyses focused on three additional data types, assessed independently: physical-habitat condition, macroinvertebrate assemblages, and water temperature.

Upslope-riparian analyses integrated finer scaled data metrics reflecting indicators of key watershed processes. These processes included (1) stream-sediment delivery

from landslides, based on road and vegetation disturbances, in addition to geology and climate attributes; (2) down-wood production and delivery, based on riparian and upland vegetation metrics; (3) riparian condition and associated processes as represented by stream temperature, streambank stability, and species-habitat provision based on riparian vegetation condition and riparian road density; (4) hydrology, focusing on peak flow, based on road and vegetation metrics; and (5) fish passage, based on stream gradient and assessment of barriers (e.g., dams, some road crossings). Using a multicriteria assessment approach, akin to analyses conducted in previous reports, attributes for a watershed were scored to a common 0 to 100 scale, reflecting an index of most-to-least deviation from least human-disturbed conditions.

The 20-year report found little change in the average upslope-riparian condition, from a score of 68 in 1993 to 69 in 2012. However, noticeable shifts were observed in the overall score distribution (fig. 7-3A). In particular, there was a noticeable increase in scores from the low to mid range (15 to 50) to a higher range (60 to 90), whereas the area with the highest scores (>90) decreased slightly. These patterns reflected a signature of federal land use allocations. The mean score in the most protected category of land use allocation (Congressionally reserved lands) decreased (−1), indicating greater disturbance, whereas averages for late-successional reserves and matrix lands increased (+2, +3), indicating less disturbance. The upward shift in the low-range scores is likely attributed to widespread vegetation regrowth and targeted road decommissioning in previously harvested watersheds. In contrast, the decrease in the high-end scores mainly followed the pattern of large fires during the assessment period, many of which occurred in wilderness areas, including the Biscuit Fire in southwest Oregon (2002), the B&B Complex fires in the central Oregon Cascade Range (2003), and numerous fires along the eastern edge of the North Cascade Range in Washington (2006).

It may seem counterintuitive that the most protected lands would show a trend toward more disturbance. Although this trend might be seen as negative because fire results in a loss of vegetation and an increase in riparian-upland disturbance, it is simplistic to consider this an




Figure 7-3—Results of the 20-year assessment of watershed conditions by the Aquatic and Riparian Effectiveness Monitoring Program (Miller et al. 2017): (A) upslope-riparian condition, (B) in-channel condition, (C) aquatic macroinvertebrates, and (D) 7-day running average of maximum water temperatures. Source: Miller et al. 2017.

adverse effect at the regional scale of forest ecosystems and their embedded watersheds. Wildfires are an integral component of long-term forest and stream ecosystem dynamics (e.g., Bisson et al. 2003, Franklin et al. 2017, Reeves et al. 1995), with direct benefits to stream habitats and biota resulting from fire (e.g., Flitcroft et al. 2016a) and related natural disturbances such as landslides (wood and sediment delivery to streams) (e.g., Benda and Dunne 1997a, 1997b; May and Gresswell 2003; Reeves et al. 2003). Aquatic-riparian ecosystems are dynamic, being multistate in space and time (Olson et al. 2017a, Penaluna et al. 2016). This recent AREMP finding highlights our nascent understanding of the range of historical aquatic-riparian conditions in the NWFP area and the cadence and extent of natural disturbance events. It also brings to the forefront the role of both passive and active management of these aquatic-riparian systems in the future to maintain and restore the dynamics of aquatic-riparian ecosystems in the region, and the importance of considering whether we need to manage for resilience. In this framework, shaped by land use allocations and trends detected therein, it is possible that development of more nuanced evaluation methods is needed to assess variation in aquatic-riparian conditions. The problem also becomes conceptually challenging, given a known shifted baseline from past anthropogenic disturbances, including the effects of fire suppression, as well as from the need to develop projections of climate change, climate extremes, and related disturbances from fire and landslides.

In-channel watershed-condition assessments conducted for the 20-year report were the first such assessments to have enough field-site revisits (about half the sample) to estimate trends. Instream conditions in subsampled watersheds in the Plan area were assessed by evaluating three separate elements: physical habitat, macroinvertebrates, and water temperature. First, for physical habitat, a composite index score (on a 0 to 100 scale, relative to unmanaged reference conditions; see below for more discussion on this topic) based on expert judgment was derived from substrate (percentage of fine substrates at a stream-reach scale), pool-tail fine substrates, and the frequency of medium- and large-size down wood.

A small but statistically significant increasing trend (from 46 to 49) in overall physical habitat condition was detected when measured both on a yearly basis with all data and from the subset of watersheds that had been revisited (fig. 7-3B). Individual components of the physical habitat varied: reach-scale fine substrate showed an increasing trend in occurrence, whereas instream wood and pool-tail fine substrates showed no significant changes.

Second, an index was also derived for macroinvertebrates (fig. 7-3C), which were assessed at the site level, then grouped into taxonomic classes and compiled into a score representing the ratio of expected species based on reference sites ($E = 1$) (see below) to occurrence of no expected species ($O = 0$). Site-scale scores were aggregated into watershed scores. A positive change in the mean score was detected for macroinvertebrate diversity, suggesting a shift toward a species composition reflective of expected reference conditions.

Third, water temperature was evaluated using an integrated model that assessed 7-day maximum temperatures collected at low points in watersheds from June to September. Water temperature showed a decreasing trend, although temperature averages were still higher than federal and state standards for salmonid habitat (fig. 7-3D). Interestingly, water temperatures in all land use categories decreased, but temperatures in the most protected category showed the smallest decline, perhaps reflecting the upslope findings, which showed a signature of disturbance owing to vegetation loss in reserves.

Reflection on adaptive processes through AREMP implementation—

Here, we outline primary changes and challenges in AREMP monitoring over the past 20 years, many of which are ongoing research-emphasis areas. These topics are common, foundational aspects of many aquatic-riparian monitoring programs, representing a larger coalition of scientists and managers addressing similar themes over diverse landscapes. The science of watershed-scale ecology has followed the development of the discipline of landscape ecology, and the challenges cited here are representative of a parallel course of design and analytical adaptive processes occurring in terrestrial forest ecosystems.

Overall, NWFP monitoring, including the AREMP (Hohler et al. 2001), was framed as an adaptive-management cycle (Mulder et al. 1999). For AREMP, four primary changes and challenges over the course of the first 20 years of the NWFP have included (1) redefining the overarching objectives of watershed-condition monitoring, shifting from a salmonid habitat focus to one that was more representative of the environmental conditions of entire watersheds, while retaining selected key elements of salmonid habitats; (2) refining the indicators used in data collection and analysis; (3) reconsidering approaches to using benchmarks or reference conditions for assessment; and (4) modifying data integration methods. These four topics are discussed further below, with comparisons to other aquatic-monitoring programs for a broader perspective.

Defining objectives—

The first step in designing the NWFP monitoring modules was to define the goals and objectives of monitoring (Mulder et al. 1999). The NWFP defined the central question for aquatic ecosystems as, “Is the ecological health of the aquatic ecosystems recovering or sufficiently maintained to support stable and well-distributed populations of fish species and stocks?” (USDA and USDI 1994a, E-7). The primary fish taxa with status of concern in the NWFP area were native salmonids, hence a taxonomic focus was present from the origin of the Plan. Although particular species (owls, murrelets, salmonids) have been a principle interest of NWFP monitoring programs, concepts developed in the monitoring plan also stated that “Because of the current wide (and justified) interest in all components of biological diversity, however, the species-centric approach is no longer sufficient” (Mulder 1999, p. 29). In this vein, the language of the ACS objectives (USDA and USDI 1994a) had also included aquatic-riparian habitat conditions and species, but with a focus on multiple processes that were known to be tied to development of salmonid habitat conditions. At the time of the NWFP, emerging science on the role of disturbance in renewal of aquatic habitats also suggested a change in focus from the assessment of narrowly specified, in-channel habitat elements (e.g., a certain number of pieces of large wood per stream length) toward the ecosystem processes that form and maintain habitats

(USDA and USDI 1994a). AREMP was perhaps the most ambitious of the monitoring modules in this regard, calling for monitoring a broad set of conditions in the upslope, riparian, and in-channel portions of watersheds that related to ecological processes tied to fish habitat, and evaluating these in comparison to broad distributions of conditions rather than solely on a watershed-by-watershed basis (Reeves et al. 2004).

The ACS was originally envisioned as a “coarse-filter” conservation effort (Hunter 2005, Noss 1987) (see additional discussion in chapter 12). The focus of the ACS was on restoring and maintaining ecological processes that created and maintained aquatic ecosystems for a suite of organisms, primarily ESA-listed fish, and for clean water and other ecosystem services (USDA and USDI 1994a). AREMP was, therefore, initially directed at the habitat of native salmonids, a primary responsibility of federal land managers and regulators in the NWFP area (Reeves et al. 2004). Habitat conditions for native salmonid fishes were initially used as metrics for watershed condition trend assessment, owing to their sensitivity to changes in several habitat features (e.g., water temperature, sediment, down wood). As with other coarse-filter assessments that use biotic indicators such as umbrella or flagship species (e.g., Raphael and Molina 2007), it was assumed that other aquatic and riparian-dependent organisms would benefit if watershed conditions for salmonids improved. Hence, salmonids were recognized as a focal species group, assuming that if their habitats were sustained or improved in condition over time, it would infer sustainability or improvement of the greater community of biota and the broader watershed-scale ecosystem functions and processes upon which they rely.

Development of aquatic monitoring programs requires a clear articulation of which biota and associated functional characteristics of habitats and ecosystems are being considered, and how they are likely to be altered as a result of the actions of interest (Carlisle et al. 2008, Palmer et al. 2005, Pont et al. 2006). Such species-based approaches may not fully account for the variation in species abundance or community composition present, given the spatiotemporal heterogeneity in ecosystem conditions generated by natural disturbances. Further, this natural variability in species and

environmental conditions may make it difficult to identify the effects of anthropogenic disturbances and recovery from those disturbances, and thus make it difficult to assess and understand the ecological consequences of detected changes (Frissell et al. 2001). However, explicitly identifying the organism(s) of interest is essential for understanding what the monitoring results mean for those species and the fauna the species represents (Wohl 2016).

The 20-year AREMP analysis shifted from the emphasis of the 10- and 15-year analyses on evaluating habitat for salmonids to characterizing more general environmental conditions. Miller et al. (2017) used the 5th and 95th percentile values from a suite of physical attributes in reference sites in systems with the least human-caused disturbance (see later discussion) to determine the favorability of conditions for biota in monitored watersheds. Based on expert judgment, the 5th percentile was considered the most favorable for some attributes (e.g., pool-tail fines, reach fines) and the 95th for others (wood). However, Miller et al. (2017) did not identify which organisms these conditions were supposed to favor, making it difficult to understand the ecological validity of these values and the consequences of any changes detected.

This switch highlights a continuing scientific debate in the monitoring and assessment community on the merits of focusing assessments on particular flagship or umbrella taxa rather than on more general environmental processes and conditions. On the one hand, umbrella species serve as meaningful “endpoints” (Suter 2001) or “final ecosystem services” (Blahna et al. 2017) that provide relevance to monitoring results. On the other hand, in aquatic-riparian ecosystems, salmonid distributions do not reach headwater streams, which make up most of the stream length in NWFP watersheds (Gomi et al. 2002). Salmonid habitat in larger streams may not be representative of the condition of the entire watershed, unless solid ties to upstream and upslope conditions can be made. Further, the adequacy of salmonids as umbrella species has not been formally assessed (Murphy et al. 2010, Simberloff 1998), and there are questions about whether one species can be an indicator of the condition of other species (Carlisle et al. 2008). Although these two

objectives for watershed-condition assessments (salmonid habitat versus watershed environmental conditions) are closely related, differences in emphasis have a ripple effect that plays out through the monitoring and assessment process, and affects how results might be interpreted relative to goals of maintaining and restoring conditions.

The original AREMP design document laid out a conceptual model that considered the interactions between upslope, riparian, and in-channel processes, all within a variable landscape (e.g., precipitation, geology) (Reeves et al. 2004). No formal assessment of relationships between upslope/riparian/in-channel indicators in the AREMP conceptual model has been published, but a number of studies are relevant to pieces of this framework. Burnett et al. (2007) developed a relative ranking, Intrinsic Potential (IP) ranging between 0 (poor) and 1 (excellent) to determine the geomorphic potential of a reach to provide habitat for coho salmon in larger streams. This work was followed by data-driven watershed-scale models of several habitat attributes important for salmon that tied upland-riparian conditions to instream habitats, including models of debris-flow-prone areas (delivering sediment to streams: Benda and Dunne 1997a, 1997b; Burnett and Miller 2007); wood recruitment (e.g., Reeves et al. 2004); and thermal loading (as a proxy to represent stream temperatures; see Reeves et al. 2016a). Several of these studies also contributed to a better understanding of instream processes connecting lower order headwater streams to higher order streams occupied by salmonids. Syntheses of these multifaceted watershed-process models have contributed to a better understanding of the multistate nature of aquatic-riparian ecosystems (Olson et al. 2017a, Penaluna et al. 2016, Reeves and Spies 2017), and integration of several of these watershed-integration models have been used to evaluate potential management options (Reeves et al. 2016a). Full incorporation of these concepts into watershed-condition assessments has been indirect to date, for example, by implicitly supporting approaches to assess upland-riparian areas and full in-channel networks from headwaters downstream. This topic deserves continued attention as AREMP procedures continue to develop, and can potentially inform the overarching objectives of the program.

Other aquatic-monitoring programs have incorporated upslope/riparian/in-channel relationships into their conceptual models, albeit in quite different ways. The National Rivers and Stream Survey (NRSA) (USEPA 2016) related four chemical and four physical habitat stressors to multi-metric macroinvertebrate and fish indices using a concept of relative risk: the likelihood of finding poor biological conditions in a river or stream when stressor concentrations are high relative to the likelihood when they are low. Indirectly, these stressors, as well as three land-use metrics (urban land cover, agricultural row-crop land cover, and dam influence), were used in selecting the reference sites used to evaluate the stressor and response metrics.

PACFISH/INFISH Biological Opinion Monitoring Program (PIBO) approaches have not directly assessed upslope indicators, but have selected a set of physical-habitat indicators based on sensitivity to land-use management intensity, using road density as a surrogate (Al-Chokhachy et al. 2010). A recent assessment of the PIBO program conceptual model by Irvine et al. (2015) examined correlations between upslope (grazing, road density, percentage forested), in-channel habitat (fine sediment, temperature), and a macroinvertebrate observed/expected index. Although they found weak to no support for causal pathways related to effects of anthropogenic drivers on biological condition, they surmised that the conceptual model was sound, and the weak correlations were due to imprecision in the measurement of drivers (grazing, roads); stressors (sediment, measured in pool tails); and responses (macroinvertebrates, measured in riffles). They cited the more general issue that regional trend monitoring is not optimized for detecting causal mechanisms. A related and broader concern is that such surveys may underestimate infrequent but high-severity events (Suter 2001). In contrast, it was notable in the AREMP 20-year report that the signatures of wildfire and road decommissioning, relatively low-frequency events, were detectable in the upslope-riparian assessment because it included a full census of watersheds, rather than a limited sample. Overall, scientific work in the past 20 years has continued to support the dynamic, disturbance-based ecology of aquatic-riparian ecosystems (see “Natural

Variability” section later in this chapter). Although the AREMP conceptual model has not changed, there have been numerous refinements to the indicators used, as well as to their combination and interpretation.

Selection of indicators—

In-channel biotic indicators—In-channel biotic metrics have proven to be particularly challenging to monitor. Although fish populations are of principal concern to managers because of regulatory requirements and their potential as umbrella or flagship species, fish-data collection was dropped from AREMP protocol in 2007 because most in-channel watershed sample sites were above salmonid habitat, and the collection of meaningful salmonid-habitat data would have required a separate and intensive effort. Similarly, streambank amphibians were dropped because of detectability issues. Variability in detection spatially within watersheds as well as temporally within the year made streambank amphibian monitoring challenging: species that were present at a site early in a season could be missed if the site is not sampled until later in the season. Further, terrestrial salamanders are fossorial (live largely subsurface) and may not have been detected even if present (e.g., Hyde and Simons 2001).

Macroinvertebrates are the only remaining in-channel biotic indicator collected by AREMP. Macroinvertebrates have come to play a central role in many aquatic-monitoring programs because of their presumed responsiveness to local environmental conditions and ease of collection (PIBO [Al-Chokhachy et al. 2010], Oregon Department of Environmental Quality [Hubler 2009], NRSA [USEPA 2016]). This commonality creates the potential for data sharing between monitoring programs, assuming sufficient similarity in the sampling methods used. However, differing results for the fish and macroinvertebrate indices in a recent national assessment emphasized the danger of relying on one taxonomic group to represent the potential responses or condition of other groups (USEPA 2016). Understanding how to reliably collect and incorporate data from taxa other than macroinvertebrates is a challenge for ongoing research; new multispecies environmental DNA methods are promising and are undergoing trial now (see app. 1).

In-channel abiotic indicators—Measurement reliability has also been a challenge with abiotic in-channel indicators. Based on quality-control sampling, AREMP dropped the evaluation of pool frequency, depth, and median particle size from the latest 20-year assessment. Measurements of these parameters are still being collected, owing to the perceived importance of these indicators for assessment of habitat conditions, and the AREMP program is actively investigating ways to make current collections of these data more reliable for use in future analyses. Remaining abiotic indicators used in the latest report were pool-tail fine substrates, reach-scale fine substrates, down wood, and water temperature (Miller et al. 2017).

Upslope-riparian indicators—The basic indicators used in the upslope-riparian portion of the AREMP assessments have changed little, but their evaluation has become more context sensitive. In the latest report, their combination was reorganized into an index with a more process-based structure. The new process-based model structure includes five processes that aligned it more directly with the conceptual model in the original AREMP plan: sediment and wood delivery, riparian shading, hydrology, and habitat connectivity. All these indicators are based primarily on road and vegetation data, the two major land-use metrics that can be traced backward in time to assess trends since the beginning of the NWFP. This estimation of historical data is a challenge that appears not to have been attempted by other broad-scale programs (Gordon 2014). Road and vegetation-management effects on aquatic systems continue to be active areas of research.

Of particular relevance to the AREMP assessment is work identifying the differential effects of sediment delivery from roads, based on landscape position (Al-Chokhachy et al. 2016, Black et al. 2012) and the incorporation of potential fish-habitat considerations into the measure of aquatic connectivity (Chelgren and Dunham 2015). Advances in satellite data and their classification have considerably expanded the available vegetation metrics and the ability to track these through yearly time steps (Kennedy et al. 2010, Ohmann et al. 2011). These capabilities should be examined in light of the disturbance-ecology paradigm discussed throughout this chapter.

Examining the freshwater assessment literature more generally, Kuehne et al. (2017) found a shift in measures used from field-based responses to landscape-stressor metrics. The use of upslope indicators is attractive to managers because this is where the most extensive management activities currently occur (vegetation, roads), and also because these measures are more easily collected via remote sensing and existing GIS data, and do not require more labor-intensive field surveys. Measuring and understanding both upslope and in-channel processes, and the relationships between them, is critical. Taking into account the difficulties encountered by other parallel aquatic-monitoring programs, more formal testing of the AREMP conceptual model is warranted.

Benchmarks for assessment—

In the data-assessment step of analysis, indicators are typically compared against some benchmark (alternatively referred to as standards, thresholds, or evaluation criteria) to come up with a measure of watershed or aquatic-habitat condition. This has proven to be one of the greatest challenges, particularly given the expanding recognition of the reliance of aquatic habitats on dynamic processes of disturbance and renewal. Such monitoring faces the fundamental challenge of using static measurements (with limited temporal frequency) to measure dynamic processes.

Stoddard et al. (2006) described a number of common approaches to choosing benchmarks: reference conditions, best professional judgment, interpreting historical condition, extrapolating from empirical models, and ambient distributions. The first two AREMP reports (10 and 15 year) relied on empirical models and expert judgment to set evaluation criteria (Gordon and Gallo 2011). To accommodate environmental heterogeneity, separate thresholds were solicited for seven aquatic provinces identified in the NWFP area. To accommodate environmental heterogeneity, separate thresholds were solicited for seven aquatic provinces identified in the NWFP area. In most provinces, experts were unwilling to commit to standards in higher gradient streams for some attributes (floodplain connectivity, pool frequency, pool-tail fines, and median substrate diameter), so they were not included in the evaluation of these sites.

The most recent (20-year) AREMP report switched to using reference conditions to set criteria for in-channel conditions and upslope vegetation. Reference criteria were chosen using a nearest-neighbor approach, which matched a site to the five to seven nearest reference sites, based not on geographic distance but rather on similarity of largely invariant site characteristics (e.g., gradient, geology) (Bates Prins and Smith 2007). This approach was more empirically based than previous assessments; it set standards for higher gradient stream reaches and established a consistent method across the whole NWFP area. The use of the reference-condition approach in a monitoring program has important associated assumptions. The selection of reference sites should match the distribution of states in the ecosystem, reflecting the spatial and temporal dynamics at play. These values or thresholds are often either a direct judgment call or a chosen percentile of the overall disturbance distribution. In more highly disturbed ecoregions, few sites may qualify, as was the case with the Oregon/Washington Coast Range and the Franciscan provinces in the latest AREMP assessment (Miller et al. 2016). The nearest-neighbor approach did not rely on provinces/ecoregions, but rather incorporated environmental variability more directly through site characteristics.

Another key consideration in development of reference distributions is including the entire natural range of conditions that an ecosystem can experience (Lisle et al. 2007, NRC 2000, Stoddard et al. 2006). Relative to the ACS and development of AREMP as originally conceived (Reeves et al. 2004), it was generally assumed that there is a given condition or limited set of conditions that supports aquatic organisms, primarily fish and macroinvertebrates (e.g., Karr and Chu 1998). The panel of scientists and managers who initially framed the ACS assumed that favorable conditions for fish were constrained to areas with cold water and structural heterogeneity provided by physical habitat components such as large down wood and coarse substrates—conditions often associated with old-growth forests—thus, these conditions and the associated old-growth forested riparian habitats were assumed to be most suitable for fish.

Recent studies, however, have demonstrated that native salmonids (Howell 2006, Rieman and Isaak 2010, Sestrich et al. 2011) and aquatic invertebrates (Minshall et

al. 1989) are capable of adapting to and being productive in a wide range of conditions, including those following major disturbances such as wildfire that affect stream conditions. Flitcroft et al. (2016a) found that although conditions for one life-history stage of salmonids may be unfavorable, other life-history stages may find the same conditions suitable, and populations may respond positively. Native salmonids may also change life-history tactics, such as by reducing age or size at maturity (Rosenberger et al. 2015). Evolving in naturally dynamic landscapes with infrequent to frequent fire (see chapter 3) and occasional landslides, these species appear to be resilient to a broad range of disturbances and environments that occurred under the natural range of variability. It is important for monitoring programs to incorporate this new perspective into the development of benchmarks and interpretation of results to better reflect the responses of aquatic organisms to both management and natural disturbances. The range of natural conditions likely spans recently disturbed sites as well as areas that have been undisturbed for hundreds of years or longer.

However, understanding the natural range of variability for an ecosystem is often difficult, owing to the extent and magnitude of anthropogenic effects (Miller et al. 2016, NRC 2000, Steel et al. 2016, Stoddard et al. 2006). This may especially be the case in dry-forest regions in the NWFP area where fire exclusion has altered forest and riparian plant composition and structure (see chapter 3); in areas where invasive species are now a dominant component of communities (app. 1); or where the signature of past human activities (Steel et al. 2016) and pervasive “press” disturbances (Yount and Neimi 1990) such as timber harvest have influenced the entire landscape so that current conditions, which may be a departure from the historical range, may now be seen as the norm (Pinnegar and Engelhard 2008), though they may have been rare or unknown in the past. Even areas that may appear to lack any sign of current or historical land use—areas with no discernable sign of recent human-caused disturbances—may no longer be considered pristine (see chapter 12). The Pacific Northwest moist coniferous forest region has recently been described as a “human-forest ecosystem,” because people are now a foundational element of the system (Olson and Van Horne 2017).

Because pristine areas may no longer exist in many, if not most, ecosystems, the reference-condition approach sometimes has been modified to use “least-disturbed conditions” as a reference (Stoddard et al. 2006). But depending on how this approach is applied, it may not include the full range of potential ecosystem conditions, especially in naturally dynamic landscapes as described above, where disturbance had been excluded. Worse, in today’s human-influenced forest landscape, there may be no locations that fit even the “least-disturbed” condition (see discussion in chapter 3).

Excluding the natural range of variability in the reference population influences the assessment of current conditions (NRC 2000). This issue can be illustrated by using the down wood data from Reeves et al. (1995), who examined three watersheds in the Oregon Coast Range that differed in the lengths of time since the last large wildfire (see details in “Attribute Integration Approaches” below). If just the values from the watersheds that were at an intermediate time point and the longest time point from disturbance were included as being in the population of reference conditions, wood values would vary from 12 to 24 pieces of wood per 100 m of stream. Twelve wood pieces per 328 ft (100 m) might be considered an extremely low value, and 24 a high value. However, if the most recently disturbed system from this dataset were included in the pool of reference conditions, the lower bound would be near 6 pieces of wood/328 ft, and the score for 12 pieces of wood/328 ft would be much higher, relatively speaking. Clearly, use of reference conditions to assess monitoring trends and their ecological relevance can be problematic if a wide range of variation is not included. Articulating how reference conditions were determined and the range of conditions they represent is essential to understanding the context for comparison, whether least-disturbed conditions, old-growth conditions, or professional judgment are used.

The relatively small number of matched sites in the reference pool used for the 20-year report is a potential drawback. Further, only one site-matching metric, quadratic mean diameter of conifers, reflected forest-ecosystem characteristics; this metric is only a limited surrogate for seral stage and does not reflect forest type, both of which

may influence stream characteristics. Concerning the reference-site approach more generally, there is also some question as to whether reference sites relatively free of human disturbance still exist, given widespread fire suppression and now climate change (see chapter 3) (Herlihy et al. 2008, Stoddard et al. 2006). To the extent that these concerns are true, it likely creates uncertainty concerning monitoring results that needs to be explored.

Other recent large-scale assessments have also used a variation of the reference-condition approach, but none have incorporated much detail on surrounding vegetation conditions. The NRSA (USEPA 2016) incorporated environmental variability more generally into their reference sites by selecting a different set of sites for each ecoregion. Thus, their reference distribution includes a larger number of sites than in the AREMP analysis, but the NRSA authors recognized that this approach might not account for fine-scale variability within ecoregions and made direct comparison of results between ecoregions problematic (Herlihy et al. 2008, USEPA 2016). They incorporated some finer scale measures of environmental variability (e.g., elevation) by including them as covariates in the multiple linear regression (MLR) equations that defined reference expectations for each indicator. The PIBO analysis incorporated environmental variability by using the MLR approach, but the only covariate related to forest condition was the percentage of the 295-ft (90-m) stream buffer in forested condition (Al-Chokhachy et al. 2010).

The final step in applying the reference-condition approach involves choosing evaluation thresholds from the distribution of reference values and comparing site values to these thresholds. AREMP and PIBO selected the 5th/95th percentiles of their reference distributions to define normalized scores, assuming that more extreme values might be outliers that could skew the scoring process. They then reported these values directly, so that scores approximated the percentile in the reference distribution. In contrast, NRSA chose to place all results into three classes (good/fair/poor) based on the less than 5th/5th–25th/greater than 25th percentiles of reference. The reference-condition approach may appear to be more empirical, but it still relies on professional judgment to set evaluation thresholds.

To promote success in ACS implementation, it is important to know how gains in watershed condition are measured. Because the natural range of variability occurring in a system over large spatial and long temporal scales occurs across a multidimensional continuum, it can be difficult, if not impossible, to incorporate into assessments (see chapter 3). One potentially useful approach is the use of “state and transition” models (e.g., Wondzell et al. 2007, 2012). Although such models can be difficult to validate, they can still be useful. It can be helpful to view aquatic ecosystems as multistate systems resulting from a variety of natural disturbances, as well as exogenous or anthropogenic processes that can alter habitat conditions, biota, and ecological processes (Penaluna et al. 2016). Having the full range of potential variability classified into discrete states provides a way to begin enumerating the ways in which variation is arrayed over large spatial scales and how it changes over long temporal scales. We suggest further exploration of reference conditions and their potential utility for analytical approaches, including consideration of how to use them in concert with state-transition models such as those developed by Wondzell et al. (2007, 2012).

The concept of the reference condition remains important in land management—not because it is a goal of management agencies to restore systems to some previous reference condition, but rather because knowledge of the historical range of variability can help inform choices about desired future conditions and, thereby, help determine management and restoration goals (chapters 3 and 12). Theoretically, departure from the reference condition would provide a relative measure to evaluate watershed conditions for managers who seek to maintain or restore ecosystem and species diversity (Nonaka et al. 2007, Safford et al. 2012). The multistate conceptual approach (Penaluna et al. 2016, Reeves et al. 1995) clearly shows that “reference conditions” are, in fact, a distribution of conditions from watersheds in various ecological states, similar to successional states in terrestrial systems (see discussion later in this chapter). As such, “departure from the reference condition” is no longer just a watershed-scale

question, but rather a regional-scale problem that considers the distribution of conditions across multiple watersheds.

Equilibrium versus spatially and temporally dynamic ecosystem concepts and the choice of benchmark reference conditions will become even greater challenges in future assessments given the increasing influence of climate change. There is growing concern about the extent to which ecosystems in the NWFP area, and elsewhere, have been affected by climate change and altered disturbance regimes, such as fire exclusion (Hessburg et al. 2005, Luce et al. 2012; also see chapter 3). We are likely seeing, or will soon see, the development of ecosystems that are different from the present and at least the near past (Hobbs et al. 2009, Luce et al. 2012) (fig. 7-4). Some have called this a new geological epoch—the “Anthropocene” (see chapter 12). The conditions




Figure 7-4—Conceptual roles for disturbance in a changing climate: (A) Disturbance could continue to operate much as it always has, with unique disturbance/recovery patterns, or (B) it could become the catalyst that forces ecosystems to shift rapidly and via alternate and uncertain pathways in response to climate. Source: Luce et al. 2012.

that result from these altered ecosystem trajectories could be very different from those that would be found in unaffected systems of the past or present, and they may not necessarily meet social or legal expectations (Luce et al. 2012). Using benchmarks based on our understanding of aquatic ecosystems today may also affect assessments of ecological consequences from natural and anthropogenic factors. We rephrase our statement from near the top of this section: the potential implications of these changes merit a primary research focus, but in the meantime it will be important that monitoring programs, whether they use reference conditions, least-disturbed conditions (e.g., Miller et al. 2017), decision-support models (e.g., Reeves et al. 2004), or other approaches, recognize and acknowledge these potential concerns in the process of analysis and the interpretation and application of results. (The topic of reference conditions is discussed in different contexts later in this report, including an expanded treatment of approaches for riparian restoration.)

Attribute integration approaches—

Integration of metrics—Watersheds and streams operate as integrated systems, and no one indicator is likely to accurately characterize their condition. A significant scientific challenge remains in how to reflect this integration in their assessment. Early efforts and ongoing regulatory guidance for assessing salmonid habitat look at a number of indicators individually, without an explicit procedure for integration (NMFS 1996, USDA FS et al. 2004). NRSA, the largest national assessment, also primarily reports on indicators separately (fish, macroinvertebrates, chemical and physical stressors), although many of their indicators are themselves composite metrics. They incorporate limited integration through their measure of relative risk, which looks at the likelihood of finding poor biological conditions in a river or stream when stressor concentrations are high, relative to the likelihood when they are low (USEPA 2016).

The AREMP monitoring plan was a pioneering attempt to integrate indicators into a composite watershed-condition index (Reeves et al. 2004). In practice, the extent of integration has declined in each of AREMP's reports. The 10-year AREMP report integrated all upslope

and in-channel variables into a single score for each watershed (Gallo et al. 2005). Trend was calculated only for the upslope portion; repeated measurements of sufficient in-channel sites were not available until the 20-year report. The 15-year assessment separated upslope and in-channel metrics for two reasons. First, the upslope data (GIS and remote sensing) covered the whole region, so there was no need to restrict that analysis to the in-channel subsample. Second, little correlation was found between the upslope and in-channel results, so it was believed that these outputs offered fundamentally different types of information. In addition, the mixing of stressors and responses has been criticized in other watershed indices (Schultz 2001).

The 20-year report maintained this upslope/in-channel split, and also reported the in-channel elements of physical habitat, macroinvertebrates, and temperature separately. Temperature was split off because it was collected under a different sample design, only at the lowest point in the watershed rather than at each site. Macroinvertebrates were separated from physical habitat because they are often considered a qualitatively different type of indicator: physical habitat as a condition or stressor, and macroinvertebrates as a response. It was also believed that reporting these indicators separately would better identify problems by not obscuring high and low values in an aggregated average. Although macroinvertebrate, physical habitat, and temperature data were not integrated in the 20-year report, additional analyses may be useful to further assess a combined metric, especially as novel techniques emerge that can address issues of spatial autocorrelation along linear stream networks (Peterson et al. 2013, Ver Hoef et al. 2014).

Other monitoring programs have built condition indices by combining indicators into a more integrated value. PIBO averages its physical channel attributes into a single index score, but maintains macroinvertebrates separately (Archer and Ojala 2016). The state of Oregon combines a number of chemical metrics into an overall water-quality index but also reports macroinvertebrates separately (Hubler 2009). The national Forest Service watershed condition

class combines upslope, riparian, and in-channel biotic and abiotic indicators into one overall watershed-condition score (Potyondy and Geier 2011).

Having the metrics for watershed condition assessed independently begs the question as to how to interpret overall condition relative to findings for instream habitat, stream temperature, and upland/riparian condition. Miller et al. (2017) acknowledged that reliance on a single biological metric can lead to erroneous interpretation of the biological condition of a watershed (e.g., Barbour et al. 1999), and suggested that the findings of the four separate stream metrics can be used as multiple lines of evidence to look at watershed-condition trends. So, when concordance among the measures differs within a given watershed, for example, if one or two parameters show an improving trend while the third does not change or declines, one can better understand which parameter may signal a potential issue and spur additional investigation. (See additional discussion of this issue later in this section.)

Interpreting long-term changes in a single metric can be complicated. For example, Miller et al. (2017) reported a broad-scale change in the distribution of water temperatures toward lower temperatures and an increase in watersheds with improving aquatic macroinvertebrate assemblages across the NWFP area. Nonetheless, 55 percent of waterbodies monitored by AREMP in Oregon still exceed state water-quality standards for these parameters (ODEQ 2012). One potential reason for this apparent lack of agreement between Miller et al. (2017) and the increase in miles of water-quality-impaired streams in Oregon (ODEQ 2012) is the lack of concordance among the indicators in a given watershed, as was shown in figure 7-5.

One parameter may be trending in a positive direction while another is outside or moves outside the acceptable range. Also, the number of streams surveyed by ODEQ for water-quality impairment has increased over the last 10

years, and differences in the way specific metrics are used may explain some of these apparent differences. More research is needed to fully assess whether such monitoring results represent favorable ecological changes over the long term.

Generally, analytical approaches and their interpretation for broad ecosystem assessments are still developing, including novel uses of individual metrics and multivariate methods. The initial NWFP monitoring strategy was based on identifying key stressors and a conceptual model that linked ecosystem and species components (Mulder et al. 1999). The decision-support system based on expert judgement used for interpretation of metrics in the 10-year report (fig. 7-6) is prone to subjectivity and uncertainty, however, there may be no real alternative given the needs of policymakers, the complexity of the




Figure 7-5—Relation between scores for overall watershed condition and condition of the macroinvertebrate communities for watersheds in the Northwest Forest Plan area in the assessment by Miller et al. (2017).




Figure 7-6—Example of a decision process for assessing the ecological condition of a watershed (Lanigan et al. 2012). AVE = average of scores; MIN = minimum; D50 = median particle size; EPT = Ephemeroptera, Plecoptera, Tricoptera index. See Reeves et al. (2004) for details.

ecological systems, and existing information gaps (Pielke 2007). The movement away from expert opinion in the 20-year report raises a new set of questions about use of single metrics interpreted independently and without an ecological framework. Carlisle et al. (2008) suggested that if one of the parameters of interest in a sampling unit (reach or watershed) is outside the threshold value for suitable conditions, the unit as a whole is outside the suitable range. A more statistically rigorous approach was described by Bowman and Somers (2006) and Collier (2009). In sum, exploration of a variety of available approaches is merited and timely.

Summarization over space and time—

A final consideration related to data integration is how to aggregate and present data over space and time. Reporting units over the spatial extents of land ownerships, land authority jurisdictions (e.g., county), and land-use allocations can address management and policy objectives, but the relationship of environmental conditions over space and time (e.g., via ecological provinces, watershed hydrological units) is also fundamental to our scientific understanding of aquatic-riparian ecosystems, particularly under the disturbance ecology paradigm (Olson et al. 2017b). Broad-scale aquatic assessments often report results by one or more aggregated spatial subunits.

Two types of units are evident in AREMP and other efforts. The first is based on ecological conditions. Similar to the other NWFP monitoring modules, AREMP reports results by seven aquatic provinces, which are similar to the physiographic zones developed in the NWFP planning process (FEMAT 1993: app. V-A). Broad ecological zones were similarly used in NRSA, both for the definition of reference conditions and the reporting of results (USEPA 2016). Hubler (2009) took a more water-centric approach, using water basins (hydrologic units, in particular HUC6, mean area $\sim 10,000 \text{ mi}^2$ [$\sim 25,900 \text{ km}^2$]) as their principal reporting unit. Studies focused on fish populations have developed units to represent genetically and demographically independent groups of fish, referred to as evolutionarily significant units (ESUs) or distinct population segments (DPS) (McClure et al. 2003).

A second type of spatial aggregation common in the aquatic-monitoring literature is by management units. AREMP reports on indicators by NWFP land-use allocations, which generally correspond to land use intensities. The National Water-Quality Assessment targeted their sample by land use disturbance levels and so displays many of its results by urban, agriculture, and mixed-use classes (Carlisle et al. 2013). Hubler (2009) reported their results by ownerships (federal, state, private industrial, private nonindustrial). PIBO's regional reports have not broken down their data by management classes directly, but they have displayed results of all sites in concert with reference distributions, indirectly reflecting management classes (Archer and Ojala 2016).

In earlier AREMP reports, the NWFP aquatic provinces were the basis of alternate evaluation criteria. However, as described above, the most recent assessment used neighborhoods based on in-channel characteristics, and upslope zones based on forest types. Because the aquatic provinces are no longer used to set environmental parameters for the assessment, other aggregations may prove more useful. For example, the NWFP provides a common set of standards and guidelines for management in the region; they are implemented via management plans developed by each agency for each of their forests and districts. AREMP may wish to consider reporting by these

unit and agency boundaries to increase their relevance to managers. Additionally, because these plans must address endangered species issues, the ESU and DPS boundaries may now be more relevant ecological units than the NWFP provinces.

AREMP's mandate is only to assess conditions on federal lands under the jurisdiction of the NWFP. However, nonfederal lands have been shown to be important, both for their effects on federal land conditions as well as for containing potentially productive fish habitat (Burnett et al. 2007, Reeves et al. 2016a, Van Horne et al. 2017). Further investigation in how to link AREMP data with assessments covering nonfederal lands could help address this gap, for an all-lands approach to watershed assessments in the moist coniferous forest ecosystem.

Because watershed conditions differ naturally over time, individual watershed ratings may be truly meaningful only when considered in some larger aggregation (Poole et al. 2004, Reeves et al. 2004). Thus, the end goal of AREMP was to look for changes in the distribution of watershed conditions in the whole NWFP area over time. As a baseline for comparison, AREMP chose to simply use conditions in the first monitoring period, because the other options considered (historical conditions, simulated natural conditions) would have been challenging to implement. Some other major programs have also used initial monitoring results as baseline conditions (Archer and Ojala 2016, USEPA 2016), whereas others have simply focused on one point in time, for example, using their most recent data (Carlisle et al. 2013, Hubler 2009). To our knowledge, historical conditions have not been estimated directly from past records, and simulation of natural conditions has been attempted only over considerably smaller spatial scales (Wondzell et al. 2007).

Finally, analytical approaches and their interpretation for broad ecosystem assessments are still developing, including novel uses of individual and multivariate methods. The rich AREMP dataset is ideal for comparison of alternative analytical approaches to assess watershed condition. Despite uncertainties and challenges faced by the 10-, 15-, and 20-year AREMP reports, reported trends support the intent of the ACS to sustain and improve conditions of federally managed watersheds in the NWFP area.

Components of the Aquatic Conservation Strategy

Riparian Reserves

Riparian reserves were intended to define and delineate the outer boundaries of the riparian ecosystem and to encompass the portions of a watershed most tightly coupled with streams and rivers (FEMAT 1993). These areas were assumed to provide the ecological functions and processes necessary to create and maintain habitat for aquatic and riparian-dependent organisms over time. This includes dispersal corridors for a variety of terrestrial and riparian-dependent organisms, and connectivity of streams within watersheds (FEMAT 1993). In 1993, the Forest Ecosystem Management Assessment Team (FEMAT) developed three management scenarios for riparian reserves along fish-bearing and non-fish-bearing streams (FEMAT 1993). Each scenario required a reserve width on fish-bearing streams of two times the height of a site-potential tree (minimum of 300 ft [91.4 m]), defined as the average maximum height the dominant tree would be expected to attain given the growing conditions at that location. On non-fish-bearing streams, the width of the riparian reserves varied from one-sixth of a site-potential tree-height (minimum of 25 ft [7.6 m]) to one-half of a site-potential tree-height to one site-potential tree-height (FEMAT 1993). One scenario was integrated into each of the 10 landscape alternatives developed and evaluated by the FEMAT (1993) scientists.

The Secretaries of the Interior and Agriculture selected FEMAT's Option 9 as their preferred option, which required a riparian-reserve network that was two site-potential tree-heights wide on fish-bearing streams and one-half of a site-potential tree-height on most non-fish-bearing streams. Interim boundaries of the riparian reserves were extended to a full site-potential tree-height on all non-fish-bearing streams between the draft and final environmental impact statements (USDA and USDI 1994a) to increase the likelihood of success of the ACS, and to provide additional protections from timber management and road building for non-fish organisms that use the area in or near streams as habitat or migratory corridors (USDA and USDI 1994a). On some fish-bearing streams, two site-potential tree-heights from the edge of a stream may not encompass the whole

floodplain, which can be an important source of large wood, making it critical to recognize and protect the entire floodplain (Latterell and Naiman 2007). This was accomplished in the ACS by requiring the boundary of the riparian reserve to extend to the edge of the 100-year floodplain (USDA and USDI 1994a). These boundaries were considered interim until a watershed analysis, which could adjust the size of the riparian reserve, was completed (USDA and USDI 1994a).

Depending on the degree of dissection of the forested landscape by streams, riparian reserves along both perennial and intermittent streams may occupy between 40 and 90 percent of the landscape (FEMAT 1993, Hohler et al. 2001). Interim riparian reserves of this magnitude, coupled with key watersheds and late-successional reserves, have provided a connected watershed-level reserve system for terrestrial, riparian, and aquatic ecosystems (Everest and Reeves 2007). However, the area of the forested landscape contained in the riparian reserves has fueled a controversy regarding riparian protection, resulting in new research to evaluate prescribed widths of riparian-management areas and a reexamination of existing scientific literature on the subject (Everest and Reeves 2007). The following sections summarize some of the recent key literature relating to the functions and size of riparian reserves.

Ecological functions—

The scientific basis for delineation of interim riparian reserves in the NWFP was derived from two sets of curves showing the relationship between various ecological functions provided by riparian zones and distance from the channel (figs. 7-7 and 7-8). These curves were developed by FEMAT scientists based on the scientific literature that was available at the time, and on professional judgment when sources of information were incomplete (see table 7-2 for original sources). The original relationships (FEMAT 1993) that were incorporated into the NWFP (USDA and USDI 1994a) suggest that most ecological functions could be maintained by reserves equal to or less than the distance of one site-potential tree-height. The functions include beneficial effects of root strength for bank stability, litterfall, shading to moderate water temperatures, and delivery of coarse wood to streams (fig. 7-7A). In addition, the majority of moderating effects on sediment delivery to streams




Figure 7-7—(A) Relation of distance from stream channel to cumulative effectiveness of riparian ecological functions (FEMAT 1993: V-27); (B) modified effectiveness curve for wood delivery to streams as a function of distance from the stream channel. The original curve was changed based on scientific literature developed since the original curve was portrayed in FEMAT (1993). Source: Spies et al. 2013.

from overland erosion associated with upland activities generally occur within a distance of one site-potential tree-height (Castelle et al. 1994, Naylor et al. 2012). The FEMAT scientists also provided a margin for error allowing for incomplete science, unknown cumulative effects, or strategic uncertainty in defining interim riparian reserves prior to watershed analysis. Everest and Reeves (2007) concluded that science published since original development of the FEMAT curves has generally supported the original assumptions and judgments.




Figure 7-8—(A) Relation of distance from stream channel to cumulative effectiveness of ecological factors influencing microclimate in riparian ecosystems (FEMAT 1993: V-27); (B) modified effectiveness curve for relative humidity as a function of distance from the stream channel. The curve was changed based on scientific literature developed since the original curve was portrayed in FEMAT (1993). Source: Reeves et al. 2016a.

Recent studies of wood recruitment suggest that changes in some of the ecological function curves may be supported. According to the graph of the relationship between the cumulative effectiveness of an ecological process and the distance for wood recruitment from the immediately adjacent riparian area in fish-bearing streams, developed in FEMAT (1993), about 60 percent of wood recruitment from the immediate riparian area along fish-bearing streams occurs within one-half of a tree-height (fig. 7-7A). This graph was based on a limited number

Table 7-2—Literature sources used to develop the original curves of ecological functions in riparian reserves in FEMAT (1993)

Function	Sources
Root strength	Burroughs and Thomas 1977 Wu et al. 1986
Wood delivery	Beschta et al. 1987 McDade et al. 1990 Van Sickle and Gregory 1990
Litterfall	Professional judgment
Shading	Beschta et al. 1987 Steinblums 1977
Microclimate	Chen 1991

of studies (McDade et al. 1990, Van Sickle and Gregory 1990) and the professional judgment of scientists involved with FEMAT. More recent studies on the sources of wood (Gregory et al. 2003, Spies et al. 2013, Welty et al. 2002) found that, at least in the Cascade Range of western Oregon and Washington, about 95 percent of the total instream wood inputs from the adjacent riparian area along fish-bearing streams came from distances of 82 to 148 ft (25 to 45 m) from the stream, representing a distance of 0.6 to 0.7 of a site-potential tree-height for this area (fig. 7-7B). The shape of this curve differs from the FEMAT curve (fig. 7-7A), which showed that 95 percent of the wood-recruitment function of the same streams occurs within a distance equal to about 0.95 of the height of a site-potential tree.

A primary purpose for the extension of the boundary of the riparian reserve of the NWFP from one site-potential tree-height to two on fish-bearing streams was to protect and enhance the microclimate of the riparian ecosystem within the first tree-height (USDA and USDI 1994a). At the time the ACS was developed, the only research on the effects of clearcutting on microclimatic conditions in adjacent forests had been done in upland forests on level terrain (Chen 1991). Those studies found that the influence of recent clearcuts (10 to 15 years old) extended from tens of yards (e.g., soil moisture and radiation) to hundreds of yards (e.g., wind velocity) into adjacent unharvested stands. Based on the initial work of Chen (1991), FEMAT (1993) hypothesized that a second tree-height could provide a considerable safety margin from

negative effects of intensive management on riparian areas, in terms of relative humidity and other microclimatic effects in the riparian reserve along fish-bearing streams (FEMAT 1993) (fig. 7-8A).

Since the ACS and associated ecological-function curves were originally formulated, a number of research efforts have examined the effects of forest management on microclimate in riparian areas. The vast majority of this work has focused on air temperature and relative humidity in small, headwater streams; few studies were conducted along larger streams (see review by Moore et al. 2005; also Olson et al. 2007, 2014). The magnitude of harvest-related changes in microclimate in riparian areas is usually inversely related to the width of the riparian buffer and the type and extent of management activities on the outer (upslope) edge. Some studies failed to show any edge effect between clearcuts and riparian buffers composed of intact mature forest (i.e., the extent of change in microclimatic conditions resulting from the presence of a clearcut on upslope edge of the riparian area) (Anderson et al. 2007, Rykken et al. 2007). Other studies have found that edge effects varied from a distance of 98.5 ft (30 m) (Anderson et al. 2007, Rykken et al. 2007) to 148 ft (45 m) (Brososke et al. 1997) from the stream. At the other extreme, Ledwith (1996; as cited by Moore et al. 2005) found that above-stream temperature decreased and relative humidity increased as buffer widths increased up to 492 ft (150 m). Rykken et al. (2007) attributed the lack of an edge effect to a “stream effect,” described by Moore et al. (2005), who noted that the stream can act as a heat sink and a source of water vapor during the day, thus keeping near-stream microclimates cooler and more humid than areas farther from the stream. Rykken et al. (2007) suggested that this stream effect might counteract edge effects of harvest on microclimate, thereby reducing the distance that harvest effects penetrate into riparian zones, relative to the distances measured in upland forest edges (e.g., from those projected by Chen et al. [1993] in uplands). Moore et al. (2005) also posited that cool, moist air might be carried by down-valley breezes, contributing to this stream effect.

The FEMAT microclimate curves were based on upland studies of forest-edge effects and thus they do not necessarily

apply to riparian areas with a strong stream effect, protected topographic position, and retention of some canopy in the adjacent managed stand. Reeves et al. (2016a) suggest that a one tree-height buffer on fish-bearing streams (fig. 7-8B) would reduce most potential effects on microclimate and water temperature in near-stream environments from timber harvest in areas on the edge of the riparian reserve, particularly when some trees are retained in the harvest unit. In general, most studies show that microclimatic changes in temperature and relative humidity seldom extend farther than one site-potential tree-height from the clearcut edge into an intact riparian buffer composed of mature forest (see review by Moore et al. 2005 and references cited therein). However, the large variety of effects measured in different studies demonstrates that substantial uncertainties remain about the size and management of riparian reserves. These uncertainties have important implications when considering changes in the width of the NWFP riparian reserves.

Increased stream temperature following forest harvest is one of the most frequently mentioned management concerns, and one that retention of riparian buffers is clearly designed to mitigate. Generally, the smaller the riparian area and the more extensive the activities, the greater the effect on stream temperature. Clearcut logging without riparian buffers usually leads to large, post-harvest increases in stream temperature, and the width of the riparian buffer needed to limit, or even eliminate, temperature increases remains uncertain (see reviews by Moore et al. 2005 and Leinenbach et al. 2013). Given these uncertainties, management prescriptions that reduce the width of the riparian reserve or allow some tree harvest within the reserve remain controversial.

The NWFP area encompasses a wide array of bioclimatic conditions, across its latitudinal span, west to east with distance from the ocean and rain-shadow effects of mountain ranges, and with increasing elevation. Given this variation, we describe only a few broad, general patterns in riparian vegetation here. In subsequent sections, we contrast these patterns with present-day patterns in areas that were previously logged, especially where logging pre-dated the establishment of current forest-practices rules and allowed harvest right up to streambanks. Although these general trends are important

considerations, we also emphasize that more detailed local knowledge will be critical for determining appropriate management goals and planning specific actions.

Riparian and upland vegetation along headwater streams in moist or wet forest types is typically dominated by conifers (Nierenberg and Hibbs 2000, Pabst and Spies 1999, Sheridan and Spies 2005) (fig. 7-9A). However, conifer density can be lower in riparian zones compared to adjacent terrestrial areas (Sheridan and Spies 2005), hardwoods are uncommon in both riparian and upslope




Figure 7-9—Conceptual representation of (A) vegetative conditions in headwater and (B) mid-order streams in the Northwest Forest Plan area.

areas, and there are no clear differences in shrubs between the two zones (Sheridan and Spies 2005). Many mosses and liverworts are also found at the wetted edges of small streams or on wood and rock in and along the channels (Hylander et al. 2002). In these wet forest types, tree canopies are often dense, limiting sunlight and therefore primary productivity. As a result, headwater streams depend on allochthonous (coming from outside the stream) inputs of litter and terrestrial invertebrates from the adjacent riparian forests, with as much as 95 percent coming from within 45 to 83 ft [13.6 to 25.3 m] of the channel (Bilby and Heffner 2016), as the primary energy source for aquatic and riparian organisms in these streams and for those lower in the network (Baxter et al. 2005, Gomi et al. 2002, Leroy and Marks 2006, Richardson et al. 2005, Wallace et al. 1997, Wipfli and Baxter 2010). Allochthonous material is exported to downstream areas as dissolved organic carbon, coarse (>1 mm, >0.04 inches) particulate organic matter, and to a much lesser extent, as fine particulate organic matter (Gomi et al. 2002, Richardson et al. 2005), which contribute to the productivity of fish-bearing streams. The structure and composition of the riparian vegetation determines the quality, quantity, and timing of the allochthonous input (Cummins et al. 1989, Frady et al. 2007), all of which influence overall stream productivity.

Forests in riparian zones and adjacent uplands become increasingly different as the size of streams increases (fig. 7-9B). In the middle portions of stream networks, the riparian forest is more diverse than along headwater streams (Acker et al. 2003, Johnson et al. 2000, Sarr and Hibbs 2007). Riparian forests along many mid-sized streams still remain dominated by conifers, but they are often mixed with deciduous trees such as alder (*Alnus* spp.), big-leaf maple (*Acer macrophyllum*), willow (*Salix* spp.), and cottonwood (*Populus* spp.). Big-leaf maple and California black oak (*Quercus kelloggii*) can be common in riparian zones in the southern portion of the NWFP area.

Some studies in mid-sized streams have shown that conifer basal area nearest the stream can be lower owing to reduced survival from disturbances such as flooding (Hibbs and Giordano 1996, Pabst and Spies 1999). Also, the availability of growing sites might be limited; conifers preferen-

tially establish on “microtopographic ridges” created by old tree-falls and behind wood jams (Fetherston et al. 1995). As a result, tree density near the stream can be about half of that of upland stands (Acker et al. 2003, Rot et al. 2000, Wimberly and Spies 2001). In contrast, Pollock et al. (2012) suggested that there is little difference in tree density between upland and riparian stands. All these studies, however, did find that the basal area of conifers in streamside stands is greater than in stands farther from the channel or in adjacent uplands. Streamside trees can be among the largest in a watershed (Poage and Tappeiner 2002), and thus can be the source of the largest down trees (conifers) found in the channel (i.e., the key pieces) (Rot et al. 2000), which are generally recruited to the channel by undercutting at high water (Abbe and Montgomery 2003, Acker et al. 2003, Benda et al. 2003).

In the Oregon Coast Range, hardwoods were most abundant in the area closest to the channel of streams in the middle portion of the stream network (Pabst and Spies 1999, Wimberly and Spies 2001), particularly in unconstrained reaches (Acker et al. 2003), and they decreased in density moving away from the channel. This mix of hardwoods and conifers is important ecologically (Sponseller and Benfield 2001, Sponseller et al. 2001) and is frequently maintained by periodic flooding (Sarr and Hibbs 2007). The vegetative diversity provides diverse habitat for a suite of terrestrial and riparian organisms; hardwoods are especially important for riparian mollusks (Foster and Ziegler 2013) and Neotropical migrant and resident bird species (Pearson and Manuwal 2001). Riparian areas dominated by hardwoods, particularly nitrogen-fixing red alder (*Alnus rubra*), have the potential to increase primary (Cornwell et al. 2008, Kominoski and Pringle 2009, Kominoski et al. 2011, Schindler and Gessner 2009, Swan et al. 2009) and secondary productivity and invertebrate diversity in adjacent streams (Piccolo and Wipfli 2002, Srivastava et al. 2009, Wipfli and Musslewhite 2004). Watersheds with mixed hardwood/coniferous riparian vegetation in the Oregon Coast Range received nearly 30 percent greater influx of terrestrial invertebrate biomass than streams with conifer-dominated riparian areas (Romero et al. 2005). The loss or reduction of deciduous litter could potentially influence the structure, composition, and productivity of riparian and aquatic biota (Wallace et al. 1997, 1999).

Although these broad general patterns hold across much of the NWFP area, we do not mean to give the impression that the riparian forests and their adjacent uplands were uniformly forested. Rather, these forests were a complex, shifting mosaic of vegetation patches, presenting a landscape with great spatial variability and temporal dynamics (see more detailed discussion in chapter 3). Wildfire was the primary factor driving forest dynamics across the Oregon Coast Range (Wimberly et al. 2000) and other parts of the region (see chapter 3), although windthrow, insects, and disease can also be important. As a consequence, upland forests, even when assessed at large spatial scales, showed substantial variation in the area and ages of forests. For example, over long periods, the proportion of upland forest in old-growth condition, when summed over areas of more than 4.9 million ac (2 million ha), ranged from 25 to 75 percent; at the scale of late-successional reserves as specified in the NWFP (~98,842 ac [40 000 ha]), the amounts of old growth could range from 0 to 100 percent (Wimberley et al. 2000).

In addition to the factors described above for upland forests, riparian forests are also influenced by fluvial and geomorphic processes such as floods, debris flows, and bank erosion. State-and-transition simulations of the natural disturbance regime in the Oregon Coast Range (Wondzell et al. 2012) showed that 51 percent of the riparian network was in mature forest (stand age of 66 to 200+ years). The simulations also showed that the long-term average forest composition was highly variable. Only 2 percent of the riparian network was in a nonforested condition, 28 percent was alder dominated, 40 percent was in mixed alder/conifer stands, and only 29 percent was in conifer-dominated stands (see table 5 in Wondzell et al. 2012). The specific results cited above pertain only to the central Oregon Coast Range, but it is clear that no single condition—defined by stand composition, structure, and tree age—can represent the full distribution of naturally occurring conditions over large areas. Rather, riparian forest conditions, when assessed over broad landscapes, showed a distribution of conditions that resulted from the combined influences of natural disturbances and plant succession.

Human impacts and restoration—

Riparian forests throughout much of the NWFP area have been changed by the land use activities that have taken place over the past century. As a consequence, the present-day forests may frequently differ in structure and composition from the presettlement forests that preceded them (McIntyre et al. 2015, Naiman et al. 2000, Swanson et al. 2011). This is particularly evident in an estimated 30 to 50 percent of the riparian ecosystems in the NWFP area that have been converted to plantations, based on the percentages of plantations in upland forests (see chapter 3). Riparian forested areas were harvested extensively, often to the edge of the stream, prior to the advent of current policies (Everest and Reeves 2007). In many cases, the riparian zones were subsequently planted with the most commercially valuable conifers, primarily Douglas-fir (*Pseudotsuga menziesii*), resulting in the development of dense, relatively uniform conifer stands and a decrease in hardwoods. In other cases, conifers were not successfully reestablished in logged riparian zones that are now dominated by alder with a dense salmonberry (*Rubus spectabilis*) understory, as observed at the reach scale by Hibbs and Giordano (1996). In watershed-scale simulations, Wondzell et al. (2012) estimated that, under historical conditions, 28 percent of the stream network in the Oregon Coast Range was in alder-dominated riparian forests, and that presently it is more than 40 percent. Fire suppression in dry forests with high- or moderate-frequency fire regimes has likely altered the structure and composition of riparian vegetation in ways similar to those described for upslope forested areas—namely an increase in the density of shade-tolerant conifers and a reduction in hardwoods (see chapter 3). In moist forests, with infrequent fire regimes, fire suppression has likely reduced the area of early-seral conditions in uplands and riparian areas (see chapter 3). Similarly, the removal of large conifers along rivers in the coast redwood (*Sequoia sempervirens*) range of northern California has been associated with increased dominance by alder (Madej et al. 2006). Clearly, the direct effects of logging on the structure and composition of present-day riparian forest can be varied, but overall, the distribution of conditions has changed dramatically relative to those under natural disturbance regimes.

Indirect effects of logging have also modified riparian forests. For example, rates of landslides and debris flows have increased in heavily roaded and logged watersheds (Goetz et al. 2015, Guthrie 2002, Jakob 2000), which has led to systematic changes in riparian vegetation. Debris-flow tracks are frequently scoured free of large wood and subsequently recolonized by red alder (Russell 2009, Villarin et al. 2009), with large wood deposited in runout zones. Further, the frequency of debris flows and landslides has contributed additional sediment to stream channels, driving more severe floods, with the combined effect of increasing the width of stream channels (Lyons and Beschta 1983). Exposed gravel bars within these channels are most often colonized by hardwoods, leading to substantial changes along the stream corridor.

Restoration challenges—

The changes to riparian forests described above create substantial challenges for restoration. For example, thinning of dense riparian Douglas-fir stands could open stands, allowing increased hardwood presence and thereby increasing the diversity of riparian vegetation, while also promoting

growth of the remaining trees to decrease the time needed to grow trees large enough to act as key structural elements in the stream channel. Although such restoration treatments may speed the restoration of some ecological functions (USDA and USDI 1994a), they also may reduce dead wood (chapter 3), and may present risks, such as development of novel conditions and loss of a particular species or ecological condition. Concerns about the tradeoffs between potential gains and potential losses, or other management issues, appear to have limited restoration activities, particularly within the first site-potential tree-height of streams. Reeves (2006) estimated that 48,000 ac (19 400 ha) of riparian reserve in the matrix of the NWFP area was treated for restoration purposes using some form of vegetation management, primarily thinning, in the first 10 years of the Plan. Between 2010 and 2015, an additional estimated 38,719 ac (15 669 ha) in the Forest Service's Pacific Northwest Region (Region 6) were commercially or noncommercially thinned (table 7-3). FEMAT (1993) estimated that 2.2 million ac (890 000 ha) of riparian reserves were outside of other reserves and congressionally withdrawn areas in the NWFP area.

Table 7-3—The estimated area of riparian reserves in the Northwest Forest Plan area in the Pacific Northwest Region of the U.S. Forest Service where active management that produced trees for commercial (primarily in the second tree-height of the riparian reserve) and restoration (noncommercial) purposes has occurred in 2010–2015

National forest	Area of riparian reserve managed			
	Commercial		Noncommercial	
	<i>Hectares</i>	<i>Acres</i>	<i>Hectares</i>	<i>Acres</i>
Deschutes	168	415	461	1,139
Fremont-Winema	0	0	0	0
Gifford-Pinchot	1031	2,548	301	744
Mount Baker–Snoqualmie	125	309	0	0
Mount Hood	674	1,665	0	0
Okanogan–Wenatchee	331	818	2150	5,313
Olympic	750	1,853	454	1,122
Rogue River–Siskiyou	142	351	616	1,522
Siuslaw	3923	9,694	203	502
Umpqua	883	2,182	622	1537
Willamette	2835	7,005	No data	No data
Total	10 862	26,841	4807	11,878

Source: USDA Forest Service Pacific Northwest Region.

Because this is not the total area of riparian reserves, it is not possible to estimate the fraction of the riparian reserve in the NWFP area that has undergone restoration. However, it is clear that the area that has been treated represents a relatively small proportion of the riparian reserves in total, and of the amount that has been altered by past activities.

Primary reasons for the limited amount of restoration activity are various and probably include (1) differing perspectives about the characterization of reference conditions, conservation, and management; (2) concerns about the potential effects of mechanical treatments on stream temperature and wood recruitment; (3) concerns about rare and little-known organisms that made managers reluctant to alter default prescriptions (Reeves 2006); and (4) trust (see chapter 12). We explore the potential challenges associated with these restoration activities below.

Reference condition versus restoring function—

Restoration activities necessarily require a “target” condition or conditions toward which the restoration activity is intended to move a system. One way to select a target for restoration goals is to identify a minimally disturbed condition and use it as a reference to which the current condition can be compared. The minimally disturbed condition is commonly called the reference condition. Although intellectually appealing, the selection of a reference condition is fraught with potential biases. For example, Pollock et al. (2012) set very stringent requirements on stand attributes that would be acceptable as a reference condition for Douglas-fir-dominated stands in riparian forests of western Washington state: choosing undisturbed, single-storied, conifer-dominated stands ranging in age from 80 to 200 years, and excluding stands dominated by hardwoods or shrubs that showed evidence of recent severe disturbance—including disturbances such as wildfire, insects, and disease—because these disturbances may themselves have been a product of fire exclusion or climate change. Also, stands with these features were assumed to be the successional climax forest for this size of stream, and such stands had been greatly reduced by logging.

The study by Pollock et al. (2012) illustrates some of the challenges inherent in finding reference conditions—they are often rare and may not represent the historical range of conditions that existed before extensive anthro-

pogenic modification of upland and riparian vegetation. Ideally, reference conditions would be identified in areas with similar potential vegetation and in relatively close proximity, or at least within the same ecoregion, so that the reference provides an appropriate comparison for similar forest stands (NRC 2000). Because of their focus on older conifer-dominated patches and the assumption that these types of stands represented the primary natural vegetation of streambanks in more confined terrain, Pollock et al. (2012) were able to identify only a small portion of the existing riparian forests in which stands meeting their criteria for dominance by older Douglas-fir trees could be found (only 117, or 3.3 percent, of the 3,521 potential sites met the filtering criteria). These stands were widely scattered, spanning a broad latitudinal and climatic range in western Washington. Further, they lumped together both upland stands and riparian stands, and the only riparian reference stands were located in the western Washington Cascade Range. Thus, the applicability of the results to other stand types and locations is very limited. Nonetheless, Pollock et al. (2012) illustrated that a stringent filtering approach to identifying reference sites could contribute to characterization of reference conditions at the patch or stand scale (e.g., stand density and tree size) for evaluating riparian-management options for Douglas-fir riparian stands in western Washington. The Douglas-fir patch-scale reference conditions could also be used in setting management goals for the entire landscapes of a larger riparian zone. A similar approach could be applied to other stand types or regions to provide a more complete system of reference conditions for riparian management in the Pacific Northwest.

Another approach to setting reference conditions (although they did not call them “reference conditions” at the time) for riparian zones was used by Pabst and Spies (1999) and Nierenberg and Hibbs (2000) in the Oregon Coast Range. They sampled along first- through fourth-order streambanks without roads or a history of logging, and having no evidence of wildfire at least 80 years. Vegetation was sampled in transects from randomly selected starting points. Hence, the samples contained areas dominated by older conifers as well as patches of hardwoods and shrubs, in proportion to their occurrence in the riparian area. Smaller,

recent areas of geomorphic disturbances, disease, and wind-throw would have been included in the samples. The studies focused in particular on how vegetation differed between streambanks and uplands at a site, controlling for differences in environment and disturbance history. Generally, these studies found that conifer dominance decreased from the uplands to the streambank, and that many areas within 53 ft (16 m) of streams were typically a mosaic of conifers, hardwoods, and shrubs even along streams in relatively confined topographic settings. Conifer-snag densities were relatively low 17 snags/ac (6.9 snags/ha) within 53 ft (16 m) of streams, and about one-quarter of the densities found at distances of more than 53 ft (16 m) from streams (Pabst and Spies 1999). These studies could be used in developing management guidelines for riparian forests in the Oregon Coast Range. Note, however, that this approach (random samples of unmanaged riparian vegetation) did not sample many areas that had grown for several centuries since wildfire and also did not sample in large areas that were recently affected by fluvial, geomorphic, and fire disturbances, which would have been an important part of the historical range of variability in these ecosystems at watershed scales (Spies et al. 2002).

Wondzell et al. (2012) used state-and-transition models to explore the range of ecological states of the riparian network of a large river network in the central Oregon Coast Range. They used GIS methods to partition the stream network and its valley floor into discrete reaches, which were classified into potential geomorphic and vegetation types. A state-and-transition model was then developed for each potential type that included all possible states that could result from succession, natural disturbance, and land use activities. Wondzell et al. (2012) found that the structure and composition of the current riparian vegetation differed from the historical; there were fewer conifers, particularly the largest (>30 inches [76.2 cm]), and more alder-dominated patches. They clearly stated that their simulation results “should be interpreted as hypotheses of likely outcomes,” and that, despite several model limitations, they can be used to “hindcast” expected historical distribution of riparian forest conditions.

Part of the debate about restoration needs for riparian areas may derive from differing views of riparian reference conditions (as a goal for restoration), and how they differ

with scale and across watersheds and the NWFP region. Although many studies (e.g., Acker et al. 2003, Hibbs and Sarr 2007, Pabst and Spies 1999) have found that riparian vegetation and upland vegetation frequently differ in structure, composition, and dynamics depending on stream size, some have noted that differences between riparian and upland vegetation may be small for some stand types, and that in some cases upland sites can supplement riparian sites to increase sample size for describing target conditions for riparian management. For example, Pollock et al. (2012) noted that, for Douglas-fir-dominated stands in western Washington, “both forest types [upslope and riparian] are generally similar, but riparian stands have more live tree wood volumes and basal areas, suggesting they may be growing on sites that are more productive.” Therefore, they concluded that riparian restoration in Douglas-fir-dominated riparian zones should aim to produce stand characteristics with densities and sizes of live and dead trees that are within the range of reference conditions (both upland and riparian). On the other hand, others (Gregory 1997, Pabst and Spies 1999, Welty et al. 2002, Wimberly and Spies 2001) have found that the type and magnitude of differences in features between upslope and riparian forests can be large, suggesting that upslope vegetation should not be assumed to be a reference for designing and assessing managed strategies for riparian vegetation in other stand types, or where riparian stands differ significantly from upland stands (e.g., in floodplains).

This variety of findings makes it difficult for managers and regulators to design and implement management actions in riparian reserves. We suggest that each of the approaches examined above—that of Pollock et al. (2012), Pabst and Spies (1999), Nierenberg and Hibbs (2000), and Wondzell et al. (2012)—offers important information that would contribute to building a “reference condition”-based strategy to examine current conditions, to project likely outcomes of planned management activities, and to help evaluate the tradeoffs between potential risks and benefits of any overall management strategy. For example, a modeling approach like the state-and-transition models of Wondzell et al. (2012) could be used to generate an expected historical distribution of states for the riparian vegetation

within a stream network. The results could be used to identify relatively little-disturbed watersheds across the ecoregion and the monitoring plots located within those watersheds; individual plots could then be compared to specific states in the state-and-transition model. Those states could be attributed with values for various metrics (e.g., cover, basal area, tree densities, snag densities, species composition), as was done by Pabst and Spies (1999). Because anthropogenically disturbed states are also included in the state-and-transition models, something similar could be done to attribute these states with empirical data. The models could then be used to hindcast the historical distribution of state classes, and descriptive metrics from the empirical data could then be linked to the historical distribution. This result could be compared to the current condition. Also, the models could be used in forward simulations, incorporating different land use choices, to project how the distribution of conditions might be expected to change over time in response to various management strategies. Whatever approach is used, it will be important for managers and regulators to understand the limitations of the research they use to design and support proposed actions, which in turn necessitates that researchers clearly identify the limitations of their research (such as how broadly or to what ecosystem type they can be applied), and recognize the large variation in the inherent structure and composition of riparian areas across the NWFP area.

Riparian thinning and water temperature—Because the current distribution of conditions of riparian forests in many stream networks is far different from the historical distribution, there is substantial interest in active restoration treatments—especially thinning dense conifer plantations (Reeves et al. 2016a) or logging hardwood-dominated stands and replanting to convert them to conifer dominance (Cristea and Janisch 2007). Although these treatments are not inconsistent with the ACS, which generally allowed thinning for ecological objectives in the area beyond 120 to 150 ft (36.6 to 45.7 m) to a distance of one site-potential tree-height, they could potentially exceed the 0.3 °C “non-degradation standard” for water-quality effects of logging. The 0.3 °C standard is important from a regulatory perspective, limiting potential cumulative effects from multiple actions,

none of which individually might be sufficient to impair water quality. Alternatively, restoration treatments might speed the attainment of desired future conditions. These decisions pose critical management challenges. Clearly, there are risks from any active restoration treatment, but choosing not to act also poses risks, not only by increasing the time needed to attain a desired future condition, but also leaving the riparian zone at greater risk of uncharacteristic disturbance—for example, dense conifer stands in dry forest zones are more prone to high-severity wildfire (see chapter 3). Also, there may be increases in primary production (Warren et al. 2016) and fish growth (Wilzbach et al. 2005) with the opening of the canopy along small and medium streams.

Reach-scale studies clearly demonstrate that solar radiation is the primary factor affecting stream-water temperatures during summer (Leinenbach et al. 2013). Thus, the likely effect of forest harvest on stream temperatures will be a function of the amount of shade lost. The largest effects will generally be seen with clearcut logging right to the streambanks, whereas retention of forested buffers tends to reduce these effects, as does thinning rather than clearcutting outside the buffer. The actual magnitude of stream-temperature increases can vary greatly and is determined by factors such as discharge, water depth, width, flow velocity, hyporheic exchange, and groundwater inflows (Janisch et al. 2012, Johnson 2004, Moore et al. 2005). Topographic shading can also influence water temperatures, particularly in small streams flowing in narrow, steep-sided valleys, as much as or perhaps more than shade from streamside forests (Zhang et al. 2017). It is important to remember that canopy removal also results in nighttime long-wave radiation loss, leading to lower water temperatures. This effect contributes to increased thermal variability, with poorly understood biological consequences.

Relatively few studies have examined the effects of riparian thinning on stream-water temperature. A few studies have examined clearcut harvesting combined with partial harvest of riparian buffers (Kreutzweiser et al. 2009, Macdonald et al. 2003, Mellina et al. 2002, Wilkerson et al. 2006). These studies, like those cited above, suggest that the effect of riparian thinning on summer stream temperatures will be correlated positively with the amount of forest

canopy removed and inversely with the distance from the stream that the activity occurs, and thus the amount of shade lost (Leinenbach et al. 2013). However, the amount of shade lost from a given thinning treatment can be highly variable, and the small number of studies makes it difficult to draw strong generalities. The amount of shade lost can be smaller than the amount of tree basal area removed, and in one study, removal of 10 to 20 percent of the basal area had no measureable effect on angular canopy density (Kreutzweiser et al. 2009). Further, any shade loss and stream-temperature increases from riparian thinning are likely to be short lived because riparian forest canopies can close relatively quickly (within 3 years) after thinning (Chan et al. 2006, Yeung et al. 2017). The potential magnitude of stream-temperature increases in response to riparian thinning will be highly dependent on forest attributes outside the riparian buffer, the buffer size, the prethinned riparian forest attributes (Leinenbach et al. 2013), the thinning prescription, and the thermal sensitivity of the stream (Janisch et al. 2012). Further research is needed to improve our understanding of the impacts of thinning, but there is some evidence that light thinning may not substantially increase stream temperatures.

Managers thus face the following question: Are there places in the stream network in which riparian thinning would help speed attainment of the reference distribution, and where present-day thermal regimes would suggest that small temperature increases would not have significant detrimental effects on fish (Groom et al. 2011) or other organisms of interest? This question tends to be investigated at the reach scale. For example, Pollock et al. (2012) examined the potential effects of a thinning treatment on the development of riparian forest-stand attributes, and Groom et al. (2011) looked at summer maximum temperatures in the treated reach. Rarely are these questions expanded to consider the context of the distribution of reference conditions across the larger watershed. If they were asked, the question would then become: Are the conditions of the treated reach overrepresented with respect to the reference distribution, or underrepresented? In the Oregon Coast Range, for example, it is clear that not all reaches would be maintained in conifer-dominated mature forest under

a natural disturbance regime (Wondzell et al. 2012). If dense, young, conifer-dominated stands are currently more abundant than expected from the reference distribution, then should some of those stands be thinned, perhaps mimicking windthrow events that open stand canopies and allow development of multistoried, mixed stands? If so, how many should be treated to better change the long-term trajectory of conditions from the current distribution toward one that is closer to the reference distribution?

Riparian thinning and large wood—The absence or diminished quantity of wood in streams throughout the NWFP area is a primary concern for managers and regulators because wood is important for creating habitat and performing other ecological functions. Thinning and other active management in plantations in riparian zones can reduce the potential amount of wood that can be delivered to streams (Beechie et al. 2000, Pollock et al. 2012) and the forest floor (Pollock and Beechie 2014, Pollock et al. 2012) if the trees are removed from the site. Additionally, thinning may negatively affect habitat, at least in the short run, for some species that are favored by dense conifer cover (see chapter 3 for more details), potentially increase water temperature (Leinenbach et al. 2013), and reduce carbon storage (D’Amore et al. 2015). However, there are also many potential benefits to thinning, including increasing structural diversity, species richness, flowering and fruiting of understory shrubs and herbs (Burton et al. 2014, Carey 2003, Hagar et al. 1996, Muir et al. 2002), and faster development of mature-forest conditions, including very large trees with thick limbs that may be used for nesting by marbled murrelets (*Brachyramphus marmoratus*) (Carey and Curtis 1996, Franklin et al. 2002, Tappeiner et al. 1997) (see chapter 5). Furthermore, variable density thinning of the overstory in the second-growth riparian forest could accelerate recovery of old-growth characteristics by promoting dominance of redwood in the southern portion of the NWFP area (Keyes and Teraoka 2014) (see chapter 3).

Considerable research on wood dynamics in the NWFP has been done in wet forests of California, Oregon, and Washington, but there has generally been less research in areas with drier forest types, including northern California. Riparian areas in redwood-dominated forests are

particularly distinctive owing to the exceptional productivity, low mortality, and slow decay of those trees (Benda et al. 2002). Benda and Bigelow (2014) compared wood volumes across four different regions of northern California, including the Coast Range, Klamath Mountains, Cascade Range, and Sierra Nevada, as well as variation associated with forest management. They noted that coastal streams had much greater wood volumes, which they attributed to greater forest biomass and higher growth rates of redwood forests, as well as slower decay of large wood pieces. They also observed that some second-growth forests along streams in that region had wood volumes comparable to those in old-growth forests, owing to heavy debris remaining from tractor-era logging before the 1970s. Although the volumes were similar, streams in old-growth areas had fewer but larger logs (Benda et al. 2002). Benda and Bigelow (2014) also found that streams in the Cascades and Sierras that they characterized as more heavily managed had larger volumes of stream wood than less intensively managed areas in the same regions. They conjectured that managed forests could have higher rates of tree mortality because of stem exclusion (successional phase characterized by the rapid growth and biomass accumulations of a particular species, and intense competition among cohorts [Oliver 1981]) than more mature, but not yet decadent, unmanaged forests with lower tree densities.

A panel of scientists from the Forest Service and the National Marine Fisheries Service recently reviewed the published literature on the effects of thinning in riparian areas (Spies et al. 2013). Their major conclusions are summarized below:

- Accurate assessment of thinning effects requires site-specific information. The effects of thinning regimes on dead-wood creation and recruitment (relative to no thinning) will depend on many factors, including initial stand conditions, particularly stand density, and thinning prescriptions.
- Conventional thinning generally produces fewer large dead trees. Thinning with removal of trees (conventional thinning) will generally produce fewer large dead trees across a range of sizes over the several decades following thinning and the lifetime of the

stand relative to equivalent stands that are not thinned.

- Thinning to develop old-growth structure is most beneficial in dense young stands less than 80 years old and especially those less than 50 years old.
- Conventional thinning can accelerate the development of very large diameter trees. In stands that are conventionally thinned, the appearance of very large diameter dead trees (greater than 40 in [102 cm]) may be accelerated by up to 20 years relative to unthinned plantations, depending on thinning intensity and initial stand conditions.
- To produce down wood immediately, thinning can leave trees that are cut as part of the restoration program (see Benda et al. 2016 for details).
- Thinning can increase the amount of pool-forming wood only when the thinned trees are larger in diameter than the average diameter of pool-forming wood (which varies with stream size).
- Effects of thinning on instream wood need to be placed in a watershed context. Assessing the relative effects of riparian thinning on instream wood loads at a site and over the long term requires an estimation of the likely wood recruitment that will occur from both the banks and downstream movement from upstream sources, and the rate of decay and downstream transport of wood from the site.
- The ecological effects of thinning on instream habitat will vary depending upon location in the stream network. Riparian-management practices can be altered to match the ecological functions of streams.
- Variation in thinning is essential to increase species diversity and heterogeneity (i.e., do not use the same prescription everywhere).

Since Spies et al. (2013) summarized the state of the science, other studies have increased our understanding of the effect of restoration thinning in riparian areas. Benda et al. (2015) simulated the idea of adding wood to channels during thinning by modeling the amount of instream wood that would result from thinning a 50- to 80-year-old Douglas-fir stand from below (i.e., removing the smallest trees to simulate suppression mortality) from 400 to 90 trees/ac (988 to 222 trees/ha), which is considered a moderate

amount of thinning, then directionally falling or pulling over varying proportions of the harvested trees into the stream (table 7-4). This wood loading was compared to the amount that would be expected in the stream if the existing stand was not thinned. Not surprisingly, the amount of wood increased above the “no-thin” level immediately after the tipping simulation in all the wood-addition options. However, the cumulative total amount of wood expected in the stream over 100 years relative to the unthinned stand varied depending on the amount of wood delivered. Adding ≤ 10 percent of the wood that would be removed during thinning resulted in less wood in the channel over time than the unthinned option (i.e., if the stand were not actively managed). When 15 to 20 percent of the volume of thinned trees from one side of the stream was directed to the stream at each entry, the total amount of dead wood in the channel exceeded the unthinned scenario over time (table 7-4). Thinning the stand again 25 years after the first thinning further increased wood levels (table 7-4). Carah et al. (2014) found that adding unanchored wood into the stream was less costly than securing the wood, and improved habitat conditions for coho salmon. Reeves et al. (2016a) included wood addition (tree-tipping) as a component of options for managing the riparian reserves on Oregon and California Railroad Revested lands of the BLM in western Oregon to accelerate attainment of restoration objectives.

Ecological tradeoffs—There are potential ecological consequences of limiting tree harvest (thinning) only to the outer portions of the riparian reserves. A myriad of ecological processes create and maintain the freshwater habitats of Pacific salmon (Bisson et al. 1997, 2009) and the ecological context in which they evolved (Frissell et al. 1997). This is especially relevant to the goals of the ACS, which are broad and include more than aquatic conditions. Holling and Meffe (1996) contended that uniform management prescriptions often fail when applied to situations in which processes are complex, nonlinear, and poorly understood, such as in aquatic ecosystems in the NWFP area, and may lead to further degradation or compromising of the ecosystems and landscapes of interest (Dale et al. 2000, Hiers et al. 2016, Rieman et al. 2006). For example, managing for a single purpose (e.g., maximizing dead wood) may compromise or retard other ecological functions, such as development of hardwoods and shrubs or growing large trees, in areas near the stream (see previous discussion), and ultimately may alter the structure of the food web (Bellmore et al. 2013). Pollock and Beechie (2014) stated that “species that utilize large-diameter live trees will benefit most from heavy thinning, whereas species that utilize large-diameter deadwood will benefit most from light or no thinning. Because far more vertebrate species utilize large deadwood rather than large live trees, allowing

Table 7-4—Predicted change in the estimated volume of wood in a stream channel under two different harvest options (single and double entry) over the simulated 100 hundred years (includes decay) when varying proportions of harvest wood are placed or felled into the stream channel

Scenario	Change from no treatment	
	Single-entry thin	Double-entry thin (25 years after first entry)
	----- Percent -----	
No treatment (reference)	0	0
Thin entire stand, no tipping	-33	-42
Thin entire stand except with a buffer 32.8 ft (10 m) no-harvest buffer	-7	-11
Thin with a buffer 32.8 ft (10 m) and tip 5 percent of the harvested wood	-15	-15
Thin with a buffer 32.8 ft (10 m) and tip 10 percent of the harvested wood	-6	+1
Thin with a buffer 32.8 ft (10 m) and tip 15 percent of the harvested wood	+1	+16
Thin with a buffer 32.8 ft (10 m) and tip 20 percent of the harvested wood	+6	+24

Source: Benda et al. 2016.

riparian forests to naturally develop may result in the most rapid and sustained development of structural features.” We agree that tradeoffs exist and that prioritization will be needed (see chapter 12 for more discussion of tradeoffs).

The choice of priority conservation targets (e.g., dead wood, plant-community diversity, large live trees, geomorphic disturbances) for riparian management is a difficult one to make, involving scientific criteria, risk assessment, and social values. Given the diversity of conditions in riparian areas at watershed and regional scales, it would make sense not to apply one-size-fits-all strategies, but rather to develop priorities based on a watershed-scale view (see “A context-dependent approach to riparian conservation and management” below). For example, Pollock and Beechie (2014) stated that “management strategies that seek to create a range of large live and dead tree densities across the landscape will help to hedge against uncertain outcomes related to unanticipated disturbances, unexpected species needs, and unknown errors in model assumptions.” It will be important to consider the full suite of ecological functions across a watershed; focusing only on one condition or metric may limit recovery of riparian ecosystems in ways that prevent full achievement of the broad objectives of the ACS. Given these broad objectives, a more comprehensive watershed- and regional-scale consideration of all ecological processes, and studies to develop new and more complete approaches, may be more fruitful than focusing on only one or two metrics.

A context-dependent approach to riparian conservation and management—

A key component of the ACS is watershed analysis (FEMAT 1993), which is supposed to provide the context of a given location for adjusting the boundaries of, and allowing activities within, riparian reserves. However, the intent of watershed analysis was never realized (Reeves et al. 2006), owing to a number of factors including cost of analysis and the need to consider a multitude of species and their ecological requirements. Neither FEMAT (1993) nor the NWFP (USDA and USDI 1994a) provided explicit criteria for changing the riparian reserve boundaries or demonstrating that proposed changes would meet or not prevent attainment of ACS objectives over the long term.

In addition, at the time, credible analytical tools to aid decisionmaking were lacking (Reeves 2006); a fixed-width approach is easy to administer and apply and is less costly than developing site-specific recommendations (Richardson et al. 2012). As a result, adjustments have proven difficult for the agencies to make (Naylor et al. 2012, Richardson et al. 2012), and interim boundaries of the riparian reserves remained intact in the vast majority of watersheds (Baker et al. 2006).

Since the development of the ACS, there has been a call in the scientific literature to allow discretion in setting site-specific activities (Kuglerová et al. 2014, Lee et al. 2004, Richardson et al. 2012), which can be economically beneficial (Tiwari et al. 2016). Greater flexibility in the management of riparian areas would depend on the “context” of the area of interest (Kondolf et al. 2006, Montgomery 2004), and the primary management objective for the specific area (Burnett and Miller 2007). However, development of such an approach has been limited because of the reliance on “off-the-shelf” and one-size-fits-all concepts and designs, rather than on an understanding of specific features and capabilities of the location of interest (Kondolf et al. 2003, Naiman et al. 2012). A mix of approaches could be undertaken, recognizing ecological and other goals such as timber harvest, especially if applied over larger spatial scales (Burnett and Miller 2007, Miller and Burnett 2008, Olson and Rugger 2007), and if consideration is given to the distribution of populations of concern and connectivity among them (Olson and Burnett 2009, Olson and Kluber 2014, Olson et al. 2007).

There have been a few attempts to design and implement a site-specific approach. Cissel et al. (1999) proposed a plan based on variation in the disturbance patterns (in this case, wildfire) in the target watershed, and called for harvest of some older trees and a revision of the interim riparian reserves for the Central Cascades Adaptive Management Area. Olson and Rugger (2007) proposed a two-tiered approach to riparian management to first identify reaches in which sensitive species occur, then manage their critical habitat elements, hence varying riparian reserve management with species distributions. Olson and Burnett (2009) applied sensitive-species filters to criteria for designations

of habitats for connectivity within and among watersheds. Interwatershed connections provided by riparian areas are critical avenues of movement to new habitats. None of these approaches have been implemented to date.

Reeves et al. (2016a) proposed a context-dependent approach for management of the riparian reserves in the matrix of federal lands in western Oregon that divided the riparian reserve into inner and outer zones, with management tailored to the specific features and characteristics of individual stream reaches (Option B of Reeves et al. 2016a). The context-dependent option was informed by new research, tools, and concepts, including:

- The influence of the width of riparian area on microclimate (see earlier discussion).
- Movement of amphibians along non-fish-bearing streams (Olson and Burton 2014, Olson et al. 2007).
- The distance to, and sources of, wood for fish-bearing streams (Spies et al. 2013).
- Intrinsic potential, a concept for assessing the capability of a given set of geomorphic conditions in a stream reach to provide habitat for selected species of Pacific salmon (Burnett et al. 2007).
- NetMap (Benda et al. 2007), a geospatial platform for watershed analysis that can, among other things, identify the location of some key ecological processes that influence aquatic and riparian ecosystems on the landscape and in the stream network.
- Concepts for managing riparian ecosystems and the activities that affect them, such as ecological forestry (Franklin and Johnson 2012) and tree-tipping (Benda et al. 2016).

Under the context-dependent option, current interim riparian reserves of two site-potential tree-heights along fish-bearing streams and one site-potential tree-height along non-fish-bearing streams would be retained in late-successional reserves and other special land designations (Reeves et al. 2016a). In lands allocated as matrix under the NWFP, the area of interest for aquatic conservation (Reeves et al. [2016a] referred to this as the riparian conservation area) extended upslope from the stream for a distance equal to the height of one site-potential tree along fish-bearing and non-fish-bearing streams. The riparian conservation area

was divided into an inner and an outer zone depending on “ecological context,” based on four characteristics of each stream reach—susceptibility to surface erosion, debris flows, thermal loading, and habitat potential for target fish species—to determine the width of the inner zone. The entire riparian conservation area of the most ecologically sensitive stream reaches along fish-bearing and non-fish-bearing streams could be managed solely for ecological goals for fish and other aquatic and riparian-dependent biota. In other fish-bearing and non-fish-bearing streams, the inner zone was 100 ft (30.5 m) and 50 ft (15.3 m) wide, respectively (Reeves et al. 2016a). Active management was limited to stands age 80 years or younger in reserves (Spies et al. 2013), and tree-tipping (Benda et al. 2016) was used throughout the riparian reserve to ensure that harvest did not negatively affect wood recruitment to the stream (table 7-4).

Using the matrix in BLM-managed lands in western Oregon to illustrate the application, Reeves et al. (2016a) estimated that an average of 46 percent of the riparian reserve in a watershed would be managed solely for ACS goals. Also, an estimated average of 36 percent would achieve ACS goals along with other potential goals, which could include timber production, and 18 percent could be managed for a variety of purposes, including wildlife and timber, in accordance with NWFP requirements (Reeves et al. 2016a). In late-successional and other reserve allocations, which cover approximately half of the BLM lands in western Oregon, interim riparian reserves would remain unchanged. Assuming that half of the interim riparian reserves on BLM lands in western Oregon would remain unchanged, and applying their study estimates of changes in matrix, Reeves et al. (2016a) estimated that about 72 percent of the interim riparian reserves would remain solely devoted to ACS goals, and 19 percent would likely meet ACS goals and could also provide opportunity for achievement of matrix goals including limited timber production. The reduction of the width of the riparian reserve along fish-bearing streams to one tree-height would return an estimated 9 percent of interim riparian reserves to matrix on these lands.

The analysis of Reeves et al. (2016a) was not intended to provide a single recommendation for managing riparian

ecosystems. The primary purpose was to reevaluate riparian-conservation strategies using the latest scientific evidence. This or other options should be viewed as working hypotheses to be tested with monitoring and adaptive-management experiments. The analysis provides an example of how a context- and landscape-dependent approach could be designed to address multiple conservation goals of the ACS, the commodity goals of the NWFP, and the significant challenges of climate change. Although new science has refined our understanding of the ecological processes in riparian ecosystems, uncertainties and information needs remain. Thus, an adaptive-management approach and further research are critical to continual improvement and evaluation of this and other options for meeting the goals of the ACS (Stankey et al. 2005).

Key Watersheds

Tier 1 key watersheds (a total of 141, covering 8,154,500 ac (3 300 000 ha) (fig. 7-10) were intended to serve as refugia for aquatic organisms or to have high potential for restoration (USDA and USDI 1994a). Tier 2 key watersheds provide sources of high-quality water, and comprised 23 watersheds covering a total of about 1,112,000 ac (405 000 ha) (fig. 7-10). Key watersheds are aligned as closely as possible with the late-successional reserves of the NWFP (areas designated to protect late-successional and old-growth ecosystems) and other officially designated reserve areas to maximize ecological efficiency (USDA and USDI 1994a), and to minimize the amount of area in which timber-harvest activities were restricted. A primary objective for tier 1 key watersheds is to aid in the recovery of ESA-listed fishes, particularly in the short term (FEMAT 1993). Tier 1 key watersheds in good condition at the time of the Plan's inception were assumed to serve as centers for potential recovery of depressed populations. Those with degraded conditions were expected to have the greatest potential for restoration and to become future sources of good habitat.

The trend in the condition of key watersheds differed among assessments. Gallo et al. (2005) reported that a greater proportion of the key watersheds had their condition scores improve than did non-key watersheds. Lanigan et al. (2012) found key watersheds to be in better condition than




Figure 7-10—Location of key watersheds in the Northwest Forest Plan area.

P. Eldred

non-key watersheds, primarily because more than twice as many miles of roads were decommissioned in key watersheds as in non-key watersheds (Gallo et al. 2005, Lanigan et al. 2012), suggesting that land management agencies appear to recognize key watersheds as priority areas for restoration. Miller et al. (2017), however, saw no statistical differences between the two groups.

Key watersheds were originally selected based on the professional judgment of the scientists involved with the development of the ACS, in consultation with fish and aquatic biologists and hydrologists from the national forests and BLM districts covered by the NWFP. Also, they were tightly aligned with late-successional/old-growth reserves, based in part on the assumption that streams in old-growth forests would be most favorable for fish (FEMAT 1993). New techniques (e.g., NetMap, Benda et al. 2007) and understanding of aquatic ecosystems now provide a different perspective on aquatic ecosystems and how they operate in space and time.

New concepts such as intrinsic potential of fish habitat (Burnett et al. 2007), projections of climate change, and new questions as to whether stream conditions in old-growth forests are actually most favorable for native salmonids (Bisson et al. 2009, Reeves and Bisson 2009, Reeves et al. 1995) are pivotal concepts that reframe our understanding of aquatic ecology and ecosystems. No formal evaluation of the potential effectiveness of the network of key watersheds was conducted during development of the NWFP, or has been undertaken since it was implemented. Fish populations in need of attention are clearly identified now, and it would be useful to investigate whether the current system is beneficial to those fish in terms of the overall distribution and the suitability of individual watersheds. Additionally, the distribution of other sensitive aquatic-riparian species (e.g., ESA-listed or petitioned herpetofauna) could contribute to this assessment.

Watershed Analysis

Watershed analysis was designed to provide the context for management activities in a particular sixth-field watershed as the basis for developing project-specific proposals and determining restoration needs. It was envisioned as an

analytical and not a decisionmaking process, involving individuals from a variety of scientific disciplines (USDA and USDI 1994a). Management agencies were expected to complete a watershed analysis before activities (other than minor ones) were initiated in key watersheds or riparian reserves (USDA and USDI 1994a). The version of watershed analysis advocated in the NWFP differs from previous versions (e.g., Washington Forest Practices Board 1993) and involves multiple disciplines and issues other than those that are specifically aquatic.

Baker et al. (2006) estimated that about 500 watershed analyses had been done by 2003, but that the quality and effectiveness of these analyses differed widely. No formal assessment of watershed analyses has been completed as of this writing, so it would be prudent to conduct a comprehensive review and evaluation, and consider incorporating new analytical tools such as NetMap (Benda et al. 2007) to help improve the process and reduce costs while increasing the usefulness of the product. The watershed analysis process could also be reexamined so that it is conducted more efficiently and considers the appropriate spatial scales, including the watershed of interest and its context within the larger basin or region. The latter could be particularly relevant for effective planning at a landscape scale and to deal with climate change.

New Perspectives on Conservation of Riparian and Aquatic Ecosystems

The ACS was premised on the view that aquatic ecosystems were dynamic in space and time, exhibiting a range of potential conditions, similar to the terrestrial systems in which they are embedded (FEMAT 1993). Aquatic ecosystems in Pacific Northwest forests are multifaceted and complex, and can be conceptualized as a set of ecological states (Penaluna et al. 2016, Reeves et al. 1995, Rieman et al. 2015), each with particular abiotic and biotic conditions, functions, and processes at any given time. The number and variety of ecological states in a domain (i.e., the range of conditions or range of natural variability for an ecosystem) is in constant flux in response to changes in local conditions, stochastic processes, legacies of past disturbance, and time since past disturbance (Beechie et al. 2010; Benda




Figure 7-11A—Examples of the range of conditions that aquatic ecosystems experience: (1) a stream 90 to 100 years after the last disturbance, (2) 160 to 180 years, and (3) more than 330 years (Reeves et al. 1995) (see table 5 for specific details).

et al. 1998; Liss et al. 2006; Miller et al. 2003; Reeves et al. 1995; Resh et al. 1988; Rieman et al. 2006, 2015; Wondzell et al. 2007). Examples of the variation that aquatic ecosystems can experience through time are shown for the central Oregon Coast Range (Reeves et al. 1995) (fig. 7-11A and table 7-5), and eastern Oregon (Wondzell et al. 2007) (fig. 7-11B). Larger streams and rivers in the lower portion of the network are less variable through time; those in the upper and middle portions are more dynamic (Naiman et al. 1992). Because of the variation in the size and asynchronous nature of disturbance events (Allen et al. 1982, Malard et al. 2002, Schindler et al. 2010, Wiens 2002), conditions will vary over time among watersheds, resulting in a mosaic of

biophysical conditions across the landscape. Unmanaged and minimally disturbed aquatic systems may actually exhibit a wider range of conditions than more heavily managed systems (Lisle 2002, Lisle et al. 2007).

A contrasting view holds that aquatic ecosystems tend to be in an equilibrium or steady state, and when disturbed, they are expected to return to predisturbance conditions relatively quickly (Resh et al. 1988, Swanson et al. 1988). Biological (Vannote et al. 1980) and physical conditions (Rosgen 1994) are presumed to be relatively constant through time and to be “good” (barring human interference) in all systems at the same time. Conditions in aquatic systems with little or no human influence and natural




Figure 7-11B—Examples of the range of conditions that aquatic ecosystems in eastern Oregon can experience through time (Wondzell et al. 2007).

Table 7-5—Features of streams from the Oregon Coast Range used in figure 7-11A

Feature/stream	Harvey Creek (1)	Franklin Creek (2)	Skate Creek (3)
Time since disturbance (years)	90–100	160–180	More than 330
Number of pieces of wood/100 m	7.9	12.3	23.5
Mean depth of pools (m)	0.9	0.35	0.1
Dominant substrate	Gravel	Gravel	Bedrock
Percent of fish assemblage juvenile coho salmon	98.0	85.0	100.0
Percent of fish assemblage juvenile steelhead	1.0	12.5	0
Percent of fish assemblage juvenile cutthroat trout	1.0	2.5	0

Source: Reeves et al. 1995.

disturbance, particularly those associated with late-successional and old-growth forest, are assumed to have the most favorable conditions for fish (Fox and Bolton 2007, Murphy and Koski 1989, Pollock and Beechie 2014, Pollock et al. 2012) and other aquatic organisms, and are most frequently used as references against which the condition of managed

streams (e.g., Index of Biotic Integrity) (Karr and Chu 1998) and effects of management actions can be assessed. Systems experiencing disturbances, such as wildfire or floods, are often immediately “restored” by attempting to reduce or eliminate erosional processes. For example, fences were placed in headwater streams following a

wildfire in Colorado to reduce the potential for erosion and debris flows (Chin et al. 2014). However, reaches lower in the stream network downcut, creating other concerns. Although this static ecosystem view is being questioned in the general ecological literature (Hiers et al. 2016, Jackson et al. 2009, Montgomery 1999, White and Jentsch 2001, Wohl et al. 2014), it is still being used to guide management and assess effects on aquatic ecosystems, and persists in environmental laws and policies developed in the 1970s, such as the Clean Water Act (Craig 2010).

Resilience is the capacity of an ecosystem to absorb change and remain within the ecosystem state and domain in the face of natural disturbances and human stressors (Desjardins et al. 2015). As ecosystems undergo larger shifts from human stressors, the ecosystem can be redefined, with a completely different set of characteristics and a compromised or altered range of conditions (Bisson et al. 2009, Reeves et al. 1995). Some ecosystem components may persist through this transition, whereas others may be new components arising from human or climatic alterations, including the development of novel states that may result in the loss of selected ecosystem services and conditions for at least some native species (Penaluna et al. 2016).

The physical aspect of these dynamics is understood conceptually (see review in Buffington 2012), but few mechanistic models currently exist to help us understand the potential effects of management on dynamic ecosystems (but see Wondzell et al. 2007), especially under climate change. As a result, consideration of dynamics remains largely conceptual, and holistic models of basin function (i.e., watershed analyses) are generally lacking, limiting the development of process-based applications of river management and restoration (Beechie et al. 2010). Also, there is also a tendency to focus on mean or median conditions while overlooking temporal variability as “noise” and losing sight of the considerable inherent variability that characterizes riparian and aquatic ecosystems (Fausch et al. 2002, Montgomery 1999), which is ecologically critical (Hiers et al. 2016). Accounting for this variability and for nonstationarity of fluvial processes is central to assessing potential effects of climate change on riverine ecosystems (Buffington 2012, Miller et al. 2003, Montgomery 1999).

However, being able to incorporate this variability into restoration and mitigation actions may be limited by social concerns (Kondolf et al. 2006) (see chapter 12).

Consideration of large spatial and temporal scales is critical to the development of management and conservation strategies for ecosystems (Dale et al. 2000, Holling and Meffe 1996), including a range of conditions for aquatic ecosystems (Fausch 2010, Fausch et al. 2002, IMST 1999, Liss et al. 2006, NRC 1996). This shift requires moving from the current focus on relatively small spatial scales, with little or no consideration of the relevance of time, to a focus that considers large spatial scales, specifically ecosystems and landscapes, over relatively long periods (tens to hundreds of years) (Bisson et al. 2009, Naiman and Latterell 2005, Poff et al. 1997, Reeves et al. 1995). An example of the importance of relations between scales can be seen in the “portfolio effect” of the behavior of populations of sockeye salmon (*O. nerka*) in Bristol Bay, Alaska, identified by Schindler et al. (2010). This study found large variation in the number of fish in any local population over time. However, the variation among the many populations was asynchronous—not all were high or low at the same time. As a result, the total number of fish was relatively constant at the landscape scale, a pattern similar to the amount of old growth historically found in the Oregon Coast Range (Wimberly et al. 2000). This pattern appears to be disrupted in heavily managed systems (Moore et al. 2010).

Both the NWFP and new Forest Service planning rule (USDA FS 2012a) require managers to consider large spatial scales in designing, implementing, and evaluating management actions. The new planning rule also emphasizes ecosystem goals based on ecological integrity. This can be daunting given the lack of scientifically sound examples of how to design and implement forest management at large temporal and spatial scales (North and Keeton 2008, Reeves and Duncan 2009, Thompson et al. 2009) and the lack of adequate tools and guidance. Shifting the management focus to the landscape level and longer time intervals requires recognition of the principles of hierarchy theory and the relation among levels of organization to increase the potential for success of future riparian policies and practices (Fausch 2010, Fausch et al. 2002).

Regulators may recognize the need to apply policies and regulations across broad areas, but may be constrained by the regulatory framework in which they are operating, and generally default to single standards that are applied across broad areas (e.g., National Marine Fisheries Service's matrix of pathway and indicators) (NMFS 1999). This premise is inappropriate for addressing complex, multifaceted landscapes, however (Allen and Starr 1982; O'Neill et al. 1986, 1989); instead, it is important to recognize that a multiwatershed landscape operates differently through time than does a single watershed, and that smaller spatial scales tend to be more variable over time than larger scales (Benda et al. 1998, Wimberly et al. 2000). Increasing levels of aggregation, especially as spatial scales increase, may obscure important system processes (Clark and Avery 1976) and may result in unrealistic expectations for ecosystems and contribute to the contention that often surrounds large-scale management proposals (Allan and Curtis 2005, O'Neill et al. 1986, Shindler et al. 2002). Also, the failure to recognize the different levels of ecological organization and the potential response of each to component parts of disturbance and management may incur unintended economic and social costs, such as repeated investment in ineffective restoration and management strategies (Caraher et al. 1999, Dale et al. 2000).

The emerging consideration of ecosystem dynamics and large spatial and temporal scales has implications for approaches to restoration of aquatic-riparian ecosystems. Many restoration efforts have focused mainly on improving habitat attributes, primarily wood placement, and to a lesser degree on shade improvement for water temperature. These efforts too often aim to bring "stability" to degraded systems, and are viewed as the final phase of restoration (see Palmer et al. 2014). The dynamic approach, not yet broadly practiced, focuses on restoring ecological processes (Beechie et al. 2009, 2010; Bernhardt and Palmer 2011), including periodic inputs or reoccurrences of these important habitat attributes. This requires a shift from reliance on striving only to develop a particular condition (e.g., number of pieces of wood per unit length) or channel classification (e.g., Rosgen 1994) to a quantitative approach based on ecological processes, theory, empirical field methods, and limited modeling (Kline and Cahoon 2010, Wohl et al. 2005).

Some researchers have pointed out that although restoration of ecological processes, such as flow, water temperature, habitat complexity, and connectivity, is a critical consideration in restoring many streams, it may not be sufficient for degraded channels, and can even worsen the ecological condition of the stream (Louhi et al. 2011, Tullos et al. 2009). For example, in restoring floodplain overflow potential, if riparian trees are removed from a previously closed-canopy stream, the underlying energy regime may change from one based on allochthonous resources to one driven by primary production. This may shift the stream farther from the desired ecological state and often toward algae-dominated streambeds and higher temperatures (Robinson 2012, Sudduth et al. 2011). Similarly, if the hydrologic regime is restored but there is no nearby source of invertebrate colonists, then the instream communities will remain altered (Sundermann et al. 2011). Finally, an overreliance on an in-channel focus (small-scale) may not address the stressor(s) that most limit recovery of the aquatic ecosystem; quite often this factor is water quality, and thus ecological recovery will not occur until the stressor is addressed (Beechie et al. 2010, Kail et al. 2012, Selvakumar et al. 2010). Examples of process-focused restoration are presented below in the section on climate change.

In addition to considering spatial complexity, temporal dynamics are particularly important to understand because many key ecological processes such as canopy closure, tree-fall, and fuel loading are related to the age of trees in riparian areas as well as time since disturbance. Temporal dynamics can be examined using models, but long-term studies and monitoring are needed to understand how systems respond over time (Hassan et al. 2005). One strategy that may be appropriate is to design monitoring to focus more on changes following major disturbances rather than focusing simply on short-term trends.

The other challenge posed by a dynamic perspective of aquatic ecosystems is the consideration of large spatial scales. Restoration efforts are generally performed at small spatial scales, with only a relatively small percentage of any watershed actually treated (Ogston et al. 2014, Roni et al. 2010). Roni et al. (2010) estimated that a minimum of 20 percent of the habitat of a given species in a watershed should

be restored to detect a 25 percent increase in smolt (salmon or steelhead) numbers, the minimum detection level for most monitoring programs. They found that floodplain restoration yielded greater increases than in-channel restoration. However, because of the large variability in numbers for most populations (Bisson et al. 1997, Schindler et al. 2010), Roni et al. (2010) suggested that 100 percent of the habitat should be restored to have a significant ecological impact.

Non-Fish-Bearing Streams

The ecological importance of headwater streams, which generally make up 70 percent or more of the stream network (Downing et al. 2012, Gomi et al. 2002), was not well known or understood at the time the ACS was developed, but it is now better established in the scientific literature (Leigh et al. 2016, Richardson and Danehy 2007). Headwaters are sources of sediment (Benda and Dunne 1997a, 1997b; May and Lee 2004; Zimmerman and Church 2001; see review by MacDonald and Coe 2007) and wood (Bigelow et al. 2007; May and Gresswell 2003, 2004; Reeves et al. 2003) for fish-bearing streams; provide habitat (Kelsey and West 1998, Olson et al. 2007) (see chapter 6) and movement corridors (Olson and Burnett 2009, Olson and Kluber 2014) for several species of native amphibians and macroinvertebrates (Alexander et al. 2011, Meyer et al. 2007), including recently discovered species (Dieterich and Anderson 2000); and may be important sources of food for fish (Kiffney et al. 2000, Wipfli and Baxter 2010, Wipfli and Gregovich 2002, Wipfli et al. 2007; also see reviews by MacDonald and Coe 2007 and Clarke et al. 2008). Wood jams in small streams are important sites of carbon storage (Beckman and Wohl 2014), and these streams export large amounts of carbon; one-third is emitted to the atmosphere and the remainder transported downstream (Argerich et al. 2016).

Tributary junctions of headwater streams with larger channels are important nodes for regulating material flows (Benda et al. 2004, Gomi et al. 2002, Montgomery et al. 2003) and cold water (Ebersole et al. 2015) in a watershed, and are the locations where site-scale effects from management activities are often observed (Richardson and Béraud 2014). These locations have unique hydrologic, geomorphic,

and biological attributes and differ in the types and amount of materials delivered to the channel, making them sites of high biodiversity (Benda et al. 2004, Danehy et al. 2012)

Headwater streams are among the most dynamic portions of aquatic ecosystems (Benda et al. 2005, Hassan et al. 2005, MacDonald and Coe 2007, Naiman et al. 1992). Headwater habitats may range from simple to complex, depending on the amount of time since disturbance (such as landslides and debris flows). Following evacuation by a debris flow, headwater depressions and channels fill with material from the surrounding hillslopes, including large wood that falls into these channels, forming obstructions behind which sediments and wood accumulate (Benda and Cundy 1990, May and Gresswell 2004), and then empty again with the next landslide or debris flow (fig. 7-12). As a result, headwater streams are likely to exhibit a range of conditions across the landscape at any point in time.

This cycle of filling and emptying results in a punctuated movement of sediment and wood to larger, fish-bearing streams (Benda et al. 1998, Naiman et al. 1992), contributing to the long-term productivity of many aquatic ecosystems (Benda et al. 2003, Hogan et al. 1998, Reeves et al. 1995). A common consequence of past clearcutting is an absence of down wood to replenish the refilling process. This lack of wood may result in a chronic movement of sediment to larger channels, which could lead to both non-fish-bearing and fish-bearing channels developing characteristics different from those that occurred before forest management. Such conditions could be outside the range of variability to which native biota are adapted (Beschta et al. 2004), limiting the effectiveness of conservation and recovery programs.

Wood enters streams via chronic and episodic processes (Bisson et al. 1987). Chronic processes, such as tree mortality and bank undercutting (Bilby and Bisson 1998, Murphy and Koski 1989), generally introduce single trees or a relatively small number of trees at frequent intervals. Wood from headwater streams, which originates from within 131 ft (40 m) of the channel (May and Gresswell 2003), is delivered to fish-bearing streams by large, infrequent events, such as windthrow (Harmon et al. 1986), wildfire (Agee 1993), severe floods, landslides, and debris flows (Benda et al. 2003; May and Gresswell 2003, 2004; Reeves




Figure 7-12—Conceptual illustration of the changes in channel morphology based on the time since the previous debris flow (May and Gresswell 2004).

et al. 2003). Geomorphic features of a watershed influence the potential contribution of upslope wood sources. Steeper, more highly dissected watersheds will likely have a greater proportion of wood coming from upslope sources than will watersheds with lower stream densities and gradients. Also, there is wide variation in the potential of headwater streams to deliver sediment and wood to fish-bearing streams, depending on channel steepness and angle of entry along the run-out track, among other factors (Benda and Dunne 1997a, 1997b; Brayshaw and Hassan 2009; Burnett and Miller 2007; May 2007). Culverts and other stream crossings can also impede wood movement from smaller to larger streams (Trombulak and Frissell 2000).

The presence of large wood from headwater streams influences the behavior of landslides and debris flows and the response of the channel to such events. Large wood in debris flows and landslides influences the run-out length of these disturbance events (Lancaster et al. 2003). Debris flows without large wood move faster and farther than those with wood, and they are less likely to stop high in the stream network. A debris flow without wood is likely to be a concentrated slurry of sediments of various sizes that can

move at relatively high speeds over long distances, scouring substrate and wood from the affected channels. These types of debris flows are more likely to negatively affect fish-bearing channels, as compared to the potentially favorable effects that result from the presence of wood. Woodless debris flows can further delay or impede the development of favorable conditions for fish and other aquatic organisms. In contrast, those containing wood can help store sediments (Bunn and Montgomery 2004) and build terraces that can persist for extended periods (Lancaster and Casebeer 2007, May and Lee 2004).

Intermittent streams, which can make up half the total length of the stream network (Datry et al. 2014), connected to larger fish-bearing streams can provide important seasonal habitats for spawning and rearing by fish (Boughton et al. 2009, Wigington et al. 2006). In the Oregon Coast Range, growth and survival of juvenile coho salmon was higher in intermittent streams than the perennial mainstem (Ebersole et al. 2006, 2009; Hance et al. 2016). Identification, protection, and restoration of these streams is important to the success of conservation efforts for native fish across the NWFP area.

A rich non-fish community inhabits headwater streams throughout the NWFP area. For example, Olson and Weaver (2007) found 3 species of fish and 12 species of amphibians in stream reaches in 12 western Oregon study sites ranging from Mount Hood to Coos Bay. In this study and Olson and Burton (2014), torrent salamanders (*Rhyacotriton* spp.) dominate intermittent streams and appear to be sensitive to thinning in narrow riparian-management areas; NWFP riparian reserves appear to be benefiting retention of this aquatic-dependent community in abundant small streams in the region. Nevertheless, two torrent salamander species are currently petitioned for listing under the ESA; both have ranges that include significant tracts of nonfederal lands.

Continuing and Emerging
Topics of Concern

Water

Federal lands are important sources of fresh water for human consumption, recreation, agriculture, and environmental needs in the United States. These lands produce an estimated 24.2 percent of the Nation’s water supply, 18 percent and 1.5 percent from Forest Service and BLM lands, respectively (Brown et al. 2008). In the West,⁵ federal lands contribute 66 percent of the mean annual water supply, 51 percent of which comes from Forest Service lands and 5.4 percent from BLM lands (Brown et al. 2008). Management strives to maintain the quality and quantity of this water. The extent of the contribution of federal lands to regional

water supplies was not well quantified at the time the NWFP and ACS were developed.

The contribution of water from federal lands specifically in the NWFP area is also important; however, exact estimates are not currently available and were beyond the scope of this review. At the state level, the majority of water in the three NWFP states (California, Oregon, and Washington) originates from federal lands (table 7-6), with the bulk coming from Forest Service lands. Within the NWFP area, the amount of water flowing from federal lands varies among national forests and watersheds. Some forests, such as the Deschutes and Willamette National Forests, make relatively large contributions, 40 percent or more, to the flow of rivers whose watersheds they include (figs. 7-13A and 7-13B, respectively). Contributions from other forests are smaller (less than 20 percent) (fig. 7-13C) but nonetheless important. See “Climate Change” below for potential future issues pertaining to water supply and stream temperatures.

Roads

Roads provide necessary motorized access for forest management, recreation, and other beneficial purposes (Gucinski et al. 2001), but they can also have detrimental effects on native biodiversity and ecosystem function. The focus of the NWFP and ACS has been to address the negative effects of roads on aquatic ecosystems through a broad program of road decommissioning and upgrading, including remediation of chronic sedimentation and barriers to aquatic organism movement. Several syntheses describe the types, causes, and effects of road networks on streams, and meta-analyses concerning the ecological effects of

⁵ The West is defined here as including the states of Arizona, California, Colorado, Idaho, Montana, New Mexico, Nevada, Oregon, Utah, Washington, and Wyoming.

Table 7-6—Contribution of federal lands and agencies to the total mean annual water supply of states in the Northwest Forest Plan area (percentage of mean annual water supply)

State	All federal lands	Forest Service	Bureau of Land Management	Other
Percent				
California	61.1	46.6	5.5	9.0
Oregon	55.3	44.0	9.4	2.0
Washington	60.2	41.5	0	18.7

Source: Brown et al. 2008.




Figure 7-13—Contribution of selected national forests in the Northwest Forest Plan area to the mean annual flow of adjacent watersheds: (A) Deschutes National Forest, central Oregon Cascades; (B) Willamette National Forest, western Oregon Cascades; (C) Siuslaw National Forest, Oregon Coast Range; and (D) Six Rivers National Forest, coastal northern California. (<https://www.fs.fed.us/rmrs/projects/national-forest-contributions-streamflow>).

roads in general and specifically the delivery of sediment from mountain roads with low maintenance standards have been published (Croke and Hairsine 2006, Forman and Alexander 1998, Jones et al. 2000, Trombulak and Frissell 2000). Reducing the effect of roads and associated infrastructure remains a challenge for federal management agencies and others.

The vulnerability of roads to hydrologic changes and the associated effects on aquatic and riparian ecosystems differ based on topography, geology, slope stability, design, location, and use. Roads can affect streams directly by:

1. Accelerating erosion and increasing sediment loading (Allan 2004, Daigle 2010, MacDonald and Coe 2008, Suttle et al. 2004).
2. Imposing barriers to the migration of aquatic organisms, including access to floodplains and off-channel habitats (Clarkin et al. 2005, Daigle 2010, Gibson et al. 2005, Sagar 2004, Trombulak and Frissell 2000).
3. Increasing stream temperatures (Wenger et al. 2011).
4. Causing changes in channel morphology (Daigle 2010, Hassan et al. 2005).
5. Introducing exotic species (Daigle 2010, McKinney 2001).
6. Increasing harvest and poaching pressure (Lee et al. 1997, Trombulak and Frissell 2000).
7. Changing hillslope hydrology and resulting peak flows (Daigle 2010, Jones and Grant 2001).

Roads penetrating remote and otherwise intact forested landscapes can have particularly significant effects on aquatic ecosystems (Forman 2003, Havlick 2002, Trombulak and Frissell 2000). The ecological consequences of these effects are shown in table 7-7.

The effects of roads differ widely depending on local features (Al-Chokhachy et al. 2016). Recently developed techniques, such as the Geomorphic Roads Analysis and Inventory Package (Black et al. 2012), can be employed to identify priority locations of sources of sediment, including culvert failures, landslides, and gullies. A modified version of this technique has been incorporated into NetMap (Benda et al. 2007) to reduce the amount of field time

needed to assess roads. Evaluating the effectiveness of new analytical approaches and focusing on treating limited lengths of roads could be a research priority.

A significant number of watershed-improvement actions implemented under the NWFP and other large-scale forest planning efforts involve decommissioning roads that have a high probability of contributing to landslides, and that are not regarded as essential to meeting local forest objectives, as well as removing road-related impediments to upstream and downstream movements of aquatic organisms (Switalski et al. 2004). The watershed-analysis component of the ACS identified forest roads where (1) drainage systems hastened runoff from storms and promoted sedimentation of streams, (2) unstable fill materials concentrated water and increased the risk of landslides, and (3) the roadbed encroached on riparian zones (Kershner 1997). Since NWFP implementation, the Aquatic and Riparian Effectiveness Monitoring Program estimated that 6.7 percent of the road network has been removed or closed (5,390 out of 80,750 mi total [8674 of 129 954 km]) in the NWFP area.⁶ Additionally, 10 percent of the road crossings that impeded the movements of aquatic and riparian organisms (209 of 2,114) have been made passable on Forest Service Region 6 lands in the NWFP area since NWFP implementation.⁷

Though restoration of fish passage is often listed as a top priority for stream restoration in the Pacific Northwest (Roni et al. 2002, USGAO 2001), recent work has contributed much to our understanding of just how complex this issue is in practice (McKay et al. 2016). Advances have been made in culvert inventory and assessment (Clarkin et al. 2005), ecosystem-based restoration approaches (USDA FS 2008), and effectiveness monitoring (Heredia et al. 2016, Hoffmann et al. 2012). Until recently, however, the ecological benefit of these efforts has been difficult to quantify beyond the level of individual projects. A new

⁶ Miller, S. 2016. Personal communication. National riparian program lead, U.S. Department of the Interior, Bureau of Land Management, 1849 C Street NW, Washington, DC 20240.

⁷ Capurso, J. 2017. Personal communication. Regional fish and aquatic program manager, U.S. Forest Service, 333 SW First Ave., Portland, OR 97204, jcapurso@fs.fed.us.

Table 7-7—Summary of effects of road on aquatic ecosystems and associated biota

Ecological effect	Habitat loss/degradation	Habitat fragmentation	Direct mortality
Low population density	✓	✓	✓
Low population reproductive rates	✓		✓
Area occupied restricted	✓	✓	✓
Decreased habitat connectivity	✓	✓	
Overharvest			✓
Changes in water quality	✓		✓
Changes in hydrologic functions	✓	✓	✓
Change in wood and sediment recruitment	✓	✓	✓

Source: Modified from Robinson et al. 2010 and Daigle 2010.

study evaluating the effectiveness of passage restoration at the level of an entire forest (the Siuslaw National Forest) (Chelgren and Dunham 2015) found that individual culvert replacements successfully increased the probability of upstream access for all fishes in the study area. Results of this work also showed that the net benefit of culvert replacements was fairly modest across the extent of the forest when expressed in terms of gains in kilometers of stream occupied or increases in fish numbers resulting from restoration. The authors hastened to add that some limitations of the study design could have influenced these findings (Chelgren and Dunham 2015), but results of this study nonetheless point to the value of programmatic (vs. project-only) evaluations of culvert restoration. This echoes more general recommendations for following the cycle of adaptive management on national forests (Marcot et al. 2012) and the recommended scales for managing watersheds (Fausch et al. 2002, Neeson et al. 2015).

Much of the guidance for fish-passage restoration on federal lands in the Pacific Northwest was issued by an assessment in 2001 (USGAO 2001). This assessment highlighted the need for larger scale assessments, as noted above, as well as the economic challenges associated with passage restoration, which are only beginning to be addressed. For example, a followup to the Chelgren and Dunham (2015) study by Reagan (2015) evaluated costs and benefits of remaining culvert replacement opportunities on the Siuslaw National Forest in relation to multiple objectives, including

benefits to fish (estimated from Chelgren and Dunham 2015), maintenance of transportation networks, and the probability of culvert failures based on culvert size and influences of floods and major erosional events in streams. The Reagan (2015) analysis explicitly quantified economic costs and benefits of restoration in relation to these objectives and their relative assumed values. This work (along with others in the region, e.g., Chelgren and Dunham 2015) has demonstrated the value of a proactive, economic analysis of multiple objectives to identify priorities for restoration investments in a programmatic context. These new tools, if applied, can more completely address standing recommendations to land management agencies in the Pacific Northwest (e.g., USGAO 2001) to more efficiently invest limited resources to benefit fisheries and other management objectives through culvert replacements.

Because road access management must take into account social, economic, and environmental objectives (Daigle 2010), the decisionmaking process for dealing with roads is complex. A decision matrix for identifying actions is shown in figure 7-14. In many cases, limited funds or socioeconomic issues may preclude closing or removing roads identified as high priority for treatment on the basis of their effects on riparian ecosystems. Also, a road network may be needed to effectively implement landscape-scale restoration projects that might involve widespread thinning and prescribed fire (Franklin and Johnson 2012), and for fire management, fuel reduction, and fire control. Studies

Road importance ↑ High ↓ Low	Preventive management: • perform regular maintenance • monitor for signs of presence effects	Mitigate presence effects: • maintain fish passage • prevent erosion • soften edge effects
	Decommission road: • abandon	Decommission road: • rip or obliterate
	Low ← Ecological impact → High	

Figure 7-14—A decision matrix for identifying potential options for managing roads. (Modified from Robinson et al. 2010).

in Redwood National Park suggest that removal of logging roads can yield carbon-storage dividends, in particular by preventing soil erosion (van Mantgem et al. 2013). The vast road system on private and state lands that abut federal lands also needs to be considered in road assessments. Understanding how to balance fire management, recreation, and other needs against potential negative aspects of roads will require a concerted cooperative effort of managers and physical, biological, and social scientists, other organizations, and the public. (See next section for additional discussion of roads.)

Climate Change

Since 1994, our knowledge of climate change in the NWFP area has greatly improved, just as dealing with climate change has become an important aspect of environmental planning in the Forest Service and BLM. Many advances have come from models that forecast trends in temperature, precipitation, and snowpack, and associations of these trends with the habitat conditions for various species. Although there is general agreement about the direction of trends in many meteorological parameters, the rates and amounts of change at specific locations in the NWFP area differ among models (Climate Impacts Group 2009; see also chapter 2). Further, other climate-related changes such as increases in forest insect and disease outbreaks and uncharacteristically severe wildfires may

accentuate the undesirable effects of meteorological and hydrological trends, resulting in threats to the integrity of both terrestrial and aquatic ecosystems. Although developing proactive measures that would ameliorate undesirable effects of climate change on forest resources was not a centerpiece of the NWFP, one of the Plan's main objectives was to restore forest ecosystems that were resilient in the face of natural and anthropogenic disturbances. The question is: how well does the NWFP address climate-related threats to native fishes and other aquatic biota as they are currently perceived? (See chapter 2 for further details.)

In this section, we focus on a review of recent advances in our knowledge of the likely effects of climate change on native fishes of the NWFP area. We examine climate-change effects on fish life cycles, with a principal focus on anadromous salmonids, a group of species that has received the most scientific attention, as well as significant conservation effort (table 7-8) (see additional discussion in app. 2). Watershed improvements undertaken through the NWFP are related to potential climate effects on fish life cycles, and to the capacity of populations to adapt and persist through time. Finally, we discuss the role that federally managed forests in the NWFP area play in conserving native fishes in a changing climate, when viewed in a broader matrix of different land ownerships and other landscape-scale uses.

Table 7-8—Species of Pacific salmonids considered in this section and their typical freshwater and marine residence times

Species		Residence time	
		Freshwater	Marine
Pink salmon	<i>Oncorhynchus gorbusha</i>	Less than 30 days	2 years
Chum salmon	<i>O. keta</i>	Less than 30 days	2 to 5 years
Sockeye salmon	<i>O. nerka</i>	Few months to 2 years	2 to 5 years
Coho salmon	<i>O. kisutch</i>	1 to 2 years	1.5 years
Chinook salmon	<i>O. tshawytscha</i>	Few months to 1 year	2 to 6 years
Steelhead	<i>O. mykiss</i>	1 to 3 years	2 to 4 years
Coastal cutthroat trout	<i>O. clarkii clarkii</i>	2 to 4 years	Short forays into nearshore environment

Climate change in the Northwest Forest Plan area—

Projected changes in climate are usually derived from models based on historical data coupled with scenarios incorporating alternative assumptions about future greenhouse gas emissions. These assumptions range from high global rates of economic development and human population growth to conservative industrial and population-growth rates and widespread implementation of “clean” technologies. Model outcomes are often displayed as incremental changes in an environmental parameter of interest such as air temperature, sea level, or precipitation over a fixed period. Projected changes in climate under different scenarios are plotted to provide a range of outcomes at a given point in time, with scenarios incorporating intermediate assumptions about future greenhouse gas emissions generally believed to represent the most realistic expectations.

Air and water temperatures—

Virtually all climate models forecast a gradual rise in air temperature by the end of this century. Recent changes in climate appear to be happening more rapidly than in at least the past 1,000 years (IPCC 2007), and have included a global average warming of 1.4 °F (0.8 °C) during the past 120 years. According to the IPCC (Intergovernmental Panel on Climate Change) (IPCC 2014), most general circulation models predict that 2 to 7 times more warming will occur by early in the next century, with projected increases in mean global surface temperatures by 2100, ranging from 2.7 to 3.6 °F (1.5 to 2.0 °C) relative to the 1850–1900 time frame, depending on carbon dioxide (CO₂) emission

scenarios (IPCC 2014). The 2014 IPCC synthesis report (IPCC 2014: 10) states:

It is **virtually certain** that there will be more frequent hot and fewer cold temperature extremes over most land areas on daily and seasonal timescales, as global mean surface temperature increases. It is **very likely** that heat waves will occur with a higher frequency and longer duration. Occasional cold winter extremes will continue to occur. [emphasis theirs]

The finding that climate change will include both gradual long-term temperature trends as well as increases in the frequency and duration of extreme events has important implications for aquatic ecosystems in the NWFP area.

Air-temperature changes in forests of the NWFP area are predicted to be generally consistent with global climate models, although somewhat more variable, with forecast increases ranging from 1 to 6.3 °F (0.5 to 3.5 °C) in the remainder of this century, depending on the greenhouse gas emission scenario used in the model and on forest location (Latta et al. 2010). Overall, these authors noted that relative temperature increases were more apparent at higher elevations than at lower elevations, and that proximity to the Pacific Ocean moderated the rate of change. Mote and Salathé (2010) examined a broad suite of IPCC climate models and found that, by the 2080s, average air temperatures in the Pacific Northwest were predicted to increase 2.9 °F (1.6 °C) under the coolest scenario and

by 10.3 °F (5.7 °C) under the warmest scenario. In most models, the greatest absolute temperature increases were projected for summer months, although warming was also forecast in other seasons. Sea-surface temperatures showed less warming over the same period than those modeled over land.

Similar to air temperatures, water temperatures are expected to rise in much of the NWFP area as a result of climate change (Isaak et al. 2011 [NorWeST model]). Modeled water temperatures were developed primarily from models of the relation between air and water temperatures, and are projected to be stressful to lethal for many native salmonids (e.g., Isaak et al. 2012, Wade et al. 2013) (see app. 2 for more details). More recent studies suggest that the extent of temperature change may not be as great as originally projected, particularly at higher elevations (Isaak et al. 2016). However, other researchers (Arismendi et al. 2014) have questioned the ability to project future water temperature from past relations between air and water temperatures. In addition, Arismendi et al. (2013a) found that recent trends in water temperature have been more variable than those reported by Isaak et al. (2012)—using empirical records, they found that water temperatures increased in some systems and decreased in others. Also, Leach et al. (2016) also found variability in water temperature in a headwater stream of the Oregon Coast Range that was not captured by the NorWeST model (Isaak et al. 2010), but noted that the model was not designed to track such small-scale effects. Although there is some uncertainty about the extent of temperature changes that streams in the NWFP area will experience, it is clear that dealing with water temperatures will be a major challenge for managers.

Potential patterns of changes in water temperature are highly variable when examined at the local scale (Leach et al. 2016, Reeves et al. 2016b, Turschwell et al. 2016) (fig. 7-15). This variability is a result of local conditions such as stream orientation, topographic shading, and elevation, and strongly influences physical and biological attributes and resultant ecosystem integrity (Gomi et al. 2002, Thorp et al. 2006). Understanding this variability will be crucial to developing effective restoration and mitigation programs and prioritizing specifically where

to target efforts. Watershed analysis tools such as Net-Map (Benda et al. 2007) can help identify areas that can provide thermal refugia and areas in which riparian restoration efforts (fig. 7-16) could help reduce water temperatures to levels less stressful or even optimal for native fish, despite climate change (Justice et al. 2017, Lawrence et al. 2014, Ruesh et al. 2012).

Hydrology—

Predicted future changes in streamflow on national forests in the Pacific Northwest are fundamentally tied to changes in the region's climate. Predicted changes in annual precipitation are much less certain, and most models project that future precipitation will remain approximately the same as it has been for the past 50 years (Salathé et al. 2007). Most predictions of changing streamflows for the Pacific Northwest therefore focus primarily on the effects of changes in temperature. Seasonal changes in precipitation are showing up in the data (Safeeq et al. 2013) but are difficult to resolve regionally, and consequently are not as well understood.

A key factor affecting both high and low streamflows in the future will be the fate of snow and the seasonal snowpack. Snowpack dynamics are important to understanding streamflow regimes because snow represents a dominant form of storage on the landscape. When precipitation falls as snow, it is not available for runoff or groundwater recharge until it melts. Similarly, the rate and timing of snowmelt are first-order controls on both peak and low streamflows, as discussed below.

A particularly crucial dimension of snowpack dynamics is the geographic location of the rain-snow transition on the landscape. This transition is controlled by elevation and determines how much of the winter precipitation falls as rain versus snow. Although often visualized as a fixed elevation, this transition is better seen as a stacked sequence of elevationally controlled zones or ranges with imprecise and regionally varying boundaries (Klos et al. 2014, Nolin and Daly 2006). In general, for any area, there is an elevation below which virtually all winter precipitation falls as rain and above which it falls as snow. Elevations in between are defined as the transitional snow zone (TSZ) that receives both rain and snow; snow and the snowpack usually will not persist all winter.





Figure 7-15—(A) Current and (B) projected (2040) summer water temperatures (°C) in the study basins in the Treaty of Olympia area (Reeves et al. 2016b).



The widely varying elevational gradients in the Pacific Northwest in general, and on national forest lands in particular, therefore impose considerable variability in the response of the landscape to changing climate. Depending on the proportion of the landscape that occupies each of these zones, a warming climate, hence a rising snow line, may transition the landscape from a zone dominated by seasonal snow accumulation and melt (snow zone) to one that receives a mixture of rain and snow (and rain-on-snow)—the TSZ.

Or it may push the landscape out of the TSZ and into the rain zone (Klos et al. 2014, Luce et al. 2014a)

The effects of a changing climate are already apparent in the snow data for the Pacific Northwest. As winter and spring temperatures have warmed over the past 50 to 70 years, spring snowpacks have been smaller (Hamlet et al. 2005, Mote 2003, Mote et al. 2005) and have melted out earlier (Hamlet and Lettenmaier 2007, Stewart et al. 2005). Moreover, the aforementioned zonal changes are already

occurring in some landscapes as snow zones transition to the TSZ, and the TSZ becomes rain dominated (Tohver et al. 2014). These trends are expected to continue across much of the region (Luce et al. 2014a).

However, snowpack dynamics alone do not determine what future streamflow regimes will look like on national forests in the NWFP area. Recent work has shown that another first-order control is the landscape-scale drainage efficiency: the inherent, geologically mediated efficiency of landscapes in converting recharge (precipitation) into discharge (Safeeq et al. 2013, 2014; Tague and Grant 2009). In essence, the drainage efficiency determines how quickly precipitation, either as rain or snowmelt, becomes streamflow. Although drainage efficiency is “hard-wired” into the landscape on millennial timescales, and thus is not changing with climate, it mediates the climate-influenced signals and therefore has to be considered in predicting future streamflow regimes. This is particularly true for low-flow regimes, but influences peak flows as well. Basically, the drainage efficiency of a landscape is determined by the rate at which water moves through the subsurface. In steep landscapes with shallow soils, water rapidly moves laterally through the subsurface via both saturated and unsaturated pathways, drainage efficiency is high, and streams respond quickly to recharge events. In flatter landscapes with deep, permeable, porous, or fractured bedrock, water moves slowly as deep groundwater, drainage efficiency is low, and streams respond slowly to recharge events but may have sustained high base flows.

Effects of climate change on peak flows—Here we broadly consider how both climate and drainage efficiency can shape predictions of future streamflows on national forest lands. We distinguish between effects on peak and low flows, as the mechanisms of streamflow generation are different in each case. Finally, we discuss how these broad predictions can be refined for individual forests, a topic beyond the scope of the current analysis.

There are several hydrologic mechanisms by which climate could increase peak flows in rivers and thus their propensity to flood. More intense or frequent rainstorms are one mechanism, and some research has suggested that a warming atmosphere may result in a more northerly storm track for the North Pacific, potentially resulting in more

intense precipitation (Salathé 2006). However, these results have large uncertainties and are not well represented in most global circulation models. A somewhat better-understood mechanism is the shifting potential for rain-on-snow (ROS) events in the Pacific Northwest as the climate warms. ROS events are known to be a potent flood-producing mechanism in steep mountain landscapes in the Pacific Northwest (Harr 1981, Marks et al. 1998, McCabe et al. 2007). In general, landscape susceptibility to ROS events is determined by climate and topography; the effects of climate warming on ROS are similarly influenced by the same controls; and climate warming may increase, decrease, or not affect the risk, depending on whether snowpacks are cold or warm (i.e., near the freezing point). As summarized by Hamlet and Lettenmaier (2007):

Cold river basins where snow processes dominate the annual hydrologic cycle (<6 °C average in midwinter) typically show reductions in flood risk due to overall reductions in spring snowpack. Relatively warm rain-dominant basins (greater than 5 °C average in midwinter) show little systematic change. Intermediate or transient basins show a wide range of effects depending on competing factors such as the relative role of antecedent snow and contributing basin area during storms that cause flooding. Warmer transient basins along the coast in Washington, Oregon, and California, in particular, tend to show increased flood risk.

A more recent analysis looked at a range of factors influencing peak flows, including ROS in Oregon and Washington, and developed a model of sensitivity to peak-flow increases based on perturbing the temperature in the model using warming scenarios from 2020 to 2080 and the A1B⁸ emissions scenario (Safeeq et al. 2015). The analysis yielded regional sensitivity maps for Oregon and Washington that can be used to characterize the risk on individual national forests and landscapes. They concluded

⁸ This scenario assumes a future world of rapid economic growth and global populations peaking in the mid-21st century, then declining with the rapid introduction of new technology, with a balance between the use of fossil fuels and non-fossil-fuel sources.

that corresponding changes in snowpack dynamics may result in large (more than 30 to 40 percent) increases in peak flows, primarily in the Cascade Range and Olympic Mountains. The North Cascades, in particular, were most vulnerable (fig. 7-17). Lower elevation areas were less likely to be affected but were still vulnerable to larger floods generated from upstream reaches in vulnerable landscapes. These watersheds are also likely more susceptible to warming (Arismendi et al. 2013a, 2013b; Poole and Berman 2001; van Vliet et al. 2011, 2013). Streams at higher elevations should retain flows; with stable, cooler water temperatures, they will be critical cool-water refugia for native fish (Isaak et al. 2012, 2015; Luce et al. 2014b; Lusardi et al. 2016; Wenger et al. 2011).

Effects of climate change on low flows—Snowpack dynamics and drainage efficiency combine to determine the sensitivity of individual landscapes to a warming climate (Safeeq et al. 2013, Tague and Grant 2009). There has been a general trend over the past 50 years for less snow in winter and earlier snowmelt, resulting in reductions of spring, early-summer, and late-summer flows in the Western United States (Leppi et al. 2012, Safeeq et al. 2013), with the lowest flows showing the greatest decreases across the Pacific Northwest (Luce and Holden 2009). Hydrologic models such as the variable infiltration capacity (VIC) model, coupled with downscaled climate simulations, have been used to generate predictions of future low flows across much of the Pacific Northwest (e.g., Hamlet et al. 2013).

However, snowpack changes are not the only factor determining future low flows. Other recent work has shown that the drainage efficiency (slow versus fast) mediates the signal from climate-induced changes in snowpack and snowmelt. Employing a simple exponential-decay model to describe the recession behavior of streams, coupled with a regional-scale estimation of variations in aquifer-drainage characteristics, Safeeq et al. (2014) developed a sensitivity map for changes in summer streamflow across Oregon and Washington. As with the VIC products and peak-flow maps previously described, these maps provide water and landscape managers with a spatially explicit representation of where future changes in

low flows are likely to be most pronounced. For example, these maps show that areas drained by young volcanic rocks with deep, slow groundwater systems, such as the High Cascades, may be particularly vulnerable to declines in summer streamflow, whereas areas with shallow subsurface aquifers and limited potential to store water are less sensitive to changes in low flows. Climate-change effects on summer low flows may be compounded by effects of forest-vegetation conditions. Perry and Jones (2017) found that average daily streamflow in smaller streams in summer in watersheds with 34- to 43-year-old plantations of Douglas-fir was 50 percent lower than streamflow from reference basins with 150- to 500-year-old forests. The change in flows is also likely to be highly variable among watersheds in a given area (fig. 7-18).

Assessing climate change effects on streamflow at the scale of individual national forests—The discussion above highlights how existing tools and models can be used to give technically sound predictions about the magnitude and timing of streamflow changes in specific landscapes. Although not a trivial exercise, any national forest can use the spatially explicit models already developed to make first-order forecasts for changes in streamflow regimes. The products to date cover most but not all forests in the area of the NWFP. Extending results to these unmapped forests (mostly in northern California) would require some extrapolation, but is well within the scope of the existing data. Tools and approaches such as the concept of “hydrologic landscapes” can expedite this process (Patil et al. 2014, Wigington et al. 2013, Winter 2001).

Furthermore, there are several examples to date of individual forests or groups of national forests and other federal and nonfederal landholders that have coordinated efforts to develop detailed assessments of likely hydrologic changes that can serve as models for other forests and regions. Specific examples include the Olympic National Forest (Halofsky et al. 2011), the Quinalt Indian Nation on the Olympic Peninsula (Reeves et al. 2016b), the Blue Mountains Adaptation Partnership (Halofsky and Peterson 2017), and the upcoming report from the South Central Oregon Adaptation Partnership.




Figure 7-17—Sixth-field hydrologic unit-scale average peak-flow sensitivities across all flood magnitudes (Q2, Q10, Q25, Q50, and Q100) under A1B emission scenario for the (A) 2020s, (B) 2040s, and (C) 2080s, in which red is more sensitive and blue is less sensitive. (Modified from Safeeq et al. 2014.)




Figure 7-18—Percentage of reduction in average (A) summer and (B) winter flow levels from current to 2040 in study basins in the Olympic Peninsula area (Reeves et al. 2016b).

Extreme events—Increased frequency of extended, severe droughts and intense winter-storm events (IPCC 2014) will also affect aquatic ecosystems and fish populations in forested landscapes (Ward et al. 2015). The pattern of changes will differ widely within and among watersheds depending on local features, making it difficult to generalize the effects. However, changes in the seasonal timing of annual hydrographs and more frequent extremely low or high flows are very likely to affect native fish populations. Changes in flows that lead to earlier spring runoff and prolonged periods of summer low flows could have important implications for the habitat of (ISAB 2007) and food chains encompassing fish and other aquatic organisms (Power et al. 2008, Wootton et al. 1996) (see discussion in app. 2 for more details). Amphibians that inhabit ephemeral ponds and streams are likely to be especially vulnerable to drought and general climate change effects (Blaustein and Olson 1991, Shoo et al. 2011).

Ocean conditions—

Over the past several decades, the importance of the marine environment to fish that spend part or all of their lives at sea has been recognized as a major factor regulating population abundance. Climate-related changes in the ocean that are potentially important to native fishes in the NWFP area include acidification (Orr et al. 2005), increased sea-surface temperatures (IPCC 2007), changes in wind and current patterns (Rykaczewski and Dunne 2010), and sea-level rise (IPCC 2007). Absorption of anthropogenic CO₂ by the upper ocean decreases pH and carbonate-ion concentrations (Hendriks et al. 2010, Orr et al. 2005), increasing acidity and inhibiting the ability of planktonic organisms to form calcium carbonate, a key component of their exoskeleton. Many of these organisms form the base of the food chain that supports anadromous fishes during the marine phase of their life cycles. The subarctic Pacific Ocean has naturally higher carbon concentrations than most other ocean basins, and the effects of acidification are expected to occur sooner and be more pronounced there (Cooley et al. 2012).

Rising sea-surface temperatures may reduce the amount of preferred thermal habitat for anadromous salmonids in the ocean and potentially limit their marine distribution (Aziz et al. 2011, Welch et al. 1995). As areas with

suitable temperatures decrease or shift northward, Pacific salmon could become concentrated in smaller foraging zones, resulting in increased competition for limited food resources (Grebmeier et al. 2006, Johnson and Schindler 2009, Mantua et al. 2009, Welch et al. 1995). Salmon may be able to partially compensate for these changes by using cooler subsurface waters; however, deeper water may provide reduced food resources, increased competition with other marine species, or greater exposure to predation (Hinke et al. 2005, Myers et al. 1996).

Other potentially important climate-related changes in the marine environment include sea-level rise (IPCC 2007) and altered patterns of coastal upwelling (Wang et al. 2015). The consequences of sea-level rise for nearshore fishes are uncertain and will be strongly influenced by local topography; new habitat could be created in some areas but lost in others (Flitcroft et al. 2013). Saltwater inundation may affect species that sometimes spawn immediately above tidewater (e.g., pink and chum salmon, *Oncorhynchus gorbuscha* and *O. keta*). Both positive and negative effects on estuaries are also possible as new land is inundated, and the degree to which estuarine productivity is altered will be influenced by the extent of human development. Where development of estuary and coastal shorelines is extensive, sea-level rise will likely result in more seawalls, channelization, and other measures to prevent flooding during storm surges (Neumann et al. 2015).

Changes in the patterns of coastal upwelling in the NWFP area could have very significant effects on anadromous fishes as well as other animals that depend on marine food webs. Wind-driven ocean currents regulate the strength of coastal upwelling along the Pacific Coast, where nutrients from deep-ocean waters fuel plankton blooms that are critical to marine food webs that support salmon (Francis and Sibley 1991). Long-term shifts in the timing and intensity of coastal currents and upwelling have accompanied climate change in the eastern Pacific Ocean, with winter and spring storm tracks gradually shifting northward (Salathé 2006) and upwelling along the coast in the NWFP area becoming more erratic and unpredictable (Bylhouwer et al. 2013). Anadromous salmonids are particularly vulnerable to changes in upwelling because survival of

fish in the first few weeks after entering the ocean depends heavily on their ability to feed and grow large enough to avoid predation (Beamish and Boullion 1993, Pearcy 1992, Walters et al. 1978).

The occurrence of interdecadal shifts in sea-surface temperatures and related weather patterns (Pacific Decadal Oscillation—PDO) from cool/wet to warm/dry conditions (Mantua et al. 1997) further complicates the productivity of marine environments along fish migration routes, with more favorable ocean conditions occurring when the NWFP area is in a cool/wet phase than in a warm/dry phase. Wang et al. (2015) used a suite of climate models to forecast upwelling over the remainder of this century and found that, by the year 2100, coastal upwelling will likely start earlier, end later, and be more intense in the northern latitudes (British Columbia, Canada, and southeast Alaska) than in southern latitudes (northern California and Oregon). Wang et al. (2015) also noted that an intensification of upwelling could actually promote plankton productivity, but in extreme cases could also result in large swaths of anoxic conditions developing over broad areas, leading to massive die-offs of marine life where such conditions develop. Taken together, the new information on climate-related PDO cycles and trends in upwelling patterns suggest that the marine environment along the Pacific Coast is becoming more variable spatially and temporally, with northern California and Oregon being somewhat more likely to exhibit unpredictable ocean conditions than more northerly latitudes. For migratory organisms such as anadromous salmonids whose life cycles are adapted to being in the right place at the right time for feeding and reproduction, introducing more variability into the part of their life cycle where most growth occurs is likely to add to population destabilization.

Climate effects on fish life cycles—

Although the extent to which a particular fish population in the NWFP area will be affected by climate change depends to a large degree on changes that occur at the local level, climate-related effects, both favorable and unfavorable, can accumulate across multiple life-history stages. Restricting an understanding of climate influences to a single life-history stage may well underestimate the total effect on the population. Further, because of the wide geographic

distributions of many native fishes and the heterogeneity of aquatic environments in which they reside, climate effects may be expressed differently across the range of a given species. Locally adapted life histories differ over broad landscapes and among different species; even stocks of the same species can exhibit dissimilar responses to similar climate trends (Schindler et al. 2010). A number of papers have investigated the potential effects of climate change on Pacific salmon, but these have primarily been overviews (e.g., Bryant 2009, ISAB 2007) or results of modeled effects on a given life-history stage (e.g., Crozier and Zabel 2006, Rand et al. 2006) and its associated habitat (e.g., O’Neal 2002). A comprehensive review of the effects of climate change on native fishes in the NWFP area across their ranges, including effects accumulated across multiple life-history stages, is lacking.

Understanding the potential consequences of altered future conditions, particularly where the perceived effects may not be lethal, requires consideration of the effects at each life-history stage (Fleming et al. 1997, ISAB 2007, Jonsson and Jonsson 2009). Changes at one life stage can cascade throughout the remaining stages, significantly altering population response. Focusing on anadromous Pacific salmonids, it is possible to examine the overall impacts of climate change by identifying effects at each life-history stage and discussing how those effects might be propagated through succeeding stages. These effects and potential management options are listed in table 7-9. It is also possible to identify attributes of Pacific salmon life cycles that promote their adaptive capacity to climate change, along with options for managers and decisionmakers to enable and enhance those attributes to mitigate potential effects of climate change in the NWFP area.

Other climate-related factors—

Climate warming will lead to an increase in the area burned by wildfires (IPCC 2014) (chapter 2). Wildfire trends in the NWFP area will be complex because the area includes a wide array of forest types, elevations, weather regimes, and forest-management histories (Hessburg and Agee 2003); hence risks of damage to native fish habitats are likely to be highly variable across the region. In addition to altering wildfire frequency and intensity, climate change will also

Table 7-9—Potential effects of climate change on anadromous salmonids of the Pacific Northwest, by life-history stage

Life stage and habitat	Potential effect of climate change	Ecological consequences	Ecological implications	Potential actions
Adults:				
Ocean	Acidification (Hendriks et al. 2010, Orr et al. 2005)	Reduced growth and survival	Lower productivity of freshwater because of reduced amount of marine-derived nutrients and eggs (Bilby et al. 1996, Cederholm et al. 2001, Garner et al. 2009, Gende et al. 2004, Helfield and Naiman 2001, Lang et al. 2006, Schindler et al. 2003)	
	Increased sea-surface temperatures (Aziz et al. 2011, IPCC 2007)	Smaller size at return	Reduced population reproductive capacity (Hankin and McKelvey 1985, Healey and Heard 1984)	
		Change in life-history expression (<i>Oncorhynchus mykiss</i>)	Loss of steelhead life history (migratory) and increase in rainbow trout life history (resident) (Benjamin et al. 2013, Quinn and Myers 2004, Rosenberger et al. 2015, Sloat and Reeves 2014)	Population monitoring with consideration of life-history types
Freshwater	Sea-level rise (IPCC 2007)	Increased estuary habitat	Increased life-history diversity (Bottom et al. 2005)	Population monitoring with consideration of life-history types
		Loss of spawning habitat in areas close to coast	Reduced population productive capacity	
		Increased flooding during surge events	Reduced egg survival	
	Increased water temperature (Isaak et al. 2010)	Increased stress	Reduced survival to spawning grounds (Martins et al. 2011, Rand et al. 2006) and population reproductive capacity (Miller et al. 2011, Pankhurst et al. 1996) Increased susceptibility to disease and parasites (Chiaromonte et al. 2016, Johnson et al. 1996, Miller et al. 2011, Ray et al. 2012)	Monitoring water temperatures in entire stream network to identify and protect areas of thermal refugia

Table 7-9—Potential effects of climate change on anadromous salmonids of the Pacific Northwest, by life-history stage (continued)

Life stage and habitat	Potential effect of climate change	Ecological consequences	Ecological implications	Potential actions
Eggs and alevins:				
Freshwater	Elevated winter water temperatures	Increased rates of development (McCullough 1999, Neuheimer and Taggart 2007)	Smaller size at emergence (Beacham and Murray 1990, Elliott and Hurley 1998)	Year-round monitoring of water temperatures
		Earlier time of emergence (Holtby 1988)	Increased growth rates, earlier timing of smolting, and smaller size at ocean entry, but decreased marine survival (Holtby and Scrivener 1989, Schindler et al. 2005)	Increase availability of floodplain and off-channel habitats
	Increased winter flows (Hamlet and Lettenmaier 2007, Hamlet et al. 2005, Tague and Grant 2009)	Increased scour of redds (Battin et al. 2007)	Reduced survival (Battin et al. 2007, Leppi et al. 2014, Shanley and Albert 2014)	Increase connection to floodplain, remove roads and infrastructure that restrict access to floodplain, with wood placement in and near low-gradient spawning areas
		Increased landslides and flooding (Dale et al. 2001, Hamlet and Lettenmaier 2007, Miller et al. 2003)	Decreased habitat quality in short term but improved conditions in long term (Bisson et al. 2009; Flitcroft et al. 2016a; Reeves et al. 1995; Rieman et al. 2006, 2015)	Road removal and culvert improvements in most susceptible areas Silvicultural treatment of plantations in most susceptible areas to increase size of trees
Juveniles:				
Freshwater	Higher spring flows (Hamlet and Lettenmaier 2007, Hamlet et al. 2005, Tague and Grant 2009)	Increased access to floodplain and off-channel habitats	Increased growth and survival if floodplains and off-channel habitats available (Brown and Hartman 1988, Moore and Gregory 1988, Peterson 1982a); decreases if not	Increased access to floodplain and off-channel habitats
	Earlier onset of low flows (Hamlet and Lettenmaier 2007, Hamlet et al. 2005, Tague and Grant 2009)	Reduced habitat availability (Battin et al. 2007, Luce and Holden 2009, Mantua et al. 2010, Stewart et al. 2005)	Reduced survival (Battin et al. 2007, Mantua et al. 2010)	Identify areas in network that are likely to be refugia during low-flow period and improve habitat conditions, including improving riparian conditions to reduce water temperature

Table 7-9—Potential effects of climate change on anadromous salmonids of the Pacific Northwest, by life-history stage (continued)

Life stage and habitat	Potential effect of climate change	Ecological consequences	Ecological implications	Potential actions
	Increased summer water temperatures (Isaak et al. 2010)	Reduced growth and survival if temperature increases are beyond favorable range (Crozier and Zabel 2006, Isaak et al. 2010, Marine and Cech 2004, Royer and Minshall 1997, Scarnecchia and Bergersen 1987)	Smaller size and reduced survival (ISAB 2007, Quinn and Petersen 1996)	Improve riparian conditions
		Increased growth if temperatures move into more favorable range		
		Altered outcomes of interactions with other species (ISAB 2007, 2012; Petersen and Kitchell 2001; Reeves et al. 1987)	Reduce growth and survival (ISAB 2007, Petersen and Kitchell 2001, Reeves et al. 1987) Change in structure and composition of fish communities (warmwater species increase) (ISAB 2012)	
	Increased nonsummer water temperatures	Increased growth rates (Ebersole et al. 2006, Sogard et al. 2010)	Increased growth and survival	Year-round monitoring of water temperatures Provide access to intermittent streams, off-channel habitats, and floodplains
Smolts:				
Freshwater	Earlier onset of low flows (Hamlet and Lettenmaier 2007, Hamlet et al. 2005, Tague and Grant 2009)	Altered timing and rate of ocean migration	Decreased marine survival (Holtby 1988, Rechisky et al. 2009)	
	Warmer water temperatures (Isaak et al. 2010)	Smaller size at ocean entry	Reduced marine survival (Holtby and Scrivener 1989, Quinn and Peterson 1989, Slaney 1988)	Improve riparian conditions
Ocean	Altered timing of ocean upwelling (Barth et al. 2007, Snyder et al. 2003)	Reduced availability of food resources (Nickelson 1986, Scheurell and Williams 2005)	Reduced marine survival (Holtby and Scrivener 1989, Quinn and Peterson 1989, Slaney 1988)	

influence outbreaks of insects and forest diseases (Spies et al. 2010) in some cases, leading to alterations of forest stands that affect aquatic habitats. Wildfires, insects, and forest diseases should not be viewed strictly as threats to native fishes, however—they may also provide benefits. They can create openings and patches along water bodies that result in more complex stream channels and greater biodiversity (Flitcroft et al. 2016a, Reeves et al. 1995, Rie- man et al. 2006). In addition, the erosional processes that accompany these disturbances are important for recruiting wood and coarse sediment that form essential habitats for many aquatic organisms (Benda et al. 2004). Thus, actions that seek to control erosion and other ecological processes that occur following wildfire may have long-term and unintended negative consequences for aquatic ecosystems (Chin et al. 2016, Harris et al. 2015).

The effects of climate change on aquatic ecosystems in the NWFP area expressed through wildfires, insects, and diseases will be complex and difficult to predict, but it will be important to examine the current responses to wildfire and consider making potential changes to allow fire to be more ecologically beneficial. Climate change will likely influence the expansion of nonnative plant and animal species in the NWFP area, while at the same time either reducing or even extirpating native species (Dale et al. 2001, Garcia et al. 2014, Urban 2015). Nonnative species include undesirable invasives, species undergoing expansion of their native ranges, and nonnative species deliberately introduced for commercial, recreational, or cultural reasons. They can occur in both terrestrial (riparian) and aquatic ecosystems. Nonnative species are not always harmful to native fishes or their habitats, but in many instances they can (1) compete with, prey upon, hybridize with, or infect native species with novel pathogens; (2) greatly alter the structure of food webs; or (3) cause habitat changes that reduce the productivity of desirable aquatic organisms. See appendix 1 for a detailed discussion of invasive species in the NWFP area.

Sanderson et al. (2009) provided a useful summary of underappreciated threats to salmon posed by nonnative vertebrates, invertebrates, and plants. They concluded that threats posed by nonnative species may equal or outweigh

threats posed by traditionally perceived causes of decline—habitat alteration, harvest, hatcheries, and hydroelectric production. Many of the nonnative fishes known to harm native fishes of the NWFP area are warmwater fish species deliberately introduced from eastern North America. In some river basins, these forms have largely displaced native fishes from dominant roles in the aquatic food webs of low-elevation, low-gradient rivers (ISAB 2012). Continued warming will favor the expansion of warm-adapted species in western North America (Rahel et al. 2008), and shrinking headwater flows resulting from longer, drier summers (Moore et al. 2007) could force cool-adapted native species lower in drainage systems, where there will be greater opportunity for unwanted interactions with established populations of introduced game fishes. Restoration of riparian areas, however, can help reduce water temperatures and the potential negative consequences of climate change related to elevated water temperatures (Justice et al. 2017, Lawrence et al. 2014)

Restoration and response to climate change under the Aquatic Conservation Strategy—

Watershed improvements implemented in the Northwest Forest Plan area—An important goal of the NWFP was to create a managed federal forest landscape in which natural ecological processes sustained healthy populations of native fish and wildlife (USDA and USDI 1994a). Architects of the ACS recognized that federally managed forests might anchor the recovery of imperiled native fishes, but because of their location relative to state and private forests as well as other types of land use (which tended to be located in lowland areas), they could not ensure that appropriate conservation measures would be applied throughout the full suite of freshwater environments to which many native species, particularly anadromous salmonids, were exposed (Sedell et al. 1997). Nevertheless, many of the aquatic-conservation actions that emerged from the NWFP were considered at the time to provide more protection to aquatic and riparian habitats than had ever before been implemented on multiple-use forests in the Pacific Northwest (NRC 1996). The region's national parks and designated wilderness areas were also considered to possess

high-quality habitats in which natural ecosystem processes could operate. However, because of their scarcity and location (Reeves et al. 2016a, Sedell et al. 1994), such areas were generally believed to be inadequate to prevent species or their distinct population segments (evolutionarily significant units) from becoming imperiled, or to hasten recovery. Given the impact of climate change on fish life cycles as discussed above, how the framework and standard of guides of future forest plans could address these potential effects merits priority for future research.

Restoration of mid- and late-seral forest stands—

Concurrent with the restoration of mid- and late-seral stands in the NWFP area, the region will likely see a reduction in large openings caused by regeneration harvests (clearcuts) and by wildfire, as a result of continuing fire suppression (see chapter 3). As forest stands grow older in the seasonally transient snow (“rain-on-snow”) zone, snowfall interception by branches will diminish the accumulation of ground-level snow and will prolong melting and runoff processes during subsequent rain events (Harr 1986). Peak flows were found to increase by as much as 20 percent in small watersheds and 30 to 100 percent in larger basins over a 50-year period in the western Cascade Range of Oregon in response to road building and clearcutting (Jones and Grant 2001). However, a recent synthesis of peak-runoff studies in western Oregon and Washington (Grant et al. 2008) concluded that the incremental contribution of clearcutting to peak flows in the transient snow zone was minor relative to other types of human disturbance, and would likely be confined to stream reaches possessing 2 percent gradients with sand and gravel substrates. In areas in which climate change results in an expansion of the transient snow zone, restoration of late-seral stands is likely to reduce the frequency and possibly duration of flows that are capable of mobilizing substrates of some fish-bearing streams, which could benefit survival of developing fish eggs and alevins as well as the abundance of amphibians and benthic macroinvertebrates.

One climate trend with important implications for native fishes is the lengthening of low-flow periods during the warm season; aquatic organisms in watersheds with reduced snowpack will be especially affected by lower

summer flows. Although not thoroughly investigated, the capture of fog by tree branches in areas with summer fog can result in “fog drip” that contributes to runoff during times when rainfall is scarce (Harr 1982). Whether climate change will alter the frequency of foggy days in the NWFP area is poorly understood, but there is preliminary evidence that the intensification of wind-driven upwelling in the California current as a result of increased CO₂ could lead to more fog and increased moisture flux along the Pacific Northwest coast during the upwelling season (Snyder et al. 2003). However, Johnstone and Dawson (2010) reported that fog frequency along the northern California coast declined by 33 percent in the 20th century. Nonetheless, restoration of late-seral stands will result in taller trees with larger limbs, which could capture more moisture and deliver some of it to streams during a season when water is in short supply.

Increasing the amount and sources of large wood will help aquatic ecosystems and associated biota meet the challenges of climate change. The progressive impoverishment of large wood in Pacific Northwest streams, particularly large-diameter, habitat-forming tree trunks and rootwads, has long been recognized (Bisson et al. 1987, Sedell and Swanson 1984). Climate change is expected to change the frequency and severity of fires and the incidence of forest-pathogen outbreaks in many parts of the NWFP area (see chapters 2 and 3). However, the ensuing recruitment of large wood to streams, a key component of fish habitat, may be limited if landslide-prone headwalls that normally deliver this material to channels during and following natural disturbances no longer contain trees of the size needed to form and maintain structural fish habitats. The importance of wood recruited to streams from unstable hillslopes is often underappreciated. For example, Reeves et al. (2003) found that 65 percent of the large wood pieces and 47 percent of the large wood volume in an Oregon coastal stream originated from upslope sources. Measures that could take advantage of this source of wood include inventorying and mapping unstable headwall areas, protecting them from forestry-related disturbance, permitting natural wood-delivery processes to occur, and allowing late-seral stands to develop in these areas where appropriate (Cissel et al. 1999).

Reducing the effects of roads and passage barriers—

Reducing the hydrological and biological effects of forest roads in the NWFP area should improve watershed resilience to the adverse effects of climate change on aquatic ecosystems. Road cuts are known to be a major contributor to accelerated runoff during storms by intercepting subsurface flow and capturing it in ditches, which rapidly deliver water and fine sediment to streams (Wemple and Jones 2003). As the intensity of storms increases with gradual warming and, in some parts of the NWFP area, with greater precipitation, the risk of streambed-mobilizing runoff events will rise as well. Reducing the exacerbating effects of road drainage networks on peak flow in watersheds where roads have been decommissioned could lessen the potential for severe storms to scour eggs and alevins in stream gravels and likewise reduce the intrusion of harmful fine sediment into spawning substrates. In addition, eliminating road-related initiation points for landslides through road decommissioning will help return the frequency of mass wasting in watersheds to more natural levels.

Road corridors can serve as important invasion routes for nonnative species, especially nonnative plants (González-Moreno et al. 2015, Heckman 1999, Menuz and Kettenring 2013), and climate change is likely to favor continued expansion of nuisance and harmful exotic herbaceous species in watersheds (Dale et al. 2001). The effect of invasive plants on riparian ecosystems in federally managed forests has received relatively little study, but some plants (e.g., Asian knotweed, *Polygonum* spp.) are capable of displacing native vegetation (Urgenson et al. 2009) and disrupting the transfer of organic material from streamside vegetation to stream channels. Invasive plant-control programs are costly, and even in riparian zones where treatments have been applied, the long-term reestablishment of native plants has been difficult to achieve (Claeson and Bisson 2013). Therefore, reducing road densities in a watershed and across large areas should help forestall the movement of unwanted nonnative plants into sensitive riparian areas and protect the integrity of native plant assemblages.

Floodplain protection—One of the key tenets of the ACS was that connections between streams and rivers and their associated floodplain and wetland habitats should be pro-

tected and, if necessary, restored (Reeves et al. 2006). In valleys where rivers are unconstrained and riparian forests are well developed, off-channel habitats such as braided streams, oxbow lakes, springs, and other floodplain features provide important seasonal rearing habitats for a wide variety of aquatic and terrestrial species and are considered to be among the most biophysically complex and diverse systems on Earth (Bayley 1995). Additionally, they can be important areas of carbon storage (Sutfin et al. 2016). Flood pulses that redistribute sediment and organic matter create a dynamic mosaic of physical habitat features on floodplains (Junk et al. 1989, Stanford et al. 2005), which support diverse and productive biological communities. In forested regions of the Pacific Northwest, flood-induced channel migration creates a variety of aquatic habitat patches that differ in age and connectivity with the main channel, from connected side channels that reside within the active flood zone to disconnected side channels that become connected only during larger flood events.

Flood-induced erosion and deposition of substrate also create dynamic and heterogeneous plant communities. Early-successional species such as alder, willow, and cottonwood are generally found on newly deposited sediments, whereas mixed-species (deciduous and coniferous) mature forests and old-growth coniferous forests are found on older and more stable floodplain surfaces (Naiman et al. 2010). This spatial heterogeneity can also create highly complex and spatially structured food webs (Bellmore et al. 2013), which may be important for mediating the strength of predator-prey interactions and promoting biodiversity and resilience (Bellmore et al. 2015).

In the context of large-scale environmental stressors such as climate change, intact floodplains may be hubs of ecological resilience. The biological and physical diversity found across floodplains may promote ecological resilience in river networks via at least two pathways. First, enhanced species diversity in floodplains may provide functional redundancy within species guilds, whereby individual species extirpations may not significantly reduce ecological function (e.g., primary/secondary production, nutrient cycling) until some critical threshold is exceeded (Walker 1992). Second, the physical heterogeneity or spatial complexity found

across floodplains may provide critical refugia for individual species (Boughton and Pike 2013, Sloat et al. 2017). For example, groundwater upwelling in floodplain springbrooks can provide cold-water thermal refugia when main-channel waters exceed thermal optimums for a particular species (Ebersole et al. 2003, Torgersen et al. 1999). Unfortunately, many river-floodplain systems have been severely altered by human disturbance, which has constrained the physical processes that create and maintain habitat heterogeneity in floodplains (Tockner and Stanford 2002), and the associated resilience these habitats may provide. Although active restoration efforts are frequently targeted at recreating specific floodplain habitats (e.g., side channels), the reestablishment of natural channel-forming processes (Beechie et al. 2013), such as the “natural flow regime” (Poff et al. 1997), may be most successful at restoring the biophysical complexity of floodplains throughout the stream network over the long term and help negate potential effects of climate change.

Winners and losers—

Climate change is projected to lead to changes in the distribution and abundance of native fishes and a host of other aquatic-riparian organisms in the NWFP area. Some species will be adversely affected by climate-mediated shifts in environmental conditions; others may actually benefit from the changes. Whether conditions will become more or less favorable for a particular species depends on physiological requirements, life-history and migratory patterns, habitat preferences, shifts in aquatic-community composition, and geographic location within the region covered by the NWFP. In general, we expect that fishes that prefer warm water and benefit from alterations in aquatic food webs and hydrologic regimes that accompany climate change will likely increase in abundance and expand their ranges. Other native fishes that prefer cool water will likely suffer losses from recently established predators and competitors; elements of their habitats that are needed at different points in their life cycles will likely decrease in abundance; and their ranges will either contract or shift northward. Population fragmentation in cool-water fishes is also likely to increase as favorable thermal conditions retreat to higher elevations, and smaller populations may suffer reduced genetic variability that threatens long-term survival (Kovach et al. 2015).

For anadromous species, survival and growth at sea will depend on how climate change alters upwelling patterns, plankton blooms, forage-fish populations, predator abundance, and other potentially limiting variables. In the fisheries management community, there is no clear consensus on whether freshwater or marine environments are “more important” to regulating the abundance of Pacific salmon, but it has become apparent that both ecosystems can exert a strong influence on run size, and that there are many uncertainties about how these two ecosystems interact to govern population viability and resilience.

In the NWFP area, climate change will lead to freshwater alterations that will be more or less favorable for some fish species relative to others. In figure 7-19, we list life-history strategies of fish that could increase vulnerability to the types of habitat change discussed earlier in this chapter. These include inflexible habitat specialization; extended freshwater rearing (1 year or more); low movement and spawning stray rates; potential for extended exposure to high water temperatures in their preferred habitats; and autumn spawning, placing them at risk of exposure to flow extremes. We also list life-history and habitat requirements that are likely to fare better in future climates. These include being able to use many different habitat types (habitat generalist); an abbreviated period in fresh water prior to seaward migration; high movement and spawning stray rates; either brief exposure to high water temperatures or a tolerance of prolonged elevated temperatures; and spring spawning that occurs after peak winter flows. Fishes are then arrayed along a risk scale, ranging from those we believe to be less vulnerable to harm from climate change to those that may be moderately vulnerable, and finally to those that may be at high risk of long-term harm. No species possesses all life-history attributes that are well adapted to thriving under predicted climate regimes, just as no species possesses only attributes that are ill-adapted to all projected future conditions. However, based on what is known about climate-related trends in freshwater habitats and on detailed knowledge of the life-history requirements of native Pacific Northwest fishes, we suggest that there will be winners and losers among fish assemblages. To some extent, the NWFP addresses many of the habitat changes likely to be associated




Figure 7-19—Life cycle and habitat-preference strategies of freshwater fishes that are considered in this report to be favored (“winning”) and disfavored (“losing”) in future climates of the Northwest Forest Plan area. Beneath the lists of winning and losing strategies is a grouping of fishes along a gradient of low to high risk from climate effects. These groupings, which are somewhat subjective, are based on current knowledge of each species’ life histories, spawning and rearing locations in watersheds, and residence time in fresh water.

with climate-related alterations in federally managed forests by creating and maintaining functional riparian areas within a watershed and focusing on road removal. However, some changes (e.g., trends in marine conditions) will not be materially affected by NWFP implementation.

The geographical distribution of native fishes and the variation in their life histories, combined with the wide range of effects of climate change on freshwater environments, make it difficult to predict which species will benefit most from NWFP aquatic-habitat protections. In figure 7-20, we divided the NWFP area into four zones: eastern, western, northern, and southern. The western zone includes watersheds draining coastal mountain ranges, whereas the eastern zone includes central lowlands of the NWFP area (Puget Sound, Willamette Valley, and California’s Central Valley) and western drainages of the Cascade Range and Siskiyou Mountains. The northern zone includes all river systems north of the Columbia River; the southern zone includes river systems southward into the Sacramento River.

The zones are not mutually exclusive because the northern and southern zones include both eastern and western areas; however, some fishes occur primarily in western coastal systems and others are found primarily in eastern portions of the NWFP area.

Based on different types of improvements to aquatic habitats from implementation of the NWFP that mitigate harmful effects of climate change as discussed above, figure 7-20 lists native salmonids that are likely to benefit in some way from the framework and standards and guidelines introduced by the NWFP. A few of the fishes (e.g., Chinook and coho salmon, steelhead [*O. mykiss*], and coastal cutthroat trout [*O. clarkii clarkii*]) are found throughout the NWFP area and therefore occur on each list; others (e.g., westslope cutthroat trout, *O. clarkii lewisi*) are limited to relatively small regions of the NWFP area. Figure 7-20 does not include nonnative species or nonsalmonids. In general, nonsalmonids (e.g., native minnows and suckers) are likely to benefit from climate warming (although see




Figure 7-20—Native salmonid fishes in the Northwest Forest Plan area that are likely to benefit in some way from environmental protections from the harmful effects of climate change, grouped by different geographical zones (see text).

Moyle et al. [2013], who suggested that this may not be true in California) and may or may not respond to NWFP aquatic-habitat improvements such as fish-passage barrier removal. Nonnative salmonids (e.g., introduced chars—brook and lake trout, *Salvelinus fontinalis* and *S. namaycush*) will probably be adversely affected by climate change, but also may or may not benefit from NWFP actions. Other introduced species, especially warmwater fishes (e.g., sunfishes and basses, Centrarchidae and *Micropterus* spp.) will likely become more abundant and may increase the risk of predation, competition, and exotic disease exposure to native fishes. However, restoration of riparian habitats may reduce water temperatures and restrict expansion of these fish (Lawrence et al. 2014). Also, the effects of hatchery fish may reduce the potential of wild populations to respond to climate change (Quiñones et al. 2014a, 2014b).

Climate refugia can also be projected for amphibian species (Shoo et al. 2011), with myriad ecological consequences. For lentic-breeding amphibians in the NWFP area, higher-elevation-adapted Cascades frogs (*R. cascadae*) may be faced with shifts in their breeding-habitat conditions. In addition, they may encounter novel interactions with species associated with warmer, lower elevation habitats, such as native northern red-legged frogs (*Rana aurora*), which may spread to higher elevations with altered climate. The low- to mid-elevation-adapted red-legged frogs may in turn encounter invasive American bullfrogs (*Lithobates catesbeianus*), which now occur at warm, low elevations. Similarly, because mountain streamflows are projected to change, torrent salamanders (*Rhyacotriton* spp.) associated with intermittent streams, could have a truncated active season, retreating below ground as small streams dry earlier in the season, possibly affecting survival and reproduction.⁹ They may also move downstream and be faced with new interactions with larger predatory salamanders or fish in perennial reaches. If they migrate downstream, their over-ridge dispersal to new watersheds may be affected, as distances between flowing water bodies increase. Hence their populations could become more isolated and vulnerable to stochastic events. For

terrestrial-breeding salamanders, we can project the consequence of warmer, drier conditions by examining the distribution of current species in the drier portion of the Northwest; these are species for which climate change has already occurred. Optimal habitat for the Siskiyou Mountains salamander (*Plethodon stormi*) is modeled to occur on the shaded side of mountain ridges and in cooler riparian areas in the dry and warm southern Oregon landscape (Suzuki et al. 2008), and the black salamander (*Aneides flavipunctatus*) appears to become a riparian associate in dry portions of its range (Nauman and Olson 2004). Hence for cool, moisture-dependent species, riparian areas and north-facing slopes with hill shading may become more important with projected changes in climate. Alternatively, as for torrent salamanders, their activity pattern may be altered, with reduced surface activities during dry times and possible consequences for survival. Range shifts for temperature- and moisture-dependent species have also been projected for pathogens of aquatic organisms, such as the amphibian chytrid fungus (*Batrachochytrium dendrobatidis*), which is predicted to increase in occurrence probability in the NWFP area with climate change (Xie et al. 2016).

Implementation of the NWFP represented a significant change in the approach to protection and management of freshwater habitats in federal forests of the Pacific Northwest. Although not directed at mitigating the negative effects of climate change on native aquatic organisms at its outset, the protections provided under the standards and guidelines of the NWFP will benefit populations of native coldwater fishes throughout their life cycles and will help maintain the diverse mosaic of habitat types on the landscape that is essential for population resilience (Beechie et al. 2013, Bisson et al. 2009). However, although many aquatic and riparian habitats in federal forests are likely to retain favorable conditions for aquatic-riparian biota or to slowly improve as watershed-restoration actions are undertaken, it is important to recognize that federally managed forests are usually embedded in a landscape that includes many different types of landowners and uses, and that the standard of environmental protection for other lands is quite different from ACS-based standards and guidelines of the NWFP (Reeves et al. 2016a). Climate-related changes

⁹ Unpublished data. On file with: Deanna Olson, Forestry Sciences Laboratory, 3200 SW Jefferson Way, Corvallis, OR 97331.

in aquatic and riparian habitats on nonfederal lands may be much less favorable for native aquatic organisms and more favorable for a variety of nonnative species.

As the biological communities of whole river systems are transformed under a changing climate, there will be a continuing need to monitor the role that federal forests play in conserving native aquatic organisms in the NWFP area. It will be critical for planners to identify vulnerabilities to climate change, and to incorporate approaches that allow management adjustments as the effects of climate change become apparent (Joyce et al. 2009). Because of the nature of environmental variability, the inevitability of novelty and surprise, and the range of management objectives and situations across the NWFP area, no single approach will fit all situations. A range of management options could include practices focused on mitigating or negating the effects of climate change by building resistance and resilience into current ecosystems, and on managing for change by enabling ecosystems and associated biota to adapt to climate change (Joyce et al. 2009, Perry et al. 2015). Better and more widespread implementation of already known practices that reduce the effects of existing stressors represents an important “no-regrets” strategy (Joyce et al. 2009). These management opportunities will require consideration of the Forest Service’s adaptive capacity, including availability of personnel with the expertise to conduct required technical analyses, and being able to work cooperatively with the public and other federal agencies to develop and implement the resulting management strategies.

The marine environment is likely to be a major challenge for Pacific salmon in the NWFP area. The predicted effects of climate change on the oceans, including acidification and increased temperatures, and their potential ecological consequences, reduced survival and size of returning adult fish, were described earlier. Pacific salmon have survived climate shifts in the past (Waples et al. 2009) and likely have the ability to persist in many areas of their current range even under more pessimistic climate change scenarios. Salmonid populations exhibit large genetic and phenotypic diversity relative to many other bony fishes (Crozier et al. 2008, Schindler et al. 2010, Waples 1991) and can adapt to changing conditions rapidly (Healey and

Prince 1995, Quinn et al. 2001). This diversity has allowed for persistence in highly dynamic and ecologically diverse environments in the past (Greene et al. 2009, Moore et al. 2014, Waples et al. 2009) and will be a key to future survival (Copeland and Vendetti 2009, Mangel 1994). However, we note that Gienapp et al. (2008) cautioned that our knowledge about the role of genetic variation and the ability of natural populations to respond adaptively to current and future environmental change is limited, and that assuming that adaptation can or will happen is risky because of the uncertain rate and extent of climate change, effects of invasive species, and altered ecological processes. The challenge to managers will be to conserve natural environmental complexity in space and time so it can provide the physical template for maintaining genotypic and phenotypic diversity in populations that are currently strong, or to restore environmental complexity where it is currently compromised.

Research Needs, Uncertainties, Information Gaps, and Limitations

The scientific basis of the ACS is still sound and is supported by new science produced since its inception by FEMAT in 1993. However, we have learned much about relationships of riparian vegetation to stream habitats and environments that has refined and modified some hypotheses that were used to develop the ACS in the early 1990s. A major knowledge gain has related to the behavior of aquatic and riparian ecosystems in space and time. At the time the ACS was developed, it was assumed that these systems were relatively stable through time. However, recent science is suggesting that these systems may be very dynamic in space and time, similar to terrestrial systems, and that aquatic organisms are adapted to this dynamism. Implementing this perspective in management actions will be challenging. It is not consistent with many current regulatory approaches, which require aquatic and riparian ecosystems to meet a given standard. Also, a dynamic perspective could be incorporated into the requirements for range of natural variability and all lands consideration of the 2012 planning rule, but will likely require close coordination between managers and researchers.

Emerging science also suggests that the absence of disturbance and management in upland terrestrial ecosystems, primarily fire, may be affecting vegetation, and combined with climate change, is likely altering these ecosystems (see chapters 2 and 3). The same trends are likely occurring in riparian and aquatic ecosystems in a manner that is not fully understood at present; this could be a useful subject for research conducted in an adaptive-management context to provide information to managers, regulators, and policy-makers in a timely manner.

Climate change is expected to affect aquatic and riparian ecosystems throughout the NWFP area, though with much uncertainty. Effects will likely differ widely within and among watersheds and geographic areas, necessitating the development of new approaches to identify this variation and help craft strategies and programs for mitigation and adaptation. Much of the focus has been on individual species. Research that focuses on understanding potential effects over the life history of species and how effects may cascade through life-history stages, as well as consideration of community-level effects, is critical. Understanding the effects on water quantity and quality is also important, particularly across spatial scales within watersheds, among watersheds, and across seasons and years. It is likely that aquatic and associated terrestrial ecosystems will change in uncertain, and maybe unpredictable, ways under a changing climate, and that this change will vary widely across the NWFP area. Having the capacity to do the needed analysis will also be critical for the involved agencies to successfully meet this challenge in a timely and effective manner, particularly in an era when budgets and personnel for federal land-management agencies are declining (see chapter 8). Thus, development of cost-effective and scientifically sound analysis procedures performed with close collaboration between research and management is key to addressing this need.

The contribution of federal lands to the conservation and recovery of ESA-listed fish continues to be important. However, federal lands alone are likely to be insufficient in geographic scope to reach the comprehensive goals of the NWFP relative to recovery of listed fish, particularly many evolutionarily significant units of Pacific salmon, as originally expected by FEMAT (1993) and the record of decision

(USDA and USDI 1994a). Although the geomorphic setting of streams on federal lands may be as capable as originally expected of providing sufficient favorable habitat, particularly for salmon, streams on state and private lands may have a much greater potential to provide habitat in many watersheds. Thus, it will be important to work closely with adjoining landowners and other interested parties to develop more comprehensive efforts across species ranges. The development of incentive programs is likely to be important to build partnerships for fish-habitat management across land ownerships. Developing an understanding of the variation in the capacity of watersheds to provide favorable conditions for fish and other aquatic biota could be critical to the success of such programs.

Monitoring the effectiveness of the ACS will continue to be important. Some meaningful uncertainties remain regarding the aquatic-riparian monitoring approaches, especially relative to whether they are capable of capturing the effects of the ACS on a wide range of ecological processes and species of aquatic and riparian ecosystems. Research is needed to test the ecological validity of individual metrics and different ways of combining metrics to represent different components of complex and diverse aquatic and riparian ecosystems and communities. It would, therefore, be prudent to compare alternative approaches in the face of the new understanding about the behavior of aquatic and riparian ecosystems in time and space and the yet-to-be-understood effects of disturbance or lack of disturbance, climate change, and novel ecosystems. A related research need is to better understand the relationship of the productivity of aquatic biota, which include organisms other than salmonids, in the context of different upland vegetation and in-channel successional stages or restoration treatments. This type of information can feed into watershed assessments to better ensure that the effects of the ACS are captured more comprehensively relative to the biota that are a key ecosystem service of aquatic-riparian ecosystems. In particular, we lack information about the amount, pattern, and type of restoration activities that have occurred in upland and riparian forests. Implementation monitoring has not been adequate to enable a sufficient understanding of the consequences of restoration actions (or lack of actions),

especially relative to how they may have altered aquatic ecosystems in space and time.

Roads and their effects will continue to be a major issue in the NWFP area. Both research on road effects and the continued development of analysis tools such as Geomorphic Roads Analysis and Inventory Package (Black et al. 2012) are important. In particular, understanding the consequences of focusing on small segments rather than the entire network should be a priority. The same is true for effects of culverts on ecological processes and the movement of aquatic biota. These are current priorities given the uncertainties of climate change. Also, understanding how to balance fire management, recreation, and other needs against potential negative aspects of roads will require a concerted cooperative effort of managers and physical, biological, and social scientists.

A key uncertainty that has emerged from our analysis is how to understand and assess the effects of “no-action” management options and tradeoffs of managing for one factor (e.g., water temperature or wood recruitment) on other ecological processes or attributes. The assumption has been that focusing on one concern would not influence other processes or attributes, and that taking no action was synonymous with having no effect. However, these assumptions are questionable and deserve increased consideration and focus by researchers.

Several other topics relating to the components of the ACS merit further research. Watershed analysis could be reexamined so that it is conducted more efficiently and considers the appropriate spatial scales, including a smaller watershed of interest and its context within a larger basin. The larger scale context is particularly relevant for effective landscape-scale planning. In addition, no formal evaluation of the potential effectiveness of the network of key watersheds was conducted during development of the NWFP, nor has such an evaluation been attempted since it was implemented. New concepts, tools, and emerging understandings about aquatic ecosystems are now available to better assess and increase the potential effectiveness of key watersheds. Our understanding of aquatic ecosystems is incomplete (though evolving) at this time, but because there could be significant implications for the productivity of these systems, they will continue to be a major focus of research.

Conclusions and Management Considerations

The goal of the ACS was to maintain and restore aquatic-riparian ecosystems on federal lands within the range of the northern spotted owl. A review of monitoring efforts and the pertinent scientific literature suggests that (1) aquatic ecosystems in the NWFP area are likely improving as expected, albeit slowly; (2) the fundamental tenets and ecological framework of the ACS are sound, and we are gaining more explicit understanding of several components that over time will have important implications for future management; and (3) opportunities exist for implementing parts of the ACS differently while continuing to achieve its goals. The third finding is particularly applicable to the riparian-reserve component of the ACS, where more active management may help to address potential concerns about the effects of the lack of natural disturbance (primarily wildfire), and climate change.

The following is a detailed summary of our main findings and conclusions. We also note to which guiding questions the conclusion applies.

Guiding Questions

1. Is the scientific foundation for the ACS valid, or does the science developed since 1993 suggest potential changes or adjustments that could be made to the ACS?
2. What is the basis of trends observed in the ACS monitoring program, and what are the limitations, uncertainties, and research needs related to monitoring?
3. What is known about variation of characteristics of unmanaged streams and riparian ecosystems in relation to the stream networks across the NWFP area?
4. What has been learned about the effects of riparian vegetation on stream habitat and environment?
5. What effects have human activities had on stream and riparian ecosystems?
6. What is the scientific basis for restoration management in riparian reserves, and how does restoration relate to the ecological goals of the ACS?
7. What is the capacity of federal lands in the NWFP area to contribute water for a suite of economic, recreational, and ecological uses?

8. What are the potential effects of climate change on aquatic ecosystems in the NWFP area, and are they adequately addressed by the ACS?

Science Foundation for the ACS (Question 1)

The scientific foundation of the ACS is generally sound.

1. It is a coarse-filter approach designed to protect and restore ecological processes that create and maintain favorable habitat conditions for native anadromous salmonids. This assumes that if conditions are favorable for these organisms, then they should be suitable for other aquatic and riparian associated organisms.
 - a. Verifying these assumptions could be a research priority.
 - b. There is growing scientific support for larger scale ecological processes acting at small-to-large watershed scales affecting salmonid habitats and populations; these include landslides delivering sediment and wood, canopy closure, and hill-shading effects on aquatic-riparian temperatures, and the contribution of headwaters to downstream conditions and populations.
 - c. The ecological process and species-habitat emphasis areas of the ACS are supported, but since 1993 additional factors have come to the forefront.
 - i. More aquatic species have been considered for listing as threatened and endangered, some requiring more focused attention than the regional scale of the ACS, and consideration of threats on a case-by-case basis.
 - ii. Aquatic invasives have emerged as an elevated concern because of their effects on native species.
 - iii. Anthropogenic disturbances from timber-harvest activities, including road building and maintenance, remain key concerns for aquatic-riparian ecosystems, but new concerns about the extent and severity of wildfire and climate change have emerged as research and monitoring priorities.
- iv. Reliance on federal lands alone cannot address the conservation need to maintain or restore well-distributed populations of all aquatic-riparian species; key salmonid habitats rely on nonfederal lands, and fragmented federal land ownerships affect aquatic-riparian-terrestrial habitat connectivity for organisms dependent upon aquatic-riparian ecosystems.
2. The scientific foundation for the riparian reserve network is valid. The riparian reserve network was intended to identify the outer boundary of the aquatic/riparian ecosystem.
 - a. Since 1993, new science supports riparian buffers to maintain aquatic-riparian processes, habitat conditions, and species.
 - b. Our ecological knowledge about non-fish-bearing streams has increased tremendously since 1993, and the approach for protecting them is supported.
 - c. However, there are suggestions that the second site-potential tree-height on fish-bearing streams may not be required to maintain microclimatic conditions within the first tree-height.
 - d. There are potential options available to move away from fixed-width riparian buffers toward riparian management that considers the variability in ecological context within the stream network and specifies management depending on ecological importance and risk.
 - e. Passive restoration approaches of riparian forests in streamside buffers have dominated management choices; active restoration might be acceptable in some locations and could accelerate achievement of goals such as growth of large trees to supply key pieces of large wood in the future.
 - f. Although science has addressed reach-scale effects of riparian reserves, research on effects of larger scale management activities (e.g., small to large watersheds) is becoming a new research priority.
 - g. Implementation, effectiveness, and validation monitoring of NWFP riparian reserves has not formally occurred, and is an emerging priority.

3. The use and structure of the key watershed network are supported by recent science.
 - a. There is emerging evidence that the key watersheds do not have the capacity to support and provide favorable habitat for ESA-listed fish to the extent that was originally assumed.
 - b. Also, the assumption that habitat conditions in old-growth forests are the most favorable for native salmonids is being questioned.
 - c. A review of the key watershed network and the criteria for selecting watersheds would be useful and timely.
 4. Watershed analysis remains an important process for developing and assessing management options.
 - a. New analytical tools and processes are available that could be used to improve these analyses and make them more cost effective; this is a research priority relative to individual watershed analysis as well as assessment of multiple watersheds across the region that may have differing contexts.
 - b. The ability of the Forest Service and other federal land managers to conduct such analyses may be limited by a declining workforce and technical competency; see chapter 8 for more detailed discussion.
 5. A tremendous amount of effort has been directed at restoring degraded watersheds and the associated aquatic and riparian ecosystems.
 - a. The vast majority of this effort has been directed toward fish-bearing streams. More effort could be directed at the non-fish-bearing portions of the stream network, which will be important to addressing potential effects of climate change.
 - b. Implementation, effectiveness, and validation monitoring is needed to assess restoration activities and contribute to adaptive-management processes.
1. The primary reasons for improvement are likely a reduction in the extent of roads, primarily in key watersheds, and an increase in the number of large trees in the riparian reserve.
 2. Regionally, there is a signature of wildfire interacting with AREMP restoration criteria in some places in the NWFP area. The ecological significance of this interaction is unclear and merits examination.
 3. Assessing watershed condition is inherently challenging, and this synthesis has highlighted a number of areas for further research and management focus:
 - a. Use of multiple independent measures of watershed attributes makes it difficult to assess the overall condition of a watershed.
 - b. Development of aquatic-riparian monitoring programs requires a clear articulation of which biota and associated functional characteristics of habitats and ecosystems are being considered, tying these to our understanding of patterns of change over space and time, and how they are likely to be altered as a result of the actions of interest.
 - i. It is important to clearly describe the ecological context of aquatic-riparian monitoring, for example, to determine to what extent AREMP should focus on environmental parameters that measure habitat for native salmonids or on other aquatic and riparian-dependent organisms.
 - ii. Within a dynamic aquatic-riparian ecosystem framework, change is anticipated, but a challenge for monitoring is to assess alterations in conditions that may reflect restoration or other trajectories of patterns in response to a variety of human actions and other events.
 - c. A key consideration in development of reference distributions for comparison with current conditions is including the entire natural range of conditions that an ecosystem can experience (natural range of variability). This is critical to be able to evaluate the implications of change.
 - i. Reference conditions that are too narrowly or broadly defined can skew the interpretation of monitoring results and introduce uncertainty

Monitoring of the ACS (Question 2)

The AREMP results suggest that the condition of aquatic and riparian ecosystems in the NWFP area is improving, albeit slowly, as was originally expected owing to the extensive amount of degradation and lengthy time needed for recovery.

into the process. This is an active area of research for the NWFP area: Are there neighborhood, provincial, or regional patterns to consider? We suggest further exploration of the use of reference conditions and their potential utility for diverse analytical approaches, including consideration of how to use them in concert with state-transition models and the potential development of novel conditions in the future.

- d. Refinement of the objectives and approaches of aquatic-riparian monitoring programs, including AREMP, is anticipated as our understanding of these ecological systems improve and new analytical tools are developed. Advances in watershed-condition assessment procedures will be important to ensure the validity and reduce the uncertainty of future results and their implications for management.

Aquatic and Riparian Ecosystems (Questions 3–6)

1. Unaltered aquatic and riparian ecosystems likely exhibit a wide range of conditions in space and time, locally and across the NWFP area, depending to a large degree on the magnitude and frequency of the associated disturbance regime. (See chapter 3 for more details.)
 - a. Headwater streams tend to be dominated primarily by conifers much of the time.
 - i. The biological processing of vegetation that falls into the stream (allochthonous material) is a primary energy source for downstream fish-bearing streams.
 - b. In the middle parts of the stream network, the riparian zone is composed of a mixture of conifers and deciduous hardwoods.
 - i. Hardwoods are important sources of high-quality allochthonous material important for system productivity.
 - ii. Hardwoods are scarce in many areas because of the conversion of riparian areas to conifer-dominated plantations. There may be important
- implications to system productivity that need to be explored.
2. Human impacts have extensively altered riparian ecosystems.
 - a. An estimated 30 to 50 percent of the riparian reserve has been converted to single-species plantations, primarily conifer, as a result of past management that harvested trees to the edge of the streams throughout the network. In headwater streams that have experienced increased rates of landsliding, riparian zones are frequently dominated by alder.
 - b. The trajectory of riparian and aquatic ecosystems has been altered as a result, reducing ecological variability across the area of the NWFP.
 - c. Additionally, fire exclusion (see chapter 3) and climate change are likely altering aquatic and riparian ecosystems in ways that are not fully recognized or appreciated at this time.
3. Restoration activities in riparian reserves have been limited because of concerns about potential negative effects, particularly increased water temperatures and decreased wood-delivery potential, and lack of trust of the Forest Service (see chapter 12).
 - a. Restoration activities have primarily been restricted to fish-bearing streams.
 - i. Assessment of these activities has been extremely limited, so it is not possible to quantify the effects.
 - ii. More active management may be needed.
 - iii. The question of whether the increased risk is sufficiently offset by the long-term gains realized from active restoration or other activities within portions of the riparian buffers is a key research need.
 - b. Passive restoration has been the dominant policy in riparian reserves.
 - c. This approach assumes that “no activity equals no effect.” However, this assumption is questionable, and “no activity” may actually compromise or eliminate key ecological processes such as development of the largest trees.

4. Non-fish-bearing streams have received little attention in terms of restoration.
 - a. These streams can be important sources of large wood for streams lower in the network. Improving stand conditions is critical to maintaining this important ecological function.

Water Contributions From Federal Lands (Question 7)

The capacity of federal lands in the NWFP area to contribute water for a suite of purposes varies widely among forests, ranging from less than 20 percent of the total flow in some basins to more than 40 percent in others.

Climate Change (Question 8)

The primary effects of climate change in the NWFP area will be increased water temperatures, decreased streamflows in summer, and increased winter streamflows. The extent of these effects will vary widely depending on location and local topographic features.

1. The ACS has the potential to meet these challenges, but it will take a focused effort to do so, including:
 - b. Conducting local-scale analyses.
 - c. Considering “all lands.”
 - d. Shifting the focus of management and restoration from increasing population sizes to increasing the life-history diversity of aquatic and riparian organisms.
 - e. Recognizing that there will be “winners” and “losers”—because of their inherent capacity to adapt, some organisms will increase while others are likely to decrease.
2. With regard to anadromous fish, changes in ocean conditions (water temperature, acidification, and timing of upwelling), which are beyond the capacity of the federal land-management agencies to influence, may exert a stronger influence on populations than changes in freshwater ecosystems.

Literature Cited

- Abbe, T.B.; Montgomery, D.R. 2003.** Patterns and processes of wood debris accumulation in the Queets River basin, Washington. *Geomorphology*. 51: 81–107.
- Achord, S.; Zabel, R.W.; Sandford, B.J. 2007.** Migration timing, growth, and estimated parr-to-smolt survival rates of wild Snake River spring-summer Chinook salmon from the Salmon River basin, Idaho, to the lower Snake River. *Transactions of the American Fisheries Society*. 136: 142–154.
- Acker, S.A.; Gregory, S.V.; Lienkaemper, G.; McKee, W.A.; Swanson, F.J.; Miller, S.D. 2003.** Composition, complexity, and tree mortality in riparian forests in the central Western Cascades of Oregon. *Forest Ecology and Management*. 173: 293–308.
- Adams, B.L.; Zaugg, W.S.; McLain, L.R. 1973.** Temperature effect on parr-smolt transformation in steelhead trout (*Salmo gairdneri*) as measured by sodium-potassium stimulated adenosine triphosphatase. *Comparative Biochemistry and Physiology*. 44A: 1333–1339.
- Agee, J.K. 1993.** Fire ecology of Pacific Northwest forests. Washington, DC: Island Press. 493 p.
- Al-Chokhachy, R.; Black, T.A.; Thomas, C.; Luce, C.H.; Rieman, B.; Cissel, R.; Carlson, A.; Hendrickson, S.; Archer, E.K.; Kershner, J.L. 2016.** Linkages between unpaved forest roads and streambed sediment: why context matters in directing road restoration. *Restoration Ecology*. 24: 589–598.
- Al-Chokhachy, R.; Roper, B.B.; Archer, E.K. 2010.** Evaluating the status and trends of physical stream habitat in headwater streams within the interior Columbia River and upper Missouri River basins using an index approach. *Transactions of the American Fisheries Society*. 139: 1041–1059.

- Alexander, L.C.; Hawthorne, D.J.; Palmer, M.A.; Lamp, W.O. 2011.** Loss of genetic diversity in the North American mayfly *Ephemerella invaria* associated with deforestation of headwater streams. *Freshwater Biology*. 56: 1456–1467. doi:10.1111/j.1365-2427.2010.02566.x.
- Allan, C.; Curtis, A. 2005.** Nipped in the bud: why regional scale adaptive management is not blooming. *Environmental Management*. 36: 414–425.
- Allan, J.D. 2004.** Landscapes and riverscapes: the influence of land use on stream ecosystems. *Annual Review of Ecology, Evolution, and Systematics*. 35: 257–284.
- Allen, D.; Dietrich, W.; Baker, P.; Ligon, F.; Orr, B. 2007.** Development of a mechanistically based, basin-scale stream temperature model: applications to cumulative effects modeling. In: Standiford, R.B.; Giusti, G.A.; Valachovic, Y.; Zielinski, W.J.; Furniss, M.J., eds. *Proceedings of the redwood region forest science symposium: What does the future hold?* Gen. Tech. Rep. PSW-GTR-194. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 11–24.
- Allen, T.F.; Starr, T.B. 1982.** *Hierarchy: perspectives for ecological complexity*. Chicago, IL: University of Chicago Press. 310 p.
- Andersen, H.; Lanigan, S. 2010.** Aquatic and Riparian Effectiveness Monitoring Program: invasive species report, 2009 field season. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, and U.S. Department of the Interior, Bureau of Land Management, Oregon State Office. <https://www.fs.fed.us/r6/reo/monitoring/reports/AREMP%20Aquatic%20Invasive%20Species%20rpt%202009.pdf>. (31 May 2017).
- Anderson, P.D.; Larson, D.J.; Chan, S.S. 2007.** Riparian buffer and density management influences on microclimate of young headwater forests of western Oregon. *Forest Science*. 53: 254–269.
- Aquatic Invasive Species Network [AISN]. 2017.** Aquatic Invasive Species Network: a collaboration to prevent the introduction and spread of aquatic invasive species. Pacific States Marine Fisheries Commission. <http://www.westernais.org/monitoring>. (6 July 2017).
- Araki, H.; Schmid, C. 2010.** Is hatchery stocking a help or harm?: evidence, limitations and future directions in ecological and genetic surveys. *Aquaculture*. 308(Suppl. 1): S2–S11.
- Archer, E.; Ojala, J.V. 2016.** Stream habitat condition for sites in the USDA Forest Service Region 6. On file with: PacFish/InFish Biological Opinion (PIBO) Monitoring Program, USDA Forest Service, 324 25th Street, Ogden, UT 84401.
- Argerich, A.; Haggerty, R.; Johnson, S.L.; Wondzell, S.M.; Dosch, N.; Corson-Rikert, H.; Ashkenas, L.R.; Pennington, R.; Thomas, C.K. 2016.** Comprehensive multiyear carbon budget of a temperate headwater stream. *Journal of Geophysical Research Biogeoscience*. 121: 1306–1315.
- Arismendi, I.; Johnson, S.L.; Dunham, J.B.; Haggerty, R. 2013a.** Descriptors of natural thermal regimes in streams and their responsiveness to change in the Pacific Northwest of North America. *Freshwater Biology*. 58: 880–894.
- Arismendi, I.; Safeeq, M.; Johnson, S.L.; Dunham, J.B.; Haggerty, R. 2013b.** Increasing synchrony of high temperature and low flow in western North American streams: double trouble for coldwater biota? *Hydrobiologia*. 712: 61–70.
- Arismendi, I.; Safeeq, M.; Dunham, J.B.; Johnson, S.L. 2014.** Can air temperature be used to project influences of climate change on stream temperature? *Environmental Research Letters*. 9(8): 084015. doi:10.1088/1748-9326/9/8/084015. (1 September 2017).

- Auliya, M.; Gardia-Moreno, J.; Schmidt, B.R.; Schmeller, D.S.; Hoogmoed, M.S.; Fisher, M.C.; Pasmans, F.; Henle, K.; Bickford, D.; Martel, A. 2016.** The global amphibian trade flows through Europe: the need for enforcing and improving legislation. *Biodiversity Conservation*. 25: 2581–2595.
- Aziz, O.; Mantua, N.J.; Myers, K.W. 2011.** Potential climate change impacts on thermal habitats of Pacific salmon in the North Pacific Ocean and adjacent seas. *Canadian Journal of Fisheries and Aquatic Sciences*. 68: 1660–1680.
- Baker, D.; Ferguson, C.; Palmer, C.; Tolle, T. 2006.** The Northwest Forest Plan—the first ten years (1994–2003): implementation monitoring: summary of regional interagency monitoring results. Tech. Pap. R6-RPM-TP-04-2005. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region. 18 p. <https://re0.gov/monitoring/reports/10yr-report/implementation/CPImpMoniFirstTenYears1-23-06.pdf>. (1 September 2017).
- Barbour, M.T.; Gerritsen, J.; Snyder, B.D.; Stribling, J.B. 1999.** Rapid bioassessment protocols for use in streams and wadeable rivers: periphyton, benthic macroinvertebrates and fish. 2nd ed. Washington, DC: U.S. Environmental Protection Agency, Office of Water. 339 p.
- Barth, J.A.; Menge, B.A.; Lubchenco, J.; Chan, F.; Bane, J.M.; Kirincich, A.R.; McManus, M.A.; Nielsen, K.J.; Pierce, S.D.; Washburn, L. 2007.** Delayed upwelling alters nearshore coastal ocean ecosystems in the northern California Current. *Proceedings of the National Academy of Sciences of the United States of America*. 104: 3719–3724.
- Bates Prins, S.C.; Smith, E.P. 2007.** Using biological metrics to score and evaluate sites: a nearest-neighbour reference condition approach. *Freshwater Biology*. 52: 98–111.
- Battin, J.; Wiley, M.W.; Ruckelshaus, M.H.; Palmer, R.N.; Korb, E.; Bartz, K.K.; Imaki, H. 2007.** Projected impacts of climate change on salmon habitat restoration. *Proceedings of the National Academy of Sciences of the United States of America*. 104: 6720–6725.
- Baxter, C.V.; Fausch, K.D.; Saunders, W.C. 2005.** Tangled webs: reciprocal flows of invertebrate prey link streams and riparian zones. *Freshwater Biology*. 50: 201–220.
- Bayley, P.B. 1995.** Understanding large river-floodplain ecosystems. *BioScience*. 45: 153–158.
- Beacham, T.D.; Murray, C.B. 1990.** Temperature, egg size, and development of embryos and alevins of fish species of Pacific salmon: a comparative analysis. *Transactions of the American Fisheries Society*. 119: 927–945.
- Beamish, R.J.; Bouillon, D.R. 1993.** Pacific salmon production trends in relation to climate. *Canadian Journal of Fisheries and Aquatic Sciences*. 50: 1002–1016.
- Beckman, N.D.; Wohl, E. 2014.** Carbon storage in mountainous headwater streams: the role of old-growth forest and logjams. *Water Resources Research*. 50: 2376–2393.
- Beechie, T.; Buhle, E.; Ruckelshaus, M.; Fullerton, A.; Holsinger, L. 2006.** Hydrologic regime and the conservation of salmon life history diversity. *Biological Conservation*. 130: 560–572.
- Beechie, T.; Imaki, H.; Greene, J.; Wade, A.; Wu, H.; Pess, G.; Roni, P.; Kimball, J.; Stanford, J.; Kiffney, P.; Mantua, N. 2013.** Restoring salmon habitat for a changing climate. *River Research and Applications*. 29: 939–960.
- Beechie, T.J.; Pess, G.; Kennard, P.; Bilby, R.E.; Bolton, S. 2000.** Modeling recovery rates and pathways for woody debris recruitment in northwestern Washington streams. *North American Journal of Fisheries Management*. 20: 436–452.

- Beechie, T.J.; Pess, G.R.; Pollock, M.M.; Ruckelshaus, M.H.; Roni, P. 2009.** Restoring rivers in the twenty-first century: science challenges in a management context. In: Beamish, R.J.; Rothschild, B.J., eds. The future of fisheries science in North America. Amsterdam: Springer: 697–717.
- Beechie, T.J.; Sear, D.A.; Olden, J.D.; Pess, G.R.; Buffington, J.M.; Moir, H.; Roni, P.; Pollock, M.M. 2010.** Process-based principles for restoring river ecosystems. *BioScience*. 60: 209–222.
- Behnke, R.J. 2002.** Trout and salmon of North America. New York: The Free Press. 359 p.
- Bellmore, J.R.; Baxter, C.V.; Connolly, P.J. 2015.** Spatial complexity reduces interaction strengths in the meta-food web of a river floodplain mosaic. *Ecology*. 96: 274–283.
- Bellmore, J.R.; Baxter, C.V.; Martens, K.; Connolly, P.J. 2013.** The floodplain food web mosaic: a study of its importance to salmon and steelhead with implications for their recovery. *Ecological Applications*. 23: 189–207.
- Benda, L.; Bigelow, P. 2014.** On the patterns and processes of wood in northern California streams. *Geomorphology*. 209: 79–97.
- Benda, L.; Dunne, T. 1997a.** Stochastic forcing of sediment supply to channel networks from landsliding and debris flow. *Water Resources Research*. 33: 2849–2863.
- Benda, L.; Dunne, T. 1997b.** Stochastic forcing of sediment routing and storage in channel networks. *Water Resources Research*. 33: 2865–2880.
- Benda, L.; Hassan, M.A.; Church, M.; May, C.L. 2005.** Geomorphology of steep-land headwaters: the transition from hillslopes to channels. *Journal of the American Water Resources Association*. 41: 835–851.
- Benda, L.; Miller, D.; Sias, J.; Martin, D.; Bilby, R.; Veldhuisen, C.; Dunne, T. 2003.** Wood recruitment processes and wood budgeting. *American Fisheries Society Symposium*. 37: 49–74.
- Benda, L.; Poff, N.L.; Miller, D.; Dunne, T.; Reeves, G.; Pollock, M.; Pess, G. 2004.** The network dynamics hypothesis: how river networks structure riverine habitats. *BioScience*. 54: 413–427.
- Benda, L.E.; Cundy, T.W. 1990.** Predicting deposition of debris flows in mountain channels. *Canadian Geotechnical Journal*. 27: 409–417.
- Benda, L.E.; Bigelow, P.; Worsley, T.M. 2002.** Recruitment of wood to streams in old-growth and second-growth redwood forests, northern California, USA. *Canadian Journal of Forest Research*. 32: 1460–1477.
- Benda, L.E.; Litschert, S.E.; Reeves, G.; Pabst, R. 2016.** Thinning and in-stream wood recruitment in riparian second growth forests in coastal Oregon and the use of buffers and tree tipping as mitigation. *Journal of Forestry Research*. 27: 821–836.
- Benda, L.; Miller, D.; Andras, K.; Bigelow, P.; Reeves, G.; Michael, D. 2007.** NetMap: a new tool in support of watershed science and resource management. *Forest Science*. 53: 206–219.
- Benda, L.E.; Miller, D.J.; Dunne, T.; Reeves, G.H.; Agee, J.K. 1998.** Dynamic landscape systems. In: Bilby, R.E.; Naiman, R.J., eds. *River ecology and management: lessons from the Pacific coastal ecoregion*. New York: Springer-Verlag: 261–288.
- Benjamin, J.R.; Connolly, P.J.; Romine, J.G.; Perry, R.W. 2013.** Potential effects of changes in temperature and food resources on life history trajectories of juvenile *Oncorhynchus mykiss*. *Transactions of the American Fisheries Society*. 142: 208–220.
- Bern Convention. 2015.** 35th meeting of the Standing Committee, Strasbourg 1–4 Dec 2015: List of decisions and adopted texts. <http://www.coe.int/en/web/bern-convention/-/35th-standing-committee-meeting>. (30 March 2017).

- Bernhardt, E.S.; Palmer, M.A. 2011.** River restoration: the fuzzy logic of repairing reaches to reverse catchment scale degradation. *Ecological Applications*. 21: 1926–1931.
- Beschta, R.L.; Bilby, R.E.; Brown, G.W.; Holtby, L.B.; Hofstra, T.D. 1987.** Stream temperature and aquatic habitat: fisheries and forestry interactions. In: Salo, E.O.; Cundy, T.W., eds. *Streamside management: forestry and fishery interactions*. Contribution No. 57. Seattle, WA: University of Washington, Institute of Forest Resources: 191–232.
- Beschta, R.L.; Rhodes, J.J.; Kauffman, J.B.; Gresswell, R.E.; Minshall, G.W.; Karr, J.R.; Perry, D.A.; Hauer, F.R.; Frissell, C.A. 2004.** Postfire management on forested public lands of the western United States. *Conservation Biology*. 18: 957–967.
- Biebighauser, T.R. 2011.** Wetland restoration and construction—a technical guide. 2nd ed. Burdett, NY: The Wetland Trust. 186 p.
- Bigelow, P.E.; Benda, L.E.; Miller, D.J.; Burnett, K.M. 2007.** On debris flows, river networks, and the spatial structure of channel morphology. *Forest Science*. 53: 220–238.
- Bigler, B.S.; Welch, D.W.; Helle, J.H. 1996.** A review of size trends among North Pacific salmon (*Oncorhynchus* spp.). *Canadian Journal of Fisheries and Aquatic Sciences*. 53: 455–465.
- Bilby, R.E.; Bisson, P.A. 1998.** Function and distribution of large woody debris. In: Naiman, R.J.; Bilby, R.E., eds. *River ecology and management: lessons from the Pacific coastal ecoregion*. New York: Springer-Verlag: 324–346.
- Bilby, R.E.; Heffner, J.T. 2016.** Factors influencing litter delivery to streams. *Forest Ecology and Management*. 369: 29–37.
- Bilby, R.E.; Fransen, B.R.; Bisson P.A. 1996.** Incorporation of nitrogen and carbon from spawning coho salmon into the trophic system of small streams: evidence from stable isotopes. *Canadian Journal of Fisheries and Aquatic Sciences*. 53: 164–173.
- Bisson, P.A.; Bilby, R.E.; Bryant, M.D.; Dolloff, C.A.; Grette, G.B.; House, R.A.; Murphy, M.L.; Koski, K.V.; Sedell, J.R. 1987.** Large woody debris in forested streams in the Pacific Northwest: past, present, and future. In: Salo, E.O.; Cundy, T.W., eds. *Streamside management: forestry and fishery interactions*. Contribution No. 57. Seattle, WA: Institute of Forest Resources, University of Washington: 143–190.
- Bisson, P.A.; Davis, G.E. 1976.** Production of juvenile Chinook salmon, *Oncorhynchus tshawytscha*, in a heated model stream. *NOAA Fisheries Bulletin*. 74: 763–774.
- Bisson, P.A.; Rieman, B.E.; Luce, C.; Hessburg, P.F.; Lee, D.C.; Kershner, J.L.; Reeves, G.H.; Gresswell, R.E. 2003.** Fire and aquatic ecosystems of the western USA: current knowledge and key questions. *Forest Ecology and Management*. 178: 213–229.
- Bisson, P.A.; Dunham, J.B.; Reeves, G.H. 2009.** Freshwater ecosystems and resilience of Pacific salmon: habitat management based on natural variability. *Ecology and Society*. 14(1): 45. <http://www.ecologyandsociety.org/vol14/iss1/art45/>. (1 September 2017).
- Bisson, P.A.; Reeves, G.H.; Bilby, R.E.; Naiman, R.J. 1997.** Watershed management and Pacific salmon: desired future conditions. In: Stouder, D.J.; Bisson, P.A.; Naiman, R.J., eds. *Pacific salmon and their ecosystems: status and future options*. New York: Springer: 447–474.
- Black, T.A.; Cissel, R.M.; Luce, C.H. 2012.** The Geomorphic Road Analysis and Inventory Package (GRAIP) Volume 1: Data collection method. Gen. Tech. Rep. RMRS-GTR-280WWW. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 110 p.
- Blahna, D.J.; Asah, S.T.; Deal, R.L. 2017.** An ecosystem services framework. In: Olson, D.H.; Van Horne, B., eds. *People, forests, and change: lessons from the Pacific Northwest*. Washington, DC: Island Press: 62–75. Chapter 5.
- Blaustein, A.R.; Olson, D.H. 1991.** Declining amphibians. *Science*. 253: 1467.

- Bond, N.A.; Cronin, M.F.; Freeland, H.; Mantua, N. 2015.** Causes and impacts of the 2014 warm anomaly in the NE Pacific. *Geophysical Research Letters*. 42: 3414–3420.
- Bothwell, M.L.; Taylor, B.W.; Kilroy, C. 2014.** The Didymo story: the role of low dissolved phosphorus in the formation of *Didymosphenia geminata* blooms. *Diatom Research*. 29: 229–236.
- Bottom, D.L.; Jones, K.K.; Cornwell, T.J.; Gray, A.; Simenstad, C.A. 2005.** Patterns of Chinook salmon migration and residency in the Salmon River estuary (Oregon). *Estuarine, Coastal and Shelf Science*. 64: 79–93.
- Boughton, D.A.; Fish, H.; Pope, J.; Holt, G. 2009.** Spatial patterning of habitat for *Oncorhynchus mykiss* in a system of intermittent and perennial streams. *Ecology of Freshwater Fish*. 18: 92–105.
- Boughton, D.A.; Pike, A.S. 2013.** Floodplain rehabilitation as a hedge against hydroclimatic uncertainty in a migration corridor of threatened steelhead. *Conservation Biology*. 27: 1158–1168.
- Bowman, M.F.; Somers, K.M. 2006.** Evaluating a novel Test Site Analysis (TSA) bioassessment approach. *Journal of the North American Benthological Society*. 25: 712–727.
- Brayshaw, D.; Hassan, M.A. 2009.** Debris flow initiation and sediment recharge in gullies. *Geomorphology*. 109: 122–131.
- Brett, J.R. 1952.** Temperature tolerances in young Pacific salmon, genus *Oncorhynchus*. *Journal of the Fisheries Research Board of Canada*. 9: 265–323.
- Brett, J.R.; Shelbourn, J.E.; Shoop, C.T. 1969.** Growth rate and body composition of fingerling sockeye salmon, *Oncorhynchus nerka*, in relation to temperature and ration size. *Journal of the Fisheries Research Board of Canada*. 26: 2362–2394.
- Brosofske, K.D.; Chen, J.; Naiman, R.J.; Franklin, J.F. 1997.** Effects of harvesting on micro-climatic gradients from streams to up-lands in western Washington, USA. *Ecological Applications*. 7: 1188–1200.
- Brown, T.C.; Froemke, P. 2010.** Risk of impaired condition of watersheds containing National Forest lands. Gen. Tech. Rep. RMRS-GTR-251. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 57 p.
- Brown, T.C.; Froemke, P. 2012.** Nationwide assessment of nonpoint source threats to water quality. *BioScience*. 62: 136–146.
- Brown, T.C.; Hobbins, M.T.; Ramirez, J.A. 2008.** Spatial distribution of water supply in the conterminous United States. *Journal of the American Water Resources Association*. 44: 1474–1487.
- Brown, T.G.; Hartman, G.F. 1988.** Contribution of seasonally flooded lands and minor tributaries to the production of coho salmon in Carnation Creek, British Columbia. *Transactions of the American Fisheries Society*. 117: 546–551.
- Bryant, M.D. 2009.** Global climate change and potential effects on Pacific salmonids in freshwater ecosystems of southeast Alaska. *Climate Change*. 95: 169–193.
- Bucciarelli, G.M.; Blaustein, A.R.; Garcia, T.S.; Kats, L.B. 2014.** Invasion complexities: the diverse impacts of nonnative species on amphibians. *Copeia*. 14: 611–632.
- Buffington, J.M. 2012.** Changes in channel morphology over human time scales. In: Church, M.; Biron, P.M.; Roy, A.G., eds. *Gravel-bed rivers: processes, tools, environments*. Chichester, United Kingdom: Wiley: 181–213.
- Bunn, J.T.; Montgomery, D.R. 2004.** Patterns of wood and sediment storage along debris-flow impacted headwater channels in old-growth and industrial forests of the western Olympic Mountains, Washington. In: Bennett, S.J.; Simon, A., eds. *Riparian vegetation and fluvial geomorphology*. Washington, DC: American Geophysical Union. 282 p.

- Burnett, K.M.; Miller, D.J. 2007.** Streamside policies for headwater channels: an example considering debris flows in the Oregon Coastal Province. *Forest Science*. 53: 239–253.
- Burnett, K.M.; Reeves, G.H.; Miller, D.J.; Clarke, S.; Vance-Borland, K.; Christiansen, K. 2007.** Distribution of salmon-habitat potential relative to landscape characteristics and implications for conservation. *Ecological Applications*. 17: 66–80.
- Burroughs, E.R.; Thomas, B.R. 1977.** Declining root strength in Douglas-fir after felling as a factor in slope stability. Res. Pap. INT-190. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station. 27 p.
- Burton, J.I.; Ganio, L.M.; Puettmann, K.J. 2014.** Multiscale spatial controls of understory vegetation in Douglas-fir–western hemlock forests of western Oregon, USA. *Ecosphere*. 5(12): art151.
- Bylhouwer, B.; Ianson, D.; Kohfeld, K. 2013.** Changes in the onset and intensity of wind-driven upwelling and downwelling along the North American Pacific coast. *Journal of Geophysical Research Oceans*. 118: 2565–2580.
- Byrne, C.J.; Poole, R.; Dillane, A.; Rogan, G.; Whalen, K.F. 2004.** Temporal and environmental influences on the variation in sea trout (*Salmo trutta* L.) smolt migration in the Burrishoole system in the west of Ireland from 1971–2000. *Fisheries Research*. 66: 85–94.
- Carah, J.K.; Blencowe, C.C.; Wright, D.W.; Bolton, L.A. 2014.** Low-cost restoration techniques for rapidly increasing wood cover in coastal coho salmon streams. *North American Journal of Fisheries Management*. 34: 1003–1013.
- Caraher, D. L.; Zack, A.C.; Stage, A.R. 1999.** Scales and ecosystem management. In: Szaro, R.C.; Johnson, N.C.; Sexton, W.T.; Malick, A.J., eds. *Ecological stewardship: a common reference for ecosystem management*, vol. 2. Oxford, United Kingdom: Elsevier Science: 343–352.
- Carey, A.B. 2003.** Biocomplexity and restoration of biodiversity in temperate coniferous forest: inducing spatial heterogeneity with variable-density thinning. *Forestry*. 76: 127–136.
- Carey, A.B.; Curtis, R.O. 1996.** Conservation of biodiversity: a useful paradigm for forest ecosystem management. *Wildlife Society Bulletin*. 24: 610–620.
- Carey, M.P.; Sanderson, B.L.; Friesen, T.A.; Barnas, K.A.; Olden, J.D. 2011.** Smallmouth bass in the Pacific Northwest: a threat to native species; a benefit for anglers. *Reviews in Fisheries Science*. 19: 305–315.
- Carlisle, D.M.; Hawkins, C.P.; Meador, M.R.; Potapova, M.; Falcone, J. 2008.** Biological assessments of Appalachian streams based on predictive models for fish, macroinvertebrate, and diatom assemblages. *Journal of the North American Benthological Society*. 27: 16–37.
- Carlisle, D.M.; Meador, M.R.; Short, T.M.; Tate, C.M.; Gurtz, M.E.; Bryant, W.L.; Falcone, J.A.; Woodside, M.D. 2013.** The quality of our Nation’s waters—ecological health in the nation’s streams, 1993–2005. Circular 1391. Reston, VA: U.S. Department of the Interior, Geological Survey. <https://pubs.usgs.gov/circ/1391/>. (8 August 2017).
- Castelle, A.J.; Johnson, A.W.; Conolly, C. 1994.** Wetland and stream buffer size requirements—a review. *Journal of Environmental Quality*. 23: 878–882.
- Cederholm, C.J.; Johnson, D.H.; Bilby, R.E.; Dominguez, L.G.; Garrett, A.M.; Graeber, W.H.; Greda, E.L.; Kunze, M.D.; Marcot, B.G.; Palmisano, J.F.; Plotnikoff, R.W.; Pearcey, W.G.; Simenstad, C.A.; Trotter, P.C. 2001.** Pacific salmon and wildlife—ecological contexts, relationships, and implications for management. In: Johnson, D.H.; O’Neil, T.A., eds. *Wildlife-habitat relationships in Oregon and Washington*. Corvallis, OR: Oregon State University Press: 628–684.

- Center for Invasive Species and Ecosystem Health. 2017.** EDDMapS: early detection & distribution mapping system. Athens, GA: University of Georgia, Center for Invasive Species and Ecosystem Health. <https://www.eddmaps.org/>. (18 April 2017).
- Chan, S.S.; Larson, D.J.; Maas-Hebner, K.G.; Emmingham, W.H.; Johnston, S.R.; Mikowski, D.A. 2006.** Overstory and understory development in thinned and underplanted Oregon Coast Range Douglas-fir stands. *Canadian Journal of Forest Research*. 36: 2696–2711.
- Chelgren, N.D.; Dunham, J.B. 2015.** Connectivity and conditional models of access and abundance of species in stream networks. *Ecological Applications*. 25: 1357–1372.
- Chen, J. 1991.** Edge effects: microclimatic pattern and biological responses in old-growth Douglas-fir forests. Seattle, WA: University of Washington, Seattle. 174 p. Ph.D. dissertation.
- Chen, J.; Franklin, J.F.; Spies, T.A. 1993.** Contrasting microclimates among clearcut, edge, and interior of old-growth Douglas-fir forest. *Agricultural and Forest Meteorology*. 63: 219–237.
- Chiaramonte, L.V.; Ray, R.A.; Corum, R.A.; Soto, T.; Hallett, S.L.; Bartholomew, J.L. 2016.** Klamath River thermal refuge provides juvenile salmon reduced exposure to the parasite *Ceratonova shasta*. *Transactions of the American Fisheries Society*. 145: 810–820.
- Chin, A.; An, L.; Florsheim, J.L.; Laurencio, L.R.; Marston, R.A.; Solverson, A.P.; Simon, G.L.; Stinson, E.; Wohl, E. 2016.** Investigating feedbacks in human–landscape systems: lessons following a wildfire in Colorado, USA. *Geomorphology*. 252: 40–50.
- Chin, A.; Florsheim, J.L.; Wohl, E.; Collins, B.D. 2014.** Feedbacks in human-landscape systems. *Environmental Management*. 53: 28–41.
- Christensen, V.; Trites, A.W. 2011.** Predation on Fraser River sockeye salmon. In: Cohen, B.I., commissioner. The uncertain future of Fraser River Sockeye. Final report. Vol. 2. Causes of the decline. Ottawa, ON: Government of Canada, Minister of Public Works and Government Services: 186–188. http://publications.gc.ca/collections/collection_2012/bcp-pco/CP32-93-2012-2-eng.pdf. (5 September 2017).
- Cissel, J.H.; Swanson, F.J.; Weisberg, P.J. 1999.** Landscape management using historical fire regimes: Blue River, Oregon. *Ecological Applications*. 9: 1217–1231.
- Claeson, S.M.; Bisson, P.A. 2013.** Passive reestablishment of riparian vegetation following removal of invasive knotweed (*Polygonum*). *Invasive Plant Science and Management*. 6: 208–218.
- Clark, W.A.V.; Avery, K.L. 1976.** The effects of data aggregation in statistical analysis. *Geographical Analysis*. 8: 428–438. doi:10.1111/j.1538-4632.1976.tb00549.x.
- Clarke, A.; Mac Nally, R.; Bond, N.; Lake, P.S. 2008.** Macroinvertebrate diversity in headwater streams: a review. *Freshwater Biology*. 53: 1707–1721.
- Clarkin, K.; Connor, A.; Furniss, M.J.; Gubernick, B.; Love, M.; Moynan, K.; Wilson Musser, S. 2005.** National Inventory and Assessment Procedure—for identifying barriers to aquatic organism passage at road-stream crossings. 7700–Transportation Mgmt. San Dimas, CA: U.S. Department of Agriculture, Forest Service, National Technology and Development Program. 61 p.
- Clayton, D.R.; Olson, D.H. 2007.** Conservation assessment for the Oregon slender salamander (*Batrachoseps wrighti*). Version 1.0. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, and U.S. Department of the Interior, Bureau of Land Management, Oregon office, Special Status Species Program. <http://www.fs.fed.us/r6/sfpnw/issssp/documents/planning-docs/ca-ha-bawr-2007-05-17.doc>. (1 September 2017).

- Climate Impacts Group. 2009.** The Washington climate change impacts assessment: evaluating Washington's future in a changing climate. Full report Seattle, WA: University of Washington, College of the Environment. <https://cig.uw.edu/resources/special-reports/2009-washington-climate-change-impact-assessment/>. (27 November 2017).
- Collier, K.J. 2009.** Linking multimetric and multivariate approaches to assess the ecological condition of streams. *Environmental Monitoring and Assessment*. 157: 113–124.
- Connor, W.P.; Burge, H.L.; Yearsley, J.R.; Bjornn, T.C. 2003.** Influence of flow and temperature on survival of wild subyearling fall Chinook salmon in the Snake River. *North American Journal of Fisheries Management*. 23: 362–375.
- Conservation Institute. 2013.** Top 100 invasive species. <http://www.conservationinstitute.org/the-top-100-invasive-species/>. (29 March 2017).
- Cooley, S.R.; Lucey, N.; Kite-Powell, H.L.; Doney, S.C. 2012.** Nutrition and income from mollusks today imply vulnerability to ocean acidification tomorrow. *Fish and Fisheries*. 13: 182–215.
- Copeland, T.; Vendetti, D.V. 2009.** Contribution of three life history types to smolt production in a Chinook salmon (*Oncorhynchus tshawytscha*) population. *Canadian Journal of Fisheries and Aquatic Sciences*. 66: 1658–1665.
- Cornwell, W.K.; Cornelissen, J.H.; Amatangelo, K.; Dorrepaal, E.; Eviner, V.T.; Godoy, O.; Hobbie, S.E.; Hoorens, B.; Kurokawa, H.; Pérez-Harguindeguy, N.; Quesed, H.M. 2008.** Plant species traits are the predominant control on litter decomposition rates within biomes worldwide. *Ecology Letters*. 11: 1065–1071.
- Coulthard, T.J.; Kirkby, M.J.; Macklin, M.G. 2000.** Modelling geomorphic response to environmental change in an upland catchment. *Hydrological Processes*. 14: 2031–2045.
- Craig, R.K. 2010.** 'Stationarity is dead'—long live transformation: five principles for climate change adaptation law. *Harvard Environmental Law Review*. 34: 9–75.
- Columbia River Basin Aquatic Invasive Species [CRBAIS]. 2011.** <http://crbais.psmfc.org/about-us>. (6 July 2017).
- Cristea, N.; Janisch, J. 2007.** Analysis of factors affecting stream temperature to assist the development of hardwood conversion guidelines for small forest landowners. Report. Olympia, WA: Washington Department of Ecology.
- Croke, J.C.; Hairsine, P.B. 2006.** Sediment delivery in managed forests: a review. *Environmental Reviews*. 14: 59–87.
- Crozier, L.G.; Hendry, A.P.; Lawson, P.W.; Quinn, T.P.; Mantua, N.J.; Battin, J.; Shaw, R.G.; Huey, R.B. 2008.** Potential response to climate change in organisms with complex life histories: evolution and plasticity in Pacific salmon. *Evolutionary Applications*. 1: 252–270.
- Crozier, L.G.; Zabel, R.W. 2006.** Climate impacts at multiple scales: evidence for differential population responses in juvenile Chinook salmon. *Journal of Animal Ecology*. 75: 1100–1109.
- Cummins, K.W.; Wilzbach, M.A.; Gates, D.M.; Perry, J.B.; Taliaferro, W.B. 1989.** Shredders and riparian vegetation. *BioScience*. 39: 24–30.
- Daigle, P. 2010.** A summary of the environmental impacts of roads, management responses, and research gaps: a literature review. *BC Journal of Ecosystems and Management*. 10: 65–89.
- Dale, V.H.; Brown, S.; Haeuber, R.A.; Hobbs, N.T.; Huntly, N.; Naiman, R.J.; Riebsame, W.E.; Turner, M.G.; Valone, T.J. 2000.** Ecological principles and guidelines for managing the use of land. *Ecological Applications*. 10: 639–670.

- Dale, V.H.; Joyce, L.A.; McNulty, S.; Neilson, R.P.; Ayres, M.P.; Flannigan, M.D.; Hanson, P.J.; Irland, L.C.; Lugo, A.E.; Peterson, C.J.; Simberloff, D.; Swanson, F.J.; Stocks, B.J.; Wotton, B.M. 2001.** Climate change and forest disturbances. *BioScience*. 51: 723–734.
- D’Amore, D.V.; Oken, K.L.; Herendeen, P.A.; Steel, E.A.; Hennon, P.E. 2015.** Carbon accretion in unthinned and thinned young-growth forest stands of the Alaskan perhumid coastal temperate rainforest. *Carbon Balance and Management*. 10: 25. <https://link.springer.com/content/pdf/10.1186%2Fs13021-015-0035-4.pdf>. (1 September 2017).
- Danehy, R.J.; Bilby, R.E.; Langshaw, R.B.; Evans, D.M.; Turner, T.R.; Floyd, W.C.; Schoenholtz, S.H.; Duke, S.D. 2012.** Biological and water quality responses to hydrologic disturbances in third-order forested streams. *Ecohydrology*. 5: 90–98.
- Datry, T.; Larned, S.T.; Tockner, K. 2014.** Intermittent rivers: a challenge for freshwater ecology. *BioScience*. 64: 229–235.
- Desjardins, E.; Barker, G.; Lindo, Z.; Dieleman, C.; Dussault, A.C. 2015.** Promoting resilience. *The Quarterly Review of Biology*. 90: 147–165.
- Dieterich, M.; Anderson, N.H. 2000.** The invertebrate fauna of summer-dry streams in western Oregon. *Archives für Hydrobiologie*. 147: 273–295.
- Downing, J.A.; Cole, J.J.; Duarte, C.A.; Middelburg, J.J.; Melack, J.M.; Prairie, Y.T.; Kortelainen, P.; Striegl, R.G.; McDowell, W.H.; Tranvik, L.J. 2012.** Global abundance and size distribution of streams and rivers. *Inland Waters*. 2: 229–236.
- Dunham, J.B.; Pilliod, D.S.; Young, M. 2004.** Assessing the consequences of nonnative trout in headwater ecosystems. *Fisheries*. 29: 18–26.
- Ebersole, J.L.; Colvin, M.E.; Wigington, P.J., Jr.; Leibowitz, S.G.; Baker, J.P.; Church, M.R.; Compton, J.E.; Cairns, M.A. 2009.** Hierarchical modeling of late-summer weight and summer abundance of juvenile coho salmon across a stream network. *Transactions of the American Fisheries Society*. 138: 1138–1156.
- Ebersole, J.L.; Liss, W.J.; Frissell, C.A. 2003.** Thermal heterogeneity, stream channel morphology and salmonid abundance in northeast Oregon streams. *Canadian Journal of Fisheries and Aquatic Sciences*. 60: 1266–1280.
- Ebersole, J.L.; Wigington, P.J.; Baker, J.P.; Cairns, M.A.; Church, M.R.; Compton, J.E.; Leibowitz, S.G.; Miller, B.; Hansen, B. 2006.** Juvenile coho salmon growth and survival across stream network seasonal habitats. *Transactions of the American Fisheries Society*. 135: 1681–1697.
- Ebersole, J.L.; Wigington, P.J., Jr.; Leibowitz, S.G.; Comeleo, R.L.; Van Sickle, J. 2015.** Predicting the occurrence of cold-water patches at intermittent and ephemeral tributary confluences with warm rivers. *Freshwater Science*. 34: 111–124.
- Edwards, N. 1999.** Anatoxin: also known as Very Fast Death Factor. Brighton, United Kingdom: University of Sussex at Brighton. <http://www.chm.bris.ac.uk/motm/antx/antx.htm>. (7 September 2017).
- Elliott, J.M. 1994.** Quantitative ecology and the brown trout. New York: Oxford University Press. 286 p.
- Elliott, J.M.; Hurley, M.A. 1998.** An individual-based model for predicting the emergence period of sea trout in a Lake District stream. *Journal of Fish Biology*. 53: 414–433.
- Elsner, M.M.; Cuo, L.; Voisin, N.; Deems, J.S.; Hamlet, A.F.; Vano, J.A.; Mickelson, K.E.B.; Lee, S.Y.; Lettenmaier, D.P. 2010.** Implications of 21st century climate change for the hydrology of Washington state. *Climatic Change*. 102: 225–260. doi:10.1007/s10584-010-9855-0. (1 September 2017).

- Everest, F.H. 1975.** Ecology and management of summer steelhead in the Rogue River. Final report, project AFS31. Fishery Research Report No. 7. Oregon State Game Commission. Corvallis, OR: Oregon State University. 48 p.
- Everest, F.H.; Reeves, G.H. 2007.** Riparian and aquatic habitats of the Pacific Northwest and southeast Alaska: ecology, management history, and potential management strategies. Gen. Tech. Rep. PNW-GTR-692. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 130 p.
- Fagen, R. 1988.** Long-term trends in maximum size of sport-caught Chinook salmon (*Oncorhynchus tshawytscha*): a data-analytic approach to weights of first-prize fish in four southeastern Alaska salmon derbies. Fisheries Research. 6: 125–134.
- Fausch, K.D. 2010.** Preface: A renaissance in stream fish ecology. In: Gido, K.B.; Jackson, D.A., eds. Community ecology of stream fishes: concepts, approaches, and techniques. American Fisheries Society Symposium 73. Bethesda, MD: American Fisheries Society: 199–206.
- Fausch, K.D.; Torgersen, C.E.; Baxter, C.V.; Li, H.W. 2002.** Landscapes to riverscapes: bridging the gap between research and conservation of stream fishes a continuous view of the river is needed to understand how processes interacting among scales set the context for stream fishes and their habitat. BioScience. 52: 483–498.
- Fetherston, K.L.; Naiman, R.J.; Bilby, R.E. 1995.** Large woody debris, physical process, and riparian forest development in montane river networks of the Pacific Northwest. Geomorphology. 13: 133–144.
- Fleming, I.A.; Gross, M.R. 1990.** Latitudinal clines: a trade-off between egg number and size in Pacific salmon. Ecology. 71: 1–11.
- Fleming, I.A.; Lamberg, A.; Jonsson, B. 1997.** Effects of early experience on the reproductive performance of Atlantic salmon. Behavioural Ecology. 8: 470–480.
- Flitcroft, R.; Burnett, K.; Christiansen, K. 2013.** A simple model that identifies potential effects of sea-level rise on estuarine and estuary-ecotone habitat locations for salmonids in Oregon, USA. Environmental Management. 52: 196–208.
- Flitcroft, R.; Hansen, B.; Capurso, J.; Christiansen, K. 2016b.** Coverage of aquatic invasive risk assessment in USFS Region 6. Report to U.S. Forest Service Region 6. Corvallis, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. https://www.fs.fed.us/pnw/lwm/aem/docs/flitcroft/2016_flitcroft_et_al_coverage_ais_assessment_region6.pdf. (31 March 2017).
- Flitcroft, R.L.; Falke, J.A.; Reeves, G.H.; Hessburg, P.F.; McNyset, K.M.; Benda, L.E. 2016a.** Wildfire may increase habitat quality for spring Chinook salmon in the Wenatchee River subbasin, WA, USA. Forest Ecology and Management. 359: 126–140.
- Forest Ecosystem Management Assessment Team [FEMAT]. 1993.** Forest ecosystem management: an ecological, economic, and social assessment. Report of the Forest Ecosystem Management Assessment Team. Portland, OR: U.S. Department of Agriculture; U.S. Department of the Interior [and others]. [Irregular pagination].
- Forman, R.T.T. 2003.** Road ecology: science and solutions. Washington, DC: Island Press. 504 p.
- Forman, R.T.T.; Alexander, L.E. 1998.** Roads and their major ecological effects. Annual Review of Ecology and Systematics. 29: 207–231.
- Foster, A.D.; Ziegltrum, J. 2013.** Riparian-associated gastropods in western Washington: Community composition and the effects of forest management. Northwest Science. 87: 243–256.
- Fox, M.; Bolton, S. 2007.** A regional and geomorphic reference for quantities and volumes of instream wood in unmanaged forested basins of Washington State. North American Journal of Fisheries Management. 27: 342–359.

- Frady, C.; Johnson, S.; Li, J. 2007.** Stream macro-invertebrate community responses as legacies of forest harvest at the H.J. Andrews Experimental Forest, Oregon. *Forest Science*. 53: 281–293.
- Francis, R.C.; Hare, S.R. 1997.** Regime scale climate forcing of salmon populations in the Northeast Pacific—some new thoughts and findings. In: Emmett, R.L.; Schiewe, M.H., eds. *Estuarine and ocean survival of northeastern Pacific salmon: proceedings of the workshop*. Tech. Memo. NMFS-NWFSC-29. Seattle, WA: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Northwest Fisheries Science Center: 113–128.
- Francis, R.C.; Sibley, T.H. 1991.** Climate change and fisheries: what are the real issues? *The Northwest Environmental Journal*. 7: 295–307.
- Franklin, J.F.; Johnson, K.N. 2012.** A restoration framework for federal forests in the Pacific Northwest. *Journal of Forestry*. 110: 429–439.
- Franklin, J.F.; Spies, T.A.; Swanson, F.J. 2017.** Setting the stage: vegetation ecology and dynamics. In: Olson, D.H.; Van Horne, B., eds. *People, forests, and change: lessons from the Pacific Northwest*. Washington, DC: Island Press: 16–32. Chapter 2.
- Franklin, J.F.; Spies, T.A.; Van Pelt, R.; Carey, A.B.; Thornburgh, D.A.; Berg, D.R.; Lindenmayer, D.B.; Harmon, M.E.; Keeton, W.S.; Shaw, D.C.; Bible, K. 2002.** Disturbances and structural development of natural forest ecosystems with silvicultural implications, using Douglas-fir forests as an example. *Forest Ecology and Management*. 155: 399–423.
- Frissell, C.A.; Liss, W.J.; Gresswell, R.E.; Nawa, R.K.; Ebersole, J.L. 1997.** A resource in crisis: changing the measure of salmon. In: Stouder, D.J.; Bisson, P.A.; Naiman, R.J., eds. *Pacific salmon and their ecosystems: status and future options*. New York: Chapman and Hall: 411–446.
- Frissell, C.A.; Poff, N.L.; Jensen, M.E. 2001.** Assessment of biotic patterns in freshwater ecosystems. In: Jensen, M.E.; Bourgeron, P.S., eds. *A guidebook for integrated ecological assessments*. New York: Springer: 390–403.
- Gallo, K.; Lanigan, S.H.; Eldred, P.; Gordon, S.N.; Moyer, C. 2005.** Northwest Forest Plan—the first 10 years (1994–2003): preliminary assessment of the condition of watersheds. Gen. Tech. Rep. PNW-GTR-647. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 133 p.
- Garcia, R.A.; Cabeza, M.; Rahbek, C.; Araújo, M.B. 2014.** Multiple dimensions of climate change and their implications for biodiversity. *Science*. 344(6183): 1247579. doi:10.1126/science.1247579.
- Garner, S.R.; Heath, J.W.; Neff, B.D. 2009.** Egg consumption in mature Pacific salmon (*Oncorhynchus* spp.). *Canadian Journal of Fisheries and Aquatic Sciences*. 66: 1546–1553.
- Gende, S.M.; Quinn, T.P.; Willson, M.F.; Heintz, R.; Scott, T.M. 2004.** Magnitude and fate of salmon-derived nutrients and energy in a coastal stream ecosystem. *Journal of Freshwater Ecology*. 19: 149–160.
- Gibson, R.J.; Haedrich, R.L.; Wernerheim, C.M. 2005.** Loss of fish habitat as a consequence of inappropriately constructed stream crossings. *Fisheries*. 30: 10–17.
- Gienapp, P.; Teplitsky, C.; Aho, J.S.; Mills, J.A.; Mertä, J. 2008.** Climate change and evolution: disentangling environmental and genetic responses. *Molecular Ecology*. 17: 167–176.
- Goetz, J.N.; Guthrie, R.H.; Brenning, A. 2015.** Forest harvesting is associated with increased landslide activity during an extreme rainstorm on Vancouver Island, Canada. *Natural Hazards Earth Systems Science*. 15: 1311–1330.

- Gomi, T.; Sidle, R.C.; Richardson, J.S. 2002.** Understanding processes and downstream linkages of headwater systems: headwaters differ from downstream reaches by their close coupling to hillslope processes, more temporal and spatial variation, and their need for different means of protection from land use. *BioScience*. 52: 905–916.
- González-Moreno, P.; Diez, J.M.; Richardson, D.M.; Vilà, M. 2015.** Beyond climate: disturbance niche shifts in invasive species. *Global Ecology and Biogeography*. 24: 360–370.
- Goode, J.R.; Buffington, J.M.; Tonina, D.; Isaak, D.J.; Thurow, R.F.; Wenger, S.; Nagel, D.; Luce, C.; Tetzlaff, D.; Soulsby, C. 2013.** Potential effects of climate change on streambed scour and risks to salmonid survival in snow-dominated mountain basins. *Hydrological Processes*. 27: 750–765.
- Gordon, S. 2014.** Use of EMDS in conservation and management planning for watersheds. In: Reynolds, K.M.; Hessburg, P.F.; Bourgeron, P.S., eds. *Making transparent environmental decisions: applications of the ecosystem management system*. Heidelberg, Germany: Springer: 63–91.
- Gordon, S.N.; Gallo, K. 2011.** Structuring expert input for a knowledge-based approach to watershed condition assessment for the Northwest Forest Plan, USA. *Environmental Monitoring and Assessment*. 172: 643–661.
- Grant, E.H.; Miller, D.A.W.; Schmidt, B.R.; Adams, M.J.; Amburgey, S.M.; Chambert, T.; Cruickshank, S.S.; Fisher, R.N.; Green, D.M.; Hossack, B.R.; Johnson, P.T.H.; Joseph, M.; Rittenhouse, T.A.G.; Ryan, M.E.; Waddle, J.H.; Walls, S.C.; Bailey, L.L.; Fellers, G.M.; Gorman, T.A.; Ray, A.M.; Pilliod, D.S.; Price, S.J.; Saenz, D.; Sadinski, W.; Muths, E. 2016.** Quantitative evidence for the effects of multiple drivers on continental-scale amphibian declines. *Nature Scientific Reports*. 6: 25625. doi:10.1038/srep25625. (5 September 2017).
- Grant, G.E.; Lewis, S.L.; Swanson, F.J.; Cissel, J.H.; McDonnell, J.J. 2008.** Effects of forest practices on peak flows and consequent channel response: a state-of-science report for western Oregon and Washington. Gen. Tech. Rep. PNW-GTR-760. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 76 p.
- Grantham, T.E.; Fesenmyer, K.A.; Peek, R.; Holmes, E.; Quiñones, R.M.; Bell, A.; Santos, N.; Howard, J.K.; Viers, J.H.; Moyle, P.B. 2017.** Missing the boat on freshwater fish conservation in California. *Conservation Letters*. 10: 77–85.
- Gray, M.J.; Duffus, A.L.J.; Haman, K.H.; Harris, R.N.; Allender, M.C.; Thompson, T.A.; Christman, M.R.; Sacerdote-Velat, A.; Sprague, L.A.; Williams, J.M.; Miller, D.L. 2017.** Pathogen surveillance in herpetofaunal populations: guidance on study design, sample collection, biosecurity, and intervention strategies. *Herpetological Review*. 48: 334–351.
- Grebmeier, J.M.; Overland, J.E.; Moore, S.E.; Farley, E.V.; Carmack, E.C.; Cooper, L.W.; Frey, K.E.; Helle, J.H.; McLaughlin, F.A.; McNutt, S.L. 2006.** A major ecosystem shift in the northern Bering Sea. *Science*. 311: 1461–1464.
- Greene, C.M.; Hall, J.E.; Guilbault, K.R.; Quinn, T.P. 2009.** Improved variability of populations with diverse life-history portfolios. *Biology Letters*. 6: 382–386. doi:10.1098/rsbl.2009.0780.
- Gregory, S.V. 1997.** Riparian management in the 21st century. In: Kohm, K.A.; Franklin, J.F., eds. *Creating a forestry for the 21st century*. Washington, DC: Island Press: 69–85.
- Gregory, S.V.; Meleason, M.A.; Sobota, D.J. 2003.** Modeling the dynamics of wood in streams and rivers. *American Fisheries Society Symposium*. 37: 315–335.
- Groom J.D.; Dent, L.; Madsen, L.; Fleuret, J. 2011.** Response of western Oregon (USA) stream temperatures to contemporary forest management. *Forest Ecology and Management*. 262: 1618–1629.

- Gruendike, P.; Lanigan, S. 2008.** Aquatic and Riparian Effectiveness Monitoring Program: invasive species report, 2007 field season. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region; U.S. Department of the Interior, Bureau of Land Management, Oregon state office. <https://www.fs.fed.us/r6/reo/monitoring/reports/InvasiveSpeciesAREMP2007.pdf>. (31 May 2017).
- Gruendike, P.; Lanigan, S. 2009.** Aquatic and Riparian Effectiveness Monitoring Program: invasive species report, 2008 field season. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region; U.S. Department of the Interior, Bureau of Land Management, Oregon state office. <https://www.fs.fed.us/r6/reo/monitoring/reports/AREMP%20Invasive%20Report%202008%208x11.pdf>. (31 May 2017).
- Gucinski, H.; Furniss, M.J.; Ziemer, R.R.; Brookes, M.H. 2001.** Forest roads: a synthesis of scientific information. Gen. Tech. Rep. PNW-GTR-509. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 103 p.
- Guthrie, R.H. 2002.** The effects of logging on frequency and distribution of landslides in three watersheds on Vancouver Island, British Columbia. *Geomorphology*. 43: 273–292.
- Hagar, J.C.; McComb, W.C.; Emmingham, W.H. 1996.** Bird communities in commercially thinned and unthinned Douglas-fir stands of western Oregon. *Wildlife Society Bulletin*. 24: 353–366.
- Halofsky, J.E.; Peterson, D.L., eds. 2017.** Climate change vulnerability and adaptation in the Blue Mountains. Gen. Tech. Rep. PNW-GTR-939. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 331 p.
- Halofsky, J.E.; Peterson, D.L.; O'Halloran, K.A.; Hawkins Hoffman, C., eds. 2011.** Adapting to climate change at Olympic National Forest and Olympic National Park. Gen. Tech. Rep. PNW-GTR-844. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 130 p.
- Hamer, A.J.; Land, S.J.; Mahony, M.J. 2002.** The role of introduced mosquitofish (*Gambusi holbrooki*) in excluding the native green and golden bell frog (*Litoria aurea*) from original habitats in south-eastern Australia. *Oecologia*. 132: 445–452.
- Hamlet, A.F.; Lettenmaier, D.P. 2007.** Effects of 20th century warming and climate variability on flood risk in the western U.S. *Water Resources Research*. 43: W06427. doi:10.1029/2006WR005099.
- Hamlet, A.F.; McGuire Elsner, M.; Mauger, G.S.; Lee, S.-Y.; Tohver, I.; Norheim, R.A. 2013.** An overview of the Columbia Basin Climate Change Scenarios Project: approach, methods, and summary of key results. *Atmosphere-Ocean*. 51: 392–415.
- Hamlet, A.F.; Mote, P.W.; Clark, M.P.; Lettenmaier, D.P. 2005.** Effects of temperature and precipitation variability on snowpack trends in the western United States. *Journal of Climate*. 18: 4545–4561.
- Hance, D.J.; Ganio, L.M.; Burtnett, K.M.; Ebersole, J.L. 2016.** Basin-scale variation in the spatial pattern of fall movement of juvenile Coho salmon in the West Fork Smith River, Oregon. *Transactions of the American Fisheries Society*. 145: 1018–1034.
- Hankin, D.G.; McKelvey, R. 1985.** Comment on fecundity of Chinook salmon and its relevance to life history theory. *Canadian Journal of Fisheries and Aquatic Sciences*. 42: 393–395.
- Harmon, M.E.; Franklin, J.F.; Swanson, F.J.; Sollins, P.; Gregory, S.V.; Lattin, J.D.; Anderson, N.H.; Cline, S.P.; Aumen, N.G.; Sedell, J.R.; Lienkaemper, G.W. 1986.** Ecology of coarse woody debris in temperate ecosystems. *Advances in Ecological Research*. 15: 133–302.
- Harr, R.D. 1981.** Some characteristics and consequences of snowmelt during rainfall in western Oregon. *Journal of Hydrology*. 53: 277–304.
- Harr, R.D. 1982.** Fog drip in the Bull Run municipal watershed, Oregon. *Water Resources Bulletin*. 18: 785–789.

- Harr, R.D. 1986.** Effects of clearcutting on rain-on-snow runoff in western Oregon: a new look at old studies. *Water Resources Research*. 22: 1095–1100.
- Harris, H.E.; Baxter, C.V.; Davis, J.M. 2015.** Debris flows amplify effects of wildfire on magnitude and composition of tributary subsidies to mainstem habitats. *Freshwater Science*. 34: 1457–1467.
- Hassan, M.A.; Hogan, D.L.; Bird, S.A.; May, C.L.; Gomi, T.; Campbell, D. 2005.** Spatial and temporal dynamics of wood in headwater streams of the Pacific Northwest. *Journal of the American Water Resources Association*. 41: 899–919.
- Havlick, D. 2002.** No place distant: roads and motorized recreation on America's public lands. Washington, DC: Island Press. 297 p.
- Hayes, M.P.; Jennings, M.R. 2005.** Bullfrog: *Rana catesbeiana*. In: Jones, L.L.C.; Leonard, W.P.; Olson, D.H., eds. *Amphibians of the Pacific Northwest*. Seattle, WA: Seattle Audubon Society: 190–193.
- Healey, M.C.; Heard, W.R. 1984.** Inter- and intra-population variation in the fecundity of Chinook salmon (*Oncorhynchus tshawytscha*) and its relevance to life history theory. *Canadian Journal of Fisheries and Aquatic Sciences*. 41: 476–483.
- Healey, M.C.; Prince, A. 1995.** Scales of variation in life history tactics of Pacific salmon and the conservation of phenotype and genotype. *American Fisheries Society Symposium*. 17: 176–184.
- Heckman, C.W. 1999.** The encroachment of exotic herbaceous plants into the Olympic National Forest. *Northwest Science*. 73: 264–276.
- Helfield, J.M.; Naiman, R.J. 2001.** Effects of salmon-derived nitrogen on riparian forest growth and implications for stream productivity. *Ecology*. 82: 2403–2409.
- Helle, J.H.; Martinson, E.C.; Eggers, D.M.; Gritsenko, O. 2007.** Influence of salmon abundance and ocean conditions on body size of Pacific salmon. *North Pacific Anadromous Fish Commission Bulletin*. 4: 289–4298.
- Hendriks, I.E.; Duarte, C.M.; Álvarez, M. 2010.** Vulnerability of marine biodiversity to ocean acidification: a meta-analysis. *Estuarine, Coastal and Shelf Science*. 86: 157–164.
- Heredia, N.; Roper, B.; Gillespie, N.; Roghair, C. 2016.** Technical guide for field practitioners: understanding and monitoring aquatic organism passage at road-stream crossings. Technical Report TR-101. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, National Stream & Aquatic Ecology Center. 35 p.
- Herlihy, A.T.; Paulsen, S.G.; Van Sickle, J.; Stoddard, J.L.; Hawkins, C.P.; Yuan, L.L. 2008.** Striving for consistency in a national assessment: the challenges of applying a reference-condition approach at a continental scale. *Journal of the North American Benthological Society*. 27: 860–877.
- Hessburg, P.F.; Agee, J.K. 2003.** An environmental narrative of Inland Northwest United States forests. *Forest Ecology and Management*. 178: 23–59.
- Hessburg, P.F.; Agee, J.K.; Franklin, J.F. 2005.** Dry forests and wildland fires of the inland Northwest USA: contrasting the landscape ecology of the pre-settlement and modern eras. *Forest Ecology and Management*. 211: 117–139.
- Hibbs, D.E.; Giordano, P.A. 1996.** Vegetation characteristics of alder-dominated riparian buffer strips in the Oregon Coast Range. *Northwest Science*. 70: 213–222.
- Hiers, J.K.; Jackson, S.T.; Hobbs, R.J.; Bernhardt, E.S.; Valentine, L.E. 2016.** The precision problem in conservation and restoration. *Trends in Ecology & Evolution*. 31: 820–830. <https://doi.org/10.1016/j.tree.2016.08.001>.

- Hinke, J.T.; Foley, D.G.; Wilson, C.; Watters, G.M. 2005.** Persistent habitat use by Chinook salmon *Oncorhynchus tshawytscha* in the coastal ocean. *Marine Ecology Progress Series*. 304: 207–220.
- Hobbs, R.J.; Higgs, E.; Harris, J.A. 2009.** Novel ecosystems: implications for conservation and restoration. *Trends in Ecology & Evolution*. 24: 599–605.
- Hodgeson, S.; Quinn, T.P.; Hilborn, R.; Francis, R.C.; Rogers, D.E. 2006.** Marine and freshwater climatic factors affecting interannual variation in the timing of return migration to fresh water of sockeye salmon (*Oncorhynchus nerka*). *Fisheries Oceanography*. 15: 1–24.
- Hoffman, R.L.; Dunham, J.B.; Hansen, B.P., eds. 2012.** Aquatic organism passage at road-stream crossings—synthesis and guidelines for effectiveness monitoring. Open-File Report 2012-1090. Reston, VA: U.S. Department of the Interior, Geological Survey. 64 p. <https://pubs.usgs.gov/of/2012/1090/pdf/ofr20121090.pdf>. (5 September 2017).
- Hogan, D.L.; Bird, S.A.; Hassan, M.L. 1998.** Spatial and temporal evolution of small coast gravel-bed streams: the influence of forest management on channel morphology and fish habitat. In: Klingman, P.C.; Beschta, R.L.; Komar, P.D.; Bradley, J.B., eds. *Gravel-bed rivers in the environment*. Proceeding of the 4th International Gravel-Bed Rivers workshop (GBR-IV). Highland Branch, CO: Water Resources Publications: 365–392.
- Hohler, D.; Sedell, J.; Olson, D. 2001.** Aquatic conservation strategy. Northwest Forest Plan research synthesis. Gen. Tech. Rep. PNW-GTR-498. In: Haynes, R.W.; Perez, G.E., tech. eds. *Northwest Forest Plan research synthesis*. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 30–39.
- Holling, C.S.; Meffe, G.K. 1996.** Command and control and the pathology of natural resource management. *Conservation Biology*. 10: 328–337.
- Holtby, L.B. 1988.** Effects of logging on stream temperatures in Carnation Creek British Columbia, and associated impacts on the coho salmon (*Oncorhynchus kisutch*). *Canadian Journal of Fisheries and Aquatic Sciences*. 45: 502–515.
- Holtby, L.B.; Scrivener, J.C. 1989.** Observed and simulated effects of climatic variability, clear-cut logging and fishing on the numbers of chum salmon (*Oncorhynchus keta*) and coho salmon (*O. kisutch*) returning to Carnation Creek, British Columbia. In: Levings, C.D.; Holtby, L.B.; Henderson, M.A., eds. *Proceedings of the national workshop on effects of habitat alterations on salmonid stocks*. Canadian Special Publication of Fisheries and Aquatic Sciences. 105: 62–81.
- Howell, P.J. 2006.** Effects of wildfire and subsequent hydrologic events on fish distribution and abundance in tributaries of North Fork John Day River. *North American Journal of Fisheries Management*. 26: 983–994.
- Hubler, S. 2009.** High level indicators of Oregon's forested streams. Hillsboro, OR: Oregon Department of Environmental Quality, Laboratory and Environmental Assessment Division. 79 p. <http://www.oregon.gov/deq/FilterDocs/10-LAB-003.pdf>. (28 August 2017).
- Hunter, M.L. 2005.** A mesofilter conservation strategy to complement fine and coarse filters. *Conservation Biology*. 19: 1025–1029.
- Hyde, E.J.; Simons, T.R. 2001.** Sampling plethodontid salamanders: sources of variability. *Journal of Wildlife Management*. 4: 624–632.
- Hylander, K.; Jonsson, B.G.; Nilsson, C. 2002.** Evaluating buffer strips along boreal streams using bryophytes as indicators. *Ecological Applications*. 12: 797–806.

Independent Multidisciplinary Science Team [IMST].

1999. Recovery of wild salmonids in western Oregon forests: Oregon Forest Practices Act rules and the measures in the Oregon Plan for Salmon and Watersheds. Tech. Report 1999-1 to the Oregon Plan for Salmon and Watersheds, Governor's Natural Resources Office, Salem, OR. Corvallis, OR: Oregon State University. http://www.krisweb.com/biblio/gen_ognro_imst_1999_1.pdf. (5 September 2017).

Intergovernmental Panel on Climate Change [IPCC].

2007. Climate change 2007: the physical science basis. Contribution of Working Group I to the fourth assessment report of the Intergovernmental Panel on Climate Change. Geneva, Switzerland. <http://www.slvwd.com/agendas/Full/2007/06-07-07/Item%2010b.pdf>. (5 September 2017).

Intergovernmental Panel on Climate Change [IPCC].

2014. Climate change 2014: synthesis report. Contribution of Working Groups I, II and III to the fifth assessment report of the Intergovernmental Panel on Climate Change. Geneva, Switzerland. 151 p. http://www.ipcc.ch/pdf/assessment-report/ar5/syr/AR5_SYR_FINAL_All_Topics.pdf. (5 September 2017).

Irvine, K.M.; Miller, S.W.; Al-Chokhachy, R.K.; Archer,

E.K.; Roper, B.B.; Kershner, J.L. 2015. Empirical evaluation of the conceptual model underpinning a regional aquatic long-term monitoring program using causal modelling. *Ecological Indicators*. 50: 8–23.

Isaak, D.J.; Luce, C.H.; Rieman, B.E.; Nagel, D.E.;

Peterson, E.E.; Horan, D.L.; Parkes, S.; Chandler, G.L. 2010. Effects of climate change and wildfire on stream temperatures and salmonid thermal habitat in a mountain river network. *Ecological Applications*. 20: 1350–1371.

Isaak, D.J.; Wenger, S.J.; Peterson, E.E.; Ver Hoef, J.M.;

Hostetler, S.; Luce, C.H.; Dunham, J.B.; Kershner, J.; Roper, B.B.; Nagel, D. 2011. NorWeST: an interagency stream temperature database and model for the northwest United States. U.S. Fish and Wildlife Service, Great Northern Landscape Conservation Cooperative Grant, Project. <https://www.fs.fed.us/rm/boise/AWAE/projects/NorWeST.html>. (5 December 2017).

Isaak, D.J.; Wollrab, S.; Horan, D.; Chandler, G.

2012. Climate change effects on stream and river temperatures across the northwest US from 1980–2009 and implications for salmonid fishes. *Climatic Change*. 113: 499–524.

Isaak, D.J.; Young, M.K.; Luce, C.H.; Hostetler, S.W.;

Wenger, S.J.; Peterson, E.E.; Ver Hoef, J.M.; Groce, M.C.; Horan, D.L.; Nagel, D.E. 2016. Slow climate velocities of mountain streams portend their role as refugia for cold-water biodiversity. *Proceedings of the National Academy of Sciences of the United States of America*. 113: 4374–4379

Isaak, D.J.; Young, M.K.; Nagel, D.E.; Horan, D.L.;

Groce, M.C. 2015. The cold-water climate shield: delineating refugia for preserving salmonid fishes through the 21st century. *Global Change Biology*. 21: 2540–2553.

Independent Scientific Advisory Board [ISAB].

2007. Climate change impacts on Columbia River Basin Fish and Wildlife. ISAB 2007-2. Portland, OR: Northwest Power and Conservation Council. <https://www.fws.gov/oregonfwo/ExternalAffairs/ClimateChange/Documents/ISAB%202007-2%20Climate%20Change.pdf>. (6 September 2017).

Independent Scientific Advisory Board [ISAB]. 2012.

Developing a broader scientific foundation for river restoration: Columbia River food webs. *Proceedings of the National Academy of Sciences of the United States of America*. 109: 21201–21207.

- Jackson, S.T.; Betancourt, J.L.; Booth, R.K.; Gray, S.T. 2009.** Ecology and the ratchet of events: climate variability, niche dimensions, and species distributions. *Proceedings of the National Academy of Sciences of the United States of America*. 106(Suppl. 2): 19685–19692.
- Jakob, M. 2000.** The impacts of logging on landslide activity at Clayoquot Sound, British Columbia. *Catena*. 38: 279–300.
- Janisch, J.E.; Wondzell, S.M.; Ehinger, W.J. 2012.** Headwater stream temperature: interpreting response after logging, with and without riparian buffers, Washington, USA. *Forest Ecology and Management*. 270: 302–313.
- Johnson, K.N.; Franklin, J.F.; Thomas, J.W.; Gordon, J. 1991.** Alternatives for management of late-successional forests of the Pacific Northwest. A report to the Agriculture Committee and the Merchant Marine and Fisheries Committee of the U.S. House of Representatives. [Place of publication unknown]: Scientific Panel on Late-Successional Forest Ecosystems (U.S.). 59 p.
- Johnson, S.C.; Blaylock, R.B.; Elphick, J.; Hyatt, K.D. 1996.** Disease induced by the sea louse (*Lepeophtheirus salmonis*) (Copepoda: Caligidae) in wild sockeye salmon (*Oncorhynchus nerka*) stocks of the Alberni Inlet, British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences*. 53: 2888–2897.
- Johnson, S.L. 2004.** Factors influencing stream temperatures in small streams: substrate effects and a shading experiment. *Canadian Journal of Fisheries and Aquatic Sciences*. 61: 913–923.
- Johnson, S.L.; Swanson, F.J.; Grant, G.E.; Wondzell, S.M. 2000.** Riparian forest disturbances by a mountain flood: the influence of floated wood. *Hydrological Process*. 14: 3031–3050.
- Johnson, S.P.; Schindler, D.E. 2009.** Trophic ecology of Pacific salmon (*Oncorhynchus* spp.) in the ocean: a synthesis of stable isotope research. *Ecological Research*. 24: 855–863.
- Johnstone, J.A.; Dawson, T.E. 2010.** Climatic context and ecological implications of summer fog decline in the coast redwood region. *Proceedings of the National Academy of Sciences of the United States of America*. 107: 4533–4538.
- Jones, J.A.; Grant, G.E. 2001.** Comment on “Peak flow responses to clearcutting and roads in small and large basins, western Cascades, Oregon: a second opinion” by R.B. Thomas and W.F. Megahan. *Water Resources Research*. 37: 175–178.
- Jones, J.A.; Swanson, F.J.; Wemple, B.C.; Snyder, K.U. 2000.** Effects of roads on hydrology, geomorphology, and disturbance patches in stream networks. *Conservation Biology*. 14: 76–85.
- Jonsson, B.; Jonsson, N. 2009.** A review of the likely effects of climate change on anadromous Atlantic salmon *Salmo salar* and brown trout *Salmo trutta*, with particular reference to water temperature and flow. *Journal of Fish Biology*. 75: 2381–2447.
- Joyce, L.A.; Blate, G.M.; McNulty, S.G.; Millar, C.I.; Moser, S.; Neilson, R.P.; Peterson, D.L. 2009.** Managing for multiple resources under climate change: national forests. *Environmental Management*. 44: 1022–1032.
- Junk, W.J.; Bayley, P.B.; Sparks, R.E. 1989.** The flood pulse concept in river-floodplain systems. In: Dodge, D.P., ed. *Proceedings of the International Large River symposium (LARS)*. Canadian Special Publication of Fisheries and Aquatic Sciences 106. Ottawa, ON: Government of Canada, Department of Fisheries and Oceans: 110–127. <http://www.dfo-mpo.gc.ca/Library/111846.pdf>. (6 September 2017).

- Justice, C.; White, S.M.; McCullough, D.A.; Graves, D.S.; Blanchard, M.R. 2017.** Can stream and riparian restoration offset climate change impacts to salmon populations? *Journal of Environmental Management*. 188: 212–227.
- Kail, J.; Arle, J.; Jahrig, S.C. 2012.** Limiting factors and thresholds for macroinvertebrate assemblages in European rivers: empirical evidence from three datasets on water quality, catchment urbanization, and river restoration. *Ecological Indicators*. 18: 63–72
- Karr, J.R.; Chu, E.W. 1998.** Restoring life in running waters: better biological monitoring. Washington, DC: Island Press. 206 p.
- Kats, L.B.; Ferrer, R.P. 2003.** Alien predators and amphibian declines: review of two decades of science and the transition to conservation. *Diversity and Distributions*. 9: 99–110.
- Kelsey, K.A.; West, S.D. 1998.** Riparian wildlife. In: Bilby, R.E.; Naiman, R.J., eds. *River ecology and management: lessons from the Pacific coastal ecoregion*. New York: Springer-Verlag: 235–260.
- Kennedy, R.E.; Yang, Z.; Cohen, W.B. 2010.** Detecting trends in forest disturbance and recovery using yearly Landsat time series: 1. LandTrendr-temporal segmentation algorithms. *Remote Sensing of Environment*. 114: 2897–2910.
- Kershner, J.L. 1997.** Setting riparian/aquatic restoration objectives within a watershed context. *Restoration Ecology*. 5(4S): 15–24.
- Keyes, C.R.; Teraoka, E.K. 2014.** Structure and composition of old-growth and unmanaged second-growth riparian forests at Redwood National Park, USA. *Forests*. 5: 256–268.
- Kiffney, P.M.; Richardson, J.S.; Feller, M.C. 2000.** Fluvial and epilithic organic matter dynamics in headwater streams of southwestern British Columbia, Canada. *Archiv für Hydrobiologie*. 149: 109–129.
- Kinziger, A.P.; Nakamoto, R.J.; Harvey, B.C. 2014.** Local-scale invasion pathways and small founder numbers in introduced Sacramento pikeminnow (*Ptchocheilus grandis*). *Conservation Genetics*. 15: 1–9.
- Kline, M.; Cahoon, B. 2010.** Protecting river corridors in Vermont. *Journal of the American Water Resources Association*. 46: 227–236.
- Klos, P.E.; Link, T.E.; Abatzoglou, J.T. 2014.** Extent of the rain-snow transition zone in the western U.S. under historic and projected climate. *Geophysics Research Letters*. 41: 4560–4568.
- Knapp, R.A.; Matthews, K.R. 2000.** Non-native fish introductions and the decline of the mountain yellow-legged frog from within protected areas. *Conservation Biology*. 14: 428–438.
- Kominoski, J.S.; Marczak, L.B.; Richardson, J.S. 2011.** Riparian forest composition affects stream litter decomposition despite similar microbial and invertebrate communities. *Ecology*. 92: 151–159.
- Kominoski, J.S.; Pringle, C.M. 2009.** Resource–consumer diversity: testing the effects of leaf litter species diversity on stream macroinvertebrate communities. *Freshwater Biology*. 54: 1461–1473.
- Kondolf, G.M.; Boulton, A.J.; O’Daniel, S.; Poole, G.C.; Rahel, F.J.; Stanley, E.H.; Wohl, E.; Bång, A.; Carlstrom, J.; Cristoni, C.; Huber, H. 2006.** Process-based ecological river restoration: visualizing three-dimensional connectivity and dynamic vectors to recover lost linkages. *Ecology and Society*. 11(2): art 5.
- Kondolf, G.M.; Montgomery, D.R.; Piégay, H.; Schmitt, L. 2003.** Geomorphic classification of rivers and streams. In: Kondolf, G.M.; Piégay, H., eds. *Tools in fluvial geomorphology*. New York: John Wiley & Sons: 171–204.

- Kovach, R.P.; Muhlfeld, C.C.; Wade, A.A.; Hand, B.K.; Whited, D.C.; DeHaan, P.W.; Al-Chokhachy, R.; Luikart, G. 2015.** Genetic diversity is related to climatic variation and vulnerability in threatened bull trout. *Global Change Biology*. 21: 2510–2524. doi:10.1111/gcb.12850.
- Kreutzweiser, D.P.; Capell, S.S.; Holmes, S.B. 2009.** Stream temperature responses to partial-harvest logging in riparian buffers of boreal mixedwood forest watersheds. *Canadian Journal of Forest Research*. 39: 497–506.
- Kuehne, L.M.; Olden, J.D.; Strecker, A.L.; Lawler, J.J.; Theobald, D.M. 2017.** Past, present, and future of ecological integrity assessment for fresh waters. *Frontiers in Ecology and the Environment*. 15: 197–205. doi:10.1002/fee.1483.
- Kuglerová, L.; Ågren, A.; Jansson, R.; Laudon, H. 2014.** Towards optimizing riparian buffer zones: ecological and biogeochemical implications for forest management. *Forest Ecology and Management*. 334: 74–84.
- Lam, J.; Chan, S.; Hansen, G.; Chapman, J.; Miller, J.; Carlton, J.; Boatner, R.; Cooper, R.; Kight, P.; Siemens, T.; Martin, K.; Corcoran, J.. 2015.** Key aquatic invasive species watch: Japanese tsunami marine debris in the eastern Pacific. ORESU-H-15-002. Corvallis, OR: Oregon Sea Grant. 12 p. <http://seagrant.oregonstate.edu/sgpubs/key-aquatic-invasive-species-watch-japanese-tsunami-marine-debris-eastern-pacific>. (30 March 2017).
- Lancaster, S.T.; Casebeer, N.E. 2007.** Sediment storage and evacuation in headwater valleys at the transition between debris-flow and fluvial processes. *Geology*. 35: 1027–1030.
- Lancaster, S.T.; Hayes, S.K.; Grant, G.E. 2003.** Effects of wood on debris flow run-out in small mountain watersheds: *Water Resources Research*. 39(6): 1168. doi:10.1029/2001WR001227.
- Lang, D.W.; Reeves, G.H.; Hall, J.D.; Wipfli, M.S. 2006.** The influence of fall-spawning coho salmon (*Oncorhynchus kisutch*) on growth and production of juvenile coho salmon rearing in beaver ponds on the Copper River Delta, Alaska. *Canadian Journal of Fisheries and Aquatic Sciences*. 63: 917–930.
- Lanigan, S.H.; Gordon, S.N.; Eldred, P.; Isley, M.; Wilcox, S.; Moyer, C.; Andersen, H. 2012.** Northwest Forest Plan—the first 15 years (1994–2008): watershed condition status and trend. Gen. Tech. Rep. PNW-GTR-856. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 155 p.
- Lapointe, M.; Eaton, B.; Driscoll, S.; Latulippe, C. 2000.** Modelling the probability of salmonid egg pocket scour due to floods. *Canadian Journal of Fisheries and Aquatic Sciences*. 57: 1120–1130.
- Latta, G.; Temesgen, H.; Adams, D.; Barrett, T. 2010.** Analysis of potential impacts of climate change on forests of the United States Pacific Northwest. *Forest Ecology and Management*. 259: 720–729.
- Latterell, J.J.; Naiman, R.J. 2007.** Sources and dynamics of large logs in a temperate floodplain river. *Ecological Applications*. 17: 1127–1141.
- LaVigne, H.R.; Huges, R.M.; Wildman, R.C.; Gregory, S.V.; Herlihy, A.T. 2008.** Summer distribution and species richness of non-native fishes in the mainstem Willamette River, Oregon, 1944–2006. *Northwest Science*. 82: 83–93.
- Lawrence, D.J.; Stewart-Koster, B.; Olden, J.D.; Ruesch, A.S.; Torgersen, C.E.; Lawler, J.J.; Butcher, D.P.; Crown, J.K. 2014.** The interactive effects of climate change, riparian management, and a nonnative predator on stream-rearing salmon. *Ecological Applications*. 24: 895–912.
- Lawson, P.W. 1993.** Cycles in ocean productivity, trends in habitat quality, and the restoration of salmon runs in Oregon. *Fisheries*. 18(8): 6–10.

- Leach, J.A.; Olson, D.H.; Anderson, P.D.; Eskelson, B.N.I. 2016.** Spatial and seasonal variability of forested headwater stream temperatures in western Oregon, USA. *Aquatic Sciences*. 79: 291–307. doi:10.1007/s00027-016-0497-9.
- Ledwith, T.S. 1996.** The effects of buffer strip width on air temperature and relative humidity in a stream riparian zone. Arcata, CA: Humboldt State University. 89 p. M.S. thesis.
- Lee, D.C.; Sedell, J.R.; Rieman, B.E.; Thurow, R.F.; Williams, J.E. 1997.** Broad-scale assessment of aquatic species and habitats. In: Quigley, T.M.; Arbelbide, S.J., tech. eds. *An assessment of ecosystem components in the interior Columbia Basin and portions of the Klamath and Great Basins*. Gen. Tech. Rep. PNW-GTR-405. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 1057–1496.
- Lee, P.; Smyth, C.; Boutin, S. 2004.** Quantitative review of riparian buffer width guidelines from Canada and the United States. *Journal of Environmental Management*. 70: 170–180.
- Leigh, C.; Boulton, A.J.; Courtwright, J.L.; Fritz, K.; May, C.L.; Walker, R.H.; Datry, T. 2016.** Ecological research and management of intermittent rivers: an historical review and future directions. *Freshwater Biology*. 61: 1181–1199.
- Leinenbach, P.; McFadden, G.; Torgersen, C. 2013.** Effects of riparian management strategies on stream temperature. Science Review Team Temperature Subgroup. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region. 22 p. On file with: U.S. Forest Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis, OR 97331.
- Leppi, J.C.; DeLuca, T.H.; Harrar, S.W.; Running, S.W. 2012.** Impacts of climate change on August stream discharge in the Central Rocky Mountains. *Climatic Change*. 112: 997–1014.
- Leppi, J.C.; Rinella, D.J.; Wilson, R.R.; Loya, W.M. 2014.** Linking climate change projections for an Alaskan watershed to future coho salmon production. *Global Change Biology*. 20: 1808–1820.
- Leroy, C.J.; Marks, J.C. 2006.** Litter quality, stream characteristics and litter diversity influence decomposition rates and macroinvertebrates. *Freshwater Biology*. 51: 605–617.
- Li, H.W.; Lamberti, G.A.; Pearsons, T.N.; Tait, C.K.; Li, J.L.; Buckhouse, J.C. 1994.** Cumulative effects of riparian disturbances along high desert trout streams of the John Day Basin, Oregon. *Transactions of the American Fisheries Society*. 123: 627–640.
- Lisle, T.E. 2002.** How much dead wood in channels is enough? In: Laudenslayer, W.F., Jr.; Shea, P.J.; Valentine, B.E.; Weatherspoon, C.P.; Lisle, T.E., eds. *Proceedings of the symposium on the ecology and management of dead wood in western forests*. Gen. Tech. Rep. PSW-GTR-181. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 58–93.
- Lisle, T.; Cummins, K.; Madej, M.A. 2007.** An examination of references for ecosystems in a watershed context: results of a scientific pulse in Redwood National and State Parks, California. In: Furniss, M.J.; Clifton, C.F.; Ronnenberg, K.L., eds. *Advancing the fundamental sciences: proceedings of the Forest Service National Earth Sciences conference*. Gen. Tech. Rep. PNW-GTR-689. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 118–130. Vol. 1.
- Liss, W.J.; Stanford, J.A.; Lichatowich, J.A.; Williams, R.N.; Coutant, C.C.; Mundy, P.R.; Whitney, R.R. 2006.** A foundation for restoration. In: Williams, R.N., ed. *Return to the river: strategies for salmon restoration in the Columbia River Basin*. Burlington, MA: Elsevier Academic Press: 51–98.

- Louhi, P.; Mykrä, H.; Paavola, R.; Huusko, A.; Vehanen, T.; Mäki-Petäys, A.; Muotka, T. 2011.** Twenty years of stream restoration in Finland: little response by benthic macroinvertebrate communities. *Ecological Applications*. 21: 1950–1961.
- Loukas, A.; Quick, M.C. 1999.** The effect of climate change on floods in British Columbia. *Nordic Hydrology*. 30: 231–256.
- Luce, C.; Morgan, P.; Dwire, K.; Isaak, D.; Holden, Z.; Rieman, B. 2012.** Climate change, forests, fire, water, and fish: building resilient landscapes, streams, and managers. Gen. Tech. Rep. RMRS-GTR-290. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 207 p.
- Luce, C.; Staab, B.; Kramer, M.; Wenger, S.; Isaak, D.; McConnell, C. 2014b.** Sensitivity of summer stream temperatures to climate variability in the Pacific Northwest. *Water Resources Research*. 50: 3428–3443.
- Luce, C.H.; Holden, Z.A. 2009.** Declining annual streamflow distributions in the Pacific Northwest United States, 1948–2006. *Geophysical Research Letters*. 36: L16401. doi:10.1029/2009GL039407.
- Luce, C.H.; Lopez-Burgos, V.; Holden, Z. 2014a.** Sensitivity of snowpack storage to precipitation and temperature using spatial and temporal analog models. *Water Resources Research*. 50: 9447–9462. doi:10.1002/2013WR014844.
- Lunetta, R.S.; Cosentino, B.L.; Montgomery, D.R.; Beamer, E.M.; Beechie, T.J. 1997.** GIS-based evaluation of salmon habitat in the Pacific Northwest. *Photogrammetric Engineering and Remote Sensing*. 63: 1219–1229.
- Lusardi, R.A.; Bogan, M.T.; Moyle, P.B.; Dahlgren, R.A. 2016.** Environment shapes invertebrate assemblage structure differences between volcanic springfed and runoff rivers in northern California. *Freshwater Science*. 35: 1010–1022.
- Lyons, J.K.; Beschta, R.L. 1983.** Land use, floods, and channel changes: Upper Middle Fork Willamette River, Oregon (1936–1980). *Water Resources Research*. 19: 463–471.
- MacDonald, L.H.; Coe, D. 2007.** Influence of headwater streams on downstream reaches in forested areas. *Forest Science*. 53: 148–168.
- MacDonald, L.H.; Coe, D.B.R. 2008.** Road sediment production and delivery: processes and management. In: *Proceedings of the First World Landslide Forum, International Programme on Landslides and International Strategy for Disaster Reduction*. Tokyo: United Nations University: 381–384. <http://ucanr.edu/sites/forestry/files/138028.pdf>. (6 September 2017).
- Macdonald, J.S.; MacIsaac, E.A.; Herunter, H.E. 2003.** The effect of variable-retention riparian buffer zones on water temperatures in small headwater streams in sub-boreal forest ecosystems of British Columbia. *Canadian Journal of Forest Research*. 33: 1371–1382.
- Madej, M.A.; Currens, C.; Ozaki, V.; Yee, J.; Anderson, D. 2006.** Assessing possible thermal rearing restrictions for juvenile coho salmon (*Oncorhynchus kisutch*) through thermal infrared imaging and in-stream monitoring, Redwood Creek, California. *Canadian Journal of Fisheries and Aquatic Sciences*. 63: 1384–1396.
- Malard, F.; Tockner, K.; Dole-Oliver, M.-J.; Ward, J.V. 2002.** A landscape perspective of surface–subsurface hydrological exchanges in river corridors. *Freshwater Biology*. 47: 621–640.
- Mangel, M. 1994.** Climate change and salmonid life history variation. *Deep Sea Research Part II: Tropical Studies in Oceanography*. 41: 75–106.
- Mantua, N.J.; Hare, S.R.; Zhang, Y.; Wallace, J.M.; Francis, R.C. 1997.** A Pacific interdecadal climate oscillation with impacts on salmon production. *Bulletin of the American Meteorological Society*. 78: 1069–1079.

- Mantua, N.J.; Taylor, N.G.; Ruggerone, G.T.; Myers, K.W.; Preikshot, D.; Augerot, X.; Davis, N.D.; Dorner, B.; Hilborn, R.; Peterman, R.M.; Rand, P.; Schindler, D.; Stanford, J.; Walker, R.V.; Walters, C.J. 2009.** The salmon MALBEC project: a North Pacific-scale study to support salmon conservation planning. North Pacific Anadromous Fish Commission [NPAFC] Doc. 1060. Seattle, WA: University of Washington, School of Aquatic and Fishery Sciences. 49 p. <https://digital.lib.washington.edu/researchworks/bitstream/handle/1773/38391/2007-5.pdf?sequence=1>. (6 September 2017).
- Mantua, N.J.; Tohver, I.; Hamlet, A. 2010.** Climate change impacts on streamflow extremes and summertime stream temperature and their possible consequences for freshwater salmon habitat in Washington state. *Climatic Change*. 102: 187–223. doi:10.1007/s10584-010-9845-2.
- Marcot, B.G.; Thompson, M.P.; Runge, M.C.; Thompson, F.R.; McNulty, S.; Cleaves, D.; Tomosy, M.; Fisher, L.A.; Bliss, A. 2012.** Recent advances in applying decision science to managing national forests. *Forest Ecology and Management*. 285: 123–132.
- Marine, K.R.; Cech, J.J. 2004.** Effects of high temperature on growth, smoltification, and predator avoidance in juvenile Sacramento River Chinook salmon. *North American Journal of Fisheries Management*. 24: 198–210.
- Marks, D.; Kimball, J.; Tingey, D.; Link, T. 1998.** The sensitivity of snowmelt processes to climate conditions and forest cover during rain-on-snow: a case study of the 1996 Pacific Northwest flood. *Hydrological Processes*. 12: 1569–1587.
- Marris, E. 2010.** The new normal. *Conservation Magazine*. <http://conservationmagazine.org/2010/the-new-normal/>. (1 May 2015).
- Martel, A.; Blooi, M.; Adriaensen, C.; Van Rooij, P.; Beukema, W.; Fisher, M.C.; Farrer, R.A.; Schmidt, B.R.; Tobler, U.; Goka, K.; Lips, K.R.; Muletz, C.; Zamudio, K.R.; Bosch, J.; Lötters, S.; Wombwell, E.; Garner, T.W.J.; Cunningham, A.A.; Spitzen-van der Sluijs, A.; Salvidio, S.; Ducatelle, R.; Nishikawa, K.; Nguyen, T.T.; Kolby, J.E.; Van Bocxlaer, I.; Bossuyt, F.; Pasmans, F. 2014.** Recent introduction of a chytrid fungus endangers Western Palearctic salamanders. *Science*. 346: 630–631.
- Martel, A.; Spitzen-van der Sluijs, A.; Blooi, M.; Bert, W.; Ducatelle, R.; Fisher, M.C.; Woeltjes, A.; Bosman, W.; Chiers, K.; Bossuyt, F.; Pasmans, F. 2013.** *Batrachochytrium salamandrivorans* sp. nov. causes lethal chytridiomycosis in amphibians. *Proceedings of the National Academy of Sciences of the United States of America*. 110: 15325–15329.
- Martins, E.G.; Hinch, S.G.; Patterson, D.A.; Hague, M.J.; Cooke, S.J.; Miller, K.M.; Lapointe, M.F.; English, K.K.; Farrell, A.P. 2011.** Effects of river temperature and climate warming on stock-specific survival of adult migrating Fraser River sockeye salmon (*Oncorhynchus nerka*). *Global Change Biology*. 17: 99–114.
- May, C. 2007.** Sediment and wood routing in steep headwater streams: an overview of geomorphic processes and their topographic signatures. *Forest Science*. 53: 119–130.
- May, C.L.; Gresswell, R.E. 2003.** Large wood recruitment and redistribution in headwater streams of the Oregon Coast Range, U.S.A. *Canadian Journal of Forest Research*. 33: 1352–1362.
- May, C.L.; Gresswell, R.E. 2004.** Processes and rates of sediment and wood accumulation in headwater streams of the Oregon Coast Range, U.S.A. *Earth Surface Processes and Landforms*. 28: 409–424.

- May, C.L.; Lee, D.C. 2004.** The relationships among in-channel sediment storage, pool depth, and summer survival of juvenile salmonids in Oregon Coast Range streams. *North American Journal of Fisheries Management*. 24: 761–774.
- McCabe, G.J.; Lauren, E.H.; Clark, M.P. 2007.** Rain-on-snow events in the western United States. *Bulletin of the American Meteorological Society*. 88: 319–328.
- McCarthy, I.D.; Houlihan, D.F. 1997.** The effect of temperature on protein synthesis in fish: the possible consequences for wild Atlantic salmon (*Salmo salar*) stocks in Europe as a result of global warming. In: Wood, C.M.; McDonald, G., eds. *Global warming: implications for freshwater and marine fish*. Cambridge, United Kingdom: Cambridge University Press: 51–77.
- McConnaha, W.E.; Williams, R.N.; Lichatowich, J.A. 2006.** The problem of the Columbia River salmon. In: Williams, R.N., ed. *Return to the river: restoring salmon back to the Columbia River*. Burlington, MA: Elsevier Academic Press: 1–28. Chapter 1.
- McCullough, D.A. 1999.** A review and synthesis of effects of alterations to the water temperature regime on freshwater life stages of salmonids, with special reference to Chinook salmon. Report No. 910-R-99-010. Seattle, WA: U.S. Environmental Protection Agency. 291 p. <https://nepis.epa.gov/Exe/ZyPDF.cgi/P1004JG1.PDF?Dockey=P1004JG1.PDF>. (28 August 2017.)
- McDade, M.H.; Swanson, F.J.; McKee, W.A.; Franklin, J.F.; Sickle, J.V. 1990.** Source distances for coarse woody debris entering small streams in western Oregon and Washington. *Canadian Journal of Forest Research*. 20: 326–330.
- McElhany, P.; Ruckelshaus, M.H.; Ford, M.J.; Wainwright, T.C.; Bjorkstedt, E.P. 2000.** Viable salmonid populations and the recovery of evolutionarily significant units. Tech. Memo. NMFS-NWFSC-42. Seattle, WA: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Northwest Fisheries Science Center. 156 p.
- McIntyre, P.J.; Thorne, J.H.; Dolanc, C.R.; Flint, A.L.; Flint, L.E.; Kelly, M.; Ackerly, D.D. 2015.** Twentieth-century shifts in forest structure in California: denser forests, smaller trees, and increased dominance of oaks. *Proceedings of the National Academy of Sciences of the United States of America*. 112: 1458–1463.
- McKay, S.K.; Cooper, A.R.; Diebel, M.W.; Elkins, D.; Oldford, G.; Roghair, C.; Wieferich, D. 2016.** Informing watershed connectivity barrier prioritization decisions: a synthesis. *River Research and Applications*. 33: 847–862. doi:10.1002/rra.3021.
- McKean, J.; Tonina, D. 2013.** Bed stability in unconfined gravel bed mountain streams: with implications for salmon spawning viability in future climates. *Journal of Geophysical Research: Earth Surface*. 118: 1227–1240.
- McKinney, M.L. 2001.** Effects of human population, area, and time on non-native plant and fish diversity in the United States. *Biological Conservation*. 100: 243–252.
- McKinney, M.L.; Lockwood, J.L. 1999.** Biotic homogenizations: a few winners replacing many losers in the next mass extinction. *Trends in Ecology & Evolution*. 14: 450–453.
- Mellina, E.; Moore, R.D.; Hinch, S.G.; Macdonald, J.S.; Pearson, G. 2002.** Stream temperature responses to clearcut logging in British Columbia: the moderating influences of groundwater and headwater lakes. *Canadian Journal of Fisheries and Aquatic Sciences*. 59: 1886–1900.
- Menuz, D.R.; Kettenring, K.M. 2013.** The importance of roads, nutrients, and climate for invasive plant establishment in riparian areas in the northwestern United States. *Biological Invasions*. 15: 1601–1612.
- Meyer, J.L.; Strayer, D.L.; Wallace, J.B.; Eggert, S.L.; Helfman, G.S.; and Leonard, N.E. 2007.** The contribution of headwater streams to biodiversity in river networks. *Journal of the American Water Resources Association*. 43: 86–103.

- Miller, D.; Luce, C.; Benda, L. 2003.** Time, space, and episodicity of physical disturbance in streams. *Forest Ecology and Management* 178: 121–140. doi:10.1016/S0378-1127(03)00057-4.
- Miller, D.J.; Burnett, K.M. 2008.** A probabilistic model of debris-flow delivery to stream channels, demonstrated for the Coast Range of Oregon, USA. *Geomorphology* 94: 184–205.
- Miller, K.M.; Li, S.; Kaukinen, K.H.; Ginther, N.; Hammill, E.; Curtis, J.M.R.; Patterson, D.A.; Sierocinski, T.; Donnison, L.; Pavlidis, P.; Hinch, S.G.; Hruska, K.A.; Cooke, S.J.; English, K.K.; Farrell, A.P. 2011.** Genomic signatures predict migration and spawning failure in wild Canadian salmon. *Science* 331: 214–217.
- Miller, S.A.; Eldred, P.; Muldoon, A.; Anlauf-Dunn, K.; Stein, C.; Hubler, S.; Merrick, L.; Haxton, N.; Larson, C.; Rehn, A.; Ode, P.; Vander Laan, J. 2016.** A large-scale, multiagency approach to defining a reference network for Pacific Northwest streams. *Environmental Management* 58: 1091–1104. doi:10.1007/s00267-016-0739-6.
- Miller, S.A.; Gordon, S.N.; Eldred, P.; Beloin, R.M.; Wilcox, S.; Raggon, M.; Andersen, H.; Muldoon, A. 2017.** Northwest Forest Plan—the first 20 years (1994–2013): watershed condition status and trends. Gen. Tech. Rep. PNW-GTR-932. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 74 p. https://www.fs.fed.us/pnw/pubs/pnw_gtr932.pdf. (11 October 2017).
- Minshall, G.W.; Brock, J.T.; Varley, J.D. 1989.** Wildfires and Yellowstone’s stream ecosystems. *BioScience* 39: 707–715.
- Montgomery, D.R. 1999.** Process domains and the river continuum. *Journal of the American Water Resources Association* 35: 397–410.
- Montgomery, D.R. 2004.** Geology, geomorphology, and the restoration ecology of salmon. *GSA Today* 14(11): 4–12. doi: 10.1130/1052-5173(2004)014<4:GGATRE>2.0.CO;2.
- Montgomery, D.R.; Buffington, J.M. 1997.** Channel-reach morphology in mountain drainage basins. *Geological Society of America Bulletin* 109: 596–611.
- Montgomery, D.R.; Buffington, J.M.; Peterson, N.P.; Schuett-Hames, D.; Quinn, T.P. 1996.** Stream-bed scour, egg burial depths, and the influence of salmonid spawning on bed surface mobility and embryo survival. *Canadian Journal of Fisheries and Aquatic Sciences* 53: 1061–1070.
- Montgomery, D.R.; MacDonald, L.H. 2002.** Diagnostic approach to stream channel assessment and monitoring. *Journal of the American Water Resources Association* 38: 1–16.
- Montgomery, D.R.; Massong, T.M.; Hawley, S.C.S. 2003.** Influence of debris flows and log jams on the location of pools and alluvial channel reaches, Oregon Coast Range. *Geological Society of America Bulletin* 115: 78–88.
- Moore, J.N.; Harper, J.T.; Greenwood, M.C. 2007.** Significance of trends toward earlier snowmelt runoff, Columbia and Missouri Basin headwaters, western United States. *Geophysical Research Letters* 34(16): L16402. doi:10.1029/2007GL031022.
- Moore, J.W.; McClure, M.; Rogers, L.A.; Schindler, D.E. 2010.** Synchronization and portfolio performance of threatened salmon. *Conservation Letters* 3: 340–348.
- Moore, R.D.; Spittlehouse, D.L.; Story, A. 2005.** Riparian microclimate and stream temperature response to forest harvesting. *Journal of the American Water Resources Association* 41: 813–834.
- Moore, J.W.; Yeakel, J.D.; Peard, D.; Lough, J.; Beere, M. 2014.** Life-history diversity and its importance to population stability and persistence of a migratory fish: steelhead in two large North American watersheds. *Journal of Animal Ecology* 83: 1035–1046.

- Moore, K.M.S.; Gregory, S.V. 1988.** Response of young-of-the-year cutthroat trout to manipulation of habitat structure in a small stream. *Transactions of the American Fisheries Society*. 117: 162–170.
- Mote, P.W. 2003.** Trends in snow water equivalent in the Pacific Northwest and their climatic causes. *Geophysical Research Letters*. 30(12): 8015. doi:10.1029/2002GL016435.
- Mote, P.W.; Hamlet, A.F.; Clark, M.; Lettenmaier, D.P. 2005.** Declining mountain snowpack in western North America. *Bulletin of the American Meteorological Society*. 86: 39–49.
- Mote, P.W.; Salathé, E.P., Jr. 2010.** Future climate in the Pacific Northwest. *Climatic Change*. 102(1–2): 29–50. doi:10.1007/s10584-010-9848-z. http://www-k12.atmos.washington.edu/~salathe/papers/published/Mote_Salathe_2010.pdf. (6 September 2017).
- Moyle, P.B.; Kiernan, J.D.; Crain, P.K.; Quiñones, R.M. 2013.** Climate change vulnerability of native and alien freshwater fishes of California: a systematic assessment approach. *PLoS One*. 8(5): e63883. doi:10.1371/journal.pone.0063883.
- Muir, P.S.; Mattingly, R.L.; Tappeiner, J.C., II; Bailey, J.D.; Elliott, W.E.; Hagar, J.C.; Miller, J.C.; Peterson, E.B.; Starkey, E.E. 2002.** Managing for biodiversity in young Douglas-fir forests of western Oregon. *Biol. Science Rep. USGS/BRD/BSR—2002-0006*. Corvallis, OR: U.S. Department of the Interior, Geological Survey, Forest and Rangeland Ecosystem Science Center. 76 p. https://fresc.usgs.gov/products/papers/mang_bio.pdf. (6 September 2017).
- Mulder, B.S.; Noon, B.R.; Spies, T.A.; Raphael, M.G.; Palmer, C.J.; Olsen, A.R.; Reeves, G.H.; Welsh, H.H., tech. coords. 1999.** The strategy and design of the effectiveness monitoring program for the Northwest Forest Plan. *Gen. Tech. Rep. PNW-GTR-437*. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 138 p.
- Murcia, C.; Aronson, J.; Kattan, G.H.; Moreno-Mateos, D.; Dixon, K.; Simberloff, D. 2014.** A critique of the ‘novel ecosystem’ concept. *Trends in Ecology & Evolution*. 29: 548–553.
- Murphy, D.D.; Weiland, P.S.; Cummins, K.W. 2011.** A critical assessment of the use of surrogate species in conservation planning in the Sacramento-San Joaquin Delta, California (U.S.A.). *Conservation Biology*. 25: 873–878.
- Murphy, M.L.; Koski, K.V. 1989.** Input and depletion of woody debris in Alaska streams and implications for streamside management. *North American Journal of Fisheries Management*. 9: 427–436.
- Myers, K.W.; Aydin, K.Y.; Walker, R.V.; Fowler, S.; Dahlberg, M.L. 1996.** Known ocean ranges of stocks of Pacific salmon and steelhead as shown by tagging experiments, 1956–1995. *NPAFC Document 192*. Seattle, WA: University of Washington, School of Fisheries, Fisheries Research Institute. 228 p. <https://digital.lib.washington.edu/researchworks/bitstream/handle/1773/4234/9614.pdf?sequence=1>. (6 September 2017).
- Naiman, R.J.; Alldredge, J.R.; Beauchamp, D.A.; Bisson, P.A.; Congleton, J.; Henny, C.J.; Huntly, N.; Lamberson, R.; Levings, C.; Merrill, E.N.; Pearcy, W.G.; Rieman, B.E.; Ruggerone, G.T.; Scarnecchia, D.; Smouse, P.E.; Wood, C.C. 2012.** Developing a broader scientific foundation for river restoration: Columbia River food webs. *Proceedings of the National Academy of Sciences of the United States of America*. 109: 21201–21207.
- Naiman, R.J.; Bechtold, J.S.; Beechie, T.J.; Latterell, J.J.; Van Pelt, R. 2010.** A process-based view of floodplain forest patterns in coastal river valleys of the Pacific Northwest. *Ecosystems*. 13: 1–31.

- Naiman, R.J.; Beechie, T.J.; Benda, L.E.; Berg, D.R.; Bison, P.A.; MacDonald, L.H.; O'Connor, M.D.; Olson, P.L.; Steel, E.A. 1992.** Fundamental elements of ecologically healthy watersheds in the Pacific Northwest coastal ecoregion. In: Naiman, R.J., ed. Watershed management: balancing sustainability and environmental change. New York: Springer-Verlag: 127–188.
- Naiman, R.J.; Bilby, R.E.; Bisson, P.A. 2000.** Riparian ecology and management in the Pacific coastal rain forest. *BioScience*. 50: 996–1011.
- Naiman, R.J.; Latterell, J.J. 2005.** Principles for linking fish habitat to fisheries management and conservation. *Journal of Fish Biology*. 67(sB): 166–185.
- National Marine Fisheries Service [NMFS]. 1999.** The habitat approach: implementation of Section 7 of the Endangered Species Act for actions affecting the habitat of Pacific anadromous salmonids. Portland, OR: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Northwest Region, Habitat Conservation and Protected Resources Divisions. 13 p. http://www.westcoast.fisheries.noaa.gov/publications/reference_documents/esa_refs/habitatapproach_081999-2.pdf. (6 September 2017).
- National Research Council [NRC]. 1996.** Upstream: salmon and society in the Pacific Northwest. Washington, DC: National Academies Press. 472 p. <https://doi.org/10.17226/4976>. (6 September 2017).
- National Research Council [NRC]. 2000.** Riparian areas: functions and strategies for management. Washington, DC: National Academies Press. 444 p. <https://doi.org/10.17226/10327>. (6 September 2017).
- National Wildfire Coordinating Group [NWCG]. 2017.** Guide to preventing aquatic invasive species transport by wildland fire operations. PMS 444. https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/fseprd530817.pdf. (5 April 2017).
- Nauman, R.S.; Olson, D.H. 2004.** Surveys for terrestrial amphibians in Shasta County, California, with notes on the distribution of Shasta salamanders (*Hydromantes shastae*). *Northwestern Naturalist*. 85: 35–38.
- Naylor, B.J.; Mackereth, R.W.; Kreutzweiser, D.P.; Sibley, P.K. 2012.** Merging END concepts with protection of fish habitat and water quality in new direction for riparian forests in Ontario: a study of science guiding policy and practice. *Freshwater Science*. 31: 248–257.
- Neeson, T.M.; Ferris, M.C.; Diebel, M.W.; Doran, P.J.; O'Hanley, J.R.; McIntyre, P.B. 2015.** Enhancing ecosystem restoration efficiency through spatial and temporal coordination. *Proceedings of the National Academy of Sciences of the United States of America*. 112: 6236–6241.
- Neuheimer, A.B.; Taggart, C.T. 2007.** The growing degree-day and fish size-at-age: the overlooked metric. *Canadian Journal of Fisheries and Aquatic Sciences*. 64: 375–385.
- Neumann, J.E.; Emanuel, K.; Ravela, S.; Ludwig, L.; Kirshen, P.; Bosma, K.; Martinich, J. 2015.** Joint effects of storm surge and sea-level rise on US Coasts: new economic estimates of impacts, adaptation, and benefits of mitigation policy. *Climatic Change*. 129: 337–349. doi:10.1007/s10584-014-1304-z.
- Nickelson, T.E. 1986.** Influences of upwelling, ocean temperature, and smolt abundance on marine survival of coho salmon (*Oncorhynchus kisutch*) in the Oregon production area. *Canadian Journal of Fisheries and Aquatic Sciences*. 43: 527–535.
- Nierenberg, T.R.; Hibbs, D.E. 2000.** A characterization of unmanaged riparian areas in the central Coast Range of western Oregon. *Forest Ecology and Management*. 129: 195–206.
- Nolin, A.W.; Daly, C. 2006.** Mapping “at risk” snow in the Pacific Northwest. *Journal of Hydrometeorology*. 7: 1164–1171.

- Nonaka, E.; Spies, T.A.; Wimberly, M.C.; Ohmann, J.L. 2007.** Historical range of variability in live and dead wood biomass: a regional-scale simulation study. *Canadian Journal of Forest Research*. 37: 2349–2364.
- North, M.P.; Keeton, W.S. 2008.** Emulating natural disturbance regimes: an emerging approach for sustainable forest management. In: Laforzezza, R.; Chen, J.; Sanesi, G.; Crow, T.R. eds. *Patterns and processes in forest landscapes: multiple use and sustainable management*. Amsterdam: Springer Netherlands: 341–372.
- Noss, R.F. 1987.** From plant communities to landscapes in conservation inventories: a look at The Nature Conservancy (USA). *Biological Conservation*. 41: 11–37.
- Ogston, L.; Gidora, S.; Foy, M.; Rosenfeld, J. 2014.** Watershed-scale effectiveness of floodplain habitat restoration for juvenile coho salmon in the Chilliwack River, British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences*. 72: 479–490.
- Ohmann, J.; Gregory, M.; Henderson, E.; Roberts, H. 2011.** Mapping gradients of community composition with nearest-neighbour imputation: extending plot data for landscape analysis. *Journal of Vegetation Science*. 22: 660–676.
- Olden, J.D.; Poff, N.L.; Douglas, M.R.; Douglas, M.E.; Fausch, K.D. 2004.** Ecological and evolutionary consequences of biotic homogenization. *Trends in Ecology & Evolution*. 19: 18–24.
- Oliver, C.D. 1981.** Forest development in North America following major disturbances. *Forest Ecology and Management*. 3: 153–168.
- Olson, D.H.; Aanensen, D.M.; Ronnenberg, K.L.; Powell, C.I.; Walker, S.F.; Bielby, J.; Garner, T.W.J.; Weaver, G.; the Bd Mapping Group; Fisher, M.C. 2013.** Mapping the global emergence of *Batrachochytrium dendrobatidis*, the amphibian chytrid fungus. *PLoS One*. 8(2): e56802.
- Olson, D.H.; Anderson, P.D.; Frissell, C.A.; Welsh, H.H., Jr.; Bradford, D.F. 2007.** Biodiversity management approaches for stream–riparian areas: perspectives for Pacific Northwest headwater forests, microclimates, and amphibians. *Forest Ecology and Management*. 246: 81–107.
- Olson, D.H.; Burnett, K.M. 2009.** Design and management of linkage areas across headwater drainages to conserve biodiversity in forest ecosystems. *Forest Ecology and Management*. 258: S117–S126.
- Olson, D.H.; Burton, J.H. 2014.** Near-term effects of repeated-thinning with riparian buffers on headwater stream vertebrates and habitats in Oregon, USA. *Forests*. 5: 2703–2729. doi:10.3390/f5112703.
- Olson, D.H.; Johnson, S.L.; Anderson, P.D.; Penulana, B.E.; Dunham, J.B. 2017a.** Aquatic-riparian systems. In: Olson, D.H.; Van Horne, B., eds., *People, forests, and change: lessons from the Pacific Northwest*. Washington, DC: Island Press: 191–206. Chapter 14.
- Olson, D.H.; Kluber, M.R. 2014.** Plethodontid salamander distributions in managed forest headwaters in western Oregon, USA. *Herpetological Conservation and Biology*. 9: 76–96.
- Olson, D.H.; Leirness, J.B.; Cunningham, P.G.; Steel, E.A. 2014.** Riparian buffers and forest thinning: effects on headwater vertebrates 10 years after thinning. *Forest Ecology and Management*. 321: 81–93.
- Olson, D.H.; Rugger, C. 2007.** Preliminary study of the effects of headwater riparian reserves with upslope thinning on stream habitats and amphibians in western Oregon. *Forest Science*. 53: 331–342.
- Olson, D.H.; Van Horne, B., eds. 2017.** *People, forests, and change: lessons from the Pacific Northwest*. Washington, DC: Island Press. 331 p.

- Olson, D.H.; Van Horne, B.; Bormann, B.T.; Deal, R.L.; DeLuca, T.H. 2017b.** The future of human-forest ecosystem sustainability. In: Olson, D.H.; Van Horne, B., eds., *People, forests, and change: lessons from the Pacific Northwest*. Washington, DC: Island Press: 275–290. Chapter 19.
- Olson, D.H.; Weaver, G. 2007.** Vertebrate assemblages associated with headwater hydrology in western Oregon managed forests. *Forest Science*. 53: 343–355.
- O’Neal, K. 2002.** Effects of global warming on trout and salmon in U.S. streams. Washington, DC: Defenders of Wildlife. 46 p. http://www.defenders.org/publications/effects_of_global_warming_on_trout_and_salmon.pdf. (6 September 2017).
- O’Neill, R.V.; DeAngelis, D.L.; Allen, T.F.H.; Waide, J.B. 1986.** A hierarchical concept of ecosystems. *Monographs in Population Biology* 23. Princeton, NJ: Princeton University Press. 262 p.
- O’Neill, R.V.; Johnson, A.R.; King, A.W. 1989.** A hierarchical framework for the analysis of scale. *Landscape Ecology*. 3: 193–205.
- Oregon Department of Environmental Quality [ODEQ]. 2012.** Water quality assessment—Oregon’s 2012 integrated report assessment database and 303(d) list. <http://www.deq.state.or.us/wq/assessment/rpt2012/search.asp>. (13 October 2017).
- Oregon Invasive Species Council [OISC]. 2015.** 100 worst list. https://static1.squarespace.com/static/58740d57579fb3b4fa5ce66f/t/5891577c579fb38e735f0f83/1485920126222/OISC_top100_2015.pdf. (28 March 2017).
- Orr, J.C.; Fabry, V.J.; Aumont, O.; Bopp, L.; Doney, S.C.; Feely, R.A.; Gnanadesikan, A.; Gruber, N.; Ishida, A.; Joos, F.; Key, R.M.; Lindsay, K.; Maier-Reimer E.; Matear, R.; Monfray, P.; Mouchet, A.; Najjar, R.G.; Plattner, G.-K.; Rodgers, K.B.; Sabine, C.L.; Sarmiento, J.L.; Schlitzer, R.; Slater, R.D.; Totterdell, I.J.; Weirig, M.-F.; Yool, A. 2005.** Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*. 437: 681–686.
- Pabst, R.J.; Spies, T.A. 1999.** Composition and structure of unmanaged riparian forests in the coastal mountains of Oregon. *Canadian Journal of Forest Research*. 29: 1557–1573.
- Palmer, M.A.; Bernhardt, E.S.; Allan, J.D.; Lake, P.S.; Alexander, G.; Brooks, S.; Carr, J.; Clayton, S.; Dahm, C.N.; Follstad Shah, J.; Galat, D.L. 2005.** Standards for ecologically successful river restoration. *Journal of Applied Ecology*. 42: 208–217.
- Palmer, M.A.; Hondula, K.L.; Koch, B.J. 2014.** Ecological restoration of streams and rivers: shifting strategies and shifting goals. *Annual Review of Ecology, Evolution, and Systematics*. 45: 247–269.
- Pankhurst, N.W.; Purser, G.J.; van der Kraak, G.; Thomas, P.M.; Forteath, G.N.R. 1996.** Effect of holding temperature on ovulation, egg fertility, plasma levels of reproductive hormones and in vitro ovarian steroidogenesis in the rainbow trout *Oncorhynchus mykiss*. *Aquaculture*. 146: 277–290.
- Paoletti, D.J.; Olson, D.H.; Blaustein, A.R. 2011.** Responses of foothill yellow-legged frog (*Rana boylei*) larvae to an introduced predator. *Copeia*. 2011: 161–168.
- Patil, S.D.; Wigington, P.J.; Leibowitz, S.G.; Comeleo, R.L. 2014.** Use of hydrologic landscape classification to diagnose streamflow predictability in Oregon. *Journal of the American Water Resources Association*. 50: 762–776.

- Pearcy, W.G. 1992.** Ocean ecology of North Pacific salmonids. Seattle, WA: Washington Sea Grant Program, University of Washington Press. 179 p.
- Pearl, C.A.; Adams M.J.; McCreary, B. 2013.** Habitat and co-occurrence of native and invasive crayfish in the Pacific Northwest, USA. *Aquatic Invasions*. 8: 171–184.
- Pearson, S.F.; Manuwal, D.A. 2001.** Breeding bird response to riparian buffer width in managed Pacific Northwest Douglas-fir forests. *Ecological Applications*. 11: 840–853.
- Penaluna, B.E.; Olson, D.H.; Flitcroft, R.L.; Weber, M.A.; Bellmore, J.R.; Wondzell, S.M.; Dunham, J.B.; Johnson, S.L.; Reeves, G.H. 2016.** Aquatic biodiversity in forests: a weak link in ecosystem services resilience. *Biodiversity and Conservation*. doi:10.1007/s10531-016-1148-0.
- Pennell, T.; Raggon, M. 2016.** Aquatic and Riparian Effectiveness Monitoring Program: invasive species report, 2015 field season. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region; U.S. Department of the Interior, Bureau of Land Management, Oregon State office. <https://reo.gov/monitoring/reports/watershed/AREMP%20Aquatic%20Invasive%20Species%20Report%202015.pdf>. (27 April 2017).
- Perry, L.G.; Reynolds, L.V.; Beechie, T.J.; Collins, M.J.; Shafroth, P.B. 2015.** Incorporating climate change projections into riparian restoration planning and design. *Ecohydrology*. 8: 863–879.
- Perry, T.D.; Jones, J.A. 2017.** Summer streamflow deficits from regenerating Douglas-fir forest in the Pacific Northwest, USA. *Ecohydrology*. doi:10.1002/eco.1790.
- Petersen, J.H.; Kitchell, J.F. 2001.** Climate regimes and water temperature changes in the Columbia River: bioenergetic implications for predators of juvenile salmon. *Canadian Journal of Fisheries and Aquatic Sciences*. 58: 1831–1841.
- Peterson, E.E.; Ver Hoef, J.M.; Isaak, D.J.; Falke, J.A.; Fortin, M.J.; Jordan, C.E.; McNyset, K.; Monestiez, P.; Ruesch, A.S.; Sengupta, A.; Som, N. 2013.** Modelling dendritic ecological networks in space: an integrated network perspective. *Ecology Letters*. 16: 707–719.
- Peterson, N.P. 1982a.** Immigration of juvenile coho salmon (*Oncorhynchus kisutch*) into riverine ponds. *Canadian Journal of Fisheries and Aquatic Sciences*. 39: 1308–1310.
- Peterson, N.P. 1982b.** Population characteristics of juvenile coho salmon (*Oncorhynchus kisutch*) overwintering in riverine ponds. *Canadian Journal of Fisheries and Aquatic Sciences*. 39: 1303–1307.
- Piccolo, J.J.; Wipfli, M.S. 2002.** Does red alder (*Alnus rubra*) in upland riparian forests elevate macroinvertebrate and detritus export from headwater streams to downstream habitats in southeastern Alaska? *Canadian Journal of Fisheries and Aquatic Sciences*. 59: 503–513.
- Pielke, R.A., Jr. 2007.** The honest broker: making sense of science in policy and politics. Cambridge, United Kingdom: Cambridge University Press. 188 p.
- Pimentel, D.; Lach, L.; Zuniga, R.; Morrison, D. 2000.** Environmental and economic costs of nonindigenous species in the United States. *BioScience*. 50: 53–65.
- Pinnegar, J.K.; Engelhard, G.H. 2008.** The ‘shifting baseline’ phenomenon: a global perspective. *Reviews in Fish Biology and Fisheries*. 18: 1–16.
- Poage, N.J.; Tappeiner, J.C., II. 2002.** Long-term patterns of diameter and basal area growth of old-growth Douglas-fir trees in western Oregon. *Canadian Journal of Forest Research*. 32: 1232–1243.
- Poff, N.L.; Allan, J.D.; Bain, M.B.; Karr, J.R.; Prestegard, K.L.; Richter, B.D.; Sparks, R.E.; Stromberg, J.C. 1997.** The natural flow regime. *BioScience*. 47: 769–784.

- Pollock, M.M.; Beechie, T.J. 2014.** Does riparian forest restoration thinning enhance biodiversity? The ecological importance of large wood. *Journal of the American Water Resources Association*. 50: 543–559.
- Pollock, M.M.; Beechie, T.J.; Imaki, H. 2012.** Using reference conditions in ecosystem restoration: an example for riparian conifer forests in the Pacific Northwest. *Ecosphere*. 3(11): art98.
- Pont, D.; Hugueny, B.; Beier, U.; Goffaux, D.; Melcher, A.; Noble, R.; Rogers, C.; Roset, N.; Schmutz, S. 2006.** Assessing river biotic condition at a continental scale: a European approach using functional metrics and fish assemblages. *Journal of Applied Ecology*. 43: 70–80.
- Poole, G.C.; Berman, C.H. 2001.** An ecological perspective on in-stream temperature: natural heat dynamics and mechanisms of human-caused thermal degradation. *Environmental Management*. 27: 787–802.
- Poole, G.C.; Dunham, J.B.; Keenan, D.M.; Sauter, S.T.; McCullough, D.A.; Mebane, C.; Lockwood, J.C.; Essig, D.A.; Hicks, M.P.; Sturdevant, D.J.; Materna, E.J. 2004.** The case for regime-based water quality standards. *BioScience*. 54: 155–161.
- Pope, K.; Brown, C.; Hayes, M.; Green, G.; Macfarlane, D., tech. cords. 2014.** Cascades frog conservation assessment. Gen. Tech. Rep. PSW-GTR-244. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 116 p.
- Potyondy, J.P.; Geier, T.W. 2011.** Watershed condition classification technical guide. FS-978. Washington, DC: U.S. Department of Agriculture, Forest Service. 48 p.
- Power, M.E.; Parker, M.S.; Dietrich, W.E. 2008.** Seasonal reassembly of a river food web: floods, droughts, and impacts of fish. *Ecological Monographs*. 78: 263–282.
- Quinn, T.P.; Myers, K.W. 2004.** Anadromy and the marine migrations of Pacific salmon and trout: Rounsefell revisited. *Reviews in Fish Biology and Fisheries*. 14: 421–442.
- Quinn, T.P.; Kinnison, M.T.; Unwin, M.J. 2001.** Evolution of Chinook salmon (*Oncorhynchus tshawytscha*) populations in New Zealand: pattern, rate, and process. *Genetica*. 112: 493–513.
- Quinn, T.P.; Petersen, N.P. 1996.** The influence of habitat complexity and fish size on over-winter survival and growth of individually marked juvenile coho salmon (*Oncorhynchus kisutch*) in Big Beef Creek, Washington. *Canadian Journal of Fisheries and Aquatic Sciences*. 53: 1555–1564.
- Quinn, T.P.; Vøllestad, L.A. 2003.** Influences of freshwater and marine growth on the egg size-egg number trade-off in coho and Chinook salmon. *Transactions of the American Fisheries Society*. 131: 591–598.
- Quiñones, R.M.; Holyoak, M.; Johnson, M.L.; Moyle, P.B. 2014a.** Potential factors affecting survival differ by run-timing and location: linear mixed-effects models of Pacific salmonids (*Oncorhynchus* spp.) in the Klamath River, California. *PLoS One*. 9(5): e98392.
- Quiñones, R.M.; Johnson, M.L.; Moyle, P.B. 2014b.** Hatchery practices may result in replacement of wild salmonids: adult trends in the Klamath basin, California. *Environmental Biology of Fishes*. 97: 233–246.
- Raggon, M. 2014.** Aquatic and Riparian Effectiveness Monitoring Program: invasive species report, 2014 field season. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region; U.S. Department of the Interior, Bureau of Land Management, Oregon State office. <https://reo.gov/monitoring/reports/watershed/AREMP%20Aquatic%20Invasive%20Species%20Report%202014.pdf>. (31 May 2017).
- Raggon, M. 2017.** Aquatic and Riparian Effectiveness Monitoring Program: invasive species report, 2016 field season. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region; U.S. Department of the Interior, Bureau of Land Management, Oregon State office. <https://reo.gov/monitoring/reports/watershed/AREMP%20Aquatic%20Invasive%20Species%20Report%202016.pdf>. (22 August 2017).

- Raggon, M.; Lanigan, S. 2011.** Aquatic and Riparian Effectiveness Monitoring Program: invasive species report, 2010 field season. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region; U.S. Department of the Interior, Bureau of Land Management, Oregon State office. <https://reo.gov/monitoring/reports/watershed/AREMP%20Aquatic%20Invasive%20Species%20Report%202010.pdf>. (31 May 2017).
- Raggon, M.; Lanigan, S. 2012a.** Aquatic and Riparian Effectiveness Monitoring Program: invasive species report, 2011 field season. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region; U.S. Department of the Interior, Bureau of Land Management, Oregon State office. <https://reo.gov/monitoring/reports/watershed/AREMP%20Aquatic%20Invasive%20Species%20Report%202011.pdf>. (31 May 2017).
- Raggon, M.; Lanigan, S. 2012b.** Aquatic and Riparian Effectiveness Monitoring Program: invasive species report, 2012 field season. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region; U.S. Department of the Interior, Bureau of Land Management, Oregon State office. <https://reo.gov/monitoring/reports/watershed/AREMP%20Aquatic%20Invasive%20Species%20Report%202011.pdf>. (31 May 2017).
- Raggon, M.; Lanigan, S. 2013.** Aquatic and Riparian Effectiveness Monitoring Program: invasive species report, 2013 field season. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region; U.S. Department of the Interior, Bureau of Land Management, Oregon State office. <https://reo.gov/monitoring/reports/watershed/AREMP%20Aquatic%20Invasive%20Species%20Report%202012.pdf>. (31 May 2017).
- Rahel, F.J.; Bierwagen, B.; Taniguchi, Y. 2008.** Managing aquatic species of conservation concern in the face of climate change and invasive species. *Conservation Biology*. 22: 551–561.
- Rand, P.S.; Hinch, S.G.; Morrison, J.; Foreman, M.G.G.; MacNutt, M.J.; Macdonald, J.S.; Healey, M.C.; Farrell, A.P.; Higgs, D.A. 2006.** Effects of river discharge, temperature, and future climates on energetics and mortality of adult migrating Fraser River sockeye salmon. *Transactions of the American Fisheries Society*. 135: 655–667.
- Raphael, M.G.; Molina, R., eds. 2007.** Conservation of rare or little-known species: biological, social, and economic considerations. Washington, DC: Island Press. 375 p.
- Ray, R.A.; Holt, R.A.; Bartholomew, J.L. 2012.** Relationship between temperature and *Ceratomyxa shasta*-induced mortality in Klamath River salmonids. *Journal of Parasitology*. 98: 520–526.
- Reagan, R.E. 2015.** Where the stream meets the road—prioritizing culvert replacement for fish passage. Corvallis, OR: Oregon State University. 69 p. M.S. thesis.
- Rechisky, E.L.; Welch, D.W.; Porter, A.D.; Jacobs, M.C.; Ladouceur, A. 2009.** Experimental measurement of hydrosystem induced delayed mortality in juvenile Snake River spring Chinook salmon (*Oncorhynchus tshawytscha*) using a large-scale acoustic array. *Canadian Journal of Fisheries and Aquatic Sciences*. 66: 1019–1024.
- Reeves, G.; Benda, L.; Dalton, M.; Chisholm-Hatfield, S. 2016b.** Freshwater aquatic environment. In: Dalton, M.M.; Benda, L.; Case, M.; Chisholm-Hatfield, S.; Cohn, N.; Conlin, M.; Lawler, J.; Mote, P.; Reeves, G.; Ruggiero, P.; Serafin, K.; Sharp, D. Climate change vulnerability assessment for the Treaty of Olympia Tribes: a report to the Quinault Indian Nation, Hoh Tribe, and Quileute Tribe. Corvallis, OR: Oregon Climate Change Research Institute. 52 p. Chapter 4. [Pagination by chapter].

- Reeves, G.H. 2006.** The Aquatic Conservation Strategy of the Northwest Forest Plan: an assessment after ten years. In: Haynes, R.W.; Bormann, B.T.; Lee, D.C.; Martin, J.R., tech. eds. 2006. Northwest Forest Plan—the first 10 years (1994– 2003): synthesis of monitoring and research results. Gen. Tech. Rep. PNW-GTR-651. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 181–217. Chapter 9.
- Reeves, G.H.; Benda, L.E.; Burnett, K.M.; Bisson, P.A.; Sedell, J.R. 1995.** A disturbance-based ecosystem approach to maintaining and restoring freshwater habitats of evolutionarily significant units of anadromous salmonids in the Pacific Northwest. American Fisheries Society Symposium. 17: 334–349.
- Reeves, G.H.; Bisson, P.A. 2009.** Fish and old-growth forests. In: Spies, T.; Duncan, S., eds. Old growth in a new world: a Pacific Northwest icon re-examined. Washington, DC: Island Press: 70–82.
- Reeves, G.H.; Burnett, K.M.; McGarry, E.V. 2003.** Sources of large wood in the main stem of a fourth-order watershed in coastal Oregon stream. Canadian Journal of Forestry Research. 33: 1363–1370.
- Reeves, G.H.; Duncan, S.L. 2009.** Ecological history vs. social expectations: managing aquatic ecosystems. Ecology and Society. 14(2): art8.
- Reeves, G.H.; Everest, F.H.; Hall, J.D. 1987.** Interactions between the redbside shiner (*Richardsonius balteatus*) and the steelhead trout (*Oncorhynchus mykiss*) in western Oregon: the influence of water temperature. Canadian Journal of Fisheries and Aquatic Sciences. 43: 1521–1533.
- Reeves, G.H.; Hohler, D.B.; Larsen, D.P.; Busch, D.E.; Kratz, K.; Reynolds, K.; Stein, K.F.; Atzet, T.; Hays, P.; Tehan, M. 2004.** Aquatic and riparian effectiveness monitoring plan for the Northwest Forest Plan. Gen. Tech. Rep. PNW-GTR-577. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 70 p.
- Reeves, G.H.; Pickard, B.R.; Johnson, K.N. 2016a.** An initial evaluation of potential options for managing riparian reserves of the Aquatic Conservation Strategy of the Northwest Forest Plan. Gen. Tech. Rep. PNW-GTR-937. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 97 p.
- Reeves, G.H.; Spies, T.A. 2017.** Watersheds and landscapes. In: Olson, D.H.; Van Horne, B., eds. People, forests, and change: lessons from the Pacific Northwest. Washington, DC: Island Press: 207–224. Chapter 15.
- Reeves, G.H.; Williams, J.E.; Burnett, K.M.; Gallo, K. 2006.** The Aquatic Conservation Strategy of the Northwest Forest Plan. Conservation Biology. 20: 319–329.
- Reid, P.C.; Hari, R.E.; Beaugrand, G.; Livingstone, D.M.; Marty, C.; Straile, D.; Barichivich, J.; Goberville, E.; Adrian, R.; Aono, Y.; Brown, R.; Foster, J.; Groisman, P.; Hélaouët, P.; Hsu, H.-H.; Kirby, R.; Knight, J.; Kraberg, A.; Li, J.; Lo, T.-T.; Myneni, R.B.; North, R.P.; Pounds, A.J.; Sparks, T.; Stübi, R.; Tian, Y.; Wiltshire, K.H.; Xiao, D.; Zhu, Z. 2016.** Global impacts of the 1980s regime shift. Global Change Biology. 22: 682–703. doi:10.1111/gcb.13106.
- Resh, V.H.; Brown, A.V.; Covich, A.P.; Gurtz, M.E.; Li, H.W.; Minshall, G.W.; Reice, S.R.; Sheldon, A.L.; Wallace, J.B.; Wissmar, R.C. 1988.** The role of disturbance in stream ecology. Journal of the North American Benthological Society. 7: 433–455.
- Reynolds, K.W.; Hessburg, P.F.; Bourgeron, P.S., eds. 2014.** Making transparent environmental management decisions: applications of the ecosystem management decision support system. Heidelberg: Springer. 337 p.
- Richardson, J.S.; Béraud, S. 2014.** Effects of riparian forest harvest on streams: a meta-analysis. Journal of Applied Ecology. 51: 1712–1721. doi:10.1111/1365-2664.12332
- Richardson, J.S.; Danehy, R.J. 2007.** A synthesis of the ecology of headwater streams and their riparian zones in temperate forests. Forest Science. 53: 131–147.

- Richardson, J.S.; Naiman, R.J.; Bisson, P.A. 2012.** How did fixed-width buffers become the standard practice for protecting water from forestry? *Freshwater Science*. 31: 232–238.
- Richardson, J.S.; Naiman, R.J.; Swanson, F.J.; Hibbs, D.E. 2005.** Riparian communities associated with Pacific Northwest headwater streams: assemblages, processes, and uniqueness. *Journal of the American Water Resources Association*. 41: 935–947.
- Richgels, K.L.D.; Russell, R.E.; Adams, M.J.; White, C.L.; Grant, E.H.C. 2016.** Spatial variation in risk and consequence of *Batrachochytrium salamandrivorans* introduction in the United States. *Royal Society Open Science* 8: 150616. <http://dx.doi.org/10.1098/rsos.150616>.
- Rieman, B.; Dunham, J.; Clayton, J. 2006.** Emerging concepts for management of river ecosystems and challenges to applied integration of physical and biological sciences in the Pacific Northwest, USA. *International Journal of River Basin Management*. 4: 85–97.
- Rieman, B.E.; Isaak, D.J. 2010.** Climate change, aquatic ecosystems, and fishes in the Rocky Mountain West: implications and alternatives for management. Gen. Tech. Rep. RMRS-GTR-250. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 46 p.
- Rieman, B.E.; Smith, C.L.; Naiman, R.J.; Ruggerone, G.T.; Wood, C.C.; Huntly, N.; Merrill, E.N.; Alldredge, J.R.; Bisson, P.A.; Congleton, J.; Fausch, K.D. 2015.** A comprehensive approach for habitat restoration in the Columbia Basin. *Fisheries*. 40: 124–135.
- Robards, M.D.; Quinn, T.P. 2002.** The migratory timing of adult summer-run steelhead in the Columbia River over six decades of environmental change. *Transactions of the American Fisheries Society*. 131: 523–536.
- Robinson, C.T. 2012.** Long-term changes in community assembly, resistance, and resilience following experimental floods. *Ecological Applications*. 22: 949–1961.
- Robinson, C.; Duinker, P.N.; Beazley, K.F. 2010.** A conceptual framework for understanding, assessing, and mitigating ecological effects of forest roads. *Environmental Reviews*. 18: 61–86.
- Romero, N.; Gresswell, R.E.; Li, J.L. 2005.** Changing patterns in coastal cutthroat trout (*Oncorhynchus clarki clarki*) diet and prey in a gradient of deciduous canopies. *Canadian Journal of Fisheries and Aquatic Sciences*. 62: 1797–1807.
- Roni, P.; Beechie, T.J.; Bilby, R.E.; Leonetti, F.E.; Pollock, M.M.; Pess, G.R. 2002.** A review of stream restoration techniques and a hierarchical strategy for prioritizing restoration in Pacific Northwest watersheds. *North American Journal of Fisheries Management*. 22: 1–20.
- Roni, P.; Pess, G.; Beechie, T.; Morely, S. 2010.** Estimating changes in coho salmon and steelhead abundance from watershed restoration: How much restoration is needed to measurably increase smolt production? *North American Journal of Fisheries Management*. 30: 1469–1484. doi:10.1577/M09-162.1.
- Roper, B.B.; Scarnecchia, D.L. 1999.** Emigration of age-0 Chinook salmon (*Oncorhynchus tshawytscha*) smolts from the upper South Umpqua River basin, Oregon, USA. *Canadian Journal of Fisheries and Aquatic Sciences*. 56: 939–946.
- Rose, A.L.; Salmon, T.P.; Lokondeh, T.; Neilan, B.A.; Waite, T.D. 2005.** Use of superoxide as an electron shuttle for iron acquisition by the marine cyanobacterium *Lyngbya majuscula*. *Environmental Science and Technology*. 39: 3708–3715.

- Rosenberg, D.; Gervais, J.; Vesely, D.; Barnes, S.; Holts, L.; Horn, R.; Swift, R.; Todd, L.; Yee, C. 2009.** Conservation assessment of the western pond turtle in Oregon (*Actinemys marmorata*). Version 1.0. Portland, OR: U.S. Department of the Interior, Bureau of Land Management; U.S. Department of Agriculture, Forest Service, Pacific Northwest Region; Oregon Department of Fish and Wildlife; City of Portland, Metro Regional Government. <http://www.fs.fed.us/r6/sfpnw/issssp/documents/planning-docs/ca-hr-actinemys-marmorata-2009-11.pdf>. (29 August 2016).
- Rosenberger, A.E.; Dunham, J.B.; Neuswanger, J.R.; Railsback, S.F. 2015.** Legacy effects of wildfire on stream thermal regimes and rainbow trout ecology: an integrated analysis of observation and individual-based models. *Freshwater Science*. 34: 1571–1584.
- Rosgen, D.L. 1994.** A classification of natural rivers. *Catena*. 22: 169–199.
- Rot, B.W.; Naiman, R.J.; Bilby, R.E. 2000.** Stream channel configuration, landform, and riparian forest structure in the Cascade Mountains, Washington. *Canadian Journal of Fisheries and Aquatic Sciences*. 57: 699–707.
- Rowe, J.C.; Garcia, T.S. 2014.** Impacts of wetland restoration efforts on an amphibian assemblage in a multi-invader community. *Wetlands*. 34: 141–153.
- Royer, T.V.; Minshall, G.W. 1997.** Temperature patterns in small streams following wildfire. *Archiv für Hydrobiologie*. 140: 237–242.
- Ruesch, A.S.; Torgersen, C.E.; Lawler, J.J.; Olden, J.D.; Peterson, E.E.; Volk, C.J.; Lawrence, D.J. 2012.** Projected climate-induced habitat loss for salmonids in the John Day River network, Oregon, USA. *Conservation Biology*. 26: 873–882.
- Russell, W. 2009.** The influence of timber harvest on the structure and composition of riparian forests in the Coastal Redwood region. *Forest Ecology and Management*. 257: 1427–1433.
- Rykaczewski, R.R.; Dunne, J.P. 2010.** Enhanced nutrient supply to the California current ecosystem with global warming and increased stratification in an earth system model. *Geophysical Research Letters*. 37(21): L21814. doi:10.1029/2010GL045019.
- Rykken, J.J.; Chan, S.S.; Moldenke, A.R. 2007.** Headwater riparian microclimate patterns under alternative forest management treatments. *Forest Science*. 53: 270–280.
- Safeeq, M.; Grant, G.E.; Lewis, S.L.; Kramer, M.G.; Staab, B. 2014.** A hydrogeologic framework for characterizing summer streamflow sensitivity to climate warming in the Pacific Northwest, USA. *Hydrology and Earth Systems Science*. 18: 3693–3710. doi:10.5194/hess-18-3693-2014.
- Safeeq, M.; Grant, G.E.; Lewis, S.L.; Staab, B. 2015.** Predicting landscape sensitivity to present and future floods in the Pacific Northwest, USA. *Hydrological Processes*. 29: 5337–5353. doi:10.1002/hyp.10553.
- Safeeq, M.; Grant, G.E.; Lewis, S.L.; Tague, C. 2013.** Coupling snowpack and groundwater dynamics to interpret historical streamflow trends in the western United States. *Hydrological Processes*. 27: 655–668.
- Safford, H.D.; Wiens, J.A.; Hayward, G.D. 2012.** The growing importance of the past in managing ecosystems of the future. In: Wiens, J.A.; Hayward, G.D.; Hugh, D.; Giffen, C., eds. *Historical environmental variation in conservation and natural resource management*. New York: John Wiley & Sons: 319–327.
- Sagar, J.P. 2004.** Movement and demography of larval coastal giant salamanders (*Dicamptodon tenebrosus*) in streams with culverts in the Oregon Coast Range. Corvallis, OR: Oregon State University. 83 p. M.S. thesis.
- Salathé, E.P., Jr. 2006.** Influences of a shift in north Pacific storm tracks on western North American precipitation under global warming. *Geophysical Research Letters*. 33: L19820.

- Salathé, E.P., Jr.; Mote, P.W.; Wiley, M.W. 2007.** Review of scenario selection and downscaling methods for the assessment of climate change impacts on hydrology in the United States Pacific Northwest. *International Journal of Climatology*. 27: 1611–1621.
- Sanderson, B.L.; Barnas, K.A.; Rub, A.M.W. 2009.** Nonindigenous species of the Pacific Northwest: an overlooked risk to endangered salmon? *BioScience*. 59: 245–256.
- Sarr, D.A.; Hibbs, D.E. 2007.** Multiscale controls on woody plant diversity in western Oregon riparian forests. *Ecological Monographs*. 77: 179–201.
- Scarnecchia, D.L.; Bergersen, E.P. 1987.** Trout production and standing crop in Colorado's small streams as related to environmental factors. *North American Journal of Fisheries Management*. 7: 315–330.
- Schade, C.B.; Bonar, S.A. 2005.** Distribution and abundance of nonnative fishes in streams of the western United States. *North American Journal of Fisheries Management*. 25: 1386–1394.
- Scheuerell, M.D.; Williams, J.G. 2005.** Forecasting climate-induced changes in the survival of Snake River spring/summer Chinook salmon (*Oncorhynchus tshawytscha*). *Fisheries Oceanography*. 14: 448–457.
- Schindler, D.E.; Hilborn, R.; Chasco, B.; Boatright, C.P.; Quinn, T.P.; Rogers, L.A.; Webster, M.S. 2010.** Population diversity and the portfolio effect in an exploited species. *Nature*. 465: 609–612.
- Schindler, D.E.; Rogers, D.E.; Scheuerell, M.D.; Abrey, C.A. 2005.** Effects of changing climate on zooplankton and juvenile sockeye salmon growth in southwestern Alaska. *Ecology*. 86: 198–209.
- Schindler, D.E.; Scheuerell, M.D.; Moore, J.W.; Gende, M.; Francis, T.B.; Palen, W.S. 2003.** Salmon and the ecology of coastal ecosystems. *Frontiers in Ecology and the Environment*. 1: 31–37.
- Schindler, D.W.; Bayley, S.E.; Barker, B.R.; Beaty, K.G.; Cruikshank, D.R.; Fee, E.J.; Schindler, E.U.; Stainton, M.P. 1990.** Effects of climate warming on properties of boreal lakes and streams at the Experimental Lakes Area, northwestern Ontario. *Limnology and Oceanography*. 41: 1004–1017.
- Schindler, M.H.; Gessner, M.O. 2009.** Functional leaf traits and biodiversity effects on litter decomposition in a stream. *Ecology*. 90: 1641–1649.
- Schloegel, L.M.; Daszak, P.; Cunningham, A.A.; Speare, R.; Hill, B. 2010.** Two amphibian diseases, chytridiomycosis and ranaviral disease, are now globally notifiable to the World Organization for Animal Health (OIE): an assessment. *Diseases of Aquatic Organisms*. 92: 101–108.
- Schtickzelle, N.; Quinn, T.P. 2007.** A metapopulation perspective for salmon and other anadromous fish. *Fish and Fisheries*. 8: 297–314.
- Schultz, M.T. 2001.** A critique of EPA's index of watershed indicators. *Journal of Environmental Management*. 62: 429–442.
- Sedell, J.R.; Reeves, G.H.; Bisson, P.A. 1997.** Habitat policy for salmon in the Pacific Northwest. In: Stouder, D.J.; Bisson, P.A.; Naiman, R.J., eds. *Pacific salmon and their ecosystems: status and future options*. New York: Chapman and Hall: 375–387.
- Sedell, J.R.; Reeves, G.H.; Burnett, K.M. 1994.** Development and evaluation of aquatic conservation strategies. *Journal of Forestry*. 92(4): 28–31.
- Sedell, J.R.; Swanson, F.J. 1984.** Ecological characteristics of streams in old-growth forests of the Pacific Northwest. In: Meehan, W.R.; Merrell, T.R., Jr.; Hanley, T.A., eds. *Fish and wildlife relationships in old-growth forests*. Juneau, AK: American Institute of Fishery Research Biologists: 9–16.

- Selvakumar, A.; Connor, T.P.O.; Struck, S.D. 2010.** Role of stream restoration on improving benthic macroinvertebrates and in-stream water quality in an urban watershed: case study. *Journal of Environmental Engineering*. 136: 127–139.
- Sestrich, C.M.; McMahon, T.E.; Young, M.K. 2011.** Influence of fire on native and nonnative salmonid populations and habitat in a western Montana basin. *Transactions of the American Fisheries Society*. 140: 136–146.
- Shanley, C.S.; Albert, D.M. 2014.** Climate change sensitivity index for Pacific salmon habitat in southeast Alaska. *PLoS One*. 9(8): e104799.
- Sheridan, C.D.; Spies, T.A. 2005.** Vegetation environment relationships in zero-order basins in coastal Oregon. *Canadian Journal of Forest Research*. 35: 340–355.
- Shindler, B.A.; Brunson, M.; Stankey, G.H. 2002.** Social acceptability of forest conditions and management practices: a problem analysis. Gen. Tech. Rep. PNW-GTR-537. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 68 p.
- Shoo, L.P.; Olson, D.H.; McMenamin, S.K.; Murray, K.A.; Van Sluys, M.; Donnelly, M.A.; Stratford, D.; Terhivuo, J.; Merino-Viteri, A.; Herbert, S.M.; Bishop, P.J.; Corn, P.S.; Dovey, L.; Griffiths, R.A.; Low, K.; Mahony, M.; McCallum, H.; Shuker, J.D.; Simpkins, C.; Skerratt, L.F.; Williams, S.E.; Hero, J-M. 2011.** Engineering a future for amphibians under climate change. *Journal of Applied Ecology*. 48: 487–492.
- Simberloff, D. 1998.** Flagships, umbrellas, and keystones: Is single-species management passé in the landscape era? *Biological Conservation*. 83: 247–257.
- Slaney, P.A. 1988.** Life history and smolt-to-adult survival of Keogh River steelhead trout (*Salmo gairdneri*) and the relationship to smolt size. *Canadian Journal of Fisheries and Aquatic Sciences*. 45: 1110–1122.
- Sloat, M.R.; Reeves, G.H. 2014.** Individual condition, standard metabolic rate, and rearing temperature influence steelhead and rainbow trout (*Oncorhynchus mykiss*) life histories. *Canadian Journal of Fisheries and Aquatic Sciences*. 71: 491–501.
- Smith, S.G.; Muir, W.D.; Williams, J.G.; Skalski, J.R. 2002.** Factors associated with travel time and survival of migrant yearling Chinook salmon and steelhead trout in the lower Snake River. *North American Journal of Fisheries Management*. 22: 385–405.
- Snover, A.K.; Hamlet, A.F.; Lettermeier, D.P. 2003.** Climate-change scenarios for water planning studies: pilot applications in the Pacific Northwest. *Bulletin of the American Meteorological Society*. 84: 1513–1518.
- Snyder, F.L.; Hilgendorf, M.B.; Garton, D.W. 1997.** Zebra mussels in North America: the invasion and its implications. Columbus, OH: Ohio Sea Grant. http://ohioseagrant.osu.edu/_documents/publications/FS/FS-045%20Zebra%20mussels%20in%20North%20America.pdf. (31 March 2017).
- Snyder, M.A.; Sloan, L.C.; Diffenbaugh, N.S.; Bell, J.L. 2003.** Future climate change and upwelling in the California Current. *Geophysical Research Letters*. 30(15): 1823. doi:10.1029/2003GL017647.
- Sogard, S.M.; Williams, T.H.; Fish, H. 2010.** Seasonal patterns of abundance, growth, and site fidelity of juvenile steelhead in a small coastal California stream. *Transactions of the American Fisheries Society*. 138: 549–563.
- Solazzi, M.F.; Nickelson, T.E.; Johnson, S.L.; Rodgers, J.D. 2000.** Effects of increasing winter rearing habitat on abundance of salmonids in two coastal Oregon streams. *Canadian Journal of Fisheries and Aquatic Sciences*. 57: 906–914.

- Soto, D.; Arismendi, I.; González, J.; Sanzana, J.; Jara, F.; Jara, C.; Guzman, E.; Lara, A. 2006.** Southern Chile, trout and salmon country: invasion patterns and threats for native species. *Revista Chilena de Historia Natural*. 79: 97–117.
- Spies, T.; Pollock, M.; Reeves, G.; Beechie, T. 2013.** Effects of riparian thinning on wood recruitment: a scientific synthesis. Science Review Team Wood Recruitment Subgroup. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region. 46 p. On file with: Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis, OR 97331.
- Spies, T.A.; Giesen, T.W.; Swanson, F.J.; Franklin, J.F.; Lach, D.; Johnson, K.N. 2010.** Climate change adaptation strategies for federal forests of the Pacific Northwest, USA: ecological, policy, and socio-economic perspectives. *Landscape Ecology*. 25: 1185–1199.
- Spies, T.A.; Hibbs, D.E.; Ohmann, J.L.; Reeves, G.H.; Pabst, R.J.; Swanson, F.J.; Whitlock, C.; Jones, J.A.; Wemple, B.C.; Parendes, L.A.; Schrader, B.A. 2002.** The ecological basis of forest ecosystem management in the Oregon Coast Range. In: Hobbs, S.D.; Hayes, J.P.; Johnson, R.L.; Reeves, G.H.; Spies, T.A.; Tappeiner, J.C., II; Wells, G.E., eds. *Forest and stream management in the Oregon Coast Range*. Corvallis, OR: Oregon State University Press: 31–67.
- Sponseller, R.A.; Benfield, E.F. 2001.** Influences of land use on leaf breakdown in southern Appalachian headwater streams: a multiple-scale analysis. *Journal of the North American Benthological Society*. 20: 44–59.
- Sponseller, R.A.; Benfield, E.F.; Valett, H.M. 2001.** Relationships between land use, spatial scale and stream macroinvertebrate communities. *Freshwater Biology*. 46: 1409–1424.
- Srivastava, D.S.; Cardinale, B.J.; Downing, A.L.; Duffy, J.E.; Jouseau, C.; Sankaran, M.; Wright, J.P. 2009.** Diversity has stronger top-down than bottom-up effects on decomposition. *Ecology*. 90: 1073–1083.
- Stanford, J.A.; Lorang, M.S.; Hauer, F.R. 2005.** The shifting habitat mosaic of river ecosystems. *Verhandlungen der Internationalen Vereinigung für Theoretische und Angewandte Limnologie*. 29: 123–136.
- Stankey, G.H.; Clark, R.N.; Bormann, B.T. 2005.** Adaptive management of natural resources: theory, concepts, and management institutions. Gen. Tech. Rep. PNW-GTR-654. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 73 p.
- Steel, E.A.; Muldoon, A.; Flitcroft, R.L.; Firman, J.C.; Anlauf-Dunn, K.J.; Burnett, K.M.; Danehy, R.J. 2016.** Current landscapes and legacies of land-use past: understanding the distribution of juvenile coho salmon (*Oncorhynchus kisutch*) and their habitats along the Oregon Coast, USA. *Canadian Journal of Fisheries and Aquatic Sciences*. 74: 546–561.
- Stefan, H.G.; Fang, X.; Eaton, J.G. 2001.** Simulated fish habitat changes in North American lakes in response to projected climate warming. *Transactions of the American Fisheries Society*. 130: 459–477.
- Steinblums, I.J. 1977.** Streamside buffer strips: survival effectiveness and design. Corvallis, OR: Oregon State University. 181 p. M.S. thesis.
- Stewart, I.T.; Cayan, D.R.; Dettinger, M.D. 2005.** Changes toward earlier streamflow timing across western North America. *Journal of Climate*. 18: 1136–1155.
- Stocking, R.W.; Holt, R.A.; Foott, J.S.; Bartholomew, J.L. 2006.** Spatial and temporal occurrence of the salmonid parasite *Ceratomyxa shasta* in the Oregon-California Klamath River basin. *Journal of Aquatic Animal Health*. 18: 194–202.
- Stoddard, J.L.; Larsen, D.P.; Hawkins, C.P.; Johnson, R.K.; Norris, R.H. 2006.** Setting expectations for the ecological condition of streams: the concept of reference condition. *Ecological Applications*. 16: 1267–1276.

- Sudduth, E.B.; Hassett, B.A.; Cada, P.; Bernhardt, E.S. 2011.** Testing the field of dreams hypothesis: functional responses to urbanization and restoration in stream ecosystems. *Ecological Applications*. 21: 1972–88
- Sundermann, A.; Stoll, S.; Haase, P. 2011.** River restoration success depends on the species pool of the immediate surroundings. *Ecological Applications*. 21: 1962–1971.
- Suter, G.W. 2001.** Applicability of indicator monitoring to ecological risk assessment. *Ecological Indicators*. 1:101–112.
- Sutfin, N.A.; Wohl, E.E.; Dwire, K.A. 2016.** Banking carbon: a review of organic carbon storage and physical factors influencing retention in floodplains and riparian ecosystems. *Earth Surface Processes and Landforms*. 41: 38–60.
- Suttle, K.B.; Power, M.E.; Levine, J.M.; McNeely, C. 2004.** How fine sediment in riverbeds impairs growth and survival of juvenile salmonids. *Ecological Applications*. 14: 969–974.
- Suzuki, N.; Olson, D.H.; Reilly, E.C. 2008.** Developing landscape habitat models for rare amphibians with small geographic ranges: a case study of Siskiyou Mountains salamanders in the western USA. *Biodiversity and Conservation*. 17: 2197–2218.
- Swan, C.M.; Gluth, M.A.; Horne, C.L. 2009.** Leaf litter species evenness influences nonadditive breakdown in a headwater stream. *Ecology*. 90: 1650–1658.
- Swanson, F.J.; Kratz, T.K.; Caine, N.; Woodmansee, R.G. 1988.** Landform effects on ecosystem patterns and processes. *BioScience*. 38: 92–98.
- Swanson, M.; Franklin, J.F.; Beschta, R.L.; Crisafulli, C.M.; DellaSala, D.A.; Hutto, R.L.; Lindenmayer, D.B.; Swanson, F.J. 2011.** The forgotten stage of forest succession: early-successional ecosystems on forest sites. *Frontiers in Ecology and the Environment*. 9: 117–125.
- Switalski, T.A.; Bissonette, J.A.; DeLuca, T.H.; Luce, C.H.; Madej, M.A. 2004.** Benefits and impacts of road removal. *Frontiers in Ecology and the Environment*. 2(1): 21–28.
- Tague, C.; Grant, G.E. 2009.** Ground water dynamics mediate low-flow response to global warming in snow-dominated alpine regions. *Water Resources Research*. 45: W07421. doi:10.1029/2008WR007179.
- Tappeiner, J.C., II; Huffman, D.W.; Marshall, D.; Spies, T.A.; Bailey, J.D. 1997.** Density, ages, and growth rates in old-growth and young-growth forests in coastal Oregon. *Canadian Journal of Forest Research*. 27: 638–648.
- Thomas, J.W.; Franklin, J.F.; Gordon, J.; Johnson, K.N. 2006.** The Northwest Forest Plan: origins, components, implementation experience, and suggestions for change. *Conservation Biology*. 20: 277–287.
- Thomas, J.W.; Raphael, M.G.; Anthony, R.G.; Forsman, E.D.; Gunderson, A.G.; Holthausen, R.S.; Marcot, B.G.; Reeves, G.H.; Sedell, J.R.; Solis, D.M. 1993.** Viability assessments and management considerations for species associated with late-successional and old-growth forests of the Pacific Northwest: the report of the Scientific Analysis Team. Portland, OR: U.S. Department of Agriculture, National Forest System, Forest Service Research. 530 p.
- Thompson, J.R.; Duncan, S.L.; Johnson, K.N. 2009.** Is there potential for the historical range of variability to guide conservation given the social range of variability? *Ecology and Society*. 14(1): art18.
- Thorp, J.H.; Thoms, M.C.; Delong, M.D. 2006.** The riverine ecosystem synthesis: biocomplexity in river networks across space and time. *River Research and Applications*. 22: 123–147.

- Tiwari, T.; Lundström, J.; Kuglerová, L.; Laudon, H.; Öhman, K.; Ågren, A.M. 2016.** Cost of riparian buffer zones: a comparison of hydrologically adapted site-specific riparian buffers with traditional fixed widths. *Water Resources Research*. 52: 1056–1069. doi: 10.1002/2015/WR018014.
- Tockner, K.; Stanford, J.A. 2002.** Riverine flood plains: present state and future trends. *Environmental Conservation*. 29: 308–330.
- Tohver, I.M.; Hamlet, A.F.; Lee, S.-Y. 2014.** Impacts of 21st-century climate change on hydrologic extremes in the Pacific Northwest region of North America. *Journal of the American Water Resources Association*. 50: 1461–1476.
- Torgersen, C.E.; Price, D.M.; Li, H.W.; McIntosh, B.A. 1999.** Multiscale thermal refugia and stream habitat associations of Chinook salmon in northeastern Oregon. *Ecological Applications*. 9: 301–319.
- Townsend, C.R. 1996.** Invasion biology and ecological impacts of brown trout *Salmo trutta* in New Zealand. *Biological Conservation*. 78: 13–22.
- Trombulak, S.C.; Frissell, C.A. 2000.** Review of ecological effects of roads on terrestrial and aquatic communities. *Conservation Biology*. 14: 18–30.
- Trotter, P.C. 1989.** Coastal cutthroat trout: a life history compendium. *Transactions of the American Fisheries Society*. 118: 463–473.
- Tullos, D.D.; Penrose, D.L.; Jennings, G.D.; Cope, W.G. 2009.** Analysis of functional traits in reconfigured channels: implications for the bioassessment and disturbance of river restoration. *Journal of the North American Benthological Society*. 28: 80–92.
- Turschwell, M.P.; Peterson, E.E.; Balcombe, S.R.; Sheldon, F. 2016.** To aggregate or not? Capturing the spatio-temporal complexity of the thermal regime. *Ecological Indicators*. 67: 39–48.
- Urban, M.C. 2015.** Accelerating extinction risk from climate change. *Science*. 348: 571–573.
- Urgenson, L.S.; Reichard, S.H.; Halpern, C.B. 2009.** Community and ecosystem consequences of giant knotweed (*Polygonum sachalinense*) invasion into riparian forests of western Washington, USA. *Biological Conservation*. 142: 1536–1541.
- U.S. Department of Agriculture [USDA]. 2017.** National Invasive Species Information Center: manager's tool kit. Washington, DC: U.S. Department of Agriculture, National Agricultural Library. <https://www.invasivespeciesinfo.gov/toolkit/detection.shtml>. (18 April 2017).
- U.S. Department of Agriculture, Forest Service [USDA FS]. 1997.** Tongass land management plan revision. Final environmental impact statement Part 1: summary, chapters 1 and 2, and chapter 3 (physical and biological environment). R10-MB-338b. Washington, DC: U.S. Department of Agriculture, Forest Service. [Irregular pagination].
- U.S. Department of Agriculture, Forest Service [USDA FS]. 2008.** Stream simulation: an ecological approach to providing passage for aquatic organisms at road-stream crossings. 7700—Transportation Management, 0877 1801—SDTDC. San Dimas, CA: U.S. Department of Agriculture, Forest Service, National Technology and Development Program, Stream Simulation Working Group. 659 p. <http://www.fs.fed.us/eng/pubs/pdf/StreamSimulation/index.shtml>. (7 September 2017).
- U.S. Department of Agriculture, Forest Service [USDA FS]. 2012a.** National Forest System Land Management Planning, Final Rule. *Federal Register* 77(68): 21162–21276 (April 9, 2012) (codified at 36 C.F.R. Part 219).
- U.S. Department of Agriculture, Forest Service [USDA FS]. 2012b.** Regional aquatic invasive species strategy and management plan. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region. 39 p. https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb5313602.pdf. (24 August 2017).

- U.S. Department of Agriculture, Forest Service [USDA FS]; U.S. Department of Commerce, National Oceanic and Atmospheric Administration; U.S. Department of Interior, Bureau of Land Management; U.S. Department of Interior, Fish and Wildlife Service. 2004.** Analytical process for developing biological assessments for federal actions affecting fish within the Northwest Forest Plan area. https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb5288191.pdf. (11 September 2017).
- U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management [USDA and USDI]. 1994a.** Record of decision for amendments to Forest Service and Bureau of Land Management planning documents within the range of the northern spotted owl. [Place of publication unknown]. 74 p. [plus attachment A: standards and guidelines].
- U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management [USDA and USDI]. 1994b.** Environmental assessment for the implementation of interim strategies (PACFISH) for managing anadromous fish-producing watersheds in eastern Oregon and Washington, Idaho, and portions of California. Washington, DC. [Pages unknown].
- U.S. Department of the Interior, Fish and Wildlife Service [USFWS]. 2015.** Endangered and threatened wildlife and plants; removing the Oregon chub from the federal list of endangered and threatened wildlife. Federal Register. 80(33): 9125–9150. (50 C.F.R. 17)
- U.S. Department of the Interior, Fish and Wildlife Service [USFWS]. 2016.** Fish and aquatic conservation: listing salamanders as injurious due to risk of salamander chytrid fungus. <https://www.fws.gov/injuriouswildlife/salamanders.html>. (28 March 2017).
- U.S. Government Accountability Office [USGAO]. 2001.** Restoring fish passage through culverts on Forest Service and BLM lands in Oregon and Washington could take decades. GAO-02-136. Washington, DC. 34 p. <http://www.gao.gov/assets/240/233076.pdf>. (5 September 2017).
- U.S. Environmental Protection Agency [USEPA]. 2016.** National rivers and streams assessment 2008–2009: a collaborative survey. EPA/841/R-16/007. Washington, DC: Office of Water and Office of Research and Development. https://www.epa.gov/sites/production/files/2016-03/documents/nrsa_0809_march_2_final.pdf. (7 September 2017).
- van den Berghe, P.E.; Gross, M.R. 1989.** Female size and nest depth in coho salmon (*Oncorhynchus kisutch*). Canadian Journal of Fisheries and Aquatic Sciences. 41: 204–206.
- van Mantgem, P.J.; Madej, M.A.; Seney, J.; Deshais, J. 2013.** Estimating ecosystem carbon stocks at Redwood National and State Parks. Park Science. 30: 20–26.
- Van Sickle, J.V.; Gregory, S.V. 1990.** Modeling inputs of large woody debris to streams from falling trees. Canadian Journal of Forest Research. 20: 1593–1601.
- van Vliet, M.T.; Franssen, W.H.; Yearsley, J.R.; Ludwig, F.; Haddeland, I.; Lettenmaier, D.P.; Kabat, P. 2013.** Global river discharge and water temperature under climate change. Global Environmental Change. 23: 450–464.
- van Vliet, M.T.H.; Ludwig, F.; Zwolsman, J.J.G.; Weedon, G.P.; Kabat, P. 2011.** Global river temperatures and sensitivity to atmospheric warming and changes in river flow. Water Resources Research. 47(2): W02544. doi:10.1029/2010WR009198.
- Vannote, R.L.; Minshall, G.W.; Cummins, K.W.; Sedell, J.R.; Cushing, C.E. 1980.** The river continuum concept. Canadian Journal of Fisheries and Aquatic Sciences. 37: 130–137.

- Ver Hoef, J.M.; Peterson, E.E.; Clifford, D.; Shah, R. 2014.** SSN: An R package for spatial statistical modeling on stream networks. *Journal of Statistical Software*. 56(3): 1–45. <https://eprints.qut.edu.au/103733/>. (8 September 2017).
- Villarin, L.A.; Chapin, D.M.; Jones, J.E. 2009.** Riparian forest structure and succession in second-growth stands of the central Cascade Mountains, Washington, USA. *Forest Ecology and Management*. 257: 1375–1385.
- Vitousek, P.M.; D’Antonio, C.M.; Loope, L.L.; Rejmanek, M.; Westbrooks, R. 1997.** Introduced species: a significant component of human caused global change. *New Zealand Journal of Ecology*. 21: 1–16.
- Wade, A.A.; Beechie, T.J.; Fleishman, E.; Mantua, N.J.; Wu, H.; Kimball, J.S.; Stoms, D.M.; Stanford, J.A. 2013.** Steelhead vulnerability to climate change in the Pacific Northwest. *Journal of Applied Ecology*. 50: 1093–1104.
- Walker, B.H. 1992.** Biodiversity and ecological redundancy. *Conservation Biology*. 6: 18–23.
- Wallace, J.B.; Eggert, S.L.; Meyer, J.L.; Webster, J.R. 1997.** Multiple trophic levels of a forest stream linked to terrestrial litter inputs. *Science*. 277: 102–104.
- Wallace, J.B.; Eggert, S.L.; Meyer, J.L.; Webster, J.R. 1999.** Effects of resource limitation on a detrital-based ecosystem. *Ecological Monographs*. 69: 409–442.
- Walters, C.J.; Hilborn, R.; Peterman, R.M.; Staley, M.J. 1978.** Model for examining early ocean limitation of Pacific salmon production. *Journal of the Fisheries Research Board of Canada*. 35: 1303–1315.
- Wang, D.; Gouhier, T.C.; Menge, B.A.; Ganguly, A.R. 2015.** Intensification and spatial homogenization of coastal upwelling under climate change. *Nature*. 518: 390–394. doi:10.1038/nature14235.
- Waples, R.S. 1991.** Pacific salmon, *Oncorhynchus* spp., and the definition of “species” under the Endangered Species Act. *Marine Fisheries Review*. 53: 11–22.
- Waples, R.S.; Beechie, T.; Pess, G.R. 2009.** Evolutionary history, habitat disturbance regimes, and anthropogenic changes: what do these mean for resilience of Pacific salmon populations? *Ecology and Society*. 14(1): 3. <http://www.ecologyandsociety.org/vol14/iss1/art3/>. (7 September 2017).
- Ward, E.J.; Anderson, J.H.; Beechie, T.J.; Pess, G.R.; Ford, M.J. 2015.** Increasing hydrologic variability threatens depleted anadromous fish populations. *Global Change Biology*. 21: 2500–2509. doi:10.1111/gcb.12847.
- Warren, C.E. 1971.** *Biology and water pollution control*. Philadelphia, PA: W.B. Saunders. 434 p.
- Warren, C.E.; Davis, G.E. 1967.** Laboratory studies on the feeding, bioenergetics, and growth of fish. In: Gerking, S.D., ed. *The biological basis of freshwater fish production*. Oxford, United Kingdom: Blackwell Science Publications: 175–214.
- Warren, D.R.; Collins, S.M.; Purvis, E.M.; Kaylor, M.J.; Bechtold, H.A. 2016.** Spatial variability in light yields colimitation of primary production by both light and nutrients in a forested stream ecosystem. *Ecosystems*. 1: 198–210.
- Washington Department of Fish and Wildlife [WDFW]. 2016.** Business and residential notice: chemical treatment for African clawed frogs (*Xenopus laevis*). Olympia, WA. <http://wdfw.wa.gov/ais/ACFCityLaceyChemicalTreatmentNotice.pdf>. (30 March 2017).
- Washington Forest Practice Board [WFPB]. 1993.** Standard methodology for conducting watershed analysis. Washington Forest Practice Act Board manual, Version 2.0. 85 p. [Note: this historical document may no longer be available.]
- Washington Invasive Species Council [WISP]. 2009.** Top priorities. Washington Invasive Species Council. <http://www.invasivespecies.wa.gov/documents/priorities/top50invasivespecies.pdf>. (28 March 2017).

- Welch, D.W.; Chigirinsky, A.I.; Ishida, Y. 1995.** Upper thermal limits on the oceanic distribution of Pacific salmon (*Oncorhynchus* spp.) in the spring. *Canadian Journal of Fisheries and Aquatic Sciences*. 52: 489–503.
- Welty, J.J.; Beechie, T.; Sullivan, K.; Hyink, D.M.; Bilby, R.E.; Andrus, C.; Pess, G. 2002.** Riparian aquatic interaction simulator (RAIS): a model of riparian forest dynamics for the generation of large woody debris and shade. *Forest Ecology and Management*. 162: 299–318.
- Wemple, B.C.; Jones, J.A. 2003.** Runoff production on forest roads in a steep, mountain catchment. *Water Resources Research*. 39(8): 1220. doi:10.1029/2002WR001744.
- Wenger, S.J.; Isaak, D.J.; Luce, C.H.; Neville, H.M.; Fausch, K.D.; Dunham, J.B.; Dauwalter, D.C.; Young, M.K.; Elsner, M.M.; Rieman, B.E.; Hamlet, A.F. 2011.** Flow regime, temperature, and biotic interactions drive differential declines of trout species under climate change. *Proceedings of the National Academy of Sciences of the United States of America*. 108: 14175–14180.
- Westerlund, S.; Ohman, P. 1991.** Iron in the water column of the Weddell Sea. *Marine Chemistry*. 35: 199–217.
- White, P.S.; Jentsch, A. 2001.** The search for generality in studies of disturbance and ecosystem dynamics. *Progress in Botany*. 62: 399–450.
- Wiens, J.A. 2002.** Riverine landscapes: taking landscape ecology into the water. *Freshwater Biology*. 47: 501–515.
- Wiens, J.D.; Anthony, R.G.; Forsman, E.D. 2014.** Competitive interactions and resource partitioning between northern spotted owls and barred owls in western Oregon. *Wildlife Monograph*. 185: 1–50.
- Wigington, P.J.; Ebersole, J.L.; Colvin, M.E.; Leibowitz, S.G.; Miller, B.; Hansen, B.; Lavigne, H.R.; White, D.; Baker, J.P.; Church, M.R.; Brooks, J.R. 2006.** Coho salmon dependence on intermittent streams. *Frontiers in Ecology and the Environment*. 4: 513–518.
- Wigington, P.J.; Leibowitz, S.G.; Comeleo, R.L.; Ebersole, J.L. 2013.** Oregon hydrologic landscapes: a classification framework. *Journal of the American Water Resources Association*. 49: 163–182. doi:10.1111/jawr.12009.
- Wilkerson, E.; Hagan, J.M.; Siegel, D.; Whitman, A.A. 2006.** The effectiveness of different buffer widths for protecting headwater stream temperature in Maine. *Forest Science*. 52: 221–231.
- Willson, M.F.; Gende, S.M.; Bisson, P.A. 2004.** Anadromous fishes as ecological links between ocean, fresh water, and land. In: Polis, G.A.; Power, M.E.; Huxel, G.R., eds. *Food webs at the landscape level*. Chicago, IL: University of Chicago Press: 284–300.
- Wilzbach, M.A.; Harvey, B.C.; White, J.L.; Nakamoto, R.J. 2005.** Effects of riparian canopy opening and salmon carcass addition on the abundance and growth of resident salmonids. *Canadian Journal of Fisheries and Aquatic Sciences*. 62: 58–67.
- Wimberly, M.C.; Spies, T.A. 2001.** Influences of environment and disturbance on forest patterns in coastal Oregon watersheds. *Ecology*. 82: 1443–1459.
- Wimberly, M.C.; Spies, T.A.; Long, C.J.; Whitlock C. 2000.** Simulating historical variability in the amount of old forests in the Oregon Coast Range. *Conservation Biology*. 14: 167–180.
- Winder, M.; Schindler, D.E. 2004.** Climate change uncouples trophic interactions in an aquatic ecosystem. *Ecology*. 85: 2100–2106.
- Winter, T.C. 2001.** The concept of hydrologic landscapes. *Journal of the American Water Resources Association*. 37: 335–349.
- Wipfli, M.S.; Baxter, C.V. 2010.** Linking ecosystems, food webs, and fish production: subsidies in salmonid watersheds. *Fisheries*. 35: 373–387.

- Wipfli, M.S.; Gregovich, D.P. 2002.** Export of invertebrates and detritus from fishless headwater streams in southeastern Alaska: implications for downstream salmonid production. *Freshwater Biology*. 47: 957–969.
- Wipfli, M.S.; Musslewhite, J. 2004.** Density of red alder (*Alnus rubra*) in headwaters influences invertebrate and detritus subsidies to downstream fish habitats in Alaska. *Hydrobiologia*. 520: 153–163.
- Wipfli, M.S.; Richardson, J.S.; Naiman, R.J. 2007.** Ecological linkages between headwaters and downstream ecosystems: transport of organic matter, invertebrates, and wood down headwater channels. *Journal of the American Water Resources Association*. 43: 72–85. doi:10.1111/j.1752-1688.2007.00007.x.
- Wohl, E. 2016.** Messy rivers are healthy rivers: the role of physical complexity. In: Constantinescu, G.; Garcia, M.; Hanes, D., eds. *River flow 2016*. New York: CRC Press: 24–29.
- Wohl, E.; Angermeier, P.L.; Bledsoe, B.; Kondolf, G.M.; MacDonnell, L.; Merritt, D.M.; Palmer, M.A.; Poff, N.L.; Tarboton, D. 2005.** River restoration. *Water Resources Research*. 41(10): W10301. doi:10.1029/2005WR003985.
- Wohl, E.; Gerlak, A.K.; Poff, N.L.; Chin, A. 2014.** Common core themes in geomorphic, ecological, and social systems. *Environmental Management*. 53: 14–27.
- Wondzell, S.M.; Hemstrom, M.A.; Bisson, P.A. 2007.** Simulating riparian vegetation and aquatic habitat dynamics in response to natural and anthropogenic disturbance regimes in the Upper Grande Ronde River, Oregon, USA. *Landscape and Urban Planning*. 80: 249–267.
- Wondzell, S.M.; Przeszlowska, A.; Pflugmacher, D.; Hemstrom, M.A.; Bisson, P.A. 2012.** Modeling the dynamic responses of riparian vegetation and salmon habitat in the Oregon Coast Range with state and transition models. In: Kerns, B.K.; Shlisky, A.J.; Daniel, C.J., tech. eds. 2012. *Proceedings of the first landscape state-and-transition simulation modeling conference*. Gen. Tech. Rep. PNW-GTR-869. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 173–196.
- Wootton, J.T.; Parker, M.S.; Power, M.E. 1996.** Effects of disturbance on river food webs. *Science*. 273: 1558–1561.
- World Organization for Animal Health [OIE]. 2017.** OIE-listed diseases, infections and infestations in force in 2017. <http://www.oie.int/animal-health-in-the-world/oie-listed-diseases-2017/>. (17 April 2017).
- Wu, T.H.; Bettadapura, D.P.; Beal, P.E.; Erb, R.T.; Hollister, D.L.; Lan, C.; Stadler, A.T. 1986.** Root geometry model and simulation. Unpublished final report. National Science Foundation Grant CEE-811253. USDA Forest Service Grant PNW-83-317. Columbus, OH: Ohio State University, Department of Civil Engineering. 62 p.
- Wurtsbaugh, W.A.; Davis, G.E. 1977.** Effects of temperature and ration level on the growth and food conversion efficiency of *Salmo gairdneri* Richardson. *Journal of Fish Biology*. 11: 87–98.
- Xie, G.Y.; Olson, D.H.; Blaustein, A.R. 2016.** Projecting the global distribution of the emerging amphibian fungal pathogen, *Batrachochytrium dendrobatidis*, based on IPCC climate futures. *PLoS One*. 11(8): e0160746.
- Yamada, B.S.; Gillespie, G.E. 2008.** Will the European green crab (*Carcinus maenas*) persist in the Pacific Northwest? *ICES Journal of Marine Science*. 65: 725–729.
- Yap, T.A.; Koo, M.S.; Ambrose, R.F.; Wake, D.B.; Vredenburg, V.T. 2015.** Averting a North American biodiversity crisis. *Science*. 349: 481–482.

Yeung, A.C.Y.; Lecerf, A.; Richardson, J.S. 2017.

Assessing the long-term ecological effects of riparian management practices on headwater streams in a coastal temperate rainforest. *Forest Ecology and Management*. 384: 100–109.

Yount, J.D.; Niemi, G.J. 1990. Recovery of lotic communities and ecosystems from disturbance—a narrative review of case studies. *Environmental Management*. 14: 547–569.

Zhang, C.; Li, S.; Qi, J.; Xing, Z.; Meng, F. 2017.

Assessing impacts of riparian buffer zones on sediment and nutrient loadings into streams at watershed scale using an integrated REMM-SWAT model. *Hydrological Processes*. 31: 916–924.

Zimmerman, A.; Church, M. 2001. Channel morphology, gradient profiles and bed stresses during a flood in a step-pool channel. *Geomorphology*. 40: 311–327.

Appendix 1: Aquatic-Riparian Invasive Species of the Northwest Forest Plan Area

*Deanna H. Olson and Rebecca L. Flitcroft*¹

Invasive species are generally considered novel species that are not native to established systems, and their introductions are harmful ecologically (Vitousek et al. 1997) or economically (Pimentel et al. 2000). Nuisance-species lists have been developed for various jurisdictions, including species that both have been or potentially could be introduced to an area with subsequent adverse effects. Priority aquatic-riparian invasive species (ARIS) include those that have the potential to greatly alter food webs or ecosystem structure, economic interests such as fisheries, and recreation opportunities or human safety—for example, by fouling waterways or affecting water transportation. Priority invasive species include pathogens that can trigger disease die-offs, predators that may restructure native communities via trophic cascades, ecosystem engineers that alter physical or biological habitat conditions, and macroinvertebrates and plants that may produce population booms in systems, altering their ecosystem structure or function.

ARIS were not raised as a priority concern during development of the Aquatic Conservation Strategy (ACS) for the Northwest Forest Plan (NWFP, or Plan) in 1993–1994. As described in the 10 ACS objectives (USDA and USDI 1994a), the focus at the time was to maintain and restore watershed, landscape, riparian, and aquatic habitat conditions to which species, populations, and communities are uniquely adapted—hence emphasis was placed on native species. More explicitly, ACS objective 10 refers to the maintenance and restoration of habitat to support well-distributed populations of **native** plant, invertebrate, and vertebrate riparian-dependent species. Since 1994, ARIS concerns have intensified, and several state and federal agency groups with species jurisdictions overlapping the range of the NWFP have been addressing ARIS. In particular, modifications to the Aquatic-Riparian Effectiveness Monitoring Program (AREMP) now address ARIS during

the program’s annual monitoring efforts. Herein, we provide an overview of ARIS that are priorities for natural resource managers in the NWFP area, highlight key science findings of recent research, and describe the development of invasive species monitoring programs.

Priority Aquatic Invasive Species

Overall, across the Plan area, we identified 63 species and species groups as top regional aquatic-riparian invasive or nuisance-species priorities (table 7-10). Of these, 31 (49 percent) species or species groups were designated as “high concern” and inventoried by AREMP in 2016. Our broader top-priority list of 63 taxa was derived from lists compiled by state government departments in the region, interagency collaborative groups such as state invasive species councils, regional U.S. Forest Service personnel, or other entities identifying nuisance species or emerging infectious diseases. Specifically, our 63 priority taxa include those aquatic-riparian species on Oregon’s “100 Worst List” (OISC 2015), Washington’s “50 Priority Species” list (WISP 2009), a focal species list for U.S. Forest Service Pacific Northwest Region (Region 6) lands (Flitcroft et al. 2016b; S. Bautista, pers. comm.²), and the AREMP list (Raggon 2017). We recognize that top priorities identified by California and the U.S. Forest Service Pacific Southwest Region include other aquatic-riparian taxa, but upon inspection, species identified only in California and not by these other sources appeared to be of lesser immediate concern in northwestern California forests in the Plan area. We acknowledge that some other important California invasive species may merit consideration if our list were to be refined further. Lastly, some pathogens were included here because of their national and international priority status from other entities (Auliya et al. 2016, Bern Convention 2015, Conservation Institute 2013, OIE 2017, Schloegel et al. 2010, USFWS 2016). Note that priority species differ between

¹ **Deanna H. Olson** is a research ecologist and **Rebecca L. Flitcroft** is a research fish biologist, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis, OR 97331.

² Bautista, S. 2017. Personal communication. Pesticide use & invasive plant coordinator, U.S. Forest Service, 1220 SW Third Ave., Portland, OR 97204, sbautista@fs.fed.us.

Table 7-10—Aquatic invasive species of concern in the Pacific Northwest states of Oregon and Washington and within the administrative boundaries of the U.S. Forest Service Pacific Northwest Region

Scientific name	Common name	Oregon	Washington	Pacific Northwest Region	AREMP
Pathogens and parasites:					
<i>Phytophthora alni</i> , <i>P. kernoviae</i> , <i>P. pluvialis</i> , <i>P. lateralis</i> ; <i>P.</i> <i>ramorum</i>	Alder root rot; <i>Phytophthora</i> taxon C; needle cast of Douglas-fir, Port Orford cedar root disease, sudden oak death	✓		✓	
<i>Batrachochytrium dendrobatidis</i> , <i>B. salamandrivorans</i>	Amphibian chytrid fungi (<i>Bd</i> , <i>Bsal</i>)				
<i>Bothriocephalus acheilognathi</i>	Asian tapeworm	✓			
<i>Orthomyxoviridae isavirus</i>	Infectious salmon anemia virus (ISAV)	✓			
<i>Ranavirus</i>	Ranavirus				
Rhabdovirus SVCV	Spring viremia of carp virus (SVCV)		✓		
<i>Novirhabdovirus</i> spp.	Viral hemorrhagic septicemia virus (VHSV)	✓	✓		
<i>Myxobolus cerebralis</i>	Whirling disease	✓			
Aquatic plants:					
<i>Lagarosiphon major</i>	African waterweed or African elodea	✓		✓	
<i>Phragmites australis</i>	Common reed	✓	✓		✓
<i>Potamogeton crispus</i>	Curly-leaf pondweed			✓	✓
<i>Butomus umbellatus</i>	Flowering rush	✓		✓	✓
<i>Salvinia molesta</i>	Giant salvinia	✓			✓
<i>Arundo donax</i>	Giant reed				✓
<i>Hydrilla verticillata</i>	Hydrilla, water thyme	✓	✓	✓	✓
<i>Myriophyllum</i> spp. including <i>M.</i> <i>spicatum</i> , <i>M. aquaticum</i>	Milfoils: Eurasian, parrotfeather		✓	✓	✓
<i>Lythrum salicaria</i> , <i>Lysimachia</i> <i>vulgaris</i>	Purple loosestrife, garden yellow loosestrife		✓	✓	✓
<i>Phalaris arundinacea</i> ; <i>P.</i> <i>arundinacea</i> var. <i>picta</i>	Reed canary grass; ribbongrass			✓	
<i>Didymosphenia geminata</i>	Rock snot (Didymo)	✓		✓	✓
<i>Chondrilla juncea</i>	Rush skeletonweed		✓	✓	
<i>Spartina</i> spp. including <i>S.</i> <i>alterniflora</i> , <i>S. densiflora</i>	Spartina (cordgrass)	✓	✓		
<i>Prymnesium parvum</i> , <i>Cylindrospermopsis raciborskii</i>	Toxic algae (golden, toxic cyanobacteria)	✓			
<i>Trapa natans</i>	Water chestnut (European)	✓	✓		

Table 7-10—Aquatic invasive species of concern in the Pacific Northwest states of Oregon and Washington and within the administrative boundaries of the U.S. Forest Service Pacific Northwest Region (continued)

Scientific name	Common name	Oregon	Washington	Pacific Northwest Region	AREMP
<i>Ludwigia</i> spp.	Water primrose			✓	✓
<i>Egeria densa</i> ; <i>Elodea nuttallii</i> , <i>E. canadensis</i> , <i>E. canadensis</i> × <i>E. nuttallii</i> hybrid	Brazilian elodea, western waterweed (<i>Elodea</i>)		✓		✓
<i>Iris pseudacorus</i>	Paleyellow iris			✓	✓
<i>Nymphoides peltata</i>	Yellow floating heart	✓		✓	✓
Riparian-terrestrial plants:					
<i>Hedera helix</i>	English ivy			✓	✓
<i>Alliaria petiolata</i>	Garlic mustard		✓		✓
<i>Geranium robertianum</i> , <i>G. lucidum</i>	geraniums (Herb-Robert, shining)			✓	✓
<i>Heracleum mantegazzianum</i>	Giant hogweed	✓	✓		✓
<i>Rubus ulmifolius</i>	Himalayan blackberry		✓	✓	✓
<i>Fallopia japonica</i> var. <i>japonica</i> ; <i>Polygonum bohemicum</i>	Knotweeds (Japanese, Bohemian)		✓	✓	✓
<i>Pueraria lobata</i>	Kudzu	✓	✓		✓
<i>Clematis vitalba</i>	Old man's beard			✓	✓
<i>Hieracium aurantiacum</i>	Orange hawkweed				✓
<i>Potentilla recta</i>	Sulphur cinquefoil			✓	
<i>Tamarix</i> spp.	Tamarix (salt cedar)		✓	✓	✓
<i>Lamium strumarium</i>	Yellow archangel				✓
Aquatic invertebrates:					
<i>Potamocorbula amurensis</i>	Asian clam	✓		✓	✓
<i>Radix auricularia</i>	Big-eared radix			✓	✓
<i>Eriocheir sinensis</i>	Chinese mitten crab	✓	✓		
<i>Cipangopaludina chinensis</i>	Chinese mystery snail			✓	✓
<i>Orconectes</i> spp., <i>Orconectes virilis</i> , <i>Procambarus</i> spp.	Crayfish (red swamp, rusty, ringed, virile, marbled, signal, northern)		✓	✓	✓
<i>Carcinus maenas</i>	European green crab		✓		
<i>Potamopyrgus antipodarum</i>	New Zealand mud snail		✓	✓	✓
<i>Philine auriformis</i>	New Zealand sea slug	✓			
<i>Bythotrephes longimanus</i> [cederstroemi], <i>Cercopagis pengoi</i>	Waterfleas	✓			
<i>Dreissena polymorpha</i> , <i>D. rostriformis bugensis</i>	Zebra and quagga mussels	✓	✓	✓	✓

Table 7-10—Aquatic invasive species of concern in the Pacific Northwest states of Oregon and Washington and within the administrative boundaries of the U.S. Forest Service Pacific Northwest Region (continued)

Scientific name	Common name	Oregon	Washington	Pacific Northwest Region	AREMP
Aquatic vertebrates:					
<i>Lithobates catesbeianus</i> (<i>Rana catesbeiana</i>)	American bullfrog		✓	✓	✓
<i>Hypophthalmichthys</i> spp., <i>Mylopharyngodon piceus</i>	Asian carp, black carp	✓	✓		
<i>Salmo salar</i>	Atlantic salmon	✓	✓		
<i>Didemnum vexillum</i>	<i>Didemnum</i> tunicate		✓		
<i>Chelydra serpentina serpentina</i>	Eastern snapping turtle	✓			
<i>Neogobius melanostomus</i> , <i>Rhinogobius brunneus</i> , <i>Tridentiger bifasciatus</i>	Goby	✓			
<i>Noteigonus crysoleucas</i>	Golden shiner	✓			
<i>Esox</i> spp.	Muskellunge/northern pike	✓			
<i>Gymnocephalus cernuus</i>	Ruffe	✓			
<i>Channa</i> spp.	Snakehead	✓	✓		
<i>Dorosoma petenense</i>	Threadfin shad (yellow tails)	✓			
Riparian-terrestrial vertebrates:					
<i>Sus scrofa</i>	Feral swine	✓	✓	✓	✓
<i>Cygnus olor</i>	Mute swan	✓			
<i>Myocaster coypus</i>	Nutria		✓	✓	✓

Note: Three aquatic pathogens (ranavirus, *Batrachochytrium* spp.) are included in this table owing to other national and international priority status (Auliya et al. 2016, Bern Convention 2015, Conservation Institute 2013, OIE 2016, Schloegel et al. 2010, USFWS 2016). Some species clustering within rows was conducted. “✓” denotes priority species from authority listed (above), not occurrence of species within jurisdiction.

Source: Flitcroft et al. 2016b; Bautista, S., personal communication (see footnote 2 on page 585); and Aquatic-Riparian Effectiveness Monitoring Program (AREMP) of the Northwest Forest Plan (Raggon 2017).

lists created by different jurisdictions or entities because of their variable selection criteria or jurisdiction-specific habitats and issues. A species listed as a priority for one jurisdiction and not another may have established populations and be of concern in both areas, yet because of different perspectives not be considered a top priority everywhere. Hence, the broader species list may be important to consider as regionally representative taxa of ecological or economic concern from an all-lands perspective across the Plan area. A few estuarine species are included and may be relevant to consider here, because tidally influenced areas are critical ecosystems interfacing with the

coastal forest land base. Other primarily marine-associated species are not included here.

Altogether, as potentially representative of the Plan area, Northwest taxa identified as regional ARIS priorities fall into six categories (table 7-10): 8 pathogens; 19 aquatic plants; 12 riparian-terrestrial plants; 10 aquatic invertebrates; 11 aquatic vertebrates; and 3 riparian-terrestrial vertebrates. Specifically for the Plan area, AREMP’s 31 invasive taxa fall into 5 categories: 13 aquatic plants; 11 riparian-terrestrial plants; 6 aquatic invertebrates; 1 aquatic vertebrate; and 2 riparian-terrestrial vertebrates.

Taxonomic Summaries

Pathogens and Parasites

Pathogens and parasites are considered invasive when their spread has been documented coincident with devastating disease effects on host species. There is heightened concern for disease-causing pathogens and parasites affecting sensitive host taxonomic groups that provide important ecosystem services (goods and services that people desire) (Blahna et al. 2017, Penaluna et al. 2016). These host taxa include culturally and economically important species such as salmonid fishes; species with broad distributions that may be central to ecosystem structural integrity and biodiversity such as Douglas-fir (*Pseudotsuga menziesii*) and alder (*Alnus* spp.); and those diseases with multiple host taxa that could affect several native species, with consequences for the organization of wild, native communities, such as the fungi that cause amphibian chytridiomycosis.

Northwest aquatic-riparian pathogens of key concern are viruses and fungi; parasites include cnidarian and cestode worms. Aquatic invasive pathogen species infect vertebrates, for example: (1) cnidarian myxosporean parasites (*Myxobolus cerebralis*) cause whirling disease in salmonid fishes (first described in Germany); (2) Asian tapeworms infect cyprinid fishes in their native range in Asia; (3) viral hemorrhagic septicemia virus (VHSV) infects salmonids, historically known from Europe; (4) ranavirus is considered an emerging infectious disease of fishes, turtles, and amphibians; and (5) chytrid fungi of the genus *Batrachochytrium* can cause the emerging infectious disease chytridiomycosis in amphibians. Riparian pathogens of key concern are fungi of the genus *Phytophthora* (table 7-10) that infect trees typical of riparian zones, such as alder, Douglas-fir, and Port Orford cedar (*Chaemaecyparis lawsoniana*).

The World Organization of Animal Health (OIE) lists species as notifiable because of the extent of their effects, the availability of diagnostic tests for detection, and the role of humans in disease spread. The United States is one of 180 member nations of OIE, hence OIE listing is relevant for consideration here. For pathogens identified here as top ARIS priorities in Oregon, Washington, and Alaska, three are OIE notifiable (OIE 2017): VHSV, ranavirus, and *B. dendrobatidis* (*Bd*). *Bd* is also listed by the Conservation

Institute (2013) on their world list of the Top 100 Invasive Species. A second amphibian chytrid fungus included on our Top 60 list, *B. salamandrivorans* (*Bsal*), was recently described in Europe (Martel et al. 2013), and early challenge experiments found numerous North American taxa to be vulnerable to disease effects, with rapid mortality after infection (Martel et al. 2014). The Bern Convention (2015) and others (Auliya et al. 2016) have endorsed legislation to forestall the spread of *Bsal*, and an interim rule to the U.S. Lacey Act (USFWS 2016) has listed host salamander species that may be susceptible to *Bsal* infection as injurious, hence restricting their transportation among jurisdictions. *Bsal* risk models (Richgels et al. 2016, Yap et al. 2015) show Oregon and Washington to be extremely vulnerable to *Bsal* introduction owing to the presence of susceptible host amphibian taxa such as the rough-skinned newt (*Taricha granulosa*), suitable *Bsal* habitat conditions, and proximity to U.S. ports of entry. The pet industry has placed a moratorium on some salamander trade imports to the United States, significantly forestalling the transmission of this pathogen.

Aquatic Plants

The 19 aquatic invasive plants of concern (table 7-10) include two algae, a diatom (rock snot or Didymo), multiple species of submerged aquatic plants (e.g., *Elodea*, *Hydrilla*, milfoils [*Myriophyllum* spp.]), emergent plants (e.g., reeds, cordgrass [*Spartina* spp.], loosestrife [*Lysimachia* spp.], rushes, *Salvinia*, reed canarygrass [*Phalaris arundinacea*], paleyellow iris [*Iris pseudacorus*], water primrose [*Ludwigia* spp.], and floating plants (e.g., curly-leaf pondweed [*Potamogeton crispus*], water chestnut [*Trapa natans*], yellow floating heart [*Nymphoides peltata*]). Once established, these taxa may affect ecosystems, water quality, human health, navigation, and recreation. Emergent plants can dominate wetland and floodplain areas, outcompeting or displacing native species, thereby reducing biodiversity and altering ecosystem functions. Toxic algae are a health concern for native vertebrates and humans because they create powerful toxins known to kill fish, ducks, geese, marine mammals, and other wildlife (Edwards 1999). The diatom commonly called rock snot (Didymo) is native to

the Pacific Northwest but is included on priority invasive species lists owing to a change in its growth habits in the mid-1980s (Bothwell et al. 2014), becoming more prolific in its distribution and affecting recreational experiences and activities. Some Northwest ARIS plants were initially brought to the region by the aquarium trade (the elodeas) or for ornamental use (reed canary grass), and then spread to other areas. Further, established populations may be spread by waterfowl as they move from one location to another, or by human vectors (e.g., boats and fishing gear/tackle).

Riparian-Terrestrial Plants

The 12 invasive riparian plants listed (table 7-10) are problematic in both upland and riparian environments. Species such as Himalayan blackberry (*Rubus armeniacus*), Japanese knotweed (*Fallopia japonica* var. *japonica*), and giant hogweed (*Heracleum mantegazzianum*) tend to shade out smaller native plants, reducing plant diversity and altering habitat and food resources for native wildlife. For example, Japanese knotweed (native to Europe and Asia) and giant hogweed (native to the Caucasus region of Eurasia) can grow as tall as 15 to 20 ft (4.5 to 6 m) and spread rapidly. Japanese knotweed is known globally as one of the world's most destructive invasive species because its large underground root system can damage structures, walls, and architectural sites, and reduce stream-channel capacity. Giant hogweed is considered a public-health hazard because it causes a phototoxic reaction when skin is exposed to sap and ultraviolet radiation. Species in the genus *Tamarix* are riparian shrubs or small trees that are aggressively invasive and well known in the Southwestern United States. These riparian trees are known to decrease streamflows, lower biodiversity, and create salinization issues, among other problems. Some of the listed invasive riparian plants were imported to the Northwest as ornamentals and have become invasive (e.g., English ivy [*Hedera helix*] and old man's beard [*Clematis vitalba*], native to the United Kingdom; garlic mustard [*Alliaria petiolate*], native to Europe and Asia). Garlic mustard was initially introduced to the east coast of North America as a medicinal herb, but it has spread through forest understories, where it competes with native species.

Aquatic Invertebrates

The 10 invertebrates on the northwest aquatic-riparian invasive species list (table 7-10) include several mollusks (Asian clam [*Potamocorbula amurensis*], big-eared radix [*Radix auricularia*], Chinese mystery snail [*Cipangopaludina chinensis*], New Zealand mud snail [*Potamopyrgus antipodarum*], New Zealand sea slug [*Philine auriformis*], zebra mussel [*Dreissena polymorpha*], quagga mussel [*D. rostriformis bugensis*]), and crustaceans (crayfish and crab species; waterfleas). Mollusks may spread rapidly and attain large population sizes that displace native species. These taxa can deplete prey resources rapidly, affecting foundation levels of food webs (algae, phytoplankton) in aquatic systems. Along with abundant populations come abundant waste products—in some systems, the tissues or waste products of zebra mussels may accumulate contaminants to 300,000 times the level available in the habitat they occupy, with subsequent effects on their environment, including their predators (Snyder et al. 1997). Another concern is that large numbers of some mollusks can foul human structures. Introductions of some species are likely tied to inadvertent human transmission, such as in ship ballast water or on boats or fishing gear (e.g., zebra/quagga mussels, waterfleas, green crabs). Deliberate introduction and consequent escape of some species is also associated with food and medical markets, biological supplies for education, and the aquarium and bait trade (Chinese mystery snails, crayfish, mitten crabs).

Aquatic Vertebrates

One frog (American bullfrog [*Lithobates catesbeianus*]), one turtle, eight fishes, and a tunicate (*Didemnum*) are included in the priority aquatic-riparian invasive species list (table 7-10). These taxa are strongly tied to human introductions. For example, American bullfrogs are native to the Eastern United States and were brought to the West to be farmed for food and out of nostalgia for their calls. Bullfrogs are carriers of the amphibian chytrid fungus *Bd* but do not always exhibit disease symptoms and hence may serve as a reservoir species of the pathogen, another invasive species of concern. Additional concerns surrounding bullfrog introductions include alterations of the native ecosystem via food-web changes, an issue associated with the snapping turtle (*Chelydra serpentina*) as well.

The invasive fishes include a mix of species introduced for human food, as bait for recreational fisheries, or from the aquarium or ornamental industry. Atlantic salmon (*Salmo salar*) are native to the North Atlantic Ocean, where they are anadromous, occurring in the ocean and returning to spawn in rivers. Farms in Washington and British Columbia are thought to be the origin of Atlantic salmon found elsewhere in the Northwest. Concerns arise in conjunction with their potential competition with native salmonids, pollution from the farms, and the potential for farm-raised animals to carry pathogens to native stocks. Gobies are of Asian origin, occurring in fresh and brackish water. They are thought to have been introduced in ballast water, and may compete with or prey upon native species. Golden shiners (*Noteigonus crysoleucas*) are from the Eastern United States and are pond-cultured fishes that are also used as bait. Where numerous, golden shiners may result in displacement of native species.

Didemnum vexillum is commonly called the carpet sea squirt, or ascidian. It is a colonial tunicate in the chordate phylum, hence is included here together with vertebrates—a sister chordate lineage. It seems to be native to Japan, and has been detected along the Washington coast since 2009, in two Oregon bays since 2010, and near Sitka, Alaska, in 2010. It is a fouling organism in marine and estuarine systems that grows rapidly to cover vast surfaces as mats, displacing native biota and encrusting dock pilings and aquatic equipment. It can be introduced in ballast water, or may hitchhike on the hulls of boats or on commercial shellfish stock or equipment.

Riparian-Terrestrial Vertebrates

The category of terrestrial vertebrates is the smallest, with only three species (table 7-10), but these can have extensive aquatic-riparian effects, ecologically and socioeconomically. Feral swine are escaped domestic pigs with rooting behavior that degrades waterway habitat, provides an invasion pathway for nonnative plants, and causes damage to agricultural crops and lands. The mute swan was introduced from New York for aesthetic enjoyment. These aggressive, large (2- to 30-lb [0.9- to 14-kg]) birds may consume significant quantities of aquatic plants, competing with

native birds for food and habitat. Nutria (*Myocastor coypus*) were initially brought to the Pacific Northwest for fur farming in the 1920s. After the collapse of this element of the fur industry, escaped and released animals subsequently spread throughout the region. Nutria burrow into the banks of streams and agricultural canals, destabilizing natural stream systems and human agricultural infrastructure, and they consume vegetable crops.

Research and Development, Monitoring, and Management

Research and Development

Invasive-species disturbance ecology has developed conceptually in the past few decades. Aquatic-riparian ecosystems, like their terrestrial counterparts, are heterogeneous in space and time, occurring in multiple states within an ecosystem domain (e.g., Penaluna et al. 2016). This domain is highly resilient to many natural disturbances, yet larger disturbances can push an ecosystem beyond its “tipping point” to a new domain, a novel ecosystem. Novel ecosystems (e.g., Hobbs et al. 2009) are developing on our planet from a variety of disturbances, including the effects of invasive species. As discussed above, invaders may engineer habitat structures and functions, or become key players in food webs and trophic cascades, altering the native community and ecosystem. Biotic homogenization may result when the variety of initial states of ecosystems becomes equalized as a result of domination of invasive species over natives (e.g., McKinney and Lockwood 1999, Olden et al. 2004). “The New Normal” is a pragmatic description of today’s ecological systems that seem to be undergoing irrevocable change, a newly developing status quo (Marris 2010). However, it may be premature to characterize such changes as irrevocable, because there are many examples of restoration successes (e.g., Murcia et al. 2014). Nevertheless, without preempting invasions, vigilance at the early stages of invasion, and concerted, often-continuous restoration efforts, transformation to a novel ecosystem can occur.

These concepts are playing out with aquatic invasive species globally. As cases of ARIS are analyzed, costs to native biodiversity are being claimed. For example, the introductions of the European brown trout (*Salmo trutta*)

into South America (Soto et al. 2006) and New Zealand (Townsend 1996), and the eastern mosquitofish (*Gambusia holbrooki*) into Australia (Hamer et al. 2002), have caused major reductions in native fishes. Similar adverse effects on native amphibians and other ecosystem components have been documented by fish-stocking practices (reviews: Dunham et al. 2004, Kats and Ferrer 2003). Despite regionwide efforts to control ARIS spread, some species are recognized as requiring continuous management, or the efficacy of control methods is low. As a result, some invasive species (e.g., bullfrogs, New Zealand mud snails, Himalayan blackberry) seem fully established in some watersheds; the specter of a pragmatic New Normal with diminished native aquatic biodiversity in forests may be realized, owing to our lack of capacity to effectively control invasions everywhere. Furthermore, a clear conflict exists between maintenance of native biodiversity and pursuit of high-value recreational fisheries through nonnative fish-stocking programs.

Nevertheless, restoration tools are being applied to maintain habitats for key native species despite nearby invasive species occurrences (Biebighauser 2011). The solution appears to be purposeful management of the multistate ecosystem across landscapes and regions beyond that which has thus far occurred, in order to designate both wild and nonwild states, in which some places retain a semblance of pristine native ecosystems, whereas in other places different ecosystem services (e.g., fishing experiences) can be fostered. This managed multistate condition is likely part of our regional, if not global, future.

In the Plan area, limited research on ARIS has been conducted recently; several examples of case studies or syntheses follow. First, in a study of invasive fishes in the Willamette River, Oregon, LaVigne et al. (2008) documented an increase in invasive fish diversity and abundance since the 1940s. They also noted the significant contribution to overall fish biomass in the river contributed by the top three most common invasive fishes (smallmouth bass [*Micropterus dolomieu*], largemouth bass [*M. salmoides*], and common carp [*Cyprinus carpio*]). They argued for increased river monitoring and the use of double-pass electrofishing as a means of fish capture and eradication. Carey et al. (2011) assessed the threat to native salmonids posed by smallmouth

bass. They described the tension between conservation of native salmon and angling opportunities provided by invasive warm-water fishes, such as smallmouth bass. They argued for more specific management that targeted locations for native fishes only, and others in which invasive species would be allowed to enhance angling opportunities. This notion supports the wild versus nonwild ecosystem-management approach described above.

Sanderson et al. (2009) completed a comprehensive assessment of the potential effect of invasive species on Pacific salmon in the Pacific Northwest. They found that invasive species may pose an even greater threat to salmonid persistence in the region than the four traditional factors generally thought to affect abundance and survival of native salmonids (habitat alteration, harvest, hatcheries, and the hydrosystem). They considered invasive-species management to be a significant component of salmonid-recovery planning.

Yamada and Gillespie (2008) described how initial assessments of the effects of European green crab (*Carcinus maenas*) invasion in the Pacific Northwest (in 1989), which predicted that the crabs would naturally die out, have been proven incorrect. Instead, they found that changing environmental conditions and coastal currents stemming from El Niño cycles have resulted in shifting habitat characteristics amenable to the crab, promoting its spread. They concluded that management for crab eradication is a more pressing issue than first thought.

Pearl et al. (2013) also showed that invasive crayfish in the Pacific Northwest displace native crayfish. In their work in the Rogue, Umpqua, and Willamette/Columbia River basins, they found that invasive crayfish (in particular, *Procambarus clarkia*) tended to be associated with anthropogenic effects on streams, and that these crayfish appeared to have a negative effect on occupancy of native crayfish. They argued that there is still time to control invasive nonnative crayfish, but that the window of opportunity for management to have a meaningful effect is closing.

Claeson and Bisson (2013) conducted a study of the efficacy of invasive knotweed removal by herbicidal application, and effects on the riparian plant community in western Washington. They found that sites where knotweed had been removed, followed by passive restoration, had

more nonnative species and vegetative cover than reference (no knotweed) sites, and reference sites had more native species. This finding was especially true for riparian areas along larger streams in their sample, as riparian areas along smaller 2nd- to 3rd-order channels had primarily native plant-species assemblages. They suggested active restoration to control secondary invaders, such as replanting native species. Also, they proposed that effectiveness monitoring of invasive species control projects could help to refine and improve restoration approaches.

Kinziger et al. (2014) described establishment pathways of an introduced fish from a nearby source area to two coastal rivers of northern California. Using genetic techniques, they reported that the Eel River invasion of Sacramento pikeminnow (*Ptychocheilus grandis*) likely was the result of only three or four founding individuals, whereas the Elk River invasion likely came from seven founders. This reflects an astounding adaptive capacity for rapid invasion, and highlights the threat posed by such close-range invaders. This species is not included on our priority list (table 7-10).

In a study of western Oregon wetlands, Rowe and Garcia (2014) reported that native anuran amphibians were negatively associated with invasive plant cover, nonnative fish presence, and invasive bullfrog counts. More generally, Bucciarelli et al. (2014) conducted a comprehensive review of the effects of nonnative species on amphibians and broader ecosystem services, painting a complex picture of negative and potentially positive effects. Their conclusion points to the need for additional research on the interactions of native and nonnative species in many ecosystems.

Globally, invasive species experts are using Web-portal technologies to expedite communication among multiple stakeholders, including natural resource management communities, research, and the public sector. Web-portal information can be used to address scientific hypotheses, aid decisionmaking regarding surveillance priorities, and support local-to-regional management actions. To aid communication about emerging invasive pathogens, for example, the Pacific Northwest (PNW) Research Station has partnered with disease and bioinformatics experts internationally to create online data

and mapping portals for ranavirus (<https://mantle.io/grrs>) and the amphibian chytrid fungus *Bd* (Olson et al. 2013) (<https://www.Bd-maps.net>). The online *Bd* database has been used in subsequent research (e.g., Grant et al. 2016, Xie et al. 2016). For land-management applications, the *Bd* point-locality and watershed-scale occurrence maps have been used during firefighting for decisions about which water sources might be used for water draws and whether water disinfection procedures may be necessary (NWCG 2017). A new portal is being populated now with data on both chytrid fungi, *Bd* and *Bsal*, including both planned and completed research and monitoring reports (www.amphibiandisease.org). In addition, EDDMapS (Center for Invasive Species and Ecosystem Health 2017) is a new, national, real-time tracking system for invasive species that employs global positioning system (GPS)-based mobile applications technology to allow users to report invasive species occurrences.

Monitoring and Management

Northwest Forest Plan implementation has required ACS monitoring, conducted by AREMP, with adjustments and modifications over time to address new knowledge and aquatic priorities. AREMP assessments for invasive species began in 2007 during annual field surveys at watersheds across the NWFP area (Gruendike and Lanigan 2008). The initial focus was on 13 species of primary concern for Northwest national forest waterways, with an additional 14 species considered of secondary concern (27 species total). The number of species assessed during annual sampling of watersheds in the Plan area has since fluctuated between 23 and 41 species, with 38 total species (i.e., 31 species groups in table 7-10) being included in the 2017 survey season. Survey methods have also been modified as needed, examining streams and adjacent banks within the bankfull width of the channel during summer low-flow periods. Surveys now include subsampling for benthic snails, mussels, crayfish, and ARIS plants. Few invasive species have been detected annually (table 7-11), with Himalayan blackberry being the most common species reported. Because the design of the AREMP surveys revisits watersheds every 8 years, some of the detections do not represent

Table 7-11—Invasive species detections by the Aquatic-Riparian Effectiveness Monitoring Program (AREMP) of the Northwest Forest Plan, 2007–2016

Year	Watersheds surveyed	Sites surveyed	Number of invasive species detections	Species detected (number of detections)	Reference
2007	31	149	7	Himalayan blackberry (7)	Gruendike and Lanigan 2008
2008	31	167	17	Himalayan blackberry (15) reed canary grass (2)	Gruendike and Lanigan 2009
2009	28	189	17	Himalayan blackberry (14) ringed crayfish (2) Japanese knotweed (1)	Andersen and Lanigan 2010
2010	28	185	7	Himalayan blackberry (4) reed canary grass (2) Robert geranium (1) Also reported common mullein (<i>Verbascum thapsus</i>) from the Region 6 invasive species list	Raggon and Lanigan 2011
2011	29	184	15	Himalayan blackberry (14) English ivy (1)	Raggon and Lanigan 2012a
2012	28	177	10	Himalayan blackberry (9) ringed crayfish (1)	Raggon and Lanigan 2012b
2013	28	187	4	Himalayan blackberry (2) English ivy (1) ringed crayfish (1)	Raggon and Lanigan 2013
2014	27	157	10	Himalayan blackberry (9) herb Robert, geranium (1)	Raggon 2014
2015	34	177	18	Himalayan blackberry (17) parrotfeather watermilfoil (1)	Pennell and Raggon 2016
2016	25	140	10	Himalayan blackberry(10)	Raggon 2017
Overall totals	289	1,712	125	8 species total	
Unique totals	225	1,376			

Note: Ten-year overall totals do not represent unique watersheds and sites, as some resampling among years was conducted as per the AREMP design. Totals of unique watersheds and sites sampled are also provided.

sampling of unique watersheds or sites. Taking unique sites and watersheds into consideration, overall, only 125 invasive species detections of 8 species have occurred across 1,376 unique sites sampled in 225 unique watersheds for the 10 years spanning 2007 to 2016.^{3 4}

³ Hirsch, C. 2017. Personal communication. Fish Program Manager, Siuslaw National Forest, Corvallis, OR 97331, chirsch@fs.fed.us

⁴ Unpublished data. On file with: Aquatic and Riparian Effectiveness Monitoring Program, 3200 SW Jefferson Way, Corvallis, OR 97331.

ARIS monitoring is also a priority for the Aquatic Invasive Species Network (AISN) (AISN 2017). Initiated by the U.S. Geological Survey and Washington State University in 2010 as the Columbia River Basin Aquatic Invasive Species group (CRBAIS 2011), the network integrates federal, state, academic, and tribal organizations over the area of the Columbia River basin, which includes parts of seven states and British Columbia, Canada, and is roughly the size of France. This region overlaps the NWFP area, in watersheds along the western Washington-Oregon border. AISN objectives are to develop an integrated monitoring

and information system, coordinate early-detection efforts, assess invasion pathways, and contribute to an evaluation of the effects of climate change on native and nonnative biota (CRBAIS 2011). Zebra and quagga mussels have been the main taxonomic emphasis; the 2016 map of monitoring sites shows no occurrences of zebra and quagga mussels in the NWFP area (AISN 2017).

In 2012, Forest Service Region 6 developed the Regional Aquatic Invasive Species Strategy and Management Plan (USDA FS 2012b). Its three goals were to (1) prevent new introductions of ARIS into waters and riparian areas of the region; (2) limit the spread of established populations of ARIS into uninfested waters; and (3) provide a cooperative environment that encourages coordinated activities among all affected parties throughout the region. The strategy to achieve these goals is multifaceted, including educational and training programs; implementation of biosecurity protocols (e.g., equipment use and cleaning, inspections); mapping of known invasive species occurrences to inform decisions for water draws for firefighting; coordination of inventory and monitoring efforts; and advance knowledge for eradication procedures. The 2012 list of focal species in this document includes 26 species and species groups listed in table 7-10; known species occurrences were mapped in 2012 (USDA FS 2012b).

As part of the Region 6 ARIS strategy (USDA FS 2012b), surveillance and management is conducted by the joint Region 6 and PNW Research Station dive team.⁵ Four specific incidences of invasive species establishment in the NWFP area have been addressed by the dive team in recent years. First, yellow floating heart infestation at a lake in Rogue River–Siskiyou National Forest, Oregon, was evaluated, and control measures were applied and monitored for efficacy. The plants were pulled out, and the area covered by a geotech-style (hardware weed-control) cloth; after 3 years, the area has remained clear of the plant. Second, Eurasian water milfoil was detected in Coldwater Lake at the Mount St. Helens National Volcanic

Monument, Washington. The plants were pulled out by the dive team and the lake has been monitored for 2 years. Third, in an estuary adjacent to the Siuslaw National Forest, an invasive tunicate was found. The dive team continues to monitor this situation, and, as yet, no control measures have been implemented. Fourth, the dive team partners with the Siuslaw National Forest to survey for freshwater mussels, work that includes documenting occurrences of the Asian clam. Additional surveillance by this regional dive team occurred in summer 2017 within the Plan area on the Deschutes National Forest, Oregon. The dive team also conducts half-day annual training sessions in aquatic-riparian invasive species identification and management for Region 6 personnel, to be applied as stream field crews conduct stream and lake inventories on the national forests.

Management actions implemented when there are known infestations of invasive species on federal lands in the Plan area differ depending on the species considered. In the best scenario, invasive species are identified when their population is small enough for control to be effective. Early Detection and Rapid Response is considered the cornerstone of effective invasive species management (USDA 2017). In situations in which identification of invasive species occurs before the species becomes overly abundant on the landscape, control techniques may be quite effective, as in the above examples implemented by the regional dive team. Additionally, across this region and elsewhere, biosecurity protocols are in effect for personal disinfection of aquatic field gear to prevent spread of emerging diseases and invasive organisms (e.g., Gray et al. 2017, NWCG 2017). In riparian and upland settings, when invasive species have long been present in the environment and are ubiquitous in landscapes surrounding Forest Service land, plans for complete eradication are less feasible. Instead, measures are taken annually to control spread of invasive species, or with the understanding that additional treatments to combat recolonization will be necessary. This is particularly true of several of the invasive riparian and wetland plant species, including reed canary grass, Himalayan blackberry, and purple loosestrife. Mechanical means of control using masticating machines are combined

⁵ Hansen, B. 2017. Personal communication. Ecologist, Pacific Northwest Research Station, Corvallis, OR 97331, bhansen@fs.fed.us

with pesticide applications for the most comprehensive and long-term control of these species.

In addition to invasive species control measures, management actions implemented by Region 6 include development of more effective monitoring frameworks, along with preventative measures to forestall ARIS invasion. To prevent the spread of invasive species, educational programs have been initiated, and biosecurity programs have been implemented. For example, changes have been made to fire equipment contracts that now require equipment such as water tanks to arrive at a fire clean and drained (see ARIS website at <https://www.fs.fed.us/r6/fire/aquatic-invasive-species/>). A two-pronged ARIS monitoring program has recently been developed by the Region 6 and the PNW Research Station. First, to leverage existing freshwater monitoring programs for ARIS detection, a review was completed of ongoing aquatic habitat monitoring, with particular focus on those programs that assess ARIS risk factors. Results of the evaluation found that wadeable stream sections were adequately represented in monitoring on lands administered by Region 6. However, non-wadeable river sections, lakes, and reservoirs that may be at highest risk for invasion by aquatic species were not well represented in existing survey programs (Flitcroft et al. 2016b). The second element was developed as a followup to this finding—a Pilot Monitoring Project of “big water,” began in the summer of 2017. The pilot project will implement multispecies environmental DNA (eDNA) methods to facilitate a consistent and rigorous sampling program. Environmental DNA refers to the residual DNA found in water that is shed by species present in (aquatic species) or near (e.g., tree fungi) the water. Water samples are filtered and then processed in the laboratory, allowing for the identification of DNA from species present in the water sample. Traditionally, eDNA has been used to identify one target species at a time. Techniques being developed by the PNW Research Station in collaboration with Region 6 will allow for up to 48 species to be identified per sample. This approach could provide a breakthrough for ARIS monitoring.

Last, in an attempt to leverage existing resources for ARIS monitoring and mitigation, Region 6 has been able to involve forest law enforcement officers (LEOs) in AIS

monitoring. LEOs interact regularly with users of Forest Service lands, making them ideal partners in ARIS monitoring. For LEOs to have authority to inspect vehicles, trailers, and boats, Region 6 completed a National Environmental Policy Act review in 2016 to activate two relevant federal regulations that prohibit the transfer of animal and plant invasive species across National Forest System lands. LEOs in the region are trained on invasive species inspection and identification. This effort parallels invasive species law enforcement at the state level, in which state fish and wildlife agencies or invasive species councils work with state police as the law enforcement entities to help remove illegal alien and invasive species. For example, in Washington state, the 2015 Report to the Legislature (<http://wdfw.wa.gov/publications/01697/>) reported results from 2011 to 2013, including (1) more than 27,000 boat inspections, with decontamination of 83 boats with aquatic invasive species, of which 19 boats had zebra or quagga mussels, and (2) six new infestations of New Zealand mud snails. The Oregon state “report card” for 2013 (<http://www.oregoninvasivespeciescouncil.org/oregons-report-card>) similarly reports the results of required boat inspections, including a 73 percent compliance rate, mandatory decontamination of ~4 percent (n = 289) of boats because of the presence of invasive plants or animals, including 17 boats decontaminated for quagga or zebra mussels. It is easy to envision how truncating the transmission pathway along our nation’s roadways can be an effective ARIS mitigation.

Future Considerations

Our focus has been on describing those invasive species that have become leading priorities for state or Forest Service management actions in the region of the NWFP, and summarizing recent research and management advances. There is a much longer list of nonnative species detected in western Oregon and Washington that are raising local to widespread concerns. Considerable attention is being paid to novel species in the region, with efforts to prevent introductions of nonnative species into new areas and to understand their potential effects. There are also species that are naturalized to the extent that they are considered long-term, somewhat intractable problems, and that do not

appear on state or other types of species lists; this situation explains why some priority species are identified in some Northwest states, but not others.

Other Nonnative Species on the Radar in the Northwest

It is important to note that species priorities change with time as new knowledge or events trigger new concerns. For example, the 2011 earthquake and tsunami in Japan resulted in an alert for alien aquatic species crossing the Pacific Ocean and reaching the eastern Pacific shores of Oregon and Washington. Heightened attention to potential introductions of aquatic nuisance species has occurred, as over 300 nonnative species have been found on debris from Japan that reached North America through the summer of 2016,⁶ with more than 100 Japanese marine species drifting across the Pacific on a single dock from Misawa, Japan, to Agate Beach, Oregon (Lam et al. 2015). The potential importance of estuaries for such invasions also supports our rationale for their inclusion here.

Barred owls (*Strix varia*) have naturally dispersed into Oregon and Washington from the south and are being recorded as having ripple effects through western forested ecosystems. In addition to interactions affecting the native northern spotted owl (*S. occidentalis caurina*), barred owls have been found to have a broader prey base, including aquatic prey such as crayfish, amphibians, and fish (Wiens et al. 2014). The effects of barred owls on aquatic ecosystems await further research.

Many nonnative fish species have been released in the Western United States for sports fisheries (Schade and Bonar 2005). Effects on native species have been implicated most frequently for amphibians in the lower 48 states (e.g., Knapp and Matthews 2000). Among nonnative fishes released in Oregon and Washington lakes and rivers are smallmouth bass (*Micropterus dolomieu*) and brook trout (*Salvelinus fontinalis*). Both are predators, and effects on native aquatic prey are a concern. For example, bass are implicated as having adverse effects on state-sensitive spe-

cies such as the foothill yellow-legged frogs (*Rana boylei*) in Oregon (Paoletti et al. 2011). Nonnative stocked fishes are an example of a conflict between ecosystem services: recreation versus native species. In the NWFP area, there have been increasing efforts to eradicate nonnative fish from wild areas, such as Crater Lake National Park, Oregon. To date, stocking continues at some historically stocked sites in Oregon and Washington, and native vertebrates in these systems appear to be persisting—yet may have declined from historical numbers or distribution. This is an evolving issue, and monitoring may be needed for sensitive native species under additional stressors such as disease.

Numerous additional species could be mentioned here. Newly identified nonnative species can gain quick attention with the hope of rapid eradication, forestalling a new invasive species gaining a foothold in the region. For example, chemical treatment for African clawed frogs (*Xenopus laevis*) was conducted in 2016 at a pond in Lacey, Washington, (WDFW 2016). The clawed frogs were eradicated, and surveillance is ongoing to continue efforts as needed. The role of human releases of nonnative species into the wild focuses attention on how these animals enter the region, to state laws and their enforcement for alien species generally, and to the pet trade more specifically.

Climate Change Projections

Climate factors (temperature and precipitation regimes) strongly affect seasonal conditions in upslope, riparian, and freshwater environments. As with native species, invasive species survival is also tied to these same parameters, and projected climate change can likewise affect them. Hence, the distributions of both native and invasive species are likely to synchronously respond to changing conditions. Effects on some species have not yet been modeled, but for many species, the effects of climate change on invasions can be assessed. For example, American bullfrogs require water temperatures greater than 20 to 21 °C (68 to 70 °F) for breeding (Hayes and Jennings 2005), and suitable breeding sites are projected to be found at higher latitudes and altitudes in the future. Similarly, a northward expansion of the relatively cold-water-adapted amphibian chytrid fungus *Bd* has been projected with a variety of climate

⁶ Chan, S. 2017. Personal communication. Extension Watersheds and Aquatic Invasive Species. Oregon Sea Grant, Oregon State University, samuel.chan@oregonstate.edu.

futures (Xie et al. 2016). Temperature and precipitation often constrain the range of many invasive plants and limit their successful establishment. With climate change, new habitat may become available, enabling plants to survive outside their historical ranges and expand beyond their current range. For plants, disturbances such as wildfire or logging can provide a “fast-track” for changes in plant communities or even type conversion. For aquatic-dependent species, coldwater refugia are now being considered as localized areas that may be used to more practically protect native species and assemblages from the projected increase in biotic homogenization that is occurring from climate change effects and warm-water species invasions. Finally, climate-related drought and flooding events are also associated with invasive species dispersal; these effects merit additional consideration for the Northwest and elsewhere.

Research and Monitoring Priorities

Despite their increasing recognition as a potentially dominant force in restructuring ecosystems, relatively few research studies have been conducted on Northwest ARIS; their effects on the composition, function, or processes of ecosystems; ecosystem services valued by people; and mitigation efficacies. As evidenced by the above selected studies, support is growing for the importance of invasive species in altering native ecosystems. For example, Sander-son et al. (2009) considered invasive species more important for native salmonids than four other leading concerns combined, including habitat alteration and overexploitation from fishing pressures (harvest).

Several areas stand out as potentially meriting additional research attention. First, aquatic-riparian pathogens and parasites appear underrepresented on Forest Service regional lists of invasive concern species. None are included in AREMP annual surveys, and only tree fungi are included in the Region 6 watch list. Recognition of pathogens and parasites as taxa for regional monitoring could enable their early detection and help forestall invasions. New eDNA techniques could aid in this regard, as detection of cryptic invaders such as pathogens and parasites may otherwise require significant time commitment or costly laboratory analyses applied after disease events are large enough to

be easily detectable. Second, only one study summarized above (Claeson and Bisson 2013) addressed effectiveness of invasive species mitigation approaches with a scientific study design. This is a topic that deserves research for all categories of invasive species in table 7-10, and likely a species-by-species comparison of approaches is needed. For example, field intervention strategies for novel species such as *Bsal* have never been attempted, and foreknowledge of fungicidal or other approaches could be vital to control spread. Nevertheless, trial of some invasive species control methods is ongoing via case-by-case management actions with monitoring, like those being conducted by the regional dive team. Although this adds significantly to our knowledge, it is critical to apply the rigor of hypothesis testing with a scientific design. Lastly, the notion of managing for wild and nonwild ecosystems has been broached, but several questions arise about how this might be developed into an effective long-term strategy. For example, relative to federal lands of the NWFP area, if reserved land use allocations are desired to be wild, can that goal be effectively achieved relative to aquatic-riparian ecosystems given that streams often are contiguous across wild and nonwild areas, potentially promoting invasive species dispersal? Furthermore, can forest restoration practices aid in forestalling nonnative species introductions, or altering the existing heterogeneity within the aquatic ecosystem to avoid crossing “tipping points” to establishment of a novel ecosystem domain (Penaluna et al. 2016)?

Appendix 2: Influence of Climate Change on Life Stages of Pacific Salmon

Peter A. Bisson, Gordon H. Reeves, Nate Mantua, and Steven M. Wondzell¹

Adults

The species of anadromous Pacific salmonids found in the Northwest Forest Plan (NWFP) area and their fresh-water and marine residence times are shown in table 7-8. The freshwater environment is used for both growth and reproduction; the marine environment is used for growth and the initiation of sexual maturity. Depending on species, fish may spend from 1 to 5 or more years in the eastern Pacific Ocean before returning to fresh water to spawn. An exception is coastal cutthroat trout (*Oncorhynchus clarkii*), which generally make limited forays into nearshore areas and typically do not range more than 65 mi (100 km) from natal rivers (Trotter 1989). Owing to marine heterogeneity, the disparate migration patterns of various stocks, and the

widely varying amount of time spent at sea, the influences of climate change on survival and growth of different populations of salmon in the ocean will differ.

Although the specific effects of climate change on marine survival and growth of salmon will depend on the location of their natal rivers and their movements at sea, some trends seem to be common to populations along the Pacific Coast. Possibly as a result of decreasing pH and increasing temperature, salmon are becoming smaller and sometimes younger upon return to fresh water, and exhibit reduced marine survival rates. The size of returning adults of most Pacific salmon species has generally trended downward over the past three decades of the 20th century (Bigler et al. 1996), although there have been multiyear periods when both sizes and abundances have increased (Helle et al. 2007). Some populations of sockeye salmon in Bristol Bay, Alaska, have returned to spawn at a younger age in the second half of the 20th century (Hodgeson et al. 2006, Robards and Quinn 2002).

In the past 69 years, the size of the largest fish caught in a Juneau, Alaska, fishing derby for Chinook salmon (*O. tshawytscha*) has declined (fig. 7-21). Although the origin of these salmon is not known with certainty, it is possible that some originated from rivers in the NWFP area, as migratory routes for some Pacific Northwest Chinook salmon

¹ **Peter Bisson** is a research fish ecologist (retired), U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 3625 93rd Ave. SW, Olympia, WA 98512; **Gordon H. Reeves** is a research fish ecologist and **Steven M. Wondzell** is a research geologist, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis, OR 97331; **Nate Mantua** is a landscape ecologist, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 110 McAllister Way, Santa Cruz, CA 95060.




Figure 7-21—Winning weights of Juneau's Golden North Salmon Derby from 1947 through 2015 and the preceding 42-month average Pacific Decadal Oscillation (PDO) index. Positive deviations in the PDO index occur in warmer-than-average PDO cycles, and negative deviations occur in cooler cycles. See also Fagen (1988) and Reid et al. (2016).

stocks include southeast Alaska. There are many possible explanations for the observed declines in fish size, but several previously published examinations of adult salmon sizes from either commercial fishing records (Helle et al. 2007) or Alaskan fishing derbies (Fagen 1988) attributed at least some of the decline to increased competition, owing to large numbers of hatchery-produced salmon (Bigler et al. 1996, Francis and Hare 1997). However, such relationships are not simple. Helle et al. (2007) analyzed data for different species and stocks from northern Alaska to Oregon and concluded that adult body size resulted from both density-dependent factors (competition) and density-independent factors (environmental conditions). Long-term trends in body size observed over time in a Juneau, Alaska, fishing derby are weakly correlated with the gradually warming 42-month average Pacific Decadal Oscillation (PDO) index observed prior to when the fish were caught (fig. 7-21), suggesting a linkage between ocean conditions and fish size that portends future size declines under a warming climate. The relationship between gradual warming and shifts in the frequency and intensity of PDO fluctuations is unclear, but significant PDO regime shifts can signal major changes in the Earth's biophysical systems (Reid et al. 2016).

Decreases in adult body size resulting from changing environmental conditions in the ocean could also lead to reduced reproductive success. In Pacific salmon, both the number of eggs (Hankin and McKelvey 1985, Healey and Heard 1984) and egg size (Quinn and Vøllestad 2003) are directly related to the weight of adult females. Reproductive capacity of populations could decline if females have fewer eggs (McElhany et al. 2000). Egg size, primarily related to yolk reserve, can also be an adaptation to the environment in which eggs develop. Fish that spawn in warmer areas tend to have larger eggs compared to those from cooler areas because the efficiency of yolk conversion to body tissue is reduced at higher temperatures (Fleming and Gross 1990). The survival and body mass at hatching of eggs incubating at warmer future temperatures could therefore be compromised if egg size does not increase as well.

Food webs in aquatic and riparian ecosystems are supported by the influx of marine-derived nutrients from returning adult salmonids (Bilby et al. 1996, Schindler et al.

2003). The productivity of many streams and rivers within the range of Pacific salmon is influenced by the quantity of marine-derived nutrients from salmon carcasses (Gende et al. 2004, Helfield and Naiman 2001, Willson et al. 2004). A reduction in the size and number of returning adult salmon could compromise the capacity of freshwater ecosystems to produce new salmon, with carryover effects on the wide variety of aquatic and terrestrial organisms that may also benefit from the consumption of eggs during the spawning period (Cederholm et al. 2001, Garner et al. 2009). The growth of juvenile salmon during the spawning season is important for their overwinter survival (Lang et al. 2006). Energy derived from eggs consumed by returning adults can also allow for longer migrations and extended spawning times (Copeland and Venditti 2009); thus fewer, smaller eggs could diminish this potential energy source.

According to climate change predictions for most rivers in the NWFP area, returning adult salmon will face warmer temperatures and lower flows if migrations take place in summer. Some species and life-history types, such as stream-type ("spring") Chinook salmon and summer steelhead in the southern and middle portions of the Pacific Coast range of Pacific salmon, return to fresh water in spring or early-summer months, and hold in rivers and streams for several months before spawning. Adults feed infrequently and usually rest in large pools with cool water. Such pools are not abundant in late summer and early autumn, with coolwater refuges likely to become even less available at those times as climate continues to warm. This circumstance suggests that holding and migrating adults may become increasingly stressed, which will diminish their reproductive potential and increase prespawning mortality. Beechie et al. (2006) believed that the loss of summer prespawn staging habitats in rivers entering Puget Sound, Washington, could result in the replacement of stream-type Chinook salmon by ocean-type Chinook salmon, whose autumn run timing avoids exposure to warm, low-flow summer conditions. For populations undertaking long upstream migrations to spawning grounds, elevated stream temperatures will incur higher metabolic costs and mortality (Rand et al. 2006), and fish that do arrive at spawning grounds may have reduced reproductive capacities (Miller et al. 2011).

Warmer temperatures may also limit gonadal development; Pankhurst et al. (1996) found that female steelhead did not ovulate when temperatures exceeded 70 °F (21 °C). The extirpation of Atlantic salmon in the southern portion of their distributional range is attributed to reproductive failure associated with elevated water temperatures in freshwater spawning areas (McCarthy and Houlihan 1997).

Elevated water temperatures during migration can have indirect effects on returning adults. Returning adults may be more vulnerable to disease and parasites if conditions are warmer in fresh water (Johnson et al. 1996, Ray et al. 2012). However, Stocking et al. (2006) found no relation between water temperature and infection of salmonids with *Ceratomyxa shasta* in the Klamath River, California. Juveniles (Chiaramonte et al. 2016) that are unable to find coolwater holding areas during migration in warmer water may be particularly vulnerable to disease because warm water will favor rapid disease transmission and virulence of warm-adapted pathogens that could lead to fish kills. For example, Miller et al. (2011) presented evidence that elevated temperatures in British Columbia's Fraser River have likely contributed to the virulence of a virus that infects adult sockeye salmon (*O. nerka*) prior to entering the Fraser River, resulting in a high incidence of prespawning mortality.

Rising sea level (IPCC 2007) may affect the reproductive success of species that spawn close to tidewater, particularly some pink (*O. gorbuscha*) and chum (*O. keta*) salmon populations. For small populations that spawn in streams just above the high-tide level, elevated sea levels could reduce the available spawning habitat if suitable spawning sites upstream are inaccessible.

The development and persistence of less favorable ocean conditions could potentially influence the degree of anadromy in populations that possess both anadromous and nonanadromous (fully freshwater-resident) life cycle options. Steelhead, the anadromous form of *O. mykiss*, persist at least in part because there is a fitness advantage associated with migrating to the ocean to feed and returning to fresh water to spawn (Quinn and Myers 2004). If this advantage is reduced or lost, residency could increase in populations, assuming that changes in the freshwater environment are suitable for the persistence of the fresh-

water life-history variant of rainbow trout (Benjamin et al. 2013, Rosenberger et al. 2015, Sloat and Reeves 2014). Other Pacific Coast populations of *O. mykiss* maintain primarily resident populations in locations where the marine environment is believed to be unfavorable for survival and growth, as in southern California (Behnke 2002).

Eggs and Alevins

Eggs and developing embryos will likely be affected by two different aspects of climate change—increased temperatures during egg incubation and altered hydrographs. Under some climate scenarios, winter temperatures are predicted to increase at faster rates than are summer temperatures for Alaska (IPCC 2007), whereas the opposite is true for the more southerly NWFP region (Mote and Salathé 2010).

Most research on climate effects on native fish has focused on the potential for elevated summer temperatures (e.g., Crozier and Zabel 2006, Isaak et al. 2010). However, the effect of elevated winter temperatures may be as, and perhaps even more, pronounced and ecologically significant than increases in summer temperatures. Increased winter temperatures in the NWFP area will result in more precipitation falling as rain rather than snow. Watersheds that historically developed a seasonal snowpack will experience a trend from snow to rain, resulting in more rapid runoff in winter and early spring when snow usually falls, and lower late-spring and early-summer flows owing to reduced snowmelt (Hamlet and Lettenmaier 2007, Hamlet et al. 2005, Tague and Grant 2009). In Washington state's transitional drainage systems that historically possessed both autumn/winter and spring/summer runoff peaks, the shift to a rain-dominant hydrograph is expected to be the most dramatic. Substantial increases are anticipated in the magnitude and frequency of extremely high-flow events in winter, coupled with substantial reductions in summer low flows (e.g., Elsner et al. 2010, Mantua et al. 2010). However, because snowpack will be reduced, rivers with snowmelt-dominated hydrographs could likely see a reduction in the magnitude of high flows during spring runoff. Loukas and Quick (1999) predicted that floods in the snowmelt-dominated continental portions of British Columbia will decrease in magnitude by 7 percent and in volume by 38 percent, and occur as many as 20 days earlier, as a result

of the snow-to-rain transition. In coastal areas, Loukas and Quick (1999) projected that there would be little change in the timing of floods, but that, on average, peak-flow magnitude (+14 percent), flood volume (+94 percent), frequency (+11 percent), and duration (+44 percent) would all increase.

High-flow events will influence egg and alevin survival, depending on the depth of the redd, the size of the female, and the location of spawning in the stream network. Eggs in shallower redds will be more susceptible to being scoured than will those in deeper redds, and smaller salmon often excavate shallower redds than larger salmon (van den Berghe and Gross 1989). It has been speculated that increased peak flows during the incubation period could result in decreased survival of eggs and embryos in populations exposed to hydrologic regimes that have become more prone to gravel-mobilizing flows (Battin et al. 2007)

Potential effects of hydrographs altered by climate change are likely to differ among species and life-history forms. In most drainages of the NWFP area, scour is likely to increase the most in small streams or in confined, steep rivers, affecting fish such as bull trout (*Salvelinus confluentus*) that spawn in the late autumn and early winter when the most severe storms tend to occur along the northwestern Pacific Coast (Isaak et al. 2012). Fish spawning in lower gradient, unconfined areas, such as coho (*O. kisutch*), Chinook, pink, and chum salmon, could be less affected. Studies that have examined potential effects of increased flows on streambed scour (Battin et al. 2007, Leppi et al. 2014, Shanley and Albert 2014) assumed a uniform relationship between flood magnitudes and the vulnerability of salmon populations and their habitat. However, the geographic range of Pacific salmon is characterized by exceptional topographic complexity and watershed dynamism (Montgomery 1999), which can generate considerable diversity in watershed- and stream reach-scale responses of habitat to flood disturbance (Buffington 2012, Montgomery and MacDonald 2002). Thus, effects of increased flows are unlikely to be similar among watersheds or even among reaches within stream networks.

Previous research has demonstrated that stream-channel response potential varies according to position within the dendritic structure of stream networks (Benda et al.

2004), variation in valley and reach-scale confinement (Coulthard et al. 2000, Montgomery and Buffington 1997), and differences among species in their use of habitats created by this physiographical complexity (Goode et al. 2013). In terms of management, floodplain connectivity may ameliorate the effects of future increases in discharge on streambed dynamics. Floodplain connectivity in unconfined reaches provides a “stress release valve” (McKean and Tonina 2013) that limits vulnerability of salmon spawning habitat even in large floods with return intervals of decades to centuries (Goode et al. 2013, Lapointe et al. 2000, McKean and Tonina 2013). In this regard, maintaining or restoring connectivity between streams and adjacent floodplains will mitigate near-term responses to increased flood magnitudes. Additionally, maintaining or restoring channel complexity and hydraulic roughness from large wood may further mitigate the effect of higher flows on salmon spawning habitat (Montgomery et al. 1996, Sloat et al. 2017).

The rate of development of eggs and the size of fish at emergence is related to water temperature. Egg development depends on the accumulation of degree days (Neuheimer and Taggart 2007). Even slight increases in temperature can accelerate rate of development and ultimately result in earlier time of emergence from the gravel (McCullough 1999) (fig. 7-22). Accelerated development leads to smaller




Figure 7-22—Changes in time of emergence of Chinook and coho salmon as a result of a 1 °C-increment increase in water temperature during egg development. From McCullough 1999.

individuals at emergence because metabolic costs decrease the efficiency of yolk use (Beacham and Murray 1990, Elliott and Hurley 1998). Upon emergence, smaller fish are more susceptible to displacement at higher flows. Some fish species may be more influenced by thermal shifts during incubation than others; Beacham and Murray (1990) suggested that coho salmon are adapted for cool water temperatures during development and could experience poorer survival under warming climate scenarios.

There are important ecological implications of climate-related changes in the time and size of fish at emergence. Earlier emergence can result in an extended growing season, a benefit that can lead to increased fitness. Holtby (1988) found that an increase of 1.3 °F (0.7 °C) in winter water temperatures following timber harvest in Carnation Creek on the west coast of Vancouver Island, British Columbia, resulted in coho salmon emerging 6 weeks earlier. Size at age increased because of the extended growing season, resulting in more fish completing their freshwater-rearing life history in one year rather than two. Coho salmon in Carnation Creek also smolted and moved to sea about 2 weeks earlier following timber harvest (which raised stream temperatures); however, marine survival declined, possibly as a result of the decoupling of the timing of smolt migration from marine plankton blooms (Holtby and Scrivener 1989). Similarly, warmer winter temperatures increased the length of the growing season of recently emerged sockeye salmon in southwest Alaska. Like coho salmon in Carnation Creek, sockeye salmon grew faster, and more underwent smolt transformation at age 1+ during warm periods rather than at age 2+ in cooler periods (Schindler et al. 2005). However, age-1+ smolts were smaller than age-2+ smolts and were expected to have decreased marine survival.

Juveniles

Juvenile Pacific salmon (defined here as recently emerged fry up to, but not including, smolts) face a number of challenges from the potential effects of climate change. These challenges will include elevated temperatures and altered streamflows, both of which can affect physical and biological aspects of stream habitats. The type and extent of flow effects will differ depending on the time of emergence. For

example, fish emerging in the late winter and early spring may experience high flows caused by earlier snowmelt. The consequences of a changing hydrograph will depend to a large degree on the geomorphic setting in which spawning and emergence occurs. In some settings, increased flooding could improve use of floodplain habitats when fish in wide, geomorphically unconstrained channels have access to habitats where floodplain vegetation is intact and secondary channels are available.

Low-gradient streams and rivers can be important areas for postemergent and seasonal growth (Brown and Hartman 1988, Moore and Gregory 1988, Peterson 1982a), and marginal areas with reduced water velocities provide refuge against downstream displacement. Fry that emerge at a smaller size if water temperature is warmer can potentially overcome their size disadvantage by gaining an early start on the growing season (Holtby 1988). Juvenile salmonids in rain-dominated hydrographic regimes often move into the lower reaches of the channel network or into off-channel habitats in autumn to seek refuge from unfavorable water velocities in the main channel (Ebersole et al. 2006, Everest 1975, Peterson 1982b, Solazzi et al. 2000). In high-elevation snowfall-dominated drainage systems, however, climate warming might not significantly increase mid-winter flood flows and facilitate access to floodplain habitats if precipitation still falls as snow.

Under several climate scenarios, the onset of the low-flow period is expected to occur up to 4 to 6 weeks earlier in most areas as a result of warming (Hamlet and Lettenmaier 2007, Hamlet et al. 2005, Tague and Grant 2009). An extended period of low discharge over the dry season would likely decrease the amount of habitat suitable to juvenile salmonids, and this effect could be most pronounced in small to mid-sized streams (Stewart et al. 2005), resulting in some reaches that formerly held surface flows throughout the year becoming intermittent or even drying completely. As noted by Battin et al. (2007), flow reductions in headwater areas during the dry season could force resident fishes downstream in the stream network, as well as compromise their ability to cope with drought, by reducing the network of connected, perennially flowing channels. Additionally, the downstream displacement of headwater-rearing fish

will expose them to warmer temperatures than those to which they are adapted, and possibly to harmful biological interactions with native and nonnative species inhabiting the lower watershed.

The consequences of climate-induced changes in low flows for juvenile salmonids such as Chinook salmon and steelhead that often rear in rivers are likely similar to those in smaller streams, although the risk of river reaches becoming intermittent is less because drainage areas are larger. Mantua et al. (2010) found widespread declines in summer discharge for many rivers in Washington state under climatic warming scenarios. Likewise, Luce and Holden (2009) examined hydrographic records from drainage systems throughout the Pacific Northwest and found that summer flows in all types of hydrologic regimes have been declining, thus providing increasingly smaller rearing areas to river-dwelling species.

In addition to lower flows, elevated summer water temperatures will likely have strong ecological effects on juvenile Pacific salmon, with the direction and magnitude of influence varying geographically, by species, and by life-history type. Water temperature influences the metabolism, food consumption, and growth of an individual (Brett et al. 1969, Warren and Davis 1967, Wurtsbaugh and Davis 1977). Age and size of individuals also influence thermal effects; younger and smaller fish are most susceptible to thermal extremes (Brett 1952) and to short-term thermal variation (Elliott 1994). There is a temperature range in which an individual performs best given a certain level of food resources, and beyond that range, metabolic costs increase such that growth declines (Warren 1971). Increased temperature could potentially affect juvenile salmonids in opposing ways (Li et al. 1994). Warmer water could enhance primary and secondary aquatic production, leading to greater food availability; however, if the increased metabolic demands of warmer temperatures reduce food-conversion efficiency or if the organisms benefiting from warmer temperatures are not preferred food items, the net effect of warming could be reduced growth (Bisson and Davis 1976). In southern portions of a species' range, elevated temperatures could reduce the suitability of rearing areas for juveniles during the summer as temperatures exceed

the point at which gains resulting from increased aquatic production are offset by physiological costs, resulting in reduced summer growth rates (Marine and Cech 2004). In contrast, growth rates of juveniles in more northern areas could increase if projected temperature changes stimulate aquatic productivity while remaining within the preferred physiological range for the species.

If the net effects of elevated temperatures resulting from climate change in southern areas reduce summer growth (Isaak et al. 2010, Royer and Minshall 1997, Scarnecchia and Bergersen 1987), juveniles will be smaller entering the winter (ISAB 2007), and overwinter survival may decrease (Quinn and Petersen 1996). However, thermal increases may be beneficial for growth during other seasons if abundant food is present. Sogard et al. (2010) found that juvenile steelhead on the central coast of California attained the most growth in the spring and autumn, and that juvenile coho salmon grew in the winter in coastal Oregon (Ebersole et al. 2006, 2009).

Outcomes of interactions between salmonids and nonsalmonids can be influenced by changing water temperatures. Rearing salmonids tend to outcompete nonsalmonids for food resources and preferred feeding areas at cooler temperatures, whereas nonsalmonids have the advantage at warmer temperatures (Petersen and Kitchell 2001, Reeves et al. 1987). The susceptibility of juvenile salmonids to disease could also increase at warmer temperatures and could be compounded by the presence of competitors that are less susceptible to the pathogens infecting salmon and trout (Reeves et al. 1987). Additionally, warmer temperatures could lead to increased predation from nonnative warmwater fish (ISAB 2007, Petersen and Kitchell 2001). The aggregate results of these indirect effects are likely to be changes in the structure and composition of fish communities in the affected stream systems (ISAB 2012), particularly in the southern portions of the NWFP area where the potential for interaction with warmwater species is greatest owing to widespread introduction and proliferation of nonnative warmwater fishes.

The effects of climate change on rearing habitats for juvenile salmon at the local level will depend, to some degree, on the geomorphic features of a particular location.

Crozier and Zabel (2006) suggested that two climate-influenced factors—stream temperature and flow—could affect habitat in different ways: narrow, confined streams were predicted to be more responsive to flow changes, and geomorphically unconfined streams would be more sensitive to temperature changes. In addition, the future quantity and quality of freshwater rearing habitat of Pacific salmon may also be influenced by predicted increases in the magnitude and frequency of large disturbances. Climate change scenarios predict an increase in exceptional flood events caused by transitions from snow to rain, accelerated glacial melt, wildfires, and forest pathogen outbreaks (Dale et al. 2001, Hamlet and Lettenmaier 2007). Frequent large floods promote landsliding and stream sedimentation in many areas (Miller et al. 2003). The effects of floods and associated erosion events on freshwater habitat will differ depending on the geomorphic setting, the magnitude and legacy of the event, the interval between succeeding disturbances, and the extent to which the affected ecosystem has been altered by past human activities (Reeves et al. 1995, Rieman et al. 2006).

Increased disturbance frequency and severity can have short-term negative consequences for fish populations, including substrate scour and fine-sediment intrusion that reduces egg and alevin survival and macroinvertebrate abundance in confined channels, displacement of juveniles downstream, and loss of surface flow in summer in reaches where porous material has been deposited in the channel. However, in functionally intact systems there is a strong potential for aquatic habitat complexity to improve with flooding because floodplain linkages can be reestablished and large wood will be recruited to the channel network (Bisson et al. 2009). Long-term changes could be favorable to rearing salmon if the cumulative effects of climate change on water temperature, fine-sediment levels, and surface flows remain within limits tolerable to juvenile salmon or exceed those thresholds only for a short duration.

Population productivity after large disturbances will also be enhanced by the presence of adjacent fish populations that provide sources of colonizers to help initiate recovery and that add to the phenotypic and genetic diver-

sity of affected populations (Schtickzelle and Quinn 2007). But it is also possible that in greatly altered watersheds, where the cumulative harmful effects of climate change exceed environmental tolerance limits, the damage caused by large-scale disturbances will be too great, and if there are no nearby populations to provide new colonists, local population extirpation will occur.

Lakes are important rearing habitats for sockeye salmon and will also be affected by climate change, although there are relatively few drainage systems in the NWFP area that support sockeye salmon runs. Potential effects will vary greatly depending on the location and features of the lake, but a primary effect will be the magnitude and seasonality of warming, with epilimnetic water and the timing of spring and autumn turnover experiencing the greatest changes (Stefan et al. 2001). Slight warming of deep lakes could lead to increased sockeye growth rates if temperatures stimulate primary and secondary production without significantly affecting the availability of cooler water during periods when the epilimnion becomes too warm for efficient metabolism. This benefit could be offset during the growing season by a reduction in the delivery of inorganic nutrients and dissolved organic carbon from terrestrial systems as a result of decreased spring and summer flows. Reduced inputs of nutrients and dissolved organic carbon from the surrounding watershed could result in diminished algal production, which would result in deeper light penetration and additional warming of the lake (Schindler et al. 1990).

The productivity of zooplankton, the principal food of juvenile sockeye salmon in lakes, will be affected by climate change, but whether or not the changes are beneficial will depend on ambient thermal and hydrologic regimes. In Alaska, warming temperatures have resulted in earlier ice melt, greater densities of zooplankton, and increasing sockeye growth rates (Schindler et al. 2005). In contrast, earlier onset of spring in western Washington's Lake Washington has advanced lake stratification by 20 days in recent years, resulting in earlier diatom blooms and a decline in cladocerans (*Daphnia* spp.), important prey species for juvenile sockeye rearing in the lake (Winder and Schindler 2004).

Smolts

Anadromous salmonids typically undergo the smolting process and move to the ocean in spring, although seaward migrations of some salmon stocks occur throughout the year. Water temperature, day length, and changes in flow are the principal cues influencing the timing of parr-smolt transformations. Environmental signals affecting smolting can be divided into regulating and controlling factors (Byrne et al. 2004). Regulating factors act on juvenile salmon before the migration and influence the physiological aspects of smolting. Controlling factors operate during migration and affect the speed of downstream movement. Water temperature and day length appear to be key regulating factors (Jonsson and Jonsson 2009). Day length is not influenced by climate change, but increased temperature will affect the onset of smoltification. For Pacific salmon, elevated winter temperatures can result in earlier migration times of smolts. Chinook salmon have been observed to migrate earlier in warmer years than in cooler years (Achord et al. 2007, Roper and Scarnecchia 1999), but Jonsson and Jonsson (2009) cite a suite of other studies on Atlantic salmon, brown trout (*Salmo trutta*), and steelhead in which water temperatures did not affect the timing of smolt migration. Under certain conditions, elevated temperatures may even inhibit parr-smolt transformation. Adams et al. (1973) found that smolting in steelhead held at 59 °F (15 °C) or warmer led to reductions of ATPase activity needed to initiate the smolt transformation process. Thus, the effect of altered temperature on timing of smolt migration remains unpredictable and likely will vary widely across populations.

To a large extent, streamflow determines the rate at which smolts move downstream (Connor et al. 2003, Smith et al. 2002). Climate model projections of stream runoff (Snover et al. 2003, Tague and Grant 2009) suggest that the onset of the low-flow period will occur 4 to 6 weeks earlier over much of the NWFP area in the next century. Projections of the annual cycle of elevated flows from melting snow for more northerly areas are not currently available, but we assume that they will be similar. The consequences of altered flows are likely to be population-specific, with the timing and smolt survival rates of those populations that

tend to migrate later or are required to move long distances likely to be the most affected by climate change.

The survival of smolts entering the ocean depends on a number of factors (Pearcy 1992). Larger smolts tend to have higher survival rates than do smaller fish (Holtby and Scrivener 1989, Quinn and Peterson 1996, Slaney 1988), possibly because they are better able to avoid predation. The size of an individual at smolting is influenced by its size at the beginning of the previous winter. Brown and Hartman (1988) found that stream and groundwater warming caused by logging in a coastal Vancouver Island watershed resulted in increased overwinter growth of presmolt coho salmon, and Holtby and Scrivener (1989) suggested that this growth advantage led to higher smolt-to-adult return rates through improved ocean survival.

Conditions in marine nearshore areas at the time of ocean entry are known to strongly influence ocean survival (Rechisky et al. 2009). In the coastal area influenced by the California current—primarily the southern half of the distributional range of many Pacific salmon species—potential changes in the timing and intensity of upwelling have important implications for smolts (Barth et al. 2007). Cold, nutrient-rich waters are pushed into nearshore areas by northerly winds in the late spring and early summer, producing favorable conditions for plankton production (Nickelson 1986, Scheuerell and Williams 2005). Under one climate change scenario, upwelling is projected to intensify but occur later in the summer (Snyder et al. 2003), decoupling the timing of smolt migration relative to plankton blooms for early-entry salmon smolts.

The abundance of predators in nearshore areas can also influence marine survival of smolts (Pearcy 1992). Coho salmon from Carnation Creek on the west coast of Vancouver Island, British Columbia, entered the ocean about 2 weeks earlier as a result of increased growth as juveniles (Holtby 1988), but survival declined compared to the timing of pre-logging smolt migration. It was believed that predation by mackerel (*Scomber japonicus*) and hake (*Merluccius productus*) contributed to the decline, as both species moved into Barkley Sound during periods of warm sea-surface temperatures. Elevated ocean temperatures could also result in the expansion of subtropical predators

such as the Humboldt squid (*Dosidicus gigas*) into Pacific Northwest waters, further increasing predation pressure on salmon smolts (Christensen and Trites 2011, ISAB 2007).

Nearshore conditions in northern portions of the NWFP area will also be influenced by climate change. In some locations, melting glaciers could increase iron levels in nearshore areas (Westerlund and Ohman 1991). Iron levels are often considered limiting to primary production in the North Pacific, and increased iron levels in freshwater plumes could potentially enhance marine food webs (Rose et al. 2005) and thus improve growth and survival of young salmon. The projected effects of climate change on the ocean ecology of Pacific salmon will therefore result from the combined influences of several factors, notably predation, food resource abundance, and both intra- and interspecific competition.

Scientific and common names of plant species identified in this report

Scientific name	Common name
<i>Abies amabilis</i> (Douglas ex Loudon) Douglas ex Forbes	Pacific silver fir
<i>Abies concolor</i> (Gord. & Glend.) Lindl. ex Hildebr.	White fir
<i>Abies grandis</i> (Douglas ex D. Don) Lindl.	Grand fir
<i>Abies lasiocarpa</i> (Hook.) Nutt.	Subalpine pine
<i>Abies magnifica</i> A. Murray bis	California red fir
<i>Abies procera</i> Rehder	Noble fir
<i>Acer circinatum</i> Pursh	Vine maple
<i>Acer macrophyllum</i> Pursh	Bigleaf maple
<i>Achlys triphylla</i> (Sm.) DC.	Sweet after death
<i>Adenocaulon bicolor</i> Hook.	American trailplant
<i>Alliaria petiolata</i> (M. Bieb.) Cavara & Grande	Garlic mustard
<i>Alnus rubra</i> Bong.	Red alder
<i>Amelanchier alnifolia</i> (Nutt.) Nutt. ex M. Roem.	Saskatoon serviceberry
<i>Anemone oregana</i> A. Gray	Blue windflower
<i>Apocynum cannabinum</i> L.	Dogbane
<i>Arbutus menziesii</i> Pursh	Madrone
<i>Arceuthobium</i> M. Bieb.	Dwarf mistletoe
<i>Arceuthobium occidentale</i> Engelm.	Gray pine dwarf mistletoe
<i>Arceuthobium tsugense</i> Rosendahl	Hemlock dwarf mistletoe
<i>Arctostaphylos nevadensis</i> A. Gray	Pinemat manzanita
<i>Brachypodium sylvaticum</i> (Huds.) P. Beauv.	False brome
<i>Brodiaea coronaria</i> (Salisb.) Engl.	Cluster-lilies
<i>Callitropsis nootkatensis</i> (D. Don) Oerst. ex D.P. Little	Alaska yellow-cedar
<i>Calocedrus decurrens</i> (Torr.) Florin	Incense cedar
<i>Cannabis</i> L.	Marijuana
<i>Carex barbarae</i> Dewey and <i>C. obnupta</i> L.H. Bailey	Sedges
<i>Centaurea solstitialis</i> L.	Yellow starthistle
<i>Chamaecyparis lawsoniana</i> (A. Murray bis) Parl.	Port Orford cedar
<i>Chimaphila menziesii</i> (R. Br. ex D. Don) Spreng.	Little prince's pine
<i>Chimaphila umbellata</i> (L.) W.P.C. Barton	Pipsissewa
<i>Clematis vitalba</i> L.	Old man's beard
<i>Clintonia uniflora</i> Menzies ex Schult. & Schult. f.) Kunth	Bride's bonnet
<i>Coptis laciniata</i> A. Gray	Oregon goldthread
<i>Corylus cornuta</i> Marshall var. <i>californica</i> (A. DC.) Sharp	California hazel
<i>Cornus canadensis</i> L.	Bunchberry dogwood
<i>Cytisus scoparius</i> (L.) Link	Scotch broom
<i>Disporum hookeri</i> (Torr.) G. Nicholson var. <i>hookeri</i>	Drops-of-gold
<i>Fallopia japonica</i> (Houtt.) Ronse Decr. var. <i>japonica</i>	Japanese knotweed
<i>Gaultheria ovatifolia</i> A. Gray	Western teaberry
<i>Gaultheria shallon</i> Pursh	Salal

Scientific name	Common name
<i>Gentiana douglasiana</i> Bong.	Swamp gentian
<i>Geranium lucidum</i> L.	Shining geranium
<i>Geranium robertianum</i> L.	Robert geranium
<i>Goodyera oblongifolia</i> Raf.	Western rattlesnake plantain
<i>Hedera helix</i> L.	English ivy
<i>Heracleum mantegazzianum</i> Sommier & Levier	Giant hogweed
<i>Hesperocyparis sargentii</i> (Jeps.) Bartel	Sargent's cypress
<i>Hieracium aurantiacum</i> L.	Orange hawkweed
<i>Ilex aquifolium</i> L.	English holly
<i>Iris pseudacorus</i> L.	Paleyellow iris
<i>Juniperus occidentalis</i> Hook.	Western juniper
<i>Lamiastrum galeobdolon</i> (L.) Ehrend. & Polatschek	Yellow archangel
<i>Lilium occidentale</i> Purdy	Western lily
<i>Linnaea borealis</i> L.	Twinflower
<i>Lithocarpus densiflorus</i> (Hook. & Arn.) Rehder	Tanoak
<i>Lonicera hispidula</i> Pursh	Honeysuckle
<i>Lupinus albicaulis</i> Douglas	Sickle-keeled lupine
<i>Lycopodium clavatum</i> L.	Running clubmoss
<i>Lythrum salicaria</i> L.	Purple loosestrife
<i>Mahonia nervosa</i> (Pursh) Nutt.	Cascade barberry
<i>Malus fusca</i> (Raf.) C.K. Schneid.	Pacific crabapple
<i>Notholithocarpus densiflorus</i> (Hook. & Arn.) P.S. Manos, C.H. Cannon, & S.H. Oh	Tanoak
<i>Notholithocarpus densiflorus</i> (Hook. & Arn.) P.S. Manos, C.H. Cannon, & S.H. Oh var. <i>echinoides</i> (R.Br. ter) P.S. Manos, C.H. Cannon & S.H. Oh	Shrub form of tanoak
<i>Nuphar polysepala</i> (Engelm.)	Yellow pond lily
<i>Nymphoides peltata</i> (S.G. Gmel.) Kuntze	Yellow floating heart
<i>Osmorhiza chilensis</i> Hook. & Arn.	Sweetcicely
<i>Phalaris arundinacea</i> L.	Reed canarygrass
<i>Picea engelmannii</i> Parry ex Engelm.	Engelmann spruce
<i>Picea sitchensis</i> (Bong.) Carrière	Sitka spruce
<i>Pinus albicaulis</i> Engelm.	Whitebark pine
<i>Pinus attenuata</i> Lemmon	Knobcone pine
<i>Pinus contorta</i> Douglas ex Loudon	Lodgepole pine
<i>Pinus contorta</i> Douglas ex Loudon var. <i>contorta</i>	Beach pine, shore pine
<i>Pinus jeffreyi</i> Balf.	Jeffrey pine
<i>Pinus lambertiana</i> Douglas	Sugar pine
<i>Pinus monticola</i> Douglas ex D. Don)	Western white pine
<i>Pinus ponderosa</i> Lawson & C. Lawson	Ponderosa pine
<i>Populus trichocarpa</i> L. ssp. <i>trichocarpa</i> (Torr. & A. Gray ex Hook) Brayshaw	Black cottonwood
<i>Potamogeton crispus</i> L.	Curly pondweed
<i>Potentilla recta</i> L.	Sulphur cinquefoil

Scientific name	Common name
<i>Prunus emarginata</i> (Douglas ex Hook. D. Dietr.)	Bitter cherry
<i>Pseudotsuga menziesii</i> (Mirb.) Franco	Douglas-fir
<i>Pteridium aquilinum</i> (L. Kuhn)	Brackenfern
<i>Pueraria montana</i> (Lour.) Merr. var. <i>lobata</i> (Willd.) Maesen & S.M. Almeida ex Sanjappa & Predeep	Kudzu
<i>Pyrola asarifolia</i> Sweet	American wintergreen
<i>Quercus agrifolia</i> Née var. <i>oxyadenia</i> (Torr.) J.T. Howell	Coastal live oak
<i>Quercus berberidifolia</i> Liebm.	Scrub oak
<i>Quercus chrysolepis</i> Liebm.	Canyon live oak
<i>Quercus douglasii</i> Hook. & Arn.	Blue oak
<i>Quercus garryana</i> Douglas ex hook.	Oregon white oak
<i>Quercus kelloggi</i> Newberry	California black oak
<i>Quercus lobata</i> Née	Valley oak
<i>Rhamnus purshiana</i> (DC.) A. Gray	Cascara
<i>Rhododendron groenlandicum</i> Oeder	Bog Labrador tea
<i>Rhododendron macrophyllum</i> D. Don ex G. Don	Pacific rhododendron
<i>Ribes lacustre</i> (Pers.) Poir.	Prickly currant
<i>Rubus armeniacus</i> Focke	Himalayan blackberry
<i>Salix exigua</i> Nutt.	Sandbar willow
<i>Senecio bolanderi</i> A. Gray	Bolander's ragwort
<i>Sequoia sempervirens</i> (Lamb. ex D. Don) Endl.	Redwood
<i>Smilacina stellata</i> (L.) Desf.	Starry false Solomon's seal
<i>Synthyris reniformis</i> (Douglas ex Benth.) Benth.	Snowqueen
<i>Taxus brevifolia</i> Nutt.	Pacific yew
<i>Thuja plicata</i> Donn ex D. Don	Western redcedar
<i>Tiarella trifoliata</i> L.	Threeleaf foamflower
<i>Trapa natans</i> L.	Water chestnut
<i>Trillium ovatum</i> Pursh	Pacific trillium
<i>Tsuga heterophylla</i> (Raf.) Sarg.	Western hemlock
<i>Tsuga mertensiana</i> (Bong.) Carrière	Mountain hemlock
<i>Typha latifolia</i> L.	Cattails
<i>Umbellularia californica</i> (Hook. & Arn.) Nutt.	California bay laurel
<i>Vaccinium alaskaense</i> Howell	Alaska blueberry
<i>Vaccinium membranaceum</i> Douglas ex Torr.	Thinleaf huckleberry, big huckleberry
<i>Vaccinium ovatum</i> Pursh	Evergreen huckleberry
<i>Vaccinium oxycoccos</i> L.	Small cranberry
<i>Vaccinium parvifolium</i> Sm.	Red huckleberry
<i>Vancouveria hexandra</i> (Hook.) C. Morren & Decne.	White insideout flower
<i>Xerophyllum tenax</i> (Pursh) Nutt.	Beargrass

Glossary

This glossary is provided to help readers understand various terms used in the Northwest Forest Plan (NWFP) science synthesis. Sources include the Forest Service Handbook (FSH), the Code of Federal Regulations (CFR), executive orders, the Federal Register (FR), and various scientific publications (see “Glossary Literature Cited”). The authors have added working definitions of terms used in the synthesis and its source materials, especially when formal definitions may be lacking or when they differ across sources.

active management—Direct interventions to achieve desired outcomes, which may include harvesting and planting of vegetation and the intentional use of fire, among other activities (Carey 2003).

adaptive capacity—The ability of ecosystems and social systems to respond to, cope with, or adapt to disturbances and stressors, including environmental change, to maintain options for future generations (FSH 1909.12.5).

adaptive management—A structured, cyclical process for planning and decisionmaking in the face of uncertainty and changing conditions with feedback from monitoring, which includes using the planning process to actively test assumptions, track relevant conditions over time, and measure management effectiveness (FSH 1909.12.5). Additionally, adaptive management includes iterative decisionmaking, through which results are evaluated and actions are adjusted based on what has been learned.

adaptive management area (AMA)—A portion of the federal land area within the NWFP area that was specifically allocated for scientific monitoring and research to explore new forestry methods and other activities related to meeting the goals and objectives of the Plan. Ten AMAs were established in the NWFP area, covering about 1.5 million ac (600 000 ha), or 6 percent of the planning area (Stankey et al. 2003).

alien species—Any species, including its seeds, eggs, spores, or other biological material capable of propagating that species, that is not native to a particular ecosystem

(Executive Order 13112). The term is synonymous with exotic species, nonindigenous, and nonnative species (see also “invasive species”).

allochthonous inputs—Material, specifically food resources, that originates from outside a stream, typically in the form of leaf litter.

amenity communities—Communities located near lands with high amenity values.

amenity migration—Movement of people based on the draw of natural or cultural amenities (Gosnell and Abrams 2011).

amenity value—A noncommodity or “unpriced” value of a place or environment, typically encompassing aesthetic, social, cultural, and recreational values.

ancestral lands (of American Indian tribes)—Lands that historically were inhabited by the ancestors of American Indian tribes.

annual species review—A procedure established under the NWFP in which panels of managers and biologists evaluate new scientific and monitoring information on species to potentially support the recommendation of changes in their conservation status.

Anthropocene—The current period (or geological epoch) in which humans have become a dominant influence on the Earth’s climate and environment, generally dating from the period of rapid growth in industrialization, population, and global trade and transportation in the early 1800s (Steffen et al. 2007).

Aquatic Conservation Strategy (ACS)—A regional strategy applied to aquatic and riparian ecosystems across the area covered by the NWFP (Espy and Babbitt 1994) (see chapter 7 for more details).

at-risk species—Federally recognized threatened, endangered, proposed, and candidate species and species of conservation concern. These species are considered at risk of low viability as a result of changing environmental conditions or human-caused stressors.

best management practices (BMPs) (for water quality)—Methods, measures, or practices used to reduce or eliminate the introduction of pollutants and other detrimental impacts to water quality, including but not limited to structural and nonstructural controls and to operation and maintenance procedures.

biodiversity—In general, the variety of life forms and their processes and ecological functions, at all levels of biological organization from genes to populations, species, assemblages, communities, and ecosystems.

breeding inhibition—Prevention of reproduction in healthy adult individuals.

bryophytes—Mosses and liverworts.

canopy cover—The downward vertical projection from the outside profile of the canopy (crown) of a plant measured in percentage of land area covered.

carrying capacity—The maximum population size a specific environment can sustain.

ceded areas—Lands that particular tribes ceded to the United States government by treaties, which have been cataloged in the Library of Congress.

climate adaptation—Management actions to reduce vulnerabilities to climate change and related disturbances.

climate change—Changes in average weather conditions (including temperature, precipitation, and risk of certain types of severe weather events) that persist over multiple decades or longer, and that result from both natural factors and human activities such as increased emissions of greenhouse gases (U.S. Global Change Research Program 2017).

coarse filter—A conservation approach that focuses on conserving ecosystems, in contrast to a “fine filter” approach that focuses on conserving specific species. These two approaches are generally viewed as complementary, with fine-filtered strategies tailored to fit particular species that “fall through the pores” of the coarse filter (Hunter 2005). See also “mesofilter.”

co-management—Two or more entities, each having legally established management responsibilities, working collaboratively to achieve mutually agreed upon, compatible objectives to protect, conserve, use, enhance, or restore natural and cultural resources (81 FR 4638).

collaborative management—Two or more entities working together to actively protect, conserve, use, enhance, or restore natural and cultural resources (81 FR 4638).

collaboration or collaborative process—A structured manner in which a collection of people with diverse interests share knowledge, ideas, and resources, while working together in an inclusive and cooperative manner toward a common purpose (FSH 1909.12.05).

community (plant and animal)—A naturally occurring assemblage of plant and animal species living within a defined area or habitat (36 CFR 219.19).

community forest—A general definition is forest land that is managed by local communities to provide local benefits (Teitelbaum et al. 2006). The federal government has specifically defined community forest as “forest land owned in fee simple by an eligible entity [local government, nonprofit organization, or federally recognized tribe] that provides public access and is managed to provide community benefits pursuant to a community forest plan” (36 CFR 230.2).

community of place or place-based community—A group of people who are bound together because of where they reside, work, visit, or otherwise spend a continuous portion of their time.

community resilience—The capacity of a community to return to its initial function and structure when initially altered under disturbance.

community resistance—The capacity of a community to withstand a disturbance without changing its function and structure.

composition—The biological elements within the various levels of biological organization, from genes and species to communities and ecosystems (FSM 2020).

congeneric—Organisms that belong to the same taxonomic genus, usually belonging to different species.

connectivity (of habitats)—Environmental conditions that exist at several spatial and temporal scales that provide landscape linkages that permit (a) the exchange of flow, sediments, and nutrients; (b) genetic interchange of genes among individuals between populations; and (c) the long-distance range shifts of species, such as in response to climate change (36 CFR 219.19).

consultation (tribal)—A formal government-to-government process that enables American Indian tribes and Alaska Native Corporations to provide meaningful, timely input, and, as appropriate, exchange views, information, and recommendations on proposed policies or actions that may affect their rights or interests prior to a decision. Consultation is a unique form of communication characterized by trust and respect (FSM 1509.05).

corticosterone—A steroid hormone produced by many species of animals, often as the result of stress.

cryptogam—An organism that reproduces by spores and that does not produce true flowers and seeds; includes fungi, algae, lichens, mosses, liverworts, and ferns.

cultural keystone species—A species that significantly shapes the cultural identity of a people, as reflected in diet, materials, medicine, or spiritual practice (Garibaldi and Turner 2004).

cultural services—A type of ecosystem service that includes the nonmaterial benefits that people obtain from ecosystems through spiritual enrichment, cognitive development, reflection, recreation, and aesthetic experiences (Sarukhán and Whyte 2005).

desired conditions—A description of specific social, economic, or ecological characteristics toward which management of the land and resources should be directed.

disturbance regime—A description of the characteristic types of disturbance on a given landscape; the frequency, severity, and size distribution of these characteristic disturbance types and their interactions (36 CFR 219.19).

disturbance—Any relatively discrete event in time that disrupts ecosystem, watershed, community, or species population structure or function, and that changes resources, substrate availability, or the physical environment (36 CFR 219.19).

dynamic reserves—A conservation approach in which protected areas are relocated following changes in environmental conditions, especially owing to disturbance.

early-seral vegetation—Vegetation conditions in the early stages of succession following an event that removes the forest canopy (e.g., timber harvest, wildfire, windstorm), on sites that are capable of developing a closed canopy (Swanson et al. 2014). A nonforest or “pre-forest” condition occurs first, followed by an “early-seral forest” as young shade-intolerant trees form a closed canopy.

ecocultural resources—Valued elements of the biophysical environment, including plants, fungi, wildlife, water, and places, and the social and cultural relationships of people with those elements.

ecological conditions—The biological and physical environment that can affect the diversity of plant and animal communities, the persistence of native species, invasibility, and productive capacity of ecological systems. Ecological conditions include habitat and other influences on species and the environment. Examples of ecological conditions include the abundance and distribution of aquatic and terrestrial habitats, connectivity, roads and other structural developments, human uses, and occurrence of other species (36 CFR 219.19).

ecological forestry—A ecosystem management approach designed to achieve multiple objectives that may include conservation goals and sustainable forest management and which emphasizes disturbance-based management and retention of “legacy” elements such as old trees and dead wood (Franklin et al. 2007).

ecological integrity—The quality or condition of an ecosystem when its dominant ecological characteristics (e.g., composition, structure, function, connectivity, and species composition and diversity) occur within the natural range of

variation and can withstand and recover from most perturbations imposed by natural environmental dynamics or human influence (36 CFR 219.19).

ecological keystone species—A species whose ecological functions have extensive and disproportionately large effects on ecosystems relative to its abundance (Power et al. 1996).

ecological sustainability—The capability of ecosystems to maintain ecological integrity (36 CFR 219.19).

economic sustainability—The capability of society to produce and consume or otherwise benefit from goods and services, including contributions to jobs and market and nonmarket benefits (36 CFR 219.19).

ecoregion—A geographic area containing distinctive ecological assemblages, topographic and climatic gradients, and historical land uses.

ecosystem—A spatially explicit, relatively homogeneous unit of the Earth that includes all interacting organisms and elements of the abiotic environment within its boundaries (36 CFR 219.19).

ecosystem diversity—The variety and relative extent of ecosystems (36 CFR 219.19).

ecosystem integrity—See “ecological integrity.”

ecosystem management—Management across broad spatial and long temporal scales for a suite of goals, including maintaining populations of multiple species and ecosystem services.

ecosystem services—Benefits that people obtain from ecosystems (see also “provisioning services,” “regulating services,” “supporting services,” and “cultural services”).

ectomycorrhizal fungi—Fungal species that form symbiotic relationships with vascular plants through roots, typically aiding their uptake of nutrients. Although other mycorrhizal fungi penetrate their host’s cell walls, ectomycorrhizal fungi do not.

endangered species—Any species or subspecies that the Secretary of the Interior or the Secretary of Commerce has

deemed in danger of extinction throughout all or a significant portion of its range (16 U.S.C. Section 1532).

endemic—Native and restricted to a specific geographical area.

El Niño Southern Oscillation (ENSO)—A band of anomalously warm ocean water temperatures that occasionally develops off the western coast of South America and can cause climatic changes across the Pacific Ocean. The extremes of this climate pattern’s oscillations cause extreme weather (such as floods and droughts) in many regions of the world.

environmental DNA (eDNA)—Genetic material (DNA) contained within small biological and tissue fragments that can be collected from aquatic, terrestrial, and even atmospheric environments, linked to an individual species, and used to indicate the presence of that species.

environmental justice populations—Groups of people who have low incomes or who identify themselves as African American, Asian or Pacific Islander, American Indian or Alaskan Native, or of Hispanic origin.

ephemeral stream—A stream that flows only in direct response to precipitation in the immediate locality (watershed or catchment basin), and whose channel is at all other times above the zone of saturation.

epicormic—Literally, “of a shoot or branch,” this term implies growth from a previously dormant bud on the trunk or a limb of a tree.

epiphyte—A plant or plant ally (including mosses and lichens) that grows on the surface of another plant such as a tree, but is not a parasite.

even-aged stand—A stand of trees composed of a single age class (36 CFR 219.19).

fecundity—The reproductive rate of an organism or population.

federally recognized Indian tribe—An Indian tribe or Alaska Native Corporation, band, nation, pueblo, village, or community that the Secretary of the Interior acknowledges

to exist as an Indian tribe under the Federally Recognized Indian Tribe List Act of 1994, 25 U.S.C. 479a (36 CFR 219.19).

fine filter—A conservation approach that focuses on conserving individual species in contrast to a “coarse filter” approach that focuses on conserving ecosystems; these approaches are generally viewed as complementary with fine-filtered strategies tailored to fit particular species that “fall through the pores” of the coarse filter (Hunter 2005). See also “mesofilter.”

fire-dependent vegetation types—A vegetative community that evolved with fire as a necessary contributor to its vitality and to the renewal of habitat for its member species.

fire exclusion—Curtailed of wildland fire because of deliberate suppression of ignitions, as well as unintentional effects of human activities such as intensive grazing that removes grasses and other fuels that carry fire (Keane et al. 2002).

fire intensity—The amount of energy or heat release during fire.

fire regime—A characterization of long-term patterns of fire in a given ecosystem over a specified and relatively long period of time, based on multiple attributes, including frequency, severity, extent, spatial complexity, and seasonality of fire occurrence.

fire regime, low frequency, high severity—A fire regime with long return intervals (>200 years) and high levels of vegetation mortality (e.g., ~70 percent basal area mortality in forested ecosystems), often occurring in large patches (>10,000 ac [4047 ha]) (see chapter 3 for more details).

fire regime, moderate frequency, mixed severity—A fire regime with moderate return intervals between 50 and 200 years and mixtures of low, moderate, and high severity; high-severity patches would have been common and frequently large (>1,000 ac [>405 ha]) (see chapter 3 for more details).

fire regime, very frequent, low severity—A fire regime with short return intervals (5 to 25 years) dominated by

surface fires that result in low levels of vegetation mortality (e.g., <20 percent basal area mortality in forested ecosystems), with high-severity fire generally limited to small patches (<2.5 ac [1 ha]) (see chapter 3 for more details).

fire regime, frequent, mixed severity—A fire regime with return intervals between 15 and 50 years that burns with a mosaic of low-, moderate-, and high-severity patches (Perry et al. 2011) (see chapter 3 for more details).

fire rotation—Length of time expected for a specific amount of land to burn (some parts might burn more than once or some not at all) based upon the study of past fire records in a large landscape (Turner and Romme 1994).

fire severity—The magnitude of the effects of fire on ecosystem components, including vegetation or soils.

fire suppression—The human act of extinguishing wild-fires (Keane et al. 2002).

floodplain restoration—Ecological restoration of a stream or river’s floodplain, which may involve setback or removal of levees or other structural constraints.

focal species—A small set of species whose status is assumed to infer the integrity of the larger ecological system to which it belongs, and thus to provide meaningful information regarding the effectiveness of a resource management plan in maintaining or restoring the ecological conditions to maintain the broader diversity of plant and animal communities in the NWPf area. Focal species would be commonly selected on the basis of their functional role in ecosystems (36 CFR 219.19).

food web—Interconnecting chains between organisms in an ecological community based upon what they consume.

Forest Ecosystem Management Assessment Team

(FEMAT)—An interdisciplinary team that included expert ecological and social scientists, analysts, and managers assembled in 1993 by President Bill Clinton to develop options for ecosystem management of federal forests within the range of the northern spotted owl (FEMAT 1993).

forest fragmentation—The patterns of dispersion and connectivity of nonhomogeneous forest cover (Riitters et al. 2002). See also “landscape fragmentation” and “habitat fragmentation” for specific meanings related to habitat loss and isolation.

frequency distribution—A depiction, often appearing in the form of a curve or graph, of the abundance of possible values of a variable. In this synthesis report, we speak of the frequency of wildfire patches of various sizes.

fuels (wildland)—Combustible material in wildland areas, including live and dead plant biomass such as trees, shrub, grass, leaves, litter, snags, and logs.

fuels management—Manipulation of wildland fuels through mechanical, chemical, biological, or manual means, or by fire, in support of land management objectives to control or mitigate the effects of future wildland fire.

function (ecological)—Ecological processes, such as energy flow; nutrient cycling and retention; soil development and retention; predation and herbivory; and natural disturbances such as wind, fire, and floods that sustain composition and structure (FSM 2020). See also “key ecological function.”

future range of variation (FRV)—The natural fluctuation of pattern components of healthy ecosystems that might occur in the future, primarily affected by climate change, human infrastructure, invasive species, and other anticipated disturbances.

gaps (forest)—Small openings in a forest canopy that are naturally formed when one or a few canopy trees die (Yamamoto 2000).

genotype—The genetic makeup of an individual organism.

glucocorticoid—A class of steroid hormones produced by many species of animals, often as the result of stress.

goals (in land management plans)—Broad statements of intent, other than desired conditions, that do not include expected completion dates (36 CFR part 219.7(e)(2)).

guideline—A constraint on project and activity decision-making that allows for departure from its terms, so long as

the purpose of the guideline is met (36 CFR section 219.15(d)(3)). Guidelines are established to help achieve or maintain a desired condition or conditions, to avoid or mitigate undesirable effects, or to meet applicable legal requirements.

habitat—An area with the environmental conditions and resources that are necessary for occupancy by a species and for individuals of that species to survive and reproduce.

habitat fragmentation—Discontinuity in the spatial distribution of resources and conditions present in an area at a given scale that affects occupancy, reproduction, and survival in a particular species (see “landscape fragmentation”).

heterogeneity (forest)—Diversity, often applied to variation in forest structure within stands in two dimensions: horizontal (e.g., single trees, clumps of trees, and gaps of no trees), and vertical (e.g., vegetation at different heights from the forest floor to the top of the forest canopy), or across large landscapes (North et al. 2009).

hierarchy theory—A theory that describes ecosystems at multiple levels of organization (e.g., organisms, populations, and communities) in a nested hierarchy.

high-severity burn patch—A contiguous area of high-severity or stand-replacing fire.

historical range of variation (HRV)—Past fluctuation or range of conditions in the pattern of components of ecosystems over a specified period of time.

hybrid ecosystem—An ecosystem that has been modified from a historical state such that it has novel attributes while retaining some original characteristics (see “novel ecosystem”).

hybrid—Offspring resulting from the breeding of two different species.

inbreeding depression—Reduced fitness in a population that occurs as the result of breeding between related individuals, leading to increased homogeneity and simplification of the gene pool.

in-channel restoration—Ecological restoration of the channel of a stream or river, often through placement of materials (rocks and wood) or other structural modifications.

individuals, clumps, and openings (ICO) method—A method that incorporates reference spatial pattern targets based upon individual trees, clumps of trees, and canopy openings into silvicultural prescriptions and tree-marking guidelines (Churchill et al. 2013).

Interagency Special Status and Sensitive Species

Program (ISSSSP)—A federal agency program, established under the U.S. Forest Service Pacific Northwest Region and Bureau of Land Management Oregon/Washington state office. The ISSSSP superseded the Survey and Manage standards and guidelines under the NWFP and also addresses other species of conservation focus, coordinates development and revision of management recommendations and survey protocols, coordinates data management between the agencies, develops summaries of species biology, and conducts other tasks.

intermittent stream—A stream or reach of stream channel that flows, in its natural condition, only during certain times of the year or in several years, and is characterized by interspersed, permanent surface water areas containing aquatic flora and fauna adapted to the relatively harsh environmental conditions found in these types of environments.

invasive species—An alien species (or subspecies) whose deliberate, accidental, or self-introduction is likely to cause economic or environmental harm or harm to human health (Executive Order 13112).

key ecological function—The main behaviors performed by an organism that can influence environmental conditions or habitats of other species.

key watersheds—Watersheds that are expected to serve as refugia for aquatic organisms, particularly in the short term, for at-risk fish populations that have the greatest potential for restoration, or to provide sources of high-quality water.

land and resource management plan (Forest Service)—A document or set of documents that provides management

direction for an administrative unit of the National Forest System (FSH 1909.12.5).

landform—A specific geomorphic feature on the surface of the Earth, such as a mountain, plateau, canyon, or valley.

landscape—A defined area irrespective of ownership or other artificial boundaries, such as a spatial mosaic of terrestrial and aquatic ecosystems, landforms, and plant communities, repeated in similar form throughout such a defined area (36 CFR 219.19).

landscape fragmentation—Breaking up of continuous habitats into patches as a result of human land use and thereby generating habitat loss, isolation, and edge effects (see “habitat fragmentation”).

landscape genetics—An interdisciplinary field of study that combines population genetics and landscape ecology to explore how genetic relatedness among individuals and subpopulations of a species is influenced by landscape-level conditions.

landscape hierarchy—Organization of land areas based upon a hierarchy of nested geographic (i.e., different-sized) units, which provides a guide for defining the functional components of a system and how components at different scales are related to one another.

late-successional forest—Forests that have developed after long periods of time (typically at least 100 to 200 years) following major disturbances, and that contain a major component of shade-tolerant tree species that can regenerate beneath a canopy and eventually grow into the canopy in which small canopy gaps occur (see chapter 3 for more details). Note that FEMAT (1993) and the NWFP also applied this term to older (at least 80 years) forest types, including both old-growth and mature forests, regardless of the shade tolerance of the dominant tree species (e.g., 90-year-old forests dominated by Douglas-fir were termed late successional).

leading edge—The boundary of a species’ range at which the population is geographically expanding through colonization of new sites.

legacy trees—Individual trees that survive a major disturbance and persist as components of early-seral stands (Franklin 1990).

legacies (biological)—Live trees, seed and seedling banks, remnant populations and individuals, snags, large soil aggregates, hyphal mats, logs, uprooted trees, and other biotic features that survive a major disturbance and persist as components of early-seral stands (Franklin 1990, Franklin et al. 2002).

lentic—Still-water environments, including lakes, ponds, and wet meadows.

longitudinal studies—Studies that include repeated observations on the same response variable over time.

lotic—Freshwater environments with running water, including rivers, streams, and springs.

low-income population—A community or a group of individuals living in geographic proximity to one another, or a set of individuals, such as migrant workers or American Indians, who meet the standards for low income and experience common conditions of environmental exposure or effect (CEQ 1997).

managing wildfire for resource objectives—Managing wildfires to promote multiple objectives such as reducing fire danger or restoring forest health and ecological processes rather than attempting full suppression. The terms “managed wildfire” or “resource objective wildfire” have also been used to describe such events (Long et al. 2017). However, fire managers note that many unplanned ignitions are managed using a combination of tactics, including direct suppression, indirect containment, monitoring of fire spread, and even accelerating fire spread, across their perimeters and over their full duration. Therefore, terms that separate “managed” wildfires from fully “suppressed” wildfires do not convey that complexity. (See “Use of wildland fire,” which also includes prescribed burning).

matrix—Federal and other lands outside of specifically designated reserve areas, particularly the late-successional

reserves under the NWFP, that are managed for timber production and other objectives.

mature forest—An older forest stage (>80 years) prior to old-growth in which trees begin attaining maximum heights and developing some characteristic, for example, 80 to 200 years in the case of old-growth Douglas-fir/western hemlock forests, often (but not always) including big trees (>50 cm diameter at breast height), establishment of late-seral species (i.e., shade-tolerant trees), and initiation of decadence in early species (i.e., shade-intolerant trees).

mesofilter—A conservation approach that “focuses on conserving critical elements of ecosystems that are important to many species, especially those likely to be overlooked by fine-filter approaches, such as invertebrates, fungi, and nonvascular plants” (Hunter 2005).

meta-analysis—A study that combines the results of multiple studies.

minority population—A readily identifiable group of people living in geographic proximity with a population that is at least 50 percent minority; or, an identifiable group that has a meaningfully greater minority population than the adjacent geographic areas, or may also be a geographically dispersed/transient set of individuals such as migrant workers or Americans Indians (CEQ 1997).

mitigation (climate change)—Efforts to reduce anthropogenic alteration of climate, in particular by increasing carbon sequestration.

monitoring—A systematic process of collecting information to track implementation (implementation monitoring), to evaluate effects of actions or changes in conditions or relationships (effectiveness monitoring), or to test underlying assumptions (validation monitoring) (see 36 CFR 219.19).

mosaic—The contiguous spatial arrangement of elements within an area. In regions, this is typically the upland vegetation patches, large urban areas, large bodies of water, and large areas of barren ground or rock. However, regional mosaics can also be described in terms of land ownership, habitat

patches, land use patches, or other elements. For landscapes, this is typically the spatial arrangement of landscape elements.

multiaged stands—Forest stands having two or more age classes of trees; this includes stands resulting from variable-retention silvicultural systems or other traditionally even-aged systems that leave residual or reserve (legacy) trees.

multiple use—The management of all the various renewable surface resources of the National Forest System so that they are used in the combination that will best meet the needs of the American people; making the most judicious use of the land for some or all of these resources or related services over areas large enough to provide sufficient latitude for periodic adjustments in use to conform to changing needs and conditions; that some land will be used for less than all of the resources; and harmonious and coordinated management of the various resources, each with the other, without impairment of the productivity of the land, with consideration being given to the relative values of the various resources, and not necessarily the combination of uses that will give the greatest dollar return or the greatest unit output, consistent with the Multiple-Use Sustained-Yield Act of 1960 (16 U.S.C. 528–531) (36 CFR 219.19).

natal site—Location of birth.

native knowledge—A way of knowing or understanding the world, including traditional ecological, and social knowledge of the environment derived from multiple generations of indigenous peoples' interactions, observations, and experiences with their ecological systems. This knowledge is accumulated over successive generations and is expressed through oral traditions, ceremonies, stories, dances, songs, art, and other means within a cultural context (36 CFR 219.19).

native species—A species historically or currently present in a particular ecosystem as a result of natural migratory or evolutionary processes and not as a result of an accidental or deliberate introduction or invasion into that ecosystem (see 36 CFR 219.19).

natural range of variation (NRV)—The variation of ecological characteristics and processes over specified scales of

time and space that are appropriate for a given management application (FSH 1909.12.5).

nested hierarchy—The name given to the hierarchical structure of groups within groups used to classify organisms.

nontimber forest products (also known as “special forest products”)—Various products from forests that do not include logs from trees but do include bark, berries, boughs, bryophytes, bulbs, burls, Christmas trees, cones, ferns, firewood, forbs, fungi (including mushrooms), grasses, mosses, nuts, pine straw, roots, sedges, seeds, transplants, tree sap, wildflowers, fence material, mine props, posts and poles, shingle and shake bolts, and rails (36 CFR part 223 Subpart G).

novel ecosystem—An ecosystem that has experienced large and potentially irreversibly modifications to abiotic conditions or biotic composition in ways that result in a composition of species, ecological communities, and functions that have never before existed, and that depart from historical analogs (Hobbs et al. 2009). See “hybrid ecosystem” for comparison.

old-growth forest—A forest distinguished by old trees (>200 years) and related structural attributes that often (but not always) include large trees, high biomass of dead wood (i.e., snags, down coarse wood), multiple canopy layers, distinctive species composition and functions, and vertical and horizontal diversity in the tree canopy (see chapter 3). In dry, fire-frequent forests, old growth is characterized by large, old fire-resistant trees and relatively open stands without canopy layering.

palustrine—Inland, nontidal wetlands that may be permanently or temporarily flooded and are characterized by the presence of emergent vegetation such as swamps, marshes, vernal pools, and lakeshores.

passive management—A management approach in which natural processes are allowed to occur without human intervention to reach desired outcomes.

patch—A relatively small area with similar environmental conditions, such as vegetative structure and composition. Sometimes used interchangeably with vegetation or forest stand.

Pacific Decadal Oscillation (PDO)—A recurring (approximately decadal-scale) pattern of ocean-atmosphere—a stream or reach of a channel that flows continuously or nearly so throughout the year and whose upper surface is generally lower than the top of the zone of saturation in areas adjacent to the stream.

perennial stream—A stream or reach of a channel that flows continuously or nearly so throughout the year and whose upper surface is generally lower than the top of the zone of saturation in areas adjacent to the stream.

phenotype—Physical manifestation of the genetic makeup of an individual and its interaction with the environment.

place attachment—The “positive bond that develops between groups or individuals and their environment” (Jorgensen and Stedman 2001: 234).

place dependence—“The strength of an individual’s subjective attachment to specific places” (Stokols and Shumaker 1982: 157).

place identity—Dimensions of self that define an individual’s [or group’s] identity in relation to the physical environment through ideas, beliefs, preferences, feelings, values, goals, and behavioral tendencies and skills (Proshansky 1978).

place-based planning—“A process used to involve stakeholders by encouraging them to come together to collectively define place meanings and attachments” (Lowery and Morse 2013: 1423).

plant association—A fine level of classification in a hierarchy of potential vegetation that is defined in terms of a climax-dominant overstory tree species and typical understory herb or shrub species.

population bottleneck—An abrupt decline in the size of a population from an event, which often results in deleterious effects such as reduced genetic diversity and increased probability of local or global extirpation.

potential vegetation type (PVT)—Native, late-successional (or “climax”) plant community that reflects the regional

climate, and dominant plant species that would occur on a site in absence of disturbances (Pfister and Arno 1980).

poverty rate—A measure of financial income below a threshold that differs by family size and composition.

precautionary principle—A principle that if an action, policy, or decision has a suspected risk of causing harm to the public or to the environment, and there is no scientific consensus that it is not harmful, then the burden of proof that it is not harmful falls on those making that decision. Particular definitions of the principle differ, and some applications use the less formal term, “precautionary approach.” Important qualifications associated with many definitions include (1) the perceived harm is likely to be serious, (2) some scientific analysis suggests a significant but uncertain potential for harm, and (3) applications of the principle emphasize generally constraining an activity to mitigate it rather than “resisting” it entirely (Doremus 2007).

prescribed fire—A wildland fire originating from a planned ignition to meet specific objectives identified in a written and approved prescribed fire plan for which National Environmental Policy Act requirements (where applicable) have been met prior to ignition (synonymous with controlled burn).

primary recreation activity—A single activity that caused a recreation visit to a national forest.

probable sale quantity—An estimate of the average amount of timber likely to be awarded for sale for a given area (such as the NWFP area) during a specified period.

provisioning services—A type of ecosystem service that includes clean air and fresh water, energy, food, fuel, forage, wood products or fiber, and minerals.

public participation geographic information system (PPGIS)—Using spatial decisionmaking and mapping tools to produce local knowledge with the goal of including and empowering marginalized populations (Brown and Reed 2009).

public values—Amenity values (scenery, quality of life); environmental quality (clean air, soil, and water); ecological

values (biodiversity); public use values (outdoor recreation, education, subsistence use); and spiritual or religious values (cultural ties, tribal history).

record of decision (ROD)—The final decision document that amended the planning documents of 19 national forests and seven Bureau of Land Management districts within the range of the northern spotted owl (the NWFP area) in April 1994 (Espy and Babbitt 1994).

recreation opportunity—An opportunity to participate in a specific recreation activity in a particular recreation setting to enjoy desired recreation experiences and other benefits that accrue. Recreation opportunities include non-motorized, motorized, developed, and dispersed recreation on land, water, and in the air (36 CFR 219.19).

redundancy—The presence of multiple occurrences of ecological conditions, including key ecological functions (functional redundancy), such that not all occurrences may be eliminated by a catastrophic event.

refugia—An area that remains less altered by climatic and environmental change (including disturbances such as wind and fire) affecting surrounding regions and that therefore forms a haven for relict fauna and flora.

regalia—Dress and special elements made from a variety of items, including various plant and animal materials, and worn for tribal dances and ceremonies.

regulating services—A type of ecosystem service that includes long-term storage of carbon; climate regulation; water filtration, purification, and storage; soil stabilization; flood and drought control; and disease regulation.

representativeness—The presence of a full array of ecosystem types and successional states, based on the physical environment and characteristic disturbance processes.

reserve—An area of land designated and managed for a special purpose, often to conserve or protect ecosystems, species, or other natural and cultural resources from particular human activities that are detrimental to achieving the goals of the area.

resilience—The capacity of a system to absorb disturbance and reorganize (or return to its previous organization) so as to still retain essentially the same function, structure, identity, and feedbacks (see FSM Chapter 2020 and see also “socioecological resilience”). Definitions emphasize the capacity of a system or its constituent entities to respond or regrow after mortality induced by a disturbance event, although broad definitions of resilience may also encompass “resistance” (see below), under which such mortality may be averted.

resistance—The capacity of a system or an entity to withstand a disturbance event without much change.

restoration economy—Diverse economic activities associated with the restoration of structure or function to terrestrial and aquatic ecosystems (Nielsen-Pincus and Moseley 2013).

restoration, ecological—The process of assisting the recovery of an ecosystem that has been degraded, damaged, or destroyed. Ecological restoration focuses on reestablishing the composition, structure, pattern, and ecological processes necessary to facilitate terrestrial and aquatic ecosystems sustainability, resilience, and health under current and future conditions (36 CFR 219.19).

restoration, functional—Restoration of dynamic abiotic and biotic processes in degraded ecosystems, without necessarily a focus on structural condition and composition.

riparian areas—Three-dimensional ecotones (the transition zone between two adjoining communities) of interaction that include terrestrial and aquatic ecosystems that extend down into the groundwater, up above the canopy, outward across the floodplain, up the near slopes that drain to the water, laterally into the terrestrial ecosystem, and along the water course at variable widths (36 CFR 219.19).

riparian management zone—Portions of a watershed in which riparian-dependent resources receive primary emphasis, and for which plans include Plan components to maintain or restore riparian functions and ecological functions (36 CFR 219.19).

riparian reserves—Reserves established along streams and rivers to protect riparian ecological functions and processes

necessary to create and maintain habitat for aquatic and riparian-dependent organisms over time and ensure connectivity within and between watersheds. The Aquatic Conservation Strategy in the NWFP record of decision included standards and guidelines that delineated riparian reserves.

risk—A combination of the probability that a negative outcome will occur and the severity of the subsequent negative consequences (36 CFR 219.19).

rural restructuring—Changes in demographic and economic conditions owing to declines in natural resource production and agriculture (Nelson 2001).

scale—In ecological terms, the extent and resolution in spatial and temporal terms of a phenomenon or analysis, which differs from the definition in cartography regarding the ratio of map distance to Earth surface distance (Jenerette and Wu 2000).

scenic character—A combination of the physical, biological, and cultural images that gives an area its scenic identity and contributes to its sense of place. Scenic character provides a frame of reference from which to determine scenic attractiveness and to measure scenic integrity (36 CFR 219.19).

science synthesis—A narrative review of scientific information from a defined pool of sources that compiles and integrates and interprets findings and describes uncertainty, including the boundaries of what is known and what is not known.

sense of place—The collection of meanings, beliefs, symbols, values, and feelings that individuals or groups associate with a particular locality (Williams and Stewart 1998).

sensitive species—Plant or animal species that receive special conservation attention because of threats to their populations or habitats, but which do not have special status as listed or candidates for listing under the Endangered Species Act.

sensitivity—In ecological contexts, the propensity of communities or populations to change when subject to disturbance, or the opposite of resistance (see “community resistance”).

sink population—A population in which reproductive rates are lower than mortality rates but that is maintained by immigration of individuals from outside of that population (see also “source population”).

social sustainability—“The capability of society to support the network of relationships, traditions, culture, and activities that connect people to the land and to one another, and support vibrant communities” (36 CFR 219.19). The term is commonly invoked as one of the three parts of a “triple-bottom line” alongside environmental and economic considerations. The concept is an umbrella term for various topics such as quality of life, security, social capital, rights, sense of place, environmental justice, and community resilience, among others discussed in this synthesis.

socioecological resilience—The capacity of socioecological systems (see “socioecological system”) to cope with, adapt to, and influence change; to persist and develop in the face of change; and to innovate and transform into new, more desirable configurations in response to disturbance.

socioecological system (or social-ecological system)—A coherent system of biophysical and social factors defined at several spatial, temporal, and organizational scales that regularly interact, continuously adapt, and regulate critical natural, socioeconomic, and cultural resources (Redman et al. 2004); also described as a coupled-human and natural system (Liu et al. 2007).

source population—A population in which reproductive rates exceed those of mortality rates so that the population has the capacity to increase in size. The term is also often used to denote when such a population contributes emigrants (dispersing individuals) that move outside the population, particularly when feeding a sink population.

special forest products—See “nontimber forest products.”

special status species—Species that have been listed or proposed for listing as threatened or endangered under the Endangered Species Act.

species of conservation concern—A species, other than federally recognized as a threatened, endangered, proposed,

or candidate species, that is known to occur in the NWFP area and for which the regional forester has determined that the best available scientific information indicates substantial concern about the species' capability to persist over the long term in the Plan area (36 CFR 219.9(c)).

stand—A descriptor of a land management unit consisting of a contiguous group of trees sufficiently uniform in age-class distribution, composition, and structure, and growing on a site of sufficiently uniform quality, to be a distinguishable unit.

standard—A mandatory constraint on project and activity decisionmaking, established to help achieve or maintain the desired condition or conditions, to avoid or mitigate undesirable effects, or to meet applicable legal requirements.

stationarity—In statistics, a process that, while randomly determined, is not experiencing a change in the probability of outcomes.

stewardship contract—A contract designed to achieve land management goals while meeting local and rural community needs, including contributing to the sustainability of rural communities and providing a continuing source of local income and employment.

strategic surveys—One type of field survey, specified under the NWFP, designed to fill key information gaps on species distributions and ecologies by which to determine if species should be included under the Plan's Survey and Manage species list.

stressors—Factors that may directly or indirectly degrade or impair ecosystem composition, structure, or ecological process in a manner that may impair its ecological integrity, such as an invasive species, loss of connectivity, or the disruption of a natural disturbance regime (36 CFR 219.19).

structure (ecosystem)—The organization and physical arrangement of biological elements such as snags and down woody debris, vertical and horizontal distribution of vegetation, stream habitat complexity, landscape pattern, and connectivity (FSM 2020).

supporting services—A type of ecosystem service that includes pollination, seed dispersal, soil formation, and nutrient cycling.

Survey and Manage program—A formal part of the NWFP that established protocols for conducting various types of species surveys, identified old-forest-associated species warranting additional consideration for monitoring and protection (see "Survey and Manage species"), and instituted an annual species review procedure that evaluated new scientific and monitoring information on species for potentially recommending changes in their conservation status, including potential removal from the Survey and Manage species list.

Survey and Manage species—A list of species, compiled under the Survey and Manage program of the NWFP, that were deemed to warrant particular attention for monitoring and protection beyond the guidelines for establishing late-successional forest reserves.

sustainability—The capability to meet the needs of the present generation without compromising the ability of future generations to meet their needs (36 CFR 219.19).

sustainable recreation—The set of recreation settings and opportunities in the National Forest System that is ecologically, economically, and socially sustainable for present and future generations (36 CFR 219.19).

sympatric—Two species or populations that share a common geographic range and coexist.

threatened species—Any species that the Secretary of the Interior or the Secretary of Commerce has determined is likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range. Threatened species are listed at 50 CFR sections 17.11, 17.12, and 223.102.

timber harvest—The removal of trees for wood fiber use and other multiple-use purposes (36 CFR 219.19).

timber production—The purposeful growing, tending, harvesting, and regeneration of regulated crops of trees to be cut into logs, bolts, or other round sections for industrial or consumer use (36 CFR 219.19).

topo-edaphic—Related to or caused by particular soil conditions, as of texture or drainage, rather than by physiographic or climatic factors within a defined region or area.

traditional ecological knowledge—“A cumulative body of knowledge, practice, and belief, evolving by adaptive processes and handed down through generations by cultural transmission, about the relationship of living beings (including humans) with one another and with their environment” (Berkes et al. 2000: 1252). See also “native knowledge.”

trailing edge—When describing the range of a species, the boundary at which the species’ population is geographically contracting through local extinction at occupied sites.

trophic cascade—Changes in the relative populations of producers, herbivores, and carnivores following the addition or removal of top predators and the resulting disruption of the food web.

uncertainty—Amount or degree of confidence as a result of imperfect or incomplete information.

understory—Vegetation growing below the tree canopy in a forest, including shrubs and herbs that grow on the forest floor.

use of wildland fire—Management of either wildfire or prescribed fire to meet resource objectives specified in land or resource management plans (see “Managing wildfire for resource objectives” and “Prescribed fire”).

variable-density thinning—The method of thinning some areas within a stand to a different density (including leaving dense, unthinned areas) than other parts of the stand, which is typically done to promote ecological diversity in a relatively uniform stand.

vegetation series (plant community)—The highest level of the fine-scale component (plant associations) of potential vegetation hierarchy based on the dominant plant species that would occur in late-successional conditions in the absence of disturbance.

vegetation type—A general term for a combination or community of plants (including grasses, forbs, shrubs, or trees), typically applied to existing vegetation rather than potential vegetation.

viable population—A group of breeding individuals of a species capable of perpetuating itself over a given time scale.

vital rates—Statistics describing population dynamics such as reproduction, mortality, survival, and recruitment.

watershed—A region or land area drained by a single stream, river, or drainage network; a drainage basin (36 CFR 219.19).

watershed analysis—An analytical process that characterizes watersheds and identifies potential actions for addressing problems and concerns, along with possible management options. It assembles information necessary to determine the ecological characteristics and behavior of the watershed and to develop options to guide management in the watershed, including adjusting riparian reserve boundaries.

watershed condition assessment—A national approach used by the U.S. Forest Service to evaluate condition of hydrologic units based on 12 indicators, each composed of various attributes (USDA FS 2011).

watershed condition—The state of a watershed based on physical and biogeochemical characteristics and processes (36 CFR 219.19).

watershed restoration—Restoration activities that focus on restoring the key ecological processes required to create and maintain favorable environmental conditions for aquatic and riparian-dependent organisms.

well-being—The condition of an individual or group in social, economic, psychological, spiritual, or medical terms.

wilderness—Any area of land designated by Congress as part of the National Wilderness Preservation System that was established by the Wilderness Act of 1964 (16 U.S.C. 1131–1136) (36 CFR 219.19).

wildlife—Undomesticated animal species, including amphibians, reptiles, birds, mammals, fish, and invertebrates or even all biota, that live wild in an area without being introduced by humans.

wildfire—Unplanned ignition of a wildland fire (such as a fire caused by lightning, volcanoes, unauthorized and accidental human-caused fires), and escaped prescribed fires.

wildland-urban interface (WUI)—The line, area, or zone where structures and other human development meet or intermingle with undeveloped wildland or vegetation fuels.

Glossary Literature Cited

- Berkes, F.; Colding, J.; Folke, C. 2000.** Rediscovery of traditional ecological knowledge as adaptive management. *Ecological Applications*. 10(5): 1251–1262.
- Brown, G.G.; Reed, P. 2009.** Public participation GIS: a new method for use in national forest planning. *Forest Science*. 55(2): 166–182.
- Carey, A.B. 2003.** Restoration of landscape function: reserves or active management? *Forestry*. 76(2): 221–230.
- Churchill, D.J.; Larson, A.J.; Dahlgreen, M.C.; Franklin, J.F.; Hessburg, P.F.; Lutz, J.A. 2013.** Restoring forest resilience: from reference spatial patterns to silvicultural prescriptions and monitoring. *Forest Ecology and Management*. 291: 442–457.
- Council on Environmental Quality [CEQ]. 1997.** Environmental justice guidance under the National Environmental Policy Act. [Updated]. https://www.epa.gov/sites/production/files/2015-02/documents/ej_guidance_nepa_ceq1297.pdf. (21 December 2017).
- Doremus, H. 2007.** Precaution, science, and learning while doing in natural resource management. *Washington Law Review*. 82: 547–580.
- Espy, M.; Babbitt, B. 1994.** Record of decision: amendments to Forest Service and Bureau of Land Management planning documents within the range of the northern spotted owl. Washington, DC: U.S. Department of Agriculture, Forest Service, and U.S. Department of the Interior, Bureau of Land Management. 74 p.
- Executive Order No. 13112.** 64 FR 6183, February 8, 1999. Invasive species. <https://www.gpo.gov/fdsys/pkg/FR-1999-02-08/pdf/99-3184.pdf>. (June 2017).
- Forest Ecosystem Management Assessment Team [FEMAT]. 1993.** Forest ecosystem management: an ecological, economic, and social assessment. Washington, DC: U.S. Government Printing Office.
- Franklin, J.F. 1990.** Biological legacies: a critical management concept from Mount St. Helens. In: Sabol, K.J., ed. Transactions of the 55th North American wildlands natural resource conference. Wildlife Management Institute: 216–219.
- Franklin, J.F.; Mitchell, R.J.; Palik, B.J. 2007.** Natural disturbance and stand development principles for ecological forestry. Gen. Tech. Rep. GTR-NRS-19. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 44 p.
- Franklin, J.F.; Spies, T.A.; Pelt, R.V.; Carey, A.B.; Thornburgh, D.A.; Berg, D.R.; Lindenmayer, D.B.; Harmon, M.E.; Keeton, W.S.; Shaw, D.C.; Bible, K.; Chen, J. 2002.** Disturbances and structural development of natural forest ecosystems with silvicultural implications, using Douglas-fir forests as an example. *Forest Ecology and Management*. 155(1–3): 399–423.
- Garibaldi, A.; Turner, N. 2004.** Cultural keystone species: implications for ecological conservation and restoration. *Ecology and Society*. 9(3): art1.
- Gosnell, H.; Abrams, J. 2011.** Amenity migration: diverse conceptualizations of drivers, socioeconomic dimensions, and emerging challenges. *GeoJournal*. 76(4): 303–322.
- Hobbs, R.J.; Higgs, E.; Harris, J.A. 2009.** Novel ecosystems: implications for conservation and restoration. *Trends in Ecology & Evolution*. 24(11): 599–605.
- Hunter, M.L. 2005.** A mesofilter conservation strategy to complement fine and coarse filters. *Conservation Biology*. 19(4): 1025–1029.
- Jenerette, G.D.; Wu, J. 2000.** On the definitions of scale. *Bulletin of the Ecological Society of America*. 81(1): 104–105.
- Jorgensen, B.S.; Stedman, R.C. 2001.** Sense of place as an attitude: lakeshore owners attitudes toward their properties. *Journal of Environmental Psychology*. 21(3): 233–248.

- Keane, R.E.; Ryan, K.C.; Veblen, T.T.; Allen, C.D.; Logan, J.A.; Hawkes, B. 2002.** The cascading effects of fire exclusion in Rocky Mountain ecosystems: a literature review. Gen. Tech. Rep. RMRS-GTR-91. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 24 p.
- Liu, J.G.; Dietz, T.; Carpenter, S.R.; Alberti, M.; Folke, C.; Moran, E.; Pell, A.N.; Deadman, P.; Kratz, T.; Lubchenco, J.; Ostrom, E.; Ouyang, Z.; Provencher, W.; Redman, C.L.; Schneider, S.H.; Taylor, W.W. 2007.** Complexity of coupled human and natural systems. *Science*. 317(5844): 1513–1516.
- Long, J.W.; Tarnay, L.W.; North, M.P. 2017.** Aligning smoke management with ecological and public health goals. *Journal of Forestry*. <https://doi.org/10.5849/jof.16-042>. (22 December 2017).
- Lowery, D.R.; Morse, W.C. 2013.** A qualitative method for collecting spatial data on important places for recreation, livelihoods, and ecological meanings: integrating focus groups with public participation geographic information systems. *Society & Natural Resources*. 26(12): 1422–1437.
- Nelson, P.B. 2001.** Rural restructuring in the American West: land use, family and class discourses. *Journal of Rural Studies*. 17(4): 395–407.
- North, M.; Stine, P.A.; O'Hara, K.L.; Zielinski, W.J.; Stephens, S.L. 2009.** An ecosystems management strategy for Sierra mixed-conifer forests, with addendum. Gen. Tech. Rep. PSW-GTR-220. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 49 p.
- Perry, D.A.; Hessburg, P.F.; Skinner, C.N.; Spies, T.A.; Stephens, S.L.; Taylor, A.H.; Franklin, J.F.; McComb, B.; Riegel, G. 2011.** The ecology of mixed severity fire regimes in Washington, Oregon, and northern California. *Forest Ecology and Management*. 262(5): 703–717.
- Pfister, R.D.; Arno, S.F. 1980.** Classifying forest habitat types based on potential climax vegetation. *Forest Science*. 26(1): 52–70.
- Power, M.E.; Tilman, D.; Estes, J.A.; Menge, B.A.; Bond, W.J.; Mills, L.S.; Daily, G.; Castilla, J.C.; Lubchenco, J.; Paine, R.T. 1996.** Challenges in the quest for keystones. *BioScience*. 46(8): 609–620.
- Proshansky, H.M. 1978.** The city and self-identity. *Environment and Behavior*. 10(2): 147–169.
- Redman, C.L.; Grove, J.M.; Kuby, L.H. 2004.** Integrating social science into the long-term ecological research (LTER) network: social dimensions of ecological change and ecological dimensions of social change. *Ecosystems*. 7(2): 161–171.
- Riitters, H.K.; Wickham, D.J.; O'Neill, V.R.; Jones, B.K.; Smith, R.E.; Coulston, W.J.; Wade, G.T.; Smith, H.J. 2002.** Fragmentation of continental United States forests. *Ecosystems*. 5(8): 815–822.
- Sarukhán, J.; Whyte, A. 2005.** Ecosystems and human well-being: synthesis. Millennium Ecosystem Assessment. Washington, DC: Island Press, World Resources Institute. 137 p.
- Stankey, G.H.; Bormann, B.T.; Ryan, C.; Shindler, B.; Sturtevant, V.; Clark, R.N.; Philpot, C. 2003.** Adaptive management and the Northwest Forest Plan: rhetoric and reality. *Journal of Forestry*. 101(1): 40–46.
- Steffen, W.; Crutzen, P.J.; McNeill, J.R. 2007.** The Anthropocene: Are humans now overwhelming the great forces of nature? *AMBIO: A Journal of the Human Environment*. 36(8): 614–621.
- Stokols, D.; Shumaker, S.A. 1982.** The psychological context of residential mobility and well-being. *Journal of Social Issues*. 38(3): 149–171.
- Swanson, M.E.; Studevant, N.M.; Campbell, J.L.; Donato, D.C. 2014.** Biological associates of early-seral pre-forest in the Pacific Northwest. *Forest Ecology and Management*. 324: 160–171.

- Teitelbaum, S.; Beckley, T.; Nadeau, S. 2006.** A national portrait of community forestry on public land in Canada. *The Forestry Chronicle*. 82(3): 416–428.
- Turner, M.G.; Romme, W.H. 1994.** Landscape dynamics in crown fire ecosystems. *Landscape Ecology*. 9(1): 59–77.
- U.S. Department of Agriculture, Forest Service [USDA FS]. 2011.** Watershed condition framework. FS-977. Washington, DC: U.S. Department of Agriculture, Forest Service. 34 p.
- U.S. Global Change Research Program. 2017.** Glossary. [Updated]. <http://www.globalchange.gov/climate-change/glossary>. (11 October 2017).
- Williams, D.R.; Stewart, S.I. 1998.** Sense of place: an elusive concept that is finding a home in ecosystem management. *Journal of Forestry*. 96(5): 18–23.
- Yamamoto, S.-I. 2000.** Forest gap dynamics and tree regeneration. *Journal of Forest Research*. 5(4): 223–229.

Acknowledgments

We thank reviewers from U.S. Forest Service Regions 5 and 6, who provided valuable input on earlier drafts of these chapters. We thank the many anonymous reviewers for their constructive comments. We also thank members of the public, Tribes, and other agencies who provided peer-reviewed literature for consideration, attended the public forums, or provided review comments for peer reviewers to consider. We thank Cliff Duke with the Ecological Society of America for organizing and coordinating the peer review process. We thank Lisa McKenzie and Ty Montgomery with McKenzie Marketing Group for coordinating and summarizing the extensive public input we received and for organizing and facilitating the public forums. We also thank Kathryn Ronnenberg for assisting with figures and editing for some chapters, as well as Keith Olsen for his work on figures. Sean Gordon is thanked for his creation and management of the NWFP literature reference database and

for formatting all the citations in the chapters, which was a very large task. We really appreciate the efforts of Rhonda Mazza, who worked with the authors to write the executive summary. We appreciate the heroic efforts of the entire Pacific Northwest Research Station communications team to get the science synthesis edited and published to meet tight deadlines, especially editors Keith Routman, Carolyn Wilson, and Oscar Johnson and visual information specialist Jason Blake. Borys Tkacz and Jane Hayes are acknowledged for their diligent policy reviews of all chapters. We want to acknowledge the leadership of Paul Anderson and Cindy Miner and Yasmeen Sands for their communications coordination. Finally, the team wishes to thank Region 5, Region 6, and the Pacific Northwest and Pacific Southwest Research Stations for significant funding and other support for this effort.

Pacific Northwest Research Station

Website	http://www.fs.fed.us/pnw/
Telephone	(503) 808-2592
Publication requests	(503) 808-2138
FAX	(503) 808-2130
E-mail	pnw_pnwpubs@fs.fed.us
Mailing address	Publications Distribution Pacific Northwest Research Station P.O. Box 3890 Portland, OR 97208-3890



Federal Recycling Program
Printed on Recycled Paper



U.S. Department of Agriculture
Pacific Northwest Research Station
1220 SW 3rd Ave.
P.O. Box 3890
Portland, OR 97208-3890

Official Business
Penalty for Private Use, \$300