

Betulaceae—Birch family

***Alnus* P. Mill.**

alder

Constance A. Harrington, Leslie Chandler Brodie, Dean S. DeBell, and C. S. Schopmeyer

Dr. Harrington and Ms. Brodie are foresters on the silviculture research team at the USDA Forest Service's Pacific Northwest Research Station, Olympia, Washington; Dr. DeBell retired from the USDA Forest Service's Pacific Northwest Research Station; Dr. Schopmeyer (deceased) was the technical coordinator of the previous manual

Growth habit and occurrence. Alder—the genus *Alnus*—includes about 30 species of deciduous trees and shrubs occurring in North America, Europe, and Asia and in the Andes Mountains of Peru and Bolivia. Most alders are tolerant of moist sites and thus are commonly found along streams, rivers, and lakes and on poorly drained soils; in addition, some species occur on steep slopes and at high elevations. The principal species found in North America are listed in table 1. Many changes in the taxonomy of alder have been made over the years; in this summary, species are referred to by their currently accepted names although in many cases the information was published originally under the synonyms (and alternative common names) listed in table 1.

Although some cultivated European alder is used commercially in the eastern United States, red alder is the largest native species. It is also the most extensively utilized of the native species. Management interest and research activity on red alders have increased dramatically during the past 2 decades, and the resulting information accounts for the majority of new information added to the previous summary on alder seeds prepared by Schopmeyer (1974).

Alders are pioneer species favored by high light levels and exposed mineral soils; in addition, their ability to fix atmospheric nitrogen facilitates establishment on geologically young or disturbed sites with low levels of soil nitrogen (Harrington and others 1994). Dense stands of naturally regenerated red alders established quickly on mudflows associated with the eruption of Mount St. Helens. The trees grew rapidly and soon overtopped other pioneer species such as poplars in the nitrogen-deficient soils (Heilman 1990). Sitka alder plays a similar role in primary succession following deglaciation in Alaska.

Use. Seedlings have been planted successfully for reforestation of coal mining spoil banks (Lowry and others 1962). Soil fertility is improved through fixation of atmospheric nitrogen by microorganisms in the root nodules

(Tarrant and Trappe 1971). Alders also have been planted for wildlife food and cover (Liscinsky 1965) and for ornamental use. European and red alders have been considered for use in biomass plantings for energy (Gillespie and Pope 1994) and are considered excellent firewood. In recent years, harvest and utilization of red alder has expanded greatly on the Pacific Coast of North America, where the species is used for paper products, pallets, plywood, paneling, furniture, veneer, and cabinetry (Harrington 1984; Plank and Willits 1994). Red alder is also used as a fuel for smoking or curing salmon and other seafood and its bark is used to make a red or orange dye (Pojar and MacKinnon 1994). The soft, even-grained wood lacks odor or taste and has been traditionally been used by native peoples, and more recently other woodworkers, to make bowls, eating utensils, and other items (Pojar and MacKinnon 1994). In addition, alder exports have grown from almost nothing in 1990 to more than 153,000 m³ (or 65 million board feet) of lumber annually (Tarrant and others 1994). Several options exist for managing alder in both mixed (Miller and Murray 1978) and pure stands (Tarrant and others 1983), and a summary of management principles and alternative strategies are available for red alder (Hibbs and DeBell 1994).

Geographic races and hybrids. Considerable geographic variation exists among populations of red (Ager and others 1993; Ager and Stettler 1994; Dang and others 1994; Hamann and others 1988; Lester and DeBell 1989), speckled (Bosquet and others 1988), American green (Bosquet and others 1987), and European alders (Funk 1990; Hall and Maynard 1979). Disjunct populations of red alder have been located in Idaho (Johnson 1968), and growth of such populations and those at the extremes of species' range differs markedly from that of most populations (Lester and DeBell 1989). Natural hybridization is common in alder, and zones of introgression between some species can occur where ranges overlap (Ager and Stettler 1994). Artificial hybridization has been conducted with numerous species, including

Table 1—*Alnus*, alder: nomenclature and occurrence

Scientific name(s) & synonyms	Common name(s)	Occurrence
<i>A. glutinosa</i> (L.) Gaertn. <i>A. alnus</i> (L.) Britt. <i>A. rotundifolia</i> Mill.; <i>A. vulgaris</i> Hill <i>Betula alnus</i> var. <i>glutinosa</i> L.	European alder, black alder, European black alder	Native of Europe, northern Africa, & Asia; naturalized locally in parts of E Canada & NE US, cultivated in E, central, & S US
<i>A. incana</i> (L.) Moench <i>Betula alnus</i> var. <i>incana</i> L.	mountain alder, European speckled alder, hoary alder, gray alder	Native of Europe & the Caucasus area; occurs in North America only under cultivation
<i>A. incana</i> ssp. <i>rugosa</i> (Du Roi) Clausen <i>A. incana</i> var. <i>americana</i> Reg. <i>A. glauca</i> Michx. <i>A. rugosa</i> (Du Roi) Spreng. var. <i>americana</i> (Reg.) Fern <i>A. rugosa</i> var. <i>tomophylla</i> (Fern.) Fern. <i>Betula alnus</i> var. <i>rugosa</i> Du Roi	speckled alder, tag alder, swamp alder, <i>aulne blanchâtre</i>	E & central Canada, N central US & in Appalachian Mtns to West Virginia & Maryland
<i>A. incana</i> ssp. <i>tenuifolia</i> (Nutt.) Breitung <i>A. incana</i> var. <i>occidentalis</i> (Dippel) Hitch. <i>A. incana</i> var. <i>virescens</i> S.Wats. <i>A. occidentalis</i> Dippel <i>A. rugosa</i> var. <i>occidentalis</i> (Dippel) Hitch. <i>A. tenuifolia</i> Nutt.	thinleaf alder, mountain alder	Yukon & Alaska S to W Montana & Oregon, in Sierra Nevada to central California, & E to Arizona & New Mexico
<i>A. maritima</i> (Marsh.) Muhl. ex Nutt. <i>A. maritima</i> ssp. <i>metoporina</i> (Furrow) E. Murr <i>A. metoporina</i> Furrow <i>Betula-alnus maritima</i> Marsh.	seaside alder, brook alder	Widely disjunct populations in Delaware, Maryland, & Oklahoma
<i>A. nepalensis</i> D. Don <i>A. boshia</i> Buch.-Hamilt. ex D. Don <i>Clethropsis nepalensis</i> (D. Don) Spach.	Nepal alder, <i>utis</i> , <i>maibao</i>	Native of India & Burma; planted in Hawaii
<i>A. oblongifolia</i> Torr.	Arizona alder, New Mexican alder, <i>aliso</i> (Mexico)	Scattered populations in high mtns of Arizona, New Mexico, & Mexico
<i>A. rhombifolia</i> Nutt. <i>A. rhombifolia</i> var. <i>bernardina</i> Munz & Johnson	white alder, Sierra alder, California alder	Interior of S British Columbia, Washington, Oregon, & Idaho; Sierra Nevada & coastal ranges in California & N Baja California
<i>A. rubra</i> Bong. <i>A. oregona</i> Nutt. <i>A. oregona</i> var. <i>pinnatisecta</i> Starker	red alder, Oregon alder, western alder, Pacific Coast alder	Pacific Coast region from SE Alaska to S California
<i>A. serrulata</i> (Ait.) Willd. <i>A. incana</i> var. <i>serrulata</i> (Ait.) Boivin <i>A. noveboracensis</i> Britt. <i>A. rubra</i> (Marsh.) Tuckerman <i>A. rugosa</i> (Du Roi) Spreng. var. <i>serrulata</i> (Ait.) Winkler <i>A. serrulata</i> var. <i>subelliptica</i> Fern. <i>Betula serrulata</i> (Ait.)	hazel alder, smooth alder, black alder	SW Nova Scotia & central Maine W to Missouri & S to E Texas & Florida

hybrids of red alder with European or mountain alders (Chiba 1966; Hall and Maynard 1979; Ljunger 1959).

Flowering and fruiting. Species in the genus are typically monoecious, with clusters of separate male and female flowers in close proximity. Flower initiation probably occurs during late June or July for both red and European alders (Ager and others 1994; Brown 1986; McVean 1955). The male and female flowers develop into catkins that elongate in late winter or early spring and mature on the previous year's twigs (table 2). For red alders, peak pollen shedding precedes peak female receptivity by only 2 to 4 days (Stettler 1978). For a specific description of staminate and pistillate catkins, see Brayshaw (1976). The strobiles of most species are 10 to 15 mm long when mature (figure 1),

but those of Nepal, red, and Sitka alders are larger, having lengths of 12 to 24 mm (Carlson and Bryan 1959; Funk 1990; Harrington 1990; Krstinic 1994; Townsend and Douglass 1994). They are produced in abundance before trees reach 10 years of age in at least 2 species. European alders can produce flowers by their second growing season, and individual red alder trees are sexually mature at 3 or 4 years. Most dominant trees in a red alder stand will produce seeds by age 6 to 8 years (Harrington and DeBell 1995; Stettler 1978). Although the majority of seeds produced are probably the result of outcrossing, both selfing and apomixis occur in red alder (Stettler 1978). Seed production resulting from selfing has been reported for European and mountain alders; however, in many cases self-fertilization results in


Table 1—*Alnus*, alder: nomenclature and occurrence (Continued)

Scientific name(s) & synonyms	Common name(s)	Occurrence
<i>A. viridis</i> (Vill.) Lam. & DC. <i>A. ovata</i> (Schr.) Lodd. <i>Alnobetula</i> (Ehrh.) K. Koch <i>Betula viridis</i> Vill.	Sitka alder	S Arctic subarctic, and N mountainous regions of North America & Asia
<i>A. viridis</i> ssp. <i>crispa</i> (Ait.) Turrill <i>A. crispa</i> (Ait.) Pursh <i>A. crispa</i> var. <i>elongata</i> Raup. <i>A. crispa</i> var. <i>harricanensis</i> Lepage <i>A. crispa</i> var. <i>mollis</i> (Fern.) Fern. <i>A. crispa</i> var. <i>stragula</i> Fern. <i>A. mollis</i> Fern. <i>A. viridis</i> var. <i>crispa</i> (Michx.) House <i>A. alnobetula</i> var. <i>crispa</i> (Michx.) Winkler <i>Betula crispa</i> (Ait.)	American green alder, green alder, mountain alder	Labrador to Alberta, S to Minnesota & New England
<i>A. viridis</i> ssp. <i>fruticosa</i> (Rupr.) Nyman* <i>A. fruticosa</i> Rupr. <i>A. viridis</i> var. <i>fruticosa</i> (Rupr.) Reg.	Siberian alder	Alaska S to British Columbia & Alberta, disjunct populations in Washington, Oregon, & N California
<i>A. viridis</i> ssp. <i>sinuata</i> (Regel) A. Löve & D. Löve <i>A. crispa</i> ssp. <i>sinuata</i> (Reg.) Hultén <i>A. sinuata</i> (Reg.) Rydb. <i>A. sitchensis</i> (Reg.) Sarg. <i>A. viridis</i> var. <i>sinuata</i> Reg.	Sitka alder, mountain alder, wavyleaf alder	Yukon & Alaska S to N California & W Montana; also in E Asia

Sources: Schopmeyer (1974), FNAEC (1997).

* In western North America, Siberian alder (*A. viridis* ssp. *fruticosa*) has long been mistaken for American green alder (*A. v. ssp. crispa*), which it closely resembles, or for Sitka alder (*A. v. ssp. sinuata*) (FNAEC 1997).

Figure 1—*Alnus*, alder: mature female catkins (strobiles) of *A. rhombifolia*, white alder (**left**); *A. serrulata*, hazel alder (**right**).



aborted ovules (Krstinic 1994). Information on the effects of management practices on reproductive processes is limited. In young red alder plantings in western Washington, flowering varied by half-sib family but overall was reduced in close spacings and by summer irrigation (Harrington and DeBell 1995). However, dry weather in spring reduced germination rates of European alder seeds, making irrigation early in the year desirable when precipitation is below normal (Hall and Nyong 1987).

Seed production varies from year to year, site to site, and tree to tree (Ager and others 1994; Brown 1985, 1986; Lewis 1985; Koski and Tallquist 1978; Krstinic 1994; McGee 1993), but good crops are borne at least once every 4 years (table 3). LaBastide and van Vredenburg (1970) reported that seed crops for European alder follow an annually alternating pattern. McVean (1955) concluded that seed crops of European alder could vary substantially from year to year, but that “boom-and-bust” patterns of seed production were not typical. Complete failure of a seedcrop is rare, but after a severe freeze in November 1955, almost no red alder seeds were produced in 1956 (Worthington 1965).

Seeds are small nuts (“nutlets”) borne in pairs on the bracts of the strobiles. The nuts of red, Siberian, and Sitka alders have broad wings about as wide as or wider than the body of the nut. In the other species included here, the wings are reduced to a narrow border (figure 2) (Fernald 1950; Sargent 1965). Seeds are without endosperm and contain only small cotyledons (figure 3). For additional information on reproductive biology of red alders, see Ager and others (1994).

The factors regulating the timing of seed dispersal in alders have not been investigated, but they are probably similar to those regulating the release of seeds from the cones of conifers; that is, once strobiles are mature, disper-

Table 2—*Alnus*, alder: phenology of flowering and fruiting*

Species	Location	Flowering	Fruit ripening	Seed dispersal
<i>A. glutinosa</i>	E US S US & England	Mar–May (can start Jan)	Sept Feb–April	Sept or Oct–early spring —
<i>A. incana</i>	Europe	Mar–May	Sept–Nov	Sept–Dec
<i>ssp. rugosa</i>	Canada, US	Mar–May	—	—
<i>ssp. tenuifolia</i>	Idaho, Montana, Oregon	Mar–Apr	Aug–Sept	—
<i>A. nepalensis</i>	Hawaii	—	Oct–Feb	Oct–Apr
<i>A. rhombifolia</i>	Oregon	Mar	Late Sept–early Oct	—
<i>A. rubra</i>	Washington, Oregon	Late winter– early spring	Aug–Oct	Sept–Dec
<i>A. serrulata</i>	—	Feb–May	Late Sept–early Oct	—
<i>A. viridis</i>				
<i>ssp. crispa</i>	E US, Alaska	Spring Apr–June	Late Aug–mid-Oct Mid Sept–early Oct	Soon after ripening Sept–early spring
<i>ssp. sinuata</i>	Alaska, W Canada, & NW US	Apr–June	Sept–Dec	—

Sources: Densmore (1979), Fernald (1950), Funk (1990), Harrington (1990), Hitchcock and others (1964), Lewis (1985), McDermott (1953), McGee (1988), McVean (1955), Schopmeyer (1974), White (1981).

* Flowering occurs during the period when leaves unfold.

Table 3—*Alnus*, alder: growth habit, height, seed-bearing age, and seedcrop frequency

Species	Growth habit	Height at maturity (m)	Year first cultivated	Minimum seed-bearing age (yrs)	Years between large seedcrops
<i>A. glutinosa</i>	Tree	to 35	1866	6–7	—
<i>A. incana</i>	Tree	to 20	—	under 25	1–4
<i>ssp. rugosa</i>	Tree or shrub	to 8	—	—	—
<i>ssp. tenuifolia</i>	Tree or shrub	1–9	1880	—	—
<i>A. nepalensis</i> (Hawaii)	Tree	15–30	1916	10	—
<i>A. rhombifolia</i>	Tree	20–25	1885	—	—
<i>A. rubra</i>	Tree	12–27	1884	3–4	3–5
<i>A. serrulata</i>	Tree or shrub	to 8	1769	—	—
<i>A. viridis</i>					
<i>ssp. crispa</i>	Shrub	to 3	1782	—	—
<i>ssp. sinuata</i>	Tree or shrub	to 12	1903	—	—

Sources: Carlson and Bryan (1959), Fernald (1950), Funk (1990), Harrington (1990), Sargent (1965), Schopmeyer (1974).

sal is determined by the occurrence of weather that dries them, thus opening scales and allowing the seeds to be released (Harrington and others 1994). In general, wet weather following dry weather closes the strobiles, thus terminating a dispersal event. Nonetheless, heavy seedfall can occur during wet weather under certain conditions (Lewis 1985), but dispersal will not occur if ice freezes the seeds in the strobile. Although most seed dispersal occurs from September or October through February to April (table 2), some red alder seedfall has been observed in all months (Lewis 1985). American green alder strobiles do not release many seeds if the weather is wet during the autumn; substantial seed dispersal onto snow can occur throughout the winter (Densmore 1979). Alder seeds are very light, and when released they are dispersed long distances by wind, and in some species by water. Seeds of European alder have

remained viable after floating for 12 months in still water (McVean 1955). In Alaska, seeds of thinleaf alder have corky, thick wings and float for long periods of time, whereas seeds of American green alder have thinner wings and sink rapidly (Densmore 1979). Birds or other animals also act as dispersal agents when moving through alder crowns and when extracting seeds from the strobiles (Harrington and others 1994).

Information on damaging agents is limited. Fungal diseases of alder catkins—caused by *Taphrina occidentalis* Ray and *T. alni* (Berk. & Broome) Gjaerum—cause enlargements of the bracts of female catkins (Mix 1949) and thus prevent or hinder normal fertilization and seed development. Jumping plant lice—*Psylla alni* (L.)—lay eggs in alder catkins in western North America (Furniss and Carolin 1977). Alder seeds are an important source of food for some

Figure 2—*Alnus*, alder: nuts (seeds); *A. viridis* ssp. *crispa*, American green alder (**top left**); *A. glutinosa*, European alder (**top right**); *A. nepalensis*, Nepal alder (**middle left**); *A. rhombifolia*, white alder (**middle center**); *A. rubra*, red alder (**middle right**); *A. serrulata*, hazel alder (**bottom left**);





Figure 3—*Alnus rubra*, red alder: longitudinal section through a nut.



bird species (White and West 1977), and presumably seed predation by birds could have significant impacts when seedcrops are small.

Collection of fruits, extraction and cleaning, and storage of seeds. Seedcrops can be assessed in mid-summer by obtaining a count of mature strobiles and filled seeds (Ager and others 1994). Filled seed count should be determined from the upper third of the crown where viability is highest (Brown 1985). Seed quality can be assessed by cutting the strobile longitudinally and counting the filled

seeds on one of the cut faces. Although the number of filled seeds on a cut face can vary from 0 to 20 or more in red alders, less than 3 or 4 seeds per cut face indicate a marginal crop (Ager and others 1994). Strobiles may be collected from standing or recently felled trees when the bracts (scales) start to separate on the most mature strobiles. In red alders, ripeness can be judged by twisting the cone along the long axis; if it twists easily and the scales part slightly, the seeds are sufficiently mature for collection (Hibbs and Ager 1989). Color is also a good indicator of maturity; immature cones are green whereas mature cones are mottled shades of green, yellow, gray, or brown (Hibbs and Ager 1989). Strobiles should be collected as soon as they are ripe, for the largest seeds with the best germinability are usually released first. Thus, both seed quality and seed yield are higher if collections are made in the fall rather than in the winter or spring (Lewis 1985; Krstinic 1994). Alder cones will open after being dried on screens or in fine mesh bags in a well-ventilated room for several weeks at ambient air temperature. They can be opened in a shorter time (2 to 7 days) by drying them in a kiln at 16 to 27 °C. Higher temperatures should not be used, as the strobiles will dry too quickly, harden and not open completely. Most of the seeds fall out of the strobiles during the drying process. The remainder, if needed, may be extracted by shaking or tumbling. Overall seed yields can be improved by either wetting cones again, placing them in a cooler for 24 hours, or spraying them with a fine water mist and then redrying (Ager and others 1994). Seeds may be cleaned by screening to remove large trash and further processing with an air column to remove small extraneous material.

Purity as high as 90% has been attained with European alder by fanning and screening seedlots. Quality, however, may be low because the light weight of alder seeds makes it difficult to separate and remove empty seeds (Ager and others 1994). Soundness in most cleaned seedlots has been between 30 and 70% (table 4). Number of seeds per weight ranges from 660,000 to 2,816,000/kg (or 300,000 to 1,277,000/lb) in lots of average quality (table 4). Except for seeds of American green alder, higher numbers may indicate a low percentage of filled seeds. Numbers ranging from 1,800,000 to 4,400,000 seeds/kg (800,000 to 2,000,000/lb) have been found in samples of Nepal, red, and thinleaf alders, but less than 5% of the seeds in these samples were full (Schopmeyer 1974). One red alder seedlot, however, was 70% sound and had 2,700,000 seeds/kg (1,224,000/lb). In a trial with red alder, the percentage of filled seeds determined by x-radiography was highly correlated ($r^2 = 0.91$)

Table 4—*Alnus*, alder: yield data and soundness

Species	Seed wt/ vol of strobiles		Cleaned seeds/ wt of strobiles		Cleaned seeds/ vol of strobiles		Cleaned seeds (x1,000)/wt of strobiles				Soundness	
	kg/hl	lb/bu	kg/100 kg	lb/100 bu	kg/hl	lb/bu	Range		Average		Samples	(%)
							/kg	/lb	/kg	/lb		
<i>A. glutinosa</i> Pennsylvania Europe	—	—	—	—	1.5	1.2	565–882 635–1,406	257–401 289–639	706 774	321 352	7 86	— 39
<i>A. incana</i> Europe	21–30	16–23	8–10	8–10	—	—	961–1,980	437–900	1,470	668	123	51
<i>A. incana</i> <i>spp. rugosa</i>	—	—	—	—	4.8	3.7	—	—	660	299	—	30–60
<i>spp. tenuifolia</i> (fresh)*	14	11	7	7	1.3	1.0	—	—	—	—	—	—
(air dry)*	8	6	13	13	1.0	0.8	—	—	1,485	673	1	—
<i>A. rhombifolia</i> (fresh)*	23	18	5	5	1.3	1.0	—	—	—	—	—	—
(air dry)*	9	7	13	13	1.3	1.3	1,349–1,511	613–687	1,430	650	2	71
<i>A. rubra</i>	9	7	14–15	14–15	0.1–1.4	0.1–1.1	1,843–2,700	350–1,400	1,712	776	5	70
<i>A. viridis</i> <i>spp. crispa</i>	—	—	—	—	—	—	1,530–4,101	694–1,860	2,816	1,277	2	42–93
<i>spp. sinuata</i>	—	—	—	—	—	—	—	—	2,200	998	1	70

Sources: Hibbs and Ager (1989), Liscinsky (1965), Mirov and Kraebel (1939), Schopmeyer (1974), Niemc and others (1995), USDA data on file at Olympia Forestry Sciences Laboratory.
* Yield data were determined on clusters of strobiles including stems.

with the actual germination percentage (Ager and others 1994).

Air-dried seeds have been stored in sealed containers at $< -2^{\circ}\text{C}$. Under these conditions, viability has been maintained for 2 years in seeds of European alder (Holmes and Buszewicz 1958) and for 10 years in speckled alder (Heit 1967). For long-term storage, however, further drying of seeds to moisture content of less than 10% has been recommended for red alder (Ager and others 1994). This can be accomplished by kiln-drying or placing seeds in a room at 27°C with less than 25% relative humidity. Red alder seeds can then be placed in moisture-proof containers and stored at $< -12^{\circ}\text{C}$ for 10 to 20 years without substantial losses in viability (Ager and others 1994).

Pregermination treatments and germination tests. The degree of dormancy appears to vary among alder species and among provenances (geographic origins) within species. Thus, percentage germination of fresh seeds of white and thinleaf alders was equally good for stratified and nonstratified seed (Schopmeyer 1974). Fresh seeds of European and mountain alders also germinated promptly without stratification; but dried seeds, at a moisture of content of 8 to 9%, were dormant (table 5) (Schalin 1967). Germination capacity of the dried seeds, after stratification for 180 days at 5°C was higher than that of fresh seeds. Maximum germination capacity, however, was obtained only when the stratification period was followed by 3 days at -20°C (table 5) (Schalin 1967). A more recent study found that fresh mountain alder seeds initially exhibited some dormancy (that is, only about half of the filled seeds germinated in the incubator), but no dormancy was observed after one winter in the soil (Granstrom 1987). Dormancy also has been encountered in occasional seedlots of speckled (Heit 1968) and American green alders (Schopmeyer 1974). Stratification for 30 to 60 days at 1 to 5°C has been recommended for these dormant lots (Schopmeyer 1974). Stratification for 30 to 90 days also has been recommended for Sitka alder (Emery 1988).

Although physiological seed dormancy is not widespread in red alder, it can exist (Elliot and Taylor 1981). Stratification at low temperature (0 to 5°C) has little or no effect on the rate or completeness of germination of red alder seeds when tested at warm germination temperatures (Elliot and Taylor 1981; Radwan and DeBell 1981; Tanaka and others 1991).

Table 5—*Alnus*, alder: stratification and germination testing data

Species	Cold stratification period* (days)	Germination test conditions			Germination rate		Germination Avg (%)	Samples	Soundness (%)
		Temp (°C)		Days	Amount (%)	Days			
		Day	Night						
A. glutinosa (Pennsylvania)	0	30	21	28	—	—	52	7	—
A. glutinosa (Finland)									
fresh seed	0	25	25	21	21	21	28	1	43
dried seed	0	25	25	21	9	5	13	1	43
dried seed	180	25	25	21	27	5	35	1	43
dried seed	180+3†	25	25	21	35	5	46	1	43
A. incana (Europe)	0	21	21	30	—	—	45	100	—
A. incana (Finland)									
fresh seeds	0	25	25	21	21	5	29	1	45
dried seeds	0	25	25	21	12	5	16	1	45
dried seeds	180	25	25	21	25	5	34	1	45
dried seeds	180+3†	25	25	21	38	5	49	1	45
A. i. ssp. tenuifolia									
fresh seeds	0	30	20	26	4	13	4	1	6
A. rhombifolia									
fresh seeds	0	30	20	30	59	14	59	1	65
A. rubra									
dried seeds	0	24	16	7	56	7	56	4	—
fresh seeds	0–60‡	30§	20	28	18	7	71	6	—
fresh seeds	0	30	20	28	21	7	75	6	87
fresh seeds	14	30	20	28	42	7	72	6	87
fresh seeds	28	30	20	28	49	7	72	6	87
fresh seeds	0	15	5	56	0	21	16	6	87
fresh seeds	14	15	5	56	17	21	63	6	87
fresh seeds	28	15	5	56	54	21	80	6	87
A. serrulata		27	23	10	27	2	36	1	—
A. viridis									
ssp. crispa	60	30	20	30–40	28	12	28	3	30–40
ssp. sinuata	14	30	20	21	5	7	14	1	—

Sources: ISTA (1993), McDermott (1953), Radwan and DeBell (1981), Schalin (1967), Schopmeyer (1974), Tanaka and others (1991), data on file at Olympia Forestry Sciences Laboratory.

Note: Day/night, 8 hrs/16 hours.

* Stratification, when used, was in a moist medium at 1 to 5 °C.

† 180 days at 5 °C, plus 3 days at 20 °C.

‡ No difference for 0, 30, or 60 days of stratification.

§ Light period was 10 hours/day at this temperature.

|| Seeds were stratified for an unspecified period.

Under cool temperatures similar to those likely to prevail during outdoor sowings in early spring, however, 2 to 4 weeks of stratification substantially enhanced rate of germination and total germination (Tanaka and others 1991) and such a period is therefore recommended (Ager and others 1994). Thinleaf and American green alder seedlots collected near Fairbanks, Alaska, also germinated well without stratification at 25 °C but only germinated well at lower temperatures (10 to 15 °C) when combined with 72 days of stratification (Densmore 1979). Studies have also indicated the potential of 3 quick pregermination treatments for red alder seeds: gibberellin (Berry and Torrey 1985), 1% captan (Berry and Torrey 1985), and 30% hydrogen peroxide (Neal and others 1967). The results from these pregermination treatments, however, were obtained under warm germination conditions and need to be tested under the cooler conditions encountered in spring sowings. The captan and peroxide treatments may have a beneficial effect by reducing the amount of disease organisms present on seedcoats. Pretreatment with gibberellic acid improved greenhouse germination (21 °C day/13 °C night) of thinleaf alder seeds from 2 sources but did not affect germination of Arizona alder seeds from a single source (Dreesen and Harrington 1997).


For germination testing, both constant temperatures and diurnally alternating temperatures have been used (table 5). Official tests of the International Seed Testing Association (ISTA 1993) call for a 21-day test at alternating temperatures of 20/30 °C, with light during the 8 hours at 30 °C. Although seeds of European alder germinated as well in continuous darkness as under normal day length (McVean 1955), recent work indicates that seed germination of many alder species is markedly affected by light regime (Berry and Torrey 1985; Boojh and Ramakrishnan 1981; Bormann 1983; Densmore 1979; Khan and Tripathi 1989). Such effects in red alder are mediated by phytochrome: red light stimulates seed germination, far-red light inhibits it, and the effect of each light treatment can be reversed by the alternative treatment (Bormann 1983). Seeds of red alder are also sensitive to amount and quality of light under field conditions, and these factors—along with soil moisture—control germination success on disturbed sites (Haeussler and Tappeiner 1993; Haeussler and others 1995).

Nursery practice. Alder seedlings have been produced by bareroot nursery (open field or bedhouse) and container methods, as well as combinations thereof (Ahrens 1994; Ahrens and others 1992; Funk 1990; Radwan and others 1992). Successful stock types for red alder are grown in 1 year and include 1+0 open-bed bareroot, 1+0 bedhouse

bareroot, 1+0 plug, and +0.5 (plug+transplant). Most nurseries sow in the spring when growing alder species (Ahrens and others 1992; Schopmeyer 1974), but fall-sowing is mentioned by Heit (1968). Spring-sowing is sometimes delayed until late spring to reduce seedling size. Sowing depths of 2 to 5 mm (.1 to .2 in) have been used for seeds of European alder and red alder (Schopmeyer 1974). In California, seeds of red alder have been mixed with 10 parts of vermiculite and drilled 1 cm (.4 in) deep (Schopmeyer 1974). In Oregon, seeds of red alder have also been sown on the soil surface and covered with peat. Seeds of Nepal alder have been mixed with sand and spread over the nursery beds. The number of plantable seedlings obtained from 1 kg (2.2 lb) of seed was 22,000 (10,000/lb) for European alder and 88,000 (40,000/lb) for hazel alder (Van Dersal 1938). Germination is epigeal (figure 4).

Alder seedlings, particularly those of red alder, grow rapidly and seedling densities should be lower than those used for conifers. Seedlings grown at open-bed densities of 60 to 180 seedlings/m² (5 to 15/ft²) or in large containers result in much better outplanting performance than those grown at greater densities or in small Styroblocks® (Ahrens 1994). Inoculation of beds or container media with the nodulating actinomycete *Frankia* can improve establishment

Figure 4—*Alnus glutinosa*, European alder: seedling development at 1 and 7 days after germination (**left**); *Alnus incana* ssp. *tenuifolia*, thinleaf alder: 2 older seedlings (**right**).



and early growth in the nursery (Berry and Torrey 1985; Hilger and others 1991) and may enhance outplanting performance (McNeill and others 1990). Diluted suspensions of pure *Frankia* cultures and homogenates of crushed, fresh root nodules have been used for inoculation (Ahrens and others 1992; Perinet and others 1985). Detailed methods of preparation and application are available (Martin and others 1991; Molina and others 1994; Zasada and others 1991).

Development of nitrogen-fixing nodules is promoted by fertilization with low to moderate applications of nitrogen; phosphorus and lime are likely to be necessary for production of high-quality stock (Hughes and others 1968; Radwan 1987; Radwan and DeBell 1994). Although alder seedlings are produced operationally, optimum combinations of fertilizer source, amount, and timing of application have not been completely worked out; some combinations have had detrimental effects on alder seedlings or their root associates. Frequent irrigation may be necessary to prevent desiccation and heat damage of surface-sown seeds or germinants during germination and early establishment (Ahrens 1994).

Direct seeding in the field has been done successfully with 2 species. Speckled alder has been established in Pennsylvania by broadcast sowing on disked areas and on sod. Seeds collected in the fall were broadcast during the following February and March. Seeding rates were 0.28

liter/10 m² (or 0.5 pint/100 ft²) on bare soil and 0.38 liter (0.7 pint) for the same area of sod (Liscinsky 1965). In England, better stocking was obtained on a shallow blanket bog with spot sowing of European alder than with broadcast sowing. About 15 viable seeds were sown in each spot and fertilized with about 60 g of rock phosphate (McVean 1959).

Seedling care. Information to guide lifting dates is very limited, even for red alder (Ahrens 1994; Ahrens and others 1992); current recommendations based on experience in southwest Washington are to lift seedlings in January. They are then stored at either +2 °C or –2 °C; the lower temperature is recommended because it prevents budbreak during storage (and possible *Botrytis* infection associated with budbreak during storage) and reduces the tendency for planted alders to break bud too soon after planting. Storage in sealed bags will prevent desiccation. Because alder stems are brittle and sensitive, seedlings must be handled carefully during storage, transport, and outplanting to avoid damage to stems, branches, and buds. At low elevations (< 300 m) in western Washington, it has been recommended that seedlings be planted between mid-March and mid-April. The spring planting period should begin when the probability of severe frost is low and end before there is appreciable soil drying (Dobkowski and others 1994).

References

- Ager AA, Heilman PE, Stettler RF. 1993. Genetic variation in red alder (*Alnus rubra*) in relation to native climate and geography. *Canadian Journal of Forest Research* 23: 1930–1939.
- Ager AA, Stettler RF. 1994. Genetics of red alder and its implications for future management. In: Hibbs DE, DeBell DS, Tarrant RF, eds. *The biology and management of red alder*. Corvallis: Oregon State University: 92–105.
- Ager AA, Tanaka Y, McGrath J. 1994. Biology, ecology, and utilization of red alder seed. In: Hibbs DE, DeBell DS, Tarrant RF, eds. *The biology and management of red alder*. Corvallis: Oregon State University: 159–169.
- Ahrens GR. 1994. Seedling quality and nursery practices for red alder. In: Hibbs DE, DeBell DS, Tarrant RF, eds. *The biology and management of red alder*. Corvallis: Oregon State University: 170–185.
- Ahrens GR, Dobkowski A, Hibbs DE. 1992. Red alder: guidelines for successful regeneration. Spec. Pub. 24. Corvallis: Oregon State University, College of Forestry. 11 p.
- Berry AM, Torrey JG. 1985. Seed germination, seedling inoculation and establishment of *Alnus* spp. in containers in greenhouse trials. *Plant and Soil* 87(1): 161–173.
- Boojh R, Ramakrishnan PS. 1981. Germination of seeds of *Alnus nepalensis* Don. *National Academy Science Letters* 4(2): 53–56.
- Bormann BT. 1983. Ecological implications of phytochrome mediated seed germination in red alder. *Forest Science* 29(4): 734–738.
- Bousquet J, Cheliak WM, Lalonde M. 1987. Genetic differentiation among 22 mature populations of green alder (*Alnus crispa*). *Canadian Journal of Forest Research* 17: 219–227.
- Bousquet J, Cheliak WM, Lalonde M. 1988. Allozyme variation within and among mature populations of speckled alder (*Alnus rugosa*) and relationships with green alder (*A. crispa*). *American Journal of Botany* 75: 1678–1686.
- Brayshaw TC. 1976. Catkin bearing plants (Amentiferae) of British Columbia. Occ. Pap. 18. Victoria: British Columbia Provincial Museum: 127–135.
- Brown SM. 1986. Sexual allocation patterns in red alder (*Alnus rubra* Bong.) along three elevational transects [MS thesis]. Seattle: University of Washington, College of Forest Resources. 241 p.
- Brown SM. 1985. A study of reproductive biology of *Alnus rubra* along three elevational transects in Washington and Oregon. On file with: USDA Forest Service, Pacific Northwest Research Station, Olympia, WA. 48 p.
- Carlson NK, Bryan LW. 1959. Hawaiian timber for the coming generations. Honolulu: Trustees for the Bernice P. Bishop Estate. 112 p.
- Chapin FS III, Walker LR, Fastie CL, Sharman LC. 1994. Mechanisms of primary succession following deglaciation at Glacier Bay, Alaska. *Ecological Monographs*, 64(2): 149–175.
- Chiba S. 1966. Studies on the tree improvement by means of artificial hybridization and polyploidy in *Alnus* and *Populus* species. Bulletin of the Oji Institute for Forest Tree Improvement [Kuriyama, Japan: Aji Paper Co.] 1: 1–165.
- Dang QL, Xie CY, Ying C, Guy RD. 1994. Genetic variation of ecophysiological traits in red alder (*Alnus rubra* Bong.). *Canadian Journal of Forest Research* 24: 2150–2156.
- Densmore RVE. 1979. Aspects of the seed ecology of woody plants of the Alaskan taiga and tundra [PhD dissertation]. Durham, NC: Duke University, Department of Botany: 34–48.
- Dobkowski A, Figueroa PF, Tanaka Y. 1994. Red alder plantation establishment. In: Hibbs DE, DeBell DS, Tarrant RF, eds. *The biology and management of red alder*. Corvallis: Oregon State University: 186–201.

- Dreesen DR, Harrington JT. 1997. Propagation of native plants for restoration projects in the southwestern U.S.: preliminary investigations. In: Landis TD, Thompson JR, tech. coords. National Proceedings, Forest and Conservation Nursery Associations. Gen. Tech. Rep. PNW-419. Portland, OR: USDA Forest Service, Pacific Northwest Research Station: 77–88.
- Elliott DM, Taylor IEP. 1981. Germination of red alder (*Alnus rubra*) seed from several locations in its natural range. Canadian Journal of Forest Research 11: 517–521.
- Emery D. 1988. Seed propagation of native California plants. Santa Barbara, CA: Santa Barbara Botanic Garden. 115 p.
- Fernald ML. 1950. Gray's manual of botany. 8th ed. New York: American Book Co. 1632 p.
- FNAEC [Flora of North America Editorial Committee]. 1997. Flora of North America north of Mexico, Volume 3, Magnoliophyta: Magnoliidae and Hamamelidae. New York: Oxford University Press. 616 p.
- Funk DT. 1990. *Alnus glutinosa* (L.) Gaertn., European alder. In: Burns RM, Honkala BH, tech. coords. Silvics of North America. Volume 2, Hardwoods. Washington, DC: USDA Forest Service: 105–115.
- Furniss R, Carolin VM. 1977. Western forest insects. Misc. Pub. 1339. Washington, DC: USDA Forest Service. 654 p.
- Gillespie AR, Pope PE. 1994. Intensive culture of European black alder in central Indiana, U.S.A.: biomass yield and potential returns to farmers. Biomass and Bioenergy 6(6): 419–430.
- Granstrom A. 1987. Seed viability of fourteen species during five years of storage in forest soil. Journal of Ecology 75(2): 321–331.
- Haeussler S, Tappeiner JC II. 1993. Effect of the light environment on seed germination of red alder (*Alnus rubra*). Canadian Journal of Forest Research 23: 1487–1491.
- Haeussler S, Tappeiner JC II, Greber BJ. 1995. Germination, survival, and early growth of red alder seedlings in the central coast range of Oregon. Canadian Journal of Forest Research 25: 1639–1651.
- Hall RB, Maynard CA. 1979. Considerations in the genetic improvement of alder. In: Gordon JC and others, eds. Symbiotic nitrogen fixation in the management of temperate forests. Corvallis: Oregon State University: 322–344.
- Hall RB, Nyong RN. 1987. Design, establishment and management of a black alder (*Alnus glutinosa* L. Gaertn.) seed orchard. In: Proceedings, 19th Southern Forest Tree Improvement Conference. 1987 June 16–18; College Station, TX. Springfield, VA: USDC National Technical Information Service: 261–268.
- Hamann A, El-Kassaby YA, Koshy MP, Namkoong G. 1998. Multivariate analysis of allozymic and quantitative trait variation in *Alnus rubra*: geographic patterns and evolutionary implications. Canadian Journal of Forest Research 28: 1557–1565.
- Harrington CA. 1984. Red alder: an American wood. FS-215. Washington, DC: USDA Forest Service. 7 p.
- Harrington CA. 1990. *Alnus rubra* Bong., red alder. In: Burns RM, Honkala BH, tech. coords. Silvics of North America. Volume 2, Hardwoods. Agric. Handbk. 654. Washington, DC: USDA Forest Service: 116–123.
- Harrington CA, DeBell DS. 1995. Effects of irrigation, spacing and fertilization on flowering and growth in young *Alnus rubra*. Tree Physiology 15: 427–432.
- Harrington CA, Zasada JC, Allen EA. 1994. Biology of red alder (*Alnus rubra* Bong.). In: Hibbs DE, DeBell DS, Tarrant RF, eds. The biology and management of red alder. Corvallis: Oregon State University: 3–22.
- Heilman PE. 1990. Growth of Douglas-fir and red alder on coal spoils in western Washington. Soil Science Society of America Journal 54: 522–527.
- Heit CE. 1967. Propagation from seed: 11. Storage of deciduous tree and shrub seeds. American Nurseryman 126(10): 12–13, 86–94.
- Heit CE. 1968. Propagation from seed: 15. Fall planting of shrub seeds for successful seedling production. American Nurseryman 128(4): 8–10, 70–80.
- Hibbs DE, Ager AA. 1989. Red alder: guidelines for seed collection, handling, and storage. Spec. Pub. 18. Corvallis: Oregon State University, Forest Research Laboratory. 6 p.
- Hibbs DE, DeBell DS. 1994. Management of young red alder. In: Hibbs DE, DeBell DS, Tarrant RF, eds. The biology and management of red alder. Corvallis: Oregon State University: 202–215.
- Hilger AB, Tanaka Y, Myrold DD. 1991. Inoculation of fumigated nursery soil increases nodulation and yield of bare-root red alder (*Alnus rubra* Bong.). New Forests 5: 35–42.
- Hitchcock CL, Cronquist A, Ownbey M, Thompson JW. 1964. Vascular plants of the Pacific Northwest: Volume 2, Salicaceae to Saxifragaceae. Seattle: University of Washington Press: 72–76.
- Holmes GD, Buszewicz G. 1958. The storage of seed of temperate forest tree species: 2. Forestry Abstracts 19: 455–476.
- Hughes DR, Gessel SP, Walker RB. 1968. Red alder deficiency symptoms and fertilizer trials. In: Trappe JM, Franklin JF, Tarrant RF, Hansen GM, eds. Biology of alder. Proceedings, 40th Annual Meeting of the Northwest Science Association; 1967 April 14–15; Pullman, WA. Portland, OR: USDA Forest Service, Pacific Northwest Forest and Range Experiment Station: 225–237.
- ISTA [International Seed Testing Association]. 1993. International rules for seed testing: rules 1993. Seed Science and Technology 21 (Suppl.): 1–259.
- Johnson FD. 1968. Disjunct populations of red alder in Idaho. In: Trappe JM, Franklin JF, Tarrant RF, Hansen GM, eds. Biology of alder. Proceedings, 40th Annual Meeting of the Northwest Scientific Association; 1967 April 14–15; Pullman, WA. Portland, OR: USDA Forest Service, Pacific Northwest Forest and Range Experiment Station: 9–22.
- Khan ML, Tripathi RS. 1989. Effect of soil moisture, soil texture and light intensity on emergence, survival and growth of seedlings of a few sub-tropical trees. Indian Journal of Forestry 12(3): 196–204.
- Koski V, Tallquist R. 1978. Results of long-time measurements of the quantity of flowering and seed crop of forest trees. Folia Forestalia: 1–344.
- Krstinic A. 1994. Genetics of black alder (*Alnus glutinosa* (L.) Gaertn.). Annales Forestales [Zagreb] 19(2): 33–72.
- la Bastide JGA, van Vredenburg CLH. 1970. The influence of weather conditions on the seed production of some forest trees in the Netherlands. Stichting Bosbouwproefstation 'De Dorschkamp' Mededeling 102: 1–12.
- Lester DT, DeBell DS. 1989. Geographic variation in red alder. Res. Pap. PNW-409. Portland: USDA Forest Service, Pacific Northwest Research Station. 8 p.
- Lewis SJ. 1985. Seedfall, germination, and early survival of red alder [MS thesis]. Seattle: University of Washington, College of Forest Resources.
- Liscinsky S. 1965. The American woodcock in Pennsylvania. Proj. W-50-R. Harrisburg: Pennsylvania Game Commission, Pittman Robertson (Federal Aid). 32 p.
- Ljunger A. 1959. Al-och alföraäling. Skogen 46(5): 115–117.
- Lowry GL, Brokow FC, Breeding CHJ. 1962. Alder for reforesting coal spoils in Ohio. Journal of Forestry 60: 196–199.
- Martin KJ, Tanaka Y, Myrold DD. 1991. Peat carrier increases inoculation success with *Frankia* on red alder (*Alnus rubra* Bong.) in fumigated nursery beds. New Forests 5: 43–50.
- McDermott RE. 1953. Light as a factor in the germination of some bottomland hardwood seeds. Journal of Forestry 51: 203–204.
- McGee A, Feller MC. 1993. Seed banks of forested and disturbed soils in southwestern British Columbia. Canadian Journal of Botany 71(12): 1574–1583.
- McNeill JD, Hollingsworth MK, Mason WL, Moffat AJ, Sheppard LJ, Wheeler CT. 1990. Inoculation of alder seedlings to improve seedling growth and field performance. Arbor. Res. Note 88. Washington, DC: USDoE, Arboricultural Advisory and Information Service. 3 p.
- McVean DN. 1955. Ecology of *Alnus glutinosa* (L.) Gaertn.: 3. Seedling establishment. Journal of Ecology 44(1): 195–218.
- McVean DN. 1959. Ecology of *Alnus glutinosa* (L.) Gaertn.: 4. Establishment of alder by direct seeding of shallow blanket bog. Journal of Ecology 47: 615–618.
- Miller RE, Murray MD. 1978. The effects of red alder on Douglas-fir growth. In: Proceedings, Utilization and Management of Alder Symposium. Gen. Tech. Rep. PNW-70. Portland, OR: USDA Forest Service, Pacific Northwest Research Station: 283–306.
- Mirov NT, Kraebel CG. 1939. Collecting and handling seeds of wild plants. For. Publ. 5. Washington, DC: USDA Forest Service, Civilian Conservation Corps. 42 p.
- Mix AJA. 1949. A monograph of the genus *Taphrina*. University of Kansas Science Bulletin 33 (1): 3–167.
- Molina R, Myrold D, Li CY. 1994. Root symbioses of red alder: technological opportunities for enhanced regeneration and soil improvement. In: Hibbs DE, DeBell DS, Tarrant RF, eds. The biology and management of red alder. Corvallis: Oregon State University: 23–46.
- Neal JL Jr, Trappe JM, Lu KC, Bollen WB. 1967. Sterilization of red alder seedcoats with hydrogen peroxide. Forest Science 13: 104–105.
- Niemic SS, Ahrens GR, Willits S, Hibbs DH. 1995. Hardwoods of the Pacific Northwest. Res. Contrib. 8. Corvallis: Oregon State University, Forestry Research Laboratory. 115 p.
- Peeler KC, DeBell DS. 1987. Variation in damage from growing season frosts among open pollinated families of red alder. Res. Note. PNW-464. Portland OR: USDA Forest Service, Pacific Northwest Research Station. 8 p.
- Perinet P, Brouillette JG, Fortin JA, LaLonde M. 1985. Large scale inoculation of actinorhizal plants with *Frankia*. Plant and Soil 87: 175–183.
- Plank ME, Willits S. 1994. Wood quality, product yield, and economic potential. In: Hibbs DE, DeBell DS, Tarrant RF, eds. The biology and management of red alder. Corvallis: Oregon State University: 243–247.

- Pojar J, MacKinnon A, comps. & eds. 1994. Plants of the Pacific Northwest Coast. Redmond, WA: Lone Pine Publishing: 44.
- Radwan MA. 1987. Effects of fertilization on growth and foliar nutrients of red alder seedlings. Res. Pap. PNW-375. Portland, OR: USDA Forest Service, Pacific Northwest Research Station. 14 p.
- Radwan MA, DeBell DS. 1981. Germination of red alder seed. Res. Note PNW-370. Portland, OR: USDA Forest Service, Pacific Northwest Forest and Range Experiment Station. 4 p.
- Radwan MA, DeBell DS. 1994. Fertilization and nutrition of red alder. In: Hibbs DE, DeBell DS, Tarrant RF, eds. The biology and management of red alder. Corvallis: Oregon State University: 216–228.
- Radwan MA, Tanaka Y, Dobkowski A, Fangen W. 1992. Production and assessment of red alder planting stock. Res. Pap. PNW-RP-450. Portland, OR: USDA Forest Service, Pacific Northwest Research Station.
- Sargent CS. 1965. Manual of the trees of North America (exclusive of Mexico). 2nd ed., corrected and reprinted. New York: Dover Publications. 934 p.
- Schalin I. 1967. Germination analysis of *Alnus incana* (L.) Moench and *Alnus glutinosa* (L.) Gaertn. seeds. Acta Oecologia Scandinavica 18: 253–260.
- Schopmeyer CS. 1974. *Alnus* B. Ehrh., alder. In: Schopmeyer CS, ed. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 206–211.
- Stettler RF. 1978. Biological aspects of red alder pertinent to potential breeding programs. In: Briggs DG, DeBell DS, Atkinson WA, eds. Utilization and management of alder. Gen. Tech. Rep. PNW-70. Portland, OR: USDA Forest Service, Pacific Northwest Forest and Range Experiment Station: 209–222.
- Tanaka Y, Brotherton PJ, Dobkowski A, Cameron PC. 1991. Germination of stratified and non-stratified seeds of red alder at two germination temperatures. New Forests 5: 67–75.
- Tarrant RF, Bormann BT, DeBell DS, Atkinson WA. 1983. Managing red alder in the Douglas-fir region: some possibilities. Journal of Forestry 81: 787–792.
- Tarrant RF, Hibbs DE, DeBell DS. 1994. Introduction: red alder and the Pacific Northwest. In: Hibbs DE, DeBell DS, Tarrant RF, eds. The biology and management of red alder. Corvallis: Oregon State University: ix–xi.
- Tarrant RF, Trappe JM. 1971. The role of *Alnus* in improving the forest environment. Plant and Soil (Spec. Vol. 1971): 335–348.
- Van Dersal WR. 1938. Native woody plants of the United States: their erosion-control and wildlife values. Misc. Pub. 303. Washington, DC: USDA. 362 p.
- White CM, West GC. 1977. The annual lipid cycle and feeding behavior of Alaskan redpolls. Oecologia 27: 227–238.
- White G. 1981. Controlled pollination of European black alder. In: Franklin EC, ed. Pollen management handbook. Agric. Handbk. 587. Washington, DC: USDA Forest Service: 89–91.
- Worthington NP. 1965. Red alder (*Alnus rubra* Bong.). In: Fowells HA, ed. Silvics of forest trees of the United States. Agric. Handbk. 271. Washington, DC: USDA Forest Service: 83–88.
- Zasada J, Sharik T, Nygren M. 1991. The reproductive process in boreal forest trees. In: Shugart HH, Leemans R, Bonan GB, eds. A systems analysis of the global boreal forest. Cambridge, UK: Cambridge University Press: 85–125.