

OPEN

Using bear rub data and spatial capture-recapture models to estimate trend in a brown bear population

Katherine C. Kendall^{1,2*}, Tabitha A. Graves³, J. Andrew Royle⁴, Amy C. Macleod⁵, Kevin S. McKelvey⁶, John Boulanger⁷ & John S. Waller⁸

Trends in population abundance can be challenging to quantify during range expansion and contraction, when there is spatial variation in trend, or the conservation area is large. We used genetic detection data from natural bear rubbing sites and spatial capture-recapture (SCR) modeling to estimate local density and population growth rates in a grizzly bear population in northwestern Montana, USA. We visited bear rubs to collect hair in 2004, 2009–2012 (3,579–4,802 rubs) and detected 249–355 individual bears each year. We estimated the finite annual population rate of change 2004–2012 was 1.043 (95% CI = 1.017–1.069). Population density shifted from being concentrated in the north in 2004 to a more even distribution across the ecosystem by 2012. Our genetic detection sampling approach coupled with SCR modeling allowed us to estimate spatially variable growth rates of an expanding grizzly bear population and provided insight into how those patterns developed. The ability of SCR to utilize unstructured data and produce spatially explicit maps that indicate where population change is occurring promises to facilitate the monitoring of difficult-to-study species across large spatial areas.

Understanding trends in population abundance is fundamental to wildlife management and can be critically important for conservation of at-risk taxa¹. Trend can be particularly challenging to quantify when population density is low, the population and its range are expanding or contracting, there is spatial variation in trend within the population, or the conservation area is large or remote; traits common to most forest carnivore populations. Grizzly bears (brown bear; *Ursus arctos*) epitomize the characteristics that make populations difficult to monitor. They occur at low densities², are solitary (except while females are accompanied by their young), have large home ranges³, often move long distances in short periods of time⁴, can exhibit strong behavioural responses to sampling that employs consumable baits and handling, and tend to only survive in remote areas⁵. Many populations are in decline throughout their Holarctic distribution but some have begun to recover and reoccupy former range in recent years^{6–10}. Estimating bear density can be difficult near such expansion fronts because large differences in densities can occur over short distances and may require fine-scale sampling over large areas¹¹.

Bears routinely rub vigorously on trees and other objects, often leaving hair behind. This ubiquitous activity is most likely a form of marking behaviour that functions as chemical signaling to other bears^{12,13}. By exploiting natural rubbing behaviour, it is possible to detect a substantial portion (36%⁸ to 85%¹³) of a bear population without the use of bait. Hair samples found at bear rub sites usually have sufficient DNA for genetic determination of species, sex and individual identity^{13–16}. Thus, periodic hair collection from rub objects produces a spatial array of repeated detections for individual bears appropriate for capture-recapture modeling¹⁷. When detection is imperfect, capture-recapture modeling provides an optimal approach for estimating population size, trend, and other demographic parameters.

¹U.S. Geological Survey, Northern Rocky Mountain Science Center, West Glacier, Montana, 59936, USA. ²Present address: Ursine Ecological, Columbia Falls, Montana, 59912, USA. ³U.S. Geological Survey, Northern Rocky Mountain Science Center, West Glacier, Montana, 59936, USA. ⁴U.S. Geological Survey, Patuxent Wildlife Research Center, Laurel, Maryland, 20708, USA. ⁵Applied Conservation Ecology, University of Alberta, Edmonton, Alberta, T6G 2H1, Canada. ⁶U.S. Forest Service, Rocky Mountain Research Station, Missoula, MT, 59801, USA. ⁷Integrated Ecological Research, Nelson, British Columbia, V1L 5T2, Canada. ⁸U.S. National Park Service, Glacier National Park, West Glacier, Montana, 59936, USA. *email: kkendall2382@gmail.com

The Northern Continental Divide Ecosystem (NCDE) in northwestern Montana, USA, is home to one of five North American grizzly bear populations persisting south of Canada. All were designated threatened in 1975⁵. Bear density in the NCDE in 2004 was highest in Glacier National Park (NP) in the north with density declining to the south and to the edges of occupied range^{8,18}. Along the southern boundary of Glacier NP, the presence of a busy highway (US 2) impacts bear crossings¹⁹. Gene flow is reduced across the corridor that contains the western portion of US 2 and the rail line that parallels it⁸. Population size in the NCDE was estimated to have increased from 765 in 2004⁸ to approximately 1,000 by 2009²⁰. Because this trend estimate was based on a small portion (2–4%) of the population, which increases the potential for bias, we sought to increase the proportion sampled and improve insight on population status by detecting individual bears from hair at natural bear rub sites. The purpose of our study was to explore this new approach to monitoring the trajectory of a low-density carnivore population at an ecosystem scale and to estimate variation in local density and growth rates within the population using a spatial capture-recapture modeling framework.

Methods

Field methods. We sampled the NCDE grizzly bear population in a 33,300-km² area dominated by the rugged and remote terrain of the Rocky Mountains (Fig. 1a). We collected hair at bear rubs sites identified by looking for the presence of hair on trees and other objects while searching along trails, roads, and utility and fence lines. Hair deposition was a result of natural behaviour; no attractant was used to draw bears to survey routes or to encourage rubbing. Rubs were fitted with 3 to 4 40-cm lengths of 4-point barbed wire to facilitate hair deposition⁸. We monitored the rubs we located in 2004 and 2009–2012 (see Supplementary Information online for additional details).

We quantified rub sampling effort as the cumulative number of days between the first collection visit (after the initial visit to clear hair that may have been deposited the previous year) and last collection of the year for each rub sampled per year (Table 1). Sampling began in early June in all years, ended in early September in 2004, and most sampling ended in early October 2009–2012.

Genetic analysis. Hair samples were analyzed by a laboratory that specialized in genotyping low quantity and quality DNA (Wildlife Genetics International; <http://www.wildlifegenetics.ca>). We used the G10J marker to distinguish between grizzly bear and black bear (*U. americanus*) samples²¹. We added 6 additional nuclear microsatellite loci to determine individual identity of the grizzly bear hair samples (G1A, G10B, G1D, G10H, G10M, and G10P)^{22–24} and the amelogenin marker to identify sex²⁵. We confirmed individuals identified using this 7-locus genotype by extending the genotypes to 9 additional loci (G10C, G10L, CXX110, CXX20, Mu50, Mu59, G10U, Mu23, and G10X^{22,23,24,26,27}), and confirmed sex with a second proprietary test developed by Wildlife Genetics International. Sub-selection of samples for analysis is described in Supplementary Information online.

Spatial capture-recapture modeling. We used maximum-likelihood spatial capture-recapture (SCR) models^{28,29} to estimate local and overall density and annual population growth rates from spatial detection histories. SCR models specify a spatially explicit link between a summary of each individual's location, henceforth their activity center, and locations where they may be detected, here, the sampled bear rubs. We treated the 5 years (2004, 2009, 2010, 2011, 2012) of detection data as independent “occasions” and ran separate models for males and females. For each year we denoted the coordinates of each rub by the vector x_{jt} for years $t = 2004$ and 2009–2012 and for $j = 1, 2, \dots, n_t$ rub locations and the activity center for individual i as s_i , a 2-dimensional coordinate. SCR models parameterize the probability of detection, $p(x_{jt}, s_i)$, of an individual at each rub by a function of distance between the rub and the putative activity center. For each year and sex, we used the half-normal model³⁰, where probability of detection at a location is assumed to decline with distance from the activity center following a half-normal distribution such that:

$$p(x_{jt}, s_i) = p_0 \exp(-\text{dist}(x_{jt}, s_i)^2 / (2\sigma^2)) \quad (1)$$

The parameters we estimated were baseline detection probability at the activity center, p_0 , and the scale of movement parameter, σ , related to the extent of space used by individuals during the period of sampling. In addition, SCR models contain a parameter representing density, D , estimated from the observed encounter history data. Model parameters were estimated by maximum likelihood (MLE³¹) using the R package oSCR^{31,32}. For all parameters of the model the standard error (SE) was from the Hessian of the likelihood function evaluated at the MLEs.

We developed our model set based on our *a priori* knowledge of grizzly bear ecology and theories about factors influencing bear behaviour. The hypotheses we considered modeled various effects on each of the 3 primary parameters; density (D), baseline detection probability (p_0), and the scale of movement parameter (σ , Table S1 online). Effects were modeled on each parameter on a link-transformed scale, with an intercept and effects modeled additively. We modeled temporal variation in D by allowing full year-specificity, $\log(D_t) = \beta_0$, or a linear trend $\log(D_t) = \beta_0 + \beta_1 \text{Year}$ where *Year* was the centered year (2008 = 0). Density and σ were modeled on the log-scale and baseline detection probability was modeled on the logit scale. We included duration of sampling interval, linear effects of date, a trap-specific behavioural response, and landscape covariate effects in all models for p_0 and evaluated hypotheses that included year-specific and quadratic effects of Julian date. We fit a behavioral model because rubbing functions as communication among individuals and thus prior marking at a rub may change the likelihood that site will be rubbed again by that or other individuals in the future. We calculated rub-specific landscape detection covariates to describe ruggedness, security level (protected land ownership), and the linear and quadratic effects of percent canopy cover within 250-m of a rub³³ (see Supplementary Information online for




Figure 1. (a) Northern Continental Divide Ecosystem study area in northwestern Montana, USA where growth rates of an expanding grizzly bear population were estimated during 2004–2012. State-space specified for the spatial capture-recapture models extended 45 km beyond the area where bear rubs were sampled. (b) Locations of naturally-occurring bear rub sites where the grizzly population was sampled by collecting hair for genetic determination of individuals 2004, 2009, 2010–2012. Maps were created from the USGS National Elevation Dataset⁵⁸ using ArcMap 10.2.

additional details). We added these covariates to test the hypotheses that detection would be higher; 1) in rugged areas because bears might be more likely to walk on trails, 2) where there was higher security, and 3) where trees were of moderate to high density. We included hypotheses that σ was constant or year-specific. We used the sample size-adjusted Akaike's Information Criterion (AIC_c) and AIC_c weights³⁴ to evaluate relative support for each candidate model. Models with cumulative weights up to 1.0 were considered supported.

To address human-bear conflicts, wildlife managers sometimes move bears large distances resulting in unnaturally large distances between detections³⁵. To assess the impact of this on our density estimates, we used the best model for each sex from the initial set of candidate models, but allowed σ and p_0 to vary by translocated individuals. These models would thus estimate the proportion of translocated individuals, defined as $\Psi_{trans} = Pr(\text{translocated} = 1)$ ³⁶.

To calculate trend, we used the most supported models. For models varying linearly with year, the trend parameter was estimated directly within the model. For models with year-specific density parameters, we characterize trend by the annual geometric mean rate of population change³⁷, λ , from 2004–2012 (8 annual intervals) according to:

$$\lambda = (D_{2012}/D_{2004}) \wedge (1/8) \quad (2)$$

Year	Rub sites	Sampling occasions	Cumulative rub sampling days ^a	Individual grizzly bears detected
2004	4,802	4	312,882	274
2009	3,579	2	289,822	249
2010	4,608	2	344,942	334
2011	4,711	2	342,157	321
2012	4,635	2	342,972	355

Table 1. Sampling effort and number of individual grizzly bears detected at natural bear rub sites monitored in the Northern Continental Divide Ecosystem in northwestern Montana, USA during 2004, 2009–2012. The number of rub sites monitored only includes those visited ≥ 1 time after the initial visit to remove hair that could have been deposited the previous year. ^aCumulative number of days between the first visit and last collection of the year for each rub sampled per year. For example, if we surveyed 1,000 rubs 1 time after surveying them 60 days earlier, and sampled 2,000 rubs 2 times at 45-day intervals and 100 rubs 3 times at 40-day intervals, rub sampling effort for that year would be $(1,000 \times 60) + (2,000 \times 45 \times 2) + (100 \times 40 \times 3) = 60,000 + 180,000 + 120,000 = 360,000$ rub sampling days. Sampling days began to accrue after the initial “clean-off” visit to each rub each year.

The MLE of λ was obtained by plugging in the MLEs of year-specific densities. The standard error (SE) of estimated rates of population change was computed using a delta approximation³⁸. The SEs were used to obtain 95% Wald type confidence interval. These growth rate estimates apply to the entire state space. We also compared our estimates to the trend in effective population size, N_e , a metric referring to the number of bears that are contributing reproductively to the population, under a set of genetic assumptions. For this we used analysis of covariance as described in Pierson *et al.*³⁹.

In SCR models, MLEs are based on the marginal likelihood of the encounter histories computed by averaging over all possible locations of the activity centers associated with detected individuals. This spatial region is called the “state space”. The state-space should be chosen to be sufficiently large such that an individual having an activity center on the periphery of the state-space has negligible probability of capture (⁴⁰ p. 132). For the half-normal detection model we used, we concluded that 3σ was sufficiently close to negligible, yielding a probability of encounter < 0.01 based on estimates from our model. In our initial model runs, male σ was 15 km so we defined our state space by a regular grid of points (4 km grid spacing) extending 45 km beyond any rub location (Fig. 1). The number of state-space points varied with locations of rubs monitored each year and ranged from 3,992 (2004) to 4,422 (2012). SCR models regard the activity center, s_i , as a latent variable that is estimated along with other parameters. We estimated the latent activity centers using the estimated best unbiased predictor⁴¹ which is the conditional mean of S_i given the data, evaluated at the MLEs. Predicted density surfaces were generated by aggregating the estimated posterior distribution of each S_i . To examine changes in local trend over time, we estimated the realized local population size (aka realized density) for each state-space point (⁴⁰ sec. 5.11), using the MLEs of the model parameters. The estimated realized population size for any state-space point $\mathbf{u}$ is defined as the estimated posterior mean of the number of activity centers associated with point $\mathbf{u}$. This is computed by adding (a) the sum of the posterior probability of each observed individuals’ activity center for $\mathbf{u}$ to (b) the expected number of individuals for that state-space point that went undetected. The former quantity, (a), is denoted by $Pr(s_i = \mathbf{u} | \mathbf{y}_i, \theta)$ (⁴⁰ ch. 6) where θ represents the collection of all model parameters for a given model, and we plug-in the MLEs for θ , $\hat{\theta}$ for this evaluation. Component (b) is denoted by $n_0(\mathbf{u} | \hat{\theta})$ which is also a function of all model parameters (⁴⁰ ch. 6). Thus, the expression for the estimated realized population size for any point $\mathbf{u}$ of the discrete state-space is given by:

$$\hat{D}(\mathbf{u}) = \left\{ \sum_{i=1}^n Pr(s_i = \mathbf{u} | \mathbf{y}_i, \hat{\theta}) \right\} + E(n_0(\mathbf{u} | \hat{\theta})) \quad (3)$$

This calculation can be done using the model parameters for each year to produce $\hat{D}_t(\mathbf{u})$, from which estimated local growth rates can be computed for each unit of the state space,

$$\lambda(\mathbf{u}) = \left(\frac{\hat{D}_{2012}(\mathbf{u})}{\hat{D}_{2004}(\mathbf{u})} \right)^{1/8}. \quad (4)$$

Local growth-rate estimates were limited to the area that was sampled in 2004 as this area was sampled in all years. We also calculated growth rates above and below Highway 2, a known barrier to gene flow⁸, by summing the estimated density of the state space points within each region in each year and calculating λ from those regional estimates. Variance estimates are not available for these sub-zones, however, because variance was calculated for the full state space and cannot be partitioned.

In one year, we detected one female bear 190 km to the south of other detections of this individual. Although grizzly bears are capable of traveling long distances, having a female range this far between detections, i.e. almost 3 times the distance of the second largest female movement detected (64 km), was highly unusual and constituted the lone extreme movement in our data set. While we did not identify any errors that could explain this large movement and it was not associated with a known translocation, in spatial modeling it is common to remove outlier locations when they are not representative of the population and leverage results⁴². We, therefore, fit all hypothesized models with and without the southern location for this individual, but consider primary results

Year	Individuals			Detections			
	Detected	Recaptured	Spatially recaptured	Total	Spatial	MMDM	Max Distance
Females							
2004	119	57	55	216	205	6.49	43
2009	94	31	29	138	136	7.16	64
2010	145	70	67	285	273	5.12	17
2011	143	55	54	257	248	5.27	30
2012	167	68	68	303	293	5.56	38
Males							
2004	155	97	95	668	626	18.43	113
2009	154	93	93	505	495	16.01	68
2010	189	103	101	554	540	17.90	143
2011	178	103	102	497	486	15.86	60
2012	188	111	111	532	523	16.68	104

Table 2. Number of female and male grizzly bears detected at bear rub sampling sites in northwest Montana 2004, 2009–2012. Individual recaptures are subsequent detections of an individual at the same or new locations. Spatial recaptures of an individual are detections in locations different from the initial and subsequent detections. The number of total detections is the total number of times bears were detected, e.g. in 2004, 119 bears were detected 216 times. Spatial detections count only detections in different locations, e.g. in 2004, 119 bears were detected in 205 unique locations. Mean maximum distance moved (MMDM) is mean distance between detections for each individual identified averaged for all bears. Max distance is the longest distance between 2 detection locations for any individual. Summary does not include 1 detection of a female in 2010 that was 190 km from other detections of that individual that year.

Model ^a	D	p0	σ	Δ AIC	Weight	CumWt
Females						
1	~linear	~year + dur + jul + jul2 + beh + spatial covs	~year	0.00	0.61	0.61
2	~linear	~year + dur + jul + beh + spatial covs	~year	1.26	0.32	0.93
3	~year	~year + dur + jul + jul2 + beh + spatial covs	~year	5.24	0.04	0.98
4	~year	~year + dur + jul + beh + spatial covs	~year	6.51	0.02	1.00
Males						
1	~linear	~year + dur + jul + jul2 + beh + spatial covs	~year	0.00	0.95	0.95
3	~year	~year + dur + jul + jul2 + beh + spatial covs	~year	5.80	0.05	1.00

Table 3. Description of most supported models (models with the lowest AIC scores with cumulative weights up to 1.0) developed to estimate grizzly bear population density using spatial capture-recapture data from the Northern Continental Divide Ecosystem in northwestern Montana, USA. Spatial detection histories were derived from genetic identification of individual bears from hair collected at natural bear rub sites. For females, a single outlier point in 2010 was removed from the dataset. See Table S1 online for descriptions of the complete set of candidate models. ^aModel notation: D: density (bears/1,000 km²). p0: baseline detection probability. σ : sigma; spatial scale parameter related to the amount of space used by each individual. Δ AIC: cumulative change in Akaike Information Criterion. Weight: measure of support for each model. CumWt: cumulative measure of support for the models. ~: “a function of”. linear: density is a linear function of time. year: parameter is year specific. dur: duration of sampling interval; number of days since previous sampling visit. jul: Julian date; linear effect of season on detection probability. jul2: Julian date squared; quadratic effect of season on detection probability. beh: behavioral response; a visit of some bear at some bear rub changes its subsequent probability of detection at that rub. spatial covs: spatial covariates: cover: percent of mean canopy cover within 250 m of the trap. cover2: percent of mean canopy cover squared. curv: standard deviation of terrain curvature within 250 m of the trap. sec: security level as determined by land ownership policies at the trap: 10 = U.S. National Park Service, 7 = U.S. Forest Service, 3 = state/other public, 1 = private).

to be those associated with models with the location removed. Results from complete models that include this location are presented in the online Supplementary Information.

Results

Field and genetic. We monitored 3,579–4,802 bear rubs annually in the NCDE in 2004, 2009–2012 (Table 1, Fig. 1b). The distribution of rubs we sampled varied between 2004, 2009 and 2010 but was nearly identical 2010–2012 (Fig. 1). See Supplementary Information online for an in-depth explanation of differences in sampling effort and extent. Despite expanding sampling east of the mountains after 2004, few bears were detected there suggesting the area was minimally occupied in 2004 (Figure S1 online). Rubs were visited on at least one occasion annually to collect hair; median = 4 visits/rub in 2004, 2 visits/rub 2009–2012, not including an initial visit to each rub

	Model	Female				Male	
		1	2	3	4	1	3
D (95% CI) Bears/1,000 km ²	Year						
	2004	7.77 (6.37–9.48)	7.76 (6.37–9.47)	7.92 (6.42–9.77)	7.90 (6.41–9.75)	5.78 (4.96–6.72)	5.75 (4.89–6.75)
	No sampling 20005–2008						
	2009	9.90 (9.00–10.90)	9.91 (9.00–10.90)	8.39 (6.32–11.15)	8.40 (6.33–11.16)	6.80 (6.34–7.30)	6.77 (5.74–7.97)
	2010	10.40 (9.34–11.57)	10.40 (9.35–11.57)	10.16 (8.39–12.29)	10.18 (8.41–12.32)	7.03 (6.54–7.56)	7.20 (6.20–8.34)
	2011	10.91 (9.60–12.40)	10.92 (9.61–12.41)	10.39 (8.51–12.68)	10.38 (8.50–12.67)	7.27 (6.69–7.88)	7.32 (6.27–8.53)
	2012	11.45 (9.80–13.38)	11.47 (9.81–13.40)	11.04 (9.21–13.22)	11.04 (9.21–13.23)	7.51 (6.81–8.27)	7.36 (6.33–8.54)
σ (95% CI) km	2004	4.40 (3.96–4.88)	4.39 (3.95–4.87)	4.39 (3.96–4.87)	4.38 (3.95–4.86)	9.30 (8.81–9.81)	9.30 (8.81–9.81)
	No sampling 2005–2008						
	2009	5.84 (4.99–6.83)	5.83 (4.98–6.82)	5.89 (5.01–6.92)	5.88 (5.00–6.90)	7.34 (6.92–7.78)	7.35 (6.93–7.79)
	2010	3.05 (2.79–3.33)	3.05 (2.79–3.33)	3.04 (2.78–3.33)	3.04 (2.78–3.33)	9.17 (8.65–9.72)	9.15 (8.63–9.71)
	2011	3.24 (2.94–3.57)	3.24 (2.94–3.58)	3.24 (2.94–3.58)	3.24 (2.94–3.58)	7.62 (7.16–8.10)	7.62 (7.16–8.10)
	2012	3.86 (3.53–4.22)	3.86 (3.53–4.22)	3.86 (3.53–4.22)	3.86 (3.53–4.22)	9.12 (8.59–9.68)	9.13 (8.59–9.69)

Table 4. Estimates (transformed) of density, D, and spatial scale of movement, σ , parameters for models developed to estimate grizzly bear population trend using spatial capture-recapture data from the Northern Continental Divide Ecosystem in northwestern Montana, USA.

to clear hair that could have been deposited the previous year (Table 1). We detected 249–355 individual grizzlies each year for a total of 394 different females and 398 males during our 5 sampling years (Table 2). Our laboratory protocols made it highly unlikely that genotyping error resulted in the creation of any false detections in our data⁸.

Spatial modeling. Our models were remarkably sensitive to the single extreme movement between detections in our data. Removal of this point in 2010 changed the most supported female models from those with year-specific density to models with a linear trend in density (compare Table S3 online, Table 3). However, estimated female population growth rates were similar: λ_{04-12} was 1.042 (CI: 1.023–1.060) versus 1.050 (95% CI: 1.010–1.089) with and without the outlier, respectively. Estimates differed primarily in 2010. Namely, σ was much larger and baseline detection and density were much smaller in 2010 than other years (Table S4 online). For the top model, estimates of density were significantly lower and estimates of σ were significantly higher than estimates without the outlier location (non-overlapping CIs: Table S4, S5 online). When the outlier point was excluded, density estimates in 2010 were more consistent with estimates for the other years.

Because this one long-range movement was not representative of the population and had such a large effect, we focus here on model results that did not include this data point. The most supported model by far for females and males defined density as increasing linearly with year, year-specific σ , and included all detection covariates (Table 3; see Table S1 online to view all candidate models). Detection increased with duration, lower canopy cover, and a higher percent of secure area, and had a quadratic function for Julian date (Table S2). Models with a translocation effect did not converge, likely due to low sample size of translocated bears within the DNA data set (12 females, 9 males).

Annual estimates of density were similar among all supported models (Table 4). Both the size and distribution of the grizzly bear population in the NCDE expanded between 2004 and 2012. Ecosystem-wide, female density increased from 7.77 bears/1,000 km² (95% CI: 6.37–9.48) in 2004 to 11.45 bears/1,000 km² (95% CI: 9.80–13.38) in 2012. Male density rose from 5.78 bears/1,000 km² (95% CI: 4.96–6.72) in 2004 to 7.51 bears/1,000 km² (6.81–8.27) in 2012. In 2004, the majority of the population resided in the northern third of the ecosystem centered on Glacier NP. As the population grew, density increased throughout the ecosystem with the most obvious changes in the southern two thirds of the NCDE where bears were absent or density was low in 2004 (Fig. 2).

The annual rate of overall population change 2004–2012 (λ_{04-12}) was 1.043 (95% CI: 1.017–1.069). The growth rates of the female and male segments of the population were not significantly different; $\lambda_{\text{female}04-12} = 1.050$ (95% CI: 1.010–1.089), $\lambda_{\text{male}04-12} = 1.033$ (95% CI: 1.007–1.060) (Table 5). Both female and male growth rates were higher south of US 2 (F: 1.051, M: 1.053) than north (F: 1.030, M: 1.010). Most of the ecosystem exhibited stable to increasing local population trends, however, some areas experienced declines for one sex or the other (Fig. 3).

Discussion

Using genetic detection data from natural bear rub sites in an SCR modeling framework, we estimated ecosystem-scale as well as local trends in a recovering grizzly bear population. Grizzly bear numbers in northwestern Montana were extremely low in the early 1900s⁴³. Park protection and high-quality habitat resulted in recovery occurring earlier in Glacier NP than in the rest of the ecosystem⁸. The annual density surfaces produced by our model graphically depict the growth of the population outside of Glacier NP and how bear distribution changed 2004–2012. As the population expanded from the core in-and-adjacent-to Glacier NP, the population filled in much of the southern two thirds of the NCDE where few or no bears were found at the start of our study (Fig. 2). While the population as a whole grew at an annual rate of 1.043 in 2004–2012, growth rates were highest south of US 2 where initial density was low (Fig. 3). Population density remained fairly stable in Glacier NP and on the Blackfeet Reservation east of the Park 2004–2012 where 2004 densities were highest.




Figure 2. Annual predicted density surfaces of the grizzly bear population in the Northern Continental Divide Ecosystem 2004, 2009–2012. Density estimates (bears/16 km²) were based on the most supported spatial capture-recapture model of detection histories at natural bear rub sites. We limited estimates to the area in which rubs were identified and sampled, which varied by year.

Model	Female				Male	
	1	2	3	4	1	3
λ	1.050	1.050	1.042	1.043	1.033	1.031
SE	0.020	0.020	0.018	0.018	0.013	0.014
LCI	1.010	1.010	1.006	1.006	1.007	1.007
UCI	1.089	1.090	1.079	1.079	1.060	1.060

Table 5. Population growth rate estimates, λ , 2004–2012 for the most supported spatial capture-recapture models based on grizzly bear rub data from northwest Montana, USA. Detections used in these estimates excluded one outlier detection of a female in 2010. These estimates use the delta method to incorporate standard error, SE. Lower and upper confidence intervals, LCI and UCI, are 95%.

Because the use of SCR to model population trend surfaces from genetic detection data is new, we confirmed that independent estimates and ancillary data were consistent with patterns we found in population density and growth rates. Our annual rate of change for females 2004–2009 was similar to a previously published estimate for the NCDE ($\lambda = 1.031$ (95% CI: 0.928–1.102)) made using female vital rates derived from telemetry data and a known-fate estimator²⁰. Our growth rate was also similar to other recovering brown bear populations (see Supplementary Information online for details). As the population grew, N_e in the NCDE also increased between 2004 and 2012 and generally tracked population trend³⁹. The number of detections as well as the number of immigrants (based on a parentage assessment) consistently increased between 2004 and 2012 in the areas with the highest growth rates and lowest initial density, i.e. central, southern, and eastern regions⁴⁴. These same areas, which exhibited lower density and diversity in 2004 (Fig. 2) also increased in genetic diversity^{8,44}. This is consistent with previous evidence that bears south of US 2 were split into several small semi-isolated groups^{8,44}. We also would expect to see bears moving into habitat that was not at carrying capacity as the population grew.

Our SCR models were highly sensitive to a spatial outlier. Even though our dataset was larger than most where SCR has been applied, we found that inclusion of a single extended movement changed the top models, evaluated




Figure 3. Annual local rates of change in grizzly bear population density within the Northern Continental Divide Ecosystem in northwestern Montana 2004–2012. Population trend was calculated as the annual geometric rate of change in grizzly bear density (D) between 2 years. Rate of population change was calculated for each 16 km² state space pixel defined by our spatial capture-recapture model. The estimate area was smaller for 2004 than in subsequent years because few bear rubs were found along the eastern edge of the study area that year. Local density should only be inferred for the area sampled. Growth rates > 1.0 denote areas of population growth; rates < 1.0 denote areas of population decline.

through AIC, versus when the outlier was removed (Table S3 online, Table 3). We fit models without this point because such a large movement within one year is extremely uncommon for a female grizzly bear and resulted in unrealistically large home range and σ estimates. Removal of outliers from spatial data where the goal is fitting a declining function (e. g. half normal) is common. For example, in distance sampling⁴² it is common to remove outliers to allow fitting a simpler model⁴⁵. This often leads to a fairly heavy right truncation (e. g. 5%), and in some cases left truncation, that is normative in distance modeling^{46–48}. The treatment of space in distance modeling conceptually matches its treatment in SCR modeling, and some level of right truncation may be appropriate. In our case, while the decision to remove the outlier affected the best models and some detection and density estimates, our primary conclusions—that the population was growing and that density was increasing fastest in the southern portion of the ecosystem—were robust. However, we have no reason to believe that this is a generalizable conclusion. Our results were consistent with Stenhouse *et al.*³⁵ in finding SCR estimates to be sensitive to an uncommon extreme movement. While SCR should be robust to common large movements³⁶, we suggest comparing results with and without extreme movements to assess the impacts on all parameters.

We have seen no discussion of this sensitivity in the SCR literature and, in distance sampling, the arguments presented to support right truncation are primarily based on practicality rather than theoretical optimization⁴⁵. Lacking any guidance in the literature we have presented results from analyses with and without the outlier, but have concentrated on the truncated dataset because it is likely more representative of the population. We believe that this issue requires additional study and that the field probably should develop rules for right truncation similar to those adopted in the distance sampling literature. Incorporation of telemetry might be used to reduce the effect of outliers, but this requires further study.

We hypothesize that non-spatial sources of heterogeneity of detection at rubs could bias density estimates. Females may limit rubbing in non-breeding years to avoid males and thereby protect dependent young. However, the proportion of females in our sample was very similar in all years except 2009 when sampling effort and baseline female capture probability were lowest. Our models accounted for heterogeneity in individual detection in space, because we believe that to be the largest source of heterogeneity. However, we could not model other potential sources of individual heterogeneity. For instance, we could not assign a history of human-bear conflict or breeding status to individuals, both of which could cause avoidance of the anthropogenic routes we sampled along and result in depressed detection rates. Nonetheless, as long as the proportion of the detectable population

sampled is constant among years this approach provides a valid estimate of the trend for the population. We expect that the biases should be similar across years, which means that our trend estimates are reliable even if there is bias in density estimates. See Supplementary Information online for discussion of sampling efficiency at rub sites.

One of the most appealing aspect of using rub trees and SCR to detect trends is that both the data and the analyses are spatially explicit. Other analyses (e. g. known-fate^{49,50}) do not consider the location of captures and recaptures, and thus require evenly and consistently distributed data collection⁵¹. Rub trees provide a huge spatial sample of individual bears at minimal cost. This allows patterns such as the infilling and range expansion we observed in this study to be directly observed. Combined with SCR analyses, the resulting density maps allow the detailed evaluation of where and when the population is increasing or declining which, in turn, permits direct checks of the data. Boulanger *et al.*⁵² similarly used SCR and density surface modelling to test whether bear population density was related to habitat value, mortality risk, or a combination of factors across the range of occupied grizzly bear habitat in Alberta, Canada. A population that is expanding based on population-wide changes in density may expand its range and increase its density in areas previously minimally occupied. We were able to demonstrate these relationships, providing increased confidence that the density shifts were appropriately related to actual population dynamics. Further, areas of population growth or decline can be linked to spatial covariates such as habitat quality or human population densities to evaluate the proximal causes of these population changes.

Fine scale information on changes in density has great utility for adaptive management, prioritizing management to preserve animal migrations³¹, and evaluating interspecific competition⁵³ and resource selection²⁹. Here we have used multiple density surfaces to assess infilling of a grizzly population in areas with previously low density. This was possible because our sample sites were distributed at a fine scale (spacings $< 3 \sigma$ in all years) across our area of inference (Fig. 1b). We are confident of change at this coarse resolution, but suggest caution in over-interpreting change at the high resolution (16 km²) scale, because no estimates of variance are currently possible at that scale and little research has evaluated the appropriate resolution for estimating changes in density for animals with large versus small home ranges.

Changes in population status can occur fairly quickly in a recovering population even for species with low reproductive rates such as grizzly bears⁴⁴. Informed management of forest carnivore populations requires insight into temporal and spatial variation in density and growth rates, particularly when large local declines can rapidly occur⁵⁴. Sampling populations, such as ours, that occupy vast, remote areas in heterogeneous landscapes and monitoring during range expansion is especially challenging. Our genetic detection sampling approach coupled with SCR modeling allowed us to estimate differential growth rates of an expanding grizzly bear population and provided insight into how those patterns developed. For populations experiencing range expansion and thus violating the assumption of geographic closure, the SCR methods we employed are ideal because unlike non-spatial models, they are able to estimate the area to which the population estimates apply. While our dataset was so large that it restricted the set of methods available to us due to computational requirements, recent developments in open SCR models will additionally provide information on recruitment and survival^{55,56}. Knowledge of spatial variation in trend can be used to design monitoring and management strategies tailored to area-specific needs. Spatial patterns revealed by our approach may identify source and sink habitats or ecological traps that merit enhanced management focus. Finally, the value of obtaining ancillary information on metrics such as gene flow, genetic diversity, population structure, the number of breeders, and fitness from monitoring population trend makes a compelling case for including a genetic sampling approach in monitoring programs.

Data availability

The genetic profiles have been deposited in ScienceBase and are publicly available⁵⁷. Because grizzly bears are a US Government Endangered Species Act-listed species and spatial detection data could be used to identify grizzly bear concentration areas, access to precise locations may be limited. Field data and computer code are available from US Geological Survey, Northern Rocky Mountain Science Center, West Glacier, Montana 59936 USA.

Received: 12 May 2019; Accepted: 23 October 2019;

Published online: 14 November 2019

References

1. Taylor, M. F., Suckling, K. F. & Rachlinski, J. J. The effectiveness of the Endangered Species Act: a quantitative analysis. *BioScience* **55**, 360–367 (2005).
2. Miller, S. D. *et al.* Brown and black bear density estimation in Alaska using radiotelemetry and replicated mark-resight techniques. *Wildlife Monographs*, 133 (1997).
3. Schwartz, C. C., Miller, S. D. & Haroldson, M. A. Grizzly bear. Second edition. [Feldhamer, G. A., Thompson, B. C. & Chapman, J. A. (eds.)] *Wild Mammals of North America: Biology, Management, And Conservation*. 556–586 (Johns Hopkins University Press, 2003).
4. Proctor, M. *et al.* Population fragmentation and inter-ecosystem movements of grizzly bears in western Canada and the northern USA. *Wildlife Monographs*, 180 (2012).
5. U.S. Fish and Wildlife Service. Grizzly Bear Recovery Plan. U.S. Fish and Wildlife Service, Missoula, Montana, USA (1993).
6. Schwartz, C. C. *et al.* Temporal, spatial, and environmental influences on the demographics of grizzly bears in the greater Yellowstone ecosystem. *Wildlife Monographs*, 161 (2006).
7. Nawaz, M. A., Swenson, J. E. & Zakaria, V. Pragmatic management increases a flagship species, the Himalayan brown bears, in Pakistan's Deosai National Park. *Biological Conservation* **141**, 2,230–2,241 (2008).
8. Kendall, K. C. *et al.* Demography and genetic structure of a recovering grizzly bear population. *J. Wildlife Management* **73**, 3–17 (2009).
9. Kindberg, J. *et al.* Estimating population size and trends of the Swedish brown bear *Ursus arctos* population. *Wildlife Biology* **17**, 114–123 (2011).

10. Karamanlidis, A. A., Hernando, M., de, G., Krambokoukis, L. & Gimenez, O. Evidence of a large carnivore population recovery: Counting bears in Greece. *J. Nature Conservation* **27**, 10–17 (2015).
11. Swenson, J. E., Sandegren, F. & Soderberg, A. Geographic expansion of an increasing brown bear population: Evidence for presaturation dispersal. *J. Animal Ecology* **67**, 819–826 (1998).
12. Clapham, M., Nevin, O. T., Ramsey, A. D. & Rosell, F. The function of strategic tree selectivity in chemical signalling of brown bears. *Animal Behaviour* **85**, 1351–1357 (2013).
13. Sawaya, M. A., Stetz, J. B., Clevenger, A. P., Gibeau, M. L. & Kalinowski, S. T. Estimating grizzly and black bear population abundance and trend in Banff National Park using noninvasive genetic sampling. *PLoS ONE* **7**, 1–12, <https://doi.org/10.1371/journal.pone.0034777> (2012).
14. Kendall, K. C. *et al.* Grizzly bear density in Glacier National Park. *Montana. J. Wildlife Management* **72**, 1693–1705 (2008).
15. Karamanlidis, A. A. *et al.* Noninvasive genetic studies of brown bears using power poles. *European J. Wildlife Research* **56**, 693–702 (2010).
16. Stetz, J. B., Kendall, K. C. & Macleod, A. C. Black bear density in Glacier National Park, Montana. *Wildlife Society Bulletin* **38**, 60–70 (2014).
17. Stetz, J. B., Kendall, K. C. & Servheen, C. Evaluation of bear rub surveys to monitor grizzly bear population trends. *J. Wildlife Management* **74**, 860–870, <https://doi.org/10.2193/2008-435> (2010).
18. Graves T. A. *et al.* Balancing precision and risk: Should multiple detection methods be analyzed separately in N-mixture models? *PLoS ONE* **7**, <https://doi.org/10.1371/journal.pone.0049410> (2012).
19. Waller, J. S. & Servheen, C. Effects of transportation infrastructure on grizzly bears in northwestern. *Montana. J. Wildlife Management* **69**, 985–1000 (2005).
20. Mace, R. D. *et al.* Grizzly bear population vital rates and trend in the Northern Continental Divide Ecosystem. *Montana. J. Wildlife Management* **76**, 119–128 (2012).
21. Mowat, G. *et al.* Grizzly *Ursus arctos* and black bear *U. americanus* densities in the interior mountains of North America. *Wildlife Biology* **11**, 31–48 (2005).
22. Paetkau, D. & Strobeck, C. Microsatellite analysis of genetic variation in black bear populations. *Molecular Ecology* **3**, 489–495 (1994).
23. Paetkau, D., Calvert, W., Stirling, I. & Strobeck, C. Microsatellite analysis of population structure in Canadian polar bears. *Molecular Ecology* **4**, 347–354 (1995).
24. Paetkau, D., C. Strobeck, C. & Shields, G. F. Gene flow between insular, coastal and interior populations of brown bears in Alaska. *Molecular Ecology* **7**, 1,283–1,292 (1998).
25. Ennis, S. & Gallagher, T. F. PCR based sex determination assay in cattle based on the bovine Amelogenin locus. *Animal Genetics* **25**, 425–427 (1994).
26. Taberlet, P. *et al.* Noninvasive genetic tracking of the endangered Pyrenean brown bear population. *Molecular Ecology* **6**, 869–876 (1997).
27. Ostrander, E. A., Sprague, G. F. Jr. & Rine, J. Identification and characterization of dinucleotide repeat (CA)_n markers for genetic mapping in dog. *Genomics* **16**, 207–213 (1993).
28. Efford, M. Density estimation in live-trapping studies. *Oikos* **106**, 598–610 (2004).
29. Royle, J. A., Chandler, R. B., Sun, C. C. & Fuller, A. K. Integrating resource selection information with spatial capture–recapture. *Methods in Ecology and Evolution* **4**, 520–530, <https://doi.org/10.1111/2041-210X.12039> (2013).
30. Borchers, D. L. & Efford, M. G. Spatially explicit maximum likelihood methods for capture–recapture studies. *Biometrics* **64**, 377–385 (2008).
31. Fuller, A. K., Sutherland, C. S., Royle, J. A. & Hare, M. P. Estimating population density and connectivity of American mink using spatial capture–recapture. *Ecological Applications* **26**, 1125–1135 (2016).
32. Sutherland, C., A. Royle, A. & Linden, D. oSCR (open-source SCR), an R package for likelihood analysis of SCR models. Version 0.30.2. URL <https://sites.google.com/site/spatialcapturecapture/osc-r-package> (2016).
33. Ironside, K. E. *et al.* Geomorphometry in Landscape Ecology: Issues of scale, physiography, and application. *Environment and Ecology Research* **6**(5), 397–412, <https://doi.org/10.13189/eeer.2018.060501> (2018).
34. Burnham, K. P. & Anderson, D. R. Model selection and multimodel inference: a practical information-theoretic approach. (Springer-Verlag, 2002).
35. Stenhouse, G. B. *et al.* Estimates of grizzly bear population size and density for the 2014 Alberta Yellowhead Population Unit (BMA 3) and south Jasper National Park Inventory Project. Report prepared for Weyerhaeuser Ltd., West Fraser Mills Ltd, Alberta Environment and Parks, and Jasper National Park. <https://frirresearch.ca/resource/estimates-grizzly-bear-population-size-and-density-final-report> (2015).
36. Royle, J. A., Sutherland, C., Fuller, A. K. & Sun, C. C. Likelihood analysis of spatial capture–recapture models for stratified or class structured populations. *Ecosphere* **6**, 1–11 (2015).
37. Link, W. A. & Sauer, J. R. Estimating population change from count data: Application to the North American Breeding Bird Survey. *Ecological Applications* **8**, 258–268 (1998).
38. Parr, W. C. A note on the jackknife, the bootstrap and the delta method estimators of bias and variance. *Biometrika* **70**, 719–722 (1983).
39. Pierson, J. C., Graves, T. A., Banks, S. C., Kendall, K. C. & Lindenmayer, D. B. Relationship between effective and demographic population size in continuously distributed populations. *Evolutionary Applications* **11**, 1162–1175, <https://doi.org/10.1111/eva.12636> (2018).
40. Royle, J. A., Chandler, R. B., Sollmann, R. B. & Gardner, B. Spatial Capture-Recapture. (Elsevier Academic Press, 2013).
41. Robinson, G. That BLUP is a good thing: The estimation of random effects. *Statistical Science* **6**, 15–32; <http://www.jstor.org/stable/2245695> (1991).
42. Thomas, L. *et al.* Distance software: design and analysis of distance sampling surveys for estimating population size. *J. Applied Ecology* **47**, 5–14 (2010).
43. Keating, K. A. Historical grizzly bear trends in Glacier National Park, Montana. *Wildlife Society Bulletin* **14**, 83–87 (1986).
44. Mickle, N., Graves, T. A., Kovach, R., Kendall, K. C. & Macleod, A. C. Demographic mechanisms underpinning genetic assimilation of remnant groups of a large carnivore. *Proc. Royal Society B* **282**; <https://doi.org/10.1098/rspb.2016.1467> PMID: 27655768 (2016).
45. Buckland, S.T. *et al.* Introduction to distance sampling: estimating abundance of biological populations. 103–108, 151–153 (Oxford University Press, 2001).
46. Giunchi, D., Gaggini, V. & Baldaccini, N. E. Distance sampling as an effective method for monitoring feral pigeon (*Columba livia* f. domestica) urban populations. *Urban Ecosystems* **10**, 397–412 (2007).
47. Liu, Z., Wang, X., Teng, L., Cui, D. & Li, X. Estimating seasonal density of blue sheep (*Pseudois nayaur*) in the Helan Mountain region using distance sampling methods. *Ecological Research* **23**, 393–400 (2008).
48. Ward, A. I., White, P. C. & Critchley, C. H. Roe deer *Capreolus capreolus* behaviour affects density estimates from distance sampling surveys. *Mammal Review* **34**, 315–319 (2004).
49. Pradel, R. Utilization of mark–recapture for the study of recruitment and population growth rate. *Biometrics* **52**, 703–709 (1996).
50. Eberhardt, L. L. A paradigm for population analysis of long-lived vertebrates. *Ecology* **83**(2), 841–854 (2002).

51. Boulanger, J., Himmer, S. & Swan, C. Monitoring of grizzly bear population trend and demography using DNA mark-recapture methods in the Owikeno Lake area of British Columbia. *Canadian Journal of Zoology* **82**, 1,267–1,277 (2004).
52. Boulanger, J., Nielsen, S. E. & Stenhouse, G. B. Using spatial mark-recapture for conservation monitoring of grizzly bear populations in Alberta. *Scientific Reports* **8**, 5204, <https://doi.org/10.1038/s41598-018-23502-3> (2018).
53. Stetz, J. B., Mitchell, M. S. & Kendall, K. C. Using spatially-explicit capture-recapture models to explain variation in seasonal density patterns of sympatric ursids. *Ecography* **42**, 1–12 (2018).
54. Laufenberg, J. S., Johnson, H. E., Doherty, P. F. Jr. & Breck, S. W. Compounding effects of human development and a natural food shortage on a large carnivore population along a human development-wildland interface. *Biological Conservation* **224**, 188–198 (2018).
55. Whittington, J. & Sawaya, M. A. A comparison of grizzly bear demographic parameters estimated from non-spatial and spatial open population capture-recapture models. *PLoS ONE*, **10**; <https://doi.org/10.1371/journal.pone.0134446> (2015).
56. Gardner, B., Sollmann, R., Kumar, N. S., Jathanna, D. & Karanth, K. U. State space and movement specification in open population spatial capture-recapture models. *Ecology and Evolution* **8**(10), 336–10,344 (2018).
57. Graves, T. A., Mickle, N. L., Kendall, K. C., and Macleod, A. C. 2016, Northern Continental Divide Ecosystem grizzly bear (*Ursus arctos*) genetic profiles (1998–2012): U.S. Geological Survey data release, <https://doi.org/10.5066/F7Q81B63> (2016).
58. U.S. Geological Survey. USGS National Elevation Dataset (NED) 1 arc-second Downloadable Data Collection from The National Map 3D Elevation Program (3DEP) - National Geospatial Data Asset (NGDA) National Elevation Data Set (NED): U.S. Geological Survey, <https://www.sciencebase.gov/catalog/item/5825a0c3e4b01fad86db66dc> (2016).

Acknowledgements

Teams of 35–240 field personnel were responsible annually for collecting the thousands of bear hair samples and associated data used in this study. Laboratory analyses were conducted by the staff of Wildlife Genetics International led by D. Paetkau; J. Benson, L. Brady, C. Harris, D. Hausleitner, J. Midigan, R. Privee, S. Snively, P. Sylvestre, J. Wastl, and S. Waterhouse. C. S. Gaulke created the final version of the maps depicting results. C. Barbouletos provided the leadership that launched this project. C. Weber sustained that management support. Major funding for this work was provided by the U.S. Forest Service and U.S. Geological Survey. Additional financial support was contributed by the University of Montana, the Glacier National Park Conservancy, and Swan Valley Connections of Condon, Montana. Research conducted to acquire the data for this study was conducted in conformance with all endangered species laws. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Author contributions

K.K. conceptualized, designed, and obtained the majority of funding for the original mark-recapture projects and wrote the majority of this paper. K.K. and A.M. were instrumental in collecting the DNA data sets along with the rest of K.K.'s research teams. A.M. was responsible for data curation and summaries. T.G. produced the landscape covariates used in detection surface modelling. T.G. and J.A.R. designed and conducted all analyses with input from the other authors and wrote the majority of the SCR analysis methods, modeling results, and modeling discussion sections. A.M. and T.G. created the maps. All authors provided input on data interpretation and contributed to editing the paper.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-019-52783-5>.

Correspondence and requests for materials should be addressed to K.C.K.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019

Using bear rub data and spatial capture-recapture models to estimate trend in a brown bear population

Katherine C. Kendall, Tabitha A. Graves, J. Andrew Royle, Amy C. Macleod, Kevin S. McKelvey, John Boulanger, and John S. Waller

SUPPLEMENTARY INFORMATION

Introduction

Although bear populations are challenging to monitor, bears possess an attribute that facilitates sampling them. Bears routinely rub on trees and other objects vigorously, often leaving hair behind. The most likely explanation for this ubiquitous activity is that it is a form of marking behaviour and functions as chemical signaling to other bears¹. This tendency to rub results in the deposition of hair from individual bears at multiple sites. Because bears often use trails to travel, rub objects along trail networks can be located visually and therefore represent entirely passive detection; the sampling process has no effect on bear behaviour. While bears with a previous history of capture and handling have lower detection rates at baited hair corrals than bears that have not been trapped², no such effect has been found at bear rubs³.

Grizzly bear and black bear (*U. americanus*) marking behaviour is well documented, if imperfectly understood^{4,5,6,7}. Bears of all sex and age classes rub throughout the active season with variable frequency⁸. Evidence of rubbing is found throughout occupied habitat off-trail as well as along maintained and wildlife trails and by roads. Bears do not appear to discriminate much in selecting objects to mark. Bear hair has been found on a wide variety of natural (e.g. trees of all sizes and taxa, stumps, rocks) and man-made objects (e.g. sign and fence posts, utility poles, bridges, cabins)^{8,9,10}. Female and subadult grizzlies rarely rub during the breeding season (May–June), however, their detection probability increases after low rates in spring and approaches male rates by late summer^{3,8,11,12}.

Many studies have demonstrated that genetic detection through hair collection is a highly effective way to sample bear populations; hair collection from a grid of baited hair corrals has been used extensively to study bear population abundance throughout the world (e.g. ^{13,14,15,16}). Kendall *et al.*^{3,8} added hair collection at natural bear rub sites as a secondary sampling approach to a primary method using an array of baited hair corrals. Because there was only partial overlap in individuals detected at hair corrals and rubs, the addition of this second sampling approach increased sample coverage and population abundance estimate precision while abundance estimates themselves were not significantly changed¹⁷. Results of subsequent studies that used concurrent sampling at hair corrals and bear rubs also found that the addition of rub site sampling was a cost-effective way to increase sample size and estimate precision^{11,15,18}. While detection rates per sample site are higher at baited hair corrals than at unbaited rub sites^{3,8,17,19,20} it is possible to sample a similar or larger portion of the population at rub sites than at baited corrals^{11,18}. Sampling at rubs can also be more cost effective than at

baited hair corrals. In most studies using both methods, more field crew time was put into sampling at baited corrals than at rubs even when rubs greatly outnumbered corrals. For example, the ratio of field crews assigned to hair corral vs. rub sampling was over 4:1, while the ratio of individuals detected at corrals and rubs ranged from 1:1 to 2.5:1 for the studies conducted by Kendall *et al.*^{3,8,18}. However, the overall efficiency of rub tree sampling depends on monitoring higher numbers of rub trees to offset the lower per site sampling efficiency of rub trees^{19,20}. Therefore, rub tree sampling is most viable in areas with greater access, such as maintained trails.

Methods

Study area

Approximately 85% of the Northern Continental Divide Ecosystem (NCDE) is in public ownership, where grizzly bear conservation is a management objective. It includes Glacier National Park (NP), parts of 4 National Forests (Flathead, Kootenai, Helena-Lewis and Clark, and Lolo), five Wilderness areas (Bob Marshall, Great Bear, Mission Mountain, Rattlesnake, and Scapegoat), three Wilderness study areas (Mount Hefty-Tuchuck, Ten Lakes, and Thompson-Seton), Bureau of Land Management, U.S. Fish and Wildlife Service, and State land, and approximately half of the Blackfeet and one-third of the Flathead Indian Reservations. Private lands comprising the remaining 15% of the NCDE occur chiefly along the eastern, southern and western peripheries, but the Clearwater and Swan River Drainages, which are interior to the NCDE, contain both extensive diffuse housing and several small towns. Waterton Lakes National Park, Alberta and Akamina-Kishinena Provincial Park and provincial Crown lands in British Columbia, Canada border the NCDE to the north.

The climate and vegetation of the NCDE are shaped by the uplifted fault-blocks which formed the Rocky Mountains and by Pleistocene era ice ages²¹. Higher elevations receive more precipitation than lower slopes and valleys with annual precipitation ranging from over 170 cm in the higher elevations on the west side of Glacier NP to less than 38 cm on the eastern border. The Pacific maritime-influenced climate west of the Continental Divide is moister and more temperate than the colder and drier continental climate on the eastern side. The rugged terrain creates an array of habitats from alpine tundra and subalpine forests to moist mountain valleys, foothill groves, and semiarid grasslands in the rain shadow of large mountain masses²². Vegetation is characterized by coniferous forest, shrub fields and alpine tundra in the mountains, mixed deciduous–coniferous woodlands and herbaceous meadows in the valleys, and aspen (*Populus tremuloides*) stands, prairie grasslands, and agricultural fields along the eastern boundary³.

Field methods

To identify bear rubs, we closely examined objects that included one or more of the following characteristics; smoothed and/or discoloured bark and branch stobs, bare ground at the base of trees/poles, entrenched bear trails leading to rubs from maintained trails, and bear bite or claw marks. We attempted to get broad and even geographic distribution of areas searched for rubs and did not prioritize areas thought to have higher densities of bears. In lower elevation areas

where there were fewer trees for bears to rub on, utility poles and fence lines provided sampling opportunities. To prevent damage to pack stock and their cargo, we used double-stranded smooth wire on trees that appeared to have been bumped by horse packs. We separated the twisted ends of smooth wire slightly to make them more effective in snagging hair. We visited rubs at least twice annually. All hair deposited on the wire was collected during each visit. The hair found on rubs during the initial visit each year was not analyzed and did not contribute to detection histories because it could have been deposited the previous year. A flame was passed under each barb and wire after sample collection, and hair between wires was removed to prevent contamination when the next bear rubbed. All hairs from a 4-point barb or wire end were considered one sample.

Rubs were initially identified and set up in 2003 for sampling in 2004. Because we removed tags and wire on monitored rubs at the end of the 2004 field season, the rub survey network was re-established in 2009. The time required for initial set up meant that rubs in some remote areas were not set up until late summer 2009 or spring 2010, leading to lower rub sampling effort and extent in 2009 than in all other years.

Genetic analysis

In 2004, we attempted genetic analysis on all hair samples collected. During 2009—2012, we subsampled hair for analysis to limit genotyping costs. When we collected more than one hair sample during a visit to a rub, we genotyped the two samples with the most and/or largest follicles to maximize the amount of DNA in the sample. If sample quality was equal, we selected non-adjacent samples to increase the probability of detecting more than one individual per site. If there were distinct differences in hair colour between samples, we selected the best 2 samples from each, if available.

Spatial capture-recapture modeling

We included several trap covariates to describe detection. We calculated security level based on the location of a trap, where National Park Service =10, Forest Service =7, State and private forest lands =3, and all other =1. We used the standard deviation of curvature within a 250-m buffer of the trap covariate based on Ironside *et al.*²³, which found it to be the most robust measure of terrain ruggedness. We also evaluated the effects of percent canopy cover based on the cover layer from Landfire, using the most recent data for that year. We updated the 2004 layer to reflect reductions in canopy cover due to some logging and extensive fires between the 2000/2001 imagery and the 2004 dataset.

Results

Rub sampling effort

Sampling primarily occurred in forested habitats, however, bears' predilection to rub on fence and utility poles ensured sampling opportunities in prairie and other treeless habitats and agricultural areas. In 2004, search effort to identify rubs was lower along the eastern edge of our study area because there were few trails or little bear sign to guide our efforts. Sampling in this area was primarily in riparian corridors. Anecdotal information also indicated that few

bears ranged east of the Rocky Mountain foothills at that time. After 2004, we expanded the number of monitored rub objects along the eastern foothills and plains as bears began to reoccupy former range. Starting in 2009, additional rub objects on the eastern edge were, therefore, added to the monitoring network. Rubs continued to be identified and added as sampling sites during early 2010 and then they generally remained in our monitoring network for the duration through 2012. A small number (<1%) of trees blew down or could not be relocated during our study; however, other monitored rubs were typically < 1 km away and in some cases, nearby new rubs were added to the monitoring network to replace them. As a result, sampling distribution and effort was nearly identical 2010—2012.

Discussion

Our point estimate for the annual rate of change, 1.043, was fairly high for an interior grizzly/brown bear population. Stable growth rates ($\lambda \sim 1.0$) have been reported in the past for a number of interior populations in the Rocky Mountains of the United States and Canada (e.g. ^{24,25}). More recently, however, several populations have grown at rates comparable to those we found in the NCDE, likely a reflection of demographic recovery in response to conservation measures^{26,27,28}. A study examining demographic mechanisms underpinning genetic assimilation of remnant groups of grizzly bears in the NCDE found that the increases in genetic diversity they identified resulted primarily from immigration²⁹. It is likely that a combination of growth from within the remnant population coupled with bears moving in from higher density areas and the mathematics of small numbers in a local area (a change from 1 to 2 bears is a 100% increase) was responsible for the highest rates of pixel-specific growth we found in the NCDE (Figure 3). Quantifying local change rates using alternate measures such as gain or loss in density per pixel may be useful in smoothing this effect.

Bear species throughout the world exhibit rubbing behaviour; e.g., grizzly³⁰, European brown bear^{31,32}, Japanese brown bear³³, American black bears³⁴, and Asiatic black bears¹⁵. A growing number of studies have taken advantage of this behaviour to study bear demography (e.g., ^{8,11,35}), dispersal²⁹, and behaviour^{7,36}. Developing a better understanding of which bears rub, why they rub, and where and when they do it will make this sampling method more powerful. The data obtained from genetic detection can inform multiple aspects of population status and mechanisms. For example, studies of habitat effects on abundance³⁸, connectivity and dispersal^{38,39}, the role of social learning³⁶, genetic assimilation in remnant subpopulations²⁹, and the relationship between effective and demographic population size in continuously distributed populations⁴⁰, were based on the genetic samples collected for this and earlier genetic detection studies in the NCDE^{3,8}.

Efficient sampling methods increase sample coverage while decreasing monitoring costs. The number of individual bears we sampled through hair collection at rub sites was an order of magnitude larger and more evenly spatially distributed, e.g. in the expansion area, than would typically be possible through live capture methods. Thus, we expect that genetic monitoring will be more sensitive to spatial and temporal changes in population demography than approaches that involve fewer individuals. Our genetic detection network yielded large sample

sizes and precise estimates of population growth rate with 5 years of rub detection data (2004, 2009–2012). Other approaches to obtaining similar spatial data, such as a grid of bear hair corrals, cost millions of dollars to implement across a region of this size⁸; repeated sampling to obtain trend estimates using these methods across areas as broad as the NCDE has to date proved unsustainable. It may be possible to monitor populations such as ours with adequate precision with less sampling effort and at lower cost than in this study. The use of multiple detection methods¹⁷, clustered and stratified sampling designs^{19,35}, adding telemetry data, or integrated population models that combine SCR data with lower resolution information^{41,42} could further reduce the sampling and, thus, expense of SCR-based studies in the future.

References

1. Clapham, M., Nevin, O. T., Ramsey, A. D. & Rosell, F. The function of strategic tree selectivity in chemical signalling of brown bears. *Animal Behaviour* **85**, 1351–1357 (2013).
2. Boulanger, J. *et al.* Use of occupancy models to estimate the influence of past live captures on detection probabilities of grizzly bears using DNA hair snagging methods. *Journal of Wildlife Management* **72**, 589–595 (2008).
3. Kendall, K. C. *et al.* Grizzly bear density in Glacier National Park, Montana. *Journal of Wildlife Management* **72**, 1693–1705 (2008).
4. Burst, T. L. & Pelton, M. R. Black bear mark tree in the Smokey Mountains. *International Conference on Bear Research and Management* **5**, 45–53 (1983).
5. Green, G. I. & Mattson, D. J. Tree rubbing by Yellowstone grizzly bears (*Ursus arctos*). *Wildlife Biology* **9**, 1–9 (2003).
6. Clapham, M., Nevin, O. T., Ramsey, A. D. & Rosell, F. Hypothetico-deductive approach to assessing the social function of chemical signalling in a non-territorial solitary carnivore. *PLoS ONE* **7**, e35404; [10.1371/journal.pone.0035404](https://doi.org/10.1371/journal.pone.0035404) (2012).
7. Clapham, M., Nevin, O. T., Ramsey, A. D. & Rosell, F. Scent-marking investment and motor patterns are affected by the age and sex of wild brown bears. *Animal Behaviour* **94**, 107–116; [10.1016/j.anbehav.2014.05.017](https://doi.org/10.1016/j.anbehav.2014.05.017) (2014).
8. Kendall, K. C. *et al.* Demography and genetic structure of a recovering grizzly bear population. *Journal of Wildlife Management* **73**, 3–17 (2009).
9. Kendall, K. C. *et al.* Density, distribution, and genetic structure of grizzly bears in the Cabinet-Yaak ecosystem. *Journal of Wildlife Management* **80**, 314–331 (2016).
10. Karamanlidis, A. A., Youlatos, D., Sgardelis, S. & Scouras, Z. Using power poles to document presence of bears in Greece. *Ursus* **18**, 54–61 (2007).

11. Sawaya, M. A., Stetz, J. B., Clevenger, A. P., Gibeau, M. L. & Kalinowski, S. T. Estimating grizzly and black bear population abundance and trend in Banff National Park using noninvasive genetic sampling. *PLoS ONE* **7**, 1–12; [10.1371/journal.pone.0034777](https://doi.org/10.1371/journal.pone.0034777) (2012).
12. Whittington, J., Hebblewhite, M. & Chandler, R. B. Generalized spatial mark–resight models with an application to grizzly bears. *Journal of Applied Ecology* **55**, 157–168 (2018).
13. Woods, J. G. *et al.* Genetic tagging of free-ranging black and brown bears. *Wildlife Society Bulletin* **27**, 616–627 (1999).
14. Taberlet, P. *et al.* Noninvasive genetic tracking of the endangered Pyrenean brown bear population. *Molecular Ecology* **6**, 869–876 (1997).
15. Latham, E., Stetz, J. B., Seryodkin, I., Miquelle, D. & Gibeau, M. L. Non-invasive genetic sampling of brown bears and Asiatic black bears in the Russian Far East. *Ursus* **23**, 145–158 (2012).
16. Murphy, S. M. *et al.* Characterizing recolonization by a reintroduced bear population using genetic spatial capture-recapture. *Journal of Wildlife Management* **80**, 1390–1407 <https://doi.org/10.1002/jwmg.21144> (2016).
17. Boulanger, J. *et al.* Use of multiple data sources to improve DNA-based mark-recapture population estimates of grizzly bears. *Ecological Applications* **18**, 577–589 (2008b).
18. Kendall, K. C. *et al.* Density, distribution, and genetic structure of grizzly bears in the Cabinet-Yaak ecosystem. *Journal of Wildlife Management* **80**, 314–331 (2016).
19. Stenhouse, G. B. *et al.* Estimates of grizzly bear population size and density for the 2014 Alberta Yellowhead Population Unit (BMA 3) and south Jasper National Park Inventory Project. Report prepared for Weyerhaeuser Ltd., West Fraser Mills Ltd, Alberta Environment and Parks, and Jasper National Park. <https://friresearch.ca/resource/estimates-grizzly-bear-population-size-and-density-final-report> (2015).
20. Lamb, C. T., Walsh, D. A. & Mowat, G. Factors influencing detection of grizzly bears at genetic sampling sites. *Ursus* **27**, 31–44 (2016).
21. Alt, D. D. & Hyndman, D. W. Roadside geology of the northern Rockies. Mountain Press Publishing Company, Missoula, MT (1972).
22. Arno, S. F. Forest Regions of Montana. USDA Forest Service Research Paper INT-218, Intermountain Forest and Range Experiment Station, Forest Service, U. S. Department of Agriculture, Ogden, UT 84401 (1979).

23. Ironside, K.E., Mattson, D.J., *et al.* Geomorphometry in Landscape Ecology: Issues of scale, physiography, and application. *Environment and Ecology Research* 6(5): 397-412 (2018). Doi: 10.13189/eer.2018.060501
24. Wielgus, R. B., Bunnell, F. L., Wakkinen, W. L., & Zager, P. E. Population dynamics of Selkirk Mountain Grizzly Bears. *J. Wildlife Management* **58**, 266—272 (1994).
25. Mace, R. D. & Waller, J. S. Demography and population trend of grizzly bears in the Swan Mountains, Montana. *Conservation Biology* **12**, 1005—1016 (1998).
26. Kindberg, J. *et al.* Estimating population size and trends of the Swedish brown bear *Ursus arctos* population. *Wildlife Biology* **17**, 114—123 (2011).
27. Garshelis, D. L., Gibeau, M. L. & Herrero, S. Grizzly bear demographics in and around Banff National Park and Kananaskis Country, Alberta. *Journal of Wildlife Management* **69**, 277—297 (2005).
28. Morehouse, A. T. & Boyce, M. S. Grizzly bears without borders: Spatially explicit capture-recapture in southwestern Alberta. *Journal of Wildlife Management* **80**, 1152—1166; [10.1002/jwmg.21104](https://doi.org/10.1002/jwmg.21104) (2016).
29. Mikle, N., Graves, T. A., Kovach, R., Kendall, K. C. & Macleod, A. C. Demographic mechanisms underpinning genetic assimilation of remnant groups of a large carnivore. *Proc. Royal Society B* **282**; [10.1098/rspb.2016.1467](https://doi.org/10.1098/rspb.2016.1467) PMID: 27655768 (2016).
30. Kendall, K. C., Metzgar, L. H., Patterson, D. A. & Steele, B. M. Power of sign surveys to monitor population trends. *Ecological Applications* **2**, 422—430 (1992).
31. Karamanlidis, A. A. *et al.* Noninvasive genetic studies of brown bears using power poles. *European Journal of Wildlife Research* **56**, 693—702 (2010).
32. De Barba, M. *et al.* 2010. Comparing opportunistic and systematic sampling methods for non-invasive genetic monitoring of a small translocated brown bear population. *Journal of Applied Ecology* **47**, 172—181 (2010).
33. Sato, Y. *et al.* Selection of rub trees by brown bears (*Ursus arctos*) in Hokkaido, Japan. *Acta Theriologica* **59**, 129—137; [10.1007/s13364-013-0143-z](https://doi.org/10.1007/s13364-013-0143-z) (2014)
34. Burst, T. L. & Pelton, M. R. Black bear mark tree in the Smokey Mountains. *International Conference on Bear Research and Management* **5**, 45—53 (1983).
35. Humm, J. M., McCowen, J. W., Scheick, B. K., & Clark, J. D. Spatially explicit population estimates for black bears based on cluster sampling. *Journal of Wildlife Management* **81**, 1187—1201. [https://doi: 10.1002/jwmg.21294](https://doi.org/10.1002/jwmg.21294) (2017).

36. Morehouse, A.T., Graves, T. A., Mikle, N. & Boyce, M. S. Nature vs. nurture: Evidence for social learning of conflict behavior in grizzly bears. *PLoS ONE* **11**, e0165425; [10.1371/journal.pone.0165425](https://doi.org/10.1371/journal.pone.0165425) (2016).
37. Graves T. A. *et al.* Balancing precision and risk: Should multiple detection methods be analyzed separately in N-mixture models? *PLoS ONE* **7**; [10.1371/journal.pone.0049410](https://doi.org/10.1371/journal.pone.0049410) (2012).
38. Graves, T. A., Chandler, R. B., Royle, J. A., Beier, P. & Kendall, K. C. Estimating landscape resistance to dispersal. *Landscape Ecology* **29**, 1201—1211 (2014).
39. Dilkina, B. *et al.* Trade-offs and efficiencies in optimal budget-constrained multispecies corridor networks. *Conservation Biology* **31**, 192—202; [10.1111/cobi.12814](https://doi.org/10.1111/cobi.12814) (2017).
40. Pierson, J. C., Graves, T. A., Banks, S. C., Kendall, K. C. & Lindenmayer, D. B. Relationship between effective and demographic population size in continuously distributed populations. *Evolutionary Applications* **11**, 1162—1175; [10.1111/eva.12636](https://doi.org/10.1111/eva.12636) (2018).
41. Schaub, M. & Abadi, F. Integrated population models: a novel analysis framework for deeper insights into population dynamics. *Journal of Ornithology* **152** (Supplement 1): S227—S237 (2011).
42. Chandler, R. B. & Clark, J. D. Spatially explicit integrated population models. *Methods in Ecology and Evolution* **5**, 1351—1360 (2014).
43. U.S. Geological Survey. USGS National Elevation Dataset (NED) 1 arc-second Downloadable Data Collection from The National Map 3D Elevation Program (3DEP) - National Geospatial Data Asset (NGDA) National Elevation Data Set (NED): U.S. Geological Survey, <https://www.sciencebase.gov/catalog/item/5825a0c3e4b01fad86db66dc> (2016).

Table S1. Description of full set of candidate models developed to estimate grizzly bear population density using spatial capture recapture data from northwestern Montana, USA. Spatial detection histories were developed through genetic identification of individual bears from hair collected at natural bear rub sites. Model 17 (= Model 8 + a term for translocated bears) and Model 18 (= Model 16 + translocations) did not converge because few individuals were identified as having been translocated in our dataset.

Model ^a	D	p_0	σ
1	~ linear	~ year + dur + jul + jul2 + beh + cover + cover2 + curv + sec	~ year
2	~ linear	~ year + dur + jul + beh + cover + cover2 + curv + sec	~ year
3	~ year	~ year + dur + jul + jul2 + beh + cover + cover2 + curv + sec	~ year
4	~ year	~ year + dur + jul + beh + cover + cover2 + curv + sec	~ year
5	~ year	~ dur + jul + beh + cover + cover2 + curv + sec	~ 1
6	~ year	~ dur + jul + beh + cover + cover2 + curv + sec	~ year
7	~ year	~ dur + jul + jul2 + beh + cover + cover2 + curv + sec	~ 1
8	~ year	~ dur + jul + jul2 + beh + cover + cover2 + curv + sec	~ year
9	~ year	~ year + dur + jul + beh + cover + cover2 + curv + sec	~ 1
10	~ year	~ year + dur + jul + jul2 + beh + cover + cover2 + curv + sec	~ 1
11	~ linear	~ dur + jul + beh + cover + cover2 + curv + sec	~ 1
12	~ linear	~ dur + jul + beh + cover + cover2 + curv + sec	~ year
13	~ linear	~ dur + jul + jul2 + beh + cover + cover2 + curv + sec	~ 1
14	~ linear	~ dur + jul + jul2 + beh + cover + cover2 + curv + sec	~ year
15	~ linear	~ year + dur + jul + beh + cover + cover2 + curv + sec	~ 1
16	~ linear	~ year + dur + jul + jul2 + beh + cover + cover2 + curv + sec	~ 1

^a Model notation:

D	density (bears/1,000 km ²)
p_0	baseline detection probability
σ	sigma; spatial scale parameter related to the amount of space used by each individual
~	“a function of”
year	parameter is year-specific
linear	density is a linear function of time
dur	duration of sampling interval; number of days since previous sampling visit
jul	Julian day; linear effect of season on detection probability
jul2	Julian day squared; quadratic effect of season on detection probability
beh	behavioural response; a visit of some bear at some bear rub changes its subsequent probability of detection at that rub
cover	percent of mean canopy cover within a 250 m radius of the trap cover2 percent of mean canopy cover squared
curv	standard deviation of terrain curvature within a 250 m radius of the trap sec security level as determined by land ownership policies at the trap: 10=NPS, 7=FS, 3= state/other, public, 1=private)

Table S2. Female and male parameter estimates (untransformed) for the most supported models (Table 3) developed to estimate trend 2004—2012 in a grizzly bear population using spatial capture-recapture data from northwestern Montana, USA.

Parameter ^a	Female			Male		
	Estimate	SE	P(> z)	Estimate	SE	P(> z)
D ₀ .(Intercept)	-1.891	0.05	0	-2.25	0.04	0
D.beta.Trend	0.048	0.019	0.011	0.033	0.013	0.011
p ₀ .(Intercept)	-5.3	0.376	0	-5.079	0.106	0
p ₀ .2009	-1.092	0.269	0	0.647	0.119	0
p ₀ .2010	0.508	0.215	0.018	0.104	0.115	0.365
p ₀ .2011	0.217	0.217	0.317	0.281	0.118	0.017
p ₀ .2012	-0.014	0.211	0.946	-0.039	0.114	0.735
p.beh	1.456	0.184	0	1.589	0.12	0
p.dur	0.476	0.1	0	0.51	0.06	0
p.jul	0.542	0.217	0.012	0.124	0.039	0.002
p.jul2	-0.059	0.033	0.075	-0.05	0.007	0
p.cover	-0.207	0.068	0.002	-0.126	0.042	0.002
p.cover2	-0.103	0.058	0.077	-0.022	0.034	0.505
p.curv	0.438	0.124	0	-0.436	0.081	0
p.sec	0.236	0.067	0	0.458	0.047	0
σ.(Intercept)	1.481	0.053	0	2.23	0.027	0
σ.2009	0.284	0.096	0.003	-0.236	0.04	0
σ.2010	-0.367	0.069	0	-0.014	0.04	0.724
σ.2011	-0.304	0.072	0	-0.199	0.041	0
σ.2012	-0.13	0.07	0.062	-0.02	0.041	0.632

^a Parameter notation:

D	density (bears/1,000 km ²)
p ₀	baseline detection probability in 2004
p ₀ .2009-12	year-specific detection probability
p.beh	behavioural response; a visit of some bear at some bear rub changes its subsequent probability of detection at that rub
p.dur	duration of sampling interval; days since previous sampling visit
p.jul	Julian day; linear effect of season on detection probability
p.jul2	Julian day squared; quadratic effect of season on detection probability
p.cover	proportion of canopy cover
p.cover2	proportion of cover squared
p.curv	standard deviation of terrain curvature
p.sec	mean security level as determined by land ownership
σ	sigma; parameter related to the amount of space used by each individual
σ.2009-12	year-specific sigma

Table S3. Model selection results for females including all detections, i.e. **including one additional detection 190 km from the other detections of that bear not included in the models shown in Table 3**. Spatial covariates (spatial covs= cover, cover2, curv) were calculated within a 250-m radius of each rub sampling site. Only models with cumulative weights up to 1 are presented.

Females - with outlier point						
Model	D	p_0	σ	ΔAIC	Weight	CumWt
3	~year	~year+dur+jul+jul2+beh+spatial covs	~year	0.00	0.51	0.51
4	~year	~year+dur+jul+beh+spatial covs	~year	0.15	0.47	0.98
1	~linear	~year+dur+jul+jul2+beh+spatial covs	~year	7.87	0.01	0.99
2	~linear	~year+dur+jul+beh+spatial covs	~year	7.92	0.01	1.00
D	density (bears/1,000 km ²)					
p_0	baseline detection probability					
σ	sigma; spatial scale parameter related to the amount of space used by each individual					
ΔAIC	cumulative change in Akaike Information Criterion					
Weight	measure of support for each model					
CumWt	cumulative measure of support for the models					
~	"a function of"					
linear	density is a linear function of time					
year	parameter is year-specific					
dur	duration of sampling interval; number of days since previous sampling visit					
jul	Julian day; linear effect of season on detection probability					
jul2	Julian day squared; quadratic effect of season on detection probability					
beh	behavioural response; a visit of some bear at some bear rub changes its subsequent probability of detection at that rub					
spatial covs	spatial covariates:					
cover	percent of mean canopy cover					
cover2	percent of mean canopy cover squared					
curv	standard deviation of terrain curvature					
sec	security level as determined by land ownership policies at the trap: 10=National Park Service, 7=Forest Service, 3= state/other public, 1=private)					

Table S4. Female parameter estimates (untransformed) for the most supported models (Table S3) using all detections, i.e. **including one additional detection 190 km from the other detections of that bear not included in the models shown in Tables 3 and 4**. Estimates for model 4 were nearly identical to model 3 and estimates for model 2 were nearly identical to model 1, so are not shown.

Females - with outlier point					
Model 3			Model 1		
Parameter ^a	Estimate	SE	Parameter ^a	Estimate	SE
D ₀ .(Int)	-2.048	0.108	D ₀ .(Int)	-2.000	0.054
D.beta.2009	0.049	0.180	D.beta.linear	0.034	0.017
D.beta.2010	-0.171	0.143			
D.beta.2011	0.271	0.147			
D.beta.2012	0.326	0.141			
p ₀ .(Int)	-5.268	0.373	p ₀ .(Int)	-5.224	0.372
p ₀ .2009	-1.010	0.281	p ₀ .2009	-1.064	0.267
p ₀ .2010	-1.641	0.21	p ₀ .2010	-1.764	0.209
p ₀ .2011	0.195	0.219	p ₀ .2011	0.207	0.216
p ₀ .2012	-0.020	0.211	p ₀ .2012	-0.009	0.209
p.beh	1.838	0.186	p.beh	1.813	0.186
p.dur	0.482	0.097	p.dur	0.485	0.097
p.jul	0.476	0.215	p.jul	0.469	0.214
p.jul2	-0.047	0.033	p.jul2	-0.046	0.033
p.cover	-0.209	0.064	p.cover	-0.208	0.064
p.cover2	-0.123	0.057	p.cover2	-0.121	0.057
p.curv	0.442	0.121	p.curv	0.438	0.121
p.sec	0.256	0.064	p.sec	0.249	0.064
σ.(Int)	1.492	0.054	σ.(Int)	1.497	0.054
σ.2009	0.286	0.098	σ.2009	0.278	0.097
σ.2010	0.890	0.072	σ.2010	0.856	0.070
σ.2011	-0.303	0.073	σ.2011	-0.299	0.073
σ.2012	-0.134	0.071	σ.2012	-0.127	0.071

^a Parameter notation:

D ₀ .(Int)	density (bears/1,000 km ²) (Intercept)
p ₀ .(Int)	baseline detection probability in 2004 (Intercept)
p ₀ .2009-12	year-specific detection probability
p.beh	behavioral response; a visit of some bear at some bear rub changes its subsequent probability of detection at that rub
p.dur	duration of sampling interval; days since previous sampling visit
p.jul	Julian day; linear effect of season on detection probability
p.jul2	Julian day squared; quadratic effect of season on detection probability
p.cover	proportion of canopy cover

$p.cover2$	proportion of cover squared
$p.curv$	standard deviation of terrain curvature
$p.sec$	mean security level as determined by land ownership
$\sigma.(Int)$	baseline sigma (Intercept); related to the amount of space used by each individual
$\sigma.2009-12$	year-specific sigma

Table S5. Point estimates (transformed) of female density (bears/1,000 km²), spatial scale of movement (σ), and baseline detection probability (p_0) parameters for models developed to estimate grizzly bear population trend using spatial capture-recapture data from the Northern Continental Divide Ecosystem in northwestern Montana, USA. a. **Estimates that include one outlier detection of a single female in 2010 that was 190 km distant from other detections of that individual.** b. Estimates that exclude the outlier detection point. The baseline detection probability reported here is for $b=0$, initial behavior covariate (prior to the first detection), and all other covariates at their mean level. When the outlier was included, the best model had year-specific density (model #3) versus a linear yearly trend in density (model #1) when the outlier point was excluded.

a. Females – with outlier point. Best model is #3

Model	Density		σ		p_0	
	3	1	3	1	3	1
2004	8.06	7.37	4.45	4.47	0.005	0.005
2009	8.47	8.75	5.92	5.90	0.002	0.002
2010	6.79	9.06	10.83	10.51	0.001	0.001
2011	10.57	9.38	3.29	3.31	0.006	0.007
2012	11.17	9.71	3.89	3.94	0.005	0.005

b. Females – excluding outlier point. Best model is #1

Model	Density		σ		p_0	
	3	1	3	1	3	1
2004	7.92	7.77	4.39	4.40	0.005	0.005
2009	8.39	9.90	5.89	5.84	0.002	0.002
2010	10.16	10.40	3.04	3.05	0.008	0.008
2011	10.39	10.91	3.24	3.24	0.006	0.006
2012	11.04	11.45	3.86	3.86	0.005	0.005

Figure S1. Locations of grizzly bear detections 2004, 2009—2012 (N=1,539) derived from sampling at natural bear rub sites in the Northern Continental Divide Ecosystem in northwestern Montana, USA. Maps were created from the USGS National Elevation Dataset⁴³ using ArcMap 10.2.

