



## Review

# Climate change effects on biodiversity, ecosystems, ecosystem services, and natural resource management in the United States



Sarah R. Weiskopf<sup>a,\*</sup>, Madeleine A. Rubenstein<sup>a</sup>, Lisa G. Crozier<sup>b</sup>, Sarah Gaichas<sup>c</sup>, Roger Griffis<sup>d</sup>, Jessica E. Halofsky<sup>e</sup>, Kimberly J.W. Hyde<sup>f</sup>, Toni Lyn Morelli<sup>g</sup>, Jeffrey T. Morissette<sup>h</sup>, Roldan C. Muñoz<sup>i</sup>, Andrew J. Pershing<sup>j</sup>, David L. Peterson<sup>e</sup>, Rajendra Poudel<sup>k</sup>, Michelle D. Staudinger<sup>g</sup>, Ariana E. Sutton-Grier<sup>l</sup>, Laura Thompson<sup>a</sup>, James Vose<sup>m</sup>, Jake F. Weltzin<sup>n</sup>, Kyle Powys Whyte<sup>o</sup>

<sup>a</sup> U.S. Geological Survey National Climate Adaptation Science Center, Reston, VA, USA

<sup>b</sup> NOAA Northwest Fisheries Science Center, Seattle, WA, USA

<sup>c</sup> NOAA Northeast Fisheries Science Center, Woods Hole, MA, USA

<sup>d</sup> NOAA National Marine Fisheries Service, Silver Spring, MD, USA

<sup>e</sup> University of Washington, School of Environmental and Forest Sciences, Seattle, WA, USA

<sup>f</sup> NOAA Northeast Fisheries Science Center, Narragansett, RI, USA

<sup>g</sup> U.S. Geological Survey Northeast Climate Adaptation Science Center, Amherst, MA, USA

<sup>h</sup> U.S. Department of the Interior, National Invasive Species Council Secretariat, Fort Collins, CO, USA

<sup>i</sup> NOAA Southeast Fisheries Science Center, Beaufort, NC, USA

<sup>j</sup> Gulf of Maine Research Institute, Portland, ME, USA

<sup>k</sup> NOAA, Silver Spring, MD, USA

<sup>l</sup> University of Maryland Earth System Science Interdisciplinary Center, College Park, MD, USA

<sup>m</sup> U.S. Forest Service Southern Research Station, Raleigh, NC, USA

<sup>n</sup> U.S. Geological Survey, Fort Collins, CO, USA

<sup>o</sup> Michigan State University, East Lansing, MI, USA

## HIGHLIGHTS

- Climate change is affecting ecosystems at multiple scales.
- Individual/species: changes in morphology and behavior, phenology, and range shifts observed
- Ecosystems: shifts in productivity, species interactions, and emergent properties observed
- Together, these changes are impacting ecosystem services and human well-being.
- Natural resource managers need proactive, flexible approaches to deal with changes.

## ARTICLE INFO

## Article history:

Received 9 December 2019

Received in revised form 28 February 2020

Accepted 5 March 2020

Available online 10 March 2020

Editor: Jay Gan

## Keywords:

Global change

Biodiversity

Ecosystems

Ecosystem services

Natural resource management

## ABSTRACT

Climate change is a pervasive and growing global threat to biodiversity and ecosystems. Here, we present the most up-to-date assessment of climate change impacts on biodiversity, ecosystems, and ecosystem services in the U.S. and implications for natural resource management. We draw from the 4th National Climate Assessment to summarize observed and projected changes to ecosystems and biodiversity, explore linkages to important ecosystem services, and discuss associated challenges and opportunities for natural resource management. We find that species are responding to climate change through changes in morphology and behavior, phenology, and geographic range shifts, and these changes are mediated by plastic and evolutionary responses. Responses by species and populations, combined with direct effects of climate change on ecosystems (including more extreme events), are resulting in widespread changes in productivity, species interactions, vulnerability to biological invasions, and other emergent properties. Collectively, these impacts alter the benefits and services that natural ecosystems can provide to society. Although not all impacts are negative, even positive changes can require costly societal adjustments. Natural resource managers need proactive, flexible adaptation strategies that consider historical and future outlooks to minimize costs over the long term. Many organizations are beginning to explore these approaches, but implementation is not yet prevalent or systematic across the nation.

© 2020 The US Geological Survey. Published by Elsevier B.V. All rights reserved. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

\* Corresponding author.

E-mail address: [sweiskopf@usgs.gov](mailto:sweiskopf@usgs.gov) (S.R. Weiskopf).

# Contents

|      |                                                                               |    |
|------|-------------------------------------------------------------------------------|----|
| 1.   | Introduction . . . . .                                                        | 2  |
| 2.   | Individuals, populations, and species . . . . .                               | 2  |
| 2.1. | Behavior and morphology . . . . .                                             | 2  |
| 2.2. | Phenology . . . . .                                                           | 3  |
| 2.3. | Geographic range shifts . . . . .                                             | 4  |
| 2.4. | Mechanisms and rate of change . . . . .                                       | 4  |
| 3.   | Ecosystems . . . . .                                                          | 4  |
| 3.1. | Primary productivity . . . . .                                                | 4  |
| 3.2. | Species interactions, emergent properties, and biological invasions . . . . . | 5  |
| 3.3. | Extreme events and ecosystem resilience . . . . .                             | 6  |
| 4.   | Ecosystem services . . . . .                                                  | 6  |
| 4.1. | Provisioning services . . . . .                                               | 7  |
| 4.2. | Regulating services . . . . .                                                 | 7  |
| 4.3. | Supporting services . . . . .                                                 | 7  |
| 4.4. | Cultural services . . . . .                                                   | 7  |
| 5.   | Vulnerability of human communities . . . . .                                  | 7  |
| 6.   | Implications for natural resource management . . . . .                        | 8  |
| 6.1. | Increasing ecosystem resilience . . . . .                                     | 8  |
| 6.2. | Leveraging technology and infrastructure . . . . .                            | 9  |
| 6.3. | Strengthening governance . . . . .                                            | 9  |
| 7.   | Conclusion . . . . .                                                          | 11 |
|      | Acknowledgements . . . . .                                                    | 11 |
|      | References . . . . .                                                          | 11 |

## 1. Introduction

Climate change is a pervasive and growing global threat to biodiversity and ecosystems (Díaz et al., 2019). Climate change affects individual species and the way they interact with other organisms and their habitats, which alters the structure and function of ecosystems and the goods and services that natural systems provide to society (Díaz et al., 2019). Understanding the direction and magnitude of ecological responses allows human communities to better anticipate these changes and adapt as necessary.

Periodic assessments of current and future climate change impacts on ecosystems are important for developing and updating natural resource management plans and evaluating adaptation actions (West et al., 2009). The National Climate Assessment (NCA) is a key assessment in the United States, required by the Global Change Research Act to summarize current and projected impacts of climate change on a variety of sectors and regions in the U.S. every four years (USGCRP, 2018). Here, we draw upon the recently published Fourth NCA (NCA4) Volume II to present the most up-to-date assessment of climate change impacts on biodiversity, ecosystems, and ecosystem services in the U.S. (USGCRP, 2018). We synthesize, extend, and integrate the NCA4 chapters focused on natural resources: “Ecosystems, Ecosystem Services, and Biodiversity” (Ch. 7); “Forests” (Ch.6); “Oceans and Marine Resources” (Ch.9); “Coastal Effects” (Ch. 8); and “Tribes & Indigenous Peoples” (Ch.15) (USGCRP, 2018).

We provide a more in-depth, technical analysis of topics of interest to scientists and practitioners, and review climate change impacts at multiple scales, including: 1) the individual organisms, populations, and species of biodiversity which comprise ecosystems; 2) the properties and processes that characterize ecosystems; and 3) the goods and services that ecosystems provide which support human economies and well-being (Fig. 1). Further, we explore natural resource management challenges posed by climate change and present examples of on-the-ground adaptation actions. Many topics covered in this review are complex and deserve a review of their own. However, by covering multiple scales in one place, we provide a holistic overview of how climate change is affecting different ecosystems and how these changes may in turn affect human well-being, including impacts to vulnerable communities, tribes, and Indigenous peoples.

## 2. Individuals, populations, and species

Although climate change impacts are widespread, they are not uniform, and accumulating evidence indicates that climate change responses vary as a function of relative vulnerability due to differences in exposure, sensitivity, and adaptive capacity (Beever et al., 2016; Foden and Young, 2016; Glick et al., 2011; Kovach et al., 2019). Below, we discuss major impacts observed at the scale of individuals, populations, and species, and review the mechanisms driving changes.

### 2.1. Behavior and morphology

One way that organisms cope with changes in their environment is by altering their behavior or morphology. Behavioral responses to climate change can result from changes in temperature and manifest before changes at the population and species level, such as distribution changes or population declines (Beever et al., 2017). Behavioral responses include seeking shade or refuge, altering feeding times, changing site use, and shifting circadian or circannual rhythms (e.g., hibernation, migration; Beever et al., 2017; Bradshaw and Holzapfel, 2007; McCann et al., 2017).

Morphological changes commonly entail changes in body size (Cheung et al., 2013; Eastman et al., 2012; Ozgul et al., 2010). For example, increasing summer temperatures have been associated with reduced body size and increased wing length in North American migratory birds (Weeks et al., 2019). In ectotherms, where metabolic rate is sensitive to temperature (Gardner et al., 2011), warmer temperatures can lead to faster growth rates but can ultimately lead to smaller body size (Atkinson, 1994). This direct impact of temperature on growth has been observed, for example, in American lobster (*Homarus americanus*; Le Bris et al., 2017) and Atlantic cod (*Gadus morhua*; Pershing et al., 2016) during recent warming in the north-west Atlantic. Morphological responses, however, are complex and highly variable: changes in phenotype may not be observed if genetic change is counteracted by environmental effects (Conover et al., 2009). Moreover, short term benefits may not be adaptive in the long term (Section 2.4).




**Fig. 1.** Climate change and non-climate stressors interact and affect ecological systems at multiple scales. These combined stressors affect individuals, populations, and species, as well as ecosystem processes and properties. The relative impact of climate change versus other stressors varies depending on the species or ecosystem. Diverse biological communities and functioning ecosystems are critical to maintaining the ecosystem services (Millennium Ecosystem Assessment, 2005) that support human well-being (Díaz et al., 2019). Natural resource management affects biodiversity, ecosystems and their services and can moderate or exacerbate climate change and non-climate stressors.

## 2.2. Phenology

Phenology, or the seasonal timing of recurring biological events, is a critical part of ecological relationships (Rudolf, 2019), and a primary indicator of species responses to climate change (Staudinger et al., 2019). Across much of the terrestrial U.S., broad changes in seasonality are evidenced by an earlier start to spring compared to 20th-century averages (Ault et al., 2015; Monahan et al., 2016). Although changes in phenology are well documented, trends are far from homogenous (Cohen et al., 2018), a result of high variability in climate drivers and phenological responses across habitat types (Chmura et al., 2019; Fu et al., 2015; Pearson, 2019). Migratory birds provide clear examples of phenological shifts, with extensive documentation of earlier migration (Lehikoinen et al., 2019) and earlier breeding (Lany et al., 2016) in response to rising temperatures and altered precipitation patterns.

Phenological shifts in marine and aquatic habitats are less well documented in comparison to terrestrial systems, largely due to difficulty detecting and tracking aquatic organisms (Staudinger et al., 2019). Nonetheless, there have been clear, directional shifts in the timing of seasonal aquatic and marine abiotic drivers, including earlier transitions from winter to spring temperatures (Thomas et al., 2017) and earlier ice melting and runoff (Post, 2017; Staudinger et al., 2019). Marine

phytoplankton can respond rapidly to such abiotic changes, resulting in altered timing of phytoplankton blooms (Wasmund et al., 2019), which in turn can create a mismatch with secondary consumers and change the food web structure (Post, 2017; Sundby et al., 2016). Phenological changes have also been observed in freshwater and riparian systems, including advances in the winter spawning phenology of coho (*Oncorhynchus kisutch*) and chum salmon (*O. keta*) in the Pacific Northwest, which has driven phenological changes in bald eagle (*Haliaeetus leucocephalus*) populations (Rubenstein et al., 2019).

Differential shifts in phenology among interacting organisms could drive population declines through reduced reproductive success and/or increased predation or competition (Visser and Gienapp, 2019; Wann et al., 2019; Zimova et al., 2016). Additionally, phenology changes in species with multiple life stages are complex and shifts that are beneficial for one life stage may be detrimental to another (Campbell et al., 2019; Schluter et al., 1991). However, few species have had documented population-level consequences of mistimed reproduction, perhaps due to mitigating effects of density-dependence and greater ability to alter prey or behavior (Dunn and Møller, 2014; Staudinger et al., 2019). Asynchronous phenological shifts have the potential to disrupt the functioning, persistence, and resilience of population dynamics, ecosystems, and ecosystem services (Asch et al., 2019; Mayor et al., 2017; Staudinger et al., 2019; Visser and Gienapp, 2019).

### 2.3. Geographic range shifts

Climate change is driving large-scale shifts in species distribution, abundance, and reorganization of terrestrial and aquatic ecosystems (Lenoir and Svenning, 2015; Pacifici et al., 2017; Staudinger et al., 2012). Geographic range shifts are widespread across taxa and ecosystems: a recent review of plant and animal species in temperate North America found 55% have either contracted the warm edge or expanded the cool edge of their range (Wiens, 2016). Documented shifts poleward, upslope, and deeper average tens of kilometers per decade (Burrows et al., 2011; Chen et al., 2011). Northern Hemisphere birds, for example, are decreasing in abundance along species' southern and lower elevational range edges (Ralston et al., 2017; Tayleur et al., 2015).

Marine organisms have also demonstrated range shifts, in some cases at faster rates than observed in terrestrial systems and at pace with climate velocities (García Molinos et al., 2015; Poloczanska et al., 2013). The majority of marine taxa surveyed in a 2013 review, for example, shifted in directions consistent with climate velocity (Pinsky et al., 2013). This response, however, is not uniform, with some fish demonstrating a lag effect, potentially due to species-specific sensitivities (Alabia et al., 2018). Arctic marine environments are experiencing changes to sea ice cover, increasing temperatures, and ocean acidification, resulting in range shifts for marine fish, arthropods, and marine mammals (Mecklenburg et al., 2016; Taylor et al., 2017; U.S. Environmental Protection Agency, 2016). Some temperate marine ecosystems are 'tropicalizing', with herbivorous tropical fish expanding poleward, causing decreases in macroalgal plant communities (Vergés et al., 2014).

Despite evidence for widespread range shifts, fewer shifts have been documented than expected from projections, and some shifts are counterintuitive to expectations from projections based solely on temperature changes (Crimmins et al., 2011; Foster and D'Amato, 2015; Morley et al., 2017; Rowe et al., 2015). Indirect effects of climate change and interactive ecological and evolutionary processes can complicate predictions (Estrada et al., 2016; MacLean and Beissinger, 2017; Pacifici et al., 2017). Microclimates, complex topography, and factors such as land use change also need to be considered to accurately predict shifts (Elsen and Tingley, 2015; Guo et al., 2018; Hannah et al., 2014; Kleisner et al., 2016, 2017; Sirami et al., 2017). Moreover, documenting range shifts is challenging, requiring baseline information on historical ranges that is missing for many species, ecosystems, and geographic areas. Additionally, life stages that are most responsive to climate may not be the focus of monitoring or might show lag effects (Alexander et al., 2018). Limited and occasionally conflicting evidence could actually indicate that some species are able to adapt in place, at least for now (Beever et al., 2015; Berg et al., 2010; Jurgens and Gaylord, 2018). Although range shifts are generally regarded as a primary mechanism by which species can effectively adapt to climate change, shifts should be considered in the context of entire ecosystems: an adaptive shift for one species may negatively impact recipient communities (Wallingford et al., in review) (Box 1).

### 2.4. Mechanisms and rate of change

An organism's response to climate change can be driven by genetic (evolutionary) or non-genetic (plastic) processes (e.g., Franks et al., 2014; Kingsolver and Buckley, 2017). This distinction is important because the mechanism determines the rate of response and whether individuals, populations, and species will be able to keep pace with rapidly changing conditions (Boutin and Lane, 2014). Plastic responses occur within an individual's lifetime and are almost immediate, whereas evolutionary change requires multiple generations (Harrisson et al., 2014; Hendry et al., 2011). Current research is starting to explore the role of epigenetic responses, wherein environmental drivers alter gene expression and can be passed to future generations, occur between generations, and are considered intermediate responses (Jeremias et al.,

2018). The distinction between plastic/epigenetic responses and evolutionary change is not always clear, as an organism's ability to respond through these mechanisms can be heritable and subject to evolutionary pressure (Banta and Richards, 2018; Grenier et al., 2016).

Some rapid responses reflect a long history of genetic adaptation to natural variability in climate, and may facilitate persistence during directional climate change by allowing populations to persist long enough for genetic adaptation to occur (Fox et al., 2019; Snell-Rood et al., 2018). While often effective at increasing survival in the short term, some plastic responses are not beneficial over the long term (Ghalambor et al., 2007): they may entail tradeoffs with fecundity (e.g., smaller body sizes typically produce fewer eggs and more boom/bust population dynamics; Waples and Audzijonyte, 2016), or they may lead to interactions with other species or habitats that ultimately lower survival (Bonamour et al., 2019; Schlaepfer et al., 2002). Negative effects of plastic responses are often delayed and difficult to measure, so tracking long-term demographic responses to ensure populations of concern are truly coping with climate change is important.

The pace of climate change often exceeds average rates of evolutionary change (De Meester et al., 2018). However, evolution can happen in very few generations if populations survive strong selection and favorable genetic variation is already present (Hendry, 2017). Strong selection typically involves high mortality, so populations face extirpation before they can effectively adapt via evolution (Bay et al., 2018). Indeed, recent meta-analyses demonstrate that even species demonstrating adaptive phenotypic responses may be adapting too slowly to keep pace with climate change (Radchuk et al., 2019).

Substantial theoretical and empirical work has focused on predicting and measuring rates of response to climate change in recent years (Bell, 2013; Carlson et al., 2014; Gomulkiewicz et al., 2018; Kopp and Matuszewski, 2014; Pelletier and Colman, 2018). Rapid trait changes are common and well documented, but most cases are consistent with plastic rather than evolutionary mechanisms (Eastman et al., 2012; Merilä and Hendry, 2014). Although examples do exist of evolutionary responses across multiple taxa (Boutin and Lane, 2014; Charmantier and Gienapp, 2014; Crozier and Hutchings, 2013; Franks et al., 2014), the comparative lack of evidence may be due to complex selection and genetic landscapes, methodological constraints that hinder measurement of genetic change (Merilä, 2012), and altered species interactions which may outpace evolutionary responses (Section 3.2). Evolutionary rates are complicated to predict because selection acts on many traits simultaneously, possibly with opposing costs and benefits in different life stages or habitats (Crozier et al., 2008). However, because responses vary in the extent to which they reduce extinction risk, tracking multiple responses and understanding their limitations is critical for successful resource management.

## 3. Ecosystems

Observed ecosystem-level changes in response to climate change are due to direct impacts from changing climate drivers and interacting effects of species- and population-level responses. Here, we focus on several key ecosystem-level characteristics and properties affected by climate change: primary production; species interactions and emergent properties, including biological invasions; and the impact of extreme events on ecosystem resilience.

### 3.1. Primary productivity

Almost all life on Earth relies on primary producers, photosynthetic organisms that are the foundation of most food webs and are responsible for producing Earth's oxygen and regulating important components of carbon cycling and sequestration. Climate change has had varying effects on primary production across spatial and temporal scales (Lipton et al., 2018). Changes in primary production are likely to be amplified at higher trophic levels (Chust et al., 2014; Lefort et al., 2015; Stock

## Box 1

Examples of climate-driven changes to ecosystems.

## Terrestrial



Photo: U.S. Geological Survey



Photo: U.S. Forest Service

The 2012–2017 drought across California resulted in direct physiological stress to trees and facilitated bark beetle outbreaks, causing an unprecedented mortality of 129 million trees in Sierra Nevada forests (Asner et al., 2016; Cal Fire, 2018; Stephens et al., 2018). High levels of mortality among ponderosa pine led to increases in the relative proportion of incense cedar (*Calocedrus decurrens*), which has increased the likelihood of high-intensity surface fires and large wildfires (Stephens et al., 2018). The combined effects of drought, insects, and altered community composition has drastically changed this ecosystem.

## Marine



Photo: NOAA



Photo: New York State Department of Environmental Conservation

Non-native invasive species such as European green crab (*Carcinus maenas*) are now present off the California coast, alongside recent "invasions" (due to climate-driven range expansion) of the Humboldt squid (*Dosidicus gigas*) (Epstein and Smale, 2017; Grosholz et al., 2000; Zeidberg and Robison, 2007). Both Humboldt squid and green crab are voracious predators that feed on a variety of native prey (Field et al., 2013; Grosholz et al., 2000). Although ultimate synergistic impact on California marine communities is not yet known, impacts from these species that have already been observed separately include prey population reductions, alteration of the resource base available to migratory shorebirds, and potential fundamental modification of ecosystem structure.

et al., 2014), resulting in further changes to ecosystem function and potentially substantial changes to entire ecosystems.

Globally, terrestrial primary production increased during the late 20th and early 21st centuries due to the fertilizing effect of increasing atmospheric CO<sub>2</sub>, nutrient additions from human activities, longer growing seasons, and forest regrowth (Campbell et al., 2017; Domke et al., 2018; Graven et al., 2013; Wenzel et al., 2016; Zhu et al., 2016). However, regional trends vary, and different components of climate change may have opposing impacts on production; while increased atmospheric CO<sub>2</sub> can increase vegetation growth (e.g., Norby et al., 2002), excess or lack of nutrients, water deficits, and air pollution can limit growth (Norby et al., 2010; Oren et al., 2001; Pan et al., 2009). Warming and increased atmospheric CO<sub>2</sub> may also affect belowground biogeochemical processes, such as carbon and nitrogen cycling (Melillo et al., 2017), which can affect terrestrial production (Campbell et al., 2009).

Climate-driven changes to forest primary productivity vary by forest type and elevation. Primary production will likely decrease in forests where soil water availability is limited during the growing season (Latta et al., 2010), but will likely increase in energy-limited forests where snow and cold temperatures restrict the growing season (Latta et al., 2010; Marcinkowski et al., 2015; Wang et al., 2017). However, even in energy-limited forests, drought and extreme temperatures

could limit growth increases (Anderegg et al., 2015; Hember et al., 2017). It is also unclear if fertilization effects will continue as forests age (Norby et al., 2010).

In marine and aquatic systems, phytoplankton is responsible for nearly all primary production and generates almost half of the total global primary production. Phytoplankton growth rates affect CO<sub>2</sub> uptake from seawater and organic carbon export to the deep ocean, and also impact fisheries productivity (Tyrrell, 2019). Like terrestrial systems, climate change impacts to marine primary production vary regionally; warming in temperate and tropical oceans can increase stratification, limiting upwelling of deep nutrients that stimulate new production (IPCC, 2013). In contrast, reduced ice cover at higher latitudes increases sunlight availability to the ocean surface, increasing phytoplankton growing seasons and annual primary production (Wasmund et al., 2019). Understanding how these changes impact the food web is crucial for maintaining sustainable fisheries.

### 3.2. Species interactions, emergent properties, and biological invasions

Variability in species' exposure and responses to climate change are primary drivers of altered species interactions. Emergent properties of ecosystems, including community characteristics such as food-web

structure and function that are mediated by species interactions, are changing as species shift their distributions (Section 2.3) and phenologies (Section 2.2) in response to climate impacts (Breeggemann et al., 2016). Higher trophic levels, for example, are expected to be more sensitive to climate change because of changes in predatory demand and search and encounter rates (Abbott et al., 2014; Daly and Brodeur, 2015; Dell et al., 2014). However, functional responses vary and depend on species composition, abiotic conditions (Davis et al., 2017; McCluney and Sabo, 2016; Verdeny-Vilalta and Moya-Laraño, 2014), body size (Ewald et al., 2013), and predator-prey interactions (Peers et al., 2014; Van Zuiden et al., 2016). Collectively, these changes are resulting in novel (i.e., species composition or function is completely transformed) or hybrid (i.e., some historical characteristics persist but structure and composition lie outside the historical range of variability) ecosystems and altered interspecific relationships with no historical analog (Hobbs et al., 2009).

Emergent ecosystem properties are more difficult to predict than direct impacts to individual species because they develop from interactions radiating throughout the system. Modeling is commonly used to evaluate changes in species interactions, but high uncertainty remains for many species and ecosystems due to lack of baseline research on biotic interactions, community structure and function, adaptive capacity, and interactions of climate and non-climate stressors (Beever et al., 2016; Blois et al., 2013; Rosenblatt and Schmitz, 2014). A growing number of studies are assessing how direct and indirect climate impacts may alter ecosystem properties. For example, fossil records have shown that ecosystems respond similarly to past climate change events by becoming dominated with generalist species (Blois et al., 2013) which may be an indicator of what could be expected under contemporary climate change. Research is also providing more nuanced evaluations of behavioral changes, including phenological, reproductive, and foraging changes (Beever et al., 2016), such as how increasing temperatures are increasing ratsnake predation on birds in the southeastern U.S. (DeGregorio et al., 2015). Finally, studies are suggesting that possessing population-specific traits (Dell et al., 2014; Rasmann et al., 2014) or local adaptations (Davis et al., 2017; Herstoff and Urban, 2014) may allow species to transition and persist in novel environments (Cannizzo and Griffen, 2016). More sophisticated models that account for multi-species interactions, community structure, dispersal, and evolution are needed to make robust predictions, although complex models may not always increase predictive capability (Alexander et al., 2016; Herstoff and Urban, 2014; Novak et al., 2011; Young, 2014).

Of additional concern is that climate change is facilitating the introduction and spread of non-native invasive species. Global economic costs of invasive species are currently estimated to be over \$1.4 trillion annually and climate change has the potential to intensify these impacts (Burgiel et al., 2014). Many non-native invasive species are opportunistic generalists that can take advantage of changing conditions, colonize disturbed areas, and out-compete species, thereby altering community composition, dominance, production, and increasing extinction risk in some cases (e.g., Schmitt et al., 2019; Yeruham et al., 2020). For example, disturbance along with warming temperatures, was found to be important for plant invasions at cold, high-altitude regions in South America and Scandinavia (Lembrechts et al., 2016). Moreover, many non-native invasive plants respond more positively than native plants to increasing CO<sub>2</sub>, nitrogen deposition, and temperature, likely increasing their competitiveness under increasing climate change (Liu et al., 2017). Stronger competitive abilities will likely lead to higher non-native invasive plant abundance and declines of native species abundances and community diversity (Bradley et al., 2019). Infiltration of non-native species into natural communities has already negatively impacted biodiversity (Bradley et al., 2019). Although new interactions resulting from invasive and native range-shifting species often negatively impact ecosystems (Carey et al., 2012; Valéry et al., 2009; Vergés et al., 2016; Wallingford et al., in review) and related services (Blois et al., 2013), species responses can have counter-intuitive

outcomes. For example, the introduction of a non-native invasive bryozoan in the Gulf of Maine is thought to have resulted in substantial expansion of a native nudibranch species due to increased availability of a novel food source (Dijkstra et al., 2013). In some cases, climate change could also indirectly benefit native species by reducing populations of invasive species (Wenger et al., 2011).

### 3.3. Extreme events and ecosystem resilience

Climate change has altered the duration, magnitude, and frequency of extreme events, including droughts, forest fires, and heatwaves (Jay et al., 2018). Many of these events have significant impacts on ecosystems and interact with other climate-driven changes, reducing ecological resilience.

More extreme droughts and wildfires, driven by rising temperatures and altered precipitation patterns, affect ecosystem structure and function, particularly in forested ecosystems. Over the last two decades, warming and more variable precipitation have increased forest drought severity in the West, Southeast, and the Lake States (Clark et al., 2016), reducing tree growth and increasing mortality (Peters et al., 2015). However, responses vary (Choat et al., 2012) and can be lagged in long-lived species (Walter et al., 2013). Drought weakens tree defenses, increasing susceptibility to other disturbances, including insects, pathogens, invasive species (Trottier et al., 2017), and wildfires (Littell et al., 2016; Logan and Powell, 2009; Weed et al., 2013; see Box 1). While drought impacts have direct long-term consequences, drought-facilitated disturbances can result in more immediate changes to forest ecosystem structure and function (Loehman et al., 2017).

Earlier spring warming (Westerling et al., 2006), increased vapor pressure deficit (Abatzoglou and Williams, 2016), and reduced summer precipitation (Dennison et al., 2014; Holden et al., 2018) have increased fire season length and area burned across the western U.S. (Abatzoglou and Kolden, 2013; Gergel et al., 2017; Luce et al., 2014; McKenzie and Littell, 2017; Westerling, 2016). Climate changes have interacted with forest management practices to create large, dense forest with high fuel loads, especially in lower-elevation ponderosa pine (*Pinus ponderosa*) and dry mixed-conifer forests in the West (Keane et al., 2002). Fire frequency and area burned will likely increase in fire-prone forests (Barbero et al., 2015; Liu et al., 2015); by the 2050s, annual area burned in the U.S. might increase by 2–6 times compared to present (Kitzberger et al., 2017; Litschert et al., 2012; Ojima et al., 2014). However, frequency and severity will depend on topography, fuel levels, and fire suppression efforts (Abt et al., 2015; Butry et al., 2010).

In the North Atlantic, storms are getting stronger due to rising ocean temperatures and sea level rise (Elsner et al., 2008; Malmstadt et al., 2010). There is also evidence that tropical cyclones are bringing more extreme rainfall, even if wind speeds have not increased (Patricola and Wehner, 2018). Increased storm intensity can impact ecosystems and human communities through extreme flooding, erosive waves, and higher storm surge, making recovery from extreme events more challenging. Rising ocean temperatures have also led to periods of extraordinarily warm conditions across the globe, known as marine heatwaves (Hobday et al., 2016, 2018). The U.S. experienced significant heatwaves in the northwest Atlantic in 2012 (Mills et al., 2013) and 2016 (Pershing et al., 2018) and the northeast Pacific in 2014–2015 (Bond et al., 2015). Rising ocean temperatures are driving widespread coral bleaching, contributing to coral cover loss, impacting fish communities, and increasing exposure of nearby shores to waves (Eakin et al., 2019).

## 4. Ecosystem services

Diverse biological communities and functioning ecosystems are critical to maintaining ecosystem services that support human well-being (Díaz et al., 2019). Therefore, climate change impacts to species, populations, and ecosystems affect the availability and delivery of ecosystem

services, including changes to provisioning, regulating, supporting, and cultural services.

#### 4.1. Provisioning services

Climate-induced changes in provisioning services, the material goods that people obtain from ecosystems and biodiversity, can have profound impacts on human economies and well-being. For example, climate impacts to forested watersheds, including increased temperatures, changes to precipitation and snowfall, and disturbances such as wildfires, are altering freshwater supply for municipalities, agriculture, and power generation (Barnett et al., 2008; Stewart et al., 2005). Surface water shortages are likely in dry years in some locations (Li et al., 2017). Rising stream temperatures also affect water quality (Warziniack et al., 2018). Similarly, wildfires can increase sediment deposition and debris in streams, lakes, and reservoirs (Luce et al., 2012). These changes will stress water supplies, potentially increasing water treatment costs (Warziniack et al., 2018).

Changes in water supply, along with other climate change impacts, can alter agricultural production. Droughts and other extreme events can decrease crop yield and quality (Gowda et al., 2018), with production declines projected for several important crop species as temperatures rise (Zhao et al., 2017). In contrast, changes to growing season length can have both positive and negative effects on crop yield and prices (U.S. Environmental Protection Agency, 2016).

In marine systems, fish and invertebrate harvesting contributes \$212 billion in sales annually to the US economy (National Marine Fisheries Service, 2018), and climate change is affecting the availability, distribution (section 2.3), and quality of commercially important species (Kleisner et al., 2016; Peer and Miller, 2014; Pershing et al., 2015; Walsh et al., 2015). Rising ocean temperatures also decrease oxygen levels, which may reduce average fish body size by 14–24% by 2050 (Cheung et al., 2013). In freshwater systems, rising stream temperatures will negatively affect some harvested species (Crozier et al., 2019; Isaak et al., 2012). Future warming is expected to reduce the catch potential of all U.S. regions except the Arctic (Lam et al., 2016).

#### 4.2. Regulating services

Biodiversity and ecosystems provide important regulation services, such as sequestering carbon, moderating the impacts of extreme events (section 3.3), maintaining soil and air quality, and controlling disease spread.

Carbon storage is an important service that will become increasingly important as climate change accelerates. Although forest area has increased nationally since 2000 (EPA, 2017; Oswalt et al., 2014), it is unclear whether carbon storage from afforestation will continue to outweigh emissions from deforestation (Coulston et al., 2015). Moreover, carbon storage in many forests will likely decrease due to higher temperatures, increased water stress and disturbances (section 3.1), and lower rates of CO<sub>2</sub> uptake in aging forests when compared to regrowth forests after past disturbances (Oswalt et al., 2014; Pugh et al., 2019; USDA, 2016; Wear and Coulston, 2015). Coastal wetlands are highly productive ecosystems that store carbon (Davis et al., 2015; Howard et al., 2017), and also provide natural defenses against erosion, waves, flooding, and storm surge (Arkema et al., 2013). As human development or sea level rise degrade coastal wetlands, their capacity to provide these services diminishes.

Ecosystems also regulate the distribution, abundance, and life cycles of disease carrying organisms (Corvalan et al., 2005). Climate change is affecting the ability of ecosystems to provide this service as species ranges (section 2.3), abundances, and habitat conditions shift. For example, *Aedes* mosquitoes, which transmit diseases such as dengue, are expanding their geographic distribution in the southern U.S., increasing disease risk (Ebi and Nealon, 2016).

#### 4.3. Supporting services

Supporting ecosystem services facilitate basic ecosystem function, such as primary productivity (see section 3.1), nutrient cycling, and maintenance of genetic diversity. As temperatures rise, decomposition of soil organic matter generally increases, potentially increasing soil carbon losses and altering C:N balances (Davidson and Janssens, 2006). These changes are impacted by biotic interactions, including indirect changes to soil microbial community composition (Crowther et al., 2011).

The same activities responsible for climate change (e.g., fossil fuel combustion) result in increased nitrogen deposition, which has significant impacts on terrestrial and aquatic ecosystems, including through eutrophication (Galloway et al., 2008). The combination of higher nutrient loading and rising temperatures is increasing the frequency, duration, and extent of cyanobacteria responsible for harmful algal blooms, which can negatively impact human and animal health (Hilborn and Beasley, 2015). Estimates suggest that by the end of this century, conditions may allow blooms of the toxic alga *Alexandrium catenella* to be up to two months earlier and persist for up to an additional month (Moore et al., 2008; Sandifer and Sutton-Grier, 2014), with resulting impacts on aquaculture, recreation, and other activities.

#### 4.4. Cultural services

Cultural services are the non-material benefits that people gain from biodiversity and ecosystems, such as cultural identity, recreation, and mental and physical health. Despite their importance to human well-being, cultural services have been understudied compared to other ecosystem services (Runting et al., 2017). Indigenous peoples were among the earliest voices connecting culture and environmental change through science, scholarship, and other forms of expression (ACIA, 2004; Maynard, 1998).

There is growing evidence that human health benefits from exposure to natural ecosystems (Donatuto et al., 2014; Sandifer et al., 2015); conversely, climate-driven extremes such as increased temperatures and storms can decrease mental and physical health (Bell et al., 2016; Obradovich et al., 2018). Indirect economic costs (such as lost livelihoods) can also cause adverse socio-psychological impacts (Becker et al., 2015; Morris and Deterding, 2016; Shen and Aydin, 2014).

Although quantifying the value of ecosystems to cultural services is difficult, studies suggest that Americans place a high cultural value on natural systems: for example, a recent survey found that nearly 49 million adults nationwide participated in ocean and coastal recreation, spending more than 1.2 billion days along the coasts and over \$141 billion in ocean recreation-related goods and services (National Marine Fisheries Service, 2018). As climate change alters the ability of ecosystems to provide jobs, recreational opportunities, and restorative experiences, communities will experience declines in mental and physical health and potential losses of nature-based tourism dollars (Sandifer and Sutton-Grier, 2014).

### 5. Vulnerability of human communities

The adaptive capacity of human communities to deal with changes in ecosystem services will partly determine the magnitude of impacts on well-being. While some human communities have been proactive in identifying and planning for changes, others are more vulnerable due to a reduced ability to adapt.

Tribes and Indigenous peoples in the U.S. (groups whose exercise of self-determination as governing entities pre-dates the establishment of the U.S.; Jantarasami et al., 2018) have over 800 climate change initiatives and have led or participated in numerous climate change studies; many have developed their own climate change plans (Jantarasami et al., 2018). Since 1998, tribes and Indigenous peoples

**Table 1**  
Example management responses to changing biodiversity, ecosystems, and ecosystem services that can increase resilience, use updated technology or infrastructure, or be incorporated into governance approaches.

|                                   | Increasing ecosystem resilience                                                                                                                            | Utilizing technology and infrastructure                                                                                                                       | Improved governance                                                                                                                                                                                                                                                  |
|-----------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Individuals, populations, species | Maintaining population sizes, connectivity, and gene flow to allow for shifting ranges and increases in evolutionary resilience of populations and species | Matching individual genotypes with future environments under projected climate change increases adaptability of species                                       | Considering climate change impacts on threatened and endangered species in listing decisions can improve overall understanding of vulnerability                                                                                                                      |
| Ecosystems                        | Reduction of non-climate stressors, such as pollution and invasive species to minimize climate change impacts                                              | Adopting use of natural and nature-based infrastructure to improve resilience of natural communities, leveraging programs in place addressing other stressors | Implementing use of boundary organizations (e.g., North American Marine Protected Areas Network) can promote dialogue of diverse stakeholders in ecosystems that cross multiple jurisdictional boundaries and look for efficiencies in addressing multiple stressors |
| Ecosystem services                | Maintain biodiversity and ecological redundancy to minimize losses in valuable services                                                                    | Forecasting environmental conditions to prepare for economic changes in a particular industry (e.g., fishery)                                                 | Promoting consideration of ecosystem services and related climate impacts within federal planning and decision frameworks                                                                                                                                            |

have collaborated on the NCA, focusing on Indigenous concerns, knowledge of vulnerability, and goals for adaptation and mitigation.

Although tribes and Indigenous peoples continue to exercise self-governance, federal policies provide uneven levels of political engagement and support. Indigenous leadership is critical to addressing climate change; however, federal, state, and local governments pose barriers to tribal and Indigenous mitigation and adaptation efforts (Jantarasami et al., 2018). Historical and contemporary land-reduction, land-use restrictions, and poorly implemented treaty rights and consultation requirements exacerbate economic and health risks, which in many cases are associated with threats to Indigenous cultural maintenance. For example, climate-driven range shifts in culturally important species pose challenges to tribes and Indigenous peoples when tribal land areas are small and have limited connectivity (Rapp et al., 2019). Indigenous peoples face risks related to resettlement due to the impacts of climate change, such as coastal erosion and sea ice loss in Alaska. Despite a history of forced relocations, there are structural barriers for Indigenous peoples to participate in current policy processes trying to plan for climate-driven resettlement (Jantarasami et al., 2018).

Across the U.S., coastal communities are also particularly vulnerable to climate change impacts from rising sea levels and more intense storms, which are exacerbating high tide and storm surges, erosion, and saltwater intrusion. High tide flooding is already forcing some cities to install costly pump stations to clear floodwaters and mobilize emergency responders to routinely close flooded streets.

Society's most vulnerable populations, including children, the elderly, economically disadvantaged, homeless, and those with preexisting mental illness tend to be the most heavily impacted by climate changes and resulting impacts to ecosystem services (Dodgen et al., 2016).

## 6. Implications for natural resource management

Natural resource management traditionally focuses on maintaining or restoring to historical conditions (e.g., National Park System Advisory Board, 2012). While historical context may still motivate management decisions, restoring to historical baselines may not be realistic as the climate changes (Stein et al., 2014). In some cases, management practices to resist change may be effective; in others, managers may choose to accept ecosystem changes or to alter management practices to direct changes in order to minimize loss of valued species and services as ecosystems transform (Aplet and Mckinley, 2017; Millar and Stephenson, 2015; Stein et al., 2013).

Adaptive and proactive approaches that are continually updated to reflect emerging and anticipated climate change impacts will be needed (Bradford et al., 2018; Holsman et al., 2019; Stein et al., 2014; Table 1). For example, the U.S. has a rigorous, science-based system for detecting changes in fish abundance, productivity, and catch, which informs fishery management decisions such as seasonal and spatial closures, annual

quotas, and stock rebuilding plans (Pinsky and Mantua, 2014). Collection of this type of information is important for future assessment and updating of management objectives (Stein et al., 2014). NOAA Fisheries has developed adaptation strategies that incorporate climate and ecosystem-related factors into fishery decision-making (Busch et al., 2016; Hare et al., 2016a; Link et al., 2015). Decision support tools, including scenario planning (Cobb and Thompson, 2012; Mahmoud et al., 2009; Peterson et al., 2003) and structured decision-making (Gregory et al., 2012) can help decision-makers explore broad scenarios of risk and develop and prioritize actions that account for uncertainty, optimize tradeoffs, and reflect institutional capacity. Below, we discuss strategies for increasing resilience of ecosystems and human communities.

### 6.1. Increasing ecosystem resilience

To create effective adaptation strategies, managers need to understand which species are most at risk and why. One way to determine relative risk is through climate change vulnerability assessments that examine species exposure, sensitivity, and adaptive capacity to climate change, and exposure to non-climate stressors (Glick et al., 2011; Hare et al., 2016b; Spencer et al., 2019; Staudinger et al., 2015). Managers can then take proactive steps to increase resilience (Table 1).

Systems that are already degraded from non-climate stressors have lower resilience; therefore, restoring and conserving areas that support valued resources are important. Many strategies for reducing other stressors are things that managers already know, such as restoring populations and habitats, increasing connectivity, and reducing stress from disease, pollution and invasive species (Box 2). For example, prescribed burning and reducing forest stand density can lower wildfire risk in some forest types and can increase resistance and resilience to drought and insect outbreaks (Bottero et al., 2017; Sohn et al., 2016; Vernon et al., 2018). Similarly, in aquatic systems, effective strategies include reconnecting floodplains and side channels, ensuring effective passage for aquatic organisms, and maintaining large trees in forested riparian areas for shade and recruitment to streams (Peterson and Halofsky, 2018; Pollock et al., 2014).

Limiting invasive species spread can help maintain biodiversity, ecosystem function, and resilience (Fischer et al., 2006; Katsanevakis et al., 2014; Oliver et al., 2015). Dialogue between managers, scientists, and policymakers can help ensure that climate change mitigation does not exacerbate invasive species spread (Beaury et al., 2019). Similarly, invasive species policies that explicitly address climate change will enable proactive management (Pyke et al., 2008). Managers may also benefit from considering potential negative effects from native species range expansions (Burgiel et al., 2014; Carey et al., 2012; Giakoumi et al., 2019; Wallingford et al., in review).

Restoring habitat and increasing connectivity to enable species to disperse across the landscape and follow physiological niches is another

## Box 2

## Examples of on-the-ground adaptation.

**Restoring meadows in the Sierra Nevada**

Photo: Toni Lyn Morelli

The Sierra Nevada region is critical to the California water supply.

Snow in the mountains melts slowly during spring and summer, providing water for ecosystems and people. Climate change is projected to cause more winter precipitation to fall as rain and earlier snowmelt, leading to decreased summer water flows (Viers et al., 2013). Increased frequency and severity of floods is also anticipated. Well-functioning mountain meadows can attenuate floods and increase groundwater storage (National Wildlife Federation, 2010), and some have been identified as climate change refugia for wildlife (Morelli et al., 2017). However, historic land use changes have degraded approximately 40–60% of Sierra meadows. In 2015, Federal, state, and NGO partners restored four meadows with high ecological value located in areas projected to experience the most significant changes in hydrology (Fair and Hunt, 2015; National Wildlife Federation, 2010).

**Nature-based infrastructure**

Photo: Linda Walters

The use of nature-based designs for coastal resilience structures like living shorelines tends to be more effective at withstanding extreme events (Powell et al., 2019). For example, during Tropical Storm Irene in 2011, stream-designed road crossings survived a category 3 hurricane while nearly 1000 traditional culverts were damaged or destroyed (Sutton-Grier et al., 2018). In another example, living shorelines fared much better under a category 1 hurricane in North Carolina than traditional bulk heads (Gittman et al., 2014).

key climate adaptation action (Anderson et al., 2015; McGuire et al., 2016; Timpone-Padgham et al., 2017). For example, the recent recovery plan for Atlantic salmon (*Salmo salar*) included identifying vacant habitats, creating redundant populations, and revisiting critical habitat designations to ensure sufficient climate resilient habitats under future conditions (USFWS and NMFS, 2019). Additionally, conserving climate change refugia (areas relatively buffered from contemporary climate change that enable persistence of valued resources), has become a focus of conservation efforts for highly valued vulnerable ecosystems and species (Keppel et al., 2015; Morelli et al., 2016).

Restoring populations can be an effective way to increase genetic diversity and potential for species to evolve and adapt to changing climatic conditions (Sgrò et al., 2011). More active approaches such as diverse seed sourcing, translocation of genes or individuals to accelerate evolution, and assisted migration may be warranted for species with limited dispersal ability or that face movement barriers (Anderson et al., 2014; Isaac-Renton et al., 2014; Whiteley et al., 2015). Unforeseen and unwanted consequences are possible for any assisted migration, but developing policies to analyze and manage potential consequences could minimize unintended outcomes (Invasive Species Advisory Committee, 2017; Schwartz et al., 2012).

## 6.2. Leveraging technology and infrastructure

Human communities may need to adapt by updating technology and infrastructure (Table 1). Extreme events often motivate adaptation. For example, the inability of the lobster supply chain to handle the sudden influx of soft-shell lobster during the 2012 heatwave (Mills et al., 2013) led to increased domestic processing capacity and expanded marketing. These adaptations allowed the fishery to achieve record value during a subsequent heatwave in 2016 (Pershing et al., 2018). To-date, most documented cases of adaptation to climate impacts on the ocean have been reactive. However, it is now possible to forecast

temperature, pH, and oxygen conditions several months in advance (Jacox et al., 2017; Siedlecki et al., 2016; Tommasi et al., 2017). Ensuring that these data products are distributed and used effectively will require considerable engagement with the user community (Hobday et al., 2019; Siedlecki et al., 2016).

In coastal ecosystems, there is growing interest in natural and nature-based infrastructure (NNBI) to increase coastal community resilience (Powell et al., 2019; Spalding et al., 2014; Sutton-Grier et al., 2015). NNBI strategies include restoring or creating coastal ecosystems like salt marsh, mangroves, oyster or coral reefs, beaches, and dunes to mitigate waves and erosion (Ferrario et al., 2014; Möller et al., 2014; Powell et al., 2019; Rodriguez et al., 2014; Zhang et al., 2012), but also include hybrid combinations of natural and built infrastructure such as living shorelines or using combinations of habitat restoration and flood walls (Box 2) (Gittman et al., 2014; Sutton-Grier et al., 2015). NNBI approaches provide co-benefits in terms of habitat, water quality, and recreation, and can be cheaper and survive extreme events better than traditional infrastructure (Gittman et al., 2014; Powell et al., 2019; Sutton-Grier et al., 2018). Although long-term planning may require some coastal communities to relocate due to prohibitively high cost or infeasibility of sea level rise protection, NNBI approaches can at least mitigate some short-term impacts and allow communities more time to consider options. NNBI approaches are often not as well known or trusted in comparison to traditional grey (e.g., seawall) approaches; therefore it is important to increase information on the performance of these techniques, communicate their ecosystem service benefits, and increase coordination and planning around shared socio-ecological goals (Powell et al., 2019).

## 6.3. Strengthening governance

Federal agencies that manage natural resources are increasingly considering climate change impacts in their management plans (e.g.

Busch et al., 2016; Link, 2016; National Fish Wildlife and Plants Climate Adaptation Partnership, 2012; National Park Service, 2013; Swanston and Janowiak, 2012; Table 1). For example, the National Marine Fisheries Service has developed guidance on how climate change information should be considered in Endangered Species Act (ESA) decisions (National Marine Fisheries Service, 2016). The U.S. Fish and Wildlife Service has also considered climate change in listing decisions, biological opinions, and proposed alternative actions under the ESA (e.g., U.S. Fish and Wildlife Service, 2008, 2010), although climate change is still


not included for many species. Even when climate change has been listed as a threat, specific management actions are often lacking (Delach et al., 2019). Federal agencies have also been directed to promote consideration of ecosystem services and related climate impacts within existing planning and decision frameworks (Executive Office of the President of the United States, 2015).

Despite progress, institutional barriers such as a focus on near-term planning, fixed policies and protocols, jurisdictional restrictions, and an established practice of managing based on historical conditions remain

### Box 3

An integrated case study: climate change impacts on salmon across the U.S.

#### Individual, population, and species level responses



Bureau of Land Management



Photo: National Park Service

#### Ecosystem level responses



#### Impacts to ecosystem services



Matt Nagle, Puyallup Tribal News

#### Implications for management



(Pictorial representation of scenarios used in Borggaard et al., 2019)

- In the Northwest, abnormally warm temperatures have led to losses of migrating and spawning salmon in the Columbia River (NOAA Fisheries, 2016).
- In Alaska, some salmon populations have benefited from warmer temperatures, earlier spring, and increased density of zooplankton prey (Schindler et al., 2005).
- Coho salmon on the west coast of the U.S. are expected to shift their range north by 2050 (Cheung et al., 2015). Under a high greenhouse gas emissions scenario, projected stream temperature increases could lead to a 22% reduction in salmon habitat in Washington by late century (Niemi et al., 2009).
- Sockeye salmon (*Oncorhynchus nerka*) in the Columbia River have been migrating earlier. Nearly two-thirds of this response was explained by evolutionary rather than plastic responses (Crozier et al., 2011).
- In the Penobscot River in the Gulf of Maine, adult Atlantic salmon (*Salmo salar*) have arrived as much as 2–3 weeks earlier to their freshwater spawning grounds between the late 1970s and early 2000s (Huntington et al., 2003; Juanes et al., 2004).

- Kodiak brown bears (*Ursus arctos middendorffi*) have responded to phenological shifts in sockeye salmon and red elderberry (*Sambucus racemosa*). During years with warmer springs, elderberry fruited earlier and overlapped with the salmon spawning run, causing bears to eat less salmon and more berries (Deacy et al., 2017).
- In the Pacific Northwest, warming stream temperatures are likely to reduce the amount of rearing habitat for Chinook salmon (*Oncorhynchus tshawytscha*), but will likely increase the range of invasive smallmouth bass (*Micropterus dolomieu*) (Lawrence et al., 2014).

- As Kodiak brown bear diet shifted away from salmon (see panel 2), this reduced the distribution of marine derived nutrients being distributed into the surrounding terrestrial landscape, with implications for ecological function (Deacy et al., 2017).
- Shifts in availability and declines of salmon and other cold-water fishes impact cherished cultural resources, particularly in the Pacific Northwest where they are important to tribal nations such as the Nez Perce Tribe as cultural, subsistence, and economic resources. Tribes view salmon as an extension of life and an indicator of environmental health, and loss of salmon is equated with loss of tribal identity and culture (Colombi, 2012).

- Atlantic salmon and many distinct Pacific salmon population groups are federally protected species. Climate change is expected to exacerbate pre-existing anthropogenic stressors, which have reduced adaptive capacity (Crozier et al., 2019; Hare et al., 2016b). The Pacific Salmon Climate Vulnerability Assessment identified life stages that are most vulnerable to climate change and established a methodology for future status reviews to monitor and update recovery needs and prioritize recovery actions (Crozier et al., 2019).
- In the Yakima Basin, water resource managers, conservation groups, and state and federal agencies are enhancing reservoir storage capacity to ensure minimum flows for fish, allow passage above dams for access to cool, historical habitat, restore a more natural hydrograph, and improve riparian conditions for natural temperature and flow stabilization to bolster salmon runs and riparian habitat during droughts. Actions to redistribute water also impacted farmers, but agreements were made to sustain conditions needed to support salmon (NOAA Fisheries, 2015).
- NOAA used Atlantic salmon as a pilot case study for Scenario Planning - a structured process that embraces uncertainty and explores plausible alternative future conditions to help manage risk and prioritize actions - to explore actions to increase resilience to changing conditions in riverine and marine environments in response to declining population trends (Borggaard et al., 2019). Some outcomes from this effort have already been incorporated into the most recent revision of the Atlantic Salmon Recovery Plan (USFWS and NMFS, 2019).

a challenge (Kemp et al., 2015; Stein et al., 2014). Even with agency-level directives for climate adaptation, implementing actions on the ground can be difficult due to lack of funding and time, negative public perceptions, and difficulty transferring science between researchers and managers (Kemp et al., 2015). Boundary organizations that work at the interface of research and management, such as the U.S. Geological Survey Climate Adaptation Science Centers and the NOAA Regional Integrated Sciences and Assessment Programs, can help address some of these issues by bringing together multiple stakeholders and working with managers to develop adaptation plans.

Promoting local climate adaptation initiatives can also be an effective management strategy. Self-determination and self-governance of tribes and Indigenous peoples, as well as involvement of communities of color or natural resource users such as anglers in particular localities, supports the use of local knowledge and cultural practices that can track, adapt to and mitigate environmental change (Maldonado et al., 2013; Norton-Smith et al., 2016; Vinyeta et al., 2015). For many tribes and Indigenous peoples, their political rights are connected to cultural responsibilities to live sustainably (Borrows, 1997; De Chavez and Tauli-Corpuz, 2009). These cultural responsibilities often involve rich scientific traditions of observation and ecosystem stewardship (Shilling and Nelson, 2018; Trosper, 2009). Tribal and Indigenous governments have emphasized the importance of Indigenous scientists and knowledge keepers working collaboratively to observe changes and create adaptation strategies (Grossman and Parker, 2012; Houser et al., 2001). These governments are in unique positions to cooperate with local, state, and federal government to coordinate regional conservation and adaptation efforts that emphasize strategic foresight and sustainability (Krakoff, 2008; Morishima, 2014; Whyte et al., 2014).

## 7. Conclusion

Climate change is a pervasive and growing threat to biodiversity, ecosystems, and ecosystem services in the U.S. Climate impacts have been and will continue to be observed at the level of individuals, populations, and species through changes in behavior and morphology, phenology, and range shifts, and at the ecosystem level through changes in primary production, species interactions and emergent properties, and extreme events. Ecosystems and biodiversity underpin important services to people, thus these changes impact provisioning, regulating, supporting, and cultural services, with implications for human well-being. Effective management will require flexible, proactive approaches that account for potential climate change impacts (Box 3). Managers are beginning to implement these strategies, but face challenges due to lack of information and institutional barriers. Widespread incorporation of climate change into natural resource management is yet to be achieved, but examples are emerging that help increase awareness and provide case studies in different sectors. Moving forward, evaluations of effectiveness and demonstrative case studies of adaptation success stories are needed to promote and guide climate-smart management.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

We thank the many authors of the original NCA chapters and the USGCRP team, as well as J. Dunham for his helpful comments. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. government.

## References

- Abatzoglou, J.T., Kolden, C.A., 2013. Relationships between climate and macroscale area burned in the western United States. *Int. J. Wildl. Fire* 22, 1003–1020. <https://doi.org/10.1017/WFI3019>.
- Abatzoglou, J.T., Williams, A.P., 2016. Impact of anthropogenic climate change on wildfire across western US forests. *Proc. Natl. Acad. Sci.* 113, 11770–11775. <https://doi.org/10.1073/pnas.1607171113>.
- Abbott, K.C., Harmon, J.P., Fabina, N.S., 2014. The challenge of predicting temperature effects on short-term predator–prey dynamics. *Popul. Ecol.* 56, 375–392.
- Abt, K.L., Butry, D.T., Prestemon, J.P., Scranton, S., 2015. Effect of fire prevention programs on accidental and incendiary wildfires on tribal lands in the United States. *Int. J. Wildl. Fire* 24, 749–762. <https://doi.org/10.1017/WFI14168>.
- ACIA, 2004. *Impacts of a Warming Arctic–Arctic Climate Impact Assessment*. Cambridge University Press.
- Alabia, I.D., García Molinos, J., Saitoh, S.I., Hirawake, T., Hirata, T., Mueter, F.J., 2018. Distribution shifts of marine taxa in the Pacific Arctic under contemporary climate changes. *Divers. Distrib.* 24, 1583–1597. <https://doi.org/10.1111/ddi.12788>.
- Alexander, J.M., Diez, J.M., Hart, S.P., Levine, J.M., 2016. When climate reshuffles competitors: a call for experimental macroecology. *Trends Ecol. Evol.* 31, 831–841. <https://doi.org/10.1016/j.tree.2016.08.003>.
- Alexander, J.M., Chalmers, L., Lenoir, J., Burgess, T.J., Essl, F., Haider, S., Kueffer, C., McDougall, K., Milbau, A., Nuñez, M.A., Pauchard, A., Rabitsch, W., Rew, L.J., Sanders, N.J., Pellissier, L., 2018. Lags in the response of mountain plant communities to climate change. *Glob. Chang. Biol.* 24, 563–579. <https://doi.org/10.1111/gcb.13976>.
- Anderegg, W.R.L., Hicke, J.A., Fisher, R.A., Allen, C.D., Aukema, J., Bentz, B.J., Hood, S., Lichstein, J.W., Macalady, A.K., McDowell, N.G., Pan, Y., Raffa, K., Sala, A., Shaw, J.D., Stephenson, N.L., Tague, C., Zeppel, M., 2015. Tree mortality from drought, insects, and their interactions in a changing climate. *New Phytol.* 208, 674–683. <https://doi.org/10.1111/nph.13477>.
- Anderson, J.H., Pess, G.R., Carmichael, R.W., Ford, M.J., Cooney, T.D., Baldwin, C.M., McClure, M.M., 2014. Planning Pacific salmon and steelhead reintroductions aimed at long-term viability and recovery. *North Am. J. Fish. Manag.* 34, 72–93. <https://doi.org/10.1080/02755947.2013.847875>.
- Anderson, M.G., Clark, M., Mcrae, B.H., 2015. *Permeable Landscapes for Climate Change the Nature Conservancy*. Eastern Conservation Science, Boston, Massachusetts.
- Aplet, G.H., McKinley, P.S., 2017. A portfolio approach to managing ecological risks of global change. *Ecosyst. Heal. Sustain.* 3, 1–15. <https://doi.org/10.1002/ehs2.1261>.
- Arkema, K.K., Guannel, G., Verutes, G., Wood, S.A., Guerry, A., Ruckelshaus, M., Kareiva, P., Lacayo, M., Silver, J.M., 2013. Coastal habitats shield people and property from sea-level rise and storms. *Nat. Clim. Chang.* 3, 913–918. <https://doi.org/10.1038/nclimate1944>.
- Asch, R.G., Stock, C.A., Sarmiento, J.L., 2019. Climate change impacts on mismatches between phytoplankton blooms and fish spawning phenology. *Glob. Chang. Biol.* 25, 2544–2559. <https://doi.org/10.1111/gcb.14650>.
- Asner, G.P., Brodrick, P.G., Anderson, C.B., Vaughn, N., Knapp, D.E., Martin, R.E., 2016. Progressive forest canopy water loss during the 2012–2015 California drought. *Proc. Natl. Acad. Sci. U. S. A.* 113, E249–E255. <https://doi.org/10.1073/pnas.1523397113>.
- Atkinson, D., 1994. Temperature and organism size – a biological law for ectotherms. *Adv. Ecol. Res.* 25, 1–58.
- Ault, T.R., Schwartz, M.D., Zurita-Milla, R., Weltzin, J.F., Betancourt, J.L., 2015. Trends and natural variability of spring onset in the coterminous United States as evaluated by a new gridded dataset of spring indices. *J. Clim.* 28, 8363–8378. <https://doi.org/10.1175/JCLI-D-14-00736.1>.
- Banta, J.A., Richards, C.L., 2018. Quantitative epigenetics and evolution. *Heredity* (Edinb) 121, 210–224. <https://doi.org/10.1038/s41437-018-0114-x>.
- Barbero, R., Abatzoglou, J.T., Larkin, N.K., Kolden, C.A., Stocks, B., 2015. Climate change presents increased potential for very large fires in the contiguous United States. *Int. J. Wildl. Fire* 24, 892–899. <https://doi.org/10.1017/WFI15083>.
- Barnett, T.P., Pierce, D.W., Hidalgo, H.G., Bonfils, C., Santer, B.D., Das, T., Bala, G., Wood, A.W., Nozawa, T., Mirin, A. a, Cayan, D.R., Dettinger, M.D., 2008. Human-induced changes United States. *Science* (80-. ) 319, 1080–1083. <https://doi.org/10.1126/science.1152538>.
- Bay, R.A., Harrigan, R.J., Buermann, W., Le Underwood, V., Gibbs, H.L., Smith, T.B., Ruegg, K., 2018. Response to comment on “Genomic signals of selection predict climate-driven population declines in a migratory bird.”. *Science* (80-. ) 361, 83–86. <https://doi.org/10.1126/science.aat7956>.
- Beaury, E.M., Fusco, E.J., Jackson, M.R., Laginhas, B.B., Morelli, T.L., Allen, J.M., Pasquarella, V.J., Bradley, B.A., 2019. Incorporating climate change into invasive species management: insights from managers. *Biol. Invasions* 6. <https://doi.org/10.1007/s10530-019-02087-6>.
- Becker, A.H., Matson, P., Fischer, M., Mastrandrea, M., 2015. Towards seaport resilience for climate change adaptation: stakeholder perceptions of hurricane impacts in Gulfport (MS) and Providence (RI). *Prog. Plann.* 99, 1–49.
- Beever, E.A., Leary, J.O., Mengelt, C., West, J.M., Julius, S., Green, N., Magness, D., Petes, L., Stein, B., Nicotra, A.B., Hellmann, J.J., 2015. Improving Conservation Outcomes with a New Paradigm for Understanding Species' Fundamental and Realized Adaptive Capacity 00. , pp. 1–7. <https://doi.org/10.1111/conl.12190>.
- Beever, E.A., O'Leary, J., Mengelt, C., West, J.M., Julius, S., Green, N., Magness, D., Petes, L., Stein, B., Nicotra, A.B., Hellmann, J.J., Robertson, A.L., Staudinger, M.D., Rosenberg, A.A., Babij, E., Brennan, J., Schuurman, G.W., Hofmann, G.E., 2016. Improving conservation outcomes with a new paradigm for understanding species' fundamental and realized adaptive capacity. *Conserv. Lett.* 9, 131–137. <https://doi.org/10.1111/conl.12190>.

- Beever, E.A., Hall, L.E., Varner, J., Loosen, A.E., Dunham, J.B., Gahl, M.K., Smith, F.A., Lawler, J.J., 2017. Behavioral flexibility as a mechanism for coping with climate change. *Front. Ecol. Environ.* 15, 299–308. <https://doi.org/10.1002/fee.1502>.
- Bell, G., 2013. Evolutionary rescue and the limits of adaptation. *Philos. Trans. R. Soc. B Biol. Sci.* 368, 1–6. <https://doi.org/10.1098/rstb.2012.0080>.
- Bell, J.E., Herring, S.C., Jantarasami, L., Adrianopoli, C., Benedict, K., Conlon, K., Escobar, V., Hess, J., Luwall, J., Garcia-Pando, C.P., Quattrocchi, D., Runkle, J., II Schreck, C.J., 2016. *Ch. 4: impacts of extreme events on human health. The Impacts of Climate Change on Human Health in the United States: A Scientific Assessment*. U.S. Global Change Research Program, Washington, D.C., USA.
- Berg, M.P., Toby Kiers, E., Driessen, G., van der Heijden, M., Kooi, B.W., Kuenen, F., Liefing, M., Verhoef, H.A., Ellers, J., 2010. Adapt or disperse: understanding species persistence in a changing world. *Glob. Chang. Biol.* 16, 587–598. <https://doi.org/10.1111/j.1365-2486.2009.02014.x>.
- Blois, J.L., Zarnetske, P.L., Fitzpatrick, M.C., Finnegan, S., 2013. Climate change and the past, present, and future of biotic interactions. *Science* (80-. ) 341, 499–504. <https://doi.org/10.1126/science.1237184>.
- Bonamour, S., Chevin, L.M., Charmanter, A., Teplitsky, C., 2019. Phenotypic plasticity in response to climate change: the importance of cue variation. *Philos. Trans. R. Soc. B Biol. Sci.* 374. <https://doi.org/10.1098/rstb.2018.0178>.
- Bond, N.A., Cronin, M.F., Freeland, H., Mantua, N., 2015. Causes and impacts of the 2014 warm anomaly in the NE Pacific. *Geophys. Res. Lett.* 42, 3414–3420. <https://doi.org/10.1002/2015GL063306>.
- Borggaard, D., Dick, D., Star, J., Alexander, M., Bernier, M., Collins, M., Damon-Randall, K., Dudley, R., Griffiths, R., Hayes, S., Johnson, M., Kircheis, D., Kocik, J., Letcher, B., Mantua, N., Morrison, W., Nislow, K., Saba, V., Saunders, R., Sheehan, T., Staudinger, M., 2019. *Atlantic salmon (Salmo salar) climate scenario planning pilot report*. Greater Atlantic Region Policy Series [19-05].
- Borrows, J., 1997. *Living between water and rocks: first nations, environmental planning and democracy*. Univ. Tor. Law J. 47, 417–468.
- Bottero, A., D'Amato, A.W., Palik, B.J., Bradford, J.B., Fraver, S., Battaglia, M.A., Asherin, L.A., 2017. Density-dependent vulnerability of forest ecosystems to drought. *J. Appl. Ecol.* 54, 1605–1614. <https://doi.org/10.1111/1365-2664.12847>.
- Boutin, S., Lane, J.E., 2014. Climate change and mammals: evolutionary versus plastic responses. *Evol. Appl.* 7, 29–41. <https://doi.org/10.1111/eva.12121>.
- Bradford, J.B., Betancourt, J.L., Butterfield, B.J., Munson, S.M., Wood, T.E., 2018. Anticipatory natural resource science and management for a changing future. *Front. Ecol. Environ.* 16, 295–303. <https://doi.org/10.1002/fee.1806>.
- Bradley, B.A., Laginhas, B.B., Whitlock, R., Allen, J.M., Bates, A.E., Bernatchez, G., Diez, J.M., Early, R., Lenoir, J., Vilà, M., Sorte, C.J.B., 2019. Disentangling the abundance-impact relationship for invasive species. *Proc. Natl. Acad. Sci. U. S. A.* 116, 9919–9924. <https://doi.org/10.1073/pnas.1818081116>.
- Bradshaw, W.E., Holzapfel, C.M., 2007. Evolution of animal photoperiodism. *Annu. Rev. Ecol. Syst.* 38, 1–25. <https://doi.org/10.1146/annurev.ecolsys.37.091305.110115>.
- Breeggemann, J.J., Kaemingk, M.A., DeBates, T.J., Paukert, C.P., Krause, J.R., Letvin, A.P., Stevens, T.M., Willis, D.W., Chippis, S.R., 2016. Potential direct and indirect effects of climate change on a shallow natural lake fish assemblage. *Ecol. Freshw. Fish* 25, 487–499. <https://doi.org/10.1111/eff.12248>.
- Burgiel, S.W., Hall, T., Adams, N., Anderson, K., Bella, E., Bierwagen, B., Boroja, M., Brusati, E., Burgett, J., Carlisle, L., Cleary, R., Cleland, D., Darling, J., Draheim, R., Forest, D., Grise-Stahlman, S., Hart, S., Hatch, K., Heimowitz, P., Hoffman, J., Holcombe, T., Johnson, D., Lassuy, D., McMartin, L., Moy, P., Muir, A., Nelson, R., Pederson, J., Plumb, G., Quinn, J., Resnick, J., Savignano, D., Schmal, N., Shaw, L., Stein, B., Winter, G., Woodson, D., Wooten, D., Wullschlegel, J., 2014. *Bioinvasions in a Changing World : A Resource on Invasive Species-climate Change Interactions for Conservation and Natural Resource Management*. The Aquatic Nuisance Species Task Force (ANSTF) and The National Invasive Species Council (NIS).
- Burrows, M.T., Schoeman, D.S., Buckley, L.B., Moore, P., Poloczanska, E.S., Brander, K.M., Brown, C., Bruno, J.F., Duarte, C.M., Halpern, B.S., Holding, J., Kappel, C.V., Kiessling, W., O'Connor, M.L., Pandolfi, J.M., Parmesan, C., Schwing, F.B., Sydeman, W.J., Richardson, A.J., 2011. The pace of shifting climate in marine and terrestrial ecosystems. *Science* (80-. ) 334, 652–655. <https://doi.org/10.1126/science.1210288>.
- Busch, D.S., Griffiths, R., Link, J., Abrams, K., Baker, J., Brainard, R.E., Ford, M., Hare, J.A., Himes-Cornell, A., Hollowed, A., Mantua, N.J., McClatchie, S., McClure, M., Nelson, M.W., Osgood, K., Peterson, J.O., Rust, M., Saba, V., Sigler, M.F., Sykora-Bodie, S., Toole, C., Thunberg, E., Waples, R.S., Merrick, R., 2016. Climate science strategy of the US National Marine Fisheries Service. *Mar. Policy* 74, 58–67. <https://doi.org/10.1016/j.marpol.2016.09.001>.
- Butry, D.T., Prestemon, J.P., Abt, K.L., Sutphen, R., 2010. Economic optimisation of wildfire intervention activities. *Int. J. Wildl. Fire* 19, 659–672. <https://doi.org/10.1071/WF09090>.
- Cal Fire, 2018. *Over 129 Million Dead Trees in California Between 2010–2017 [Story Map] [WWW Document]*.
- Campbell, E.Y., Dunham, J.B., Reeves, G.H., Wondzell, S.M., 2019. Phenology of hatching, emergence, and end-of-season body size in young-of-year coho salmon in thermally contrasting streams draining the Copper River Delta, Alaska. *Can. J. Fish. Aquat. Sci.* 76, 185–191. <https://doi.org/10.1139/cjfas-2018-0003>.
- Campbell, J.E., Berry, J.A., Seibt, U., Smith, S.J., Montzka, S.A., Launois, T., Belviso, S., Bopp, L., Laine, M., 2017. Large historical growth in global terrestrial gross primary production. *Nature* 544, 84–87. <https://doi.org/10.1038/nature22030>.
- Campbell, J.L., Rustad, L.E., Boyer, E.W., Christopher, S.F., Driscoll, C.T., Fernandez, I.J., Groffman, P.M., Houle, D., Kiebusch, J., Magill, A.H., Mitchell, M.J., Ollinger, S.V., 2009. Consequences of climate change for biogeochemical cycling in forests of northeastern North America. *Can. J. For. Res.* 39, 264–284. <https://doi.org/10.1139/X08-104>.
- Cannizzo, Z.J., Griffen, B.D., 2016. Changes in spatial behaviour patterns by mangrove tree crabs following climate-induced range shift into novel habitat. *Anim. Behav.* 121, 79–86. <https://doi.org/10.1016/j.anbehav.2016.08.025>.
- Carey, M.P., Sanderson, B.L., Barnas, K.A., Olden, J.D., 2012. Native invaders - challenges for science, management, policy, and society. *Front. Ecol. Environ.* 10, 373–381. <https://doi.org/10.1890/110060>.
- Carlson, S.M., Cunningham, C.J., Westley, P.A.H., 2014. Evolutionary rescue in a changing world. *Trends Ecol. Evol.* 29, 521–530. <https://doi.org/10.1016/j.tree.2014.06.005>.
- Charmanter, A., Gienapp, P., 2014. Climate change and timing of avian breeding and migration: evolutionary versus plastic changes. *Evol. Appl.* 7, 15–28. <https://doi.org/10.1111/eva.12126>.
- Chen, I.-C., Hill, J.K., Ohlemüller, R., Roy, D.B., Thomas, C.D., 2011. Rapid range shifts of species of climate warming. *Science* (80-. ) 333, 1024–1026.
- Cheung, W.W.L., Sarmiento, J.L., Dunne, J., Frölicher, T.L., Lam, V.W.Y., Palomares, M.L.D., Watson, R., Pauly, D., 2013. Shrinking of fishes exacerbates impacts of global ocean changes on marine ecosystems. *Nat. Clim. Chang.* 3, 254–258. <https://doi.org/10.1038/nclimate1691>.
- Cheung, W.W.L., Brodeur, R.D., Okey, T.A., Pauly, D., 2015. Projecting future changes in distributions of pelagic fish species of Northeast Pacific shelf seas. *Prog. Oceanogr.* 130, 19–31. <https://doi.org/10.1016/j.pocean.2014.09.003>.
- Chmura, H.E., Kharouba, H.M., Ashander, J., Ehman, S.M., Rivest, E.B., Yang, L.H., 2019. The mechanisms of phenology: the patterns and processes of phenological shifts. *Ecol. Monogr.* 89, 1–22. <https://doi.org/10.1002/ecm.1337>.
- Choat, B., Jansen, S., Brodribb, T.J., Cochard, H., Delzon, S., Bhaskar, R., Bucci, S.J., Feild, T.S., Gleason, S.M., Hacke, U.G., Jacobsen, A.L., Lens, F., Maherali, H., Martínez-Vilalta, J., Mayr, S., Mencuccini, M., Mitchell, P.J., Nardini, A., Pittermann, J., Pratt, R.B., Sperry, J.S., Westoby, M., Wright, I.J., Zanne, A.E., 2012. Global convergence in the vulnerability of forests to drought. *Nature* 491, 752–755. <https://doi.org/10.1038/nature11688>.
- Chust, G., Allen, J.L., Bopp, L., Schrum, C., Holt, J., Tsiaras, K., Zavatarelli, M., Chifflet, M., Cannaby, H., Dadou, I., Daewel, U., Wakelin, S.L., Machu, E., Pushpadas, D., Butenschon, M., Artioli, Y., Petihakis, G., Smith, C., Garçon, V., Goubanova, K., Le Vu, B., Fach, B.A., Salihoğlu, B., Clementi, E., Irigoien, X., 2014. Biomass changes and trophic amplification of plankton in a warmer ocean. *Glob. Chang. Biol.* 20, 2124–2139. <https://doi.org/10.1111/gcb.12562>.
- Clark, J.S., Iverson, L., Woodall, C.W., Allen, C.D., Bell, D.M., Bragg, D.C., D'Amato, A.W., Davis, F.W., Hersh, M.H., Ibanez, I., Jackson, S.T., Matthews, S., Pederson, N., Peters, M., Schwartz, M.W., Waring, K.M., Zimmermann, N.E., 2016. The impacts of increasing drought on forest dynamics, structure, and biodiversity in the United States. *Glob. Chang. Biol.* 22, 2329–2352. <https://doi.org/10.1111/gcb.13160>.
- Cobb, A.N., Thompson, J.L., 2012. Climate change scenario planning: a model for the integration of science and management in environmental decision-making. *Environ. Model. Softw.* 38, 296–305. <https://doi.org/10.1016/j.envsoft.2012.06.012>.
- Cohen, J.M., Lajeunesse, M.J., Rohr, J.R., 2018. A global synthesis of animal phenological responses to climate change. *Nat. Clim. Chang.* 8, 224–228. <https://doi.org/10.1038/s41558-018-0067-3>.
- Colombi, B.J., 2012. Salmon and the adaptive capacity of Nimiipuu (Nez Perce) culture to cope with change. *Am. Indian Q.* 36, 75. <https://doi.org/10.5250/amerindian.36.1.0075>.
- Conover, D.O., Duffy, T.A., Hice, L.A., 2009. The covariance between genetic and environmental influences across ecological gradients: reassessing the evolutionary significance of countergradient and cogenetic variation. *Ann. N. Y. Acad. Sci.* <https://doi.org/10.1111/j.1749-6632.2009.04575.x>.
- Corvalan, C., Hales, S., McMichael, A., Butler, C., Campbell-Lendrum, D., Confalonieri, U., Leitner, K., Lewis, N., Patz, J., Polson, K., Scheraga, J., Woodward, A., Younes, Maged, 2005. *Ecosystems and Human Well-being: Health Synthesis, a Report of the Millennium Ecosystem Assessment*.
- Coulston, J.W., Wear, D.N., Vose, J.M., 2015. Complex forest dynamics indicate potential for slowing carbon accumulation in the southeastern United States. *Sci. Rep.* 5, 1–6. <https://doi.org/10.1038/srep08002>.
- Crimmins, S.M., Dobrowski, S.Z., Greenberg, J. a, Abatzoglou, J.T., Mynsberge, A.R., 2011. Changes in climatic water balance drive downhill shifts in plant species' optimum elevations. *Science* 331, 324–327. <https://doi.org/10.1126/science.1199040>.
- Crowther, T.W., Boddy, L., Jones, T.H., 2011. Outcomes of fungal interactions are determined by soil invertebrate grazers. *Ecol. Lett.* 14, 1134–1142. <https://doi.org/10.1111/j.1461-0248.2011.01682.x>.
- Crozier, L.G., Hutchings, J.A., 2013. Plastic and Evolutionary Responses to Climate Change in Fish. <https://doi.org/10.1111/eva.12135>.
- Crozier, L.G., Hendry, A.P., Lawson, P.W., Quinn, T.P., Mantua, N.J., Battin, J., Shaw, R.G., Huey, R.B., 2008. Potential responses to climate change in organisms with complex life histories: evolution and plasticity in Pacific salmon. *Evol. Appl.* 1, 252–270. <https://doi.org/10.1111/j.1752-4571.2008.00033.x>.
- Crozier, L.G., Scheuerell, M.D., Zabel, R.W., 2011. Using time series analysis to characterize evolutionary and plastic responses to environmental change: a case study of a shift toward earlier migration date in sockeye salmon. *Am. Nat.* 178, 755–773. <https://doi.org/10.1086/662669>.
- Crozier, L.G., McClure, M.M., Beechie, T., Bograd, S.J., Boughton, D.A., Carr, M., Cooney, T.D., Dunham, J.B., Greene, C.M., Haltuch, M.A., Hazen, E.L., Holzer, D.M., Huff, D.D., Johnson, R.C., Jordan, C.E., Kaplan, I.C., Lindley, S.T., Mantua, N.J., Moyle, P.B., Myers, J.M., Nelson, M.W., Spence, B.C., Weitkamp, L.A., Williams, T.H., Willis-Norton, E., 2019. Climate vulnerability assessment for Pacific salmon and steelhead in the California Current Large Marine Ecosystem. *PLoS One* <https://doi.org/10.1371/journal.pone.0217711>.
- Daly, E.A., Brodeur, R.D., 2015. Warming ocean conditions relate to increased trophic requirements of threatened and endangered salmon. *PLoS One* 10, 1–23. <https://doi.org/10.1371/journal.pone.0144066>.

- Davidson, E.A., Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 440, 165–173. <https://doi.org/10.1038/nature04514>.
- Davis, C.L., Miller, D.A.W., Walls, S.C., Barichivich, W.J., Riley, J.W., Brown, M.E., 2017. Species interactions and the effects of climate variability on a wetland amphibian metacommunity. *Ecol. Appl.* 27, 285–296. <https://doi.org/10.1002/eap.1442>.
- Davis, J.L., Currin, C.A., O'Brien, C., Raffenburg, C., Davis, A., 2015. Living shorelines: coastal resilience with a blue carbon benefit. *PLoS One* 10, 1–18. <https://doi.org/10.1371/journal.pone.0142595>.
- De Chavez, R., Tauli-Corpus, V., 2009. *Guide on Climate Change and Indigenous Peoples*. Tebtebba Foundation, Baguio City, Philippines.
- De Meester, L., Stoks, R., Brans, K.J., 2018. Genetic adaptation as a biological buffer against climate change: potential and limitations. *Integr. Zool.* 13, 372–391. <https://doi.org/10.1111/1749-4877.12298>.
- Deacy, W.W., Armstrong, J.B., Leacock, W.B., Robbins, C.T., Gustine, D.D., Ward, E.J., Erlenbach, J.A., Stanford, J.A., 2017. Phenological synchronization disrupts trophic interactions between Kodiak brown bears and salmon. *Proc. Natl. Acad. Sci.* 114, 10432–10437.
- DeGregorio, B.A., Westervelt, J.D., Weatherhead, P.J., Sperry, J.H., 2015. Indirect effect of climate change: shifts in ratsnake behavior alter intensity and timing of avian nest predation. *Ecol. Model.* 312, 239–246. <https://doi.org/10.1016/j.ecolmodel.2015.05.031>.
- Delach, A., Caldas, A., Edson, K.M., Krehbiel, R., Murray, S., Theoharides, K.A., Vorhees, L.J., Malcom, J.W., Salvo, M.N., Miller, J.R.B., 2019. Agency plans are inadequate to conserve US endangered species under climate change. *Nat. Clim. Chang.* 9, 999–1004. <https://doi.org/10.1038/s41558-019-0620-8>.
- Dell, A.I., Pawar, S., Savage, V.M., 2014. Temperature dependence of trophic interactions are driven by asymmetry of species responses and foraging strategy. *J. Anim. Ecol.* 83, 70–84. <https://doi.org/10.1111/1365-2656.12081>.
- Dennison, P.E., Brewer, S.C., Arnold, J.D., Moritz, M.A., 2014. Large wildfire trends in the western United States, 1984–2011. *Geophys. Prospect.* 41, 2928–2933. <https://doi.org/10.1002/2014GL061184>. Received.
- Díaz, S., Settele, J., Brondizio, E., Ngo, H.T., Guèze, M., Agard Trinidad, J., Arneth, A., Balvanera, P., Brauman, K., Butchart, S., Chan, K., Garibaldi, L., Ichii, K., Liu, J., Subramanian, S.M., Midgley, G., Miloslavich, P., Molnar, Z., Obura, D., Pfaff, A., Polasky, S., Purvis, A., Razaque, J., Reyers, B., Chowdhury, R.R., Shin, Y.-J., Visseren-Hamakers, I., Willis, K., Zayas, C., 2019. *Summary for Policymakers of the Global Assessment Report on Biodiversity and Ecosystem Services of the Intergovernmental Science-policy Platform on Biodiversity and Ecosystem Services*. Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, Bonn, Germany.
- Dijkstra, J.A., Lambert, W.J., Harris, L.G., 2013. Introduced species provide a novel temporal resource that facilitates native predator population growth. *Biol. Invasions* 15, 911–919. <https://doi.org/10.1007/s10530-012-0339-1>.
- Dodgen, D., Donato, D., Kelly, N., La Greca, A., Morganstein, J., Reser, J., Ruzek, J., Schweitzer, S., Shimamoto, M.M., Tart, K.T., Ursano, R., 2016. *Ch. 8: mental health and well-being. He Impacts of Climate Change on Human Health in the United States: A Scientific Assessment*. U.S. Global Change Research Program, Washington, D.C., USA, pp. 217–246.
- Domke, G., Williams, C.A., Birdsey, R., Coulston, J., Finzi, A., Gough, C., Haight, B., Hicke, J., Janowiak, M., Jong, B. de, Kurz, W.A., Lucas, M., Ogle, S., Olguin-Álvarez, M., Pan, Y., Skutsch, M., Smyth, C., Swanston, C., Templer, P., Wear, D., Woodall, C.W., 2018. *For-ests*. In: Cavallaro, N., Shrestha, G., Birdsey, R., Mayes, M.A., Najjar, R.G., Reed, S.C., Romero-Lankao, P., Zhu, Z. (Eds.), *Second State of the Carbon Cycle Report (SOCCR2)*. U.S. Global Change Research Program, Washington, D.C., USA.
- Donato, J., Grossman, E.E., Konovsky, J., Grossman, S., Campbell, L.W., 2014. Indigenous community health and climate change: integrating biophysical and social science indicators. *Coast. Manag.* 42, 355–373. <https://doi.org/10.1080/08920753.2014.923140>.
- Dunn, P.O., Möller, A.P., 2014. Changes in breeding phenology and population size of birds. *J. Anim. Ecol.* 83, 729–739. <https://doi.org/10.1111/1365-2656.12162>.
- Eakin, C.M., Sweatman, H.P.A., Brainard, R.E., 2019. The 2014–2017 global-scale coral bleaching event: insights and impacts. *Coral Reefs* 38, 539–545. <https://doi.org/10.1007/s00338-019-01844-2>.
- Eastman, L.M., Morelli, T.L., Rowe, K.C., Conroy, C.J., Moritz, C., 2012. Size increase in high elevation ground squirrels over the last century. *Glob. Chang. Biol.* 18, 1499–1508. <https://doi.org/10.1111/j.1365-2486.2012.02644.x>.
- Ebi, K.L., Nealon, J., 2016. Dengue in a changing climate. *Environ. Res.* 151, 115–123. <https://doi.org/10.1016/j.envres.2016.07.026>.
- Elsen, P.R., Tingley, M.W., 2015. Global mountain topography and the fate of montane species under climate change. *Nat. Clim. Chang.* 5, 5–10. <https://doi.org/10.1038/nclimate2656>.
- Elsner, J.B., Kossin, J.P., Jagger, T.H., 2008. The increasing intensity of the strongest tropical cyclones. *Nature* 455, 92–95. <https://doi.org/10.1038/nature07234>.
- EPA, 2017. *Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990–2015*, EPA 430-P-17-001. Washington, D.C. <https://doi.org/10.6009/jirt.2015.jsrt.71.10.1027>.
- Epstein, G., Smale, D.A., 2017. *Undaria Pinnatifida* : A Case Study to Highlight Challenges in Marine Invasion Ecology and Management . pp. 8624–8642. <https://doi.org/10.1002/eece3.3430>.
- Estrada, A., Morales-Castilla, I., Caplat, P., Early, R., 2016. Usefulness of species traits in predicting range shifts. *Trends Ecol. Evol.* 31, 190–203. <https://doi.org/10.1016/j.tree.2015.12.014>.
- Ewald, N.C., Hartley, S.E., Stewart, A.J.A., 2013. Climate change and trophic interactions in model temporary pond systems: the effects of high temperature on predation rate depend on prey size and density. *Freshw. Biol.* 58, 2481–2493. <https://doi.org/10.1111/fwb.12224>.
- Executive Office of the President of the United States, 2015. *M-16-01: Incorporating Ecosystem Services Into Federal Decision Making*.
- Fair, J., Hunt, L., 2015. *Headwater Meadow Restoration in the Sierra Nevada: Adapting to Climate Change [WWW Document]*. *Clim. Chang. Resour. Cent.*
- Ferrario, F., Beck, M.W., Storazzi, C.D., Micheli, F., Shepard, C.C., Airolidi, L., 2014. The effectiveness of coral reefs for coastal hazard risk reduction and adaptation. *Nat. Commun.* 5, 1–9. <https://doi.org/10.1038/ncomms4794>.
- Field, J.C., Elliger, C., Baltz, K., Gillespie, G.E., Gilly, W.F., Ruiz-Cooley, R.I., Pearse, D., Stewart, J.S., Matsubu, W., Walker, W.A., 2013. Foraging ecology and movement patterns of jumbo squid (*Dosidicus gigas*) in the California Current System. *Deep. Res. Part II Top. Stud. Oceanogr.* 95, 37–51. <https://doi.org/10.1016/j.dsr2.2012.09.006>.
- Fischer, J., Lindenmayer, D.B., Manning, A.D., 2006. *Biodiversity, Ecosystem Function, and Resilience : Ten Guiding Principles for Commodity Production Landscapes*.
- Foden, W.B., Young, B.E., 2016. *IUCN SSC Guidelines for Assessing Species' Vulnerability to Climate Change*. Cambridge, England and Gland, Switzerland. <https://doi.org/10.1038/scientificamerican0315-S1>.
- Foster, J.R., D'Amato, A.W., 2015. Montane forest ecotones moved downslope in north-eastern USA in spite of warming between 1984 and 2011. *Glob. Chang. Biol.* 21, 4497–4507. <https://doi.org/10.1111/gcb.13046>.
- Fox, R.J., Donelson, J.M., Schunter, C., Ravasi, T., Gaitán-Espitia, J.D., 2019. Beyond buying time: the role of plasticity in phenotypic adaptation to rapid environmental change. *Philos. Trans. R. Soc. B Biol. Sci.* 374. <https://doi.org/10.1098/rstb.2018.0174>.
- Franks, S.J., Weber, J.J., Aitken, S.N., 2014. Evolutionary and plastic responses to climate change in terrestrial plant populations. *Evol. Appl.* 7, 123–139. <https://doi.org/10.1111/eva.12112>.
- Fu, Y.H., Zhao, H., Piao, S., Peaucelle, M., Peng, S., Zhou, G., Ciais, P., Song, Y., Vitis, Y., Zeng, Z., Janssens, I. a, Huang, M., Menzel, A., Pen, J., 2015. Declining global warming effects on the phenology of spring leaf unfolding. *Nature* 526, 104–107. <https://doi.org/10.1038/nature15402>.
- Galloway, J.N., Townsend, A.R., Erisman, J.W., Bekunda, M., Cai, Z., Freney, J.R., Martinelli, L. a, Seitzinger, S.P., Sutton, M. a, 2008. *Transformation of the nitrogen cycle: recent trends, questions, and potential solutions*. *Science* (80- ) 320, 889–892.
- García Molinos, J., Halpern, B.S., Schoeman, D.S.S., Brown, C.J.J., Kiessling, W., Moore, P.J.J., Pandolfi, J.M.M., Poloczanska, E.S.S., Richardson, A.J.J., Burrows, M.T.T., García Molinos, J., Halpern, B.S.S., Schoeman, D.S.S., Brown, C.J.J., Kiessling, W., Moore, P.J.J., Pandolfi, J.M.M., Poloczanska, E.S.S., Richardson, A.J.J., Burrows, M.T.T., 2015. Climate velocity and the future global redistribution of marine biodiversity. *Nat. Clim. Chang.* 6, 4–11. <https://doi.org/10.1038/nclimate2769>.
- Gardner, J.L., Peters, A., Kearney, M.R., Joseph, L., Heinsohn, R., 2011. Declining body size: a third universal response to warming? *Trends Ecol. Evol.* 26, 285–291. <https://doi.org/10.1016/j.tree.2011.03.005>.
- Gergel, D.R., Nijssen, B., Abatzoglou, J.T., Lettenmaier, D.P., Stumbaugh, M.R., 2017. Effects of climate change on snowpack and fire potential in the western USA. *Clim. Chang.* 141, 287–299. <https://doi.org/10.1007/s10584-017-1899-y>.
- Ghalambor, C.K., McKay, J.K., Carroll, S.P., Reznick, D.N., 2007. Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Funct. Ecol.* 21, 394–407. <https://doi.org/10.1111/j.1365-2435.2007.01283.x>.
- Giakoumi, S., Katsanevakis, S., Albano, P.G., Azzurro, E., Cardoso, A.C., Cebrian, E., Deidun, A., Edelist, D., Francour, P., Jimenez, C., Mačić, V., Occhipinti-Ambrogi, A., Rilov, G., Sghaier, Y.R., 2019. Management priorities for marine invasive species. *Sci. Total Environ.* 688, 976–982. <https://doi.org/10.1016/j.scitotenv.2019.06.282>.
- Gittman, R.K., Popowich, A.M., Bruno, J.F., Peterson, C.H., 2014. Marshes with and without sills protect estuarine shorelines from erosion better than bulkheads during a category 1 hurricane. *Ocean Coast. Manag.* 102, 94–102. <https://doi.org/10.1016/j.ocecoaman.2014.09.016>.
- Glick, P., Stein, B.A., Edelson, N.A., 2011. *Scanning the Conservation Horizon: A Guide to Climate Change Vulnerability Assessment*.
- Gomulkiewicz, R., Kingsolver, J.G., Carter, P.A., Heckman, N., 2018. Variation and evolution of function-valued traits. *Annu. Rev. Ecol. Syst.* 49, 139–164. <https://doi.org/10.1146/annurev-ecolsys-110316-022830>.
- Gowda, P., Steiner, J.L., Olson, C., Boggess, M., Farrigan, T., Grusak, M.A., 2018. *Agriculture and rural communities*. In: Reidmiller, D.R., Avery, C.W., Easterling, D.R., Kunkel, K.E., Lewis, K.L.M., Maycock, T.K., Stewart, B.C. (Eds.), *Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment. II*. U.S. Global Change Research Program, Washington, D.C., USA, pp. 391–437.
- Graven, H.D., Keeling, R.F., Piper, S.C., Patra, P.K., Stephens, B.B., Wofsy, S.C., Santoni, G.W., Bent, J.D., 2013. *Enhanced Seasonal Exchange of CO<sub>2</sub> by Northern Ecosystems Since 1960*. 146 pp. 1085–1090.
- Gregory, R., Failing, L., Harstone, M., Long, G., McDaniels, T., Ohlson, D., 2012. *Structuring environmental management choices. Structured Decision Making: A Practical Guide to Environmental Management Choices*. John Wiley & Sons, Ltd, Chichester, UK.
- Grenier, S., Barre, P., Litrico, I., 2016. Phenotypic plasticity and selection: nonexclusive mechanisms of adaptation. *Scientifica (Cairo)* 2016. <https://doi.org/10.1155/2016/7021701>.
- Grosholz, E.D., Ruiz, G.M., Dean, C.A., Shirley, K.A., Maron, J.L., Connors, P.G., 2000. The impacts of a nonindigenous marine predator in a California bay. *Ecology* 81, 1206–1224. [https://doi.org/10.1890/0012-9658\(2000\)081\[1206:TIOANM\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[1206:TIOANM]2.0.CO;2).
- Grossman, Z., Parker, A., 2012. *Asserting Native Resilience: Pacific Rim Indigenous Nations Face the Climate Crisis*. Oregon State University Press.
- Guo, F., Lenoir, J., Bonebrake, T.C., 2018. Land-use change interacts with climate to determine elevational species redistribution. *Nat. Commun.* 9, 1–7. <https://doi.org/10.1038/s41467-018-03786-9>.
- Hannah, L., Flint, L., Syphard, A.D., Moritz, M.A., Buckley, L.B., McCullough, I.M., 2014. Fine-grain modeling of species' response to climate change: holdouts, stepping-stones, and microrefugia. *Trends Ecol. Evol.* 29, 390–397. <https://doi.org/10.1016/j.tree.2014.04.006>.

- Hare, J.A., Borggaard, D.L., Friedland, K.D., Anderson, J., Burns, P., Chu, K., Clay, P.M., Collins, M.J., Cooper, P., Fratanoti, P.S., Johnson, M.R., Manderson, J.P., Milke, L., Miller, T.J., Orphanides, C.D., Saba, V.S., 2016a. Northeast Regional Action Plan - NOAA Fisheries Climate Science Strategy.
- Hare, J.A., Morrison, W.E., Nelson, M.W., Stachura, M.M., Teeters, E.J., Griffiths, R.B., Alexander, M.A., Scott, J.D., Alade, L., Bell, R.J., Chute, A.S., Curti, K.L., Curtis, T.H., Kirches, D., Kocik, J.F., Lucey, S.M., McCandless, C.T., Milke, L.M., Richardson, D.E., Robillard, E., Walsh, H.J., McManus, M.C., Marancik, K.E., Griswold, C.A., 2016b. A vulnerability assessment of fish and invertebrates to climate change on the northeast U.S. continental shelf. *PLoS One* 11, 1–30. <https://doi.org/10.1371/journal.pone.0146756>.
- Harrison, K.A., Pavlova, A., Telonis-Scott, M., Sunnucks, P., 2014. Using genomics to characterize evolutionary potential for conservation of wild populations. *Evol. Appl.* 7, 1008–1025. <https://doi.org/10.1111/eva.12149>.
- Hember, R.A., Kurz, W.A., Coops, N.C., 2017. Relationships between individual-tree mortality and water-balance variables indicate positive trends in water stress-induced tree mortality across North America. *Glob. Chang. Biol.* 23, 1691–1710. <https://doi.org/10.1111/gcb.13428>.
- Hendry, A.P., 2017. *Eco-evolutionary Dynamics*. Princeton University Press, Princeton, NJ.
- Hendry, A.P., Kinnison, M.T., Heino, M., Day, T., Smith, T.B., Fitt, G., Bergstrom, C.T., Oakeshott, J., Jørgensen, P.S., Zalucki, M.P., Gilchrist, G., Southerton, S., Sih, A., Strauss, S., Denison, R.F., Carroll, S.P., 2011. Evolutionary principles and their practical application. *Evol. Appl.* 4, 159–183. <https://doi.org/10.1111/j.1752-4571.2010.00165.x>.
- Herstoff, E., Urban, M.C., 2014. Will pre-adaptation buffer the impacts of climate change on novel species interactions? *Ecography (Cop.)* 37, 111–119. <https://doi.org/10.1111/j.1600-0587.2013.00116.x>.
- Hilborn, E.D., Beasley, V.R., 2015. One health and cyanobacteria in freshwater systems: animal illnesses and deaths are sentinel events for human health risks. *Toxins (Basel)* 7, 1374–1395. <https://doi.org/10.3390/toxins7041374>.
- Hobbs, R.J., Higgins, E., Harris, J.A., 2009. Novel ecosystems: implications for conservation and restoration. *Trends Ecol. Evol.* 24, 599–605. <https://doi.org/10.1016/j.tree.2009.05.012>.
- Hobday, A.J., Alexander, L.V., Perkins, S.E., Smale, D.A., Straub, S.C., Oliver, E.C.J., Benthuyssen, J.A., Burrows, M.T., Donat, M.G., Feng, M., Holbrook, N.J., Moore, P.J., Scannell, H.A., Sen Gupta, A., Wernberg, T., 2016. A hierarchical approach to defining marine heatwaves. *Prog. Oceanogr.* 141, 227–238. <https://doi.org/10.1016/j.pcean.2015.12.014>.
- Hobday, A.J., Oliver, E.C.J., Gupta, A., Sen, Benthuyssen, J.A., Burrows, M.T., Donat, M.G., Holbrook, N.J., Moore, P.J., Thomsen, M.S., Wernberg, T., Smale, D.A., 2018. Categorizing and naming marine heatwaves. *Oceanography* 31, 162–173. <https://doi.org/10.5670/oceanog.2018.205>.
- Hobday, A.J., Hartog, J.R., Manderson, J.P., Mills, K.E., Oliver, M.J., Pershing, A.J., Siedlecki, S., 2019. Ethical considerations and unanticipated consequences associated with ecological forecasting for marine resources. *ICES J. Mar. Sci.* <https://doi.org/10.1093/icesjms/fsy210>.
- Holden, Z.A., Swanson, A., Luce, C.H., Jolly, W.M., Maneta, M., Oyler, J.W., Warren, D.A., Parsons, R., Affleck, D., 2018. Decreasing fire season precipitation increased recent western US forest wildfire activity. *Proc. Natl. Acad. Sci. U. S. A.* 115, E8349–E8357. <https://doi.org/10.1073/pnas.1802316115>.
- Holsman, K.K., Hazen, E.L., Haynie, A., Gouguet, S., Hollowed, A., Bograd, S.J., Samhouri, J.F., Aydin, K., 2019. Towards climate resiliency in fisheries management. *ICES J. Mar. Sci.* <https://doi.org/10.1093/icesjms/fsz031>.
- Houser, S., Teller, V., MacCracken, M., Gough, R., Spears, P., 2001. *Potential consequences of climate variability and change for native peoples and homelands. Climate Change Impacts on the United States*. U.S. Global Change Research Program, Washington, D.C., USA.
- Howard, J., Sutton-Grier, A., Herr, D., Kleypas, J., Landis, E., Mcleod, E., Pidgeon, E., Simpson, S., 2017. Clarifying the role of coastal and marine systems in climate mitigation. *Front. Ecol. Environ.* 15, 42–50. <https://doi.org/10.1002/fee.1451>.
- Huntington, T.G., Hodgkins, G.A., Dudley, R.W., 2003. Historical trend in river ice thickness and coherence in hydroclimatological trends in Maine. *Clim. Chang.* 61, 217–236. <https://doi.org/10.1023/A:1026360615401>.
- Invasive Species Advisory Committee, 2017. *Managed Relocation : Reducing the Risk of Biological Invasion*. pp. 1–6.
- IPCC, 2013. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Isaac-Renton, M.G., Roberts, D.R., Hamann, A., Spiecker, H., 2014. Douglas-fir plantations in Europe: a retrospective test of assisted migration to address climate change. *Glob. Chang. Biol.* 20, 2607–2617. <https://doi.org/10.1111/gcb.12604>.
- Isaak, D.J., Wollrab, S., Horan, D., Chandler, G., 2012. Climate change effects on stream and river temperatures across the northwest U.S. from 1980–2009 and implications for salmonid fishes. *Clim. Chang.* 113, 499–524. <https://doi.org/10.1007/s10584-011-0326-z>.
- Jacox, M.G., Alexander, M.A., Stock, C.A., Hervieux, G., 2017. On the skill of seasonal sea surface temperature forecasts in the California Current System and its connection to ENSO variability. *Clim. Dyn.* 0, 1–15. <https://doi.org/10.1007/s00382-017-3608-y>.
- Jantarassami, L.C., Novak, R., Delgado, R., Marino, E., McNeeley, S., Narducci, C., Raymond-Yakoubian, J., Singletary, L., Whyte, K.P., 2018. Tribes and indigenous Peoples. In: Reidmiller, D.R., Avery, C.W., Easterling, D.R., Kunkel, K.E., Lewis, K.L.M., Maycock, T.K., Stewart, B.C. (Eds.), *Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment, Volume II*. U.S. Global Change Research Program, Washington, D.C., USA.
- Jay, A., Reidmiller, D.R., Avery, C.W., Barrie, D., DeAngelo, B.J., Dave, A., Dzaugis, M., Kolan, M., Lewis, K.L.M., Reeves, K., Winner, D., 2018. Overview. In: Reidmiller, D.R., Avery, C.W., Easterling, D.R., Kunkel, K.E., Lewis, K.L.M., Maycock, T.K., Stewart, B.C. (Eds.), *Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment. II. U.S. Global Change Research Program, Washington, D.C., USA*, pp. 33–71.
- Jeremias, G., Barbosa, J., Marques, S.M., Asselman, J., Gonçalves, F.J.M., Pereira, J.L., 2018. Synthesizing the role of epigenetics in the response and adaptation of species to climate change in freshwater ecosystems. *Mol. Ecol.* 27, 2790–2806. <https://doi.org/10.1111/mec.14727>.
- Juanes, F., Gephard, S., Beland, K.F., 2004. Long-term changes in migration timing of adult Atlantic salmon (*Salmo salar*) at the southern edge of the species distribution. *Can. J. Fish. Aquat. Sci.* 61, 2392–2400. <https://doi.org/10.1139/f04-207>.
- Jurgens, L.J., Gaylor, B., 2018. Physical effects of habitat-forming species override latitudinal trends in temperature. *Ecol. Lett.* 21, 190–196. <https://doi.org/10.1111/ele.12881>.
- Katsanevakis, S., Wallentinus, I., Zenetos, A., Leppäkoski, E., Çinar, M.E., 2014. Impacts of invasive alien marine species on ecosystem services and biodiversity: a pan-European review. *Acquat. Invasions* 9, 391–423. <https://doi.org/10.3391/ai.2014.9.4.01>.
- Keane, R.E., Ryan, K.C., Veblen, T.T., Allen, C.D., Logan, J., Hawkes, B., 2002. *Cascading effects of fire exclusion in rocky mountain ecosystems: a literature review*. USDA For. Serv. - Gen. Tech. Rep. RMRS-GTR-1–27.
- Kemp, K.B., Blades, J.J., Klos, P.Z., Hall, T.E., Force, J.E., Morgan, P., Tinkham, W.T., 2015. Managing for climate change on federal lands of the western United States: perceived usefulness of climate science, effectiveness of adaptation strategies, and barriers to implementation. *Ecol. Soc.* 20. <https://doi.org/10.5751/ES-07522-200217>.
- Keppel, G., Mokany, K., Wardell-Johnson, G.W., Phillips, B.L., Welbergen, J.A., Reside, A.E., 2015. The capacity of refugia for conservation planning under climate change. *Front. Ecol. Environ.* 13, 106–112. <https://doi.org/10.1890/140055>.
- Kingsolver, J.G., Buckley, L.B., 2017. Quantifying thermal extremes and biological variation to predict evolutionary responses to changing climate. *Philos. Trans. R. Soc. B Biol. Sci.* 372. <https://doi.org/10.1098/rstb.2016.0147>.
- Kitzberger, T., Falk, D.A., Westerling, A.L., Swetnam, T.W., 2017. Direct and indirect climate controls predict heterogeneous early-mid 21st century wildfire burned area across western and boreal North America. *PLoS One* 12, 1–24. <https://doi.org/10.1371/journal.pone.0188486>.
- Kleisner, K.M., Fogarty, M.J., McGee, S., Barnett, A., Fratanoti, P., Greene, J., Hare, J.A., Lucey, S.M., McGuire, C., Odell, J., Saba, V.S., Smith, L., Weaver, K.J., Pinsky, M.L., 2016. The effects of sub-regional climate velocity on the distribution and spatial extent of marine species assemblages. *PLoS One* 11, 1–21. <https://doi.org/10.1371/journal.pone.0149220>.
- Kleisner, K.M., Fogarty, M.J., McGee, S., Hare, J.A., Moret, S., Perretti, C.T., Saba, V.S., 2017. Marine species distribution shifts on the U.S. Northeast Continental Shelf under continued ocean warming. *Prog. Oceanogr.* 153, 24–36. <https://doi.org/10.1016/j.pcean.2017.04.001>.
- Kopp, M., Matuszewski, S., 2014. Rapid evolution of quantitative traits: theoretical perspectives. *Evol. Appl.* 7, 169–191. <https://doi.org/10.1111/eva.12127>.
- Kovach, R.P., Dunham, J.B., Al-Chokhachy, R., Snyder, C.D., Letcher, B.H., Young, J.A., Beever, E.A., Pederson, G.T., Lynch, A.J., Hitt, N.P., Konrad, C.P., Jaeger, K.L., Rea, A.H., Sepulveda, A.J., Lambert, P.M., Stoker, J., Giersch, J.J., Muhlfeld, C.C., 2019. An integrated framework for ecological drought across riverscapes of North America. *Bioscience* 69, 418–431. <https://doi.org/10.1093/biosci/biz040>.
- Krakoff, S., 2008. *American Indians, climate change, and ethics for a warming world*. Denver Univ. Law Rev. 85, 865–898.
- Lam, V.W.Y., Cheung, W.W.L., Reygondeau, G., Rashid Sumaila, U., 2016. Projected change in global fisheries revenues under climate change. *Sci. Rep.* 6, 6–13. <https://doi.org/10.1038/srep32607>.
- Lany, N.K., Ayres, M.P., Stange, E.E., Sillett, T.S., Rodenhouse, N.L., Holmes, R.T., 2016. Breeding timed to maximize reproductive success for a migratory songbird: the importance of phenological asynchrony. *Oikos* 125, 656–666. <https://doi.org/10.1111/oik.02412>.
- Latta, G., Temesgen, H., Adams, D., Barrett, T., 2010. Analysis of potential impacts of climate change on forests of the United States Pacific Northwest. *For. Ecol. Manag.* 259, 720–729. <https://doi.org/10.1016/j.foreco.2009.09.003>.
- Lawrence, D.J., Stewart-Koster, B., Olden, J.D., Ruesch, A.S., Torgersen, C.E., Lawler, J.J., Butcher, D.P., Crown, J.K., 2014. The interactive effects of climate change, riparian management, and a nonnative predator on stream-rearing salmon. *Ecol. Appl.* 24, 895–912. <https://doi.org/10.1890/13-0753.1>.
- Le Bris, A., Pershing, A.J., Gaudette, J., Pugh, T.L., Reardon, K.M., 2017. Multi-scale quantification of the effects of temperature on size at maturity in the American lobster (*Homarus americanus*). *Fish. Res.* 186, 397–406. <https://doi.org/10.1016/j.fishres.2016.09.008>.
- Lefort, S., Aumont, O., Bopp, L., Arsouze, T., Gehlen, M., Maury, O., 2015. Spatial and body-size dependent response of marine pelagic communities to projected global climate change. *Glob. Chang. Biol.* 21, 154–164. <https://doi.org/10.1111/gcb.12679>.
- Lehikoinen, A., Lindén, A., Karlsson, M., Andersson, A., Crewe, T.L., Dunn, E.H., Gregory, G., Karlsson, L., Kristiansen, V., Mackenzie, S., Newman, S., Roer, J.E., Sharpe, C., Sokolov, L.V., Steinholtz, A., Stervander, M., Tirri, I.S., Tjørnløv, R.S., 2019. Phenology of the avian spring migratory passage in Europe and North America: asymmetric advancement in time and increase in duration. *Ecol. Indic.* 101, 985–991. <https://doi.org/10.1016/j.ecolind.2019.01.083>.
- Lembrechts, J.J., Pauchard, A., Lenoir, J., Nuñez, M.A., Geron, C., Ven, A., Bravo-Monasterio, P., Teneb, E., Nijis, I., Milbau, A., 2016. Disturbance is the key to plant invasions in cold environments. *Proc. Natl. Acad. Sci. U. S. A.* 113, 14061–14066. <https://doi.org/10.1073/pnas.1608980113>.

- Lenoir, J., Svenning, J.C., 2015. Climate-related range shifts - a global multidimensional synthesis and new research directions. *Ecography* (Cop.) 38, 15–28. <https://doi.org/10.1111/ecog.00967>.
- Li, D., Wrzesien, M.L., Durand, M., Adam, J., Lettenmaier, D.P., 2017. How much runoff originates as snow in the western United States, and how will that change in the future? *Geophys. Res. Lett.* 44, 6163–6172. <https://doi.org/10.1002/2017GL073551>.
- Link, J., 2016. *Ecosystem-based Fisheries Management Policy of the National Marine Fisheries Service*. National Oceanic and Atmospheric Administration.
- Link, J.S., Griffis, R., Busch, S., 2015. *Noaa Fisheries Climate Science Strategy* (doi:NOAA Technical Memorandum NMFS-F/SPO-155).
- Lipton, D., Rubenstein, M.A., Weiskopf, S.R., Carter, S., Peterson, J., Crozier, L., Fogarty, M., Gaichas, S., Hyde, K.J.W., Morelli, T.L., Morissette, J., Moustahfid, H., Muñoz, R., Poudel, R., Staudinger, M.D., Stock, C., Thompson, L., Waples, R., Weltzin, J.F., 2018. Ecosystems, ecosystem services, and biodiversity. In: Reidmiller, D.R., Avery, C.W., Easterling, D.R., Kunkel, K.E., Lewis, K.L.M., Maycock, T.K., Stewart, B.C. (Eds.), *Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment, Volume II*. U.S. Global Change Research Program, Washington, D.C., USA.
- Litschert, S.E., Brown, T.C., Theobald, D.M., 2012. Historic and future extent of wildfires in the Southern Rockies Ecoregion, USA. *For. Ecol. Manag.* 269, 124–133. <https://doi.org/10.1016/j.foreco.2011.12.024>.
- Littell, J.S., Peterson, D.L., Riley, K.L., Liu, Y., Luce, C.H., 2016. A review of the relationships between drought and forest fire in the United States. *Glob. Chang. Biol.* 22, 2353–2369. <https://doi.org/10.1111/gcb.13275>.
- Liu, Y., Oduor, A.M.O., Zhang, Z., Manea, A., Tooth, I.M., Leishman, M.R., Xu, X., van Kleunen, M., 2017. Do invasive alien plants benefit more from global environmental change than native plants? *Glob. Chang. Biol.* 23, 3363–3370. <https://doi.org/10.1111/gcb.13579>.
- Liu, Z., Wimberly, M.C., Lamsal, A., Sohl, T.L., Hawbaker, T.J., 2015. Climate change and wildfire risk in an expanding wildland–urban interface: a case study from the Colorado Front Range Corridor. *Landsc. Ecol.* 30, 1943–1957. <https://doi.org/10.1007/s10980-015-0222-4>.
- Loehman, R.A., Keane, R.E., Holsinger, L.M., Wu, Z., 2017. Interactions of landscape disturbances and climate change dictate ecological pattern and process: spatial modeling of wildfire, insect, and disease dynamics under future climates. *Landsc. Ecol.* 32, 1447–1459. <https://doi.org/10.1007/s10980-016-0414-6>.
- Logan, J.A., Powell, J.A., 2009. *Ecological consequences of climate change altered forest insect disturbance regimes*. In: Wagner, F.H. (Ed.), *Climate Change in Western North America: Evidence and Environmental Effects*. University of Utah Press, Salt Lake City, Utah, USA, pp. 98–109.
- Luce, C., Morgan, P., Dwire, K., Isaak, D., Holden, Z., Rieman, B., 2012. Climate change, forests, fire, water, and fish: building resilient landscapes, streams, and managers. *USDA For. Serv. - Gen. Tech. Rep. RMRS-GTR 1-155*.
- Luce, C.H., Lopez-Burgos, V., Holden, Z., 2014. Sensitivity of snowpack storage to precipitation and temperature using spatial and temporal analog models. *Water Resour. Res.* 50, 9447–9462. <https://doi.org/10.1111/j.1752-1688.1969.tb04897.x>.
- MacLean, S.A., Beissinger, S.R., 2017. Species' traits as predictors of range shifts under contemporary climate change: a review and meta-analysis. *Glob. Chang. Biol.* 23, 4094–4105. <https://doi.org/10.1111/gcb.13736>.
- Mahmoud, M., Liu, Y., Hartmann, H., Stewart, S., Wagener, T., Semmens, D., Stewart, R., Gupta, H., Dominguez, D., Dominguez, F., Hulse, D., Letcher, R., Rashleigh, B., Smith, C., Street, R., Ticehurst, J., Twery, M., Delden, H. Van, Waldick, R., White, D., Winter, L., 2009. A formal framework for scenario development in support of environmental decision-making. *Environ. Model. Softw.* 24, 798–808. <https://doi.org/10.1016/j.envsoft.2008.11.010>.
- Maldonado, J.K., Pandya, R.E., Colombi, B.J., 2013. Climate change and indigenous peoples in the United States: impacts, experiences, and actions. *Clim. Chang.* 120, 509–682.
- Malmstadt, J.C., Elsner, J.B., Jagger, T.H., 2010. Risk of strong hurricane winds to Florida cities. *J. Appl. Meteorol. Climatol.* 49, 2121–2132. <https://doi.org/10.1175/2010JAMC2420.1>.
- Marcinkowski, K., Peterson, D.L., Ettl, G.J., 2015. Nonstationary temporal response of mountain hemlock growth to climatic variability in the north cascade range, Washington, USA. *Can. J. For. Res.* 45, 676–688. <https://doi.org/10.1139/cjfr-2014-0231>.
- Maynard, N.G., 1998. *Native Peoples-native Homelands Climate Change Workshop: Lessons Learned. Final Report* (Washington, D.C., USA).
- Mayor, S.J., Guralnick, R.P., Tingley, M.W., Otegui, J., Withey, J.C., Elmendorf, S.C., Andrew, M.E., Leyk, S., Pearce, I.S., Schneider, D.C., 2017. Increasing phenological asynchrony between spring green-up and arrival of migratory birds. *Sci. Rep.* 7, 1902. <https://doi.org/10.1038/s41598-017-02045-z>.
- McCann, E.L., Johnson, N.S., Pangle, K.L., 2017. Corresponding long-term shifts in stream temperature and invasive fish migration. *Can. J. Fish. Aquat. Sci.* 778, 1–7. <https://doi.org/10.1139/cjfas-2017-0195>.
- McCluney, K.E., Sabo, J.L., 2016. Animal water balance drives top-down effects in a riparian forest - implications for terrestrial trophic cascades. *Proc. R. Soc. B* 283, 20160881.
- McGuire, J.L., Lawler, J.J., Mcrae, B.H., Nuñez, T.A., Theobald, D.M., 2016. Achieving Climate Connectivity in a Fragmented Landscape. , p. 113. <https://doi.org/10.1073/pnas.1602817113>.
- McKenzie, D., Littell, J.S., 2017. Climate change and the eco-hydrology of fire: will area burned increase in a warming western USA. *Ecol. Appl.* 27, 26–36. <https://doi.org/10.1002/eap.1420>.
- Mecklenburg, C.W., TA, M., Sheiko, B.A., Steinke, D., 2016. *Pacific Arctic marine fishes. Monitoring Series Report No 23. Conservation of Arctic Flora and Fauna*.
- Melillo, J.M., Frey, S.D., DeAngelis, K.M., Werner, W.J., Bernard, M.J., Bowles, F.P., Pold, G., Knorr, M.A., Grandy, A.S., 2017. Long-term pattern and magnitude of soil carbon feedback to the climate system in a warming world. *Science* (80-. ) 358, 101–105. <https://doi.org/10.1126/science.aan2874>.
- Merilä, J., 2012. Evolution in response to climate change: in pursuit of the missing evidence. *BioEssays* 34, 811–818.
- Merilä, J., Hendry, A.P., 2014. Climate change, adaptation, and phenotypic plasticity: the problem and the evidence. *Evol. Appl.* <https://doi.org/10.1111/eva.12137>.
- Millar, C.L., Stephenson, N.L., 2015. Temperate forest health in an era of emerging megadisturbance. *Science* (80-. ) 349, 823–826.
- Millennium Ecosystem Assessment, 2005. *Ecosystems and Human Well-being: Synthesis*. Island Press, Washington, D.C.
- Mills, K.E., Pershing, A.J., Brown, C.J., Chen, Y., Chiang, F.-S., Holland, D.S., Lehuta, S., Nye, J.A., Sun, J.C., Thomas, A.C., 2013. Fisheries management in a changing climate: lessons from the 2012 ocean heat wave in the Northwest Atlantic. *Oceanography* 26, 191–195.
- Möller, I., Kudella, M., Rupprecht, F., Spencer, T., Paul, M., Van Wesenbeeck, B.K., Wolters, G., Jensen, K., Bouma, T.J., Miranda-Lange, M., Schimmels, S., 2014. Wave attenuation over coastal salt marshes under storm surge conditions. *Nat. Geosci.* 7, 727–731. <https://doi.org/10.1038/NGEO2251>.
- Monahan, W.B., Rosemartin, A., Gerst, K.L., Fisichelli, N.A., Ault, T., Schwartz, M.D., Gross, J.E., Weltzin, J.F., 2016. Climate change is advancing spring onset across the U.S. national park system. *Ecosphere* 7, 1–17. <https://doi.org/10.1002/ecs2.1465>.
- Moore, S.K., Trainer, V.L., Mantua, N.J., Parker, M.S., Laws, E.A., Backer, L.C., Fleming, L.E., 2008. Impacts of climate variability and future climate change on harmful algal blooms and human health. *Environ. Heal. A Glob. Access Sci. Source* 7, 1–12. <https://doi.org/10.1186/1476-069X-7-S2-S4>.
- Morelli, T.L., Daly, C., Dobrowski, S.Z., Dulen, D.M., Ebersole, J.L., Jackson, S.T., Lundquist, J.D., Millar, C.L., Maher, S.P., Monahan, W.B., Nydick, K.R., Redmond, K.T., Sawyer, S.C., Stock, S., Beissinger, S.R., 2016. Managing climate change refugia for climate adaptation. *PLoS One* 11, e0159909. <https://doi.org/10.1371/journal.pone.0159909>.
- Morelli, T.L., Maher, S.P., Lim, M.C.W., Kastely, C., Eastman, L.M., Flint, L.E., Flint, A.L., Beissinger, S.R., Moritz, C., 2017. Climate change refugia and habitat connectivity promote species persistence. *Clim. Chang. Responses* 4, 1–12. <https://doi.org/10.1186/s40665-017-0036-5>.
- Morishima, G., 2014. *Climate Change and Indigenous Peoples: A Primer*.
- Morley, J.W., Batt, R.D., Pinsky, M.L., 2017. Marine assemblages respond rapidly to winter climate variability. *Glob. Chang. Biol.* 23, 2590–2601. <https://doi.org/10.1111/gcb.13578>.
- Morris, K.A., Detering, N.M., 2016. The emotional cost of distance: geographic social network dispersion and post-traumatic stress among survivors of Hurricane Katrina. *Soc. Sci. Med.* 165, 56–65.
- National Fish Wildlife and Plants Climate Adaptation Partnership, 2012. *National Fish, Wildlife, and Plants Climate Adaptation Strategy* (WWW Document).
- National Marine Fisheries Service, 2016. *National Marine Fisheries Service Procedural Instruction 02-110-18: Guidance for Treatment of Climate Change in NMFS Endangered Species Act Decisions*.
- National Marine Fisheries Service, 2018. *Fisheries Economics of the United States, 2016: Economics and Sociocultural Status and Trends Series*. NOAA Tech. Memo. NMFS-F/SPO-187a. <https://doi.org/10.1017/CBO9781107415324.004>.
- National Park Service, 2013. *Resource Stewardship Strategy: Catcotin Mountain Park*.
- National Park System Advisory Board, 2012. *Revisiting Leopold : Resource Stewardship in the National Parks*.
- National Wildlife Federation, 2010. *Business Plan SIERRA NEVADA MEADOW*.
- Niemi, E., Buckley, M., Neculae, C., Reich, S., 2009. *An Overview of Potential Economic Costs to Washington of a Business-As-Usual Approach to Climate Change*. University of Oregon, Program on Climate Economics of the Climate Leadership Initiative, Eugene, OR.
- NOAA Fisheries, 2015. *Farmers reroute water for fish: Yakima irrigators use canals to keep streams flowing in drought* [WWW document]. NOAA Fish. West Coast Reg.
- NOAA Fisheries, 2016. *2015 Adult Sockeye Salmon Passage Report*.
- Norby, R.J., Hanson, P.J., O'Neill, E.G., Tschaplinski, T.J., Weltzin, J.F., Hansen, R.A., Cheng, W., Wullschlegel, S.D., Gunderson, C.A., Edwards, N.T., Johnson, D.W., 2002. Enriched deciduous forest and the implications for carbon storage. *Ecol. Appl.* 12, 1261–1266. [https://doi.org/10.1890/1051-0761\(2002\)1012\[1261:NPPAC\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2002)1012[1261:NPPAC]2.0.CO;2).
- Norby, R.J., Warren, J.M., Iversen, C.M., Medlyn, B.E., McMurtrie, R.E., 2010. CO<sub>2</sub> enhancement of forest productivity constrained by limited nitrogen availability. *Proc. Natl. Acad. Sci. U. S. A.* 107, 19368–19373. <https://doi.org/10.1073/pnas.1006463107>.
- Norton-Smith, K., Lynn, K., Chief, K., Cozzetto, K., Donatuto, J., Redsteer, M.H., Kruger, L.E., Maldonado, J., Viles, C., Whyte, K.P., 2016. *Climate Change and Indigenous Peoples : A Synthesis of Current Impacts and Experiences*. Gen. Tech. Rep. PNW-GTR-944. pp. 1–142.
- Novak, M., Wootton, J.T., Doak, D.F., Emmerson, M., Estes, J.A., Tinker, T., 2011. Predicting community responses to perturbations in the face of imperfect knowledge and network complexity. *Ecology* 92, 836–846.
- Obradovich, N., Migliorini, R., Paulus, M.P., Rahwan, I., 2018. Empirical evidence of mental health risks posed by climate change. *Proc. Natl. Acad. Sci. U. S. A.* 115, 10953–10958. <https://doi.org/10.1073/pnas.1801528115>.
- Ojima, D.S., Iverson, L.R., Sohngen, B.L., Vose, J.M., Woodall, C.W., Domke, G.M., Peterson, D.L., Littell, J.S., Matthews, S.N., Prasad, A.M., Peters, M.P., Yohe, G.W., Friggens, M.M., 2014. Risk assessment. In: Peterson, D.L., Vose, J.M., Patel-Weyand, T. (Eds.), *Climate Change and United States Forests*. Springer, Dordrecht, The Netherlands, pp. 223–244. <https://doi.org/10.1007/978-94-007-7515-2>.
- Oliver, T.H., Heard, M.S., Isaac, N.J.B., Roy, D.B., Procter, D., Eigenbrod, F., Freckleton, R., Hector, A., Orme, C.D.L., Petchey, O.L., Proença, V., Raffaelli, D., Suttle, K.B., Mace, G.M., Martín-López, B., Woodcock, B.A., Bullock, J.M., 2015. Biodiversity and resilience

- of ecosystem functions. *Trends Ecol. Evol.* 30, 673–684. <https://doi.org/10.1016/j.tree.2015.08.009>.
- Oren, R., Ellsworth, D.S., Johnsen, K.H., Phillips, N., Ewers, B.E., Maier, C., Schäfer, K.V.R., McCarthy, H., Hendrey, G., McNulty, S.G., Katul, G.G., 2001. Soil fertility limits carbon sequestration by forest ecosystems in a CO<sub>2</sub>-enriched atmosphere. *Nature* 411, 469–472. <https://doi.org/10.1038/35078064>.
- Oswalt, S.N., Smith, W.B., Miles, P.D., Pugh, S.A., 2014. *Update of the 2010 RPA Assessment Forest Resources of the United States*. 2012 p. 228.
- Ozgul, A., Childs, D.Z., Oli, M.K., Armitage, K.B., Blumstein, D.T., Olson, L.E., Tuljapourkar, S., Coulson, T., 2010. Coupled dynamics of body mass and population growth in response to environmental change. *Nature* 466, 482–485. <https://doi.org/10.1038/nature09210>.
- Pacifici, M., Visconti, P., Butchart, S.H.M., Watson, J.E.M., Cassola, F.M., Rondinini, C., 2017. Species' traits influenced their response to recent climate change. *Nat. Clim. Chang.* 7, 205–208. <https://doi.org/10.1038/nclimate3223>.
- Pan, Y., Birdsey, R., Hom, J., McCullough, K., 2009. Separating effects of changes in atmospheric composition, climate and land-use on carbon sequestration of U.S. mid-Atlantic temperate forests. *For. Ecol. Manag.* 259, 151–164. <https://doi.org/10.1016/j.foreco.2009.09.049>.
- Patricola, C.M., Wehner, M.F., 2018. Anthropogenic influences on major tropical cyclone events. *Nature* 563, 339–346. <https://doi.org/10.1038/s41586-018-0673-2>.
- Pearson, K.D., 2019. Spring- and fall-flowering species show diverging phenological responses to climate in the Southeast USA. *Int. J. Biometeorol.* 63, 481–492. <https://doi.org/10.1007/s00484-019-01679-0>.
- Peer, A.C., Miller, T.J., 2014. Climate change, migration phenology, and fisheries management interact with unanticipated consequences. *North Am. J. Fish. Manag.* 34, 94–110. <https://doi.org/10.1080/02755947.2013.847877>.
- Peers, M.J.L., Wehrtje, M., Thornton, D.H., Murray, D.L., 2014. Prey switching as a means of enhancing persistence in predators at the trailing southern edge. *Glob. Chang. Biol.* 20, 1126–1135. <https://doi.org/10.1111/gcb.12469>.
- Pelletier, F., Colman, D.W., 2018. Will human influences on evolutionary dynamics in the wild pervade the Anthropocene? *BMC Biol.* 16, 1–10. <https://doi.org/10.1186/s12915-017-0476-1>.
- Pershing, A.J., Alexander, M.A., Hernandez, C.M., Kerr, L.A., Bris, A. Le, Mills, K.E., Nye, J.A., Record, N.R., Scannell, H.A., Scott, J.D., Sherwood, G.D., Thomas, A.C., 2015. Slow adaptation in the face of rapid warming leads to collapse of the Gulf of Maine cod fishery. *Science* (80- ) 350, 350. <https://doi.org/10.1126/science.aae0463>.
- Pershing, A.J., Alexander, M.A., Hernandez, C.M., Kerr, L.A., Bris, A. Le, Mills, K.E., Nye, J.A., Record, N.R., Scannell, H.A., Scott, J.D., Sherwood, G.D., Thomas, A.C., 2016. Response to comments on “slow adaptation in the face of rapid warming leads to collapse of the Gulf of Maine cod fishery”. *Science* (80- ) 352, 423. <https://doi.org/10.1126/science.aae0463>.
- Pershing, A.J., Mills, K.E., Dayton, A.M., Franklin, B.S., Kennedy, B.T., 2018. Evidence for adaptation from the 2016 marine heatwave in the Northwest Atlantic Ocean. *Oceanography* 31, 152–161. <https://doi.org/10.5670/oceanog.2018.213>.
- Peters, M.P., Iverson, L.R., Matthews, S.N., 2015. Long-term droughtiness and drought tolerance of eastern US forests over five decades. *For. Ecol. Manag.* 345, 56–64. <https://doi.org/10.1016/j.foreco.2015.02.022>.
- Peterson, D.L., Halofsky, J.E., 2018. Adapting to the effects of climate change on natural resources in the Blue Mountains, USA. *Clim. Serv.* 10, 63–71. <https://doi.org/10.1016/j.cliser.2017.06.005>.
- Peterson, G.D., Cumming, G.S., Carpenter, S.R., 2003. Scenario planning: a tool for conservation in an uncertain world. *Conserv. Biol.* 17, 358–366. <https://doi.org/10.1046/j.1523-1739.2003.01491.x>.
- Pinsky, M.L., Mantua, N.J., 2014. Emerging adaptation approaches for climate-ready fisheries management. *Oceanography* 27, 147–159. <https://doi.org/10.5670/oceanog.2014.93>.
- Pinsky, M.L., Worm, B., Fogarty, M.J., Sarmiento, J.L., Levin, S.A., 2013. *Marine taxa track local climate velocities*. *Science* (80- ) 341, 1239–1242.
- Pollock, M.M., Beechie, T.J., Wheaton, J.M., Jordan, C.E., Bouwes, N., Weber, N., Volk, C., 2014. Using beaver dams to restore incised stream ecosystems. *Bioscience* 64, 279–290. <https://doi.org/10.1093/biosci/biu036>.
- Poloczanska, E.S., Brown, C.J., Sydeman, W.J., Kiessling, W., Schoeman, D.S., Moore, P.J., Brander, K., Bruno, J.F., Buckley, L.B., Burrows, M.T., Duarte, C.M., Halpern, B.S., Holding, J., Kappel, C.V., O'Connor, M.I., Pandolfi, J.M., Parmesan, C., Schwing, F., Thompson, S.A., Richardson, A.J., 2013. Global imprint of climate change on marine life. *Nat. Clim. Chang.* 3, 919–925. <https://doi.org/10.1038/Nclimate1958>.
- Post, E., 2017. Implications of earlier sea ice melt for phenological cascades in arctic marine food webs. *Food Webs* 13, 60–66. <https://doi.org/10.1016/j.fooweb.2016.11.002>.
- Powell, E.J., Tyrrell, M.C., Milliken, A., Tirpak, J.M., Staudinger, M.D., 2019. A review of coastal management approaches to support the integration of ecological and human community planning for climate change. *J. Coast. Conserv.* 23. <https://doi.org/10.1007/s11852-018-0632-y>.
- Pugh, T.A.M., Lindeskog, M., Smith, B., Poulter, B., Arneth, A., Haverd, V., Calle, L., 2019. Role of forest regrowth in global carbon sink dynamics. *Proc. Natl. Acad. Sci. U. S. A.* 116, 4382–4387. <https://doi.org/10.1073/pnas.1810512116>.
- Pyke, C.R., Thomas, R., Porter, R.D., Hellmann, J.J., Dukes, J.S., Lodge, D.M., Chavarria, G., 2008. Current practices and future opportunities for policy on climate change and invasive species. *Conserv. Biol.* 22, 585–592. <https://doi.org/10.1111/j.1523-1739.2008.00956.x>.
- Radchuk, V., Reed, T., Teplitsky, C., van de Pol, M., Charmantier, A., Hassall, C., Adamík, P., Adriansen, F., Ahola, M.P., Arcese, P., Miguel Avilés, J., Balbontin, J., Berg, K.S., Borrás, A., Burthe, S., Clobert, J., Dehnhard, N., de Lope, F., Dhondt, A.A., Dingemanse, N.J., Doi, H., Eeva, T., Fickel, J., Filella, I., Fossøy, F., Goodenough, A.E., Hall, S.J.G., Hansson, B., Harris, M., Hasselquist, D., Hickler, T., Joshi, J., Kharouba, H., Martínez, J.G., Mihoub, J.-B., Mills, J.A., Molina-Morales, M., Moksnes, A., Ozgul, A., Parejo, D., Pilard, P., Poisbleau, M., Rousset, F., Rödel, M.-O., Scott, D., Senar, J.C., Stefanescu, C., Stokke, B.G., Kusano, T., Tarka, M., Tarwater, C.E., Thonicke, K., Thorley, J., Wilting, A., Tryjanowski, P., Merilä, J., Sheldon, B.C., Pape Møller, A., Matthysen, E., Janzen, F., Dobson, F.S., Visser, M.E., Beissinger, S.R., Courtiol, A., Kramer-Schadt, S., 2019. Adaptive responses of animals to climate change are most likely insufficient. *Nat. Commun.* 10, 1–14. <https://doi.org/10.1038/s41467-019-10924-4>.
- Ralston, J., DeLuca, W.V., Feldman, R.E., King, D.L., 2017. Population trends influence species ability to track climate change. *Glob. Chang. Biol.* 23, 1390–1399. <https://doi.org/10.1111/gcb.13478>.
- Rapp, J.M., Lutz, D.A., Huish, R.D., Dufour, B., Ahmed, S., Morelli, T.L., Stinson, K.A., 2019. Finding the sweet spot: shifting optimal climate for maple syrup production in North America. *For. Ecol. Manag.* 448, 187–197. <https://doi.org/10.1016/j.foreco.2019.05.045>.
- Rasmann, S., Pellissier, L., Defosse, E., Jactel, H., Kunstler, G., 2014. Climate-driven change in plant-insect interactions along elevation gradients. *Funct. Ecol.* 28, 46–54. <https://doi.org/10.1111/1365-2435.12135>.
- Rodriguez, A.B., Fodrie, F.J., Ridge, J.T., Lindquist, N.L., Theuerkauf, E.J., Coleman, S.E., Grabowski, J.H., Brodeur, M.C., Gittman, R.K., Keller, D.A., Kenworthy, M.D., 2014. Oyster reefs can outpace sea-level rise. *Nat. Clim. Chang.* 4, 493–497. <https://doi.org/10.1038/nclimate2216>.
- Rosenblatt, A.E., Schmitz, O.J., 2014. Interactive effects of multiple climate change variables on trophic interactions: a meta-analysis. *Clim. Chang. Responses* 1, 8. <https://doi.org/10.1186/s40665-014-0008-y>.
- Rowe, K.C., Rowe, K.M.C., Tingley, M.W., Koo, M.S., 2015. Spatially heterogeneous impact of climate change on small mammals of Montane California. *Proc. R. Soc. London B Biol. Sci.* 282, 20141857. <https://doi.org/10.1098/rspb.2014.1857>.
- Rubenstein, M.A., Christophersen, R., Ransom, J.L., 2019. Trophic implications of a phenological paradigm shift: bald eagles and salmon in a changing climate. *J. Appl. Ecol.* 56, 769–778. <https://doi.org/10.1111/1365-2664.13286>.
- Rudolf, V.H.W., 2019. The role of seasonal timing and phenological shifts for species coexistence. *Ecol. Lett.* 22, 1324–1338. <https://doi.org/10.1111/ele.13277>.
- Runting, R.K., Bryan, B.A., Dee, L.E., Maseyk, F.J.F., Mandle, L., Hamel, P., Wilson, K.A., Yetka, K., Possingham, H.P., Rhodes, J.R., 2017. Incorporating climate change into ecosystem service assessments and decisions: a review supplementary materials. *Glob. Chang. Biol.* 23, 28–41. <https://doi.org/10.1111/gcb.13457>.
- Sandifer, P.A., Sutton-Grier, A.E., 2014. Connecting stressors, ocean ecosystem services, and human health. *Nat. Resour. Forum* 38, 157–167. <https://doi.org/10.1111/1477-8947.12047>.
- Sandifer, P.A., Sutton-Grier, A.E., Ward, B.P., 2015. Exploring connections among nature, biodiversity, ecosystem services, and human health and well-being: opportunities to enhance health and biodiversity conservation. *Ecosyst. Serv.* 12, 1–15. <https://doi.org/10.1016/j.ecoser.2014.12.007>.
- Schindler, D.E., Rogers, D.E., Scheuerell, M.D., Abrey, C.A., 2005. Effect of changing climate on zooplankton and juvenile sockeye salmon growth in Southwestern Alaska. *Ecology* 86, 198–209. <https://doi.org/10.1890/03-0408>.
- Schlaepfer, M.A., Runge, M.C., Sherman, P.W., Sherman, P.W., 2002. Ecological and evolutionary traps. *Trends Ecol. Evol.* 17, 474–480. [https://doi.org/10.1016/0197-3975\(91\)90016-E](https://doi.org/10.1016/0197-3975(91)90016-E).
- Schluter, D., Price, T.D., Rowe, L., 1991. Conflicting selection pressures and life history trade-offs. *Proc. R. Soc. B Biol. Sci.* 246, 11–17. <https://doi.org/10.1098/rspb.1991.0118>.
- Schmitt, J.D., Peoples, B.K., Castello, L., Orth, D.J., 2019. Feeding ecology of generalist consumers: a case study of invasive blue catfish *Ictalurus furcatus* in Chesapeake Bay, Virginia, USA. *Environ. Biol. Fish.* 102, 443–465. <https://doi.org/10.1007/s10641-018-0783-6>.
- Schwartz, M.W., Hellmann, J.J., McLachlan, J.M., Sax, D.F., Borevitz, J.O., Brennan, J., Camacho, A.E., Ceballos, G., Clark, J.R., Doremus, H., Early, R., Etterson, J.R., Fielder, D., Gill, J.L., Gonzalez, P., Green, N., Hannah, L., Jamieson, D.W., Javeline, D., Minter, B.A., Odenbaugh, J., Polasky, S., Richardson, D.M., Root, T.L., Safford, H.D., Sala, O., Schneider, S.H., Thompson, A.R., Williams, J.W., Vellend, M., Vitt, P., Zellmer, S., 2012. Managed relocation: integrating the scientific, regulatory, and ethical challenges. *Bioscience* 62, 732–743. <https://doi.org/10.1525/bio.2012.62.8.6>.
- Sgrò, C.M., Lowe, A.J., Hoffmann, A.A., 2011. Building evolutionary resilience for conserving biodiversity under climate change. *Evol. Appl.* 4, 326–337. <https://doi.org/10.1111/j.1752-4571.2010.00157.x>.
- Shen, G., Aydin, S.G., 2014. Highway freight transportation disruptions under an extreme environmental event: the case of Hurricane Katrina. *Int. J. Environ. Sci. Technol.* 11, 2387–2402.
- Shilling, D., Nelson, M. (Eds.), 2018. *Traditional Ecological Knowledge: Learning From Indigenous Methods for Environmental Sustainability*. Cambridge University Press, Cambridge, UK.
- Siedlecki, S.A., Kaplan, I.C., Hermann, A.J., Nguyen, T.T., Bond, N.A., Newton, J.A., Williams, G.D., Peterson, W.T., Alin, S.R., Feely, R.A., 2016. Experiments with seasonal forecasts of ocean conditions for the Northern region of the California current upwelling system. *Sci. Rep.* 6, 1–18. <https://doi.org/10.1038/srep27203>.
- Sirami, C., Caplat, P., Popy, S., Clamens, A., Arlettaz, R., Jiguet, F., Brotons, L., Martin, J.L., 2017. Impacts of global change on species distributions: obstacles and solutions to integrate climate and land use. *Glob. Ecol. Biogeogr.* 26, 385–394. <https://doi.org/10.1111/geb.12555>.
- Snell-Rood, E.C., Kobiela, M.E., Sikkink, K.L., Shephard, A.M., 2018. Mechanisms of plastic rescue in novel environments. *Annu. Rev. Ecol. Syst.* 49, 331–354. <https://doi.org/10.1146/annurev-ecolsys-110617-062622>.
- Sohn, J.A., Saha, S., Bauhus, J., 2016. Potential of forest thinning to mitigate drought stress: a meta-analysis. *For. Ecol. Manag.* 380, 261–273. <https://doi.org/10.1016/j.foreco.2016.07.046>.

- Spalding, M.D., Ruffo, S., Lacambra, C., Meliane, I., Hale, L.Z., Shepard, C.C., Beck, M.W., 2014. The role of ecosystems in coastal protection: adapting to climate change and coastal hazards. *Ocean Coast. Manag.* 90, 50–57. <https://doi.org/10.1016/j.ocecoaman.2013.09.007>.
- Spencer, P.D., Nelson, M.W., Hollowed, A.B., Sigler, M.F., Hermann, A.J., 2019. Trait - based climate vulnerability assessments in data - rich systems : an application to eastern Bering Sea fish and invertebrate stocks. *Glob. Chang. Biol.*, 1–18 <https://doi.org/10.1111/gcb.14763>.
- Staudinger, M.D., Grimm, N.B., Staudt, A., Carter, S.L., Chapin, I.I.F.S., Kareiva, P., Ruckelshaus, M., Stein, B.A., 2012. *Impacts of Climate Change on Biodiversity, Ecosystems, and Ecosystem Services: Technical Input to the 2013 National Climate Assessment*. Coop. Rep. To 2013 Natl. Clim. Assess (296 p).
- Staudinger, M.D., Morelli, T.L., Bryan, A.M., 2015. *Integrating Climate Change Into Northeast and Midwest State Wildlife Action Plans* (Amherst, MA).
- Staudinger, M.D., Mills, K.E., Stamieszkin, K., Record, N.R., Hudak, C.A., Allyn, A., Diamond, A., Friedland, K.D., Golet, W., Henderson, M.E., Hernandez, C.M., Huntington, T.E., Ji, R., Johnson, C.L., Johnson, D.S., Jordaan, A., Kocik, J., Li, Y., Liebman, M., Nichols, O.C., Pendleton, D., Richards, R.A., Robben, T., Thomas, A.C., Walsh, H.J., Yakola, K., 2019. It's about time: a synthesis of changing phenology in the Gulf of Maine ecosystem. *Fish. Oceanogr.*, 532–566 <https://doi.org/10.1111/fog.12429>.
- Stein, B.A., Staudt, A., Cross, M.S., Dubois, N.S., Enquist, C., Griffis, R., Hansen, L.J., Hellmann, J.J., Lawler, J.J., Nelson, E.J., Pairis, A., 2013. Preparing for and managing change: climate adaptation for biodiversity and ecosystems. *Front. Ecol. Environ.* 11, 502–510. <https://doi.org/10.1890/120277>.
- Stein, B.A., Glick, P., Edelson, N., Staudt, A., 2014. *Climate-smart Conservation: Putting Adaptation Principles Into Practice*. National Wildlife Federation, Washington, D.C.
- Stephens, S.L., Collins, B.M., Fetting, C.J., Finney, M.A., Hoffman, C.M., Knapp, E.E., North, M.P., Safford, H., Wayman, R.B., 2018. Drought, tree mortality, and wildfire in forests adapted to frequent fire. *Bioscience* 68, 77–88. <https://doi.org/10.1093/biosci/bix146>.
- Stewart, I.T., Cayan, D.R., Dettinger, M.D., 2005. Changes toward earlier streamflow timing across western North Am. *J. Clim.* 18, 10.
- Stock, C.A., Dunne, J.P., John, J.G., 2014. Drivers of trophic amplification of ocean productivity trends in a changing climate. *Biogeosciences* 11, 7125–7135. <https://doi.org/10.5194/bg-11-7125-2014>.
- Sundby, S., Drinkwater, K.F., Kjesbu, O.S., 2016. The North Atlantic spring-bloom system—where the changing climate meets the winter dark. *Front. Mar. Sci.* 3, 28.
- Sutton-Grier, A.E., Wowk, K., Bamford, H., 2015. Future of our coasts: the potential for natural and hybrid infrastructure to enhance the resilience of our coastal communities, economies and ecosystems. *Environ. Sci. Pol.* 51, 137–148. <https://doi.org/10.1016/j.envsci.2015.04.006>.
- Sutton-Grier, A.E., Gittman, R.K., Arkema, K.K., Bennett, R.O., Benoit, J., Blitch, S., Burks-Copes, K.A., Colden, A., Dausman, A., DeAngelis, B.M., Hughes, A.R., Scyphers, S.B., Grabowski, J.H., 2018. Investing in natural and nature-based infrastructure: building better along our coasts. *Sustain* 10, 1–11. <https://doi.org/10.3390/su10020523>.
- Swanston, C., Janowiak, M. (Eds.), 2012. *Forest Adaptation Resources : Climate Change Tools and Approaches for Land Managers*. U.S. Department of Agriculture, Forest Service, Northern Research Station, Newtown Square, PA Gen. Tech. Rep. NRS-87.
- Taylor, C., Caplat, P., Massimino, D., Johnston, A., Jonzén, N., Smith, H.G., Lindström, Å., 2015. Swedish birds are tracking temperature but not rainfall: evidence from a decade of abundance changes. *Glob. Ecol. Biogeogr.* 24, 859–872. <https://doi.org/10.1111/geb.12308>.
- Taylor, P.C., Maslowski, W., Perlwitz, J., Wuebbles, D.J., 2017. Arctic changes and their effects on Alaska and the rest of the United States. *Fourth National Climate Assessment*. I, pp. 303–332. <https://doi.org/10.7930/J00863GK>.
- Thomas, A.C., Pershing, A.J., Friedland, K.D., Nye, J.A., Mills, K.E., Alexander, M.A., Record, N.R., Weatherbee, R., Henderson, M.E., 2017. Seasonal trends and phenology shifts in sea surface temperature on the North American northeastern continental shelf. *Elem. Sci. Anth* 5.
- Timpane-Padgham, B.L., Beechie, T., Klinger, T., 2017. A systematic review of ecological attributes that confer resilience to climate change in environmental restoration. *PLoS One* 12, 1–23. <https://doi.org/10.1371/journal.pone.0173812>.
- Tommasi, D., Stock, C.A., Hobday, A.J., Methot, R., Kaplan, I.C., Evson, J.P., Holsman, K., Miller, T.J., Gaichas, S., Gehlen, M., Pershing, A., Vecchi, G.A., Msadek, R., Delworth, T., Eakin, C.M., Haltuch, M.A., Séférian, R., Spillman, C.M., Hartog, J.R., Siedlecki, S., Samhuri, J.F., Muhling, B., Asch, R.G., Pinsky, M.L., Saba, V.S., Kapnick, S.B., Gaitan, C.F., Rykaczewski, R.R., Alexander, M.A., Xue, Y., Pegion, K.V., Lynch, P., Payne, M.R., Kristiansen, T., Lehodey, P., Werner, F.E., 2017. Managing living marine resources in a dynamic environment: the role of seasonal to decadal climate forecasts. *Prog. Oceanogr.* 152, 15–49. <https://doi.org/10.1016/j.pocan.2016.12.011>.
- Trosper, R.L., 2009. *Resilience, Reciprocity and Ecological Economics: Northwest Coast Sustainability*. Routledge, New York, New York.
- Trottier, N., Groeneveld, E., Lavoie, C., 2017. Giant hogweed at its northern distribution limit in North America: experiments for a better understanding of its dispersal dynamics along rivers. *River Res. Appl.* 33, 1098–1106. <https://doi.org/10.1002/rra.3149>.
- Tyrrell, T., 2019. Redfield Ratio. *Encycl. Ocean Sci.*, pp. 461–472. <https://doi.org/10.1016/B978-0-12-409548-9.11281-3>.
- U.S. Environmental Protection Agency, 2016. *Climate Change Indicators in the United States*. Fourth edition. (doi:EPA 430-R-16-004).
- U.S. Fish and Wildlife Service, 2008. *Endangered and threatened wildlife and plants; determination of threatened status for the polar bear (Ursus maritimus) throughout its range*. *Fed. Regist.* 73, 28211–28303.
- U.S. Fish and Wildlife Service, 2010. *12-Month finding on a petition to list the American Pika as threatened or endangered*. *Fed. Regist.* 75, 6437–6471.
- USDA, 2016. *USDA Integrated Projections for Agriculture and Forest Sector Land Use Land-use Change, and GHG Emissions and Removals: 2015 to 2060* (Washington, D.C.).
- USFWS and NMFS, 2019. *Recovery Plan for the Gulf of Maine Distinct Population Segment of Atlantic Salmon (Salmo salar)*.
- USGCRP, 2018. *Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment. II. U.S. Global Change Research Program*, Washington, D.C., USA.
- Valéry, L., Fritz, H., Lefeuve, J.C., Simberloff, D., 2009. Invasive species can also be native.... *Trends Ecol. Evol.* 24, 585. <https://doi.org/10.1016/j.tree.2009.06.017>.
- Van Zuiden, T.M., Chen, M.M., Stefanoff, S., Lopez, L., Sharma, S., 2016. Projected impacts of climate change on three freshwater fishes and potential novel competitive interactions. *Divers. Distrib.* 22, 603–614. <https://doi.org/10.1111/ddi.12422>.
- Verdeny-Vilalta, O., Moya-Laraño, J., 2014. Seeking water while avoiding predators: moisture gradients can affect predator-prey interactions. *Anim. Behav.* 90, 101–108. <https://doi.org/10.1016/j.anbehav.2014.01.027>.
- Vergés, A., Steinberg, P.D., Hay, M.E., Poore, A.G.B., Campbell, A.H., Ballesteros, E., Heck, K.L., Booth, D.J., Coleman, M.A., Feary, D.A., Figueira, W., Langlois, T., Marzinelli, E.M., Mizerek, T., Mumby, P.J., Nakamura, Y., Roughan, M., van Sebille, E., Gupta, A. Sen, Smale, D.A., Tomas, F., Wernberg, T., Wilson, S.K., 2014. The tropicalization of temperate marine ecosystems: climate-mediated changes in herbivory and community phase shifts. *Proc. R. Soc. B Biol. Sci.* 281 <https://doi.org/10.1098/rspb.2014.0846>.
- Vergés, A., Doropoulos, C., Malcolm, H.A., Skye, M., Garcia-Pizá, M., Marzinelli, E.M., Campbell, A.H., Ballesteros, E., Hoey, A.S., Vila-Concejo, A., Bozec, Y.-M., Steinberg, P.D., 2016. Long-term empirical evidence of ocean warming leading to tropicalization of fish communities, increased herbivory, and loss of kelp. *Proc. Natl. Acad. Sci.* 113, 13791–13796. <https://doi.org/10.1073/pnas.1610725113>.
- Vernon, M.J., Sherriff, R.L., van Mantgem, P., Kane, J.M., 2018. Thinning, tree-growth, and resistance to multi-year drought in a mixed-conifer forest of northern California. *For. Ecol. Manag.* 422, 190–198. <https://doi.org/10.1016/j.foreco.2018.03.043>.
- Viers, J.H., Purdy, S.E., Peek, R.A., Fryjoff-Hung, A., Santos, N.R., Katz, J.V.E., Emmons, J.D., Dolan, D.V., Yarnell, S.M., 2013. *Montane meadows in the Sierra Nevada*. *Cent. Watershed Sci. Tech. Rep.* 63.
- Vinyeta, K., Whyte, K.P., Lynn, K., 2015. *Climate change through an intersectional lens: gendered vulnerability and resilience in indigenous communities in the United States*. *USDA For. Serv. - Gen. Tech. Rep. PNW-GTR-923*, pp. 1–74.
- Visser, M.E., Gienapp, P., 2019. Evolutionary and demographic consequences of phenological mismatches. *Nat. Ecol. Evol.* 3, 879–885.
- Wallingford, P.D., Morelli, T.L., Allen, J.M., Beaury, E.M., Blumenthal, D.M., Bradley, B.A., Dukes, J.S., Early, R., Fusco, E.J., Goldberg, D.E., Ibáñez, I., Laginhas, B.B., Vilá, M., Sorte, C.J.B., n.d. Assessing impacts of climate-driven range shifts through the lens of invasion biology. *Nat. Clim. Chang. Chang.* (In review).
- Walsh, H.J., Richardson, D.E., Marancik, K.E., Hare, J.A., 2015. Long-term changes in the distributions of larval and adult fish in the northeast U.S. shelf ecosystem. *PLoS One* 10, 1–31. <https://doi.org/10.1371/journal.pone.0137382>.
- Walter, J., Jentsch, A., Beierkuhnlein, C., Kreyling, J., 2013. Ecological stress memory and cross stress tolerance in plants in the face of climate extremes. *Environ. Exp. Bot.* 94, 3–8. <https://doi.org/10.1016/j.envexpbot.2012.02.009>.
- Wang, W.J., He, H.S., Thompson, F.R., Fraser, J.S., Dijk, W.D., 2017. Changes in forest biomass and tree species distribution under climate change in the northeastern United States. *Landsc. Ecol.* 32, 1399–1413. <https://doi.org/10.1007/s10980-016-0429-z>.
- Wann, G.T., Aldridge, C.L., Seglund, A.E., Oyler-McCance, S.J., Kondratieff, B.C., Braun, C.E., 2019. Mismatches between breeding phenology and resource abundance of resident alpine ptarmigan negatively affect chick survival. *Ecol. Evol.*, 7200–7212 <https://doi.org/10.1002/ecs3.5290>.
- Waples, R.S., Audzijonyte, A., 2016. Fishery-induced evolution provides insights into adaptive responses of marine species to climate change. *Front. Ecol. Environ.* 14, 217–224. <https://doi.org/10.1002/fee.1264>.
- Warziniack, T.W., Elmer, M.J., Miller, C.J., Dante-Wood, S.K., Woodall, C.W., Nichols, M.C., Domke, G.M., Stockmann, K.D., Proctor, J.G., Borchers, A., 2018. *Effects of climate change on ecosystem services*. In: Halofsky, J.E., Peterson, D.L., Ho, J., Little, N.J., Joyce, L.A. (Eds.), *Climate Change Vulnerability and Adaptation in the Intermountain Region [Part 2]*. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fort Collins, CO, pp. 376–403.
- Wasmund, N., Nausch, G., Gerth, M., Busch, S., Burmeister, C., Hansen, R., Sadkowiak, B., 2019. Extension of the growing season of phytoplankton in the western Baltic Sea in response to climate change. *Mar. Ecol. Prog. Ser.* 622, 1–16. <https://doi.org/10.3354/meps12994>.
- Wear, D.N., Coulston, J.W., 2015. From sink to source: regional variation in U.S. forest carbon futures. *Sci. Rep.* 5, 1–11. <https://doi.org/10.1038/srep16518>.
- Weed, A.S., Ayres, M.P., Hicke, J.A., 2013. Consequences of climate change for biotic disturbances in North American forests. *Ecol. Monogr.* 83, 441–470. <https://doi.org/10.1890/13-0160.1>.
- Weeks, B.C., Willard, D.E., Ellis, A.A., Witynski, M.L., Winger, B.M., 2019. Shared morphological consequences of global warming in North American migratory birds. *Ecol. Lett.* <https://doi.org/10.1111/1365-3113.10329>.
- Wenger, S.J., Isaak, D.J., Luce, C.H., Neville, H.M., Fausch, K.D., Dunham, J.B., Dauwalter, D.C., Young, M.K., Elsner, M.M., Rieman, B.E., Hamlet, A.F., Williams, J.E., 2011. Flow regime, temperature, and biotic interactions drive differential declines of trout species under climate change. *Proc. Natl. Acad. Sci. U. S. A.* 108, 14175–14180. <https://doi.org/10.1073/pnas.1103097108>.
- Wenzel, S., Cox, P.M., Eyring, V., Friedlingstein, P., 2016. Projected land photosynthesis constrained by changes in the seasonal cycle of atmospheric CO<sub>2</sub>. *Nature* 538, 499–501. <https://doi.org/10.1038/nature19772>.
- West, J.M., Julius, S.H., Kareiva, P., Enquist, C., Lawler, J.J., Petersen, B., Johnson, A.E., Shaw, M.R., 2009. U.S. natural resources and climate change: concepts and approaches for

- management adaptation. *Environ. Manag.* 44, 1001–1021. <https://doi.org/10.1007/s00267-009-9345-1>.
- Westerling, A.L., 2016. Increasing western US forest wildfire activity: sensitivity to changes in the timing of spring. *Philos. Trans. R. Soc. B Biol. Sci.*, 371 <https://doi.org/10.1098/rstb.2015.0178>.
- Westerling, A.L., Hidalgo, H.G., Cayan, D.R., Swetnam, T.W., 2006. Warming and earlier spring increase western U.S. forest wildfire activity. *Science* (80-. ) 313, 940–943. <https://doi.org/10.1126/science.1128834>.
- Whiteley, A.R., Fitzpatrick, S.W., Funk, W.C., Tallmon, D.A., 2015. Genetic rescue to the rescue. *Trends Ecol. Evol.* 30, 42–49. <https://doi.org/10.1016/j.tree.2014.10.009>.
- Whyte, K.P., Dockry, M.J., Baule, W., Fellman, D., 2014. Supporting tribal climate change adaptation planning through community participatory strategic foresight scenario development. In: Brown, D., Baule, W., Briley, L., Gibbons, E. (Eds.), *Project Reports. Great Lakes Integrated Sciences and Assessments Center, Ann Arbor, MI*.
- Wiens, J.J., 2016. Climate-related local extinctions are already widespread among plant and animal species. *PLoS Biol.* 14. <https://doi.org/10.1371/journal.pbio.2001104>.
- Yeruham, E., Shpigel, M., Abelson, A., Rilov, G., 2020. Ocean warming and tropical invaders erode the performance of a key herbivore. *Ecology* 101, 1–13. <https://doi.org/10.1002/ecy.2925>.
- Young, K.R., 2014. Biogeography of the Anthropocene: novel species assemblages. *Prog. Phys. Geogr.* 38, 664–673. <https://doi.org/10.1177/0309133314540930>.
- Zeidberg, L.D., Robison, B.H., 2007. Invasive range expansion by the Humboldt squid, *Dosidicus gigas*, in the eastern North Pacific. *Proc. Natl. Acad. Sci. U. S. A.* 104, 12948–12950. <https://doi.org/10.1073/pnas.0702043104>.
- Zhang, K., Liu, H., Li, Y., Xu, H., Shen, J., Rhome, J., Smith, T.J., 2012. The role of mangroves in attenuating storm surges. *Estuar. Coast. Shelf Sci.* 102–103, 11–23. <https://doi.org/10.1016/j.ecss.2012.02.021>.
- Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D.B., Huang, Y., Huang, M., Yao, Y., Bassu, S., Ciais, P., Durand, J.L., Elliott, J., Ewert, F., Janssens, I.A., Li, T., Lin, E., Liu, Q., Martre, P., Müller, C., Peng, S., Peñuelas, J., Ruane, A.C., Wallach, D., Wang, T., Wu, D., Liu, Z., Zhu, Y., Zhu, Z., Asseng, S., 2017. Temperature increase reduces global yields of major crops in four independent estimates. *Proc. Natl. Acad. Sci. U. S. A.* 114, 9326–9331. <https://doi.org/10.1073/pnas.1701762114>.
- Zhu, Z., Piao, S., Myneni, R.B., Huang, M., Zeng, Z., Canadell, J.G., Ciais, P., Sitch, S., Friedlingstein, P., Arneeth, A., Cao, C., Cheng, L., Kato, E., Koven, C., Li, Y., Lian, X., Liu, Y., Liu, R., Mao, J., Pan, Y., Peng, S., Peñuelas, J., Poulter, B., Pugh, T.A.M., Stocker, B.D., Viovy, N., Wang, X., Wang, Y., Xiao, Z., Yang, H., Zaehle, S., Zeng, N., 2016. Greening of the Earth and its drivers. *Nat. Clim. Chang.* 6, 791–795. <https://doi.org/10.1038/nclimate3004>.
- Zimova, M., Mills, L.S., Nowak, J.J., 2016. High fitness costs of climate change induced camouflage mismatch in a seasonally colour moulting mammal. *Ecol. Lett.* 19, 299–307. <https://doi.org/10.1111/ele.12568>.