Chapter 12
Fire and Forests in the 21st Century: Managing Resilience Under Changing Climates and Fire Regimes in USA Forests

James M. Vose, David L. Peterson, Christopher J. Fettig, Jessica E. Halofsky, J. Kevin Hiers, Robert E. Keane, Rachel Loehman, and Michael C. Stambaugh

Abstract Higher temperatures, lower snowpacks, drought, and extended dry periods have contributed to increased wildfire activity in recent decades. Climate change is expected to increase the frequency of large fires, the cumulative area burned, and fire suppression costs and risks in many areas of the USA. Fire regimes are likely to change due to interactions among climate, fire, and other stressors and disturbances; resulting in persistent changes in forest structure and function. The remainder of the twenty-first century will present substantial challenges, as natural resource manag-

J. M. Vose (✉)
USDA Forest Service, Southern Research Station, Research Triangle Park, NC, USA
e-mail: jmvose@ncsu.edu

D. L. Peterson
School of Environmental and Forest Sciences, University of Washington, Seattle, WA, USA

C. J. Fettig
USDA Forest Service, Pacific Southwest Research Station, Ecosystem Function and Health Program, Davis, CA, USA

J. E. Halofsky
Western Wildlands Environmental Threat Assessment Center, Corvallis, WA, USA

USDA Forest Service, Pacific Northwest Research Station, Olympia, WA, USA

J. K. Hiers
Tall Timbers Research Station, Tallahassee, FL, USA

R. E. Keane
USDA Forest Service, Rocky Mountain Research Station (retired), Missoula, MT, USA

R. Loehman
US Geological Survey, Alaska Science Center, Anchorage, AK, USA

M. C. Stambaugh
School of Natural Resources, University of Missouri, Columbia, MO, USA

© This is a U.S. government work and not under copyright protection in the U.S.; foreign copyright protection may apply 2021

C. H. Greenberg, B. Collins (eds.), *Fire Ecology and Management: Past, Present, and Future of US Forested Ecosystems*, Managing Forest Ecosystems 39, https://doi.org/10.1007/978-3-030-73267-7_12

ers are faced with higher fire risk and the difficult task of maintaining ecological function in a rapidly changing biophysical and social landscape. Fuel treatments will continue to be important for minimizing the undesirable ecological effects of fire, and for enhancing firefighter safety; however, treatments must be implemented strategically across large areas. Collaboration among agencies, private landowners, and other organizations will be critical for ensuring resilience and sustainable forest management.

Keywords Wildfire risk · Prescribed fire · Ecological function · Drought · Stress complexes · Adaptation · Fuels treatment · Future range and variation

12.1 Introduction

Fire is perhaps the most influential natural and anthropogenic disturbance affecting the distribution, structure, and function of terrestrial ecosystems around the world (Bond and Keeley 2005; Bowman et al. 2009; Krawchuk et al. 2009; Scott et al. 2014). Many plants and animals depend on fire for their continued existence; those that do not, such as rainforest and tundra plants, are largely intolerant of burning (DeBano et al. 1998; Cochrane 2010). Spatial and temporal patterns of fire and resulting ecosystem effects, termed "fire regimes," are governed by complex interactions among climate, fuels, vegetation, and ignition pattern and frequency across multiple scales (Agee 1998; Bradstock et al. 2012; Keane 2013; Abatzoglou and Williams 2016; Abatzoglou et al. 2017) (Fig. 12.1). A sound understanding of the interactions between biotic and abiotic ecosystem elements of fire regimes, and the effects of fire regimes on structure and function, facilitates projections of the effects of climate change on fire as an ecological process (Agee 1993; Agee 1997; Mitchell et al. 2014; Scott et al. 2014; Halofsky et al. 2020).

Higher temperatures, lower snowpacks, drought, and extended dry periods have contributed to increased wildfire activity in recent decades, particularly in western USA forests (Freeborn et al. 2016; Holden et al. 2018; Vose et al. 2018). Drier and warmer conditions have also contributed to a longer fire season in some areas (Riley and Loehman 2016). For example, the duration of the wildfire season has increased by 80 days in some parts of the western USA as a result of increased temperature (Jolly et al. 2015; Westerling 2016). A longer burning season combined with dry fuels will promote larger (and longer-duration) wildfires relative to historical wildfire activity (Riley and Loehman 2016). Earlier onset of snowmelt in spring may also contribute to lower fuel moistures at higher elevations (McKenzie et al. 2004). Warmer and drier conditions (annual, intra-annual, and interannual) can also alter vegetation and fuel characteristics (i.e., flammability and spatial distribution) (Flannigan and Van Wagner 1991; Keane et al. 2018; Syphard et al. 2018; Hessburg et al. 2019). For example, wet periods preceding dry periods can increase fuels in dry years in the American Southwest (Grissino-Mayer and Swetnam 2000). Hence,



Fig. 12.1 An example of the complex interactions of a small set of ecosystem disturbances on vegetation with feedbacks. Although climate directly influences vegetation, most climate-mediated changes to forested landscapes will result from these interactions

based on recent observations over the past few decades and model projections for the coming century, we anticipate that climate change will increase fire frequency and annual area burned compared with historical wildfire patterns in many areas of the USA (e.g., Bachelet et al. 2000; Krawchuk et al. 2009, Liu et al. 2010).

The remainder of the twenty-first century is likely to present substantial challenges for natural resource managers as they deal with the task of maintaining ecological function and increasing resilience in a rapidly changing biophysical and social landscape. For example, changing drought regimes will make it more difficult to control wildfire and to use prescribed fire as a management tool (Mitchell et al. 2014; Kupfer et al. 2020). Options to increase resilience include activities that address engineering resilience (recovery to a previous condition), ecological resilience (remaining within the prevailing system domain through maintaining important ecosystem processes and functions or shifting to an alternative ecological domain), and socio-ecological resilience (the capacity to reorganize and adapt through multi-scale interactions between social and ecological components of the system) (Seidl et al. 2016). Some approaches for achieving resilience focus on maintaining or re-establishing historical conditions (Keane et al. 2018), but changing biophysical conditions and disturbance regimes may make it difficult to maintain ecosystem structure and function in emerging, no-analog environments (Falk et al. 2019).

In this chapter, we examine how climate change could affect the structure and function of forest ecosystems in the USA through changes in fire regimes. We discuss (1) observed and anticipated changes in climate; (2) the effects of climate change on fire regimes; (3) the effects of changing fire regimes on ecosystem structure and function; (4) interactions of future fire regimes with other stressors and

disturbances, and; (5) management options that can increase resilience to future fire regimes. We synthesize existing science and discuss how observed responses and model projections can be used to inform natural resource management and planning.

12.2 The Fire Regime

Fire *regimes* are often described by (1) type of fire (ground, surface, crown); (2) mean and variance in fire frequency, intensity, severity, and seasonality, and; (3) areal extent and pattern of a burn (Agee 1993; Keane 2016; Bond and Keane 2017). Fire frequency describes how often a fire burns in an area, commonly measured by mean fire-return interval (FRI) or fire rotation period associated with a particular landscape (Moritz et al. 2005; Falk et al. 2011). Fire *severity* is a general term describing the effects of a fire on an ecosystem (Morgan et al. 2014) that can be estimated from the amount of plant biomass consumed and/or killed (Keeley 2006). It is often confused with fire *intensity*, which is a physical measure that describes the energy released from a fire.

Fire severity in forests differs greatly across a given landscape, depending on scale, weather conditions, and the prefire condition and composition of live and dead biomass. Although the extent, pattern, and shifting mosaic of fire severity can be complex, fire regimes are commonly classified with percentages of (1) low-severity; (2) mixed-severity, and; (3) replacement- (high) severity fires (in reference to vegetation). In many ecosystems (e.g., dry forests), low-severity fires were historically frequent, comprising surface fires with low intensities and causing low levels of mortality (<10%) of dominant vegetation. Replacement-severity fires kill the majority of the dominant vegetation (grass, shrubs, or trees) (Brown 1995) regardless of fire type (Agee 1993). Mixed-severity fires contain elements of both low-severity and replacement-severity fires (Arno et al. 2000; Perry et al. 2011).

Fire regimes are often characterized as having proportions of low-, mixed-, and replacement-severity events. Passive crown fires, patchy stand-replacement fires, and low-intensity understory burns are common in mixed-severity fire regimes (Marcoux et al. 2015). Typically, mixed-severity is used to describe patterns of patchy burn-severity in an area created during one fire event. However, mixed-severity fire regimes can also describe mixed-severity fires over time (e.g., nonlethal surface fire followed by stand-replacement fire) (Shinneman and Baker 1997). Groundfires burn within surface and subsurface organic layers, typically occurring in forests with high soil organic matter, such as boreal forests (Chap. 7), pocosin swamps of the eastern USA (Chap. 3), and dry forests where fire suppression or exclusion has led to uncharacteristically high accumulations of litter and soil organic matter (Allen et al. 2002; Turetsky and Louis 2006).

Fire regimes are created by the interactions of bottom-up and top-down controls (Heyerdahl et al. 2001). Bottom-up influences, such as vegetation, wildland fuels, topography, and patch distributions dictate fire spread, intensity, and severity at fine scales (Skinner and Chang 1996). Some bottom-up controls can be manipulated

through land management activities (Keane 2015). Top-down controls are mostly driven by climate and weather that dictate fire frequency, duration, and synchrony (Swetnam 1990; Guyette et al. 2012). These top-down controls constrain the extent to which bottom-up controls, such as vegetation and fuel loading, can be manipulated to alter fire regimes.

Climate and weather trends are often embedded in atmospheric teleconnections interacting with fire regimes (Swetnam and Betancourt 1990; Duffy et al. 2005). For example, drought-induced wildfires have been associated with global circulation anomalies, such as the El Niño-Southern Oscillation (Veblen et al. 2000; Chen et al. 2011) and Pacific Decadal Oscillation (Heyerdahl et al. 2002). As a result, a fire regime is actually an aggregation of spatial disturbances that does not follow discrete mapping units. Attempts to characterize fire regimes solely from past fires (Westerling et al. 2006), vegetation (Frost 1998), fuels (Olson 1981), or topography (Keane et al. 2004) have only partially succeeded, primarily because they failed to account for the complex spatial and temporal relationships of fire and the multiscale interactions that control fire dynamics (Morgan et al. 2014) (Fig. 12.1).

The role of ignition in fire regimes is often misunderstood (Balch et al. 2017). Lightning ignitions are distributed somewhat randomly across landscapes over long temporal scales (Barrows et al. 1977; Fuquay 1980; Balch et al. 2017), whereas human ignitions are associated with locations where human activity is present (Balch et al. 2017; Keeley and Syphard 2018). Lightning ignitions in moist, productive ecosystems can be constrained by fuel moisture (Barrows et al. 1977; Fuquay 1980). However, in these same ecosystems, human ignitions can occur year-round, potentially causing fires during short periods of dry weather (e.g., Chap. 1, Fig. 1.2, Fig. 1.4, Table 1.1).

Landscape fragmentation due to human development can also reduce fire frequencies in some fire-prone ecosystems (Gill and Williams 1996). For example, land abandonment in European countries has led to successional changes producing contiguous, highly flammable vegetation (Moreira et al. 2001; Moreno and Oechel 2012), increasing area burned in recent years (Pausas and Vallejo 1999). In many USA forests, exclusion of fires by active fire suppression and various land uses has led to an increase in understory trees and an accumulation of surface fuels, which can facilitate a transition from low-severity surface fires to mixed- or high-severity crown fires (Keane et al. 2002; Stambaugh et al. 2014).

12.3 Observed and Anticipated Changes in Climate

Most areas of the USA experienced warmer air temperatures over a recent 20-year period (1986–2015) compared to 1901–1960 (Vose et al. 2017). Changes in annual temperature have been highest (>1.5 °F [0.8 °C]) in the western USA, upper Midwest, northeastern USA, and Alaska, with winter temperatures showing even stronger warming patterns than in other seasons. The primary exception is the mid-South (Alabama, Arkansas, Louisiana, Mississippi), where temperatures have

generally been unchanged or even slightly cooler, especially during summer (Vose et al. 2017), although nighttime temperatures over much of the southern USA have been increasing (Carter et al. 2018). Temperature extremes are also increasing over most of the USA, with a greater frequency of record high temperatures in the last 20 years (Vose et al. 2017).

Observed changes in precipitation indicate drying in many areas of the western USA, especially the Southwest, where recent (1986–2015) annual precipitation has decreased by 15% compared to 1901–1960 (Easterling et al. 2017). Lower annual precipitation has also been observed in the Interior West and parts of the southern USA, whereas the Midwest, Lake States, and Northeast have generally been wetter in recent years (Easterling et al. 2017). Altered seasonal patterns are also evident, with lower winter precipitation in the western USA and higher winter precipitation in the southeastern USA. Many regions are experiencing greater precipitation extremes (Easterling et al. 2017). For example, in the southern USA, heavy rainfall events (e.g., precipitation events >7.5 cm) have been increasing (Easterling et al. 2017; Carter et al. 2018). Less snow (Wehner et al. 2017) and earlier spring melting of snow is exacerbating summer drought conditions and drying fuels in some Western landscapes (Mote et al. 2018).

Model projections of future climate rely on assumptions about future greenhouse gas emissions and atmospheric concentrations. Consistent with the Fourth National Climate Assessment (USGCRP 2017), we focus on describing projections from a lower (RCP4.5 = peak in CO_2 equivalent concentration around 2040, with a value of ~550 ppm by 2100; where RCP = Representative Concentration Pathway) and higher (RCP8.5 = CO_2 concentration continues to increase, with a value of ~1250 ppm CO_2 by 2100) greenhouse gas concentration scenario (van Vuuren et al. 2011). Projections indicate continued warming across the USA, with increased average annual temperature of 8–10 °F (4.4–5.6 °C) for much of the USA by the end of the century under RCP8.5 (Fig. 12.2). Increases of 2–6 °F (1.1–3.3 °C) are projected by the mid-twenty-first century under RCP4.5. A 2–6 °F temperature increase would have significant effects on wildfire occurrence and fire regimes, in many cases accelerating changes already observed (Freeborn et al. 2016). Furthermore, both emission scenarios project an increase in extreme temperatures, as high as 14 °F (7.8 °C) on the warmest days.

Although the capacity of global climate models to project temperature is more certain than for precipitation, projections suggest generally higher precipitation (by as much as a 30% increase) in the upper Midwest, Northeast, Southeast and Pacific Northwest (especially in the winter months) under RCP8.5 by the end of the twenty-first century. Much drier conditions (30% decrease) are projected for the Interior West and Southwest and moderately drier conditions for much of the USA in summer (Fig. 12.3).

Few studies have examined the periodicity and magnitude of future drought events, due primarily to uncertainties in future climatic and environmental conditions that cause drought (Cook et al. 2015; Ryu and Hayhoe 2017; Wehner et al. 2017). Peters and Iverson (2019) used the Cumulative Drought Severity Index (CDSI) derived from a self-calibrated Palmer Drought Severity Index and future



Fig. 12.2 Projected changes in average annual temperatures (°F). Changes are the difference between the average for mid-century (2036–2065; top) or late-century (2070–2099, bottom) and the average for near-present (1976–2005). Each map depicts the weighted multimodel mean. Increases are statistically significant in all areas (>50% of the models show a statistically significant change, and >67% agree on the sign of the change). (Figure source: CICS-NC and NOAA NCEI; USGCRP 2017, Fig. 6.7 in Vose et al. (2017))

temperature to project drought conditions in the USA. The CDSI suggested that more frequent and/or intense droughts are likely in the middle to latter parts of the twenty-first century across all regions of the USA. Similarly, Wehner et al. (2017) projected soil moisture conditions, finding decreased soil moisture over most of the



Fig. 12.3 Projected change (%) in total seasonal precipitation from CMIP5 simulations for 2070–2099. The values are weighted multimodel means (RCP8.5) and expressed as the percent change relative to the 1976–2005 average. Stippling indicates that changes are assessed to be large compared to natural variations. Hatching indicates that changes are assessed to be small compared to natural variations. Blank regions are where projections are assessed to be inconclusive. Data source: World Climate Research Program's (WCRP's) Coupled Model Intercomparison Project. (Fig. source: NOAA NCEI; USGCRP 2017; Fig. 7.5 in Easterling et al. (2017))

conterminous USA in all seasons, with the largest decreases in the Southwest and Interior West.

12.4 Effects of Future Fire Regimes on Ecosystems

The combination of higher temperature, reduced precipitation in some areas, and increased drought implies that wildfire risk will increase substantially in the future (Liu et al. 2010), resulting in more and larger fires. Associated changes in fire regimes are likely to affect the composition and structure of vegetation and forest function, a process that will probably require many decades for substantial changes to occur across large landscapes (Halofsky et al. 2020). Repeated fires (lightning- and human-caused) that occur on fire-prone landscapes over time create a shifting mosaic of plant and animal communities and structures that often reflect species ability to survive fire or to colonize burned areas shortly after fire (Agee 1998; McKenzie et al. 2011). In general, plant species with fire-adapted survival traits (e.g., thick bark, high crowns, serotinous cones, buried seeds, epicormic sprouting, and deep roots) tend to dominate landscapes with frequent fires, in the absence of co-occurring stressors such as extreme droughts, high-severity fires, and invasive species that can transform the landscape to non-forested conditions (Allen et al. 2002). In contrast, more frequent fire will likely decrease the abundance of shade-tolerant species, species with thin bark, and slow invaders after fire (Nowacki and Abrams 2008; Chmura et al. 2011; Vose and Elliott 2016). Fire-sensitive vegetation may be lost from a site if unable to regenerate in burned areas (Harvey et al. 2016).

In many forests, tree species respond to differences in fire frequency, seasonality, and fire severity (Gill and Williams 1996). Different fire intervals influence population trends and patterns of succession (Agee 1993; Schoennagel et al. 2003). In forests adapted to crown fire regimes, the effects of fire depend on demographic attributes of the species. For example, population size of nonsprouting species may fluctuate more than that of sprouting species, and local extinction of nonsprouting species may be common after a single fire (Keeley et al. 2011; Pausas et al. 2018). In frequent surface-fire regimes, lengthening fire intervals can quickly affect functional group dominance and long-term survival of resprouting species (Knapp et al. 2015). Species that are slow to mature are particularly vulnerable to more frequent fire, because populations killed by fire before they have flowered and set seed may be unable to regenerate at small spatial scales (Schlaepfer et al. 2014).

Tree populations are also negatively affected where intervals between fires promote dominance of more competitive species or exceed the lifespan of a species or its seedbank. Seasonality of fire can have a profound effect on forests, because phenological state at the time of fire influences the ability of a species to regenerate (Flannigan et al. 2000; Arthur et al. 2012). Where intensity of fires exceeds the capacity of dominant prefire species to survive or regenerate, dominance may shift to other species, such as the increase in deciduous species over conifers observed in

some boreal and temperate forests (Barrett et al. 2011; Hollingsworth et al. 2013; Stambaugh et al. 2017).

Variation in fire regime characteristics affects vegetation structure at multiple spatial scales (Reilly et al. 2018). At the stand scale, frequent fire tends to reduce forest height, favor shrublands over woodlands, promote flammable or fire-tolerant species and communities, and reduce tree density and biomass (Whelan 1995; Scott et al. 2014). For example, species composition across eastern USA forests has been shifting from fire-tolerant xerophytic species (oaks, hickories, and pines) to less fire-tolerant mesophytic species (maples and yellow – poplar (*Liriodendron tulipifera*)) (Nowacki and Abrams 2008). Reasons for this shift are complex (McEwan et al. 2011; Nowacki and Abrams 2015), but fewer fires in recent decades may be an important driver (Kreye et al. 2013; Hanberry et al. 2020). Climate change may alter future fire regimes in these systems to create forests with lower overall densities and higher proportions of fire-tolerant species (Jenkins et al. 2011; Vose and Elliott 2016).

Fire can also directly influence the spatial mosaic of forest patches across large landscapes (Box 12.1), and climate-mediated changes in disturbance regimes and management interactions may cause landscape heterogeneity to increase (e.g., Marlon et al. 2009). For example, more frequent low- and mixed-severity fires may reduce fuels in drier forest ecosystems (e.g., dry mixed conifer), leading to lower-intensity fires, and a finer-scale patch mosaic (Loudermilk et al. 2019). However, factors that are both directly and indirectly related to climate change could decrease landscape heterogeneity. Continued fire exclusion, coupled with a warmer, drier climate, may promote landscape conditions in which late-seral communities dominate but are stressed from competition and drought (van Mantgem et al. 2009; Stephens et al. 2018). Simulations from global vegetation models suggest that forests might at least double in extent in the absence of fire, particularly in the flammable savanna biome (Daly et al. 2000). Wildfires that eventually burn these landscapes may become large and burn intensely, potentially creating large patches of homogeneous postburn conditions (Haugo et al. 2019) that may convert to savanna, shrub, or grassland communities, especially in areas that are too dry for tree establishment or where seed sources are eliminated (Kuenzi et al. 2008; Stroh et al. 2018).

The complexity of feedbacks among climate change, fire regime, and forest structure and function makes it challenging to project the effects of climate change on future forests. However, examining how historical and contemporary fire regimes influence forest dynamics provides some insight into how forests may respond to climate-altered fire regimes. In the following sections, we examine some of these feedbacks using examples of climate-fire relationships in boreal forests of Alaska and Canada, mesic pine forests of the Southeast, and dry conifer forests of the southwestern USA.

Box 12.1 Landscape Heterogeneity, Restoration, and Resilience

Settlement, development, and land management practices have altered the temporal and spatial characteristics of most USA forests during the past 200 years. Timber management practices have modified vegetation patch shape and structure, and fire exclusion has modified patch size and diversity. Homogeneous Western landscapes generally have low resilience to disturbance (Keane et al. 2002, 2018) and may have little ability to buffer potential climate impacts because of the high tree densities and an abundance of shade-tolerant trees (Vanderwel and Purves 2014). Wildfires and harvest activities over the last decade have returned some heterogeneity to forest landscapes, especially in wilderness areas (Campbell and Shinneman 2017; Hessburg et al. 2019). However, most are well outside of the historical range of variability in landscape structure.

It is challenging for land managers to identify an appropriate level of heterogeneity for a given landscape. Reference conditions can help prioritize and design actions that facilitate landscape heterogeneity and mitigate climate change effects. One method for estimating optimal heterogeneity uses the *historical range and variation* (HRV) of landscape characteristics as a reference (Morgan et al. 1994; Keane et al. 2009). Although the HRV of landscape metrics may not be representative of future conditions (Millar and Woolfenden 1999), it can approximate landscape conditions under which ecosystems have developed over the past thousand years. It is reasonable to assume that these historical conditions produced resilient systems and landscapes (Landres et al. 1999).

The departure of landscapes from historical compositions and structures must be known in order to design effective restoration actions (Keane et al. 2009, 2019; Dickinson 2014). Simulation modeling can be used to generate both HRV time series and future management effects (Loehman et al. 2020). For example, simulation modeling has been used to determine the efficacy of wildland fire suppression activities in the future by comparing with past HRV time series (Keane et al. 2019).

Increased stress associated with climate change—higher temperatures, more droughts, more disturbances—means that restoration (based on past conditions) may need to transition to *resilience building*. This can be accomplished by considering the *future range and variation* (FRV) of landscape characteristics as altered by climate change (Keane et al. 2018, Peterson and Halofsky 2019). FRV can be informed by climate projections integrated with ecosystem modeling to understand the environment in which future forests will regenerate and grow. Land managers can then consider which species, genotypes, structures, and spatial patterns are likely to be resilient several decades into the future.

12.4.1 Boreal Forests

Fire is the dominant disturbance regime in boreal forests (Chapin et al. 2006; Johnstone et al. 2010; Brown and Johnstone 2012; Hollingsworth et al. 2013). Paleorecords indicate that millennial-scale fire-climate relationships are highly influenced by vegetation dynamics (Hu et al. 2006). For example, during cooler, wetter climatic conditions in the Holocene (ca. 4000–7000 ybp), fires occurred relatively frequently compared with preceding millennia, likely because these conditions favored establishment of black spruce (*Picea mariana*) over forests of *Populus* species or white spruce (*P. glauca*) (Lynch et al. 2004; Hoecker et al. 2020). Black spruce forests are more flammable than white spruce or mixed white spruce-deciduous forests due to lower crown-base heights, persistent fine branches, and association with flammable species of understory vegetation (Hu et al. 2006). Little is known about how fire regimes changed in response to major biome shifts before the establishment of black spruce, but following establishment of the modern boreal forest, high-intensity crown fires occurred at intervals of 80–150 years (de Groot et al. 2013).

Under historical climate and fire regimes, boreal forest regeneration following wildfires occurs as (1) self-replacement of the same dominant prefire tree species in the postfire environment, or (2) multiple successions of tree species that assume postfire canopy dominance (Kurkowski et al. 2008). Boreal tree species adaptations for postfire regeneration include cone serotiny (black spruce), rapid resprouting (birch and poplar species), and wind-dispersed seeds (white spruce) (de Groot et al. 2013; Greene and Johnson 2000). Successional pathways depend on soil conditions, prefire species composition, and fire severity and frequency, all of which are affected by variation in temperature and precipitation (Kasischke et al. 2000; Johnstone et al. 2011).

The boreal region of North America is warming twice as fast as the global average (Scheffer et al. 2012; IPCC 2014). Wildfires and seasons with heightened wildfire activity have increased in boreal Alaska over the past 40 years, in association with above-average summer temperatures and prolonged drought (Xiao and Zhuang 2007). Climate change is expected to increase fire season length, increase area burned, and intensify fire weather (Zhang and Chen 2007). Climate-induced shifts in boreal fire regimes can lead to a third successional pathway: species replacement, in which atypically frequent or severe wildfires extirpate prefire canopy species, allowing for new species to dominate (Kurkowski et al. 2008; Johnstone et al. 2010). Increased fire severity may fundamentally reshape future landscapes of Alaska, because high-severity fires can promote conversion of coniferous spruce ecosystems to deciduous hardwoods, leading to broad-scale transformations of forest community composition and structure that may persist for centuries (Johnstone et al. 2010). A shift from conifer forest to less flammable, deciduous-dominated forest could result in a negative feedback to future fire or a shift from canopy- to surface-dominated fire regimes (Johnstone et al. 2011; Hudspith et al. 2015; Hart et al. 2019), unless a warmer climate reduces fuel moisture sufficiently to carry fire in deciduous fuel types (Flannigan et al. 2005).

12.4.2 Fire-Prone Forests in the Southeast

Fire-prone forests in the southeastern USA (Chap. 3) have among the shortest FRIs in the world (1–4 year) (Christensen 1981), but forest change in the past two centuries has been driven largely by forest conversion, development, landscape fragmentation, and reduced fire frequency through fire exclusion (Landers et al. 1995). Fire regimes in this region are now dominated by human ignitions in the form of prescribed fire (Melvin 2018). Maximum temperatures have not risen significantly regionally (Carter et al. 2018), but models project temperatures will increase by the mid-twenty-first century, potentially causing more wildfires (Prestemon et al. 2016).

In this human-dominated landscape, climate change influences on fire regimes are mediated through the filters of human intervention. Managed fire regimes target FRIs of 1–5 years; some studies suggest that managed fire regimes should include months during which lightning ignitions were historically dominant (Platt et al. 1988; Slocom et al. 2007). Climate change is projected to reduce prescribed burning opportunities in the spring and summer due to elevated risks of escape (Kupfer et al. 2020), so the trajectory of ecosystem change in frequently burned communities may increasingly be driven by wildfire. For example, drier and hotter weather conditions will alter fire potential across most ecoregions in the southern USA, which may increase fire season length (Liu et al. 2014) and reduce the number of days when weather conditions are within prescribed fire prescriptions (Kupfer et al. 2020).

Altered prescribed fire frequency has feedbacks on fuel loading, which will increase as the interval between burning widens. Compared to current prescribed fire behavior, prescribed fires in the future may be more severe, kill more and larger trees, and produce more smoke (Mitchell et al. 2014). Although fuel accumulations and consecutive drought years have increased wildfire in some areas (e.g., Georgia) (Terando et al. 2016), other areas in the Southeast may become less flammable as fire is removed from the landscape (Glitzenstein et al. 2012). This alternate stable state in the absence of prescribed fire produces a more mesic forest with longer fire-return intervals and altered ecosystem properties that may perpetuate reduced flammability (McKay and Parker 2001; Kreye et al. 2013; Carpenter et al. 2020) under future climates, although this transition is uncertain (Vose and Elliott 2016).

12.4.3 Fire-Prone Forests in the Southwest

Fire regimes in fire-prone forests of the southwestern USA (Chap. 11) have been altered by a combination of land management and drought. Historically, low-severity fires in dry forests occurred frequently, spreading in grassy surface fuels coincident with an alternating pattern of increased cool-season precipitation (resulting in fuel growth) followed by cool-season drought (resulting in fuel flammability) (Swetnam and Betancourt 1998; Margolis et al. 2017). A hundred-plus years of livestock grazing, logging, and fire exclusion have led to less frequent fire in dry

conifer forests compared with the presettlement period, creating increased surface fuel loads, dense forests, and reduced structural and spatial heterogeneity of vegetation (Allen et al. 2002; Reynolds et al. 2013). Fires in these forests can be more intense with larger patches of high-severity fire than occurred historically (Allen 2016; Singleton et al. 2019), reducing biodiversity, eliminating tree seed sources, and altering ecological function and resilience (Guiterman et al. 2018; Haffey et al. 2018).

Climate change has increased the length of fire seasons and lowered fuel moistures, making large portions of the landscape flammable for longer periods of time (Riley and Loehman 2016), and widespread, regional fire years have been associated with prolonged droughts (Heyerdahl et al. 2008). However, at some point in the future, increased fire activity in these low-productivity ecosystems may be self-limiting, such that fuel consumption limits frequency, extent, and effects of subsequent fires (Collins et al. 2007; Parks et al. 2015).

12.5 Interactions of Future Fire Regimes and Other Stressors and Disturbances

Direct and indirect interactions among fire and other stressors and disturbances will result in persistent changes in forest composition and structure under climate change (Fig. 12.1). In the context of future fire regimes, these changes are especially important for how they influence fuel loads, fire severity, and flammability.

12.5.1 Native Insects and Diseases

Insects and diseases influence forest ecosystem structure and function in complex ways, regulating primary production, nutrient cycling, ecological succession, and the distribution and abundance of plants and animals (Mattson and Addy 1975; Castello et al. 1995). The effects of insects and diseases are most influenced by abiotic factors (e.g., weather, wildfires, avalanches, windstorms) and management activities.

Climate change is expected to increase the effects of some insects and diseases in USA forests and reduce the effects of others (Vose et al. 2018). The effects of insects and diseases can be low level/chronic or acute/episodic. Many USA states have at least 10% of their forested landscapes at risk to forest insects and diseases, in which at least 25% of standing live basal area >2.5 cm in diameter will be killed in the next 15 years in the absence of remediation (management actions applied to increase resistance and resilience). Based on the latest National Insect and Disease Forest Risk Assessment, four tree species are expected to suffer substantial mortality across their extent, including redbay (*Persea borbonia*) (90% of total basal area),



Fig. 12.4 Bark beetles and wildfire are principal drivers of change in conifer forests. This figure shows the area affected by native bark beetles in the western USA, where impacts are most severe, 2000–2016. About 10.3 million ha were affected by mountain pine beetle (*Dendroctonus ponderosae*) alone, which represents almost half of the total area for all bark beetles combined in the western USA (Fettig et al. 2021). Values represent the effect observed each individual year and should not be summed across years (i.e., there may be overlap in areas affected from year to year). Note that the line for *D. jeffreyi* is overlapped by the line for *D. pseudotsugae*. Fig. developed using data from the National Insect and Disease Survey database, USDA Forest Service (Fettig *unpubl. data*)

whitebark pine (*Pinus albicaulis*) (58%), limber pine (*P. flexilis*) (44%), and lodgepole pine (*P. contorta* var. *latifolia*) (39%) (Krist et al. 2014).

Bark beetles cause most of this tree mortality, and several species have undergone notable outbreaks in recent years (Fig. 12.4; Chaps. 8, 9, 10). This is primarily due to climate and human-caused changes in forest structure and composition (Fettig et al. 2007; Bentz et al. 2010; Kolb et al. 2016), which influence host susceptibility, and beetle survival and population growth. For example, increased densities of grand fir (*Abies grandis*) and white fir (*A. concolor*) in the Cascade Range and Sierra Nevada, which are largely due to fire exclusion and selective harvest of pines, have led to increased impacts by fir engraver (*Scolytus ventralis*), especially during periods of elevated temperatures and drought (Fettig et al. 2021).

Increased wildfire and bark beetle outbreaks raise concerns about the effects of wildfires on bark beetles, and the effects of bark beetle outbreaks on fuels and fire regimes (Hicke et al. 2012; Fettig et al. 2020). Following fire, tree mortality can be immediate, or it can occur over multiple years as a result of fire injuries to the crown, bole, or roots (Hood et al. 2018). Levels of delayed tree mortality caused by bark beetles depend on tree species, tree size, tree phenology, degree of fire-caused injuries, initial and postfire levels of tree vigor, the postfire environment, and the scale, severity and composition of bark beetle populations and other tree mortality agents. Trees moderately injured by fire are generally most susceptible to bark

Fig. 12.5 Much of California experienced a severe drought in 2012–2015, inciting a large tree mortality event in the central and southern Sierra Nevada. Much of the tree mortality was attributed to western pine beetle (*Dendroctonus brevicomis*) (Fettig et al. 2019). Bark beetle outbreaks can alter fuels sufficiently to affect fire regimes and firefighter safety (photo L.M. Mortenson, USDA Forest Service, Pacific Southwest Research Station)

beetles. High-severity fires kill large numbers of potential host trees, often reducing bark beetle populations (Fettig et al. 2020). Bark beetles routinely cause additional tree mortality after low- to moderate-severity fires.

Altered fire behavior in forests that have experienced beetle outbreaks are manifest through changes in fuel loadings (Fig. 12.5), depending on the severity of the outbreak (the proportion of trees colonized and killed by bark beetles) and the amount of time since the outbreak occurred. Altered foliar moisture following beetle colonization affects flammability, but this change is short-lived. For example, the needles and twigs of lodgepole pine killed by mountain pine beetle lose 80–90% of their water content within one year. Loss of moisture increases flammability by shortening time to ignition, lowering temperature at ignition, and raising heat yields when burned (Page et al. 2012). Over time, canopy bulk density and canopy cover decline as beetle-killed trees shed needles, then small twigs and larger branches (Fig. 12.5). Bark beetle outbreaks can increase wildfire rates of spread, with crown fire potential peaking 1–3 years after outbreaks (Fig. 12.6), although outbreaks have little effect on the extent of area burned (Hart et al. 2015) or likelihood of wildfire occurrence (Bebi et al. 2003; Meigs et al. 2015).

Defoliating insects can influence fire behavior and severity (Hummel and Agee 2003), but these relationships are still being explored. Fire potential in balsam fir (*A. balsamea*) forests affected by eastern spruce budworm (*Choristoneura*



Fig. 12.6 Conceptual diagram showing changes in fuel profiles and wildfire behavior following severe drought and a bark beetle outbreak in a Sierra Nevada mixed-conifer forest without (**a) and with (b**) periodic fuel reduction treatments. Similar trends may be observed for other disturbances that cause similar patterns and levels of tree mortality. A greater potential for "mass fires" now exists in the central and southern Sierra Nevada, driven by the amount, size, and continuity of dry combustible woody fuels. Adapted from Stephens et al. (2018)

fumiferana) was shown to be significantly higher for several years following tree mortality (Stocks 1987). Fire potential peaked 5–8 years after tree mortality and decreased gradually as surface fuels began to decompose. Flower et al. (2014) proposed that if there is a relationship between defoliators and fire, it is a subtle synergistic relationship in which climate determines the probability of occurrence of each disturbance, and each disturbance dampens the severity but does not alter the probability of occurrence of other disturbances. Forest diseases can also affect fire regimes, although this relationship is not well studied. In Oregon, fire reduced the area infested by laminated root rot (*Phellinus weirii*) by favoring less susceptible host species and reducing the modal size of dead roots and logs, but infestations enhanced the probability of stand-replacing fires (Dickman and Cook 1989).

12.5.2 Invasive (Nonnative) Plants

Although most studies indicate that climate change will accelerate establishment of non-native plants in the USA (Finch et al. 2021), projecting how individual taxa will respond to different climate change scenarios and interact with fire regimes is difficult. Invasive plants, which may exert important effects on fire regimes, generally respond more favorably to elevated CO_2 than native plants (Bradley et al. 2010a), whereas responses to altered temperature and precipitation are variable (Dukes et al. 2009). Many invasive species arrive as stowaways in cargo ships. In the USA, current inspection of cargo ships for invasive species involves examining a low percentage of imports (<2%) for a small subset of federally-listed species (Finch et al. 2021). Hence, it is likely that the influx of invasive species will continue, although their effects on future fire regimes is uncertain.

Cheatgrass (*Bromus tectorum*) is an annual species that dies and dries out in spring, increasing the flammability and continuity of surface fuels (Brooks et al. 2004). Its rapid growth rate, ability to photosynthesize at low temperatures (Rice et al. 1992; Chatterton et al. 1993), and competitiveness under drought stress (Melgoza and Nowak 1991) contribute to its dominance. Cover by cheatgrass is now >15% on over 210,000 km^2 in the Intermountain West (Bradley et al. 2018). These lands are twice as likely to burn as those with a low abundance of cheatgrass, and four times more likely to burn multiple times within a 15-year period. Even small amounts of cheatgrass (1–5% cover) increase fire risk (Balch et al. 2013; Bradley et al. 2018), which is concerning given that the density and distribution of cheatgrass are projected to increase with climate change (Zelikova et al. 2013; Finch et al. 2021).

In the southeastern USA, cogongrass (*Imperata cylindrica*) is an invasive grass that can alter fire regimes by increasing fine-fuel loads, expanding horizontal fuel continuity, and increasing fuel depth, causing higher fire temperature at greater heights (Lippincott 2000). The current distribution of cogongrass is limited (at least in part) by cold; however, higher temperatures may expand its range northward and

westward, increasing its role in altering fire regimes in more areas of the eastern USA (Bradley et al. 2010b).

A related concern is the increased demand for nonnative plants with greater tolerances to warmer temperatures and drought for use in landscaping (Bradley et al. 2012). The risk of these plants escaping and becoming invasive is increasing with urbanization (Marco et al. 2010). These concerns are most relevant to wildfire in the West, especially in areas with a Mediterranean climate where fires can be frequent (Chap. 9). In the eastern USA (especially in Southern forests) where the majority of prescribed burning occurs, nonnative and native invasive trees, shrubs, and grasses can confound restoration activities that use prescribed fire.

12.6 Managing for Resilience

Managing for ecological resilience, defined broadly as the ability of a system to recover after disturbance, is a strategy intended to mitigate the environmental challenges caused by climate change, shifting fire regimes, nonnative flora and fauna, industrialization, urbanization, and their interactions (Seidl et al. 2016; Keane et al. 2018). Some strategies for achieving resilience focus on maintaining or re-establishing historical conditions (Keane et al. 2018). However, rapid climate change and altered disturbance regimes will make it difficult to maintain ecosystem functionality in emerging, no-analog environments (Falk et al. 2019).

Resilience management focuses on improving the long-term response of a system by focusing on specific attributes or drivers, and developing principles for human actions (Benson and Garmestani 2011). Management actions that can increase resilience in forest landscapes typically balance multiple and often competing objectives. Managed fire regimes are a critical element of these activities for conservation outcomes and ecosystem resilience (Stephens et al. 2018; Freeman et al. 2019). Specifically, management options can be crafted to conserve fire-dependent species, reduce wildfire risks, and address climate-related effects and influences (Carter et al. 2018; Vose et al. 2018).

Planning and sequencing of management activities at relevant scales are critical for managed fire regimes, enhancing resilience in fire-prone ecosystems (Krofcheck et al. 2018). For example, forest thinning can increase tree vigor as well as reduce fuels, thus minimizing the intensity and severity of future wildfires (Reinhardt et al. 2008; Hurteau et al. 2019). Thinning guidelines may need to be revised with fewer residual trees for some cover types (e.g., ponderosa pine [*Pinus ponderosa*]) to maintain adequate levels of resilience to drought and disturbances (e.g., bark beetle outbreaks) that influence fire regimes (Peterson et al. 2011a, b). Fuel reduction treatments may need to be more aggressively applied in the future at large spatial scales to yield effective outcomes in treated stands. In addition, more extreme fire weather conditions associated with climate change (Collins 2014) will likely require higher biomass removals to achieve resilience and fuel management objectives.

The near-term challenge in using restoration or fuel treatments is to apply thinning and managed fires in landscapes of highest priority. For example, aggregation and prioritization of treatments in the Lake Tahoe Basin (California) have reached a stage that, according to model results, will alter large wildfire potential (Loudermilk et al. 2014). However, documentation of treatment effectiveness for modifying wildfire potential at large spatial scales in Western landscapes is minimal (with the exception of protecting communities with focused treatments), and few landscapes have been treated at sufficient levels to affect long-term resilience.

In the eastern USA, the intersection of prescribed fire, conservation, and restoration is well developed, particularly in the southeastern USA Coastal Plain (Mitchell et al. 2009; Chap. 3). Prescribed fire use in the southern Great Plains and the Southeast is comparable to or greater than the areal extent of wildfire in the West (Melvin 2018; Hiers et al. 2020). However, climate change is expected to challenge current practice and threaten established restoration management practices (Mitchell et al. 2014; Kupfer et al. 2020). Although prescribed fire is an effective mitigation strategy for reducing wildfire risk (Addington et al. 2015), increasing the capacity of resource managers to safely use this treatment is often constrained by degraded air quality and smoke emissions (Kobziar et al. 2015; Chiodi et al. 2018).

Reducing the effects of existing non-climatic stressors on ecosystems, such as landscape fragmentation and invasive plants, will also increase ecosystem resilience to climatic changes (Joyce et al. 2008). For example, tactics to minimize establishment and spread of invasive species include early detection/rapid response for new invasions, implementing weed-free policies, preventing invasive plant introductions during projects, and planting locally-adapted species of native vegetation to compete with invasive plants (Table 12.1).

12.6.1 Managing Wildfire

Wildfire management involves a broad range of ecological considerations and interactions (Table 12.1, Fig. 12.5, Box 12.1). Wildfires can be actively suppressed (uncontrolled wildfires) or they can be allowed to burn under a prescribed set of weather conditions while being observed (wildfires for resource benefits). The latter option is rarely used, so fuels tend to accumulate well beyond what would occur under a "natural" fire cycle, thus perpetuating elevated fuels and fire hazard, especially in dry forests. In many landscapes, the unconsumed fuels remaining after a wildfire may be insufficient to support the spread of future fires, so previously burned areas act as a firebreak and impede fire spread (Ricotta et al. 1999; Agee et al. 2000; Peterson 2002; McKenzie et al. 2011). As a result, the continuity of flammable vegetation at large spatial scales influences the spread of future fires, and the pattern and amount of area burned influence fuel continuity (McKenzie et al. 2011; Keane et al. 2012).

Wildfire suppression tactics, such as retardant and water drops, backfiring, and fireline construction, can also be used to protect infrastructure, property, and

Table 12.1 Key climate change vulnerabilities of forests related to fire and associated adaptation strategies and on-the-ground tactics. These adaptation options were developed by resource managers in adaptation workshops across the western USA, and are summarized in the Climate Change Adaptation Library (http://adaptationpartners.org/library.php)

Climate change vulnerability	Adaptation strategy	Adaptation tactic
Increased temperatures, drier summers, and lower snowpack will result in more fire (larger areal extent and more high-severity patches).	Plan and prepare for more area burned.	Incorporate climate change into fire management plans.
		Anticipate more opportunities to use wildfire for resource benefit.
		Plan postfire response for large fires.
		Consider using prescribed fire to facilitate transition to a new fire regime in drier forests.
		Consider planting fire-tolerant tree species after fire in areas with increasing fire frequency.
		Manage forest restoration for future range of variability.
	Increase resilience of existing vegetation by reducing hazardous fuels and forest density and maintaining low densities.	Thin and burn to reduce hazardous fuels in the wildland-urban interface.
		Increase intentional use of lightning-ignited fires and management of re-ignition of lightning-ignited fires.
		Consider using more prescribed fire where scientific evidence supports change to more frequent fire regime.
		Use prescribed fire to maintain structure and promote fire-tolerant conifer species.
		Increase interagency coordination and shared risk.
		Conduct thinning treatments (precommercial and commercial).
		Use regeneration and planting to influence forest structure.
	Increase resilience through postfire management.	Consider climate change in postfire rehabilitation.
		Determine where native seed may be needed for postfire planting.
		Anticipate greater need for seed sources and propagated plants.
		Experiment with planting native grass species to compete with invasive species after fire.
		Increase postfire monitoring in areas not currently monitored.

(continued)

Tabel 12. 1 (continued)

Climate change vulnerability	Adaptation strategy	Adaptation tactic
		Identify, prioritize, and protect values at risk; initiate programs to assess values and determine best protective actions.
		Conduct prefire planning to improve response time and efficiency, prioritizing key areas at risk to geologic hazard.
		Conduct postfire vegetation management and prevent invasives.
	Manage forest vegetation to reduce severity and patch size; protect refugia (e.g., old trees).	Map fire refugia.
		Include gaps in silvicultural prescriptions.
		Identify processes and conditions that create fire refugia.
	Use high-severity wildfires as opportunities to modify ecosystem structure.	Use postfire timber harvest to prevent uncharacteristic reburns.
		Allow some burned areas to regenerate naturally.
	Manage forest landscapes to encourage fire to play a natural role.	Implement fuel breaks at strategic locations.
		Create incentives to encourage managed wildland fire.
		Implement strategic density management through forest thinning.
		Push boundaries of prescribed burning (e.g., burn earlier in spring, later in summer).
Disturbances will alter ecosystem structure, species distribution, and species abundance across large landscapes.	Increase knowledge of patterns, characteristics, and rates of change in species distributions.	Expand long-term monitoring programs.
	Create landscape patterns that are resilient to expected disturb-ance regimes.	Continue research on expected future disturbance regimes; evaluate potential transitions and thresholds.
		Improve communication across boundaries.
		Manage for diversity of structure and patch size with fire and mechanical treatments.
Lack of disturbance has caused shifts in species composition and structure in dry mixed-conifer forest, creating a risk of high-severity fire with climate change.	Maintain and restore species and age class diversity.	Identify and map highest risk areas across large landscapes to provide context for prioritization.
		Reduce stand density and shift composition toward species that are more fire adaptive and drought-tolerant.
		Restore age class diversity while protecting legacy trees.

(continued)

Tabel 12. 1 (continued)

Climate change vulnerability	Adaptation strategy	Adaptation tactic
The frequency and scale of disturbance will likely increase with climate change.	Promote disturbance-resilient species.	Thin to favor disturbance-resilient species.
		Plant disturbance-resilient species.
		Promote disturbance-resilient species with prescribed fire and/or managed wildland fire.
Areas with limited species and genetic diversity will likely be more susceptible to climate change stressors.	Promote species and genetic diversity.	Plant potential microsites with a mix of species.
		Maintain species diversity during thinning.
		Interplant to supplement natural regeneration and genetic diversity.
Current dry forest conditions (overstocked stands with more shade-tolerant species as a result of fire suppression) increase vulnerability to drought and wildfire.	Actively manage dry forest areas that are susceptible to fire and drought.	Conduct more intensive thinning.
		Introduce frequent fire.
		Promote ponderosa pine by favoring frequent fires.
Higher-elevation forests may burn more frequently with climate change.	Increase resilience of vegetation types at high elevations.	Increase landscape heterogeneity with prescribed fire.
		Use fire behavior and spatial modeling to identify high-priority areas to reduce or maintain fuels.
		Use silvicultural practices (e.g., prescribed fire, thinning) to reduce fire hazard.
Climate change may increase disturbance interactions, compounding effects.	Increase post-disturbance planning, management, and treatment implementation.	Create a strategy and develop criteria to prioritize areas that are more likely to recover after disturbance (e.g., critical habitats, population served by disturbed habitat).
		Promote climate-adapted species (species resistant and resilient to disturbance) and genotypes.
		Identify sites more susceptible to compounding disturbances (e.g., with high fuel loads, beetle-caused tree mortality, invasives); monitor disturbance occurrence; prioritize seed sources to preserve some sites; map sites across landscapes; conduct proactive treatments in areas more resistant to disturbance.

wildlife habitat (Reinhardt et al. 2008). Postfire activities that mitigate adverse wildfire impacts include erosion control, site stabilization, and planting trees when fire has killed the majority of seed sources. These actions can be implemented to protect ecosystem components that were burned by fire and to restore or maintain resilience. The application of new coupled fire-atmospheric modeling tools to managed wildfires may offer firefighters and communities greater opportunities to use wildfires to increase resilience (Hiers et al. 2020; Linn et al. 2020).

Managing the resilience of fire-prone ecosystems in the USA will likely require more prescribed fire (Hiers et al. 2020). Unlike in the Southeast where prescribed burning is widespread, managed wildfire provides a flexible option for managing fire regimes and increasing resilience in the West. Expanding the use of managed wildfire in the West will be challenging due to risk aversion in decision making (relative to protecting human communities and values) and due to sociopolitical pressure to suppress fires (Hiers et al. 2020).

12.6.2 Wildland Treatments and Effects

In the USA, prescribed fires dominate fire management east of the Rocky Mountains, with nearly five million ha burned annually for a variety of objectives (Melvin 2018; Chap. 8). In the western USA, fuel treatments designed to reduce wildfire intensity and the severity of effects are rarely implemented at scales that are large enough to mitigate undesirable wildfire impacts (Hiers 2017; Kolden 2019). Efforts to reduce fuel loads at meaningful scales are critical but may promote invasive plants in some cases (Schwilk et al. 2009).

Despite the challenges of large-scale implementation of fuel treatments, proactive treatments will undoubtedly play an increasingly important role in reducing wildfire impacts on living trees and in restoring the role of fire in forest ecosystems. However, treating fuels is only one of many objectives for planning and implementing these treatments. Care should be given to maintain stand structures and compositions that facilitate the survival of fire-adapted plant species after wildfires and promote other desired outcomes, such as habitat quality for specific animals.

Eliminating vegetation that competes with fire-adapted trees can improve tree vigor. This will be increasingly important as the climate warms, because low soil moisture, insect outbreaks, and other stressors (including wildfire) will be more prevalent. Increased vigor may allow trees to remain on the landscape for a longer period of time, as well as increase the quality and possibly the quantity (e.g., frequency of cone crops) of reproductive propagules.

Removal of fuels with mechanical treatments and prescribed fire can target specific portions of large landscapes, optimizing the social and ecological value of treatments (Table 12.1). Wildfires are often considered undesirable from a social perspective, but they can in some cases quickly create large-scale heterogeneity that would be difficult to achieve with prescribed fire and mechanical fuel treatments.

Fuel treatments involve reducing canopy fuels by cutting, masticating, or burning living trees, and reducing surface fuels by burning or mechanical removal (Stephens et al. 2012; McIver et al. 2013). Reducing fuels in or near stands that contain ecologically valuable trees can be an important hedge against losing them to future wildfires. However, some fuel treatments, such as mastication, are not designed with ecological relationships in mind. Conversely, restoration treatments that do not remove a large quantity of fuels (including some live trees) may result in unnecessary overstory tree mortality and loss of seed sources when wildfires occur.

Tree harvests are used to remove competing species and to reduce stand density (Table 12.1). It is important that a large number of shade-tolerant trees be removed to optimize the growing environment of fire-tolerant, overstory species. Some cutting treatments are followed with prescribed burning to kill residual (especially small in size and resprouting species) shade-tolerant trees and leave the more fire-tolerant individuals. However, prescribed fires are often highly variable across a given site, missing parts of the stand and severely burning other parts (including desirable trees). Although prescribed fires may be a more ecologically beneficial and less expensive treatment, mechanical removal of trees and fuels is more precise. Combining prescribed burning and mechanical removal is an effective method of achieving multiple desired outcomes (Stephens et al. 2012), and appropriate scheduling can enhance the effectiveness of treatments.

12.6.3 Planting Trees

To mitigate loss of fire-adapted tree species due to climate-mediated changes in disturbance regimes, planting key species might be appropriate when seed sources have been lost because of large, severe fires (Stevens-Rumann et al. 2018) (Table 12.1). Hotter, drier conditions will likely result in postfire regeneration failures in some locations in the future (Littlefield 2019). Increasing postfire seed sources by reducing fire severity (through fuel treatments) and increasing the number of live residual trees can increase natural postfire regeneration in some dry forests (Dodson and Root 2013). Regeneration in the driest topographic locations will generally be slower in a warmer climate than in the past (Boag et al. 2020). Some areas are likely to convert from forest to non-forest vegetation, particularly in ecotones and drier ecoregions.

Resource managers may want to supplement natural regeneration after fire in some locations, for example, in locations farther than 200 m from living trees, and where costs are not prohibitive because of remoteness or topography (North et al. 2019). Potential adaptation strategies include varying planting densities by site microclimate, and creating spatial discontinuity in fuels with variable tree densities.

Fire and large-scale tree mortality events provide opportunities to plant diverse species and genotypes (including genotypes adapted to drought) and to modify forest structure. Through postfire management, managers may be able to help transition ecosystems to warmer conditions by promoting species that will be adapted to the likely future conditions in a particular forest type and setting (Halofsky et al. 2020)

or alter fuel characteristics that affect fire regimes (Stephens et al. 2012; McIver et al. 2013). Managers may want to consider moving seed zones to elevation- and temperature-appropriate locations (within a current species distribution), and modifying genetic movement guidelines to allow more flexibility (Kilkenny et al. 2013). Care should be taken to identify species hybridization and its potential influence on the assumed fire adaptations and tolerance of planted trees (Tauer et al. 2012). Tools such as the Seedlot Selection Tool (https://seedlotselectiontool.org/sst) can help identify seedling stock that will be adapted to a given site in the present and future.

12.6.4 Working Across Boundaries

Fire and other stressors associated with climate change cross boundaries, so implementing climate-smart practices across different ownerships is needed for effective adaptation (Peterson et al. 2011a,b) (Table 12.1). Agencies can coordinate by aligning budgets, priorities, and work, and by better communicating about projects on adjacent lands (Halofsky and Peterson 2016). Agencies and landowners can also coordinate and share monitoring data, which can detect changes in plant species regeneration, growth, and mortality, and help determine adaptation treatment effectiveness (Joyce et al. 2008). With the uncertainty of climate change and its influence on managed fire regimes, collaborative learning will be critical for adapting management strategies to increase resilience.

12.7 Conclusions

Climate change is expected to increase the frequency of large fires, cumulative area burned, and fire suppression costs and risks in many areas of the USA (McKenzie et al. 2004; Vose et al. 2018). Hence, the remainder of the twenty-first century is likely to present substantial challenges, as natural resource managers are challenged by higher fire risk (Fig. 12.7) and the difficult task of maintaining ecological function and increasing resilience in a rapidly changing biophysical and social landscape. Furthermore, no-analog climatic conditions are likely to strain the applicability of current climate-fire regime concepts (e.g., Flatley and Fulé 2016), creating novel ecosystems and forest conditions (Vose et al. 2018; Falk et al. 2019). Uncertain feedbacks may result from the historical legacy of altered fire regimes (e.g., McKay and Parker 2001; Kreye et al. 2013; Carpenter et al. 2020).

The notion of “managing wildfire” to achieve desired outcomes is an emerging concept (Hiers et al. 2020); however, social acceptance of allowing wildfire and associated risks may limit widespread application, especially in the western USA. As a result, fuel treatments will likely play an increasingly important role for minimizing the undesirable ecological effects of fire, and for enhancing firefighter safety (Reinhardt et al. 2008). However, except in the Southeast where prescribed

Fig. 12.7 Extreme droughts and extended dry periods are likely to increase fire risk in many ecosystems in the USA. The photo is from a 2011 wildfire in the Okefenokee Swamp in southeast Georgia/northern Florida. The fire was ignited by lightning during a period of extreme drought (photo credit: US Fish and Wildlife Service)

burning is institutionalized and relatively inexpensive, financial resources for fuel treatments will likely never be sufficient to reduce fuels across large landscapes with elevated fuel loadings. Therefore, fuel treatments must be implemented strategically across large spatial and temporal scales to optimize spatial patterns that protect valued resources and confer resilience to climate change. Collaboration among agencies, private landowners, and other organizations will be increasingly important to ensure large-scale resilience and long-term sustainable management of forest resources in the USA.

References

Abatzoglou JT, Williams AP (2016) Impact of anthropogenic climate change on wildfire across western US forests. Proc Natl Acad Sci U S A 113:11770–11775

Abatzoglou JT, Kolden CA, Williams AP, Lutz JA, Smith AMS (2017) Climatic influences on interannual variability in regional burn severity across western US forests. Int J Wildland Fire 26:269–275

Addington RN, Hudson SJ, Hiers JK, Hurteau MD, Hutcherson TF, Matusick G, Parker JM (2015) Relationships among wildfire, prescribed fire, and drought in a fire-prone landscape in the South-Eastern United States. Int J Wildland Fire 24:778–783

Agee JK (1993) Fire ecology of Pacific Northwest forests. Island Press, Washington, DC

Agee JK (1997) Fire management for the 21st century. In: Kohm KA, Franklin JF (eds) Creating a forestry for the 21st century: the science of ecosystem management. Island Press, Washington, DC, pp 191–201

Agee JK (1998) The landscape ecology of western forest fire regimes. Northwest Sci 72:24–34

Agee JK, Bahro B, Finney MA, Omi PN, Sapsis DB, Skinner CN, van Wagtendonk JW, Phillip Weatherspoon C (2000) The use of shaded fuelbreaks in landscape fire management. For Ecol Manag 127:55–66

Allen CD (2016) Forest ecosystem reorganization underway in the southwestern US: a preview of widespread forest changes in the Anthropocene? In: Sample VA, Bixler RP, Miller C (eds) Forest conservation and management in the Anthropocene: adaptation of science, policy and practices. University Press of Colorado, Boulder, pp 103–122

Allen CD, Savage M, Falk DA, Suckling KF, Swetnam TW, Schulke T, Stacey PB, Morgan P, Hoffman M, Klingel JT (2002) Ecological restoration of southwestern ponderosa pine ecosystems: a broad perspective. Ecol Appl 12:1418–1433

Arno SF, Parsons DJ, Keane RE (2000) Mixed-severity fire regimes in the northern Rocky Mountains: consequences of fire exclusion and options for the future. In: Cole DN, McCool SF, Borrie WT, O'Loughlin J (comps) Wilderness Science in a Time of Change Conference, vol 5. USDA Forest Service Proceed RMRS-P-15-Vol-5, Ogden, Utah, pp 225–232

Arthur MA, Alexander HD, Dey DC, Schweitzer CJ, Loftis DL (2012) Refining the oak-fire hypothesis for management of oak-dominated forests of the eastern United States. J For 110:257–266

Bachelet D, Lenihan JM, Daly C et al (2000) Interactions between fire, grazing and climate change at Wind Cave National Park, SD. Ecol Model 134:229–244

Balch JK, Bradley BA, Antonio CM et al (2013) Introduced annual grass increases regional fire activity across the arid western USA (1980–2009). Glob Chang Biol 19:173–183

Balch JK, Bradley BA, Abatzoglu JT et al (2017) Human-started wildfires expand the fire niche across the United States. Proc Natl Acad Sci U S A 114(11):2946–2951

Barrett K, McGuire AD, Hoy EE et al (2011) Potential shifts in dominant forest cover in interior Alaska driven by variations in fire severity. Ecol Appl 21:2380–2396

Barrows JS, Sandberg DV, Hart JD (1977) Lightning fires in Northern Rocky Mountain forests. USDA Forest Service Intermountain Fire Sciences Laboratory, On file, USDA Forest Service, Intermountain Fire Sciences Laboratory, P.O. Box 8089, Missoula, Montana

Bebi P, Kulakowski D, Veblen TT (2003) Interactions between fire and spruce beetles in a subalpine Rocky Mountain forest landscape. Ecology 84:362–371

Benson MH, Garmestani AS (2011) Can we manage for resilience? The integration of resilience thinking into natural resource management in the United States. Environ Manag 48(3):392–399

Bentz BJ, Régnière J, Fettig CJ, Hansen EM, Hayes JL, Hicke JA, Kel-sey RG, Negrón JF, Seybold SJ (2010) Climate change and bark beetles of the western United States and Canada: direct and indirect effects. Bioscience 60:602–613

Boag AE, Ducey MJ, Palace MW, Hartter J (2020) Topography and fire legacies drive variable post-fire juvenile conifer regeneration in eastern Oregon, USA. For Ecol Manag. https://doi.org/10.1016/j.foreco.2020.118312

Bond WJ, Keane RE (2017) Ecological effects of fire, vol 2. Elsevier. https://doi.org/10.1016/B978-0-12-809633-8.02098-7

Bond WJ, Keeley JE (2005) Fire as a global 'herbivore': the ecology and evolution of flammable ecosystems. Trends Ecol Evol 20:387–394

Bowman DMJS, Blach JK, Artaxo P et al (2009) Fire in the earth system. Science 324:481–484

Bradley BA, Blumenthal DM, Wilcove DS, Ziska LH (2010a) Predicting plant invasions in an era of global change. Trends Ecol Evol 25:310–318

Bradley BA, Wilcove DS, Oppenheimer M (2010b) Climate change increases risk of plant invasion in the eastern United States. Biol Invasions 12:1855–1872

Bradley BA, Blumenthal DM, Early R, Grosholz ED, Lawler JJ, Miller LP, Sorte CJ, D'Antonio CM, Diez JM, Dukes JS, Ibanez I (2012) Global change, global trade, and the next wave of plant invasions. Fron Ecol Environ 10:20–28

Bradley BA, Curtis CA, Fusco EJ, Abatzoglou JT, Balch JK, Dadashi S, Tuanmu MN (2018) Cheatgrass (*Bromus tectorum*) distribution in the intermountain Western United States and its relationship to fire frequency, seasonality, and ignitions. Biol Invasions 20:1493–1506

Bradstock R, Malcom Gill A, Williams R (2012) Flammable Australia: fire regimes, biodiversity and ecosystems in a changing world. CSIRO Publishing Collingwood https://doi.org/10.1111/aec.12026

Brooks ML, D'antonio CM, Richardson DM, Grace JB, Keeley JE, DiTo-maso JM, Hobbs RJ, Pellant M, Pyke D (2004) Effects of invasive alien plants on fire regimes. Bioscience 54:677–688

Brown JK (1995) Fire regimes and their relevance to ecosystem management. In: Managing forests to meet peoples' needs: Proceedings of the 1994 Society of American Foresters/Canadian Institute of Forestry Convention, 18–22 September 1994, Anchorage, Alaska. Society of American Foresters, Bethesda, Maryland, pp 171–178

Brown CD, Johnstone JF (2012) Once burned, twice shy: repeat fires reduce seed availability and alter substrate constraints on *Picea mariana* regeneration. For Ecol Manag 266:34–41

Campbell JL, Shinneman DJ (2017) Potential influence of wildfire in modulating climate-induced forest redistribution in a central Rocky Mountain landscape. Ecol Proc 6:7

Carpenter DO, Taylor MK, Callaham MA et al (2020) Benefit or liability? The ectomycorrhizal association may undermine tree adaptations to fire after long-term fire exclusion. Ecosystems https://doi.org/10.1007/s10021-020-00568-7

Carter L, Terando AJ, Dow K, Hiers K, Kunkel KE, Lascurain AR, Marcy D, Osland MJ, Schramm P (2018) Southeast. In: Reidmiller DR, Avery CW, Easterling DR, Kunkel KE, Lewis KL, Maycock TK, Stewart BC (eds) Impacts, risks, and adaptation in the United States: 4th national climate assessment, vol 2. US Global Change Research Program, Washington, DC, pp 743–808

Castello JD, Leopold DL, Smallidge PJ (1995) Pathogens, patterns, and processes in forest ecosystems. Bioscience 45:16–24

Chapin FS, Oswod MW, Van Cleve K, Viereck LA, Verbyla DL (2006) Alaska's changing boreal forest. Oxford University Press, New York

Chatterton NJ, Harrison PA, Thornley WR, Bennett JH (1993) Structures of fructan oligomers in cheatgrass (*Bromus tectorum* L.). New Phytol 124:389–396

Chen Y, Randerson JT, Morton DC, DeFries RS, Collatz GJ, Kasibhatla PS, Giglio L, Jin Y, Marlier ME (2011) Forecasting fire season severity in South America using sea surface temperature anomalies. Science 334:787–791

Chiodi AM, Larkin NS, Varner JM (2018) An analysis of southeastern US prescribed burn weather windows: seasonal variability and El Niño associations. Int J Wildland Fire 27:176–189

Chmura DJ, Anderson PD, Howe GT, Harrington CA, Halofsky JE, Peter-son DL, Shaw DC, Clair JB (2011) Forest responses to climate change in the northwestern United States: Ecophysiological foundations for adaptive management. For Ecol Manag 261:1121–1142

Christensen NL (1981) Fire regimes in southeastern ecosystems. In: Mooney HA, Bonnicksen TM, Christensen NL, Lotan JE, Reiners WA (tech coords) Proceedings of the conference fire regimes and ecosystem properties, USDA Forest Service Gen Tech Rep WO-26, Washington, DC, pp. 112–136

Cochrane M (2010) Tropical fire ecology: climate change, land use and ecosystem dynamics. Springer, Berlin

Collins BM (2014) Fire weather and large fire potential in the northern Sierra Nevada. Agric For Meteorol 189–190:30–35

Collins BM, Kelly M, van Wagtendonk JW, Stephens SL (2007) Spatial patterns of large natural fires in Sierra Nevada wilderness areas. Landsc Ecol 22:545–557

Cook BI, Ault TR, Smerdon JE (2015) Unprecedented 21st century drought risk in the American Southwest and Central Plains. Sci Adv 1(1):e1400082

Daly C, Bachelet D, Lenihan JM, Neilson RP, Parton W, Ojima D (2000) Dynamic simulation of tree-grass interactions for global change studies. Ecol Appl 10:449–469

de Groot WJ, Flannigan MD, Cantin AS (2013) Climate change impacts on future boreal fire regimes. For Ecol Manag 294:35–44

DeBano LF, Neary DG, Ffolliott PF (1998) Fire's effect on ecosystems. Wiley, New York

Dickinson Y (2014) Landscape restoration of a forest with a historically mixed-severity fire regime: what was the historical landscape pattern of forest and openings? For Ecol Manag 331:264–271

Dickman A, Cook S (1989) Fire and fungus in a mountain hemlock forest. Can J Bot 67:2005–2016

Dodson EK, Root HT (2013) Conifer regeneration following stand-replacing wildfire varies along an elevation gradient in a ponderosa pine forest, Oregon. For Ecol Manag 302:163–170

Duffy PA, Walsh JE, Graham JM, Mann DH, Rupp TS (2005) Impacts of large-scale atmospheric-ocean variability on Alaskan fire season severity. Ecol Appl 15:1317–1330

Dukes JS, Pontius J, Orwig D, Garnas JR, Rodgers VL, Brazee N, Cooke B, Theoharides KA, Stange EE, Harrington R, Ehrenfeld J (2009) Responses of insect pests, pathogens, and invasive plant species to climate change in the forests of northeastern North America: what can we predict? Can J For Res 39:231–248

Easterling DR, Arnold J, Knutson T, Kunkel K, LeGrande A, Leung LR, Vose R, Waliser D, Wehner M (2017) Precipitation change in the United States. In: Wuebbles DJ, Fahey DW, Hibbard KA, Dokken DJ, Stewart BC, Maycock TK (eds) Climate science special report: 4th national climate assessment, vol1. US Global Change Research Program, Washington, DC, pp 207–230

Falk DA, Heyerdahl EK, Brown PM, Farris C, Fulé PZ, McKenzie D, Swetnam TW, Taylor AH, Van Horne ML (2011) Multi-scale controls of historical forest-fire regimes: new insights from fire-scar networks. Front Ecol Environ 9:446–454

Falk DA, Watts AC, Thode AE (2019) Scaling ecological resilience. Front Ecol Evol. https://doi.org/10.3389/fevo.2019.00275

Fettig CJ, Klepzig KD, Billings RF, Munson AS, Nebeker TE, Negrón JF, Nowak JT (2007) The effectiveness of vegetation management practices for prevention and control of bark beetle outbreaks in coniferous forests of the western and southern United States. For Ecol Manag 238:24–53

Fettig CJ, Mortenson LA, Bulaon BM, Foulk PB (2019) Tree mortality following drought in the central and southern Sierra Nevada, California, U.S. For Ecol Manag 432:164–178

Fettig CJ, Progar RA, Paschke J, Sapio FJ (2020) Forest insects. In: Robertson G, Barrett T (eds) Disturbance and sustainability in forests of the western United States. USDA Forest Service Gen Tech Rep PNW-992. Portland, Oregon, pp 81-122

Fettig CJ, Hood SM, Runyon JB, Stalling CM (2021) Bark beetle and fire interactions in western coniferous forests: research findings. Fire Manag Today 79(1):14–23

Finch DM, Butler JL, Runyon JB, Fettig CJ, Kilkenny FF, Jose S, Frankel SJ, Cushman SA, Cobb RC, Dukes JS, Hicke JA, Amelon SK (2021) Effects of climate change on invasive species. In: Poland TM, Patel-Weynand T, Finch D, Miniat CF, Lopez V (eds) Invasive species in forests and grasslands of the United States: a comprehensive science synthesis for the United States forest sector. Springer, Berlin, pp 57–83

Flannigan MD, Van Wagner CE (1991) Climate change and wildfire in Canada. Can J For Res 21:66–72

Flannigan MD, Stocks BJ, Wootton BM (2000) Climate change and forest fires. Sci Total Environ 262(3):221–229

Flannigan MD, Amiro BD, Logan KA, Stocks BJ, Wotton BM (2005) Forest fires and climate change in the 21st century. Mitig Adapt Strateg Glob Chang 11:847–859

Flatley WT, Fule PZ (2016) Are historical fire regimes compatible with future climate? Ecosphere 7(10):e01471. https://doi.org/10.1002/ecs2.1471

Flower A, Gavin DG, Heyerdahl EK, Parsons RA, Cohn GM (2014) Western spruce budworm outbreaks did not increase fire risk over the last three centuries: a dendrochronological analysis of inter-disturbance synergism. PLoS One 9(12):e114282

Freeborn PH, Jolly WM, Cochrane MA (2016) Impacts of changing fire weather conditions on reconstructed trends in U.S. wildland fire activity from 1979 to 2014. J Geophys Res Biogeosci 121:2856–2876

Freeman JE, Kobziar LN, Leone EH, Williges K (2019) Drivers of plant functional group richness and beta diversity in fire-dependent savannas. Divers Distrib 25:1024–1044

Frost CC (1998) Presettlement fire frequency regimes of the United States: a first approximation. Tall Timbers Fire Ecol Conf 20:70–81

Fuquay DM (1980) Lightning that ignites forest fires. In: Proceedings, sixth conference on Fire and Meteorology, Seattle, WA., USA, April 22–24, 1980. Society of American Foresters, Washington, DC, pp. 109–112

Gill AM, Williams JE (1996) Fire regimes and biodiversity: the effects of fragmentation of south-east Australian eucalypt forests by urbanization, agriculture and pine plantations. For Ecol Manag 85:261–278

Glitzenstein JS, Streng DR, Masters RE, Robertson KM, Hermann SM (2012) Fire-frequency effects on vegetation in North Florida pinelands: another look at the long-term Stoddard fire research plots at tall timbers Research Station. For Ecol Manag 264:197–209

Greene DF, Johnson EA (2000) Tree recruitment from burn edges. Can J For Res 30(8):1264–1274

Grissino-Mayer HD, Swetnam TW (2000) Century scale climate forcing of fire regimes in the American southwest. The Holocene 10(2):213–220

Guiterman CH, Margolis EQ, Allen CD, Falk DA, Swetnam TW (2018) Long-term persistence and fire resilience of oak shrubfields in dry conifer forests of northern New Mexico. Ecosystems 21(5):943–959

Guyette RP, Stambaugh MC, Dey DC, Muzika R-M (2012) Predicting fire frequency with chemistry and climate. Ecosystems 15:322–335

Haffey C, Sisk TD, Allen CD, Thode AE, Margolis EQ (2018) Limits to ponderosa pine regeneration following large high-severity forest fires in the United States Southwest. Fire Ecol 14(1):143–163

Halofsky JE, Peterson DL (2016) Climate change vulnerabilities and adaptation options for forest vegetation management in the northwestern USA. Atmos 7(3):46

Halofsky JE, Peterson DL, Harvey BJ (2020) Changing wildfire, changing forests: the effects of climate change on fire and vegetation in the Pacific Northwest. Fire Ecol 16(1):4

Hanberry BB, Abram MD, Arthur MA, Varner JM (2020) Reviewing fire, climate, deer, and foundation species as drivers of historically open oak and pine forests and transition to closed forests. Front For Glob Chang 3:56

Hart SJ, Schoennagel T, Veblen TT, Chapman TB (2015) Area burned in the Western United States is unaffected by recent mountain pine beetle outbreaks. Proc Natl Acad Sci U S A 112:4375–4380

Hart SJ, Henkelman J, McLoughlin PD, Nielsen SE, Truchon-Savard A, Johnstone JF (2019) Examining forest resilience to changing fire frequency in a fire-prone region of boreal forest. Glob Chang Biol 25(3):869–884

Harvey BJ, Donato DC, Turner MG (2016) High and dry: post-fire tree seedling establishment in subalpine forests decreases with post-fire drought and large stand-replacing burn patches. Glob Ecol Biogeogr 25:655–669

Haugo RD, Kellogg BS, Cansler CA, Kolden CA, Kemp KB, Robertson JC, Metlen KL, Vaillant NM, Restaino CM (2019) The missing fire: Quantifying human exclusion of wildfire in Pacific Northwest forests, USA. Ecosphere 10:e02702

Hessburg PF, Miller CL, Povak NA, Taylor AH, Higuera PE, Prichard SJ, North MP, Collins BM, Hurteau MD, Larson AJ, Allen CD (2019) Climate, environment, and disturbance history govern resilience of western North American forests. Front Ecol Evol 7:239

Heyerdahl EK, Brubaker LB, Agee JK (2001) Spatial controls of historical fire regimes: a multiscale example from the interior west, USA. Ecology 82:660–678

Heyerdahl EK, Brubaker LB, Agee JK (2002) Annual and decadal climate forcing of historical fire regimes in the interior Pacific Northwest, USA. The Holocene 12:597–604

Heyerdahl EK, Morgan P, Riser JP (2008) Multi-season climate synchronized historical fires in dry forests (1650–1900), northern Rockies, USA. Ecology 89:705–716

Hicke JA, Johnson MC, Hayes JL, Preisler HK (2012) Effects of bark beetle-caused tree mortality on wildfire. For Ecol Manag 271:81–90

Hiers JK (2017) A century of wildland fire research: contributions to long-term approaches for wildland fire management. Proceedings of a workshop, National Academies of sciences, engineering, and medicine, National Academies Press, pp 65–72

Hiers JK, O'Brien JJ, Varner JM, Butler BW, Dickinson M, Furman J, Gallagher M, Godwin D, Goodrick SL, Hood SM, Hudak A (2020) Prescribed fire science: the case for a refined research agenda. Fire Ecol 16(11):1–15

Hoecker TJ, Higuera PE, Kelly R, Hu FS (2020) Arctic and boreal paleofire records reveal drivers of fire activity and departures from Holocene variability. Ecology:e03096

Holden ZA, Swanson A, Luce CH, Jolly WM, Maneta M, Oyler JW, Warren DA, Parsons R, Affleck D (2018) Decreasing fire season precipitation increased recent western US wildfire activity. Proc Natl Acad Sci U S A 36:E8349–E8357

Hollingsworth TN, Johnstone JF, Bernhardt EL, Chapin FS III (2013) Fire severity filters regeneration traits to shape community assembly in Alaska's boreal forest. PLoS One 8(2):e56033

Hood S, Varner M, van Mantgem P, Cansler CA (2018) Fire and tree death: understanding and improving modeling of fire-induced tree mortality. Environ Res Lett 13:113004

Hu FS, Brubaker LB, Gavin DG, Higuera PE, Lynch JA, Rupp TS, Tinner W (2006) How climate and vegetation influence the fire regime of the Alaskan boreal biome: the Holocene perspective. Mitig Adapt Strateg Glob Chang 11(4):829–846

Hudspith VA, Belcher CM, Kelly R, Hu FS (2015) Charcoal reflectance reveals early Holocene boreal deciduous forests burned at high intensities. PLoS One 10(4):e0120835

Hummel S, Agee JK (2003) Western spruce budworm defoliation effects on forest structure and potential fire behavior. Northwest Sci 7:159–169

Hurteau MD, Liang S, Westerling AL, Wiedinmyer C (2019) Vegetation-fire feedback reduces projected area burned under climate change. Sci Rep 9:2838

IPCC (International Panel on Climate Change) (2014) Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Edenhofer O, Pichs-Madruga R, Sokona Y, Farahani E, Kadner S, Seyboth K, Adler A, Baum I, Brunner S, Eickemeier P, Kriemann B, Savolainen J, Schlömer S, von Stechow C, Zwickel T,Minx JC (eds) Cambridge University Press, Cambridge UK/New York

Jenkins MA, Klein RN, McDaniel VL (2011) Yellow pine regeneration as a function of fire severity and post-burn stand structure in the southern Appalachian Mountains. For Ecol Manag 262(4):681–691

Johnstone J, Hollingsworth TN, Chapin FS III, Mack MC (2010) Changes in fire regime break the legacy lock on successional trajectories in Alaskan boreal forest. Glob Chang Biol 16:1281–1295

Johnstone JF, Rupp TS, Olson M, Verblya D (2011) Modeling impacts of fire severity on successional trajectories and future fire behavior in Alaskan boreal forests. Landsc Ecol 26(4):487–500

Jolly WM, Cochrane MA, Freeborn PH, Holden ZA, Brown TJ, Williamson GJ, Bowman DM (2015) Climate-induced variations in global wildfire danger from 1979 to 2013. Nat Commun 6:7537

Joyce LA, Blate GM, Littell JS, McNulty SG (2008) National forests. In: Julius SH, West JM (eds) Preliminary review of adaptation options for climate-sensitive ecosystems and resources. Synthesis and Assessment Product 4.4. US Environmental Protection Agency, Washington, DC, pp 19–84

Kasischke ES, French NH, O'Neill KP, Richter DD, Bourgeau-Chavez LL, Harrell PA (2000) Influence of fire on long-term patterns of forest succession in Alaskan boreal forests. In: Fire, climate change, and carbon cycling in the boreal forest. Springer, New York, pp 214–235

Keane RE (2013) Disturbance regimes and the historical range of variation in terrestrial ecosystems. In: Simon AL (ed) Encyclopedia of biodiversity, 2nd edn. Academic Press, Waltham, pp 568–581

Keane RE (2015) Wildland fuel fundamentals and applications. Springer, New York

Keane RE (2016) Disturbance regimes and the historical range and variation in terrestrial ecosystems. In: Reference module in life sciences, Encyclopedia of biodiversity, 2nd edn. https://doi.org/10.1016/B978-0-12-809633-8.02397-9

Keane RE, Ryan KC, Veblen TT, Allen CD, Logan JA, Hawkes B, Barron J (2002) The cascading effects of fire exclusion in the Rocky Mountains. In: Baron J (ed) Rocky Mountain futures: an ecological perspective. Island Press, Washington, DC, pp 133–153

Keane RE, Parsons R, Rollins MG (2004) Predicting fire regimes across multiple scales. In: Perera A, Buse L (eds) Emulating natural disturbances: concepts and techniques. Cambridge University Press, Cambridge, pp 55–69

Keane RE, Hessburg PF, Landres PB, Swanson FJ (2009) A review of the use of historical range and variation (HRV) in landscape management. For Ecol Manag 258:1025–1037

Keane RE, Gray K, Bacciu V (2012) Spatial variability of wildland fuel characteristics in northern Rocky Mountain ecosystems. USDA Forest Service Res Pap RMRS-RP-98, Fort Collins, Colorado

Keane RE, Loehman RA, Holsinger LM (2018) Use of landscape simulation modeling to quantify resilience for ecological applications. Ecosphere 9:e02414

Keane RE, Gray K, Davis B, Holsinger LM, Loehman R (2019) Evaluating ecological resilience across wildfire suppression levels under climate and fuel treatment scenarios using landscape simulation modelling. Int J Wildland Fire 28:533–549

Keeley JE (2006) Fire management impacts on invasive plants in the western United States. Conserv Biol 20:375–384

Keeley JE, Syphard AD (2018) Historical patterns of wildfire ignition sources in California ecosystems. Int J Wildland Fire 27:781–799

Keeley JE, Pausas JG, Rundel PW, Bond WJ, Bradstock RA (2011) Fire as an evolutionary pressure shaping plant traits. Trends Plant Sci 16:406–411

Kilkenny F, St. Clair B, Horning, M (2013) Climate change and the future of seed zones. In: Haase DL, Pinto JR, Wilkinson KM (tech cords) National Proceedings: Forest and Conservation Nursery Associations - 2012. USDA Forest Service Proc RMRS-P-69, Fort Collins, Colorado, pp 87–89

Knapp BO, Stephan K, Hubbart JA (2015) Structure and composition of an oak-hickory forest after over 60 years of repeated prescribed burning in Missouri, USA. For Ecol Manag 344:95–109

Kobziar LN, Godwin D, Taylor L, Watts AC (2015) Perspectives on trends, effectiveness, and impediments to prescribed burning in the southern US. Forests 6(3):561–580

Kolb TE, Fettig CJ, Ayres MP, Bentz BJ, Hicke JA, Mathiasen R, Stewart JE, Weed AS (2016) Observed and anticipated impacts of drought on forest insects and diseases in the United States. For Ecol Manag 380:321–334

Kolden CA (2019) We're not doing enough prescribed fire in the Western United States to mitigate wildfire risk. Fire 2(2):30

Krawchuk MA, Moritz MA, Parisien MA, Van Dorn J, Hayhoe K (2009) Global pyrogeography: the current and future distribution of wildfire. PLoS One 4:e5102

Kreye JK, Varner JM, Hiers JK, Mola J (2013) Toward a mechanism for eastern North America forest mesophication: differential litter drying across 17 species. Ecol Appl 23:1976–1986

Krist FJ, Ellenwood JR, Woods ME, McMahan JE, Cowardin JP, Ryerson DE, Sapio F, Zweifler MO, Romero SA (2014) 2013–2027 national insect and disease forest risk assessment. USDA Forest Service Forest Health Technology Enterprise Team Report 14-01, Fort Collins, Colorado

Krofcheck DJ, Hurteau MD, Scheller RM, Loudermilk EL (2018) Prioritizing forest fuels treatments based on the probability of high-severity fire restores adaptive capacity in Sierran forests. Glob Chang Biol 24:729–737
Kuenzi AM, Fule PZ, Sieg CH (2008) Effects of fire severity and pre-fire stand treatment on plant community recovery after a large wildfire. For Ecol Manag 255:855–865
Kupfer JA, Terando AJ, Gao P, Teske C, Hiers JK (2020) Climate change projected to reduce prescribed burning opportunities in the southeastern United States. Int J Wildland Fire 29:764–778
Kurkowski TA, Mann DH, Rupp TS, Verblya DL (2008) Relative importance of different secondary successional pathways in an Alaskan boreal forest. Can J For Res 38(7):1911–1923
Landers JL, Van Lear DH, Boyer WD (1995) The longleaf pine forests of the southeast: requiem or renaissance? J For 93(11):39–44
Landres PB, Morgan P, Swanson FJ (1999) Overview of the use of natural variability concepts in managing ecological systems. Ecol Appl 9(4):1179–1188
Linn RR, Goodrick SL, Brambilla S, Brown MJ, Middleton RS, O'Brien JJ, Hiers JK (2020) Quic-fire: a fast simulation tool for prescribed fire planning. Environ Model Softw 125:104616
Lippincott CL (2000) Effects of *Imperata cylindrical* (L.) Beauv. (Cogongrass) invasion on fire regime in Florida Sandhill (USA). Nat Areas J 20(2):140–149
Littlefield CE (2019) Topography and post-fire climatic conditions shape spatio-temporal patterns of conifer establishment and growth. Fire Ecol 15(1):34
Liu Y, Stanturf J, Goodrick S (2010) Wildfire potential evaluation during a drought event with a regional climate model and NDVI. Ecol Infomatics 5:418–428
Liu Y, Goodrick S, Heilman WE (2014) Wildland fire emissions, carbon, and climate: wildfire-climate interactions. For Ecol Manag 317:80–96
Loehman RA, Keane RE, Holsinger LM (2020) Simulation modeling of complex climate, wildfire, and vegetation dynamics to address wicked problems in land management. Front For Glob Chang 3. https://doi.org/10.3389/ffgc.2020.00003
Loudermilk EL, Stanton A, Scheller RM, Dilts TE, Weisberg PJ, Skinner C, Yang J (2014) Effectiveness of fuels treatments for mitigating wildfire risk and mitigating forest carbon: a case study in the Lake Tahoe Basin. For Ecol Manag 323:114–125
Loudermilk EL, Dyer L, Pokswinski S (2019) Simulating groundcover community assembly in a frequently burned ecosystem using a simple neutral model. Front Plant Sci 10:1107
Lynch JA, Hollis JL, Hu FS (2004) Climatic and landscape controls of the boreal forest fire regime: Holocene records from Alaska. J Ecol 92(3):477–489
Marco A, Lavergne S, Dutoit T, Bertaudiere-Montes V (2010) From the backyard to the backcountry: how ecological and biological traits explain the escape of garden plants into Mediterranean old fields. Biol Invasions 12:761–779
Marcoux HM, Daniels LD, Gergel SE, Da Silva E, Gedalof ZE, Hessburg PF (2015) Differentiating mixed-and high-severity fire regimes in mixed-conifer forests of the Canadian Cordillera. For Ecol Manag 341:45–58
Margolis EQ, Woodhouse CA, Swetnam TW (2017) Drought, multi-seasonal climate, and wildfire in northern New Mexico. Clim Chang 142(3–4):433–446
Marlon JR, Bartlein PJ, Walsh MK, Harrison SP, Brown KJ, Edwards ME, Higuera PE, Power MJ, Anderson RS, Briles C, Brunelle A, Carcaillet C, Daniels M, Hu FS, Lavoie M, Long C, Minckley T, Richard PJH, Scott AC, Shafer DS, Tinner W, Umbanhowar CE, Whitlock C (2009) Wildfire responses to abrupt climate change in North America. Proc Natl Acad Sci U S A 106:219–2524
Mattson WJ Jr, Addy ND (1975) Phytophagous insects as regulators of forest primary production. Science 90:515–522
McEwan RW, Dyer JM, Pederson N (2011) Multiple interacting ecosystem drivers: toward an encompassing hypothesis of oak forest dynamics across eastern North America. Ecography 34:244–256
McIver JD, Stephens SL, Agee JK, Barbour J, Boerner RE, Edminster CB, Erickson KL, Farris KL, Fettig CJ, Fiedler CE, Haase S (2013) Ecological effects of alternative fuel reduction

treatments: highlights of the national fire and fire surrogate study (FFS). Int J Wildland Fire 22:63–82
McKay DH, Parker KC (2001) The role of vegetation dynamics in Choctawhatchee sand pine invasion of longleaf pine forests. Phys Geogr 22(6):467–482
McKenzie D, Gedalof Z, Peterson DL, Mote P (2004) Climatic change, wildfire, and conservation. Conserv Biol 18:890–902
McKenzie D, Miller C, Falk DA (2011) The landscape ecology of fire. Springer, Dordrecht
Meigs GW, Campbell JL, Zald HS, Bailey JD, Shaw DC, Kennedy RE (2015) Does wildfire likelihood increase following insect outbreaks in conifer forests? Ecosphere 6:art118
Melgoza G, Nowak RS (1991) Competition between cheatgrass and two native species after fire: implications from observations and measurements of root distribution. J Rangel Manag 44:27–33
Melvin MA (2018) National prescribed fire use survey report. Tech Rep 03-18. Coalition of Prescribed Fire Councils Inc., Newton, Georgia
Millar CI, Woolfenden WB (1999) The role of climate change in interpreting historical variability. Ecol Appl 9:1207–1216
Mitchell RJ, Hiers JK, O'Brien J, Starr G (2009) Ecological forestry in the Southeast: understanding the ecology of fuels. J For 107(8):391–397
Mitchell RJ, Liu Y, O'Brien JJ (2014) Future climate and fire interactions in the southeastern region of the United States. For Ecol Manag 327:316–326
Moreira F, Rego F, Ferreira P (2001) Temporal (1958–1995) pattern of change in a cultural landscape of northwestern Portugal: implications for fire occurrence. Landsc Ecol 16:557–567
Moreno JM, Oechel WC (2012) The role of fire in Mediterranean-type ecosystems, vol 107. Springer, New York
Morgan P, Aplet GH, Haufler JB et al (1994) Historical range of variability: a useful tool for evaluating ecosystem change. J Sustain For 2.1–2:87–111. https://doi.org/10.1300/J091v02n01_04
Morgan P, Keane RE, Dillon GK, Jain TB, Hudak AT, Karau EC, Sikkink PG, Holden ZA, Strand EK (2014) Challenges of assessing fire and burn severity using field measures, remote sensing and modelling. Int J Wildland Fire 23:1045–1060
Moritz MA, Morais ME, Summerell LA, Carlson JM, Doyle J (2005) Wildfires, complexity, and highly optimized tolerance. Proc Natl Acad Sci U S A 102:17912–17917
Mote PW, Li S, Lettenmaier DP, Xiao M, Engel R (2018) Dramatic declines in snowpack in the western US. NPJ Clim Atmos Sci 1:2
North MP, Stevens JT, Greene DF, Coppoletta M, Knapp EE, Latimer AM, Restaino CM, Tompkins RE, Welch KR, York RA, Young DJ (2019) Tamm review: reforestation for resilience in dry western U.S. forests. For Ecol Manag 432:209–224
Nowacki GJ, Abrams MD (2008) The demise of fire and "mesophication" of forests in the eastern United States. Bioscience 58:123–138
Nowacki GJ, Abrams MD (2015) Is climate an important driver of post-European vegetation change in the eastern United States? Glob Chang Biol 21:314–334
Olson J (1981) Carbon balance in relation to fire regimes. In: Mooney HA, Bonnicksen TM, Christensen NL, Lotan JE, Reiners WA (tech coords) Proceedings of the conference fire regimes and ecosystem properties. USDA Forest Service Gen Tech Rep WO-26, Honolulu, Hawaii, pp 327–378
Page WG, Jenkins MJ, Runyon JB (2012) Mountain pine beetle attack alters the chemistry and flammability of lodgepole pine foliage. Can J For Res 42:1631–1647
Parks SA, Holsinger LM, Miller C, Nelson CR (2015) Wildland fire as a self-regulating mechanism: the role of previous burns and weather in limiting fire progression. Ecol Appl 25:1478–1492
Pausas J, Vallejo VR (1999) The role of fire in European Mediterranean ecosystems. In: Chuvieco E (ed) Remote sensing of large wildfires. Springer, Berlin Heidelberg, pp 3–16
Pausas JG, Lamont BB, Paula S, Appezzato-da-Glória B, Fidelis A (2018) Unearthing belowground bud banks in fire-prone ecosystems. New Phytol 217:1435–1448

Perry DA, Hessburg PF, Skinner CN, Spies TA, Stephens SL, Taylor AH, Franklin JF, McComb B, Riegel G (2011) The ecology of mixed-severity fire regimes in Washington, Oregon, and Northern California. For Ecol Manag 262:703–717

Peters MP, Iverson LR (2019) Projected drought for the conterminous United States in the 21st century. In: Vose JM, Peterson DL, Luce CH, Patel-Weynand, T (eds) Effects of drought on forests and rangelands in the United States: translating science into management responses. USDA Forest Service Gen Tech Rep WO-98, Washington, DC, pp 19–39

Peterson GD (2002) Contagious disturbance, ecological memory, and the emergence of landscape pattern. Ecosystems 5:329–338

Peterson DL, Halofsky JE (2019) Moving the target: managing Pacific northwest forests in a changing climate. Douglasia 43:2–6

Peterson DL, Halofsky J, Johnson MC (2011a) Managing and adapting to changing fire regimes in a warmer climate. In: McKenzie D, Miller C, Falk DA (eds) The landscape ecology of fire. Springer, New York, pp 249–267

Peterson DL, Millar CI, Joyce LA, Furniss MJ, Halofsky JE, Neilson RP, Morelli TL (2011b) Responding to climate change in national forests: a guidebook for developing adaptation options. USDA Forest Service Gen Tech Rep PNW-855, Portland, Oregon

Platt WJ, Evans GW, Davis MM (1988) Effects of fire season on flowering of forbs and shrubs in longleaf pine forests. Oecologia 76(3):353–363

Prestemon JP, Shankar U, Xiu A, Talgo K, Yang D, Dixon E, McKenzie D, Abt KL (2016) Projecting wildfire area burned in the South-Eastern United States 2011–60. Int J Wildland Fire 25(7):715–729

Reilly MJ, Elia M, Spies TA, Gregory MJ, Sanesi G, Lafortezza R (2018) Cumulative effects of wildfires on forest dynamics in the eastern Cascade Mountains, USA. Ecol Appl 28:291–308

Reinhardt ED, Keane RE, Calkin DE, Cohen JD (2008) Objectives and considerations for wildland fuel treatment in forested ecosystems of the interior western United States. For Ecol Manag 256(12):1997–2006

Reynolds RT, Meador AJ, Youtz JA, Nicolet T, Matonis MS, Jackson PL, DeLorenzo DG, Graves AD (2013) Restoring composition and structure in southwestern frequent-fire forests. USDA Forest Service Gen Tech Rep RMRS-310, Fort Collins, Colorado

Rice KJ, Black RA, Radamaker G, Evans RD (1992) Photosynthesis, growth, and biomass allocation in habitat ecotypes of cheatgrass (*Bromus tectorum*). Funct Ecol 6:32–40

Ricotta C, Avena G, Marchetti M (1999) The flaming sandpile: Self-organized criticality and wildfires. Ecol Model 119:73–77

Riley KL, Loehman RA (2016) Mid-21st-century climate changes increase predicted fire occurrence and fire season length, Northern Rocky Mountains, United States. Ecosphere 7(11):e01543

Ryu J-H, Hayhoe K (2017) Observed and CMIP5 modeled influence of large-scale circulation on summer precipitation and drought in the South-Central United States. Clim Dyn 49:4293–4310

Scheffer M, Hirota M, Holmgren M, Van Nes EH, Chapin FS (2012) Thresholds for boreal biome transitions. Proc Natl Acad Sci U S A 109(52):21384–21389

Schlaepfer DR, Lauenroth WK, Bradford JB (2014) Natural regeneration processes in big sagebrush (*Artemisia tridentata*). Rangel Ecol Manag 67(4):344–357

Schoennagel T, Turner MG, Romme WH (2003) The influence of fire interval and serotiny on postfire lodgepole pine density in Yellowstone National Park. Ecology 84(11):2967–2978

Schwilk DW, Keeley JE, Knapp EE, McIver J, Bailey JD, Fettig CJ, Fiedler CE, Harrod RJ, Moghaddas JJ, Outcalt KW, Skinner CN (2009) The National Fire and fire surrogate study: effects of alternative fuel reduction methods on forest structure. Ecol Appl 19:285–304

Scott AC, Bowman DMJS, Bond WJ, Pyne SJ, Alexander ME (2014) Fire on Earth: an introduction. Wiley, Chichester

Seidl R, Spies TA, Peterson DL, Stephens SL, Hicke JA (2016) Searching for resilience: addressing the impacts of changing disturbance regimes on forest ecosystem services. J Appl Ecol 53(1):120–129

Shinneman DJ, Baker WL (1997) Nonequilibrium dynamics between catastrophic disturbances and old-growth forests in ponderosa pine landscapes of the Black Hills. Conserv Biol 11:1276–1288

Singleton MP, Thode AE, Meador AJ, Iniguez JM (2019) Increasing trends in high-severity fire in the southwestern USA from 1984 to 2015. For Ecol Manag 433:709–719

Skinner CN, Chang C-R (1996) Fire regimes, past and present. University of California, Davis

Slocum MG, Platt WJ, Beckage B, Panko B, Lushine JB (2007) Decoupling natural and anthropogenic fire regimes: a case study in Everglades National Park, Florida. Nat Areas J 27(1):41–55

Stambaugh MC, Marschall JM, Guyette RP (2014) Linking fire history to successional changes of xeric oak woodlands. For Ecol Manag 320:83–95

Stambaugh MC, Creacy G, Sparks J, Rooney M (2017) Three centuries of fire and forest vegetation transitions preceding Texas' most destructive wildfire: lost pines or lost oaks? For Ecol Manag 396:91–101

Stephens SL, McIver JD, Boerner RE, Fettig CJ, Fontaine JB, Hartsough BR, Kennedy PL, Schwilk DW (2012) Effects of forest fuel-reduction treatments in the United States. Bioscience 62:549–560

Stephens SL, Collins BM, Fettig CJ, Finney MA, Hoffman CM, Knapp EE, North MP, Safford H, Wayman RB (2018) Drought, tree mortality, and wildfire in forests adapted to frequent fire. Bioscience 68:77–88

Stevens-Rumann CS, Kemp KB, Higuera PE, Harvey BJ, Rother MT, Donato DC, Morgan P, Veblen TT (2018) Evidence for declining forest resilience to wildfires under climate change. Ecol Lett 21(2):243–252

Stocks BJ (1987) Fire potential in the spruce budworm-damaged forests of Ontario. For Chron 63:8–14

Stroh ED, Struckhoff MA, Stambaugh MC, Guyette RP (2018) Fire and climate suitability for woody vegetation communities in the south central United States. Fire Ecol 14:106–124

Swetnam TW (1990) Fire history and climate in the southwestern United States. In: Krammes JS (ed) Effects of fire management of southwestern natural resources. USDA Forest Service Gen Tech Rep RM-191, Fort Collins, Colorado, pp 6–17

Swetnam TW, Betancourt JL (1990) Fire-southern oscillation relations in the southwestern United States. Science 249:1017–1020

Swetnam TW, Betancourt JL (1998) Mesoscale disturbance and ecological response to decadal climatic variability in the American southwest. J Clim 11:3128–3147

Syphard AD, Sheehan T, Rustigian-Romsos H, Ferschweiler K (2018) Mapping future fire probability under climate change: does vegetation matter? PLoS One 13:e0201680

Tauer CG, Stewart JF, Will RE, Lilly CJ, Guldin JM, Nelson CD (2012) Hybridization leads to loss of genetic integrity in shortleaf pine: unexpected consequences of pine management and fire suppression. J For 110(4):216–224

Terando AJ, Reich B, Pacifici K, Costanza J, McKerrow A, Collazo JA (2016) Uncertainty quantification and propagation for projections of extremes in monthly area burned under climate change: A case study in the Coastal Plain of Georgia, USA. Natural hazard uncertainty assessment: Modeling and decision support, pp 223–245

Turetsky MR, Louis VLS (2006) Disturbance in boreal peatlands. In: Wieder RK, Vitt DH (eds) Boreal peatland ecosystems. Springer, Berlin, Heidelberg, pp 359–379

USGCRP (US Global Change Research Program) (2017) Climate science special report: 4th national climate assessment, vol 1. US Global Change Research Program, Washington, DC

Van Mantgem PJ, Stephenson NL, Byrne JC, Daniels LD, Franklin JF, Fulé PZ, Harmon ME, Larson AJ, Smith JM, Taylor AH, Veblen TT (2009) Widespread increase of tree mortality rates in the western United States. Science 323:521–524

Van Vuuren DP, Edmonds J, Kainuma M, Riahi K, Thomson A, Hibbard K, Hurtt GC, Kram T, Krey V, Lamarque JF, Masui T (2011) The representative concentration pathways: an overview. Clim Chang 109:5

Vanderwel MC, Purves D (2014) How do disturbances and environmental heterogeneity affect the pace and forest distribution shifts under climate change? Ecography 37:10–20

Veblen TT, Kitzberger T, Donnegan J (2000) Climatic and human influences on fire regimes in ponderosa pine forests in the Colorado front range. Ecol Appl 10:1178–1195

Vose JM, Elliott KJ (2016) Oak, fire, and global change in the eastern USA: what might the future hold? Fire Ecol 12(2):160–179

Vose RS, Easterling DR, Kunkel KE (2017) Temperature changes in the United States. In: Wuebbles DJ, Fahey DW, Hibbard KA, Dokken DJ, Stewart BC, Maycock TK (eds) Climate science special report: 4th National Climate Assessment, vol 1. US Global Change Research Program, Washington, DC, pp 185–206

Vose JM, Peterson DL, Domke GM, Fettig CJ, Joyce LA, Keane RE, Luce CH, Prestemon JE (2018) Forests. In: Reidmiller DR, Avery CW, Easterling DR, Kunkel KE, Lewis KLM, Maycock TK, Stewart BC (eds) Impacts, risks, and adaptation in the United States: 4th national climate assessment, vol 2. US Global Change Research Program, Washington, DC, pp 223–258

Wehner MF, Arnold JR, Knutson AN (2017) Droughts, floods, and wildfires. In: Wuebbles DJ, Fahey DW, Hibbard KA, Dokken DJ, Stewart BC, Maycock TK (eds) Climate science special report: 4th national climate assessment, vol 1. US Global Change Research Program, Washington, DC, pp 231–256

Westerling AL (2016) Increasing western US forest wildfire activity: sensitivity to changes in the timing of spring. Phil Trans R Soc B37120150178. https://doi.org/10.1098/rstb.2015.0178

Westerling AL, Hidalgo HG, Cayan DR, Swetman TW (2006) Warming and earlier spring increase western US forest wildfire activity. Science 313:940–943

Whelan RJ (1995) The ecology of fire. Cambridge University Press, Cambridge

Xiao J, Zhuang Q (2007) Drought effects on large fire activity in Canadian and Alaskan forests. Environ Res Lett 2(4):044003

Zelikova TJ, Hufbauer RA, Reed SC, Wertin T, Fettig C, Belnap J (2013) Eco-evolutionary responses of *Bromus tectorum* to climate change: implications for biological invasions. Ecol Evol 3:1374–1387

Zhang Q, Chen W (2007) Fire cycle of the Canada's boreal region and its potential response to global change. J For Res 18. https://doi.org/10.1007/s11676-007-0010-3