

RESEARCH ARTICLE

Impacts of insect frass and cadavers on soil surface litter decomposition along a tropical forest temperature gradient

Bernice C. Hwang¹  | Christian P. Giardina² | Creighton M. Litton³ |
Kainana S. Francisco² | Cody Pacheco² | Naneaikalaula Thomas² | Tyler Uehara² |
Daniel B. Metcalfe^{1,4} 

¹Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden

²Pacific Southwest Research Station, USDA Forest Service, Institute of Pacific Islands Forestry, Hilo, Hawaii, USA

³Department of Natural Resources and Environmental Management, University of Hawai'i at Mānoa, Honolulu, Hawaii, USA

⁴Department of Ecology and Environmental Science, Umeå University, Umeå, Sweden

Correspondence

Bernice C. Hwang, Department of Physical Geography and Ecosystem Science, Lund University, Sölvegaten 12, Lund, 223 62, Sweden.

Email: bernice.hwang@nateko.lu.se

Funding information

H2020 European Research Council, Grant/Award Number: 682707; Hatch and McIntire-Stennis Programs, Grant/Award Number: AW00132-H, HAW01127-H, HAW00188-M and HAW01123-M; USDA Forest Service, Pacific Southwest Research Station

Abstract

Insect herbivores play important roles in shaping many ecosystem processes, but how climate change will alter the effects of insect herbivory are poorly understood. To address this knowledge gap, we quantified for the first time how insect frass and cadavers affected leaf litter decomposition rates and nutrient release along a highly constrained 4.3°C mean annual temperature (MAT) gradient in a Hawaiian montane tropical wet forest. We constructed litterbags of standardized locally sourced leaf litter, with some amended with insect frass + cadavers to produce treatments designed to simulate ambient (Control = no amendment), moderate (Amended-Low = 2×Control level), or severe (Amended-High = 11×Control level) insect outbreak events. Multiple sets of these litterbags were deployed across the MAT gradient, with individual litterbags collected periodically over one year to assess how rising MAT altered the effects of insect deposits on litter decomposition rates and nitrogen (N) release. Increased MAT and insect inputs additively increased litter decomposition rates and N immobilization rates, with effects being stronger for Amended-High litterbags. However, the apparent temperature sensitivity (Q_{10}) of litter decomposition was not clearly affected by amendments. The effects of adding insect deposits in this study operated differently than the slower litter decomposition and greater N mobilization rates often observed in experiments which use chemical fertilizers (e.g., urea, ammonium nitrate). Further research is required to understand mechanistic differences between amendment types. Potential increases in outbreak-related herbivore deposits coupled with climate warming will accelerate litter decomposition and nutrient cycling rates with short-term consequences for nutrient cycling and carbon storage in tropical montane wet forests.

KEYWORDS

insect herbivory, nitrogen mineralization, nutrient cycling, Q_{10}

TAXONOMY CLASSIFICATION

Biogeochemistry, Ecosystem ecology, Soil ecology

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.

1 | INTRODUCTION

Litter decomposition is a central process regulating carbon (C) and nutrient cycling within ecosystems, with impacts on both plant productivity and species composition (Knorr et al., 2005; Prescott, 2002). Understanding the mechanisms that control litter production and decomposition is, therefore, important to efforts seeking to accurately model C and N cycling. Litter quality strongly influences decomposition rates (Prescott, 2002), particularly in the initial phases of the decomposition process (Melillo et al., 1982). Furthermore, a global study of 70 published studies determined that C:N and total nutrient concentration (the sum of N, phosphorus, potassium, calcium, and magnesium concentrations) of litter explained 70.2% of the variability in litter decomposition rates, while MAT explained an additional 7.9% of the variability (Zhang et al., 2008). Theoretically, litter C:N should regulate net N release rates during litter decomposition because microbial decomposers release N only after their own N requirements have been met (Berg, 1986). Ultimately, patterns in net N release and immobilization may depend on the C:N of the decomposer organisms relative to that of the substrate, as well as N availability in the environment (Parton et al., 2007; and references therein).

Along with litter quality, climate has also long been shown to be a major driver of litter decomposition (Knorr et al., 2005, and references therein). Temperature modulates the rate at which microbial communities decompose leaf litter and mineralize nutrients through direct effects on their activity and indirect effects on community composition (García-Palacios et al., 2013; Strickland et al., 2015; Wall et al., 2008). Many studies show that litter decomposition rates increase with rising temperatures, with the proportional change per every 10°C increase in temperature being described by Q_{10} . Across studies, Q_{10} typically ranges from 2 to 3, with an additional variation often ascribed to the variation in litter quality, moisture availability, and microbial community composition (García-Palacios et al., 2013). Exogenous N may stimulate chemical reactions between polyphenols and amino compounds, forming recalcitrant chemicals with a higher activation energy that subsequently slows decomposition (Fog, 1988). In this context, the temperature sensitivity of litter decomposition should increase as the net activation energy required for decomposition increases (Fierer et al., 2005). Slowing of microbial activity at cooler temperatures may inhibit decomposition even further. However, most temperature effect studies have been single-factor investigations (Bothwell et al., 2014; Knorr et al., 2005), while multi-factor studies examining interactions between temperature and other variables are less common (Salinas et al., 2011). One important set of understudied interactions involves climate-mediated herbivore abundance and activity effects on litter decomposition (Kristensen et al., 2018).

Climate change and insect folivory can have significant consequences on soil processes in forests. Empirical studies and models predict insect outbreaks will intensify in response to climate change and other anthropogenic disturbances, including in tropical forests (Anderegg et al., 2015; Dyer et al., 2012; McDowell et al., 2018).

While climate can have direct effects on insect performance (Bale et al., 2002), rising temperatures can have highly variable indirect effects on background levels of insect herbivory, reducing consumption in some studies while increasing consumption in others (Lemoine et al., 2014, and references therein). Such changes in ambient and outbreak levels of insect abundance and herbivory will affect the amount of foliar material consumed by insects and so production of frass, cadavers, leachate, and greenfall (Schowalter et al., 2011). Through these pathways, even background levels of herbivory by invertebrates can release as much or more of some nutrients as other major sources, with the potential to alter soil processes (Metcalf et al., 2014). For example, frass can boost microbial growth (Frost & Hunter, 2004), thereby increasing litter decomposition rates (Zimmer & Topp, 2002), N mineralization rates, and N immobilization rates (Frost & Hunter, 2007; Lovett & Ruesink, 1995). Furthermore, relatively high concentrations of N and phosphorus (P) can be found in insect cadavers compared with frass (Kos et al., 2017; Ohmart et al., 1983). Given that the effect of invertebrate cadavers on plant growth and densities of aboveground and belowground organisms can be stronger than their living counterparts, the inclusion of insect cadavers in ecosystem process studies is important for understanding ecosystem level impacts (Kos et al., 2017).

Initial efforts to study temperature effects on herbivory rates have been species- and community-specific (Lemoine et al., 2014, and references therein). Comparatively less is known about how insect herbivores collectively affect ecosystem-level processes (Lindquist et al., 2012, but see Frost & Hunter, 2004). Preliminary work primarily in temperate regions provides compelling insights into the effects of insect herbivores and climate change on ecosystems such as reducing primary productivity and exacerbating invasive species problems (Couture et al., 2015; Ward & Masters, 2007). Multi-factor in situ research is now needed to understand how insect herbivore frass and cadavers impact ecosystem processes in the understudied tropics. To address this gap, our study examined the effects of insect deposits (frass + cadavers) on litter decomposition rates across a highly constrained MAT gradient within a native-dominated tropical montane wet forest in Hawai'i. To the best of our knowledge, this is the first study to investigate the effects of paired insect frass plus cadavers on litter decomposition across a temperature gradient. We used seven of nine previously established research plots spanning a 4.3°C MAT, 666-m elevation gradient (Giardina et al., 2014; Litton et al., 2020). Instead of manipulating the ambient temperature of entire forest stands, an elevation gradient can provide a more tractable approach to assess the equilibrated response of entire, intact systems to climate change (Moser et al., 2011). While multiple environmental variables often co-vary with temperature and elevation (Wood et al., 2012), the MAT gradient used in this study is ideally suited for asking questions about the effects of long-term, whole-forest temperature differences on ecosystem processes because potential drivers of ecosystem metabolism other than temperature (i.e., geological substrate, soil type and age, disturbance history, vegetation composition, soil moisture, and solar radiation) are nearly constant across elevation (Giardina et al., 2014; Litton et al., 2011; Selmants et al., 2016).

Within these seven MAT gradient plots, we examined for one year the decay of *Metrosideros polymorpha* leaf litter, a tree species that dominates native forests in this study system and across much of Hawai'i. Litterbags represent sets of three treatments were deployed in each of the seven plots, with three levels of insect deposits (control, low and high amounts of frass + cadavers). Our goal was to investigate how rates of leaf litter decomposition and N release from decomposing litter in a tropical montane wet forest may vary with rising temperature and additions of insect deposits.

According to a meta-analysis by Knorr et al. (2005) across several ecosystem types, N additions inhibited litter decomposition when fertilization rates were 2–20 times greater than the anthropogenic level of N deposition, when ambient N deposition was 5–10 kg ha⁻¹ yr⁻¹, or when litter contained high levels of lignin. In contrast, decomposition rates increased at field sites exposed to low ambient N deposition (<5 kg ha⁻¹ yr⁻¹) and for low-lignin litters. Based on this meta-analysis, we hypothesized that while higher MAT would stimulate litter decomposition (Bothwell et al., 2014; Salinas et al., 2011), higher amendments of insect deposits would inhibit litter decomposition (Knorr et al., 2005). Specifically, we expected that litter decomposition would slow at our site where ambient levels of N deposition are high at 17 kg ha⁻¹ yr⁻¹ (Carrillo et al., 2002), additions are at least 20 times the ambient level of N deposition, and litter quality is low to medium (Hobbie & Vitousek, 2000). Further, we expected that the decelerating effect of insect deposits on litter decomposition would be greater for high versus low amendment amounts. Finally, we hypothesized that N mineralization would increase with temperature (Bothwell et al., 2014; Salinas et al., 2011), while C:N of remaining plant material would remain stable for the unamended, control litter but would decrease for the amended litters. Given the strong influence of temperature on litter decomposition rates (Coûteaux et al., 1995), we expected MAT to interact with exogenous N availability to significantly influence litter decomposition rates. A significant MAT × amendment interaction might result in significant differences in temperature sensitivity of litter decomposition between amendments.

2 | STUDY SITE

We examined the effects of MAT and insect deposits on leaf litter decomposition along a highly constrained MAT gradient on the windward side of Maunakea Volcano, Hawai'i (Giardina et al., 2014; Litton et al., 2011, 2020). Airborne light detection and ranging measurements of forest structure and intensive ground-based survey techniques were used to select plots representing maximum aboveground biomass at a given elevation while maintaining dominant tree species, geology, soil type, soil moisture, and disturbance history constant (Table 1). Plots were considered centered within larger stands of similarly sized forest. Typical on the windward side of Hawai'i Island, all plots along the gradient are classified as aggrading *M. polymorpha* – *Acacia koa* forests (Asner et al., 2009; Litton et al., 2020). Together, high-canopy *M. polymorpha* and mid-canopy *Cheirodendron trigynum* tree species comprise 84–97% of tree basal area across the plots (Litton et al., 2011; Selmants et al., 2014). Soils across the gradient fall under the tephra-derived, moderate to well-drained hydrous, ferrihydritic/amorphous, isothermic/isomesic Acrudoxic Hydruclands classification (Giardina et al., 2014). The Akaka, Honokaa, Maile, and Pihihouna soil series and soil age of ~20,000 years found across all plots are moderately common for Hawai'i Island (Litton et al., 2020). Furthermore, soil pH, bulk density, base saturation, cation exchange, and soil water content are relatively constant across the plots (Litton et al., 2020; Lyu et al., 2021). Although mean annual rainfall changes across the MAT gradient (Table 1), mean monthly soil water content is nearly identical at all plots because declining rainfall is balanced by decreasing evapotranspiration driven by lower air temperatures with increasing elevation (Selmants et al., 2016). See Litton et al. (2011) for more site selection details. As a subset of the full MAT gradient, the seven plots used in this study span from 934 to 1600 m in elevation (13.0–17.3°C), with plots at the low end of the gradient in the Laupāhoehoe Unit of the Hawai'i Experimental Tropical Forest (19°56'15.3"N, 155°16'02.6"W) and at the high end in the Hakalau Forest National Wildlife Refuge (19°50'31.3"N,

TABLE 1 Site characteristics across a 4.3°C mean annual temperature gradient in closed-canopy tropical montane wet forests on the Island of Hawai'i.

Elevation (m)	Air temp. (°C) ^a	Rainfall (mm yr ⁻¹) ^b	Potential evapotranspiration (mm yr ⁻¹) ^c	Soil temp. (°C) ^a	Soil moisture (m ³ m ⁻³) ^a	Solar radiation (W m ⁻²) ^c
934	17.3	4292	2232	17.3	0.55	200.9
1024	16.7	3975	2214	16.3	0.57	202.4
1116	16.1	3433	2137	15.6	0.51	210.1
1204	15.5	3181	2211	15.5	0.40	214.5
1274	15.1	3101	2234	14.9	0.51	216.2
1468	13.8	4119	1888	13.6	0.55	202.6
1600	13.0	3282	1961	12.6	0.57	213.1

^aLitton et al. (2011).

^bGiambelluca et al. (2013).

^cGiambelluca et al. (2014).

155°17'35.2"W). The 4.3°C temperature change falls within the long-term projected change for these ecosystems under the RCP-8.5 scenario (Collins et al., 2013), and is close to the 4.4°C change projected for 2081–2100 under the SSPS-8.5 scenario described in the IPCC 2021 summary report (Masson-Delmotte et al., 2021). All seven permanent 20m×20m plots are within 11 km of each other. Insect outbreaks are known to occur periodically in the Hawaiian Islands, which can result in massive defoliation events with subsequent extensive N deposition (Banko et al., 2016).

2.1 | Experimental design

To minimize variation in litter nutrient concentrations, we used well-mixed litter composed entirely of *M. polymorpha* leaves collected monthly from litterfall trays installed at the mid-elevation (1116m) between 2010 and 2014 along the full MAT gradient (see Giardina et al., 2014). Litter was oven-dried (70°C), sorted, and stored in double-sealed bags. To generate the large amount of insect frass and cadavers required for this study, spongy moths (*Lymantria dispar*) were raised on a standard artificial diet at the Otis Laboratory, United States Department of Agriculture in Buzzards Bay, Massachusetts to produce dried and sterilized frass + cadaver treatment mixtures. Total N of *L. dispar* frass at the start of the experiment was 65.80 mg g⁻¹ with a C:N of 6.29. Total N of *L. dispar* cadavers was 151.00 mg g⁻¹ with a C:N of 3.32. Initial N concentration for our standardized *M. polymorpha* leaf litter was 9.20 ± 0.19 mg g⁻¹ with a C:N of 53.27.

We used a synthesis of insect energetics data (Wiegert & Petersen, 1983) to estimate two levels of frass + cadaver additions. Specifically, we assumed that 66% and 13% of insect ingested C would be transferred to frass and cadavers, respectively, resulting in our amendments being administered in a 5:1 ratio of insect frass to cadavers. Amendment levels represented ambient (Control), moderate (Amended-Low), and severe (Amended-High) natural insect outbreak events. We calculated the Amended-Low amount (331 kg⁻¹ ha⁻¹) using data for actual litterfall rates of 1000 g m⁻² y⁻¹ of live foliage at our 1116m plot (Litton et al., 2020) with the 34.6% foliage consumption rate being twice the background level of herbivory predicted for the sites (latitude ±1°, Zhang et al., 2016). We based the Amended-High application (1818 kg⁻¹ ha⁻¹) on what might be expected during a major outbreak (100% foliage consumed), while relying on the same assumptions about insect energetics.

We constructed litterbags (11 cm × 11 cm) out of 1.0 mm × 1.5 mm mesh fiberglass screen filled with 2.5 g of well-mixed senescent *M. polymorpha* leaves. Each of the seven MAT plots received 35 Control treatment litterbags (no amendments) and 35 litterbags receiving the Amendment-Low treatment (low level additions of insect frass + bodies). Due to limited resources, only a subset of plots (at elevations 934, 1204, and 1600m) received another set of 35 litterbags treated with an Amendment-High treatment. We placed a cluster of five litterbags (one litterbag per collection period) with each of

the two or three treatments tethered to a central stake in seven 5 m × 5 m subplots within each MAT plot, resulting in seven replicates for each treatment, collection period, and plot. Litterbags were placed at least 20 cm apart from one another.

We followed the decomposition rates of leaf litter in the seven MAT plots between March 2020 and March 2021, by collecting one litterbag of each treatment from each subplot at approximately 1, 3, 6, 9, and 12 months post-deployment. After each collection, we carefully brushed off any debris from litter (including any insect deposits), then oven-dried at 70°C, weighed, and finely ground litter for chemical analysis. We determined total C and N of initial, undecomposed leaf litter and of decomposed litter from the five collection periods by combustion using a Costech Elemental Analyzer at Lund University, Sweden (Costech Analytical Technologies; Valencia, CA USA).

2.2 | Data analysis

We performed all statistical analyses in the R 4.2.1 statistical environment. Using the lme4 package (Bates et al., 2015) and lmer Test package (Kuznetsova et al., 2017), we generated linear mixed models to perform hypothesis testing using type III ANOVA and Tukey HSD analyses. We used amendment treatment (fixed) and MAT (random) as factors to determine whether leaf litter decomposition rate (*k*) per year varied significantly as a function of MAT and amendment level (*n* = 7, Appendix A) after approximately 12 months. Steps were taken to check that normality assumptions were met before proceeding with analyses. For each treatment, we estimated *k* rates for the seven replicate litterbag groups within each MAT plot by fitting a single exponential decay function to the litter mass data (Olson, 2007) using nonlinear regression with initial mass fixed at the initial value of 2.5 g (Adair et al., 2010):

$$M_t = M_0 e^{kt} \quad (1)$$

where *M_t* is mass remaining at time *t* (in days) and *M₀* is initial mass. We fit the exponential decay function to the time series of litter mass loss using Excel Solver, by freely varying *k* until the root-mean-square error (RMSE) between predicted and observed litter mass over the five sampling intervals for each treatment and subplot was minimized (Appendix B). This procedure is identical to any commercial curve fitting software, but allowed us to explicitly specify the metric of fit between observed and predicted values and generate a single estimate of fit for all the data and not just the individual curves.

We calculated the temperature coefficient (*Q₁₀*) for rates of litter decomposition across the MAT gradient as

$$Q_{10} = (R_2/R_1)^{10/(T_2-T_1)} \quad (2)$$

where *Q₁₀* is the proportional change in *k* due to a 10°C increase in MAT, *R₁* and *R₂* are regression-derived estimates of *k* at the lowest and

highest MAT plots, and T_1 and T_2 are MAT values of the lowest (13°C) and highest (17.3°C) MAT plots.

To estimate the proportion of initial N remaining, we divided the mass of N in litter by the mass of N in the initial undecomposed litter.

3 | RESULTS

3.1 | Effects of MAT and insect deposit amendments on litter decomposition

For both litter Amendment and Control treatments, k increased with rising MAT (Figure 1a). Control litter k ranged from $1.73 \times 10^{-3} \text{ d}^{-1}$ to $2.94 \times 10^{-3} \text{ d}^{-1}$ across the MAT gradient and k was a positive function of MAT ($R^2 = .65$, $F = 21.522$, $\text{Ndf} = 1$, $\text{Ddf} = 47$, $p < .001$). Amended-Low k ranged from $2.45 \times 10^{-3} \text{ d}^{-1}$ to $3.85 \times 10^{-3} \text{ d}^{-1}$ across MAT, with k being a positive function of MAT ($R^2 = .73$, $F = 25.851$, $\text{Ndf} = 1$, $\text{Ddf} = 47$, $p < .008$). And k for the Amended-High treatment ranged from $1.83 \times 10^{-3} \text{ d}^{-1}$ to $3.38 \times 10^{-3} \text{ d}^{-1}$ across MAT, with k again being a positive function of MAT ($R^2 = .93$, $F = 12.771$, $\text{Ndf} = 1$, $\text{Ddf} = 19$, $p < .001$). Differences between treatments on litter mass remaining (averaged across MAT plots) over time can be found in Appendix C.

Decomposition rates increased additively with higher MAT and Amendment treatments with no interaction between Amendment treatments and MAT. Significant differences in k values were detected among treatments ($F = 15.361$, $\text{Ndf} = 2$, $\text{Ddf} = 102$, $p = .003$) and MAT ($F = 14.765$, $\text{Ndf} = 6$, $\text{Ddf} = 102$, $p < .001$), but not for treatment $\times$ MAT interactions ($F = 0.503$, $\text{Ndf} = 8$, $\text{Ddf} = 102$, $p = .852$). Specifically, after nearly 12 months of incubation (Figure 1a), the Amended-High litter lost 15.9% more litter mass than Control litter ($F = 6.569$, $\text{Ndf} = 3$, $\text{Ddf} = 36$, $p = .001$) and 11.4% more litter mass than Amended-Low litter ($F = 6.569$, $\text{Ndf} = 3$, $\text{Ddf} = 36$, $p = .058$) averaged across the lowest, middle, and highest MATs (Appendix D). Amended-Low litter, however, had lost only 6.8% more mass than Control litter after nearly 12 months averaged across MAT ($F = 6.569$, $\text{Ndf} = 7$, $\text{Ddf} = 84$, $p = .220$, Figure 2a). Yet, Amended-Low litter underwent larger cumulative litter mass loss at higher MATs, whereas mass loss for Amended-High litter was more variable across MAT (Figure 2a).

We found that additional decomposition caused by the Amended-High treatment was not proportional to the amount of additional insect deposit. That is, the Amended-High treatment received five times the amount of frass + cadaver than the Amended-Low treatment but additional decomposition in the Amended-High treatment was less than four-fold higher than Amended-Low decomposition (Figure 2b).

3.2 | Q_{10} of litter decomposition

Temperature sensitivity of litter was variable across treatments. The estimated Q_{10} values of k (mean $\pm$ SE) did not differ across the three treatments; Q_{10} was 2.66 ± 0.45 , 2.99 ± 0.45 , and 2.87 ± 0.41

for Control, Amended-Low, and Amended-High litter, respectively. Decomposition rates for Control, Amended-Low, and Amended-High treated litter increased significantly by $0.22 \times 10^{-3} \text{ d}^{-1}$, $0.28 \times 10^{-3} \text{ d}^{-1}$, and $0.33 \times 10^{-3} \text{ d}^{-1}$ for each 1°C increase in MAT, respectively (Figure 1a).

3.3 | Nitrogen dynamics in decomposing litter

Nitrogen was immobilized in litter across MAT in all Amendment treatments. Litter amended with insect deposits had accumulated significantly more N than Control litter by the end of the year ($F = 20.759$, $\text{Ndf} = 2$, $\text{Ddf} = 102$, $p < .001$), where Amended-High litter amassed the most N after 1 year (Figure 1b). The degree of decomposition-associated N accumulation significantly increased with rising MAT (Figure 1b, $F = 5.423$, $\text{Ndf} = 2$, $\text{Ddf} = 102$, $p < .001$), with no treatment $\times$ MAT interaction ($F = 1.138$, $\text{Ndf} = 8$, $\text{Ddf} = 102$, $p = .345$).

The C:N of litter amended at both levels was lower than control litter overall but did not decline with MAT at the High-Amended level. After 12 months, the C:N of Amended-High litter was significantly lower than Amended-Low litter ($F = 19.089$, $\text{Ndf} = 3$, $\text{Ddf} = 36$, $p < .001$). The C:N for both Amended-Low ($F = 19.089$, $\text{Ndf} = 7$, $\text{Ddf} = 84$, $p = .017$) and Amended-High ($F = 19.089$, $\text{Ndf} = 3$, $\text{Ddf} = 36$, $p < .001$) litter were significantly lower than C:N of the Control treatment. However, Amended-High litter C:N was relatively constant across the MAT gradient (Figure 1c). The effect of the Amendment treatments on litter C:N did not interact with MAT after twelve months of decomposition ($F = 1.402$, $\text{Ndf} = 8$, $\text{Ddf} = 102$, $p = .205$).

4 | DISCUSSION

Our elevation gradient provided a near-ideal design for investigating temperature-mediated effects on ecosystem processes because geological substrate, soil type and age, disturbance history, vegetation composition, soil moisture, and solar radiation were nearly constant across sites, allowing us to focus on rising temperatures. The consistency of these characteristics across plots is invaluable as in situ variation would confound study results. In line with prior studies across these sites (Bothwell et al., 2014; Giardina et al., 2014), we found that rising MAT stimulated litter decomposition rates (Figure 1a), which also aligned with prior findings of MAT-driven increases in soil-surface CO_2 efflux and litterfall along this gradient (Giardina et al., 2014; Litton et al., 2011, 2020). We found that insect deposits had little apparent effect on Q_{10} values for litter decomposition, with the lack of a significant increase in the temperature sensitivity of *M. polymorpha* litter decomposition. Our results suggest that the litter microbial community along the gradient has a wide thermal tolerance and are insensitive to amendments.

Our hypothesis that additional insect deposits would inhibit litter decomposition (Knorr et al., 2005) was not supported.




FIGURE 1 Insect frass + cadaver treatment and mean annual temperature (MAT) effects on: (a) litter decomposition rate (k), (b) N content, and (c) C:N across a 4.3°C MAT gradient after approximately 12 months of decomposition. Solid shapes denote means; bars denote ± 1 SE, $n = 7$. Dashed lines denote best-fitted lines and bands represent 95% confidence intervals.

However, the Knorr et al. (2005) meta-analysis did not examine the role of amendment type in litter decomposition across study systems. Hobbie and Vitousek (2000) conducted an amendment experiment in a nearby forest using similar litter as in our study, where they applied N chemically at $100 \text{ kg ha}^{-1} \text{ yr}^{-1}$, and found the additions had no significant effect on litter decomposition. They discussed that the demand for N may be low relative to supply because of the poor C quality of local litter, but also N fertilization may inhibit lignin degradation (see also Fog, 1988) and thereby offset stimulatory effects of N addition. Alternatively, some other factor may have constrained decomposition more strongly than N availability. That adding insect deposits accelerated litter decomposition in our study suggests that nutrient amendment as

fertilizer versus insect frass + cadavers can lead to different stimulatory effects and that these differences may be site-specific. Our study highlights that when nutrients are bound in organic forms, they can influence litter decomposition and nutrient release differently than when in inorganic fertilizer form, which is commonly used to study nutrient supplementation effects on ecosystem processes. Insect frass and cadavers can supply nutrients other than N, such as P, that also play a role in litter decomposition (Ohmart et al., 1983), and the passage of foliage through the gut of insects may alter nutrient quality or bacterial assemblages (Hammer et al., 2017; Lovett & Ruesink, 1995) that may then interact with litter, soil, and decomposers differently than do chemical fertilizers. A logical next step would be to conduct a fertilizer




FIGURE 2 The (a) variability and (b) proportionality of additional leaf litter mass loss relative to control treatment due to two levels of insect frass + cadavers addition after 12 months of decomposition along a 4.3°C mean annual temperature gradient. Solid shapes denote means, and error bars denote ± 1 SE, $n = 7$. Solid, dashed and dotted lines in panel b denote the 5:1, 10:1, and 30:1 lines for insect frass + cadavers mass added: Additional litter mass loss.

versus insect deposit study, where a comprehensive nutrient and microbial analysis of amendments and litter would be used to compare changes to litter decomposition processes.

Interestingly, the overall effects of MAT and our amendments were additive and not synergistic. Other studies on this gradient have demonstrated that rising MAT increases ecosystem C fluxes and soil nitrate availability (Giardina et al., 2014; Pierre et al., 2017), and alters the cycling, availability, and ecological stoichiometry of a suite of macro- and micronutrients (Litton et al., 2020). In agreement with the meta-analysis by Knorr et al. (2005), the lack of a significant Amendment interaction in our study indicates that MAT may set boundaries for but not consistently alter the effect of N enrichment on litter decomposition.

In contrast to this study, other studies along our MAT gradient and in nearby *M. polymorpha* dominated forests have not observed N immobilization with increasing MAT (Bothwell et al., 2014) or MAT plus fertilization treatments (Hobbie & Vitousek, 2000). Greater rainfall during this study (Hawaii Permanent Plot Network, 2022) compared to the Bothwell et al.'s (2014) study may have made more nutrients available to microbes or altered microbial community composition (but see Selmants et al., 2016) and thus litter decomposition rates. In addition, it may be relevant that our litterbags were filled with more *M. polymorpha* leaf litter per area (0.23 g cm^{-2}) than either the Bothwell et al.'s (2014) study (0.17 g cm^{-2}) or the Hobbie and Vitousek's (2000) study ($0.10\text{--}0.20 \text{ g cm}^{-2}$). It is possible that the amount of leaf litter in our study crossed a threshold above which microbial activity and thus immobilization would be stimulated (Liu et al., 2017). Initial N concentration of leaf litter in our study (0.92% N) is low for senescent leaves of most forest species (Parton et al., 2007). At these levels, observed N immobilization aligns with findings from a global synthesis of litter chemistry and decomposition, and based on initial litter chemistry, our litter may not have decomposed sufficiently to begin releasing substantial amounts of N (Parton et al., 2007). This is line with other studies that have observed N immobilization during early stages of decomposition, sometimes followed by mineralization at later stages of

decomposition (Zhou et al., 2018). Because the litter decomposition process is lengthy (Parton et al., 2007) and differences among treatments can be dynamic and complex (Chen et al., 2013; Melillo et al., 1982), a complete understanding of the fate of insect-derived inputs requires longer tracer-based experiments.

The influence of insect inputs on litter decomposition in our study may not be entirely explained by N availability and other factors such as limitation of other nutrients or decomposer community dynamics may co-vary with N to influence litter decomposition (Hobbie, 2005). For example, differences in P supply and litter N:P can influence litter decomposition, microbial communities, and litter nutrient dynamics (Güsewell & Gessner, 2009). In fact, a recent study along this gradient demonstrated that MAT affects the cycling and availability of essential macro- and micronutrients, as well as the ecological stoichiometry of live foliage and litterfall (Litton et al., 2020). Ohmart et al. (1983) demonstrated that insect frass can be a source of not only N, but also P and potassium. The quality of insect frass can also affect soil nutrient availability, with effects spanning from positive to negative (Kagata & Ohgushi, 2012; Valdés-Correcher et al., 2019). Such variation across MAT indicates that plot-specific insect material may bring other factors to an already complex decomposition process. We therefore recommend future studies of frass and cadaver effects on nutrient dynamics use insect inputs from resident insect populations feeding on resident vegetation. Further, it would be valuable to study the circumstances under which nutrients from insect inputs are or are not mobilized by microbial activity or retained within recalcitrant organic matter. Separating the effects of frass from cadavers in future studies could also help refine our understanding of insect herbivore impacts on soil processes.

In contrast to a pattern of decreasing C:N for control and Amended-Low treatments, C:N for Amended-High litter was stable across the three low, medium, and high MAT plots in which the treatment was tested (Figure 1c). For Amended-Low litter, the addition of exogenous N from insect deposits to litter may have stimulated microbes that were initially limited by N, leading to

greater litter decomposition and litter N immobilization but lower C:N – especially in the first month of decomposition (Appendix E). That C:N was relatively constant across MAT for Amended-High litter, suggests that C:N alone cannot explain accelerated rates of litter decomposition. When N is no longer limiting, MAT may have played a greater role in stimulating microbial activity and litter decomposition (Hobbie, 2005). Our results highlight the varying effect of MAT on litter decomposition processes under conditions of N limitation versus insect-mediated enrichment, and underscore the complexity of predicting changes in soil processes under a changing climate.

Our study focused on the effects of insect deposits and MAT on surface leaf litter decomposition along a well-constrained gradient for which confounding factors are largely controlled. However, because standing litter (litter that has not reached the soil surface) can contain more dissolved organic C than surface litter, the standing phase can play a critical role in litter decomposition and soil C storage (Wang et al., 2017). In one semi-arid forest, standing litter decomposed faster than soil surface litter due to high dissolved organic C and leaf litter moisture content derived from relatively high night-time humidity (Wang et al., 2017). Because of their importance especially in dry ecosystems, standing litter-insect interactions also deserve attention in future multi-factor studies.

This study highlights how insects and a warmer climate can combine to significantly impact ecosystem processes in tropical forests. Using a well-constrained gradient, our results confirm that rising MAT increases litter decomposition rates in a tropical montane wet forest. Further, they demonstrate that herbivores can accelerate decomposition rates, and that leaf litter decomposition responses to insect frass and cadavers may operate additively to those of MAT. Finally, our results indicate that increased insect deposits immobilize nutrients in tropical wet montane forests and can operate differently from chemical fertilizers, warranting further investigation into the mechanisms responsible for observed differences among control and amendment types. An understanding of insect responses to multiple and potentially interacting natural and anthropogenic changes is necessary to improve our understanding of ecosystem responses to environmental change.

AUTHOR CONTRIBUTIONS

Bernice C. Hwang: Conceptualization (lead); data curation (lead); formal analysis (lead); investigation (lead); methodology (lead); project administration (supporting); resources (supporting); software (lead); visualization (lead); writing – original draft (lead); writing – review and editing (lead). **Christian P. Giardina:** Funding acquisition (supporting); investigation (supporting); project administration (supporting); resources (supporting); supervision (supporting); validation (supporting); writing – review and editing (supporting). **Creighton M. Litton:** Funding acquisition (supporting); project administration (supporting); resources (supporting); writing – review and editing (supporting). **Kainana S. Francisco:** Data curation (supporting); investigation (supporting); project administration (supporting);

supervision (supporting); writing – review and editing (supporting). **Cody Pacheco:** Data curation (supporting); investigation (supporting); project administration (supporting). **Naneaikalaula Thomas:** Data curation (supporting); investigation (supporting); project administration (supporting). **Tyler Uehara:** Data curation (supporting); investigation (supporting); project administration (supporting). **Daniel B. Metcalfe:** Conceptualization (supporting); formal analysis (supporting); funding acquisition (lead); investigation (supporting); project administration (supporting); resources (lead); supervision (supporting); writing – review and editing (supporting).

ACKNOWLEDGMENTS

We thank the United States Department of Agriculture Forest Service and the State of Hawai'i Department of Land and Natural Resources, Division of Forestry and Wildlife for access to the Hawai'i Experimental Tropical Forest, and the United States Fish and Wildlife Service for access to the Hakalau Forest National Wildlife Refuge. We acknowledge the lands on which this study was conducted, lands that are sacred to Kānaka 'Ōiwi, the Native Hawaiian people. Many of these lands were ceded to the US Federal Government and then the State of Hawai'i without consent or compensation to Kānaka 'Ōiwi or their descendants. We also thank Karl Ljung at the Lund University Department of Geology for his assistance with running nutrient analyses, and Hannah Nadel at the USDA Otis Laboratory for providing insect frass and bodies. And the project would not have been possible without the on-the-ground support of Rebekah Ohara and the Akaka Foundation for Tropical Forests, and their research technicians, Chantilly Kelihoomalau, Kaulimakahikina Thomas, and Kaianui Andaya, who assisted with lab and field work. We also thank Alealani Evangelista, Chloe Martins-Kelihoomalau, and Kirie Santos, Kupu Conservation Leadership Development Program members with the USDA Forest Service, who assisted with field work. The establishment of the original MAT gradient was funded by the College of Tropical Agriculture and Human Resources, University of Hawai'i at Mānoa via the USDA National Institute of Food and Agriculture, Hatch and McIntire-Stennis Programs (HAW00132-H, HAW01127-H, HAW00188-M, and HAW01123-M). Finally, we thank the European Research Council (ECOHERR, no. 682707) for funding the study and the USDA Forest Service, Pacific Southwest Research Station for additional financial and logistical support.

CONFLICT OF INTEREST

The authors have no competing interests or conflict of interest to declare.

DATA AVAILABILITY STATEMENT

Data are publicly available at <https://doi.org/10.6084/m9.figshare.19085612.v1>.

ORCID

Bernice C. Hwang  <https://orcid.org/0000-0002-3744-9771>

Daniel B. Metcalfe  <https://orcid.org/0000-0001-8325-9269>

REFERENCES

- Adair, E. C., Hobbie, E. S., & Hobbie, R. K. (2010). Single-pool exponential decomposition models: Potential pitfalls in their use in ecological studies. *Ecology*, 91, 1225–1236.
- Anderegg, W. R., Hicke, J. A., Fisher, R. A., Allen, C. D., Aukema, J., Bentz, B., Hood, S., Lichstein, J. W., Macalady, A. K., McDowell, N., & Pan, Y. (2015). Tree mortality from drought, insects, and their interactions in a changing climate. *New Phytologist*, 20, 674–683.
- Asner, G. P., Hughes, R. F., Varga, T. A., Knapp, D. E., & Kennedy-Bowdoin, T. (2009). Environmental and biotic controls over aboveground biomass throughout a tropical rain forest. *Ecosystems*, 12, 261–278.
- Bale, J. S., Masters, G. J., Hodkinson, I. D., Awmack, C., Bezemer, T. M., Brown, V. K., Butterfield, J., Buse, A., Coulson, J. C., Farrar, J., & Good, J. E. (2002). Herbivory in global climate change research: Direct effects of rising temperature on insect herbivores. *Global Change Biology*, 8, 1–6.
- Banko, P., Peck, R., Yelenik, S., Paxton, E., Bonaccorso, F., Montoya-Aiona, K., & Foote, D. (2016). *Dynamics and ecological consequences of the 2013–2014 Koa moth outbreak at Hakalau Forest National Wildlife Refuge. Hawaii cooperative studies unit technical report HCSU-058*. University of Hawaii Hilo.
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67, 1–48.
- Berg, B. (1986). Nutrient release from litter and humus in coniferous forest soils—A mini review. *Scandinavian Journal of Forest Research*, 1, 359–369.
- Bothwell, L. D., Selman, P. C., Giardina, C. P., & Litton, C. M. (2014). Leaf litter decomposition rates increase with rising mean annual temperature in Hawaiian tropical montane wet forests. *PeerJ*, 2, e685.
- Carrillo, J. H., Hastings, M. G., Sigman, D. M., & Huebert, B. J. (2002). Atmospheric deposition of inorganic and organic nitrogen and base cations in Hawaii. *Global Biogeochemical Cycles*, 16, 24–1–24–16.
- Chen, H., Dong, S., Liu, L., Ma, C., Zhang, T., Zhu, X., & Jiangming, M. (2013). Effects of experimental nitrogen and phosphorus addition on litter decomposition in an old-growth tropical forest. *PLoS One*, 8, e8410.
- Collins, M., Knutti, R., Arblaster, J., Dufresne, J. L., Fichefet, T., Friedlingstein, P., Gao, X., Gutowski, W. J., Johns, T., Krinner, G., & Shongwe, M. (2013). Long-term climate change: Projections, commitments and irreversibility. In T. F. Stocker, D. Qin, G.-K. Plattner, M. M. B. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, & P. M. Midgley (Eds.). *Climate change 2013-the physical science basis: Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change* (pp. 1056). Cambridge University Press.
- Coûteaux, M., Böttner, P., & Berg, B. (1995). Litter decomposition, climate, and litter quality. *Trends in Ecology and Evolution*, 10, 63–66.
- Couture, J. J., Meehan, T. D., Kruger, E. L., & Lindroth, R. L. (2015). Insect herbivory alters impact of atmospheric change on northern temperate forests. *Nature Plants*, 1, 150–160.
- Dyer, L. A., Carson, W. P., & Leigh, E. J., Jr. (2012). Insect outbreaks in tropical forests: Patterns, mechanisms, and consequences. *Insect Outbreaks Revisited*, 10, 219–245.
- Fierer, N., Craine, J. M., McLaughlin, K., & Schimel, J. P. (2005). Litter quality and the temperature sensitivity of decomposition. *Ecology*, 86, 320–326.
- Fog, K. (1988). The effect of added nitrogen on the rate of decomposition of organic matter. *Biological Reviews*, 63, 433–462.
- Frost, C. J., & Hunter, M. D. (2004). Insect canopy herbivory and frass deposition affect soil nutrient dynamics and export in oak mesocosms. *Ecology*, 85, 3335–3347.
- Frost, C. J., & Hunter, M. D. (2007). Recycling of nitrogen in herbivore feces: Plant recovery, herbivore assimilation, soil retention, and leaching losses. *Oecologia*, 151, 42–53.
- García-Palacios, P. F., Maestre, T., Kattge, J., Wall, D. H., & Klironomos, J. (2013). Climate and litter quality differently modulate the effects of soil fauna on litter decomposition across biomes. *Ecology Letters*, 16, 1045–1053.
- Giambelluca, T. W., Chen, Q., Frazier, A. G., Price, J. P., Chen, Y.-L., Chu, P.-S., Eischeid, J. K., & Delparte, D. M. (2013). Online rainfall atlas of Hawai'i. *Bulletin of the American Meteorological Society*, 94, 313–316.
- Giambelluca, T. W., Shuai, X., Barnes, M. L., & Alliss, R. J. (2014). *Evapotranspiration of Hawaii: Final report. US Army Corps of Engineers - Honolulu District and the commission on water resource management*. State of Hawaii.
- Giardina, C. P., Litton, C. M., Crow, S. E., & Asner, G. P. (2014). Warming-related increases in soil CO₂ efflux are explained by increased below-ground C flux. *Nature Climate Change*, 4, 822–827.
- Güsewell, S., & Gessner, M. O. (2009). N:P ratios influence litter decomposition and colonization by fungi and bacteria in microcosms. *Functional Ecology*, 23, 211–219.
- Hammer, T. J., Janzen, D. H., Hallwachs, W., Jaffe, S. P., & Fierer, N. (2017). Caterpillars lack a resident gut microbiome. *Proceedings of the National Academy of Sciences of the United States of America*, 114, 9641–9646.
- Hawaii Permanent Plot Network (2022). Climatological data summaries. Retrieved from Ostertag, R., Cordell, S., Giambelluca, T., Giardina, C., Litton, C., Nullet, M., Sack, L. on 9 February, 2022.
- Hobbie, S. E. (2005). Contrasting effects of substrate and fertilizer nitrogen on the early stages of litter decomposition. *Ecosystems*, 8, 644–656.
- Hobbie, S. E., & Vitousek, P. M. (2000). Nutrient limitation of decomposition in Hawaiian forests. *Ecology*, 81, 1867–1877.
- Kagata, H., & Ohgushi, T. (2012). Positive and negative impacts of insect frass quality on soil nitrogen availability and plant growth. *Population Ecology*, 54, 75–82.
- Knorr, M., Frey, S. D., & Curtis, P. S. (2005). Nitrogen additions and litter decomposition: A meta-analysis. *Ecology*, 86, 3252–3257.
- Kos, M., Jing, J., Keesmaat, I., Declerck, S. A. J., Wagenaar, R., & Bezemer, T. M. (2017). After-life effects: Living and dead invertebrates differentially affect plants and their associated above- and belowground multitrophic communities. *Oikos*, 126, 888–899.
- Kristensen, J. A., Metcalfe, D. B., & Rousk, J. (2018). The biogeochemical consequences of litter transformation by insect herbivory in the subarctic: A microcosm simulation experiment. *Biogeochemistry*, 138, 323–336.
- Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, 82, 1–26.
- Lemoine, N. P., Burkepille, D. E., & Parker, J. D. (2014). Variable effects of temperature on insect herbivory. *PeerJ*, 2, e376.
- Lindquist, E. J., D'Annunzio, R., Gerrand, A., MacDicken, K., Achard, F., Beauchle, R., Bringk, A., Eva, H. D., Mayau, P., San-Miguel-Ayaz, J., & Stibig, H. J. (2012). *Global forest land-use change 1990–2005*. FAO Forestry Paper No. 169. Food and Agriculture Organization of the United Nations and European Commission Joint Research Centre.
- Litton, C. M., Giardina, C. P., Albano, J. K., Long, M. S., & Asner, G. P. (2011). The magnitude and variability of soil-surface CO₂ efflux increase with mean annual temperature in Hawai'ian tropical Montane wet forests. *Soil Biology and Biochemistry*, 43, 2315–2323.
- Litton, C. M., Giardina, C. P., Freeman, K. R., Selman, P. C., & Sparks, J. P. (2020). Impact of mean annual temperature on nutrient availability in a tropical montane wet forest. *Frontiers in Plant Science*, 11, 784.
- Liu, X.-J. A., Sun, J., Mau, R. L., Finley, B. K., Compson, Z. G., van Gestel, N., Brown, J. R., Schwartz, E., Dijkstra, P., & Hungate, B. A. (2017). Labile carbon input determines the direction and magnitude of the priming effect. *Applied Soil Ecology*, 109, 7–13.
- Lovett, G. M., & Ruesink, A. E. (1995). Carbon and nitrogen mineralization from decomposing gypsy moth frass. *Oecologia*, 104, 133–138.

- Lyu, M., Giardina, C. P., & Litton, C. M. (2021). Interannual variation in rain-fall modulates temperature sensitivity of carbon allocation and flux in a tropical montane wet forest. *Global Change Biology*, 16, 3824–3836.
- Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., & Zhou, B. (2021). Climate Change 2021: the Physical Science Basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change. 2021:2.
- McDowell, N., Allen, C. D., Anderson-Teixeira, K., Brando, P., Brien, R., Chambers, J., Christofferson, B., Davies, S., Doughty, C., Duque, A., & Espirito-Santo, F. (2018). Drivers and mechanisms of tree mortality in moist tropical forests. *New Phytologist*, 219, 851–869.
- Melillo, J. M., Aber, J. D., & Muratore, J. F. (1982). Nitrogen and lignin control of hardwood leaf litter decomposition dynamics. *Ecology*, 63, 621–626.
- Metcalfe, D. B., Asner, G. P., Martin, R. E., Silva Espejo, J. E., Huasco, W. H., Farfán Amézquita, F. F., Carranza-Jimenez, L., Galiano Cabrera, D. F., Baca, L. D., Sinca, F., & Huaraca Quispe, L. P. (2014). Herbivory makes major contributions to ecosystem carbon and nutrient cycling in tropical forests. *Ecology Letters*, 17, 324–332.
- Moser, G., Leuschner, C., Hertel, D., Graefe, S., Soethe, N., & Iost, S. (2011). Elevation effects on the carbon budget of tropical mountain forests (S Ecuador): The role of the belowground compartment. *Global Change Biology*, 17, 2211–2226.
- Ohmart, C. P., Stewart, L. G., & Thomas, J. R. (1983). Leaf consumption by insects in three eucalyptus forest types in southeastern Australia and their role in short-term nutrient cycling. *Oecologia*, 59, 322–330.
- Olson, J. S. (2007). Energy storage and the balance of producers and decomposers in ecological systems. *Ecology*, 44, 322–331.
- Parton, W., Silver, W. L., Burke, I. C., Grassens, L., Harmon, M. E., Currie, W. S., King, J. Y., Adair, E. C., Brandt, L. A., Hart, S. C., & Fasth, B. (2007). Global-scale similarities in nitrogen release patterns during long-term decomposition. *Science*, 315, 361–364.
- Pierre, S., Hewson, U., Sparks, J. P., Litton, C. M., Giardina, C. P., Groffman, P. M., & Fahey, T. J. (2017). Ammonia oxidizer populations vary with nitrogen cycling across a tropical montane mean annual temperature gradient. *Ecology*, 98, 1896–1907.
- Prescott, C. E. (2002). The influence of the forest canopy on nutrient cycling. *Tree Physiology*, 22, 1193–1200.
- Salinas, N., Malhi, Y., & Meir, P. (2011). The sensitivity of tropical leaf litter decomposition to temperature: Results from a large-scale leaf translocation experiment along an elevation gradient in Peruvian forests. *New Phytologist*, 189, 967–977.
- Schowalter, T. D., Fonte, S. J., Geaghan, J., & Wang, J. (2011). Effects of manipulated herbivore inputs on nutrient flux and decomposition in a tropical rainforest in Puerto Rico. *Oecologia*, 4, 1141–1149.
- Selmants, P. C., Litton, C. M., Giardina, C. P., & Asner, G. P. (2014). Ecosystem carbon storage does not vary with mean annual temperature in Hawaiian tropical montane wet forests. *Global Change Biology*, 20, 2927–2937.
- Selmants, P. C., Adair, K. L., Litton, C. M., Giardina, C. P., & Schwartz, E. (2016). Increases in mean annual temperature do not alter soil bacterial community structure in tropical montane wet forests. *Ecosphere*, 7(4), e01296.
- Strickland, M. S., Keiser, A. D., & Bradford, M. A. (2015). Climate history shapes contemporary leaf litter decomposition. *Biogeochemistry*, 122, 165–174.
- Valdés-Correcher, E., Sitters, J., Wassen, M., Brion, N., & Venterink, H. O. (2019). Herbivore dung quality affects plant community diversity. *Scientific Reports*, 9, 1–6.
- Wall, D. H., Bradford, M. A., St. John, M. G., Trofymow, J. A., Behan-Pelletier, V., Bignell, D. E., Dangerfield, J. M., Parton, W. J., Rusek, J., Voigt, W., & Wolters, V. (2008). Global decomposition experiment shows soil animal impacts on decomposition are climate-dependent. *Global Change Biology*, 14, 2661–2677.
- Wang, J., Liu, L., Wang, X., Yang, S., Zhang, B., Li, P., Qiao, C., Deng, M., & Liu, W. (2017). High night-time humidity and dissolved organic carbon content support rapid decomposition of standing litter in a semi-arid landscape. *Functional Ecology*, 31, 1659–1668.
- Ward, N. L., & Masters, G. J. (2007). Linking climate change and species invasion: An illustration using insect herbivores. *Global Change Biology*, 13, 1605–1161.
- Wiegert, R. G., & Petersen, C. E. (1983). Energy transfer in insects. *Annual Review of Entomology*, 28, 455–486.
- Wood, T. E., Cavaleri, M. A., & Reed, S. C. (2012). Tropical forest carbon balance in a warmer world: A critical review spanning microbial- to ecosystem-scale processes. *Biological Reviews*, 87, 912–927.
- Zhang, D., Hui, D., Luo, Y., & Guoyi, Z. (2008). Rates of litter decomposition in terrestrial ecosystems: Global patterns and controlling factors. *Journal of Plant Ecology*, 1, 85–93.
- Zhang, S., Zhang, Y., & Ma, K. (2016). Latitudinal variation in herbivory: hemispheric asymmetries and the role of climatic drivers. *Journal of Ecology*, 104, 1089–1095.
- Zhou, G., Zhang, J., Qiu, X., Wei, F., & Xu, X. (2018). Decomposing litter and associated microbial activity responses to nitrogen deposition in two subtropical forests containing nitrogen-fixing or non-nitrogen-fixing tree species. *Scientific Reports*, 8, 12934.
- Zimmer, M., & Topp, W. (2002). The role of coprophagy in nutrient release from feces of phytophagous insects. *Soil Biology and Biogeochemistry*, 34, 1093–1099.

How to cite this article: Hwang, B. C., Giardina, C. P., Litton, C. M., Francisco, K. S., Pacheco, C., Thomas, N., Uehara, T., & Metcalfe, D. B. (2022). Impacts of insect frass and cadavers on soil surface litter decomposition along a tropical forest temperature gradient. *Ecology and Evolution*, 12, e9322. <https://doi.org/10.1002/ece3.9322>


APPENDIX A

ANOVA and Tukey HSD results for litter decomposition (k) using entire time series. Statistically significant changes (95%) are indicated in bold.

ANOVA	Ndf	Ddf	F-value	P-value
Treatment	2	102	15.361	0.003
Temperature	6	102	14.765	< 0.001
Treatment x temperature	8	102	0.503	0.852
Tukey HSD between treatments	Ndf	Ddf		Adjusted P-value
Low : control	7	84		0.024
High : control	4	36		< 0.001
High : low	4	36		0.002
Tukey HSD between temperatures	Ndf	Ddf		Adjusted P-value
13.8 : 13.0	2	24		1.000
15.1 : 13.0	2	24		0.110
15.5 : 13.0	3	36		< 0.001
16.1 : 13.0	2	24		0.034
16.7 : 13.0	2	24		< 0.001
17.3 : 13.0	3	36		< 0.001
15.1 : 13.8	2	24		0.109
15.5 : 13.8	2	24		< 0.001
16.1 : 13.8	2	24		0.037
16.7 : 13.8	2	24		< 0.001
17.3 : 13.8	2	24		< 0.001
15.5 : 15.1	2	24		0.057
16.1 : 15.1	2	24		1.000
16.7 : 15.1	2	24		0.051
17.3 : 15.1	2	24		0.065
16.1 : 15.5	2	24		0.170
16.7 : 15.5	2	24		1.000
17.3 : 15.5	3	36		1.000
16.7 : 16.1	2	24		0.143
17.3 : 16.1	2	24		0.189
17.3 : 16.7	2	24		1.000


APPENDIX B

Relation between litter mass observed and predicted with the exponential decay model fit the pattern of mass loss over five sampling intervals for each treatment and subplot. The solid black line indicates a 1:1 relationship; the dotted black line indicates a linear regression ($R^2 = .91$, RMSE = 0.17).




APPENDIX C

Treatment effects on litter mass remaining averaged across all MAT plots over time. Solid shapes denote means; bars denote ± 1 SE, $n = 7$. Solid lines denote best fitted lines.



APPENDIX D

Treatment differences in leaf litter mass remaining a 4.3°C MAT gradient. Solid shapes denote means; bars denote ± 1 SE, $n = 7$. Solid lines denote best-fitted lines and bands represent 95% confidence intervals.



ANOVA and Tukey HSD results for N remaining (% of initial) and C:N at each time collection. Significant differences (95% confidence level) indicated in bolded print.

ANNEX-1500 days (N = initial)				C.N.				
Treatment	Nat	Diff	Pos/Nat	Pos/Nat	Treatment	Nat	Diff	Pos/Nat
Temperature	3	302	22.61	+4.00E-1	Temperature	6	302	4.0
Treatment + temperature	3	302	1.139	+0.00E	Treatment + temperature	6	302	1.0
Treatment	Nat	Diff	Pos/Nat	Pos/Nat	Treatment	Nat	Diff	Pos/Nat
Tally/Ty to Etem	Nat	Diff	Pos/Nat	Pos/Nat	Tally/Ty to Etem	Nat	Diff	Pos/Nat
High control	4	36	+8.00E	+4.00E	High control	7	36	+8.00E
Low control	4	36	+8.00E	+4.00E	Low control	7	36	+8.00E
High level	4	36	+8.00E	+4.00E	High level	7	36	+8.00E
Tally/Ty to Etem	Nat	Diff	Pos/Nat	Pos/Nat	Tally/Ty to Etem	Nat	Diff	Pos/Nat
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	1.00	15.5.15.0	3	36	1.00	
15.8.15.0	2	24	1.00	15.8.15.0	2	24	1.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	
15.10.15.0	2	24	0.00	15.10.15.0	2	24	0.00	
15.5.15.0	3	36	0.00	15.5.15.0	3	36	0.00	
15.8.15.0	2	24	0.00	15.8.15.0	2	24	0.00	