

SOIL NUTRIENT AND MICROBIAL RESPONSE TO PRESCRIBED FIRE IN AN OAK-PINE ECOSYSTEM IN EASTERN KENTUCKY

Beth A. Blankenship and Mary A. Arthur¹

Abstract—On the Cumberland Plateau, periodic fire may be necessary to maintain oak-dominated stands on xeric ridgetops. However, thin organic horizons and nutrient poor soils may limit the beneficial effects of fire by further reducing nutrient pools. The USDA Forest Service has reintroduced fire to oak-pine ridgetops in the Daniel Boone National Forest in the form of single, late winter prescribed burns. Objectives of this study were to document effects of single prescribed fires conducted in March 1995 and March 1996 on soil nutrients and microbial biomass. In 1996, forest floor mass was determined before and after fire. In 1995 and 1996, total and available nitrogen, total carbon, pH, extractable cations, and microbial biomass were measured pre-burn and throughout the year after burning on burned and unburned sites. Thirty-two percent of the litter layer (Oi) was combusted in 1996, while no loss of the Oe and Oa horizons was found. A transitory increase in available nitrogen was detected in burned mineral soils. Increases in pH by 0.2-0.3 units were measured in the burned organic horizons. Lower concentrations of extractable cations were measured in burned soils than in unburned soils. Fire had a positive effect on active bacterial biomass, but no effect on fungal biomass. Our study suggests that single, late winter prescribed fire had minimal effects on belowground resources in these ridgetop ecosystems. Since repeated burning might be necessary to promote oak regeneration, future research must address the effects of repeated burning on soil resources.

INTRODUCTION

Fire has played an integral role in maintaining oak-pine forest communities in the southern Appalachians for as long as 3000 years BP (Delcourt and Delcourt 1997). Aboriginal use of fire was common and settlers of European descent continued fire use in the area; however, as forest use altered, fire regimes changed (Pyne 1982, Pyne and others 1996). This resulted in shifting forest composition, most noticeably as an increasing dominance of red maple (*Acer rubrum* L.) and decreasing regeneration of oak species (*Quercus* sp.) (Arends and McCormick 1987, Lorimer 1993). Ecologists and managers are increasingly promoting the use of fire to address problems of oak decline and flourishing fire-sensitive competitors like red maple.

In the Daniel Boone National Forest (DBNF) in eastern Kentucky, active fire suppression since the 1940s has successfully excluded fire on most oak-pine ridgetops (Martin 1989). Increasing regeneration of white pine (*Pinus strobus* L.) has been documented and ascribed to fire suppression (Wehner 1991), while charcoal and pollen analyses have demonstrated a growing presence of red maple and blackgum (*Nyssa sylvatica* Marshall) over the past 100 years (Delcourt and Delcourt 1997). Because of such changes in forest composition, the USDA Forest Service on the Stanton Ranger District has begun conducting late winter prescribed fires to restore fire to these ridgetop ecosystems (Richardson 1995). We have documented the effectiveness of late winter prescribed fire in reducing competition by fire-sensitive competitors (Arthur and others 1998, Blankenship [In Press]), and studies from other areas have recommended prescribed fire to improve oak regeneration (Barnes and Van Lear 1998, Brose and Van Lear 1998, Nyland and others 1983, Thor and Nichols 1974, Van Lear and Waldrop 1989).

Complicating the use of prescribed fire in eastern Kentucky is the problem of accidental and incendiary fires burning large acreage during the spring and fall seasons each year (Environmental Quality Commission 1992). Consequently, the use of prescribed fire on public lands is perceived by some to send a misleading message about fire to the general public. Because of thin organic soil horizons and low soil nutrient concentrations on ridgetops in the Red River Gorge, fire potentially could, if too hot, contribute to a loss of nutrients already in low supply. The effects of fire on soil microbial biomass may also be important because of the role fungi and bacteria play in mediating nutrient mineralization and availability. An improved understanding of the soil nutrient and biological response, as well as the plant community response, to prescribed fire is necessary to elucidate the ecological differences between the effects of prescribed fire and the potentially hotter and more frequent incendiary fires. This type of information is essential to the public debate regarding the role of prescribed fire in the management of public forestlands.

The objectives of this study were to document effects of a single, late-winter prescribed fire on soil nutrients and microbial biomass as a first step in addressing impacts of different fire regimes in these ridgetop ecosystems. Despite the plethora of studies regarding effects of fire on soil nutrients and microorganisms in forests throughout the United States and elsewhere, few studies have examined effects of burning on soil resources in oak-pine forests in the southern Appalachians (Clinton and others 1996, Vose and others [In Press], Vose and Swank 1993) and none on the Cumberland Plateau in eastern Kentucky.

¹ Graduate Student and Assistant Professor, respectively, Department of Forestry, University of Kentucky, Lexington, KY 40546-0073.

METHODS

Site Description

Permanent study plots are located on three noncontiguous ridgetops in the Stanton Ranger District of the DBNF in the Red River Gorge Geological Area. The Red River Gorge is located in the Cliff Section of the Cumberland Plateau (Braun 1950). The geological substrate is composed of shales and siltstones of the Upper and Lower members of the Breathitt Formation and the Corbin Sandstone of the Lee Formation (Weir and Richards 1974). The ridgetops, Pinch-Em-Tight Ridge, Whittleton Ridge and Klaber Ridge, are located in Powell, Wolfe and Menifee counties of eastern Kentucky. Study plots are located in stands mostly dominated by scarlet oak (*Q. coccinea* Muenchh.) and chestnut oak (*Q. prinus* L.), with some white (*Q. alba* L.) and black oak (*Q. velutina* Lam.). Pines (*P. rigida* Mill., *P. virginiana* Mill., and *P. echinata* Mill.) dominate the most xeric areas. Red maple is abundant in mid- to overstory positions. Sourwood (*Oxydendrum arboreum* (L.) DC), blackgum and dogwood (*Cornus florida* L.) are found in the understory. Eastern white pine seedlings and saplings are plentiful and a few large white pines occur in the overstory. Heath shrubs such as mountain laurel (*Kalmia latifolia* L.) and blueberry (*Vaccinium* spp.) are common throughout.

Soils at Whittleton Ridge are composed of Gilpin silt loam, a moderately deep, well drained soil with a lower subsoil of silty clay loam of the subgroup Typic Hapludult (Hayes 1993). Soils at Klaber Ridge are similar and classified as Latham-Shelock silt loam, moderately deep, moderately well drained, slowly permeable clayey soils and of the subgroups Typic and Aquic Hapludults (Avers and others 1974). Soils at Pinch-Em-Tight are of Alticrest-Ramsey-Rock outcrop complex, moderately deep and shallow, well drained with surface layer and subsoil of sandy loam, and are Typic and Lithic Dystrochrepts (Hayes 1993, SCS 1975). The region has a temperate, humid and continental climate with an evenly distributed average annual precipitation of about 113 centimeters. Mean daily maximum and minimum temperatures in January are 6°C and -6°C, and in July are 30°C and 17°C. Mean annual temperature is 12°C (Hill 1976).

Experimental Design

Each ridge (Klabe, Pinch-Em-Tight and Whittleton) was divided into three areas: no burning, burned in March 1995, and burned in March 1996. This yielded two treatments (burned and unburned) per year of study (1995 and 1996). Study plots were located at random, with the condition that any location within 15 meters of the slope break, a trail, or any human disturbance was rejected. We installed eight 0.04 hectare plots per treatment except at Pinch-Em-Tight Ridge. That site is topographically narrow and traversed with trails and campsites, and we could fit only 6 plots per treatment. The 1995 burn treatment on Pinch-Em-Tight was especially narrow and these plots were laid out systematically in a line, 30 meters apart along the top of the ridge. Centers of all plots were permanently marked, and two trees in each plot were tagged and located (azimuth and distance to plot center) for future reference. While plots were located at random within each treatment

area, treatments were applied nonrandomly to treatment areas because of the necessity of burning a contiguous finger ridge not in contact with privately owned land.

Fire Prescription

USDA Forest Service personnel of the Stanton Ranger District conducted the 1995 and 1996 prescribed fires. The fires were ignited with drip torch by firing line from the highest point and ridges first. From the ridges, strips were pulled downslope into the wind. If backing and flanking fires were not of sufficient intensity (>0.3 m flamelength), point-source and strip firing were used to increase intensity to acceptable levels (Richardson 1995). Wind speeds during the 1995 fires were from 0 to 3.2 km/h, and in 1996 were around 1.6-4.8 km/h. Flame heights for all burns were 0.3-0.9 meters (Richardson 1995, Richardson 1996). In 1996, the prescribed burning planned for Whittleton Ridge could not be conducted due to unsuitable wind direction that would have allowed smoke to blow over an adjacent highway.

Temperatures of prescribed fires were measured using 6 different Tempilac[®] temperature-sensitive paints chosen to represent a temperature range of 93°C to 576°C (Cole and others 1992). A stripe of each paint was applied on aluminum tags and vertically down strips of mica. The aluminum tags were stapled to stakes at heights of 15, 45 and 75 centimeters. The stakes and the mica sheets were placed at 4 meters from each plot center on north, south, east, and west bearings. The mica sheets were inserted into the ground to a depth of 9 centimeters below the litter surface.

Forest Floor Sampling

Forest floor was collected on all sites in February 1996 prior to burning, and in May 1996 following prescribed burning. One 27.5 x 27.5 centimeter sample was collected along three randomly selected azimuths in each plot (3 samples per plot, mean of the three used for analysis). Litter (Oi) was separated from the Oe and Oa horizons in the field and measured separately.

Soil Sampling

Soil samples for nutrient and microbial biomass analyses were taken within a 0.04 hectare plot by sampling along two randomly selected 11 meter radii originating at the center point of each permanently marked plot. Soil samples were stored at 4°C until processed. Nutrients (C, Ca, K, Mg, N, and P) were analyzed on soil samples from a composite of two 4.0 centimeter cores taken at 6.0 meters along each radius. The organic horizon (Oea) was separated from the mineral horizon in the field with mineral soils sampled to a depth of 5 centimeters. Samples for nutrient analyses were taken within 30 days pre-burn, after the first rainfall post-burn (1 week post-burn in 1995 and 2 weeks post-burn 1996), and in September.

For microbial biomass, two soil cores of 2.0 centimeter diameter were taken at 1.5 meter intervals along each radius beginning at 3.0 meters from the center for a total of 24 cores collected at 12 sampling points. Mineral soils were sampled to a depth of 5 centimeters. Organic and

mineral horizons were separated and soil cores composited into a single sample for analysis from each plot. Microbial biomass samples were collected more frequently than nutrient samples because of the expectation of an immediate (1 day) response to fire as well as an attenuation of the effect with time. In 1995, samples for microbial biomass were collected within 30 days pre-burn, the day after the burn, 1 week post-burn (followed first rainfall post-burn), and every 4 weeks until July, then every 6 weeks thereafter through November. Sampling frequency was reduced after July 1995 following examination of results from earlier in 1995. In 1996, after sampling 2 weeks post-burn (following first rainfall post-burn), microbial biomass samples were collected post-burn only every 6 weeks through September 1996.

Available nitrogen (ammonium- and nitrate-nitrogen) was measured on mineral soil samples collected for microbial biomass estimates. In 1995 no pre-burn measurements of available nitrogen were made; both pre- and post-burn measurements of available nitrogen were made in 1996.

Laboratory Analysis

Forest floor samples were oven-dried at 60°C and then weighed. Mehlich III-extractable P, Ca, Mg, and K were determined with the Mehlich III reagent method (Tran and Simard 1993), and analysis by an inductive coupled plasma spectrophotometer. Organic horizon pH was measured in a 1:10 soil:CaCl₂ mixture, and a subset of these measured in a 1:2 soil:water mixture. Statistical analysis was conducted on the organic pH values from the CaCl₂ method. The mineral soil pH was measured in a 1:2 soil:water mixture (Hendershot and Lalande 1993). Total C and N were analyzed using a Leco C/N analyzer. Determination of ammonium- and nitrate-nitrogen in the mineral soil was made using a Technicon II auto-analyzer following KCl extraction (Maynard and Kalra 1993).

Active and total fungal and bacterial biomass were calculated from direct counts of hyphal length (fungi) and number of organisms (bacteria) using direct count epifluorescent microscopy (Ingham and Klein 1984, Ingham and others 1991, Lodge and Ingham 1991). Fluorescein diacetate was used to stain metabolically active fungi and

bacteria; total fungal hyphae (metabolically active and inactive) were counted using direct light source. Direct counts of fungi and bacteria were carried out within one week of sample collection. Based upon the assumption that hyphal tissue density averages 410 mg cm⁻³ and bacterial tissue density averages 330 mg cm⁻³ (Ingham and others 1991), and using estimates of hyphal and bacterial diameters, active fungal and bacterial biomass were calculated from the volume of fungi or bacteria in 1 gram of dry forest floor.

Statistical Analysis

We analyzed the data in two ways: (1) by comparing pre-burn to post-burn within each treatment (burned or unburned) using a t-test, and (2) by comparing the burned treatment to the unburned by analysis of variance. A covariate analysis using pre-burn parameters as the covariate was used for the nutrient data to compare burned to unburned. Statistical analysis of pH was performed on [H⁺] and results were converted to pH values for presentation. On microbial biomass and available nitrogen, Levene's (1960) homogeneity of variance test revealed that often group variances within a treatment were significantly different, so the Welch ANOVA for means was used (Brown and Forsyth 1974, Welch 1951).

Initially, the three ridgetops were selected as replicates for treatment for each year with treatment area as the experimental unit (n=6; 3 burned and 3 not burned), but in 1996 one ridgetop was not burned as planned, so analysis of the 1996 data had only 2 replicates (n=4; 2 burned, 2 not burned). As the study progressed we became aware of inherent site differences. Therefore, statistical analyses were also conducted on a pseudo-replicated design (Hurlburt 1984) using the 8 (or 6) plots per treatment area as the experimental unit with ridge as block and significance determined at p<0.10 (Table 1). If these analyses revealed significant site-by-treatment interactions in addition to significant treatment effects, subsequent analyses were conducted on reduced models comparing the burned treatment to the unburned reference at each ridgetop location (Klaber and Whittleton: n=16; Pinch-Em-Tight: n=12). All statistical

Table 1—Experimental design for study of prescribed burns conducted in March 1995 and March 1996 in the Daniel Boone National Forest, Kentucky

1995			1996		
Ridge/block ^a	Plots	Treatments	Ridge/block	Plots	Treatments
Klaber Ridge	8	Burned/unburned	Klaber Ridge	8	Burned/unburned
Pinch-Em-Tight Ridge	6	Burned/unburned	Pinch-Em-Tight Ridge	6	Burned/unburned
Whittleton Ridge	8	Burned/unburned			
Total: 3	22	2	2	14	2

^a Experimental unit was the ridge (n=3 in 1995, n=2 in 1996) or the plot with ridge as block (n=22 in 1995, n=14 in 1996).

tests were conducted with the JMP® statistical package (SAS Institute 1995).

RESULTS

Pre-Burn Soil Chemistry

Thin, acidic, nutrient-poor organic horizons characterized these ridges prior to burning. Organic horizons (Oea) ranged from <1 to 4-5 centimeters in thickness with a mean pH of 3.70. Total nitrogen averaged 0.66 percent and total carbon 54 percent. Mineral soil pH was slightly higher at 4.24, with total nitrogen averaging 0.14 percent and carbon 4.9 percent.

Fire Temperatures

Surface temperatures ranged from 316°C to 398°C in 1995 and from 204°C to 315°C in 1996. Temperatures within the Oea horizons at 0.5 centimeters below the surface ranged from 204-315°C in 1995 and from 93-203°C in 1996.

Forest Floor Mass

Pre-burn forest floor mass averaged 98.9 grams/meter² for the Oi layer and 184 grams/meter² for Oea layers. The prescribed fires of 1996 combusted an average of 32 percent of the litter layer (Oi) without combusting the underlying Oe and Oa layers. The decrease in Oi because of the fires was highly significant ($p < 0.01$). Forest floor mass measurements were not made in 1995.

Soil Nitrogen

The inorganic nitrogen in the mineral soils primarily took the form of $\text{NH}_4\text{-N}$. Little, if any, $\text{NO}_3\text{-N}$ was measured. In 1995, available nitrogen concentration was significantly higher in the burned treatments than in the unburned treatments 1 day after the burns on Klaber Ridge (12.5 vs. 1.89 $\mu\text{g/g}$ dry wt soil; $p < 0.05$) and Whittleton Ridge (8.43 vs. 4.89 $\mu\text{g/g}$ dry wt soil; $p < 0.10$), but not on Pinch-Em-Tight Ridge (Table 2). One week later there were no significant differences.

In 1996, available nitrogen concentration declined from the pre-burn sampling to the post-burn sampling dates, significantly so in the unburned areas (3.23 to 1.25 $\mu\text{g/g}$ dry wt soil; $p < 0.0001$), but not significantly in the burned treatments based on the pre- vs. post-burn t-tests. Two weeks later, available nitrogen concentrations were significantly lower in the burned treatments (1.28 $\mu\text{g/g}$ dry wt soil) than in the unburned (2.43 $\mu\text{g/g}$ dry wt soil; $p < 0.10$) using covariate analysis. In September 1996, total nitrogen in the mineral horizon was significantly lower in the burned treatments (0.09 pct. vs. 0.15 pct; $p < 0.01$). No consistent effect of fire on total carbon was found.

pH

O horizon pH was significantly higher in the 1995 burned treatments one week post-burn ($p < 0.05$) and in September 1995 ($p < 0.10$), but there were significant site by treatment interactions ($p < 0.05$). Analysis on the reduced models

Table 2—Mean available N ($\text{NH}_4\text{-N} + \text{NO}_3\text{-N}$; $\mu\text{g/g}$ dry wt soil) in mineral horizon of burned and unburned treatments on Klaber, Whittleton, and Pinch-Em-Tight Ridges, Daniel Boone National Forest, Kentucky, following 1995 prescribed fires and prior to and following 1996 prescribed fires

Date	Klaber		Whittleton		Pinch-Em-Tight	
	Burned	Unburned	Burned	Unburned	Burned	Unburned
1 day post-burn 1995	12.5 ^a (3.1) ^d	1.89 (0.66)	8.43 ^a (1.7)	4.89 (0.92)	3.59 (0.99)	4.17 (2.0)
1 wk post-burn 1995	3.72 (0.44)	3.37 (0.22)	4.90 (0.39)	4.86 (0.14)	4.14 (0.57)	3.46 (0.13)
Pre-burn 1996	1.95 (0.38)	2.37 (0.14)			2.79 (0.16)	4.10 (0.43)
1 day post-burn 1996 ^b	1.60 (0.62)	1.24 (0.23)			2.21 (0.11)	1.26 (0.13)
2 wk post-burn 1996	1.49 ^c (0.57)	3.01 (0.48)			1.40 ^c (0.34)	1.64 (0.33)

^a Burned treatment significantly different from unburned treatment of same ridge at $p < 0.10$.

^b Pre-burn vs. post-burn t-tests showed burned treatments not significantly different but unburned treatments significantly different at $p < 0.0001$.

^c Burned treatments significantly different from unburned treatments by covariate analysis at $p < 0.10$.

^d Standard errors (in parentheses) are based upon mean within a treatment.

revealed that burned treatments had significantly higher pH 1 week post-burn at Whittleton Ridge (3.64 vs. 3.18; $p < 0.001$) and in September at Klaber Ridge (3.64 vs. 3.23; $p < 0.01$) (Table 3). The 1995 fires did not affect pH in the upper mineral soils.

The 1996 fires significantly increased organic horizon pH from pre-burn values to 2 weeks post-burn (3.01 to 3.24; $p < 0.10$), even though the pH of the burned treatments did not significantly differ from the unburned treatments 2 weeks post-burn ($p = 0.79$). In September 1996, we found significant differences in pH between burned and unburned treatments in both the organic and mineral horizons with significant site by treatment interactions. The reduced model analyses showed significantly lower pH in Klaber burned organic horizon (3.20 vs. 3.53; $p < 0.01$), while Pinch-Em-Tight burned treatments had significantly higher pH in organic (3.17 vs. 3.08; $p < 0.001$) and mineral horizons (4.29 vs. 3.89; $p < 0.05$).

Cations

Burned treatments contained significantly lower extractable cation concentrations than unburned treatments later in the season. Extractable potassium concentrations were significantly lower in burned organic horizons in September 1995 (362 vs. 439 $\mu\text{g/g}$ dry wt soil; $p < 0.10$) and September 1996 (362 vs. 490 $\mu\text{g/g}$ dry wt soil; $p < 0.10$) (Table 4). In September 1996, magnesium and calcium concentrations were also lower in burned organic horizons (164 vs. 246

$\mu\text{g/g}$ dry wt soil; $p < 0.01$; 1240 vs. 2500 $\mu\text{g/g}$ dry wt soil; $p < 0.01$, respectively).

Extractable cations in the burned upper mineral soil increased or were higher soon after the fires, but by September tended to be significantly lower than in unburned soils. In 1995, significantly higher potassium concentrations were found in the burned treatments one week post-burn (86.7 vs. 84.0 $\mu\text{g/g}$ dry wt soil; $p < 0.05$). In the 1996 burned treatments, significant increases of magnesium (19.4 to 25.2 $\mu\text{g/g}$ dry wt soil; $p < 0.05$) and calcium (100 to 135 $\mu\text{g/g}$ dry wt soil; $p < 0.05$) concentrations from pre-burn to 2 weeks post-burn were found. In September 1995, potassium was significantly lower in burned treatments (62.4 vs. 71.7 $\mu\text{g/g}$ dry wt soil; $p < 0.10$), and in September 1996 lower in the burned treatment of Klaber Ridge (64.6 vs. 89.6 $\mu\text{g/g}$ dry wt soil; $p < 0.10$). In September 1995 and 1996, burned treatments had significantly less magnesium than unburned treatments (1995: 21.6 vs. 25.0 $\mu\text{g/g}$ dry wt soil; $p < 0.05$; 1996: 16.9 vs. 28.7 $\mu\text{g/g}$ dry wt soil; $p < 0.10$). Calcium concentrations were also significantly lower in burned treatments than in unburned in September 1996 (84.5 vs. 169 $\mu\text{g/g}$ dry wt soil; $p < 0.05$).

Microbial Biomass

In the O horizon, microbial biomass was dominated by fungi (32 $\mu\text{g/g}$ dry wt soil) rather than bacteria (14 $\mu\text{g/g}$ dry wt soil), whereas in the mineral horizon, microbial biomass of bacteria (11.3 $\mu\text{g/g}$ dry wt soil) and fungi (11.4 $\mu\text{g/g}$ dry wt soil) were similar.

Table 3—Mean pH values for O horizon of burned and unburned treatments of Klaber, Whittleton, and Pinch-Em-Tight Ridges, Daniel Boone National Forest, Kentucky, prior to and following March 1995 and March 1996 prescribed fires

Date	Site					
	Klaber		Whittleton		Pinch-Em-Tight	
	Burned	Unburned	Burned	Unburned	Burned	Unburned
Pre-burn 1995	3.77	3.69	3.20	3.31	3.09	3.05
1 wk post-burn 1995	3.93	3.58	3.64	3.18 ^a	3.21	3.16
September 1995	3.64	3.23 ^a	3.05	2.93	2.88	2.85
Pre-burn 1996	3.10	3.30			2.94	3.10
2 wk post-burn 1996	3.32 ^b	3.63			3.25 ^b	3.14
September 1996	3.20	3.53			3.17	3.08

^a Covariate analysis of $[\text{H}^+]$ shows burned treatment significantly different from unburned within ridge at $p < 0.01$ using pre-burn $[\text{H}^+]$ as covariate.

^b Analysis of pre-burn and post-burn $[\text{H}^+]$ shows post-burn significantly different from pre-burn at $p < 0.10$.

Table 4—Mean extractable cations ($\mu\text{g/g}$ dry wt soil) in organic and mineral horizons of Burned and Unburned treatments of three ridgetops in Daniel Boone National Forest, Kentucky, prior to and following March 1995 and March 1996 prescribed fires

	O horizon				Mineral soil			
	1995		1996		1995		1996	
	Burned	Unburned	Burned	Unburned	Burned	Unburned	Burned	Unburned
K								
Pre-burn	495 (30) ^a	469 (29)	467 (39)	510 (39)	80.5 (6.7)	85.9 (6.3)	75.5 (5.3)	84.3 (4.9)
Post-burn	513 (35)	558 (34)	473 (47)	433 (47)	86.7 (5.9)	84.0 (5.9)	95.8 (6.3)	102 (6.0)
September	362 (25)	439 (24)	362 (22)	490 (23)	62.4 (5.0)	71.7 (4.7)	61.1 (4.7)	75.3 (4.4)
Mg								
Pre-burn	198 (93)	340 (90)	190 (20)	226 (20)	25.3 (2.7)	29.1 (2.0)	19.4 (3.0)	22.2 (6.0)
Post-burn	228 (12)	216 (12)	209 (22)	204 (22)	24.7 (1.8)	25.6 (1.3)	25.2 (1.5)	29.5 (1.4)
September	165 (13)	190 (12)	164 (13)	246 (14)	21.6 (1.4)	25.0 (1.7)	16.9 (2.5)	28.7 (2.4)
Ca								
Pre-burn	2010 (168)	2040 (163)	1440 (281)	2460 (133)	148 (12)	172 (14)	100 (7.8)	147 (22)
Post-burn	2310 (182)	2060 (173)	1560 (263)	1800 (262)	144 (11)	130 (7.5)	135 (11)	159 (11)
September	1900 (129)	1560 (123)	1240 (180)	2500 (188)	134 (14)	127 (8.0)	84.5 (23)	169 (23)

^a Standard errors (in parentheses) are based upon mean within a treatment.

The prescribed fires of 1995 and 1996 did not consistently affect active fungal biomass, but active bacterial biomass in the organic horizon appeared to be positively affected by both the 1995 and 1996 burns. In the 1995 burned treatments of Whittleton and Pinch-Em-Tight Ridges, active bacterial biomass was higher than the unburned for almost every sampling date post-burn. Burned treatment active bacterial biomass was significantly higher in July 1995 (13 vs. 10 $\mu\text{g/g}$ dry wt soil; $p < 0.05$), in September 1995 (15 vs. 9 $\mu\text{g/g}$ dry wt soil; $p < 0.01$), and in November 1995 (20 vs. 16 $\mu\text{g/g}$ dry wt soil; $p < 0.10$) (Table 5). In May 1995, active bacterial biomass was significantly lower in the burned treatments (17 vs. 20 $\mu\text{g/g}$ dry wt soil; $p < 0.10$). In the 1996 burned treatments, active bacterial biomass was significantly higher 2 weeks post-burn (17 vs. 12 $\mu\text{g/g}$ dry wt soil; $p < 0.05$) and in August 1996 (7.3 vs. 5.2 $\mu\text{g/g}$ dry wt soil; $p < 0.05$). No significant effect on active bacterial biomass was found in the mineral soils.

DISCUSSION

Increases in nutrient availability following fire have been found in some systems and fire regimes and not in others. More intense fires of logging slash in the southern Appalachians have combusted some of the organic layer

without significant loss of C or N from the Oea horizon (Vose and Swank 1993), and have also increased available soil nitrogen (Knoepp and Swank 1993). A stand replacement fire in this region resulted in no detectable change in soil chemistry (Vose and others [In Press]). Other fire studies in other regions show significant increases in soil nutrient concentrations following burning (Austin and Baisinger 1955, Prieto-Fernandez and others 1993).

The oak-pine ridgetops in the Red River Gorge are characterized by a thin litter layer (Oi) atop a very acidic, thin, nutrient-poor organic horizon (Oea). As none of the organic layer (Oea) combusted in these prescribed fires, the fires caused no immediate nutrient loss from the Oea horizon, and with only a fraction of the litter layer consumed, the increase of nutrient concentrations was very small. Also on such a nutrient-poor site, any addition of nutrients may have been quickly taken up by plants or immobilized by microbes (Hungerford and others 1990). While cations in the O horizon did not significantly change soon after burning, significant increases of some cations in the mineral horizon after burning suggest increased mineralization and possibly leaching of nutrients from the

Table 5—Mean active microbial biomass ($\mu\text{g/g}$ dry wt soil) in the O horizon of Burned and Unburned treatments of three ridgetops in Daniel Boone National Forest, Kentucky, following March 1995 and March 1996 prescribed fires

	O horizon			
	1995		1996	
	Burned	Unburned	Burned	Unburned
1 d post-burn	15.5 (0.99) ^a	14.9 (0.97)	10.4 (2.1)	9.36 (1.1)
1 wk post-burn	15.9 (1.7)	13.2 (1.3)	—	—
2 wk post-burn	—	—	17.4 (0.90)	11.8 (1.2)
April	25.2 (2.1)	20.5 (2.0)	—	—
May	16.5 (0.93)	19.8 (1.6)	13.0 (1.0)	11.7 (2.9)
June	5.53 (1.2)	11.7 (1.3)	11.4 (0.59)	10.4 (0.63)
July/August	12.8 (1.1)	10.1 (0.73)	7.33 (0.45)	5.16 (0.90)
September	14.7 (1.7)	8.66 (0.69)	8.27 (0.53)	7.84 (2.2)
November	20.3 (2.0)	15.6 (1.3)	—	—

^a Standard errors (in parentheses) are based upon mean within a treatment.

ash through the thin organic horizon to the mineral soil. We detected a transient increase in available nitrogen, and we know that some increased nutrient availability occurs as evidenced in increased seedling foliar N, P, and K concentrations following burning (Gilbert and others 1998, Reich and others 1990). In another acidic, nutrient-poor ecosystem, the southeastern Coastal Plain pine flatwoods, herb layer vegetation had significant increases in tissue nutrient concentrations for N, P and K following winter fire (Gilliam 1991).

Availability of nutrients for plant uptake depends in part on soil pH, which often changes after burning as oxides in ash react with hydrogen ions to raise pH (Agee 1993, Ahlgren and Ahlgren 1960). We found significantly higher pH in burned soils both years. Higher pH may also have favored bacterial populations (Ahlgren 1974) which increased following burning. While many studies show decreased microbial populations after fire (Ahlgren 1974, Borchers and Perry 1990, Pietikäinen and Fritze 1993), other studies have found bacterial populations to increase following burning (Ahlgren and Ahlgren 1965, Jurgensen and others 1981). Lower concentrations of available cations measured

late in the season may reflect increased immobilization by increasing bacterial biomass (Fritze and others 1993).

Fire is being reintroduced to these ridgetops in an attempt to maintain a community diversity that has been decreasing since fire suppression. Successful regeneration of oak species is integral to this diversity. Our data (Arthur and others 1998) as well as other studies (Barnes and Van Lear 1998, Thor and Nichols 1974, Van Lear and Waldrop 1989) suggest that multiple fires will be necessary to realize this objective. This study suggests that low intensity single fires where little, if any, organic matter is lost have little effect on soil nutrient status. Future research should examine whether repeated fire on these sites has similar effects on soil nutrients and microbial biomass or whether impacts of repeated burning on soil resources further degrade already nutrient-poor sites.

ACKNOWLEDGMENTS

For assistance in the field and lab, we thank Mark Schuster, Laurie Taylor, Kim Feeman, Peter Hadjiev, Jamie Winders, Carol Miller, and Melinda Hamilton. We also thank Frank Gilliam, Wayne Swank, and David Van Lear for their time and comments in reviewing the manuscript. The research was made possible through collaboration with USDA Forest Service Stanton Ranger District personnel, especially District Ranger Donnie Richardson, Bill Luhn, and Rita Wehner. This work was partially funded by the E.O. Robinson Forest Trust Fund and a Challenge Cost-share Agreement with the Daniel Boone National Forest, Winchester, KY. This study (#98-09-141) is connected with a project of the Kentucky Agricultural Experiment Station and is published with the approval of the Director.

REFERENCES

- Agee, J.K. 1993. Fire ecology of Pacific Northwest forests. Washington, DC: Island Press. 493 p.
- Ahlgren, I.F. 1974. The effect of fire on soil organisms. In: Kozlowski, T.T.; Ahlgren, C.E., eds. Fire and ecosystems. New York: Academic Press: 47-71.
- Ahlgren, I.F.; Ahlgren, C.E. 1960. Ecological effects of forest fires. Botanical Review. 26: 483-533.
- Ahlgren, I.F.; Ahlgren C.E. 1965. Effects of prescribed burning on soil microorganisms in a Minnesota jack pine forest. Ecology. 46: 304-310.
- Arends, E.; McCormick, J.F. 1987. Replacement of oak-chestnut forests in the Great Smoky Mountains. In: Hay, R.L.; Woods, F.W.; DeSelm, H., eds. Proceedings of the central hardwood forest conference VI; 1987 February 24-26; Knoxville, TN. Knoxville: University of Tennessee: 305-315.
- Arthur, M.A.; Paratley, R.D.; Blankenship, B.A. 1998. Single and repeated fires affect survival and regeneration of woody and herbaceous species in an oak-pine forest. Journal of the Torrey Botanical Society. 125: 225-236.
- Austin, R.C.; Baisinger, D.H. 1955. Some effects of burning on forest soils of western Oregon and Washington. Journal of Forestry. 53: 275-80.
- Avers, P.E.; Austin, J.S.; Long, J.K. [and others]. 1974. Soil survey of Menifee and Rowan Counties and northwestern Morgan County, Kentucky. Washington, DC: U.S. Department of Agriculture, Soil Conservation Service. 88 p.

- Barnes, T.A.; Van Lear, D.H.** 1998. Prescribed fire effects on advanced regeneration in mixed hardwood stands. *Southern Journal of Applied Forestry*. 22: 138-142.
- Blankenship, B.A.; Arthur, M.A.** [In press]. Prescribed fire affects eastern white pine recruitment and survival on eastern Kentucky ridgetops. *Southern Journal of Applied Forestry*.
- Borchers, J.G.; Perry, D.A.** 1990. Effects of prescribed fire on soil organisms. In: Walstad, J.D.; Radosovich, S.R.; Sandberg, D.V., eds. *Natural and prescribed fire in the Pacific Northwest*. Corvallis, OR: Oregon State University: 143-157.
- Braun, E.L.** 1950. *Deciduous forests of Eastern North America*. New York: Hafner Press. 536 p.
- Brose, P.H.; Van Lear, D.H.** 1998. Responses of hardwood advance regeneration to seasonal prescribed fires in oak-dominated shelterwood stands. *Canadian Journal of Forest Research*. 28: 331-339.
- Brown, M.B.; Forsythe, A.B.** 1974. Robust tests for the equality of variances. *Journal of the American Statistical Association*. 69: 364-367.
- Clinton, B.D.; Vose, J.M.; Swank, W.T.** 1996. Shifts in aboveground and forest floor carbon and nitrogen pools after felling and burning in the Southern Appalachians. *Forest Science*. 42: 431-441.
- Cole, K.L.; Klick, K.F.; Pavlovic, N.B.** 1992. Fire temperature monitoring during experimental burns at Indiana Dunes National Lakeshore. *Natural Areas Journal*. 12: 177-183.
- Delcourt, P.A.; Delcourt, H.R.** 1997. Report of paleoecological investigations. Cliff Palace Pond, Jackson County, Kentucky, in the Daniel Boone National Forest. Forest Service Contract 43-531A-6-0389. 61 p.
- Environmental Quality Commission.** 1992. *State of Kentucky's environment: a report of progress and problems*. Frankfort, KY: Commonwealth of Kentucky.
- Fritze, H.; Pennanen, T.; Pietikäinen, J.** 1992. Recovery of soil microbial biomass and activity from prescribed burning. *Canadian Journal of Forest Research*. 23: 1286-1290.
- Gilbert, N.L.; Arthur, M.A.; Johnson, S.L. [and others].** 1998. Prescribed fire effects on foliar nutrients, photosynthesis, and growth of ridgetop maple and oak seedlings [Poster presentation]. In: Annual meetings of the Ecological Society of America; 1998 August 1-6; Baltimore, MD.
- Gilliam, F.S.** 1991. The significance of fire in an oligotrophic forest ecosystem. In: Nodvin, S.C.; Waldrop, T.A., eds. *Fire and the environment: ecological and cultural perspectives: Proceedings of an international symposium*; 1990 March 20-24; Knoxville, TN. Gen. Tech. Rep. SE-69. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 113-122.
- Hayes, R.A.** 1993. Soil survey of Powell and Wolfe Counties, Kentucky. Washington, DC: U.S. Department of Agriculture, Soil Conservation Service. 173 p.
- Hendershot, W.H.; Lalonde, H.** 1993. Soil reaction and exchangeable acidity. In: Carter, M.R., ed. *Soil sampling and methods of analysis*. Ann Arbor: Lewis Publishers: 141-145.
- Hill, J.D.** 1976. *Climate of Kentucky*. Progress Rep. 221. Lexington, KY: University of Kentucky, College of Agriculture, Agricultural Experiment, Department of Agronomy. 88 p.
- Hungerford, R.D.; Harrington, M.G.; Frandsen, W.H. [and others].** 1990. Influence of fire on factors that affect site productivity. In: Harvey, A.E.; Neuenschwander, L.F., comps. *Proceedings, management and productivity of western-montane forest soils*; 1990 April 10-12; Boise, ID. Gen. Tech. Rep. INT-280. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 32-50.
- Hurlburt, S.H.** 1984. Pseudoreplication and the design of ecological field experiments. *Ecological Monographs*. 54: 187-211.
- Ingham, E.R.; Griffiths, R.P.; Cromack, K.; Entry, J.A.** 1991. Comparison of direct vs. fumigation incubation microbial biomass estimates from ectomycorrhizal mat and non-mat soils. *Soil Biology and Biochemistry*. 23: 465-471.
- Ingham, E.R.; Klein, D.A.** 1984. Soil fungi: Relationships between hyphal activity and staining with fluorescein diacetate. *Soil Biology and Biochemistry*. 16: 273-278.
- Jurgensen, M.F.; Harvey, A.E.; Larsen, M.J.** 1981. Effects of prescribed fire on soil nitrogen levels in a cutover Douglas-fir/western larch forest. Res. Pap. INT-275. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Research Station. 6 p.
- Knoepp, J.D.; Swank, W.T.** 1993. Site preparation burning to improve Southern Appalachian pine-hardwood stands: nitrogen responses in soil, soil water, and streams. *Canadian Journal of Forest Research*. 23: 2263-2270.
- Levene, H.** 1960. Robust tests for the equality of variances. In: Olkin, I., ed. *Contributions to probability and statistics*. Stanford: Stanford University Press. 517 p.
- Lodge, D.J.; Ingham, E.R.** 1991. A comparison of agar film techniques for estimating fungal biovolumes in litter soil. *Agriculture, Ecosystems and Environment*. 5: 131-144.
- Lorimer, C.G.** 1993. Causes of the oak regeneration problem. In: Loftis, D.; McGee, C.E., eds. *Oak regeneration: serious problems, practical recommendations, symposium proceedings*; 1992 September 8-10; Knoxville, TN. Gen. Tech. Rep. SE-84. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 14-39.
- Martin, W.H.** 1989. *The role and history of fire in the Daniel Boone National Forest*. Winchester, KY: U.S. Department of Agriculture, Forest Service. 131 p.
- Maynard, D.G.; Kalra, Y.P.** 1993. Nitrate and exchangeable ammonium nitrogen. In: Carter, M.R., ed. *Soil sampling and methods of analysis*. Ann Arbor: Lewis Publishers: 25-38.
- Nyland, R.D.; Abrahamson, L.P.; Adams, K.B.** 1983. Use of prescribed fire for regenerating white and red oak in New York. In: *America's hardwood forests - opportunities unlimited*. Proceedings of the 1992 convention of the Society of American Foresters; 1982 September 19-22; Cincinnati, OH. Bethesda, MD: Society of American Foresters: 163-167.
- Pietikäinen, J.; Fritze, H.** 1993. Microbial biomass and activity in the humus layer following burning: short-term effects of two different fires. *Canadian Journal of Forest Research*. 23: 1275-1285.
- Prieto-Fernandez, A.; Villar, M.C.; Carballas, M.; Carballas, T.** 1993. Short-term effects of a wildfire on the nitrogen status and its mineralization kinetics in an Atlantic forest soil. *Soil Biology and Biochemistry*. 25: 1657-1664.
- Pyne, S.J.** 1982. *Fire in America: a cultural history of wildland and rural fire*. Princeton: Princeton University Press. 654 p.

- Pyne, S.J.; Andrews, P.L.; Laven, R.D.** 1996. Introduction to wildland fire. 2d ed. New York: John Wiley. 769 p.
- Reich, P.B.; Abrams, M.D.; Ellsworth, D.S. [and others].** 1990. Fire affects ecophysiology and community dynamics of central Wisconsin oak forest regeneration. *Ecology*. 71: 2179-2190.
- Richardson, D.M.** 1995. Prescribed fire plan for Klaber Ridge, Pinch-Em-Tight Ridge, and Whittleton Ridge, Daniel Boone National Forest, Kentucky. R8-FS-5100-6. U.S. Department of Agriculture, Forest Service.
- Richardson, D.M.** 1996. Prescribed fire plan for Klaber Ridge and Pinch-Em-Tight Ridge, Daniel Boone National Forest, Kentucky. R8-FS-5100-6. U.S. Department of Agriculture, Forest Service.
- SAS Institute Inc.** 1995. JMP® User's guide. Cary, NC: SAS Institute, Inc. 593 p.
- Soil Conservation Service.** 1975. Soil taxonomy: a basic system of soil classification for making and interpreting soil surveys. Agric. Handb. 436. Washington, DC: U.S. Department of Agriculture, Soil Conservation Service. 754 p.
- Thor, E.; Nichols, G.M.** 1974. Some effects of fire on litter, soil and hardwood regeneration. In: Proceedings of Tall Timbers Fire Ecology Conference 13; 1973 March 22-23; Tallahassee: Tall Timbers Research Station: 317-329.
- Tran, T.S.; Simard, R.R.** 1993. Mehlich III extractable elements. In: Carter, M.R., ed. Soil sampling and methods of analysis. Ann Arbor: Lewis Publishers: 43-39.
- Van Lear, D.H.; Waldrop, T.A.** 1989. History, uses, and effects of fire in the Appalachians. Gen. Tech. Rep. SE-54. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 20 p.
- Vose, J.M.; Swank, W.T.** 1993. Site preparation burning to improve Southern Appalachian pine-hardwood stands: aboveground biomass, forest floor mass, and nitrogen and carbon pools. *Canadian Journal of Forest Research*. 23: 2255-2262.
- Vose, J.M.; Swank, W.T.; Clinton, B.D. [and others].** [In press]. Using stand replacement fires to restore Southern Appalachian pine-hardwood ecosystems: effects on mass, carbon, and nutrient pools. *Forest Ecology and Management*.
- Wehner, R.R.** 1991. The status of white pine (*Pinus strobus*) in the Clifty Wilderness, Kentucky. Lexington, KY: University of Kentucky. 67 p. M.S. thesis.
- Weir, G.W.; Richards, P.W.** 1974. Geologic map of the Pomeroyton quadrangle, east-central Kentucky. Reston, VA: U.S. Geological Survey.
- Welch, B.L.** 1951. On the comparison of several mean values: an alternative approach. *Biometrika*. 38: 330-336.