

ARTICLE

Methods, Tools, and Technologies

Integrating forest structural diversity measurement into ecological research

Jeff W. Atkins^{1,2}  | Parth Bhatt³  | Luis Carrasco⁴  | Emily Francis^{5,6}  |
 James E. Garabedian¹  | Christopher R. Hakkenberg⁷  | Brady S. Hardiman^{8,9}  |
 Jinha Jung¹⁰  | Anil Koirala¹¹  | Elizabeth A. LaRue¹²  | Sungchan Oh⁸  |
 Gang Shao⁹  | Guofan Shao⁸  | H. H. Shugart¹³  | Anna Spiers¹⁴  |
 Atticus E. L. Stovall^{15,16}  | Thilina D. Surasinghe¹⁷  | Xiaonan Tai¹⁸  |
 Lu Zhai¹⁹  | Tao Zhang²⁰  | Keith Krause²¹ 

¹USDA Forest Service Southern Research Station, New Ellenton, South Carolina, USA²Department of Biology, Virginia Commonwealth University, Richmond, Virginia, USA³College of Forest Resources and Environmental Science, Michigan Technological University, Houghton, Michigan, USA⁴National Institute for Mathematical and Biological Sciences, University of Tennessee, Knoxville, Tennessee, USA⁵Department of Biology, University of New Mexico, Albuquerque, New Mexico, USA⁶Department of Integrative Biology, University of Texas at Austin, Austin, Texas, USA⁷School of Informatics, Computing, and Cyber Systems, Northern Arizona University, Flagstaff, Arizona, USA⁸Department of Forestry and Natural Resources, Purdue University, West Lafayette, Indiana, USA⁹Libraries and School of Information Studies, Purdue University, West Lafayette, Indiana, USA¹⁰Lyles School of Civil Engineering, Purdue University, West Lafayette, Indiana, USA¹¹Warnell School of Forestry and Natural Resources, University of Georgia, Athens, Georgia, USA¹²Department of Biological Sciences, University of Texas: El Paso, El Paso, Texas, USA¹³Department of Environmental Sciences, University of Virginia, Charlottesville, Virginia, USA¹⁴Department of Ecology & Evolutionary Biology, University of Colorado Boulder, Boulder, Colorado, USA¹⁵Biospheric Sciences Laboratory, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA¹⁶Department of Geographical Science, University of Maryland, College Park, Maryland, USA¹⁷Bridgewater State University, Bridgewater, Massachusetts, USA¹⁸Department of Biological Sciences, New Jersey Institute of Technology, Newark, New Jersey, USA¹⁹Department of Natural Resource Ecology & Management, Oklahoma State University, Stillwater, Oklahoma, USA²⁰Department of Forest Resources, University of Minnesota, St. Paul, Minnesota, USA²¹Battelle, National Ecological Observatory Network (NEON), Boulder, Colorado, USA

Correspondence

Jeff W. Atkins

Email: jeffrey.atkins@usda.gov

Abstract

The measurement of forest structure has evolved steadily due to advances in technology, methodology, and theory. Such advances have greatly increased

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Ecosphere* published by Wiley Periodicals LLC on behalf of The Ecological Society of America. This article has been contributed to by U.S. Government employees and their work is in the public domain in the USA.

Funding information

National Science Foundation (NSF),
Grant/Award Number: 1924942

Handling Editor: Charles D. Canham

our capacity to describe key forest structural elements and resulted in a range of measurement approaches from traditional analog tools such as measurement tapes to highly derived and computationally intensive methods such as advanced remote sensing tools (e.g., lidar, radar). This assortment of measurement approaches results in structural metrics unique to each method, with the caveat that metrics may be biased or constrained by the measurement approach taken. While forest structural diversity (FSD) metrics foster novel research opportunities, understanding how they are measured or derived, limitations of the measurement approach taken, as well as their biological interpretation is crucial for proper application. We review the measurement of forest structure and structural diversity—an umbrella term that includes quantification of the distribution of functional and biotic components of forests. We consider how and where these approaches can be used, the role of technology in measuring structure, how measurement impacts extend beyond research, and current limitations and potential opportunities for future research.

KEYWORDS

forest ecology, forest structure, forestry, landscape ecology, lidar, measurement, remote sensing, spatial sampling, structural diversity

INTRODUCTION

Our ability to measure forest structure has greatly progressed in recent years due to technological and theoretical breakthroughs, primarily in the field of remote sensing, which have provided the long-awaited capability to precisely measure multiple dimensions of forest structure across resolutions and over vast extents. The broad availability of a suite of accessible tools and approaches to measure forest structural diversity (FSD) has fostered with novel opportunities to integrate FSD into research. Previously, the use of many remote sensing tools and associated data products have been restricted to only the most resource-rich or tech-savvy individuals, limiting the impact of such advancements. However, we are now on the precipice of broad inclusion of tools such as terrestrial lidar scanners (TLS), unoccupied aerial vehicles/systems (UAV/UAS), and radar for ecological research. While this portends the possibility of exciting, transformative research, understanding the basics of how forest structure and structural diversity can be quantified via remote sensing can guide research applications. Here we present a review and introduction to how these new technologies can be integrated into ecological research.

Forest structure and forest structural diversity

Broadly, forest structure refers to the distribution of individual trees or biomass in space within a forest

(Goff & Zedler, 1968; Zenner et al., 2012); FSD specifically refers to the patterns of these distributions (LaRue et al., 2023; McRoberts et al., 2008), or more precisely, the multidimensional characterization of measurable forest structural attributes including stand structure, height, cover, volume, heterogeneity, arrangement, and the distribution of functional and biotic components (Hakkenberg & Goetz, 2021; LaRue et al., 2023). Accordingly, FSD measurement has broad implications for science, policy, and management (Beland et al., 2019; Eitel et al., 2016; Kruys et al., 2013; Meng et al., 2016). Accurate and precise quantification of FSD is necessary for understanding environmental drivers of community organization and composition (Hakkenberg et al., 2018; Heidrich et al., 2020) including species presence, abundance, and distribution (Borcard et al., 1992; Vihervaara et al., 2015); sustainable forest management, wildlife conservation, and restoration initiatives (Almeida, Stark, et al., 2019; Garabedian et al., 2017; McNeil et al., 2023; Parker & Russ, 2004; Valbuena et al., 2020; Wales et al., 2020); detecting forest disturbance (Atkins et al., 2019; Jucker, 2021; Zhai et al., 2022) and system recovery (Almeida et al., 2021; Meng et al., 2018); characterizing habitat (Hernando et al., 2012) and microclimates (Ehbrecht et al., 2017; Zellweger et al., 2020) and mapping land cover change (Erb et al., 2018).

Outside of the field of applied forestry, the study of forest structure has historically focused on tree species composition and abundance, with limited emphasis on physical structure—despite the recognized potential importance of structure to ecological pattern and process

(Long & Shaw, 2009; Shugart et al., 2010; Wiens, 1974). The key contributors to this oversight have been the logistical, theoretical, financial, and technological constraints on measuring the physical structure. Advances in technology and theory have expanded our ability to measure FSD; however, the application of FSD measurements in ecological research relies on understanding the range of available FSD metrics, how they are derived, their biological interpretation, and their potential applications and limitations. To be broadly useful, any measures of FSD should be temporally sensitive and complementary to compositional measures (Proença et al., 2017) and serve as reliable proxies for predicting stand- to global-scale biodiversity (Cavender-Bares et al., 2020; Jetz et al., 2019) and functional diversity (Asner et al., 2017). Here, we review: (1) the range of available FSD metrics and the structural attributes they assess; (2) how FSD measurements are derived; (3) considerations for the application of FSD in research and applications; (4) implications of the use of FSD for science, policy, and management; and (5) current limitations and potential future directions.

STRUCTURAL ATTRIBUTES MEASURED BY FSD METRICS

There are several structural attributes of forests that may be quantified using FSD metrics, which we have grouped as: *stand structure* (e.g., tree size and stem diameter distributions, trees per acre, canopy volume); *height, cover, and openness* (e.g., gap fraction, canopy cover); *heterogeneity* (also vertical stratification or canopy structural complexity); *vegetation area and density* (e.g., leaf area index); and *structural heterogeneity of traits or taxa* (e.g., diversity/distribution of foliar traits; Fahey et al., 2019; Matasci et al., 2018; Sheldon et al., 2006) (Figure 1; full descriptions in Table 1).

FSD measurements most often are distilled down to singular metrics describing specific structural attributes of forests. Metrics may range in their level of complexity from the straightforward such as stand density—the number of trees for a given area (e.g., trees per acre or trees per hectare)—to highly derived metrics such as foliage height diversity—the distribution of canopy layers within the vertical plane. Distillation is a key advantage of FSD metrics, facilitating extensive spatially explicit datasets to be condensed or aggregated down to a single, information-rich metric or metrics that can then be used in modeling or analysis efforts. The range of complexity of FSD metrics is based on the dimensionality of the metric, data needed for calculation, and the level of processing or derivation required (Table 1). The choice of which FSD metric(s) to use for a given application should

be informed by: (1) the structural attributes under investigation; (2) spatial and temporal resolution and extent based on the scale of the system of study; and (3) the ability to obtain, access, process, and analyze the necessary data.

FSD measurement

The structural attributes of forests quantified by FSD metrics can be measured via multiple approaches, platforms, and methods. While a complete, detailed understanding of each method of measurement and its associated nuances may be unnecessary for many practicing ecologists and researchers, a cursory understanding of how FSD measurements are made is beneficial. FSD measurement is influenced by several technological or theoretical factors—measurement precision, instrument resolution and extent in both time and space, speed and ease of measurement acquisition, logistics of data acquisition and storage, ease of data usage, access to software and tools to work with data, and the existence of pipelines for processing and analyzing data. To discuss these points in further detail, we have chosen to group our discussion by traditional field-based methods and by remote-sensing-based approaches.

Traditional field-based approaches

Forestry has always been a field of measurement. The need for sustainable timber production necessitated the rise of forest management practices underpinned by standardized data on the abundance, volume, and distribution of trees to meet this need (Avery & Burkhart, 2015). This led to the rise of the subdiscipline of forest mensuration, which is focused solely on the measurement of forests and forest structure. Technology evolved over time to meet these needs, giving us tools such as dbh tapes, relascopes, clinometers, calipers, the Biltmore stick, and methodologies such as allometries and yield tables. Traditional methods capture several, but not all, forest structural attributes listed above (i.e., stand structure, height, vegetation area and density, and volume and surface area).

Traditional methods are often individual-, point-, or plot-based (e.g., forest inventory plot, common stand exam) and require extensive time and financial investment to acquire new data. This results in workforce capacity—either in the form of time or money or both—as the primary limiting factor on data acquisition via traditional methods. These logistical constraints result in fewer samples, coarser grains, and/or smaller spatial extents than many remote sensing approaches (Figure 2).




FIGURE 1 Conceptual diagrams of structural attributes (Table 1). (A) Three tree species distinguished by color but could also represent some trait (e.g., nitrogen content); (B) internal and external structural heterogeneity, based on the arrangement of all vegetated elements in 3D space; (C) three measures of canopy height including mean outer canopy height (MOCH) or local outer canopy height (LOCH), mean canopy height, and maximum canopy height; (D) shown in overhead 2D, the darker colors indicate greater leaf area/density; (E) canopy cover shown as absence or presence; (F) stem distribution map showing relative dbh based on size of circle as well as a dbh distribution curve at top right.

However traditional measurement does provide granular information at the level of the individual that is not possible with many remote sensors, creating a trade-off in extent versus resolution and precision versus accuracy. dbh, for example, is mostly measured using calibrated measurement tapes (dbh tapes) or calipers, each with a typical precision of 0.1 cm and associated measurement error of ± 0.2 – 0.5 cm (Luoma et al., 2017). Tree height can be measured using a standard transect tape with either a clinometer or hypsometer coupled with basic trigonometry, with precision ranging from 0.1 to 1 m and associated measurement error up to $\pm 10\%$ (Andersen et al., 2006; Luoma et al., 2017). Other FSD attributes such as basal area or volume can be measured using basal area prisms or tools such as a Biltmore

stick (Hovind & Rieck, 1961), tools calibrated to give direct estimates of these variables as opposed to estimates scaled up from dbh tapes using allometries (Jenkins et al., 2003). First-order measurements taken via traditional measurements (e.g., diameter, height, distributions) can be used to calculate advanced second-order metrics invaluable in both forestry and ecology (e.g., dissimilarity indexes, site index).

One of the key major advantages of traditional FSD measurement approaches is the lower investment costs in both time and money in data processing and analysis. FSD data acquired via traditional methods (e.g., dbh, basal area, tree height) tend to be intuitive and readily analyzable via standard database and spreadsheet software environments. However, this ease of use does not

TABLE 1 Categories of structural attributes with accompanying information, including detailed descriptions of each attribute, dimensions of the associated data, and example metrics and approaches toward quantification.

Structural attribute	Description	Data dimensions	Example metrics	Measurement approaches
Stand structure	Distribution of tree size or tree stem diameters, including basal area and tree volume	1D, 2D, 3D	Variance in diameter distributions, Gini coefficient, Weibull profile, power function scaling exponent, stand density, stand volume, board feet, fractal geometry, and ecosystem volume	Standard forestry methods include dbh tapes, Biltmore sticks, relascopes, angle gauges, etc. Remote sensing methods include terrestrial laser scanning or inference from aerial lidar or orthoimagery via statistical means (e.g., crown segmentation, imputation). Additionally volume from allometries and yield tables (e.g., stand volume, board feet). Require a low to high level of processing complexity.
Height	Tree or vegetation height, as well as height profiles, from ground to emergent canopy and aggregated height (e.g., mean, median, or local outer height)	1D, 2D, 3D	Lorey's height, mean outer canopy height, vegetation height percentile distribution, canopy relief ratio, and vertical distribution index	Traditional methods include hypsometers, relascopes, and clinometers. Remote sensing approaches include structure-from-motion, lidar, radar, and microwave. Height metrics vary from low to medium complexity processing levels.
Cover and openness	Relative proportion of canopy versus exposed forest floor	2D, 3D	Gap fraction and canopy cover	Hemispherical imagery, orthoimagery, and lidar. Cover and openness vary from low to medium processing.
Heterogeneity/vertical stratification/canopy structural complexity	Variation in the 3D arrangement of canopy elements	3D	Rumple, rugosity, foliage height diversity (FHD), and structural complexity index (SCI)	Currently only estimated from terrestrial, aerial lidar, and spaceborne lidar and tend to require a high level of processing complexity.
Vegetation area and density	The no. and density of vegetation surface layers	2D, 3D	Vegetation area index (VAI), effective no. layers (ENL), leaf area density (LAD), porosity, leaf area index (LAI), wood area index, and plant area index	Orthoimagery, hemispherical imagery, lidar, radar—though ENL and porosity are only from aerial and terrestrial lidar currently. These metrics require a medium to high level of processing complexity.
Structural heterogeneity of traits or biotic identity	Variation in the 3D arrangement of functional or biotic elements	2D, 3D	Height heterogeneity of different species, foliar or spectral diversity	Typically can only be inferred statistically from other stand structural data or via integration of structural data with trait or other data and require a high level of processing.

believe their utility. Detailed information acquired from forest inventory plots or common stand exams has been and will continue to be vital to research. Additionally, since traditional methods are well established within the research and management communities, they tend to be available for a broad range of systems and forests and are well established in the literature.

FSD measurement via remote sensing

While transformative to how we study forests and the Earth system, FSD measurement via remote sensing also carries specific advantages, considerations, and limitations regarding its use. The remote sensing platforms available for measuring FSD range from small handheld




FIGURE 2 Sensors and tools used to make forest structural diversity measurements vary in the spatial extent or resolution at which they can provide continuous structural information. Spaceborne instruments may provide near-global coverage but lack fine-scale spatial resolution; terrestrial-based instrumentation or traditional forestry tools might provide fine-scale, highly precise information at the local scale but require far too much time and effort for broader coverage. Unoccupied aerial vehicles/systems (UAV/UAS) and airborne platforms provide information at moderate resolution and extent.

devices (e.g., personal mobile device) to spaceborne platforms (Figure 2). The spatial extent and grain of a given sensor inherently influences the types of structural features it can quantify and determines the accuracy and precision of which it is capable (Atkins et al., 2023; Saatchi et al., 2011).

Broadly there are two types of remote sensing, passive and active. Passive remote sensing relies on recording energy (i.e., radiation) reflected or emitted from an object illuminated by an external energy source, most often the Sun. Recorded energy is measured as reflectance—the ratio of the amount of energy emitted from a target to the amount of energy striking that target. Passive remote sensing is an important tool, but it is not capable of directly estimating forest structure because light is scattered within a forest canopy, which affects the measurement of spectral response by confounding the relationship between the signal and any leaf or canopy-level scattering properties (Knyazikhin et al., 2013). Reflectance data are used to predict chemical (e.g., leaf nitrogen content) and functional (e.g., phenology, evapotranspiration) ecosystem properties and are used to calculate spectral indices—ratios of one or more spectral bands

to another—which are important for everything from estimating plant stress to tracking deforestation. There are more than a hundred ecologically relevant spectral indices (Kriegler et al., 1969; Xue & Su, 2017; Zeng et al., 2022) with broad applications. While many structural attributes of forests cannot be directly estimated from reflectance data without unconstrained error (Fisher et al., 2018; LaRue et al., 2019; Zeng et al., 2022), passive remote sensing is vital in ecological research to determine forest cover classifications based on vegetation composition and community dominance (Bhatt et al., 2022; Wulder et al., 2004); and to estimate crown closure and crown gaps (Frolking et al., 2009; Wang et al., 2004); aboveground forest biomass (Yan et al., 2015); and stand structure (e.g., leaf area, stem density, height, volume, and basal area), stand maturity (Cohen & Goward, 2004), and successional stage (Hall et al., 1991; Song & Woodcock, 2002). Spectral variation can even be used as a proxy for species richness (see spectral variation hypothesis; Rocchini et al., 2004; Schmidtlein & Fassnacht, 2017) and function (Schweiger et al., 2018).

Active remote sensing surmounts many of the limitations of passive systems by recording how emitted energy interacts with the environment without the confounding influence of an external radiation source, providing direct estimation of FSD (Coops et al., 2021; Lefsky et al., 2002). Active remote sensing such as lidar sensors measure the time required for an emitted light pulse to intercept an object and return to the instrument. Given the speed of light is constant and the known sensor position and beam trajectory, relative distances from objects to the sensor can be calculated directly. In forest applications, lidar produces a three-dimensional characterization of the forest as either discrete points (e.g., point clouds) in the case of time-of-flight measurement, or as a continuous waveform in the case of waveform lidar. While several sensor-specific attributes constrain or bias measurement (e.g., laser beam spot size, beam divergence, pulse density, system range, and the speed with which the sensor motor spins), lidar-based estimates of forest structure have been found to excel in terms of precision, accuracy, consistency, and spatial extent (Coops et al., 2021), making lidar the leading tool in FSD measurement for the foreseeable future.

Radar and microwave remote sensing rely on sensing objects roughly larger than the wavelength of the emitted signal (Henderson & Lewis, 1998). As a result, many radar bands can pass through cloud cover and long wavelength bands (e.g., P band), can even pass through fine vegetation elements (Imhoff, 1995). Radar backscatter largely reflects water distributed in a mixture of soil and vegetation properties, thereby capturing forest structure. Vegetation parameters, including some structural properties (e.g., vegetation height, shape, and orientation of

plant elements), can be retrieved from radar data using a model inversion approach that employs a specific discrete model based on radar polarimetry, but is most applicable in lower density forests before reaching the “saturation” point (Joshi et al., 2017), beyond which return energy does not vary with increasing biomass. For stands with higher density, the performance of radar is often poor because the sensitivity of backscatter to forest structural properties decreases dramatically (Saatchi et al., 2011). Microwave bands (C and X bands) can penetrate rather deeply into the forest canopy, potentially providing a more detailed characterization of internal canopy structure than air- or spaceborne lidar, avoiding signal interference from occlusion (Disney et al., 2006). Microwave bands are also sensitive to changes in foliage and leaf area (Du et al., 2019), and to moisture content, indicating the ability to provide robust estimates of leaf area (Tanase et al., 2019).

Close-range or proximal instruments such as TLS (Calders et al., 2020), mobile phones (Tatsumi et al., 2022), photogrammetry (Mokroš et al., 2018) and tripod-based instruments are capable of mm- to cm-scale spatial resolution and precision, making it possible to collect information on individual leaves/needles, stems, branches, boughs, and trunks (Moorthy et al., 2019). TLS has been found effective in estimating biomass (Calders et al., 2015; Wilkes et al., 2018), forest function (Atkins et al., 2018; Fotis, Morin, et al., 2018; Fotis, Murphy, et al., 2018), species habitat (Blakey et al., 2017), and canopy structure (Ehbrecht et al., 2016; Fahey et al., 2019; Walter et al., 2021). 3D point clouds can be processed using algorithms to identify points that belong to the same individual tree and geometric shapes can be fitted to the points to model individual structural elements within each tree (Disney et al., 2018; Raunonen et al., 2013; Stovall & Shugart, 2018; Wilkes et al., 2017). Close-range and proximal instruments offer highly resolved data that is not possible with other approaches to FSD measurement. However, adopting close-range and proximal instruments to measure FSD comes with a trade-off, either in the ability to gather detailed information for a single tree or structural variability across several trees, or in how much area can be sampled—as proximal sensors can only sample a small spatial extent (10s–100s of m^{-2}) due to data collection times and necessary computational resources (Wilkes et al., 2017).

UAV/UAS-based sensors can outpace both satellite and piloted platforms with ultrahigh horizontal and vertical resolutions as well as shorter time between visits (Singh & Frazier, 2018). UAV imagery is less affected by the atmosphere given the proximity between the sensor and the Earth surface—beneath the cloud layer. Flexibility across different applications and decreasing

platform costs are at least partly responsible for the increasing adoption of UAVs (Tang & Shao, 2015). Significant advancements have been made in recent years in the use of UAV platforms for forest structural assessments (Almeida et al., 2021; Almeida, Broadbent, et al., 2019). Standards need to be determined on acceptable horizontal and vertical resolutions, flight altitudes, minimum imagery overlap, placement and number of ground control or reference points, optimal flight times with respect to solar elevation angle, photogrammetric processing methods, data reporting, metadata archival, and appropriate minimum mapping units, particularly for using UAV-based lidar for FSD measurement (Jozkow et al., 2016; Pádua et al., 2017; Petrie, 2013).

Airborne remote sensing platforms cover a greater spatial extent than either terrestrial or UAV/UAS platforms, but at lower spatial resolution due to factors like flying altitude and speed. The small-footprint sampling of airborne platforms (i.e., several laser pulses per square meter) is suitable for measuring vegetation clumping, canopy gaps, and individual tree properties. Models of canopy height, shape, roughness, fractional cover, biomass, and other structural properties can be generated from aerial lidar at the spatial grain of interest for a given ecosystem function or structural property. 3D point clouds can also be used to analyze canopy layering or stratification between understory, mid-story, and upper canopy vegetation, and describe the horizontal and vertical variability in vegetation structure among individuals across the landscape (Jung et al., 2013; Whitehurst et al., 2013). Canopy height profiles generated from airborne lidar have been shown to describe the vertical distribution of canopy structure and correlate with forest age and composition (Hakkenberg et al., 2018), and they can also be used to characterize floristic diversity (Cavender-Bares et al., 2020; Hakkenberg & Goetz, 2021). Individual tree detection (ITD) and crown segmentation algorithms are advanced computational approaches that allow for quantifying FSD at the individual tree level from lidar point clouds or canopy height models (Silva et al., 2022; Valbuena et al., 2014) and are useful for estimating biophysical parameters such as biomass, volume, and canopy cover (Silva et al., 2016; Stark et al., 2012).

Spaceborne lidar sensors such as GEDI, a full-waveform lidar sensor with a 25-m ground-level footprint, and ICESat-2, a photon-counting lidar sensor with a 10-m ground-level footprint, provide 10-return data at approximately the same scale as typical field-sampling plots (Abdalati et al., 2010). Several trees are combined within a footprint and differences in structure in the horizontal dimension are lost. Canopy height, canopy cover, and vertical layering can also be analyzed. Using lidar data with radiative transfer models allows for the extraction of

more advanced structural information such as vertical optical gap probability and vertical leaf area density. Spaceborne lidar missions achieve near-global coverage but do not have the wall-to-wall sampling that is common from airborne systems, with small gaps between sequential laser pulses (60 m for GEDI) and larger gaps between laser ground tracks (600 m for GEDI) (Dubayah et al., 2020), but interpolation methods based on fusion with ancillary datasets like Landsat (Li et al., 2020; Narine et al., 2019; Potapov et al., 2020) or TanDEM-X (Qi et al., 2019; Stovall et al., 2020) are enabling high-resolution wall-to-wall maps of forest structure (Hakkenberg et al., 2023; Lang et al., 2022; Wang et al., 2022).

Spaceborne sensors have a large extent, but coarse resolution, and are therefore more applicable at regional (Smart et al., 2012) or global (Tang et al., 2019) scales; FSD measurements derived from such sensors are a reliable surrogate for other branches of biodiversity (e.g., taxonomic, trait, or phylogenetic) while capturing the variability of diversity among multiple taxa (Pommerening, 2002). Attributed to faster return rate of some orbiting satellites (MODIS Terra and Aqua), temporally sensitive FSD measurements (e.g., greenness for phenology) can be derived from such spaceborne sensors. Consequently, these FSD measurements can help researchers understand successional dynamics, impacts of disturbances, community/ecosystem resilience, post-perturbation community recovery, and unravel landscape patterns and habitat connectivity (Francis et al., 2023; Garabedian et al., 2017; Newman, 2019; Newman et al., 2019; Possingham et al., 2005; Romme, 2005).

Considerations for using remote sensing for FSD measurement

The diverse array of remote sensing approaches leads to data products that vary in spatial and temporal extent. There are inherent trade-offs in spatial extent versus spatial resolution for remote sensing approaches—large spatial extents, coarse resolution from by air- and spaceborne remote sensing versus small extent, very fine spatial resolution from proximal remote sensing (Figures 2 and 3). Several factors constrain the trade-off between spatial resolution and extent including sensor specifications and signal-to-noise ratio, sensor altitude, uncertainty in geolocation, sensor pointing precision, as well as signal attenuation and ground finding issues in dense canopies. Proximal remote sensing instruments such as TLS or UAV-based systems are expensive, with high instrument costs ranging into 100s of thousands of

USD, along with subsequent high costs for proprietary software licenses. Most air- and spaceborne remote sensing platforms are far too expensive for a singular lab or research group to afford and are typically supported by for-profit companies, research networks, federal agencies, countries, or consortiums. While data generated from platforms owned by for-profit companies (e.g., Planet Labs) require purchase, data from platforms owned or operated by other means tend to be publicly available (e.g., Landsat, MODIS, GEDI). Thus, the use of remote sensing for research relies heavily on large grants or existing publicly available data. Another consideration with the use of remote sensing data is the necessity for specialized training and software. Software environments for working with remote sensing data require a significant investment in training, whether it be for proprietary (e.g., ENVI, ArcGIS) or open-source solutions (e.g., Google Earth Engine, R, Python). Remote sensing work often requires more advanced computing solutions, though cloud computing services (e.g., Google Earth Engine, Jupyter Notebooks) are helping address this issue.

SCIENCE, POLICY, AND MANAGEMENT IMPLICATIONS

FSD measurement has profound implications beyond research, extending into social, political, and economic spheres. Standardized FSD measurement could strengthen environmental policies and be used to assess progress toward achieving biodiversity, climate, and natural resource management goals (Jetz et al., 2019) and forest structure–wildlife fitness relationships to guide endangered species recovery (Garabedian et al., 2021). Increasing the availability, consistency, and utility of FSD measurements for compliance with forest carbon offset programs (e.g., California's forest carbon offset program) can further bolster the effectiveness of policy implementation and help scientifically informed, data-driven policy reforms. For instance, FSD measurements could be integrated directly as essential biodiversity variables (EBV) or indirectly used as environmental predictors of species distributions through occupancy modeling or ecological niche modeling (Jetz et al., 2019). FSD metrics yield information on the status and trends in key biodiversity targets, which then can be used to gauge the impacts of international treaties (e.g., Convention of Biological Diversity) that strive toward biodiversity conservation targets like the UN's Sustainable Development Goals. One necessary step toward the integration of FSD into research and management is to increase the availability and visibility of FSD data. Such precedents exist. The




FIGURE 3 Graphical depictions of real data taken from many different platforms, perspectives, and sensors used in forest structural diversity metric calculation. Traditional measurements of dbh (dbh tape) and stem locations (compass and transect tape) used to make stem maps and tree size distributions (data from Harvard Forest, NEON plot HARV_036; NEON, 2022a). Terrestrial measurements of canopy structure from a 3D terrestrial lidar scanner and hemispherical camera (FoRTE Project, University of Michigan Biological Station; Atkins et al., 2021). Spaceborne data examples including a GEDI waveform (HARV_036; Dubayah et al., 2020) and backscatter from synthetic aperture radar (African mangrove; ALOS PALSAR; Shimada et al., 2014). Airborne data showing a canopy height model, point cloud, and segmented forest canopy (HARV_036; NEON, 2022b) and hyperspectral signature (HARV_036; NEON, 2023), all from NEON AOP data. UAV structure-from-motion point cloud (Pace Estate; A. Stovall) and RGB orthoimagery (Pace Estate, VA; Atkins et al., 2020).

Landsat Archive, made available freely and openly by the United States Geological Survey (USGS), has profoundly enabled our ability to understand global ecosystem dynamics (Cohen & Goward, 2004; Wulder et al., 2012). Related

progress whereby more FSD data are made available (e.g., National Ecological Observatory Network–NEON), as well as the tools and training to work with those data, could create a similar data revolution.

Robust FSD measurements that effectively capture changes in the 3D structure of forests beyond gross biomass assessments—such as forest regeneration, reforestation, degradation, deforestation, plant invasions, forest maturity, and successional stages, as well as tree size classes and differentiation between biomass sources (above/belowground, deciduous/coniferous, woody/non-woody)—can also help implement and sustain international policies geared toward greenhouse gas reductions, such as REDD+. Defensible measurement of forest carbon is one of the major challenges of carbon accounting (Badgley et al., 2021) and, therefore, is a potential barrier to more widespread implementation of forest carbon offset programs. Airborne-, satellite-, and UAV-based lidar could address some of the challenges in monitoring carbon for forest offset programs and increase the accuracy of carbon monitoring, helping to address bias created by scaling point measurements to landscapes (Disney et al., 2019; Marvin et al., 2014; Muller-Landau, 2009; Stovall & Shugart, 2018).

CURRENT LIMITATIONS AND POTENTIAL FUTURE DIRECTIONS OF FSD MEASUREMENT

A century or more of advancement in theory and instrumentation has generated a remarkable array of ecologically meaningful FSD measurements, yet there remain key gaps in our current ability to measure FSD, namely sufficient data collection and the ability to fuse traditional and remote sensing approaches, a need for robust theory underpinning FSD measurement and applications, standardized methodologies for deriving FSD, robust comparisons of FSD derivation and the underlying statistics and assumptions being made, approaches to intercomparison and scaling, technological constraints, data availability, and the ability to acquire and process data.

There is a need for high-quality, high-density data for the entire globe. However, perhaps even more important is the need to formulate novel, reproducible analytical approaches to accommodate comparisons among existing and future data products that vary in spatial, temporal, and spectral resolution to ensure backward compatibility when detecting temporal trends of FSD. The same challenge manifests at larger spatial extents, such as latitudinal or cross-continental comparisons of FSD that may require integration of spatially extensive remote sensing data of variable quality, collected from differential sensors or platforms into comparable measurement standards (Valbuena et al., 2020). Data integration or data fusion between passive and active remote sensing, between spaceborne and airborne remote sensing, and

between from-above and ground-based remote sensing holds promise to increase FSD detection precision and accuracy (e.g., Hakkenberg et al., 2023) as well as addressing issues of scale. Cross-platform comparisons of measurements across sensors, platforms, and scale are needed.

Repeat, high-resolution estimates of canopy structure enabled by remote sensing methods such as lidar and radar could transform our understanding of forest canopy dynamics, building off multiple decades of research enabled by long-term forest monitoring plots. Measurements of forest dynamics that have been made from traditional methods (e.g., forest inventories) have enabled insight into tree diameter growth, mortality, and gap disturbances, and how those dynamics are changing with climate change (Van Mantgem et al., 2009). However, scaling inferences from tree diameters to canopies requires simplifying assumptions about tree allometry and 3D space-filling of forest canopies, which may introduce error (Bohlman & Pacala, 2011; Farrior et al., 2016). Novel and creative means will be necessary to fuse these approaches. ITD methods, whereby algorithms are used with remote sensing data to identify individual trees via crown segmentation, may fill a crucial gap here, linking point measurements on the ground with specific individuals identified from remote sensing data taken above the canopy (Silva et al., 2022). New measurements of canopy structure from airborne and terrestrial lidar, collected systematically at the same site over comparable time frames to tree lifespans, are needed to understand the dynamics of forest structure and biomass, and how they are responding to global change. Presently, airborne lidar datasets exist for many sites from single overflights. However, many of these datasets lack repeat measurements, and when repeat measurements exist, they rarely span more than a decade (e.g., Dalagnol et al., 2021; Kellner & Asner, 2014). Repeat measurements taken over multiple decades will enable understanding of competitive interactions within forest ecosystems, and how forest structure and composition are changing in response to global and local influences in climate, nutrient availability, and land use.

The theoretical limitations facing FSD measurement are nontrivial and have practical ramifications. To our knowledge, no comprehensive list of all FSD measurements has been published to date. While this effort would be arduous and time-consuming given the range of sensors and algorithms employed and the pace at which advancement occurs, such a list would be a valuable tool to guide future research as well as aid the standardization of FSD derivation. Lacking a thorough inventory of the landscape of FSD measurement, creating consensus and standards is difficult. Currently, there are disjunct reports on various approaches to FSD measurement focused on

specific subfields, thus hindering applications of FSD in future research. There is also no consensus about the spatial scale or grain size at which many FSD measurements should be taken, nor is there a robust statistical treatment or analysis of the role of scale (Garabedian et al., 2014). FSD metrics are often calculated at the scale of interest of a given function (Beland et al., 2019) without explicit regard given to the stability or scale-dependency of those FSD metrics (Atkins et al., 2023). There is a clear need to understand how spatial grain and extent factor into FSD measurement. FSD measurements that capture spatial and temporal change in forest structure are needed to track changes in structural diversity equivalent to alpha, beta, and gamma diversity (Fotis, Morin, et al., 2018; Fotis, Murphy, et al., 2018; Latifi & Valbuena, 2019). It is also necessary to invest efforts in the identification, classification, and detection of attributes of structural diversity that are rare, uncommon, or critically important, but difficult to measure as well as the development of indices that are sensitive to changes of such rare structural features.

Advances in FSD theory will also catalyze work to characterize broad structural patterns across biomes (Atkins et al., 2022; Ehbrecht et al., 2021; Fahey et al., 2019; Hakkenberg & Goetz, 2021), codify FSD metric derivation (Walter et al., 2021), and incorporate theory from disparate fields into FSD measurement (Shiklomanov et al., 2019; Verbeeck et al., 2019). Integrating FSD measurement with existing field survey data and in situ sensor networks has a huge potential to advance research, as does integration with mechanistic modeling approaches (e.g., use of FSD measurements to test structural assumptions of ecosystem models and/or to create more parsimonious models using integrative FSD metrics). Measurements of forest dynamics made from forest inventory data have provided insight into tree diameter growth, mortality, and gap disturbances, and how those dynamics are changing with climate change (Van Mantgem et al., 2009). However, scaling inferences from individuals to canopies requires simplifying assumptions about tree allometry and 3D space-filling of forest canopies (Farrior et al., 2016; Fisher et al., 2018), which may introduce error. Additionally, many remote sensing data products can be fused to create more robust measures of FSD, yet more work is needed to refine algorithms (Silva et al., 2021). Data fusion of such heterogeneous data streams may alleviate these issues but may require further advances in modeling, information-theoretic approaches, and cyber infrastructure. Composite metrics or indices could be created by combining disparate data, providing further understanding of the natural world. While technological and theoretical advances will move FSD measurement forward in the coming decade, there are current needs and

opportunities within and across disciplines that can be addressed in the near term (Valbuena et al., 2020).

One key issue is the limitations that exist in the ability to access, process, and use FSD data. Development of global, national, and regional spatial data infrastructure initiatives aims to improve sustainable natural resource and environmental management by optimizing cross-national, multiagency partnerships. Infrastructure initiatives should fundamentally focus on facilitation and coordination of spatial data exchange among stakeholders from different jurisdictions and applied fields, fostering partnerships among governmental agencies, intergovernmental partnerships, nonprofit environmental organizations, commercial sector, and academic and research institutes (Rajabifard & Williamson, 2001). The NEON, NASA, OpenTopography, and the USGS 3D Elevation Program are examples of free resources for remote sensing data that range from site-level to global data coverage. We must also consider the necessity of increased abundance of and access to training and education resources to work with FSD data and tools. Open-source platforms that can be used to process and quantify FSD data do exist (e.g., QGIS, Fusion, Google Earth Engine, Python, R, CloudCompare). While educational resources providing free tutorials and webinars (e.g., [NEONscience.org](https://neonscience.org), earthdatascience.org, earthdata.nasa.gov, Data Carpentry) for accessing and working with FSD data also exist, allocation of greater financial resources to support these efforts is needed. The future of FSD measurement lies at the nexus of people and technology. The ability to work with these data must not be limited by privilege or access (Dwivedi et al., 2022).

CONCLUSIONS

Advances in technology, data distribution, and theory have driven the evolution of FSD measurement; however, further work is needed to realize its full potential. Grounding FSD measurement in theory and creating standardized approaches will expand our ability to understand forest and ecosystem dynamics and make cross-scale and -system comparisons. We face an abundance of data coupled with the need to develop rigorous methods and theories to work with that data. We also must empower and enable the community to work with these data. While there is much work to do, thoughtful consideration of how FSD can be used to broaden the scope of ecological research may be transformative for all branches of ecological and environmental research.

ACKNOWLEDGMENTS

This manuscript was conceived at a National Science Foundation (NSF)-funded workshop, “Exploring New

Dimensions of Forest Ecosystems with Structural Diversity.” NSF award number 1924942 was granted to Principal Investigators/Co-Authors Drs. Elizabeth A. LaRue and Brady S. Hardiman and PI Dr. Songlin Fei.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

Datasets utilized for demonstration purposes are available in: Shimada et al. (2014) (<https://doi.org/10.1016/j.rse.2014.04.014>), Atkins et al. (2021) (<https://doi.org/10.5194/essd-13-943-2021>), NEON (2022a, 2022b) (<https://doi.org/10.48443/73zn-k414>, <https://doi.org/10.48443/ymp-fr93>), and NEON (2023) (<https://doi.org/10.48443/ehnm-xd88>).

ORCID

Jeff W. Atkins  <https://orcid.org/0000-0002-2295-3131>

Parth Bhatt  <https://orcid.org/0000-0001-9311-7870>

Luis Carrasco  <https://orcid.org/0000-0003-4283-7725>

Emily Francis  <https://orcid.org/0000-0002-7123-781X>

James E. Garabedian  <https://orcid.org/0000-0002-3136-016X>

Christopher R. Hakkenberg  <https://orcid.org/0000-0002-6579-5954>

Brady S. Hardiman  <https://orcid.org/0000-0001-6833-9404>

Anil Koirala  <https://orcid.org/0000-0002-4286-569X>

Elizabeth A. LaRue  <https://orcid.org/0000-0002-9535-0630>

Sungchan Oh  <https://orcid.org/0000-0003-2337-9693>

Atticus E. L. Stovall  <https://orcid.org/0000-0001-9512-3318>

Thilina D. Surasinghe  <https://orcid.org/0000-0001-8758-072X>

Tao Zhang  <https://orcid.org/0000-0001-7135-1762>

REFERENCES

- Abdalati, W., H. J. Zwally, R. Bindenschadler, B. Csatho, S. L. Farrell, H. A. Fricker, D. Harding, et al. 2010. “The ICESat-2 Laser Altimetry Mission.” *Proceedings of the IEEE* 98: 735–751.
- Almeida, D. R. A., E. N. Broadbent, A. M. A. Zambrano, B. E. Wilkinson, M. E. Ferreira, R. Chazdon, P. Meli, et al. 2019. “Monitoring the Structure of Forest Restoration Plantations with a Drone-Lidar System.” *International Journal of Applied Earth Observation and Geoinformation* 79: 192–98. <https://doi.org/10.1016/j.jag.2019.03.014>.
- Almeida, D. R. A., E. N. Broadbent, M. P. Ferreira, P. Meli, A. M. A. Zambrano, E. B. Gorgens, A. F. Resende, et al. 2021. “Monitoring Restored Tropical Forest Diversity and Structure through UAV-Borne Hyperspectral and Lidar Fusion.” *Remote Sensing of Environment* 264: 112582. <https://doi.org/10.1016/j.rse.2021.112582>.
- Almeida, D. R. A., S. C. Stark, R. Chazdon, B. W. Nelson, R. G. Cesar, P. Meli, E. B. Gorgens, et al. 2019. “The Effectiveness of Lidar Remote Sensing for Monitoring Forest Cover Attributes and Landscape Restoration.” *Forest Ecology and Management* 438: 34–43. <https://doi.org/10.1016/j.foreco.2019.02.002>.
- Andersen, H. E., S. E. Reutebuch, and R. J. McGaughey. 2006. “A Rigorous Assessment of Tree Height Measurements Obtained Using Airborne Lidar and Conventional Field Methods.” *Canadian Journal of Remote Sensing* 32(5): 355–366.
- Asner, G. P., R. E. Martin, D. E. Knapp, R. Tupayachi, C. B. Anderson, F. Sinca, N. R. Vaughn, and W. Llactayo. 2017. “Airborne Laser-Guided Imaging Spectroscopy to Map Forest Trait Diversity and Guide Conservation.” *Science* 355(6323): 385–89.
- Atkins, J. W., E. Agee, A. Barry, K. M. Dahlin, K. Dorheim, M. S. Grigri, L. T. Haber, et al. 2021. “The Fortedata R Package: Open-Science Datasets from a Manipulative Experiment Testing Forest Resilience.” *Earth System Science Data* 13(3): 943–952.
- Atkins, J. W., G. Bohrer, R. T. Fahey, B. S. Hardiman, T. H. Morin, A. E. Stovall, N. Zimmerman, and C. M. Gough. 2018. “Quantifying Vegetation and Canopy Structural Complexity from Terrestrial LiDAR Data Using the ForestR Package.” *Methods in Ecology and Evolution* 9(10): 2057–66.
- Atkins, J. W., B. Bond-Lamberty, R. T. Fahey, B. S. Hardiman, L. Haber, E. Stuart-Haëntjens, E. LaRue, et al. 2019. “Multidimensional Structural Characterization Is Required to Detect and Differentiate among Moderate Disturbance Agents.” *Ecosphere* 11(6): e03156.
- Atkins, J. W., J. Costanza, K. M. Dahlin, M. P. Dannenberg, A. J. Elmore, M. C. Fitzpatrick, C. R. Hakkenberg, et al. 2023. “Scale Dependency of Lidar-Derived Forest Structural Diversity.” *Methods in Ecology and Evolution* 14(2): 708–723. <https://doi.org/10.1111/2041-210x.14040>.
- Atkins, J. W., A. E. Stovall, and X. Yang. 2020. “Mapping Temperate Forest Phenology Using Tower, UAV, and Ground-Based Sensors.” *Drones* 4(3): 56.
- Atkins, J. W., J. A. Walter, A. E. Stovall, R. T. Fahey, and C. M. Gough. 2022. “Power Law Scaling Relationships Link Canopy Structural Complexity and Height across Forest Types.” *Functional Ecology* 36(3): 713–726. <https://doi.org/10.1111/1365-2435.13983>.
- Avery, T. E., and H. E. Burkhardt. 2015. *Forest Measurements*. Long Grove, IL: Waveland Press.
- Badgley, G., J. Freeman, J. J. Hamman, B. Haya, A. T. Trugman, W. R. Anderegg, and D. Cullenward. 2021. “Systematic Over-Crediting in California’s Forest Carbon Offsets Program.” bioRxiv.
- Beland, M., G. Parker, B. Sparrow, D. Harding, L. Chasmer, S. Phinn, A. Antonarakis, and A. Strahler. 2019. “On Promoting the Use of Lidar Systems in Forest Ecosystem Research.” *Forest Ecology and Management* 450: 117484.
- Bhatt, P., A. Maclean, Y. Dickinson, and C. Kumar. 2022. “Fine-Scale Mapping of Natural Ecological Communities Using Machine Learning Approaches.” *Remote Sensing* 14(3): 563. <https://doi.org/10.3390/rs14030563>.
- Blakey, R. V., B. S. Law, R. T. Kingsford, and J. Stoklosa. 2017. “Terrestrial Laser Scanning Reveals Below-Canopy Bat Trait Relationships with Forest Structure.” *Remote Sensing of Environment* 198: 40–51.

- Bohlman, S., and S. Pacala. 2011. "A Forest Structure Model That Determines Crown Layers and Partitions Growth and Mortality Rates for Landscape-Scale Applications of Tropical Forests." *Journal of Ecology* 100(2): 508–518. <https://doi.org/10.1111/j.1365-2745.2011.01935.x>.
- Borcard, D., P. Legendre, and P. Drapeau. 1992. "Partialling Out the Spatial Component of Ecological Variation." *Ecology* 73(3): 1045–55.
- Calders, K., J. Adams, J. Armston, H. Bartholomeus, S. Bauwens, L. P. Bentley, J. Chave, et al. 2020. "Terrestrial Laser Scanning in Forest Ecology: Expanding the Horizon." *Remote Sensing of Environment* 251: 112102.
- Calders, K., G. Newnham, A. Burt, S. Murphy, P. Raunonen, M. Herold, D. Culvenor, et al. 2015. "Nondestructive Estimates of Above-Ground Biomass Using Terrestrial Laser Scanning." *Methods in Ecology and Evolution* 6: 198–208. <https://doi.org/10.1111/2041-210X.12301>.
- Cavender-Bares, J., J. A. Gamon, and P. A. Townsend. 2020. *Remote Sensing of Plant Biodiversity*. Cham: Springer. <https://doi.org/10.1007/978-3-030-33157-3>.
- Cohen, W. B., and S. N. Goward. 2004. "Landsat's Role in Ecological Applications of Remote Sensing." *BioScience* 54(6): 535–545.
- Coops, N. C., P. Tompalski, T. R. H. Goodbody, M. Queinnec, J. E. Luther, D. K. Bolton, J. C. White, M. A. Wulder, O. R. van Lier, and T. Hermosilla. 2021. "Modelling Lidar-Derived Estimates of Forest Attributes over Space and Time: A Review of Approaches and Future Trends." *Remote Sensing of Environment* 260: 112477.
- Dalagnol, R., F. H. Wagner, L. S. Galvão, A. S. Streher, O. L. Phillips, E. Gloor, T. A. M. Pugh, J. P. H. B. Ometto, and L. E. O. C. Aragão. 2021. "Large-Scale Variations in the Dynamics of Amazon Forest Canopy Gaps from Airborne Lidar Data and Opportunities for Tree Mortality Estimates." *Scientific Reports* 11: 1–14.
- Disney, M., A. Burt, K. Calders, C. Schaaf, and A. Stovall. 2019. "Innovations in Ground and Airborne Technologies as Reference and for Training and Validation: Terrestrial Laser Scanning (TLS)." *Surveys in Geophysics* 40: 937–958. <https://doi.org/10.1007/s10712-019-09527-x>.
- Disney, M., P. Lewis, and P. Saich. 2006. "3D Modelling of Forest Canopy Structure for Remote Sensing Simulations in the Optical and Microwave Domains." *Remote Sensing of Environment* 100(1): 114–132.
- Disney, M. I., V. M. Boni, A. Burt, K. Calders, S. L. Lewis, P. Raunonen, and P. Wilkes. 2018. "Weighing Trees with Lasers: Advances, Challenges and Opportunities." *Interface Focus* 8: 20170048. <https://doi.org/10.1098/rsfs.2017.0048>.
- Du, J., J. S. Kimball, I. Velicogna, M. Zhao, L. A. Jones, J. D. Watts, and Y. Kim. 2019. "Multicomponent Satellite Assessment of Drought Severity in the Contiguous United States from 2002 to 2017 Using AMSR-E and AMSR2." *Water Resources Research* 55(7): 5394–5412.
- Dubayah, R., J. B. Blair, S. Goetz, L. Fatoyinbo, M. Hansen, S. Healey, M. Hofton, et al. 2020. "The Global Ecosystem Dynamics Investigation: High-Resolution Laser Ranging of the Earth's Forests and Topography." *Science of Remote Sensing* 1: 1–14.
- Dwivedi, D., A. L. D. Santos, M. A. Barnard, T. M. Crimmins, A. Malhotra, K. A. Rod, K. S. Aho, et al. 2022. "Biogeosciences Perspectives on Integrated, Coordinated, Open, Networked (ICON) Science." *AGU Earth and Space Science* 9(3): e2021EA002119.
- Ehbrecht, M., P. Schall, C. Ammer, and D. Seidel. 2017. "Quantifying Stand Structural Complexity and Its Relationship with Forest Management, Tree Species Diversity and Microclimate." *Agricultural and Forest Meteorology* 242: 1–9.
- Ehbrecht, M., P. Schall, J. Juchheim, C. Ammer, and D. Seidel. 2016. "Effective Number of Layers: A New Measure for Quantifying Three-Dimensional Stand Structure Based on Sampling with Terrestrial LiDAR." *Forest Ecology and Management* 380: 212–223.
- Ehbrecht, M., D. Seidel, P. Annighöfer, H. Kreft, M. Köhler, D. C. Zemp, K. Puettmann, et al. 2021. "Global Patterns and Climatic Controls of Forest Structural Complexity." *Nature Communications* 12(1): 1–12.
- Eitel, J. U., B. Höfle, L. A. Vierling, A. Abellán, G. P. Asner, J. S. Deems, C. L. Glennie, et al. 2016. "Beyond 3-D: The New Spectrum of Lidar Applications for Earth and Ecological Sciences." *Remote Sensing of Environment* 186: 372–392.
- Erb, K.-H., T. Kastner, C. Plutzer, A. L. S. Bais, N. Carvalhais, T. Fetzl, S. Gingrich, et al. 2018. "Unexpectedly Large Impact of Forest Management and Grazing on Global Vegetation Biomass." *Nature* 553(7686): 73–76.
- Fahey, R. T., J. W. Atkins, C. M. Gough, B. S. Hardiman, L. E. Nave, J. M. Tallant, K. J. Nadehoffer, et al. 2019. "Defining a Spectrum of Integrative Trait-Based Vegetation Canopy Structural Types." *Ecology Letters* 22(12): 2049–59.
- Farrior, C. E., S. A. Bohlman, S. Hubbell, and S. W. Pacala. 2016. "Dominance of the Suppressed: Power-Law Size Structure in Tropical Forests." *Science* 351: 155–57.
- Fisher, R. A., C. D. Koven, W. R. L. Anderegg, B. O. Christofferson, M. C. Dietze, C. E. Farrior, J. A. Holm, et al. 2018. "Vegetation Demographics in Earth System Models: A Review of Progress and Priorities." *Global Change Biology* 24: 35–54.
- Fotis, A. T., T. H. Morin, R. T. Fahey, B. S. Hardiman, G. Bohrer, and P. S. Curtis. 2018. "Forest Structure in Space and Time: Biotic and Abiotic Determinants of Canopy Complexity and their Effects on Net Primary Productivity." *Agricultural and Forest Meteorology* 250: 181–191.
- Fotis, A. T., S. J. Murphy, R. D. Ricart, M. Krishnadas, J. Whitacre, J. W. Wenzel, S. A. Queenborough, and L. S. Comita. 2018. "Above-Ground Biomass Is Driven by Mass-Ratio Effects and Stand Structural Attributes in a Temperate Deciduous Forest." *Journal of Ecology* 106(2): 561–570.
- Francis, E. J., J. A. Lutz, and C. E. Farrior. 2023. "Elevated Mortality Rates of Large Trees Allow for Increased Frequency of Intermediate Trees: A Hypothesis Supported by Demographic Model Comparison with Plot and LIDAR Data." *Forest Ecology and Management* 540: 121035. <https://doi.org/10.1016/j.foreco.2023.121035>.
- Frolking, S., M. W. Palace, D. B. Clark, J. Q. Chambers, H. H. Shugart, and G. C. Hurtt. 2009. "Forest Disturbance and Recovery: A General Review in the Context of Spaceborne Remote Sensing of Impacts on Aboveground Biomass and Canopy Structure." *Journal of Geophysical Research: Biogeosciences* 114(G2).
- Garabedian, J. E., R. J. McGaughey, S. E. Reutebuch, B. R. Parresol, J. C. Kilgo, C. E. Moorman, and M. N. Peterson. 2014. "Quantitative Analysis of Woodpecker Habitat Using

- High-Resolution Airborne LiDAR Estimates of Forest Structure and Composition.” *Remote Sensing of Environment* 145: 68–80.
- Garabedian, J. E., C. E. Moorman, M. N. Peterson, and J. C. Kilgo. 2017. “Use of LiDAR to Define Habitat Thresholds for Forest Bird Conservation.” *Forest Ecology and Management* 399: 24–36.
- Garabedian, J. E., C. E. Moorman, M. N. Peterson, and J. C. Kilgo. 2021. “Group Size Mediates Effects of Intraspecific Competition and Forest Structure on Productivity in a Recovering Social Woodpecker Population.” *Animal Conservation* 25(3): 438–452. <https://doi.org/10.1111/acv.12757>.
- Goff, F. G., and P. H. Zedler. 1968. “Structural Gradient Analysis of Upland Forests in the Western Great Lakes Area.” *Ecological Monographs* 38(1): 65–86. <https://doi.org/10.2307/1948537>.
- Hakkenberg, C. R., and S. J. Goetz. 2021. “Climate Mediates the Relationship between Plant Biodiversity and Forest Structure across the Contiguous United States.” *Global Ecology and Biogeography*. 30: 2245–58.
- Hakkenberg, C. R., R. K. Peet, D. L. Urban, and C. Song. 2018. “Modeling Plant Composition as Community Continua in a Forest Landscape with LiDAR and Hyperspectral Remote Sensing.” *Ecological Applications* 28(1): 177–190.
- Hakkenberg, C. R., H. Tang, P. Burns, and S. J. Goetz. 2023. “Canopy Structure from Space Using GEDI Lidar.” *Frontiers in Ecology and the Environment*. 21: 55–56.
- Hall, F. G., D. B. Botkin, D. E. Strebel, K. D. Woods, and S. J. Goetz. 1991. “Large-Scale Patterns of Forest Succession as Determined by Remote Sensing.” *Ecology* 72(2): 628–640.
- Heidrich, L., S. Bae, S. Levick, S. Seibold, W. Weisser, P. Krzystek, P. Magdon, et al. 2020. “Heterogeneity-Diversity Relationships Differ between and within Trophic Levels in Temperate Forests.” *Nature Ecology and Evolution* 4: 1204–12.
- Henderson, F. M., and A. J. Lewis. 1998. *Principles and Applications of Imaging Radar. Manual of Remote Sensing: Volume 2*. New York: Wiley and Sons.
- Hernando, A., L. A. Arroyo, J. Velázquez, and R. Tejera. 2012. “Objects-Based Image Analysis for Mapping Natura 2000 Habitats to Improve Forest Management.” *Photogrammetric Engineering & Remote Sensing* 78(9): 991–99.
- Hovind, H. J., and C. E. Rieck. 1961. “Basal Area and Point-Sampling.” *Wisconsin Department of Natural Resources Technical Bulletin* 23(16): 23–30.
- Imhoff, M. L. 1995. “Radar Backscatter and Biomass Saturation: Ramifications for Global Biomass Inventory.” *IEEE Transactions on Geoscience and Remote Sensing* 33(2): 511–18.
- Jenkins, J. C., D. C. Chojnacky, L. S. Heath, and R. A. Birdsey. 2003. “National-Scale Biomass Estimators for United States Tree Species.” *Forest Science* 49(1): 12–35.
- Jetz, W., M. A. McGeoch, R. Guralnick, S. Ferrier, J. Beck, M. J. Costello, M. Fernandez, et al. 2019. “Essential Biodiversity Variables for Mapping and Monitoring Species Populations.” *Nature Ecology & Evolution* 3(4): 539–551.
- Joshi, N., E. T. A. Mitchard, M. Brolly, J. Schumacher, A. Fernández-Landa, V. K. Johannsen, M. Marchamalo, and R. Fensholt. 2017. “Understanding ‘Saturation’ of Radar Signals over Forests.” *Scientific Reports* 7(1): 3505. <https://doi.org/10.1038/s41598-017-03469-3>.
- Jozkow, G., C. Totha, and D. Grejner-Brzezinska. 2016. “UAS Topographic Mapping with Velodyne LiDAR Sensor.” *ISPRS Annals of Photogrammetry, Remote Sensing & Spatial Information Sciences* 3: 201–8.
- Jucker, T. 2021. “Deciphering the Fingerprint of Disturbance on the Three-Dimensional Structure of the World’s Forests.” *New Phytologist* 233(2): 612–17. <https://doi.org/10.1111/nph.17729>.
- Jung, J., B. K. Pekin, and B. C. Pijanowski. 2013. “Mapping Gaps in an Old-Growth, Secondary-Growth, and Selectively-Logged Tropical Rainforest Using Discrete Return LIDAR.” *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* 6(6): 2453–61.
- Kellner, J. R., and G. P. Asner. 2014. “Winners and Losers in the Competition for Space in Tropical Forest Canopies.” *Ecology Letters* 17(5): 556–562.
- Knyazikhin, Y., M. A. Schull, P. Stenberg, M. Möttus, M. Rautiainen, Y. Yang, A. Marshak, et al. 2013. “Hyperspectral Remote Sensing of Foliar Nitrogen Content.” *Proceedings of the National Academy of Sciences of the United States of America* 110(3): E185–E192.
- Kriegler, F., W. Malila, R. Nalepka, and W. Richardson. 1969. “Preprocessing Transformations and Their Effect on Multispectral Recognition.” In *Proceedings of the 6th International Symposium on Remote Sensing of Environment*, 97–131. Ann Arbor, MI: University of Michigan.
- Kruys, N., J. Fridman, F. Götmark, P. Simonsson, and L. Gustafsson. 2013. “Retaining Trees for Conservation at Clearcutting Has Increased Structural Diversity in Young Swedish Production Forests.” *Forest Ecology and Management* 304: 312–321.
- Lang, N., N. Kalischek, J. Armston, K. Schindler, R. Dubayah, and J. D. Wegner. 2022. “Global Canopy Height Regression and Uncertainty Estimation from GEDI LIDAR Waveforms with Deep Ensembles.” *Remote Sensing of Environment* 268: 112760.
- LaRue, E. A., R. T. Fahey, B. C. Alvishere, J. W. Atkins, P. Bhatt, B. Buma, A. Chen, et al. 2023. “A Theoretical Framework for the Ecological Role of Three-Dimensional Structural Diversity.” *Frontiers in Ecology and the Environment* 21(1): 4–13. <https://doi.org/10.1002/fee.2587>.
- LaRue, E. A., B. S. Hardiman, J. M. Elliott, and S. Fei. 2019. “Structural Diversity as a Predictor of Ecosystem Function.” *Environmental Research Letters* 14(11): 114011.
- Latifi, H., and R. Valbuena. 2019. “Current Trends in Forest Ecological Applications of Three-Dimensional Remote Sensing: Transition from Experimental to Operational Solutions?” *Forests* 10(10): 891.
- Lefsky, M. A., W. B. Cohen, G. G. Parker, and D. J. Harding. 2002. “Lidar Remote Sensing for Ecosystem Studies: Lidar, an Emerging Remote Sensing Technology that Directly Measures the Three-Dimensional Distribution of Plant Canopies, Can Accurately Estimate Vegetation Structural Attributes, and Should Be of Particular Interest to Forest, Landscape, and Global Ecologists.” *BioScience* 52: 19–30.
- Li, Y., M. Li, C. Li, and Z. Liu. 2020. “Forest Aboveground Biomass Estimation Using Landsat 8 and Sentinel-1A Data with Machine Learning Algorithms.” *Scientific Reports* 10(1): 9952. <https://doi.org/10.1038/s41598-020-67024-3>.
- Long, J. N., and J. D. Shaw. 2009. “The Influence of Compositional and Structural Diversity on Forest Productivity.” *Forestry* 83(2): 121–28. <https://doi.org/10.1093/forestry/cpp033>.

- Luoma, V., N. Saarinen, M. A. Wulder, J. C. White, M. Vastaranta, M. Holopainen, and J. Hyypä. 2017. "Assessing Precision in Conventional Field Measurements of Individual Tree Attributes." *Forests* 8(2): 38.
- Marvin, D. C., G. P. Asner, D. E. Knapp, C. B. Anderson, R. E. Martin, F. Sinca, and R. Tupayachi. 2014. "Amazonian Landscapes and the Bias in Field Studies of Forest Structure and Biomass." *Proceedings of the National Academy of Sciences of the United States of America* 111: E5224–E5232.
- Matasci, G., T. Hermosilla, M. A. Wulder, J. C. White, N. C. Coops, G. W. Hobart, D. K. Bolton, P. Tompalski, and C. W. Bator. 2018. "Three Decades of Forest Structural Dynamics over Canada's Forested Ecosystems Using Landsat Time-Series and Lidar Plots." *Remote Sensing of Environment* 216: 697–714.
- McNeil, D. J., G. Fisher, C. J. Fiss, A. J. Elmore, M. C. Fitzpatrick, J. W. Atkins, J. Cohen, and J. L. Larkin. 2023. "Using Aerial LiDAR to Assess Regional Availability of Potential Habitat for a Conservation Dependent Forest Bird." *Forest Ecology and Management* 540: 121002.
- McRoberts, R. E., S. Winter, G. Chirici, E. Hauk, D. R. Pelz, W. K. Moser, and M. A. Hatfield. 2008. "Large-Scale Spatial Patterns of Forest Structural Diversity." *Canadian Journal of Forest Research* 38(3): 429–438. <https://doi.org/10.1139/x07-154>.
- Meng, J., S. Li, W. Wang, Q. Liu, S. Xie, and W. Ma. 2016. "Estimation of Forest Structural Diversity Using the Spectral and Textural Information Derived from SPOT-5 Satellite Images." *Remote Sensing* 8(2): 125.
- Meng, R., J. Wu, F. Zhao, B. D. Cook, R. P. Hanavan, and S. P. Serbin. 2018. "Measuring Short-Term Post-Fire Forest Recovery across a Burn Severity Gradient in a Mixed Pine-Oak Forest Using Multi-Sensor Remote Sensing Techniques." *Remote Sensing of Environment* 210: 282–296.
- Mokroš, M., X. Liang, P. Surový, P. Valent, J. Čerňava, F. Chudý, D. Tunák, Š. Saloň, and J. Merganič. 2018. "Evaluation of Close-Range Photogrammetry Image Collection Methods for Estimating Tree Diameters." *ISPRS International Journal of Geo-Information* 7: 93. <https://doi.org/10.3390/ijgi7030093>.
- Moorthy, S. M. K., K. Calders, M. B. Vicari, and H. Verbeeck. 2019. "Improved Supervised Learning-Based Approach for Leaf and Wood Classification from LiDAR Point Clouds of Forests." *IEEE Transactions on Geoscience and Remote Sensing* 58(5): 3057–70.
- Muller-Landau, H. C. 2009. "Sink in the African Jungle." *Nature* 457: 969–970. <https://doi.org/10.1038/457969a>.
- Narine, L. L., S. C. Popescu, and L. Malambo. 2019. "Synergy of ICESat-2 and Landsat for Mapping Forest Aboveground Biomass with Deep Learning." *Remote Sensing* 11(12): 1503. <https://doi.org/10.3390/rs11121503>.
- NEON (National Ecological Observatory Network). 2022a. "Vegetation Structure, RELEASE-2022 (DP1.10098.001)." <https://doi.org/10.48443/re8n-tm87>.
- NEON (National Ecological Observatory Network). 2022b. "Discrete Return LiDAR Point Cloud, RELEASE-2022 (DP1.30003.001)." <https://doi.org/10.48443/ca22-1n03>.
- NEON (National Ecological Observatory Network). 2023. "Spectrometer Orthorectified at-Sensor Radiance—Flightline (DP1.30008.001), RELEASE-2023." <https://doi.org/10.48443/ehnm-xd88>. Dataset. <https://data.neonscience.org>.
- Newman, E. A. 2019. "Disturbance Ecology in the Anthropocene." *Frontiers in Ecology and Evolution* 7: 147.
- Newman, E. A., M. C. Kennedy, D. A. Falk, and D. McKenzie. 2019. "Scaling and Complexity in Landscape Ecology." *Frontiers in Ecology and Evolution* 7: 293.
- Pádua, L., J. Vanko, J. Hruška, T. Adão, J. J. Sousa, E. Peres, and R. Morais. 2017. "UAS, Sensors, and Data Processing in Agroforestry: A Review towards Practical Applications." *International Journal of Remote Sensing* 38: 2349–91.
- Parker, G. G., and M. E. Russ. 2004. "The Canopy Surface and Stand Development: Assessing Forest Canopy Structure and Complexity with Near-Surface Altimetry." *Forest Ecology and Management* 189(1–3): 307–315.
- Petrie, G. 2013. "Commercial Operation of Lightweight UAVs for Aerial Imaging and Mapping." *GEOInformatics* 16: 28–39.
- Pommerening, A. 2002. "Approaches to Quantifying Forest Structures." *Forestry: An International Journal of Forest Research* 75(3): 305–324.
- Possingham, H. P., J. Franklin, K. Wilson, and T. J. Regan. 2005. "The Roles of Spatial Heterogeneity and Ecological Processes in Conservation Planning." *Ecosystem Function in Heterogeneous Landscapes*: 389–406.
- Potapov, P., X. Li, A. Hernandez-Serna, A. Tyukavina, M. C. Hansen, A. Kommareddy, A. Pickens, et al. 2020. "Mapping Global Forest Canopy Height through Integration of GEDI and Landsat Data." *Remote Sensing of Environment* 253: 112165. <https://doi.org/10.1016/j.rse.2020.112165>.
- Proença, V., L. J. Martin, H. M. Pereira, M. Fernandez, L. McRae, J. Belnap, M. Böhm, et al. 2017. "Global Biodiversity Monitoring: From Data Sources to Essential Biodiversity Variables." *Biological Conservation* 213: 256–263.
- Qi, W., S. Saarela, J. Armston, G. Ståhl, and R. Dubayah. 2019. "Forest Biomass Estimation over Three Distinct Forest Types Using TanDEM-X InSAR Data and Simulated GEDI Lidar Data." *Remote Sensing of Environment* 232: 111283.
- Rajabifard, A., and I. P. Williamson. 2001. "Spatial Data Infrastructures: Concept, SDI Hierarchy and Future Directions."
- Raunonen, P., M. Kaasalainen, M. Åkerblom, S. Kaasalainen, H. Kaartinen, M. Vastaranta, M. Holopainen, M. Disney, and P. Lewis. 2013. "Fast Automatic Precision Tree Models from Terrestrial Laser Scanner Data." *Remote Sensing* 5(2): 491–520.
- Rocchini, D., A. Chiarucci, and S. A. Loiselle. 2004. "Testing the Spectral Variation Hypothesis by Using Satellite Multispectral Images." *Acta Oecologica* 26(2): 117–120.
- Romme, W. H. 2005. "The Importance of Multiscale Spatial Heterogeneity in Wildland Fire Management and Research." In *Ecosystem Function in Heterogeneous Landscapes*, edited by G. M. Lovett, M. G. Turner, C. G. Jones, and K. C. Weathers, 353–366. New York: Springer.
- Saatchi, S., M. Marlier, R. L. Chazdon, D. B. Clark, and A. E. Russell. 2011. "Impact of Spatial Variability of Tropical Forest Structure on Radar Estimation of Aboveground Biomass." *Remote Sensing of Environment* 115(11): 2836–49. <https://doi.org/10.1016/j.rse.2010.07.015>.
- Schmidt, S., and F. E. Fassnacht. 2017. "The Spectral Variability Hypothesis Does Not Hold across Landscapes." *Remote Sensing of Environment* 192: 114–125.
- Schweiger, A. K., J. Cavender-Bares, P. A. Townsend, S. E. Hobbie, M. D. Madritch, R. Wang, D. Tilman, and J. A. Gamon. 2018. "Plant Spectral Diversity Integrates Functional and Phylogenetic

- Components of Biodiversity and Predicts Ecosystem Function.” *Nature Ecology & Evolution* 2(6): 976–982.
- Sheldon, S. L., R. O. Dubayah, D. B. Clark, M. A. Hofton, and J. Blair. 2006. “Analysis of Tropical Forest Vertical and Spatial Structural Dynamics Using Large-footprint Lidar.” AGU Fall Meeting Abstracts, Vol. 2006, B41A-0163.
- Shiklomanov, A. N., B. A. Bradley, K. M. Dahlin, A. M. Fox, C. M. Gough, F. M. Hoffman, E. Middleton, S. P. Serbin, L. Smallman, and W. K. Smith. 2019. “Enhancing Global Change Experiments through Integration of Remote-Sensing Techniques.” *Frontiers in Ecology and the Environment* 17(4): 215–224.
- Shimada, M., T. Itoh, T. Motooka, M. Watanabe, T. Shiraishi, R. Thapa, and R. Lucas. 2014. “New Global Forest/Non-Forest Maps from ALOS PALSAR Data (2007–2010).” *Remote Sensing of Environment* 155: 13–31.
- Shugart, H. H., S. Saatchi, and F. G. Hall. 2010. “Importance of Structure and Its Measurement in Quantifying Function of Forest Ecosystems.” *Journal of Geophysical Research: Biogeochemistry* 115: G00E13. <https://doi.org/10.1029/2009JG000993>.
- Silva, C. A., L. Duncanson, S. Hancock, A. Neuenschwander, N. Thomas, M. Hofton, L. Fatoyinbo, et al. 2021. “Fusing Simulated GEDI, ICESat-2 and NISAR Data for Regional Aboveground Biomass Mapping.” *Remote Sensing of Environment* 253: 112234.
- Silva, C. A., A. T. Hudak, L. A. Vierling, R. Valbuena, A. Cardil, M. Mohan, D. R. de Almeida, et al. 2022. “Treetop: A Shiny-Based Application and R Package for Extracting Forest Information from LiDAR Data for Ecologists and Conservationists.” *Methods in Ecology and Evolution*. 13: 1164–76.
- Silva, C. A., C. Klauber, A. T. Hudak, L. A. Vierling, V. Liesenberg, S. P. E. Carvalho, and L. C. Rodriguez. 2016. “A Principal Component Approach for Predicting the Stem Volume in Eucalyptus Plantations in Brazil Using Airborne LiDAR Data.” *Forestry: An International Journal of Forest Research* 89(4): 422–433.
- Singh, K. K., and A. E. Frazier. 2018. “A Meta-Analysis and Review of Unmanned Aircraft System (UAS) Imagery for Terrestrial Applications.” *International Journal of Remote Sensing* 39(15–16): 5078–98.
- Smart, L. S., J. J. Swenson, N. L. Christensen, and J. O. Sexton. 2012. “Three-Dimensional Characterization of Pine Forest Type and Red-Cockaded Woodpecker Habitat by Small-Footprint, Discrete-Return Lidar.” *Forest Ecology and Management* 281: 100–110. <https://doi.org/10.1016/j.foreco.2012.06.020>.
- Song, C., and C. E. Woodcock. 2002. “The Spatial Manifestation of Forest Succession in Optical Imagery: The Potential of Multiresolution Imagery.” *Remote Sensing of Environment* 82(2–3): 271–284.
- Stark, S. C., V. Leitold, J. L. Wu, M. O. Hunter, C. V. de Castilho, F. R. Costa, S. M. McMahon, et al. 2012. “Amazon Forest Carbon Dynamics Predicted by Profiles of Canopy Leaf Area and Light Environment.” *Ecology Letters* 15(12): 1406–14.
- Stovall, A. E. L., D. Lagomasino, S.-K. Lee, M. Simard, N. M. Thomas, C. Trettin, and T. Fatoyinbo. 2020. “Evaluating Current and Future Sensor-Specific Biomass Calibration in the Tallest Mangrove Forest on Earth.” IGARSS 2020–2020 IEEE International Geoscience and Remote Sensing Symposium, 5964–67.
- Stovall, A. E. L., and H. H. Shugart. 2018. “Improved Biomass Calibration and Validation with Terrestrial LiDAR: Implications for Future LiDAR and SAR Missions.” *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* 11: 3527–37.
- Tanase, M. A., L. Villard, D. Pitar, B. Apostol, M. Petrila, S. Chivulescu, S. Leca, I. Borlaf-Mena, I. Pascu, and A. Dobre. 2019. “Synthetic Aperture Radar Sensitivity to Forest Changes: A Simulations-Based Study for the Romanian Forests.” *Science of the Total Environment* 689: 1104–14.
- Tang, H., J. Armston, S. Hancock, S. Marselis, S. Goetz, and R. Dubayah. 2019. “Characterizing Global Forest Canopy Cover Distribution Using Spaceborne Lidar.” *Remote Sensing of Environment* 231: 111262.
- Tang, L., and G. Shao. 2015. “Drone Remote Sensing for Forestry Research and Practices.” *Journal of Forestry Research* 26: 791–97.
- Tatsumi, S., K. Yamaguchi, and N. Furuya. 2022. “Forest Scanner: A Mobile Application for Measuring and Mapping Trees with Lidar-Equipped iPhone and iPad.” *Methods in Ecology and Evolution* 14(7): 1603–9. <https://doi.org/10.1111/2041-210X.13900>.
- Valbuena, R., B. O’Connor, F. Zellweger, W. Simonson, P. Vihervaara, M. Maltamo, C. A. Silva, et al. 2020. “Standardizing Ecosystem Morphological Traits from 3D Information Sources.” *Trends in Ecology & Evolution* 35(8): 656–667.
- Valbuena, R., J. Vauhkonen, P. Packalen, J. Pitkänen, and M. Maltamo. 2014. “Comparison of Airborne Laser Scanning Methods for Estimating Forest Structure Indicators Based on Lorenz Curves.” *ISPRS Journal of Photogrammetry and Remote Sensing* 95: 23–33. <https://doi.org/10.1016/j.isprsjprs.2014.06.002>.
- Van Mantgem, P. J., N. L. Stephenson, J. Byrne, L. D. Daniels, J. F. Franklin, P. Z. Fulé, M. E. Harmon, et al. 2009. “Widespread Increase of Tree Mortality Rates in the Western US.” *Science* 323: 521–24.
- Verbeeck, H., M. Bauters, T. Jackson, A. Shenkin, M. Disney, and K. Calders. 2019. “Time for a Plant Structural Economics Spectrum.” *Frontiers in Forests and Global Change* 2: 43.
- Vihervaara, P., L. Mononen, A. P. Auvinen, R. Virkkala, Y. Lü, I. Pippuri, P. Packalen, R. Valbuena, and J. Valkama. 2015. “How to Integrate Remotely Sensed Data and Biodiversity for Ecosystem Assessments at Landscape Scale.” *Landscape Ecology* 30(3): 501–516.
- Wales, S. B., M. R. Kreider, J. Atkins, C. M. Hulshof, R. T. Fahey, L. E. Nave, K. J. Nadelhoffer, and C. M. Gough. 2020. “Stand Age, Disturbance History and the Temporal Stability of Forest Production.” *Forest Ecology and Management* 460: 117865.
- Walter, J. A., A. E. Stovall, and J. W. Atkins. 2021. “Vegetation Structural Complexity and Biodiversity in the Great Smoky Mountains.” *Ecosphere* 12: e03390. <https://doi.org/10.1002/ecs2.3390>.
- Wang, C. J., A. Elmore, I. Numata, M. Cochrane, S. Lei, C. Hakkenberg, Y. Li, Y. Zhao, and Y. Tian. 2022. “A Framework for Improving Wall-to-Wall Canopy Height Mapping by Integrating GEDI LiDAR.” *Remote Sensing* 14(15): 3618.
- Wang, L., P. Gong, and G. S. Biging. 2004. “Individual Tree-Crown Delineation and Treetop Detection in High-Spatial-Resolution Aerial Imagery.” *Photogrammetric Engineering & Remote Sensing* 70(3): 351–57.

- Whitehurst, A. S., A. Swatantran, J. B. Blair, M. A. Hofton, and R. Dubayah. 2013. "Characterization of Canopy Layering in Forested Ecosystems Using Full Waveform Lidar." *Remote Sensing* 5(4): 2014–36.
- Wiens, J. A. 1974. "Habitat Heterogeneity and Avian Community Structure in North American Grasslands." *American Midland Naturalist* 91: 195–213.
- Wilkes, P., M. Disney, M. B. Vicari, K. Calders, and A. Burt. 2018. "Estimating Urban Above Ground Biomass with Multi-Scale LiDAR." *Carbon Balance and Management* 13: 10. <https://doi.org/10.1186/s13021-018-0098-0>.
- Wilkes, P., A. Lau, M. Disney, K. Calders, A. Burt, J. Gonzalez de Tanago, H. Bartholomeus, B. Brede, and M. Herold. 2017. "Data Acquisition Considerations for Terrestrial Laser Scanning of Forest Plots." *Remote Sensing of Environment* 196: 140–153. <https://doi.org/10.1016/j.rse.2017.04.030>.
- Wulder, M. A., R. J. Hall, N. C. Coops, and S. E. Franklin. 2004. "High Spatial Resolution Remotely Sensed Data for Ecosystem Characterization." *BioScience* 54(6): 511–521.
- Wulder, M. A., J. G. Masek, W. B. Cohen, T. R. Loveland, and C. E. Woodcock. 2012. "Opening the Archive: How Free Data Has Enabled the Science and Monitoring Promise of Landsat." *Remote Sensing of Environment* 122: 2–10. <https://doi.org/10.1016/j.rse.2012.01.010>.
- Xue, J., and B. Su. 2017. "Significant Remote Sensing Vegetation Indices: A Review of Developments and Applications." *Journal of Sensors* 2017: 1–17. <https://doi.org/10.1155/2017/1353691>.
- Yan, F., B. Wu, and Y. Wang. 2015. "Estimating Spatiotemporal Patterns of Aboveground Biomass Using Landsat TM and MODIS Images in the Mu Us Sandy Land, China." *Agricultural and Forest Meteorology* 200: 119–128.
- Zellweger, F., P. De Frenne, J. Lenoir, P. Vangansbeke, K. Verheyen, M. Bernhardt-Römermann, L. Baeten, et al. 2020. "Forest Microclimate Dynamics Drive Plant Responses to Warming." *Science* 368(6492): 772–75. <https://doi.org/10.1126/science.aba6880>.
- Zeng, Y., D. Hao, A. Huete, B. Dechant, J. Berry, J. M. Chen, J. Joiner, et al. 2022. "Optical Vegetation Indices for Monitoring Terrestrial Ecosystems Globally." *Nature Reviews Earth & Environment* 3(7): 477–493. <https://doi.org/10.1038/s43017-022-00298-5>.
- Zenner, E. K., Y. L. Dickinson, and J. E. Peck. 2012. "Recovery of Forest Structure and Composition to Harvesting in Different Strata of Mixed Even-Aged Central Appalachian Hardwoods." *Annals of Forest Science* 70(2): 151–59. <https://doi.org/10.1007/s13595-012-0242-z>.
- Zhai, L., D. R. Coyle, D. Li, and A. Jonko. 2022. "Fire, Insect and Disease-Caused Tree Mortalities Increased in Forests of Greater Structural Diversity during Drought." *Journal of Ecology* 110(3): 673–685. <https://doi.org/10.1111/1365-2745.13830>.

How to cite this article: Atkins, Jeff W., Parth Bhatt, Luis Carrasco, Emily Francis, James E. Garabedian, Christopher R. Hakkenberg, Brady S. Hardiman, et al. 2023. "Integrating Forest Structural Diversity Measurement into Ecological Research." *Ecosphere* 14(9): e4633. <https://doi.org/10.1002/ecs2.4633>