

RESEARCH ARTICLE

Declining tree growth resilience mediates subsequent forest mortality in the US Mountain West

Antoine Cabon^{1,2}  | R. Justin DeRose³ | John D. Shaw⁴ | William R. L. Anderegg^{1,2} 

¹Wilkes Center for Climate Science and Policy, University of Utah, Salt Lake City, Utah, USA

²School of Biological Sciences, University of Utah, Salt Lake City, Utah, USA

³Department of Wildland Resources and Ecology Center, Utah State University, Logan, Utah, USA

⁴USDA Forest Service, Rocky Mountain Research Station, Logan, Utah, USA

Correspondence

Antoine Cabon, Eidgenössische Forschungsanstalt WSL, Zürcherstrasse 111, 8903 Birmensdorf, Switzerland.
Email: antoine.cabon@wsl.ch

Funding information

Alan T. Waterman Award, Grant/Award Number: IOS-2325700; David and Lucille Packard Foundation; National Science Foundation, Grant/Award Number: 1714972, 1802880, 2044937 and 2003017; USDA National Institute of Food and Agriculture, Agricultural and Food Research Initiative Competitive Programme, Ecosystem Services and Agro-Ecosystem Management, Grant/Award Number: 2018-67019-27850

Abstract

Climate change-triggered forest die-off is an increasing threat to global forests and carbon sequestration but remains extremely challenging to predict. Tree growth resilience metrics have been proposed as measurable proxies of tree susceptibility to mortality. However, it remains unclear whether tree growth resilience can improve predictions of stand-level mortality. Here, we use an extensive tree-ring dataset collected at ~3000 permanent forest inventory plots, spanning 13 dominant species across the US Mountain West, where forests have experienced strong drought and extensive die-off has been observed in the past two decades, to test the hypothesis that tree growth resilience to drought can explain and improve predictions of observed stand-level mortality. We found substantial increases in growth variability and temporal autocorrelation as well declining drought resistance and resilience for a number of species over the second half of the 20th century. Declining resilience and low tree growth were strongly associated with cross- and within-species patterns of mortality. Resilience metrics had similar explicative power compared to climate and stand structure, but the covariance structure among predictors implied that the effect of tree resilience on mortality could partially be explained by stand and climate variables. We conclude that tree growth resilience offers highly valuable insights on tree physiology by integrating the effect of stressors on forest mortality but may have only moderate potential to improve large-scale projections of forest die-off under climate change.

KEYWORDS

drought, forest mortality, growth, insect outbreak, Lloret indices, resilience, tree-ring

1 | INTRODUCTION

Rising global temperatures due to human greenhouse gas emissions pose a threat to Earth's forests. Notably, increasing temperature has been linked to higher frequency and intensity of drought-induced forest background mortality and die-off (Allen et al., 2015; Breshears et al., 2009; Williams et al., 2012). Large-scale forest die-off is of particular concern due to its dramatic disruption of forest function, with large consequences on biodiversity (Betts et al., 2017; Feng et al., 2021), ecosystem goods and services,

and forests' prominent role in the global carbon cycle (Anderegg, Trugman, Badgley, Anderson, et al., 2020). The western US has experienced extensive drought- and insect-induced tree mortality over the past decades, associated with strong drying trends in the area (Van Mantgem et al., 2009; Williams et al., 2012; Zhang et al., 2021). These disturbances have resulted in the decline of several major tree species in the region (Stanke et al., 2021).

A large diversity of interacting drivers and pathways mediate climate-induced tree mortality. Drought-induced mortality results from a complex and interacting set of failures across the hydraulic

continuum and tree carbon economy (McDowell et al., 2022), which has been typically studied as hydraulic failure and depletion of carbohydrate stores (McDowell, 2011). Hydraulic failure, that is, the accumulation of emboli in the xylem past a threshold after which water transport cannot be recovered, disrupts water supply, leading to cell death by dehydration (Sperry & Tyree, 1988). Longer term, declines in carbon balance can eventually trigger mortality through cell failure to maintain base metabolism or osmoregulation, almost always interacting with hydraulic failure and/or biotic agents (Sevanto et al., 2014). Tree radial growth is a major process integrating water and carbon processes involved in mortality (Preisler et al., 2021). Previous to drought, tree growth favors carbon assimilation by sustaining photosynthesis and tree development. On the other hand, larger allocation to conducting tissues can be at the expense of allocation to osmoregulation or defense (de la Mata et al., 2017; Huot et al., 2014) or translate into structural overshoot rendering trees more vulnerable to drought (Jump et al., 2017). Following drought, new xylem growth allows trees to recover hydraulic function and may influence delayed mortality (Trugman et al., 2018).

Biotic agents, including diseases and insects can also drive large-scale forest mortality, often in interaction with climate stress, which challenges disentangling their relative importance (Anderegg, Hicke, et al., 2015). Notably, western US forests have been particularly prone to extensive bark beetle outbreaks in the last decades, where drought was a key predisposing factor in many species (Chapman et al., 2012; Meddens et al., 2015; Raffa et al., 2008; Van Mantgem et al., 2009). Drought may facilitate biotic outbreaks by impairing tree defenses against attacks such as resin and defense chemical production, hence favoring infestations and eventually mortality (Gaylord et al., 2013; Kolb et al., 2019). In turn, infestations can weaken trees' water and carbon economies (Frank et al., 2014) through direct damage to the bark and xylem or defoliation, which may increase tree vulnerability to drought (Anderegg & Callaway, 2012; Anderegg, Hicke, et al., 2015; Paine et al., 1997). Finally, stand factors such as host availability, density, age, and size can play a substantial role in determining whether infestations reach epidemic levels and eventually lead to extensive die-off events (Raffa et al., 2008). However, predisposing factors are often highly species specific (Reed & Hood, 2021).

Climate-induced tree mortality remains challenging to predict because of the wide array of involved processes. Success of physiology-based models has been largely limited to specific contexts and/or small scales, where environmental variations and the number of processes are limited, in contrast to regional applications (e.g., Venturas et al., 2018, 2021; see also Benito Garzón et al., 2018; Trugman et al., 2021). The challenge of accounting for such complexity has led to the adoption of simpler frameworks based on tree vigor proxies (Bigler & Bugmann, 2003; Lloret et al., 2011). Growth-based models of mortality are one such approach, wherein, based on the empirical assertion that tree growth declines previous to tree mortality (Cailleret et al., 2016), tree growth is assumed to integrate constraints on tree physiological

status. Growth-based mortality models use historical tree growth observations, for example, derived from tree ring or forest inventory, to parameterize empirical models of tree mortality, often in combination with cofactors such as tree diameter (Bigler & Bugmann, 2003; Hülsmann et al., 2018). These models have garnered relatively good success due to their predictive power and simplicity but might be limited in that they assume a static relationship between tree growth and mortality. Such assumption may allow one to simulate smooth variations in stand background mortality, but it is unclear whether tree growth can capture the effect of drought and biotic agent perturbations in a changing climate, as such events can be uncoupled from it (e.g., epidemic insect outbreak) or instead select for slower growing trees (de la Mata et al., 2017; Jump et al., 2017).

By acknowledging highly nonlinear responses, complex system theory may provide a useful framework to predict stand-level mortality. In this framework a system approaching a tipping point is denoted by critical slowing down that manifests in declining system resilience to perturbations, that is, the system's capacity to remain in a reference regime (Scheffer et al., 2001, 2009). Declining resilience and associated increasing state variability and autocorrelation, can hence theoretically be used as early warning signals (EWS) of impending system transition to an alternative stable state (Hammond, 2020; Scheffer et al., 2009). Stand or regional-scale forest mortality can be viewed as one such tipping point, where a perturbation might induce an abrupt transition toward a fundamentally different regime (e.g., change in stand structure, species composition, or transition to a different cover type) (Hammond, 2020). Consequently, a body of literature has focused on applying critical slowing down theory to forest mortality by developing EWS based on radial tree growth (Camarero et al., 2015; Lloret et al., 2011). Recent syntheses notably show that large tree growth variability—in the case of gymnosperms—and low resilience (Cailleret et al., 2019; DeSoto et al., 2020) are associated with subsequent tree-level mortality, suggesting that these could predict future forest mortality. Implications are vast given observation of widespread declining tree resilience to drought (Forzieri et al., 2022; Zheng et al., 2021, 2023). Several uncertainties nevertheless remain that hinder the application of EWS to predict forest mortality. First and foremost, previous studies mostly investigated the relationship between EWS and mortality at the tree level (i.e., differences among individual trees growing in the same stand), but it is unclear if such a relationship scales to the stand or landscape level. At large spatial scales, the role of covarying environmental and genetic factors (e.g., stand structure, climate, species identity) might complicate or overshadow the relationship between growth resilience and mortality observed at the individual level (Kannenberget al., 2019). Second, previous analyses typically reported qualitatively on the existence of such relationship but lacked quantification of their predictive power. Finally, because EWS metrics have often been investigated individually, it is not clear how different metrics, including tree growth, variability, autocorrelation, and resilience might be complementary to or on the contrary be redundant with stand and climate predictors of mortality.

Here, in order to address the link between tree growth resilience to drought and stand-level mortality, we leverage a large tree-ring dataset comprised of over 7000 records from ~3000 national forest inventory plots in the US Mountain West, where extensive mortality has been reported in recent decades. Specifically, we (1) test the hypothesis that low tree growth resilience and associated metrics are related to subsequent stand-level mortality, (2) investigate the regional stand and climatic drivers of mortality and their covariation with tree growth and resilience, and finally (3) quantify the relative power of stand, climate, growth, and resilience drivers to capture spatial patterns of stand mortality in explicative versus predictive modeling contexts.

2 | MATERIALS AND METHODS

2.1 | Forest inventory, stand structure, and mortality

The USDA Forest Service Forest Inventory and Analysis (FIA) program monitors forest structure, growth, and health in a systematic way based on an extensive network of permanent plots distributed across the United States (Bechtold & Scott, 2005). Here we use a subset of 3028 FIA plots from five states of the US Mountain West where tree cores were sampled in addition to the FIA standard sampling effort (DeRose et al., 2017). Following the FIA sampling design, four circular subplots of a radius of 7.3 m comprise each plot, where the diameter at breast height (DBH) of all trees with a diameter superior to 12.7 cm are measured. Each subplot contains a circular microplot of 2.1 m, within which trees with a diameter inferior to 12.7 cm are measured. Plots included in this study were selected on the basis that no silvicultural treatment was applied, and that mortality was attributed either to no agent (unknown agent), insect, drought (weather), or disease.

Trees are classified into live, recently dead, and older dead based on canopy status. On repeated inventory plots (~35% of all plots), "recent dead" is assigned to the trees that were live during the previous inventory and dead in current inventory. On first-visit plots, "recent dead" is assigned to trees that appear to have died during the previous 5 years, assessed by the census crew based on canopy and bark status (Shaw et al., 2005). This initial-visit approach has been commonly and successfully used in tree mortality research in this region (e.g., Venturas et al., 2021). "Old dead" trees are ignored in the following analysis and "recently dead" trees are referred to as "dead" hereafter.

We computed species-specific per hectare sum of basal area of individual trees in live and dead categories (BA). Mortality was subsequently defined at the stand level as the BA of dead in percent of total BA (live and dead). Tree density was calculated as the sum of live and dead stems per hectare. Tree species diversity was estimated based on the Shannon index where species abundance is taken as species total basal area.

2.2 | Climate data

Several climatic variables have been previously shown to be closely related to tree mortality across the United States (Venturas et al., 2021). Monthly minimum and maximum temperature, precipitation, Palmer's drought severity index (PDSI), vapor pressure deficit, and climatic water deficit were retrieved from the TerraClimate database (Abatzoglou et al., 2018) for each of the plots on the period 1958–2019 (TerraClimate variable names: tmin, tmax, ppt, PDSI, vpd, and def, respectively). Mean annual temperature (MAT, computed as the average of min and max temperature), precipitation (MAP), vapor pressure deficit, and climatic water deficit were calculated in the same period. We further calculated plot-level historical temporal trends of these climatic variables by means of linear regression. Drought intensity, frequency, and heat stress during the FIA census period were calculated at each plot, respectively, as the minimum growing season PDSI (averaged over April through September), the number of years with growing season PDSI < -1.5, and the difference between maximum and average growing season max temperature.

2.3 | FIA tree-ring collection and growth metrics

Generally, one (about 75% of the plots) but up to 16 and an average of 2.2 tree cores per plot were collected (~1.6 cores per plot and per species), yielding a total of 7281 cores. Tree cores were initially collected to determine stand age and site index, based on the average age per species within 12.7 cm diameter classes and the dominant size class, respectively. As a result, tree selection consisted exclusively of live trees and was typically skewed toward the dominant size cohort. Cores were collected during the period 2000–2022, but over 90% of the cores were collected after 2010. The cores were processed, and ring width was measured following standard dendrochronological methodology (DeRose et al., 2017). Cross-dating was performed using nearby chronologies from the International Tree-Ring Data Bank (ITRDB), as well as adjacent FIA plots as the tree-ring collection grew. Past tree basal area increment (BAI) was retrieved backward from tree DBH at core collection. A detrended ring width index (RWI) was calculated by dividing RW series by splines with 50% cutoff at 30 years fitted on individual ring width series. In order to investigate the potential of past tree growth temporal patterns as early warning metrics of mortality, BAI and RWI series were used to calculate a set of six metrics to characterize average and trends in past tree growth and resilience to drought events, during the 40 years prior to the mortality census period.

Growth metrics included BAI, as well as RWI autocorrelation (1 year lag; ar) and RWI variation coefficient (VC) calculated on 10-year windows, which have been previously proposed as early warning metrics of tree mortality (Cailleret et al., 2019). Resilience metrics included tree growth resilience (RI) to past droughts and its resistance (R_s) and recovery (R_c) components. Resilience metrics

were calculated for each RWI series and drought years. Drought years were defined as years for which growing season PDSI (averaged over April–September) was inferior to the first decile at a given plot. RI, Rs and Rc were calculated as differences, analogous to Lloret's ratios (Lloret et al., 2011):

$$Rs = RWI_{\text{drought}} - RWI_{\text{pre}} \quad (1)$$

$$Rc = RWI_{\text{post}} - RWI_{\text{drought}} \quad (2)$$

$$RI = RWI_{\text{post}} - RWI_{\text{pre}} = Rs + Rc \quad (3)$$

where RWI_{drought} is RWI during the drought year, RWI_{pre} is the average RWI during the 4 years prior, and RWI_{post} is the average RWI during the 4 years posterior to the drought year, following (Anderegg, Schwalm, et al., 2015). The use of differences instead of ratios was motivated by the distribution of RWI, which included zeroes and values close to zero.

In order to investigate tree-ring metric temporal variations and their effect on tree mortality, all six growth and resilience metrics were decomposed at the core level into a temporal trend and a period-independent average. To do so we fitted two sets of mixed linear models. First, each metric was modeled as a function of species identity, including one random intercept per core as well as a random effect of year per species, such that:

$$Y_{ijk} = \beta_{0,i} + u_{0,j} + u_{1,ik} + \epsilon_{ijk} \quad (4)$$

where Y_{ijk} is the metric value of the i th species, j th core, and k th year, $\beta_{0,i}$ is a fixed species intercept, $u_{0,j}$ is a random intercept per core, $u_{1,ik}$ is the random effect of year per species, and ϵ is an error term. Based on this model we estimated period-independent mean values per core for each metric (hereafter metric name plus "mean") such as $\beta_{0,i} + u_{0,j}$. Second, we fitted a mixed linear model which included fixed species and year per species effects, as well as a random intercept and slope per core, nested within species:

$$Y_{ij} = \beta_{0,i} + \beta_{1,i} \text{Year} + u_{0,j} + u_{1,j} \text{Year} + \epsilon_{ij} \quad (5)$$

where, $\beta_{1,i}$ is the slope for the fixed effect of year. From this model we extracted core-level linear temporal trends for each metric (hereafter metric name plus "trend") such as $\beta_{1,i} + u_{1,j}$. When relevant, previous to model fitting we normalized leptokurtic-distributed metrics using the Lambert W function in the R package "LambertW." Homoskedasticity and normality of residuals were visually checked for each model and metric.

2.4 | Stand mortality models

We considered the role of a total of ~40 unique variables as predictors of stand mortality, that were classified into one of stand, climate, growth, or resilience categories. Stand structure predictors included overall and species-specific basal area, tree density, average and

maximum height, and DBH, as well as stand age, aspect, slope, and diversity (Shannon index). Climate variables included MAT, MAP, mean annual vapor pressure and climatic water deficit, temporal trends of the previous, as well as drought intensity, frequency, and heat stress during the census period. Growth variables consisted in the overall mean and temporal trends of BAI, BAI autocorrelation, and BAI variation coefficient. Resilience variables consisted in overall mean and temporal trends of RI, Rs, and Rc.

In order to perform a balanced comparison between the effects of stand, climate, growth, and resilience factors on stand mortality and to limit the total number of model parameters to avoid overfitting, we performed a preliminary model selection to restrict the number of predictors per category to four. To do so, for each predictor category (i.e., stand, climate, growth, and resilience), we fitted logistic models of stand-level mortality with all combinations of up to four predictors per category. For each category we then identified the best model based on the Akaike information criterion (AIC) and retained the corresponding set of predictors. When the AIC difference between models was <2 , we retained the most parsimonious and biologically meaningful model (Tables S1–S4). The covariance structure of selected predictors was assessed by calculating pairwise Pearson's correlation between each predictor.

Species-specific basal area mortality occurrence, intensity (i.e., proportion of conspecific mortality basal area at plots where mortality occurred), and overall basal area mortality (hereafter "mortality") were subsequently modeled both as a function of one predictor at a time (univariate case) and as a function of all predictors at the same time (multivariate case). In all cases, the three components of mortality were modeled using logistic regressions with binomial residual distribution in the case of mortality occurrence and overall mortality, and beta distribution in the case of intensity (generalized linear models; GLMs). FIA census repeat interval was found to have only a small effect on mortality but was nevertheless included in all models as a covariable. Species-specific tree density was similarly found to induce a positive bias on mortality occurrence detection (increased likeliness to observe at least one mortality tree with large sample size; Figure S1a) as well as a negative bias on mortality intensity (increased likeliness of mortality to make up for a large proportion of species BA with low sample size; Figure S1b), but these biases canceled out in the case of overall mortality (Figure S1c). Hence, we included species-specific density as a second covariable in mortality occurrence and intensity models. GLMs had the form:

$$\begin{aligned} \text{logit}(M) = & \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \dots + \beta_n X_n \\ & + \beta_{n+1} \text{CensusPeriod} + (\beta_{n+2} \text{SpeciesDensity}) + \epsilon \end{aligned} \quad (6)$$

where M is mortality occurrence, intensity or overall mortality β_0 is the intercept, $\beta_1, \beta_2, \dots, \beta_n$ are the coefficient estimates for the included $X_1, X_2, \dots, X_n$ variables, β_{n+1} is the effect of repeat census period, β_{n+2} is the effect of species density (only for mortality occurrence and intensity models), and ϵ is an error term.

In the univariate case we used generalized linear mixed models (GLMMs) that had the same general form as Equation 5 but further

included a random intercept per species, in order to account for different mortality pathways between species and evaluate the predictive power of each variable *within* in addition to *across* species. Finally, multivariate models' variance was decomposed into stand, climate, growth, and resilience contributions based on the Lindeman, Merenda, and Gold (LMG) and proportional marginal variance decomposition (PMVD) metrics (Grömping, 2007). Both metrics are similar in that they fully decompose total model variance into non-negative shares between predictors irrespective of their order in the model. However, by attributing equal weights to all variables, LMG informs on the explicative power of predictors ("marginal" perspective), whereas PMVD provides an estimate of variable usefulness in a predictive modeling setting by giving larger weights to variables that capture a larger proportion of the variance in combination with a smaller number of variables ("conditional" perspective). Last, we calculated the unique variance carried by each predictor category as the additional amount of explained variance by the full model compared to the model without this predictor category.

3 | RESULTS

3.1 | Mortality spatial distribution

About 1200 of the ~3000 FIA plots considered here exhibited some degree of mortality. In these plots, mortality affected $25\% \pm 1\%$ (mean $\pm$ SE) of basal area on average (6%, 16%, and 38% for first, second, and third quartiles). In the field, ~31% of observed basal area mortality was associated with insect outbreaks, 12% were associated with diseases, and only 1% with drought, due to the limited field evidence and use of this disturbance code by crews. The remaining 56% of all mortality was listed as "no disturbance" but can most likely be attributed to drought (see Section 4). Stand mortality varied widely spatially and across species (Figure 1). Dry and warm, low elevation, pinyon-juniper dominated areas exhibited the lowest mortality levels, whereas colder and wetter subalpine areas exhibited extensive mortality, in relation with insect outbreaks, such as mountain pine beetles and spruce bark beetles. Subalpine pines, spruces, and firs (*Abies lasiocarpa*, *Picea engelmannii*, *Pinus albicaulis*, *P. contorta*, and *P. flexilis*) hence experienced mortality of ~20% basal area on average, of which 46% were associated with insect disturbance. Mortality averaged only 5% for the remaining conifer species, 20% of which were associated with insects. Quaking aspen (*Populus tremuloides*), which stands out as the only angiosperm species in our dataset, also experienced substantial mortality (16% on average). *P. tremuloides* was further unlike other species in the study in that its mortality agent was largely associated with diseases (38%) induced by pathogens such as *Cytospora* canker (Marchetti et al., 2011), in addition to insects (35%).

3.2 | Tree growth resilience decline

We observed that tree-level BAI of all species significantly increased between 1958 and 2010 (Figure 2a). Tree ring-based BAI estimates

were found to be representative of FIA plot-level BAI on the same period ($r = .68$, $p < .0001$). We found that this temporal increase in BAI was matched for most species by an increase in growth variability (11 of 13 species with a significantly positive trend) and, to a lower extent, autocorrelation (5 of 13 species with a significantly positive trend). We further observed a general tendency of growth drought resilience to decline over time (Figure 2b; overall mean $\pm$ SE: $-1.40 \pm 0.52 \times 10^{-3} \text{ year}^{-1}$, $p < .05$), with 7 of 13 species that exhibited a significant decline, and only one species with a significant increase (*Juniperus scopulorum*), although this last result should be interpreted cautiously due to lower sample size for *J. scopulorum*. The resilience decline appeared to be driven overall by a decrease in drought resistance (on average $-1.26 \pm 0.42 \times 10^{-3} \text{ year}^{-1}$, $p < .05$), whereas trends in drought recovery were inconsistent across species and overall did not compensate for decreased resistance ($p = .84$).

3.3 | Mortality predictor selection

Preliminary category-wise variable selection yielded three to four best predictors of forest mortality per predictor category. The best stand structure predictors were conspecific and other species basal area, species maximum tree height and overall maximum DBH. Selected climate predictors were MAT, MAP, drought intensity, and heat stress. Growth predictors comprised mean BAI, mean, and temporal trend of BAI variation coefficient and mean BAI autocorrelation. Resilience predictors were mean RI, Rs trend, and Rt trend. RI trend was not selected, likely due to its high correlation to Rs and Rt trends but was nevertheless included in univariate analyses for comparison.

3.4 | Univariate models of stand mortality

When considering the relationship between individual predictors and stand mortality *across* species (generalized linear models, black lines in Figure 3), we found a significant effect of all predictors, except RI mean and max DBH. Best individual predictors of stand mortality were MAT, RI trend, heat stress, and species max height, in that order. We observed a negative relationship between MAT or drought frequency and mortality, which was consistent with observed geographic distribution of mortality that was strongly skewed toward high mountain areas of Colorado and Wyoming (Figure 1). However, the positive effect of heat stress during the census interval, indicated that given MAT, positive summer temperature anomalies were associated with higher mortality. Finally, rather than absolute tree growth resilience to drought, its decline over time was most strongly associated with mortality across species. The association between mortality and resilience decline across species appeared to be carried mostly by its resistance component.

When looking at *within*-species patterns (colored lines in Figure 3, GLMMs) we found similar effects of MAT, heat stress, max species height, and RI trend compared to *cross*-species level. The effect of




FIGURE 1 (a) Geographic and (b) climatic distribution of stand-level mortality, in percent of conspecific basal area, at sampled forest inventory plots. (c) Number of cores collected per species. The bars color indicates species-level mortality average. (d) Average mortality per species and reported proximal disturbance agent based on field evidence of disease, insect, or drought mortality. The absence of observable dominating disturbance agent is reported as “no disturbance.” Mortality is calculated for each core as a plot-level conspecific basal area proportion. [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/terms-and-conditions)]

some variables which was significant cross-species was found to fade or disappear within species, showing that some or most of the cross-species effect was driven by between-species variations. This was the case of mean growth variation coefficient and growth autocorrelation, and to some extent, that of other species BA and MAP as well. By contrast, we found a stronger effect of RI mean on mortality at the within-species level, which indicated that species-specific differences in resilience blurred the cross-species pattern. Individual predictors of mortality intensity were somewhat similar to that of overall mortality (Figures S2 and S3), whereas the drivers of mortality occurrence more largely differed. Most strikingly, mortality intensity was most strongly related to resilience and growth metrics, whereas mortality occurrence was best described by stand structure metrics.

3.5 | Covariation of stand mortality drivers

Substantial covariation was observed between the predictors of mortality considered here (Figure 4). Notably MAT and MAP were strongly correlated with each other and with a range of other variables, including species max height, mean growth variation coefficient and autocorrelation, as well as to a lesser extent, trends in resistance and recovery. Namely, colder- and wetter-than average climates, were associated with lower tree growth variation coefficient and higher autocorrelation, as well as stronger declines in drought resistance and recovery.

Trees that experienced increasing resistance and recovery to drought also tended to exhibit higher resilience overall. We found a negative association between R_s and R_c trends, suggesting that at




FIGURE 2 Temporal trend of (a) tree growth and (b) resilience metrics. From top to bottom and left to right, basal area increment (BAI), RWI autocorrelation (ar), RWI variation coefficient (VC), growth resilience (RI), resistance (Rs), and recovery (Rc). Temporal trends were estimated from mixed linear models including a random intercept and slope per core. Variables were transformed to approximate a gaussian distribution when necessary. Solid and transparent dots represent significant and insignificant species-level temporal trends, respectively ($\alpha = .05$). [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/terms-and-conditions)]

the tree level, decline in growth resistance to drought was partially mitigated by increased recovery. We did not find that fast-growing trees were more resilient to drought. However, resistance and recovery trends were positively associated with mean growth variability, which was in turn negatively correlated with growth autocorrelation.

3.6 | Multivariate stand mortality models

Full models (including all 15 predictors and one to two covariables—census return time and species density when applicable) explained 40%, 20%, and 24% of observed mortality occurrence, intensity, and overall mortality, respectively (Figure 5a). Variance decomposition

showed that the weight of the different predictors strongly differed between models. Mortality occurrence appeared to be mostly driven by stand structure, followed by climate and denoted little relative importance of growth and resilience metrics. Mortality intensity displayed an opposite pattern, where growth and resilience metrics had the largest relative importance overall, followed by climate and a small contribution of stand structure. These effects balanced each other when considering overall mortality, leading to a roughly similar relative importance of the different predictor categories. Variance decomposition nevertheless gave somewhat different results depending on the perspective. Notably, the marginal perspective (LMG) attributed substantially higher importance to resilience and lower importance to stand and climate overall, compared to the




FIGURE 3 Univariate logistic models of stand mortality (in percent of conspecific basal area). Both generalized (thick black lines) and mixed linear models including a random species intercept (thin colored lines) were fitted to the data. All models include the effect of census period duration, but its contribution to model variance is little (<1%). Predictors are transformed to approximate a normal distribution. Corresponding model marginal or conditional R^2 (GLMMs: R_m^2 , R_c^2) and R^2 (GLMs) and significance (* $p < .05$; ** $p < .01$; *** $p < .001$) are indicated on the top left and right corner of each panel, respectively. Significant effect of x-axis variable is further denoted by solid lines ($p < .05$). Note that RI trend is included here to illustrate its effect on mortality but has been discarded from multivariate analyses in favor of its resistance and recovery components. [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/terms-and-conditions)]

conditional perspective (PMVD). This result was related to the fact that predictors shared a substantial amount of variance, depending on the model. Predictor unique variance in the mortality occurrence model averaged about 68% of overall predictor explained variance (unique/marginal variance), but that number fell under half for intensity and overall mortality models.

Coefficient estimates showed substantial consistency across occurrence, intensity, and overall mortality models (Figure 5b). This was mostly true for climate, growth, and resilience coefficients, although less so in the case of stand coefficients. Namely, we found a strong positive effect of conspecific BA and max height on occurrence and overall mortality, but all stand coefficients were nonsignificant in the




FIGURE 4 Pearson's correlation matrix of the selected mortality predictors. Correlations which absolute value is $>.25$ ($p < .0001$) are displayed on top of the corresponding cells. Outlined cells denote correlations between variables of a same group: that is, stand (green), climate (blue), tree growth (brown), and resilience variables (yellow). [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1111/gcb.16826)]

intensity model. Max DBH had no effect on all models. In contrast, we found consistently strong negative effect of resistance and recovery trends, as well as a positive effect of mean resilience on the three components of mortality. We also found a consistent negative effect of mean BAI across models, but other growth variables had small and inconsistent effect. Last, the effect of all climate variables was found to be positive across models, except MAT which was insignificant in the case of overall mortality.

4 | DISCUSSION

4.1 | Drought and biotic imprint on widespread Mountain West stand mortality

We report widespread elevated mortality rates, though highly variable, across many taxa and five states of the US Mountain West in the past two decades, consistent with previous reports in the region (McNellis et al., 2021; Stanke et al., 2021). Mortality was most pronounced in species found in high mountain regions of the Rockies, in Colorado and Wyoming. This clear distribution toward cool and moist forests contrasts with previous large-scale pinyon pine die-off in warm and dry lowlands of the region during the 2000s, where climate change type drought had been identified as a primary triggering factor (Breshears et al., 2005; Greenwood & Weisberg, 2008; Shaw et al., 2005). Pathogens and insects—notably bark beetles—have been found to be a primary disturbance for a third of mortality reported here (Figure 1d), while about half of the time no

disturbance was reported. Our results nevertheless suggest a strong contribution of drought and heat to observed mortality (Figures 3 and 5). Attribution of mortality to drought by the crews in the field is notoriously challenging because of the lack of clear diagnosis elements (Anderegg, Hicke, et al., 2015), but a broad body of literature in this region reveals that most of observed mortality is primarily driven by drought and with frequent insect interactions (Breshears et al., 2005; Shaw et al., 2005; Stanke et al., 2021; Williams et al., 2012; Worrall et al., 2010). Overall, biotic and climate factors may interact through various processes, and it is often challenging to disentangle the two as mortality drivers. For example, droughted trees are much more likely to suffer and succumb to biotic attacks, for example, because of reduced production of resin and defense compounds against biotic agents (Rissanen et al., 2020; Turtola et al., 2003). On the other hand, insects and pathogens may cause direct damage to trees' vascular system and thus compromise trees' drought resistance (Anderegg, Hicke, et al., 2015). Such interactions make it difficult in many cases to identify the primary agent of mortality and blur the role climate and biotic factors in shaping the spatial distribution of mortality.

Our results revealed that the observed decreased probability of mortality occurrence with temperature was essentially driven by between-species variations, whereas within species, this probability on the contrary tended to increase with temperature (Figure S2). This reflects larger susceptibility of high-elevation species to perturbations. Pronounced increase in overall mortality with colder temperature within species (Figure 3) nevertheless shows that despite lower probability of mortality occurrence, cool versus warm populations of a given species were considerably more




FIGURE 5 Multivariate models of stand mortality occurrence, intensity, and overall mortality ("mortality"). (a) Variance decomposition analysis. Variance is partitioned between four previously defined predictor groups (stand, climate, growth, and resilience variables) based on the LMG (dark colors) and PMVD (light colors) metrics (Grömping, 2007). LMG gives an estimate of each predictor contribution to explain observed variance, whereas PMVD is a metric of the usefulness of each variable for predictive purposes. Marginal and conditional R^2 of corresponding models are indicated on the top left of each panel. The dashed line indicates the theoretical metric value if all variables equally participated to model variance (0.25). (b) Model coefficient estimates. Coefficients are standardized. Points and intervals indicate mean and standard error. Significant ($p < .05$) coefficients are denoted by filled points. Diamonds indicate consistent coefficient sign and significance across models. Colors indicate variable groups: stand (green), climate (blue), growth (brown), and resilience (yellow). [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/terms-and-conditions)]

susceptible to mortality events reaching high intensity (Figure S3). Multivariate analyses revealed that the negative effect of temperature on was underlain by several covarying variables. Notably, cool forests in the region, which also tend to be wetter and more productive, generally exhibit larger basal area and maximum tree height under colder climates. Tall trees—because of their hydraulic architecture—may be particularly sensitive to drought, which might be exacerbated by important competition for resources, including water, in stands with large basal area (Grote et al., 2016; McDowell & Allen, 2015). On the other hand, lower stand basal area might reflect competition release or enhanced microsite suitability in areas previously affected by mortality, such as lowland pinyon-juniper communities (Greenwood & Weisberg, 2008). Finally, biomass availability and the presence of large trees is critical for insect infestations to develop sufficiently to reach epidemic levels (Raffa

et al., 2008). Consistently with all these potential processes, here we find strong positive effects of both species basal area and max height on overall mortality and occurrence. Surprisingly though, we did not find any effect of stand factors on mortality intensity. This is odd considering that higher drought and insect sensitivity conferred by a given stand structure would be expected to similarly drive mortality occurrence and intensity, but this result might be explained by the contrasting properties entailed by stand structure. For example, large basal area and max tree height may also reflect large competition asymmetry, whereby dominated individuals may be more vulnerable to mortality, while dominant trees, which make up for most of plot basal area, benefit from large resource availability (Pretzsch & Biber, 2010).

Cool forests also tended to experience more relative heat stress (i.e., larger summer temperature anomaly). Heat stress together

with drought intensity were found to be strongly related to mortality, hence suggesting an important role of recent hotter droughts in triggering mortality (Hammond et al., 2022). Positive temperature anomalies also favor bark beetle population development and outbreaks (Raffa et al., 2008), especially at cold species boundaries where warming might allow biotic agents to infest new tree populations (Deutsch et al., 2008; McDowell et al., 2020). Hence, tree species vulnerability, stand history, and current structure, as well as recent climate extremes all help explain why cool forests were found to be more susceptible to tree mortality than their warm counterparts. We further observed higher growth autocorrelation, as well as stronger declines in growth resistance and recovery to drought in cool forests, suggesting deteriorating tree physiology and a role in mediating observed mortality patterns along the temperature gradient.

4.2 | Historic decline of tree growth resilience to drought

Over the study period, we found that BAI of sampled trees increased over time, and this held for all species. Because increment core sampling was biased toward dominant crowns within diameter classes, this result is likely inflated (Duchesne et al., 2019). Positive tree growth trends were nevertheless largely confined to relatively young stands (<100 years, Figure S4), consistent with expected stand development. We further observed that tree growth tended to increase faster in cool and moist high-elevation forests, which hinted that besides stand development positive tree growth trends could be partly explained by alleviation of cold limitation (Gao et al., 2022).

Despite positive growth trends, we observed parallel increases in growth autocorrelation and variability for a majority of the study species. On the theoretical basis that system state autocorrelation and variability should increase close to a tipping point (Scheffer et al., 2009), these metrics have been proposed as early warning signals of mortality (Cailleret et al., 2019; Camarero et al., 2015; Hereş et al., 2012). This expectation relies on the common hypothesis that tree growth is a reliable proxy of system state, that is, tree physiological status. However, growth autocorrelation and variability might integrate other signals on top of tree physiology. Notably, autocorrelation and variability of tree growth are partly driven by that of climate and hydrology (Bowers et al., 2013; Coulthard et al., 2020). For example, PDSI is a strong determinant of tree growth in the region and was substantially more autocorrelated at colder sites ($r = -.45$, $p < .001$), hence potentially explaining the strong negative correlation observed between MAT and growth autocorrelation. Autocorrelation can further be determined by species functional traits such as leaf or carbon reserves lifespan (Zweifel & Sterck, 2018) and tree ontogeny (Zweifel et al., 2006). Here, we find strong association between stand structure (tree height), climate and tree growth autocorrelation and variability. Such dependences might thus substantially

blur potential relationships between tree growth autocorrelation, variability, and mortality, which could explain inconsistent reports (Cailleret et al., 2019; Tai et al., 2023).

In parallel to increasing growth autocorrelation and variability, we observed declining growth resilience to drought across several species. Despite mathematical connection between growth variability, autocorrelation, and resilience, these metrics were only weakly correlated, suggesting that they encoded different signals. Decline in tree growth resilience to drought has been previously reported at the local scale (Gazol et al., 2018) and across temperate and boreal forests (Zheng et al., 2021, 2023). The decline observed here was primarily a result of decreasing resistance, which was not compensated for by increasing recovery (Figure 2). Declining drought resistance is often symptomatic of increasing drought intensity and/or competition (Castagneri et al., 2021; Gazol et al., 2018). Although increasing temperature and drought intensity over part of the study area (Andreadis & Lettenmaier, 2006), as well as increasing stand density due to fire suppression (Noss et al., 2006) reflected in current stand structure, could indeed explain some of the decline in tree growth resilience, most of the spatial variability in RI decline was nevertheless related to annual temperature, which was ultimately related to species distribution (Figures S5 and S6; Table S5). Like mortality, species dependence of observed RI decline suggests either species-specific variation in drought vulnerability and/or host-biotic agent interactions. Notably, increasing temperatures may have allowed insect infestation of naive species at higher latitude and elevations (Raffa et al., 2008), causing decline in tree growth resilience to subsequent droughts in these populations. Overall, several factors, including increasing stand competition, drought, and insect outbreaks, are likely responsible for the observed decline in tree growth resilience to drought. In this regard, temporal resilience trends most likely carry information on past stand history and tree physiological damage, which could influence stand vulnerability to subsequent perturbations.

Despite their differences, temporal trends of tree growth autocorrelation, variability and resilience to drought converge to suggest tree physiological decline in the US Mountain West during the second half of the 20th century. This result contrasts with a context of increasing basal area growth and highlights a divergence between tree productivity and physiological status. Furthermore, while tall, productive forests located in moist and cool environments exhibited the most positive temporal growth trends, they were also the most exposed to physiological weakening denoted by declining growth resilience. Progressive lifting of cold limitation might thus benefit tree growth while at the same time exposing trees to larger vulnerability to drought and insect outbreaks, for example, because of structural overshoot and insect population development (Jump et al., 2017; Raffa et al., 2008).

4.3 | Declining resilience mediates stand mortality

Our results reveal strong association between stand-level mortality and tree growth resilience to drought. Long-term average resilience

was negatively associated with mortality, but only within species. While variations in mean resilience between species might reflect different exposition to drought, it may not reflect different susceptibility to mortality. Resilience levels are in fact in large part dependent on growth temporal autocorrelation (Klesse et al., 2022), which is notably related to species functional traits such as leaf and carbon lifespan (Song et al., 2022; Zweifel et al., 2020; Zweifel & Sterck, 2018) or wood density (Serra-Maluquer et al., 2022). Unexpectedly, we found a positive effect of mean resilience on mortality in multivariate models, indicating that all else being equal, trees with higher resilience level were associated with higher stand mortality, similar to a previous remote-sensing study (Tai et al., 2023). This likely reflects larger vulnerability of trees historically little exposed to drought, for example, because of structural overshoot (Jump et al., 2017).

Univariate and multivariate models both indicated that resilience trends were strong predictors of stand mortality, superior to resilience means in that regard. This indicates that trends in resilience successfully capture the integrated effect of past stresses on tree physiological status and susceptibility to mortality. Hence in refinement of previous report on the potential of growth resilience to drought as early warning metric (DeSoto et al., 2020), we propose that declining resilience, rather than absolute resilience, is indicative of unhealthy stands, which are likely more vulnerable to perturbations, regardless of the agent. The observed relationship between declining resilience and mortality further suggests that damage build up induced by repeat perturbations may be an important component of observed stand mortality patterns, consistent with regional and global analyses (Anderegg et al., 2013; Anderegg, Trugman, Badgley, Konings, & Shaw, 2020). For example, declining resilience might be indicative of drought-induced cavitation fatigue, that is, larger hydraulic vulnerability of the xylem induced by previous drought, which is a major factor of subsequent *Populus tremuloides* mortality following drought in the region (Anderegg et al., 2013). More generally, by hindering trees' capacity to regrow functional vascular tissue, declining growth resilience to drought might fundamentally alter trees' water and carbon economies. Processes include reduced hydraulic conductance and increased vulnerability to embolism, reduced carbon assimilation, and shifting C allocation away from resin and defense molecule production, which may compromise tree vulnerability to subsequent biotic and abiotic perturbations (McDowell, 2011; Raffa et al., 2008; Rissanen et al., 2020). At the forest scale, the processes that underlie loss of resilience likely destabilize the system and increase the likelihood that future perturbations may lead forests to transition to an alternate stable state (Hammond, 2020; Johnson et al., 2022).

On the other hand, the capacity to grow new xylem rapidly would confer trees reduced vulnerability to following droughts. Despite resilience level being a weak predictor of mortality, we found that rapid tree growth overall was associated with lower stand mortality rates, consistent with a widely observed pattern (Bigler & Bugmann, 2003; Hülsmann et al., 2018). Hence, rather than the capacity to quickly resume baseline wood growth as denoted by average resilience the ability to maintain resilience to drought and rapid overall tree

growth appears as a key factor to mitigate mortality (Anderegg, Schwalm, et al., 2015), despite rapid tree growth might come at the expense of allocation to drought resistance and pathogen defense (de la Mata et al., 2017; Huot et al., 2014). Multivariate models show that decomposing resilience into resistance and recovery components improved predictions, and further indicate similar importance of growth resistance and recovery components of resilience trends in mediating stand mortality (Figure 5; Table S4). This suggests that stress avoidance and tolerance strategies (Oliveira et al., 2021) can both be successful in mitigating mortality. Our results illustrate the role of resilience in denoting past physiological decline and mediating subsequent stand mortality.

4.4 | Modeling stand mortality

Here, we assess the relative importance of stand, climate, growth, and resilience factors to model stand-level mortality. We make the distinction between explicative (i.e., how much variance can a variable explain overall) and predictive power (i.e., how much does a model improve by adding a new variable) denoted by the marginal and conditional decomposition of model variance, respectively (Grömping, 2007). Moderate model performance overall was expected on the basis of the complexity of processes involved and variability between species, but was within the range of previous studies (Trugman et al., 2021; Venturas et al., 2021). As expected from longstanding literature on the topic, stand and climate factors played an important role in explaining stand mortality (Dietze & Moorcroft, 2011; Neumann et al., 2017; Ruiz-Benito et al., 2013). Stand variables notably most largely contributed to explain the occurrence of mortality observations, related to stand basal area and tree height, but had strikingly little effect on mortality intensity. This is likely related to stand dynamics, where mature stands are more likely to exhibit low intensity mortality (Franklin, 2002). Tree growth and resilience to drought nevertheless had similarly large importance, explaining together about half of observed spatial variations in overall stand-level mortality. Tree growth and resilience performed particularly well to explain mortality intensity, explaining over two thirds of the variance. These results upscale previous tree-level evidence (Bigler & Bugmann, 2003; Cailleret et al., 2016; DeSoto et al., 2020), and demonstrate the potential of growth and resilience metrics to encapsulate tree physiological status and help explain mortality over broad scales, despite large complexity diversity of pathways across species (Lloret et al., 2011).

On the other hand, covariation between tree growth and resilience, climate and stand factors (Lloret et al., 2011; Serra-Maluquer et al., 2022) is important to take into account as it may partially underlie their respective effect on mortality. This is particularly the case for tree growth variation coefficient and autocorrelation, which despite showing a strong effect on mortality across species, is mostly driven by other climate and stand variables, hence yielding nonsignificant parameters in multivariate models (Figure 5b). To a lesser extent, the difference between the conditional and marginal

decomposition of variance (Figure 5a) also suggests that part of the resilience effect on mortality could be explained by other variables in the multivariate model, notably stand and climate factors. This view is supported by the fact that resilience exhibited limited unique model variance. Although this result does not necessarily imply non-causality between resilience and mortality, it does suggest modest gains from adding resilience in a model for prediction purposes. The relative importance of tree growth on the other hand appeared to be robust to the adopted perspective, indicating limited redundancy with other variables in the model. The potential of past tree growth to predict future mortality that we observe here supports its common use in vegetation models (Hülsmann et al., 2018). The existence of a survivorship and dominance bias in our dataset (mostly live and often dominant trees were sampled (DeRose et al., 2017)) might have led to underestimating the growth- and resilience-mortality relationships. Analyses based on a more representative sampling design might thus conclude to a larger relative importance of these metrics to explain and predict stand mortality.

Our results highlight the strengths but also some of the limitations of tree growth-based resilience as early warning signal of stand mortality. Models, including that of forest mortality, need to strike a balance between predictive power versus the number of parameters and associated costs (Bentler & Mooijaart, 1989). Although model selection criterion (AIC) suggested that the benefit of including resilience outweighs statistical costs, further costs associated with enabling resilience-based forest mortality models need to be considered. Notably, such a model would require extensive and annually resolved tree growth monitoring. Existing national forest inventory programs generally lack temporal resolution and despite adding core sampling to inventories would provide necessary resolution (DeRose et al., 2017; Evans et al., 2022), repeat sampling and processing to enable updated predictions might reveal costly. Solving this issue would require lower costs and/or higher benefits of including resilience metrics in mortality models. Remote sensing of canopy status thus seems a promising avenue in this regard, although mortality inferences have been restricted in time so far, which limits the capacity to take action (Liu et al., 2019; Rogers et al., 2018). Automated tree measurement networks such as TreeNet (Zweifel et al., 2021) could also show potential, in that they might enable exploring new proxies of tree physiological status in addition to reducing the return time to forest plots. Spatial and temporal limitations to such networks yet currently hinders testing these applications. Despite limitations, tree ring-based estimates of forest resilience hold great value for retrospective analysis of past mortality events and to gain insights on underlying mechanisms. Tree ring-based resilience proved to be a reliable proxy of tree physiological status and forest vulnerability to perturbations. In that sense, widespread decline in tree growth resilience to drought (Zheng et al., 2021) adds to the growing evidence of increasing risk of pervasive forest mortality at the global scale (Allen et al., 2015; Hammond et al., 2022).

AUTHOR CONTRIBUTIONS

Antoine Cabon and William R. L. Anderegg designed the study. R. Justin DeRose and John D. Shaw developed the enhanced FIA core

collection protocols and compiled and processed the data. Antoine Cabon conducted the analyses. All coauthors wrote the manuscript.

ACKNOWLEDGMENTS

We acknowledge funding from USDA National Institute of Food and Agriculture, Agricultural and Food Research Initiative Competitive Programme, Ecosystem Services and Agro-Ecosystem Management, grant no. 2018-67019-27850. WA acknowledges support from the David and Lucille Packard Foundation and US National Science Foundation grants no. 1714972, 1802880, 2044937, 2003017, and the Alan T. Waterman Award, IOS-2325700. This research was supported by the Utah Agricultural Experiment Station, Utah State University, and approved as journal paper number 9651.

CONFLICT OF INTEREST STATEMENT

The author declare that they have no conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study can be accessed from Figshare at [10.6084/m9.figshare.23390036](https://figshare.com/10.6084/m9.figshare.23390036).

ORCID

Antoine Cabon  <https://orcid.org/0000-0001-6426-1726>

William R. L. Anderegg  <https://orcid.org/0000-0001-6551-3331>

REFERENCES

- Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A., & Hegewisch, K. C. (2018). TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958–2015. *Scientific Data*, 5, 1–12.
- Allen, C. D., Breshears, D. D., & McDowell, N. G. (2015). On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere*, 6, art129.
- Anderegg, W. R. L., & Callaway, E. S. (2012). Infestation and hydraulic consequences of induced carbon starvation. *Plant Physiology*, 159, 1866–1874.
- Anderegg, W. R. L., Hicke, J. A., Fisher, R. A., Allen, C. D., Aukema, J., Bentz, B., Hood, S., Lichstein, J. W., Macalady, A. K., McDowell, N., Pan, Y., Raffa, K., Sala, A., Shaw, J. D., Stephenson, N. L., Tague, C., & Zeppel, M. (2015). Tree mortality from drought, insects, and their interactions in a changing climate. *New Phytologist*, 208, 674–683.
- Anderegg, W. R. L., Plavcová, L., Anderegg, L. D. L., Hacke, U. G., Berry, J. A., & Field, C. B. (2013). Drought's legacy: Multiyear hydraulic deterioration underlies widespread aspen forest die-off and portends increased future risk. *Global Change Biology*, 19, 1188–1196.
- Anderegg, W. R. L., Trugman, A. T., Badgley, G., Anderson, C. M., Bartuska, A., Ciais, P., Cullenward, D., Field, C. B., Freeman, J., Goetz, S. J., Hicke, J. A., Huntzinger, D., Jackson, R. B., Nickerson, J., Pacala, S., & Randerson, J. T. (2020). Climate-driven risks to the climate mitigation potential of forests. *Science*, 368, eaaz7005.
- Anderegg, W. R. L., Trugman, A. T., Badgley, G., Konings, A. G., & Shaw, J. (2020). Divergent forest sensitivity to repeated extreme droughts. *Nature Climate Change*, 10(12), 1091–1095.
- Anderegg, W. R. L., Schwalm, C., Biondi, F., Camarero, J. J., Koch, G., Litvak, M., Ogle, K., Shaw, J. D., Shevliakova, E., Williams, A. P., Wolf, A., Ziaco, E., & Pacala, S. (2015). Pervasive drought legacies in forest ecosystems and their implications for carbon cycle models. *Science*, 349, 528–532.
- Andreadis, K. M., & Lettenmaier, D. P. (2006). Trends in 20th century drought over the continental United States: U.S. drought trends. *Geophysical Research Letters*, 33, L10403.

- Bechtold, W. A., & Scott, C. T. (2005). The forest inventory and analysis plot design. In W. A. Bechtold & P. L. Patterson (Eds.), *The enhanced forest inventory and analysis program: National sampling design and estimation procedures* (Vol. 1, pp. 27–67). USDA Forest Service, Southern Research Station.
- Benito Garzón, M., González Muñoz, N., Wigneron, J. P., Moisy, C., Fernández-Manjarrés, J., & Delzon, S. (2018). The legacy of water deficit on populations having experienced negative hydraulic safety margin. *Global Ecology and Biogeography*, 27, 346–356.
- Bentler, P. M., & Mooijart, A. (1989). Choice of structural model via parsimony: A rationale based on precision. *Psychological Bulletin*, 106, 315–317.
- Betts, M. G., Wolf, C., Ripple, W. J., Phalan, B., Millers, K. A., Duarte, A., Butchart, S. H. M., & Levi, T. (2017). Global forest loss disproportionately erodes biodiversity in intact landscapes. *Nature*, 547, 441–444.
- Bigler, C., & Bugmann, H. (2003). Growth-dependent tree mortality models based on tree rings. *Canadian Journal of Forest Research*, 33, 210–221.
- Bowers, M. C., Gao, J. B., & Tung, W. W. (2013). Long range correlations in tree ring chronologies of the USA: Variation within and across species. *Geophysical Research Letters*, 40, 568–572.
- Breshears, D. D., Cobb, N. S., Rich, P. M., Price, K. P., Allen, C. D., Balice, R. G., Romme, W. H., Kastens, J. H., Floyd, M. L., Belnap, J., Anderson, J. J., Myers, O. B., & Meyer, C. W. (2005). Regional vegetation die-off in response to global-change-type drought. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 15144–15148.
- Breshears, D. D., Myers, O. B., Meyer, C. W., Barnes, F. J., Zou, C. B., Allen, C. D., McDowell, N. G., & Pockman, W. T. (2009). Tree die-off in response to global change-type drought: Mortality insights from a decade of plant water potential measurements. *Frontiers in Ecology and the Environment*, 7, 185–189.
- Cailleret, M., Dakos, V., Jansen, S., Robert, E. M. R., Aakala, T., Amoroso, M. M., Antos, J. A., Bigler, C., Bugmann, H., Caccianiga, M., & Camarero, J. J. (2019). Early-warning signals of individual tree mortality based on annual radial growth. *Frontiers in Plant Science*, 9, 1–14.
- Cailleret, M., Jansen, S., Robert, E. M. R., Desoto, L., Aakala, T., Antos, J. A., Beikircher, B., Bigler, C., Bugmann, H., Caccianiga, M., Čada, V., Camarero, J. J., Cherubini, P., Cochard, H., Coyea, M. R., Čufar, K., das, A. J., Davi, H., Delzon, S., ... Martínez-Vilalta, J. (2016). A synthesis of radial growth patterns preceding tree mortality. *Global Change Biology*, 23, 1–16.
- Camarero, J. J., Gazol, A., Sangüesa-Barreda, G., Oliva, J., & Vicente-Serrano, S. M. (2015). To die or not to die: Early warnings of tree dieback in response to a severe drought. *Journal of Ecology*, 103, 44–57.
- Castagneri, D., Vacchiano, G., Hacket-Pain, A., DeRose, R. J., Klein, T., & Bottero, A. (2021). Meta-analysis reveals different competition effects on tree growth resistance and resilience to drought. *Ecosystems*, 25, 30–43.
- Chapman, T. B., Veblen, T. T., & Schoennagel, T. (2012). Spatiotemporal patterns of mountain pine beetle activity in the southern Rocky Mountains. *Ecology*, 93, 11–2185.
- Coulthard, B. L., St. George, S., & Meko, D. M. (2020). The limits of freely-available tree-ring chronologies. *Quaternary Science Reviews*, 234, 106264.
- de la Mata, R., Hood, S., & Sala, A. (2017). Insect outbreak shifts the direction of selection from fast to slow growth rates in the long-lived conifer *Pinus ponderosa*. *Proceedings of the National Academy of Sciences*, 114, 7391–7396.
- DeRose, R. J., Shaw, J. D., & Long, J. N. (2017). Building the forest inventory and analysis tree-ring data set. *Journal of Forestry*, 115, 283–291.
- DeSoto, L., Cailleret, M., Sterck, F., Jansen, S., Kramer, K., Robert, E. M. R., Aakala, T., Amoroso, M. M., Bigler, C., Camarero, J. J., Čufar, K., Gea-Izquierdo, G., Gillner, S., Haavik, L. J., Hereş, A. M., Kane, J. M., Kharuk, V. I., Kitzberger, T., Klein, T., ... Martínez-Vilalta, J. (2020). Low growth resilience to drought is related to future mortality risk in trees. *Nature Communications*, 11, 545.
- Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin, P. R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences*, 105, 6668–6672.
- Dietze, M. C., & Moorcroft, P. R. (2011). Tree mortality in the eastern and Central United States: Patterns and drivers. *Global Change Biology*, 17, 3312–3326.
- Duchesne, L., Houle, D., Ouimet, R., Caldwell, L., Gloor, M., & Brienens, R. (2019). Large apparent growth increases in boreal forests inferred from tree-rings are an artefact of sampling biases. *Scientific Reports*, 9, 6832.
- Evans, M. E. K., DeRose, R. J., Klesse, S., Girardin, M. P., Heilman, K. A., Alexander, M. R., Arsenault, A., Babst, F., Bouchard, M., Cahoon, S. M. P., Campbell, E. M., Dietze, M., Duchesne, L., Frank, D. C., Giebk, C. L., Gómez-Guerrero, A., García, G. G., Hogg, E. H., Metsaranta, J., ... Gaytán, S. A. V. (2022). Adding tree rings to North America's National Forest Inventories: An essential tool to guide drawdown of atmospheric CO₂. *Bioscience*, 72, 233–246.
- Feng, X., Merow, C., Liu, Z., Park, D. S., Roehrdanz, P. R., Maitner, B., Newman, E. A., Boyle, B. L., Lien, A., Burger, J. R., Pires, M. M., Brando, P. M., Bush, M. B., McMichael, C. N. H., Neves, D. M., Nikolopoulos, E. I., Saleska, S. R., Hannah, L., Breshears, D. D., ... Enquist, B. J. (2021). How deregulation, drought and increasing fire impact Amazonian biodiversity. *Nature*, 597, 516–521.
- Forzieri, G., Dakos, V., McDowell, N. G., Ramdane, A., & Cescatti, A. (2022). Emerging signals of declining forest resilience under climate change. *Nature*, 608, 534–539.
- Frank, J. M., Massman, W. J., Ewers, B. E., Huckaby, L. S., & Negrón, J. F. (2014). Ecosystem CO₂/H₂O fluxes are explained by hydraulically limited gas exchange during tree mortality from spruce bark beetles. *Journal of Geophysical Research: Biogeosciences*, 119, 1195–1215.
- Franklin, J. (2002). Disturbances and structural development of natural forest ecosystems with silvicultural implications, using Douglas-fir forests as an example. *Forest Ecology and Management*, 155, 399–423.
- Gao, S., Liang, E., Liu, R., Babst, F., Camarero, J. J., Fu, Y. H., Piao, S., Rossi, S., Shen, M., Wang, T., & Peñuelas, J. (2022). An earlier start of the thermal growing season enhances tree growth in cold humid areas but not in dry areas. *Nature Ecology and Evolution*, 6, 397–404.
- Gaylord, M. L., Kolb, T. E., Pockman, W. T., Plaut, J. A., Yezpe, E. A., Macalady, A. K., Pangle, R. E., & McDowell, N. G. (2013). Drought predisposes piñon-juniper woodlands to insect attacks and mortality. *New Phytologist*, 198, 567–578.
- Gazol, A., Camarero, J. J., Vicente-Serrano, S. M., Sánchez-Salguero, R., Gutiérrez, E., de Luis, M., Sangüesa-Barreda, G., Novak, K., Rozas, V., Tiscar, P. A., Linares, J. C., Martín-Hernández, N., Martínez del Castillo, E., Ribas, M., García-González, I., Silla, F., Camisón, A., Génova, M., Olano, J. M., ... Galván, J. D. (2018). Forest resilience to drought varies across biomes. *Global Change Biology*, 24, 1–16.
- Greenwood, D. L., & Weisberg, P. J. (2008). Density-dependent tree mortality in piñon-juniper woodlands. *Forest Ecology and Management*, 255, 2129–2137.
- Grömping, U. (2007). Estimators of relative importance in linear regression based on variance decomposition. *American Statistician*, 61, 139–147.
- Grote, R., Gessler, A., Hommel, R., Poschenrieder, W., & Priesack, E. (2016). Importance of tree height and social position for drought-related stress on tree growth and mortality. *Trees*, 30, 1467–1482.

- Hammond, W. M. (2020). A matter of life and death: Alternative stable states in trees, from xylem to ecosystems. *Frontiers in Forests and Global Change*, 3, 560409.
- Hammond, W. M., Williams, A. P., Abatzoglou, J. T., Adams, H. D., Klein, T., López, R., Sáenz-Romero, C., Hartmann, H., Breshears, D. D., & Allen, C. D. (2022). Global field observations of tree die-off reveal hotter-drought fingerprint for Earth's forests. *Nature Communications*, 13, 1761.
- Hereş, A.-M., Martínez-Vilalta, J., & Claramunt López, B. (2012). Growth patterns in relation to drought-induced mortality at two scots pine (*Pinus sylvestris* L.) sites in NE Iberian Peninsula. *Trees*, 26, 621–630.
- Hülsmann, L., Bugmann, H., Cailleret, M., & Brang, P. (2018). How to kill a tree: Empirical mortality models for 18 species and their performance in a dynamic forest model. *Ecological Applications*, 28, 522–540.
- Huot, B., Yao, J., Montgomery, B. L., & He, S. Y. (2014). Growth–defense tradeoffs in plants: A balancing act to optimize fitness. *Molecular Plant*, 7, 1267–1287.
- Johnson, D. M., Katul, G., & Domec, J.-C. (2022). Catastrophic hydraulic failure and tipping points in plants. *Plant, Cell & Environment*, 45, 2231–2266.
- Jump, A. S., Ruiz-Benito, P., Greenwood, S., Allen, C. D., Kitzberger, T., Fensham, R., Martínez-Vilalta, J., & Lloret, F. (2017). Structural overshoot of tree growth with climate variability and the global spectrum of drought-induced forest dieback. *Global Change Biology*, 23, 3742–3757.
- Kannenbergh, S. A., Novick, K. A., Alexander, M. R., Maxwell, J. T., Moore, D. J. P., Phillips, R. P., & Anderegg, W. R. L. (2019). Linking drought legacy effects across scales: From leaves to tree rings to ecosystems. *Global Change Biology*, 25, 2978–2992.
- Klesse, S., Babst, F., Evans, M. E. K., Hurley, A., Pappas, C., & Peters, R. L. (2022). Legacy effects in radial tree growth are rarely significant after accounting for biological memory. *Journal of Ecology*, 1365–2745, 14045.
- Kolb, T., Keefover-Ring, K., Burr, S. J., Hofstetter, R., Gaylord, M., & Raffa, K. F. (2019). Drought-mediated changes in tree physiological processes weaken tree defenses to bark beetle attack. *Journal of Chemical Ecology*, 45, 888–900.
- Liu, Y., Kumar, M., Katul, G. G., & Porporato, A. (2019). Reduced resilience as an early warning signal of forest mortality. *Nature Climate Change*, 9, 880–885.
- Lloret, F., Keeling, E. G., & Sala, A. (2011). Components of tree resilience: Effects of successive low-growth episodes in old ponderosa pine forests. *Oikos*, 120, 1909–1920.
- Marchetti, S. B., Worrall, J. J., & Eager, T. (2011). Secondary insects and diseases contribute to sudden aspen decline in southwestern Colorado, USA. *Canadian Journal of Forest Research*, 41, 2315–2325.
- McDowell, N. G. (2011). Mechanisms linking drought, hydraulics, carbon metabolism, and vegetation mortality. *Plant Physiology*, 155, 1051–1059.
- McDowell, N. G., & Allen, C. D. (2015). Darcy's law predicts widespread forest mortality under climate warming. *Nature Climate Change*, 5, 669–672.
- McDowell, N. G., Allen, C. D., Anderson-Teixeira, K., Aukema, B. H., Bond-Lamberty, B., Chini, L., Clark, J. S., Dietze, M., Grossiord, C., Hanbury-Brown, A., Hurtt, G. C., Jackson, R. B., Johnson, D. J., Kueppers, L., Lichstein, J. W., Ogle, K., Poulter, B., Pugh, T. A. M., Seidl, R., ... Xu, C. (2020). Pervasive shifts in forest dynamics in a changing world. *Science*, 368, eaaz9463.
- McDowell, N. G., Sapes, G., Pivovarov, A., Adams, H. D., Allen, C. D., Anderegg, W. R. L., Arend, M., Breshears, D. D., Brodribb, T., Choat, B., Cochard, H., De Cáceres, M., De Kauwe, M. G., Grossiord, C., Hammond, W. M., Hartmann, H., Hoch, G., Kahmen, A., Klein, T., ... Xu, C. (2022). Mechanisms of woody-plant mortality under rising drought, CO₂ and vapour pressure deficit. *Nature Reviews Earth & Environment*, 3, 294–308.
- McNellis, B. E., Smith, A. M. S., Hudak, A. T., & Strand, E. K. (2021). Tree mortality in western U.S. forests forecasted using forest inventory and random Forest classification. *Ecosphere*, 12, e03419.
- Meddens, A. J. H., Hicke, J. A., Macalady, A. K., Buotte, P. C., Cowles, T. R., & Allen, C. D. (2015). Patterns and causes of observed piñon pine mortality in the southwestern United States. *New Phytologist*, 206, 91–97.
- Neumann, M., Mues, V., Moreno, A., Hasenauer, H., & Seidl, R. (2017). Climate variability drives recent tree mortality in Europe. *Global Change Biology*, 23, 4788–4797.
- Noss, R. F., Franklin, J. F., Baker, W. L., Schoennagel, T., & Moyle, P. B. (2006). Managing fire-prone forests in the western United States. *Frontiers in Ecology and the Environment*, 4, 481–487.
- Oliveira, R. S., Eller, C. B., Barros, F. d. V., Hirota, M., Brum, M., & Bittencourt, P. (2021). Linking plant hydraulics and the fast-slow continuum to understand resilience to drought in tropical ecosystems. *New Phytologist*, 230, 904–923.
- Paine, T. D., Raffa, K. F., & Harrington, T. C. (1997). Interactions among Scolytid bark beetles, their associated fungi, and live host conifers. *Annual Review of Entomology*, 42, 179–206.
- Preisler, Y., Tatarinov, F., Grünzweig, J. M., & Yakir, D. (2021). Seeking the “point of no return” in the sequence of events leading to mortality of mature trees. *Plant, Cell & Environment*, 44, 1315–1328.
- Pretzsch, H., & Biber, P. (2010). Size-symmetric versus size-asymmetric competition and growth partitioning among trees in forest stands along an ecological gradient in Central Europe. *Canadian Journal of Forest Research*, 40, 370–384.
- Raffa, K. F., Aukema, B. H., Bentz, B. J., Carroll, A. L., Hicke, J. A., Turner, M. G., & Romme, W. H. (2008). Cross-scale drivers of natural disturbances prone to anthropogenic amplification: The dynamics of bark beetle eruptions. *Bioscience*, 58, 501–517.
- Reed, C. C., & Hood, S. M. (2021). Few generalizable patterns of tree-level mortality during extreme drought and concurrent bark beetle outbreaks. *Science of the Total Environment*, 750, 141306.
- Rissanen, K., Vanhatalo, A., Salmon, Y., Bäck, J., & Hölttä, T. (2020). Stem emissions of monoterpenes, acetaldehyde and methanol from scots pine (*Pinus sylvestris* L.) affected by tree–water relations and cambial growth. *Plant Cell and Environment*, 43, 1751–1765.
- Rogers, B. M., Solvik, K., Hogg, E. H., Ju, J., Masek, J. G., Michaelian, M., Berner, L. T., & Goetz, S. J. (2018). Detecting early warning signals of tree mortality in boreal North America using multiscale satellite data. *Global Change Biology*, 24(6), 2284–2304.
- Ruiz-Benito, P., Lines, E. R., Gómez-Aparicio, L., Zavala, M. A., & Coomes, D. A. (2013). Patterns and drivers of tree mortality in Iberian forests: Climatic effects are modified by competition (a Hector, Ed.). *PLoS ONE*, 8, e56843.
- Scheffer, M., Bascompte, J., Brock, W. A., Brovkin, V., Carpenter, S. R., Dakos, V., Held, H., van Nes, E. H., Rietkerk, M., & Sugihara, G. (2009). Early-warning signals for critical transitions. *Nature*, 461, 53–59.
- Scheffer, M., Carpenter, S., Foley, J. A., Folke, C., & Walker, B. (2001). Catastrophic shifts in ecosystems. *Nature*, 413, 591–596.
- Serra-Maluquer, X., Gazol, A., Anderegg, W. R. L., Martínez-Vilalta, J., Mencuccini, M., & Camarero, J. J. (2022). Wood density and hydraulic traits influence species' growth response to drought across biomes. *Global Change Biology*, 28, 3871.
- Sevanto, S., McDowell, N. G., Dickman, L. T., Pangle, R., & Pockman, W. T. (2014). How do trees die? A test of the hydraulic failure and carbon starvation hypotheses. *Plant, Cell & Environment*, 37, 153–161.
- Shaw, J. D., Steed, B. E., & DeBlander, L. T. (2005). Forest inventory and analysis (FIA) annual inventory answers the question: What is happening to pinyon-Juniper woodlands? *Journal of Forestry*, 103, 280–285.

- Song, Y., Sterck, F., Zhou, X., Liu, Q., Kruijt, B., & Poorter, L. (2022). Drought resilience of conifer species is driven by leaf lifespan but not by hydraulic traits. *New Phytologist*, 235, 978–992.
- Sperry, J. S., & Tyree, M. T. (1988). Mechanism of water stress-induced xylem embolism. *Plant Physiology*, 88, 581–587.
- Stanke, H., Finley, A. O., Domke, G. M., Weed, A. S., & MacFarlane, D. W. (2021). Over half of western United States' most abundant tree species in decline. *Nature Communications*, 451, 1–11.
- Tai, X., Trugman, A. T., & Anderegg, W. R. L. (2023). Linking remotely sensed ecosystem resilience with forest mortality across the continental United States. *Global Change Biology*, 29(4), 1096–1105.
- Trugman, A. T., Anderegg, L. D. L., Anderegg, W. R. L., Das, A. J., & Stephenson, N. L. (2021). Why is tree drought mortality so hard to predict? *Trends in Ecology and Evolution*, 36, 520–532.
- Trugman, A. T., Detto, M., Bartlett, M. K., Medvigy, D., Anderegg, W. R. L., Schwalm, C., Schaffer, B., & Pacala, S. W. (2018). Tree carbon allocation explains forest drought-kill and recovery patterns. *Ecology Letters*, 21, 1552–1560.
- Turtola, S., Manninen, A.-M., Rikala, R., & Kainulainen, P. (2003). Drought stress alters the concentration of wood Terpenoids in scots pine and Norway spruce seedlings. *Journal of Chemical Ecology*, 29, 1981–1995.
- Van Mantgem, P. J., Stephenson, N. L., Byrne, J. C., Daniels, L. D., Franklin, J. F., Fulé, P. Z., Harmon, M. E., Larson, A. J., Smith, J. M., Taylor, A. H., & Veblen, T. T. (2009). Widespread increase of tree mortality rates in the Western United States. *Science*, 323, 521–524.
- Venturas, M. D., Sperry, J. S., Love, D. M., Frehner, E. H., Allred, M. G., Wang, Y., & Anderegg, W. R. L. (2018). A stomatal control model based on optimization of carbon gain versus hydraulic risk predicts aspen sapling responses to drought. *New Phytologist*, 220, 836–850.
- Venturas, M. D., Todd, H. N., Trugman, A. T., & Anderegg, W. R. L. (2021). Understanding and predicting forest mortality in the western United States using long-term forest inventory data and modeled hydraulic damage. *New Phytologist*, 230, 1896–1910.
- Williams, A. P., Allen, C. D., Macalady, A. K., Griffin, D., Woodhouse, C. A., Meko, D. M., Swetnam, T. W., Rauscher, S. A., Seager, R., Grissino-Mayer, H. D., Dean, J. S., Cook, E. R., Gangadagamage, C., Cai, M., & McDowell, N. G. (2012). Temperature as a potent driver of regional forest drought stress and tree mortality. *Nature Climate Change*, 3, 292–297.
- Worrall, J. J., Marchetti, S. B., Egeland, L., Mask, R. A., Eager, T., & Howell, B. (2010). Effects and etiology of sudden aspen decline in south-western Colorado, USA. *Forest Ecology and Management*, 260, 638–648.
- Zhang, F., Biederman, J. A., Dannenberg, M. P., Yan, D., Reed, S. C., & Smith, W. K. (2021). Five decades of observed daily precipitation reveal longer and more variable drought events across much of the Western United States. *Geophysical Research Letters*, 48, 1–11.
- Zheng, T., Martínez-Vilalta, J., García-Valdés, R., Gazol, A., Camarero, J. J., & Mencuccini, M. (2021). Disentangling biology from mathematical necessity in twentieth-century gymnosperm resilience trends. *Nature Ecology and Evolution*, 5, 733–735.
- Zheng, T., Martínez-Vilalta, J., García-Valdés, R., Gazol, A., Camarero, J. J., Mu, C., & Mencuccini, M. (2023). Growth plasticity of conifers did not avoid declining resilience to soil and atmospheric droughts during the 20th century. *Forest Ecosystems*, 10, 100107.
- Zweifel, R., Etzold, S., Basler, D., Bischoff, R., Braun, S., Buchmann, N., Conedera, M., Fonti, P., Gessler, A., Haeni, M., Hoch, G., Kahmen, A., Köchli, R., Maeder, M., Nievergelt, D., Peter, M., Peters, R. L., Schaub, M., Trotsiuk, V., ... Eugster, W. (2021). TreeNet-the biological drought and growth indicator network. *Frontiers in Forests and Global Change*, 4, 1–14.
- Zweifel, R., Etzold, S., Sterck, F., Gessler, A., Anfodillo, T., Mencuccini, M., von Arx, G., Lazzarin, M., Haeni, M., Feichtinger, L., Meusburger, K., Knuesel, S., Walther, L., Salmon, Y., Bose, A. K., Schoenbeck, L., Hug, C., de Girardi, N., Giuggiola, A., ... Rigling, A. (2020). Determinants of legacy effects in pine trees—implications from an irrigation-stop experiment. *New Phytologist*, 227, 1081–1096.
- Zweifel, R., & Sterck, F. (2018). A conceptual tree model explaining legacy effects on stem growth. *Frontiers in Forests and Global Change*, 1, 1–9.
- Zweifel, R., Zimmermann, L., Zeugin, F., & Newbery, D. M. (2006). Intra-annual radial growth and water relations of trees: Implications towards a growth mechanism. *Journal of Experimental Botany*, 57, 1445–1459.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Cabon, A., DeRose, R. J., Shaw, J. D., & Anderegg, W. R. L. (2023). Declining tree growth resilience mediates subsequent forest mortality in the US Mountain West. *Global Change Biology*, 29, 4826–4841. <https://doi.org/10.1111/gcb.16826>