

Review

Seasonality of forest insects: why diapause matters

Martin Schebeck^{1,10,*,@}, Philipp Lehmann^{2,3,4,10}, Mathieu Laparie⁵, Barbara J. Bentz⁶, Gregory J. Ragland⁷, Andrea Battisti⁸, and Daniel A. Hahn⁹

Insects have major impacts on forest ecosystems, from herbivory and soil-nutrient cycling to killing trees at a large scale. Forest insects from temperate, tropical, and subtropical regions have evolved strategies to respond to seasonality; for example, by entering diapause, to mitigate adversity and to synchronize lifecycles with favorable periods. Here, we show that distinct functional groups of forest insects; that is, canopy dwellers, trunk-associated species, and soil/litter-inhabiting insects, express a variety of diapause strategies, but do not show systematic differences in diapause strategy depending on functional group. Due to the overall similarities in diapause strategies, we can better estimate the impacts of anthropogenic change on forest insect populations and, consequently, on key ecosystems.

Seasonal adaptations and diapause

Most forest insects experience predictable periods of seasonal stress every year [1,2]. Numerous species use dormancy to mitigate stressful times and synchronize lifecycles with times of the year favorable for development and reproduction [3,4]. Often dormancy is associated with overwintering and cold conditions in temperate habitats, but dormancy can also occur in tropical or subtropical climates in response to hot or dry conditions [1,4]. Dormancy as a life-history strategy can be broken down into a suite of tactics (Box 1). Diapause is a programmed alternative developmental trajectory, a state of developmental suppression that insects enter prior to the onset of stressful conditions [1]. Diapause can either be induced facultatively in response to predictable seasonal cues (e.g., photoperiod, temperature, or humidity) or can be obligate, induced every generation regardless of environmental conditions [5]. Not all insects use diapause to survive unfavorable seasons. Some species become quiescent, wherein the normal lifecycle is slowed down or suppressed directly by the immediate environment [4]. Some insects even use a combination of tactics over the course of an unfavorable season; for example, by beginning their dormant period in a programmed diapause, but ending that diapause before the onset of the growing season and remaining quiescent until favorable conditions return [1,6,7]. Ending dormancy in quiescence allows insects to immediately resume their lifecycle once stressful conditions end, to exploit the growing season and synchronize individuals across a population.

Forests are complex habitats for dormancy

Forests offer variable habitats across space and time (Box 2) [8]. The vertical profile of forest layers [9], including soil, litter, herbaceous stratum, shrubs, trunks (including wood cavities or galleries), canopy, and internal gradients [10,11], modulates light penetration, radiation, airflow, and consequently temperature and humidity [12]. All these factors likely contribute to the stratification of insect fauna [13] (Figure 1). Although the complexity and height of vertical gradients can vary depending on latitude, elevation, tree species richness, and density, the averages and magnitudes of microclimatic conditions can vary within each stratum [12,14]. For example, the canopy

Highlights

Diapause is a strategy in insects to mitigate seasonal stress and synchronize lifecycles with favorable times.

Forests are 3D, dynamic ecosystems with contrasting microhabitats where numerous insects enter diapause and the duration of diapause affects their phenology.

Ecologically important forest insects, like defoliators, bark breeders, or soil/litter-dwelling species have diverse diapause strategies.

Anthropogenic change, like climate change and urbanization, has and will have substantial impacts on forest insects. Some species may increase their impact on ecosystems, while that of others may be reduced.

Understanding how climate change or urbanization factors, like artificial light at night and the urban-heat-island effect, affect diapause may help develop generalizable rules to predict which species will show population expansion and which are likely to contract.

¹Institute of Forest Entomology, Forest Pathology and Forest Protection, Department of Forest and Soil Sciences, BOKU University, A-1190 Vienna, Austria

²Department of Animal Physiology, Zoological Institute and Museum, University of Greifswald, D-17489 Greifswald, Germany

³Department of Zoology, Stockholm University, SE-10691 Stockholm, Sweden

⁴Bolin Centre for Climate Research, SE-10691 Stockholm, Sweden

⁵INRAE, URZF, F-45075 Orleans, France

⁶US Department of Agriculture, Forest Service, Rocky Mountain Research Station, Logan, UT 84321, USA



Box 1. Key concepts for insect diapause

Many insects incorporate diapause as part of their lifecycles [1], although research effort is often skewed towards pest species. Dormancy is an umbrella term to describe any major slowdown in insect lifecycles. Dormancy can either be induced directly by harsh conditions in the environment, a state termed quiescence, or programmed to begin preemptively before conditions become unfavorable for growth or reproduction, a state termed diapause. Prevention of growth or reproduction by environmentally induced quiescence is common in lifecycles of many insects. From cold winter temperatures or hot summer conditions to lack of food sources, insects enter quiescence rapidly and also exit from quiescence rapidly to reengage in growth or reproduction as soon as conditions become permissive again. By contrast, diapause is a pre-programmed form of dormancy insects enter prior to the onset of adversity. Whereas quiescence can be entered in any lifecycle stage in response to the environment, diapause is a distinct developmental trajectory within the lifecycle that follows a stereotypical series of developmental phases and is typically only expressed in particular, species-specific life stages (Figure 1). These phases include induction, a developmental window when the insect becomes responsive to environmental cues that will cause it to enter the diapause trajectory. Preparation is a phase that might overlap with induction where the insect changes its physiology to accumulate additional nutrients or induce mechanisms of stress hardness. Diapause initiation corresponds to the phase when the insect will enter the state of developmental slowdown. Diapause maintenance is a state in which progression of the lifecycle is arrested. During the maintenance phase, insects are refractory to environmental cues that cause them to resume growth or reproduction. During diapause termination, insects become receptive to environmental cues that end diapause and induce growth or reproduction (followed by postdiapause). For some insects, diapause only ends once favorable times have already begun. Other insects will end diapause during unfavorable times (e.g., in mid-winter) and remain dormant due to environmentally induced quiescence but can readily resume their direct development or reproductive maturation as soon as conditions become permissive again. In multivoltine species, diapause is typically a facultative plastic response, induced by cues that predict oncoming unfavorable conditions (e.g., short-day photoperiods that portend the onset of winter). Diapause is an obligate part of the lifecycles of most univoltine insects and is expressed every generation regardless of environmental cues, thus obligate-diapausing insects lack the induction phase.

⁷Department of Integrative Biology, University of Colorado-Denver, Denver, CO 80204, USA

⁸Department of Agronomy, Food, Natural Resources, Animals and Environment (DAFNAE), University of Padova, I-35020 Legnaro, Italy

⁹Department of Entomology and Nematology, University of Florida, Gainesville, FL 32611-0620, USA

¹⁰These authors contributed equally to this work.

*Correspondence: martin.schebeck@boku.ac.at (M. Schebeck).

@Twitter: @martin_schebeck




Figure 1. Insect diapause is an alternative developmental trajectory with a series of physiologically distinct sequential phases that define the extent to which an insect is both dormant and receptive or refractory to cues that would end dormancy and restart the lifecycle.

intercepts and absorbs most of the sunlight while being exposed to the highest airflow, whereas ground level strata receive much less light or wind and can collect moisture from the strata above. A diverse range of contrasting habitats at a relatively small spatial scale means that seemingly sympatric forest insects can experience distinct environments even when they share the same plant in the same forest, depending on their **phenology** (see [Glossary](#)), lifespan, age, typical feeding substrates and microrefugia, and their ability to behaviorally navigate along the vertical gradient to mitigate immediate conditions [8,13,15].

Forests are long-lived ecosystems whose biophysical characteristics change cyclically over mid- and long-term timescales. In temperate deciduous forests, for example, autumnal foliage loss and associated changes in evapotranspiration alter light and microclimatic conditions from canopy to ground, while accumulation of dead leaves can modify the sheltering properties and **microclimate** of litter [16]. Finally, forests undergo profound changes over decades to centuries [17] due to tree growth, mortality, natural succession, and regeneration, as well as human interventions [18] (Box 2). Seasonal and long-term variation alters vertical gradients of forests by either exacerbating or disrupting microclimatic properties and structure of particular strata.

The multidimensionality of forests in space and time can modulate environmental factors that cue diapause initiation and termination, influencing the timing and duration of dormant and non-dormant lifecycle stages. Consideration of the multidimensionality of subhabitats within forests is therefore crucial to understand the ecology of forest insects [2], especially because insects can in turn play a major role in altering microclimate regulation of forests by defoliating or killing trees.

Significance of forest insects

Insects play multiple roles in forest ecosystems. They feed as herbivores on plant tissues, pollinate plants, disperse seeds and pollen, decompose organic matter, or prey on/parasitize other species [19,20]. Additionally, some insects, particularly herbivorous species, interfere with anthropogenic interests for certain ecosystem services and resources provided by forests [21].

From an ecological perspective, three functional groups including species from numerous taxa are of high significance in forests: (i) canopy-dwelling insects feeding on leaves/needles; (ii) trunk-associated species living on the phloem/xylem of trees; and (iii) litter/soil-dwelling insects (Figure 1). The first group includes lepidopteran larvae and larvae of hymenopteran sawflies as well as hemipterans that can degrade the canopy and affect trees through defoliation. The second group includes phloem-feeding insects, particularly **bark beetles**. Some conifer bark

Glossary

Abiotic stressor: variation in the non-living world that can negatively influence biological processes, for example, high or low temperature, strong wind or low humidity.

Bark beetle: subfamily within weevils, that is, Curculionidae, Scolytinae. The term bark beetle can be confusing as it is used taxonomically (i.e., subfamily Scolytinae) and ecologically, describing their main habitat and food source (i.e., phloem of trees). Apart from phloem-feeding/inhabiting bark beetles, other feeding modes and microhabitats are found, like xylem-inhabiting and fungus-farming species (i.e., ambrosia beetles), or species utilizing herbs, seeds, wood, pith or fungi. So far, >6300 bark beetle species have been described worldwide, with the majority living in dead/dying plants. A small number of species are able to colonize hardly stressed plants and are therefore known as pests.

Microclimate: climate variation on small scales, important for insects. For instance, the south or north side of a stone, or tree, can have dramatically different temperature and humidity.

Phenological traps: lifecycle-timing mismatches with the biotic and abiotic environment, resulting in fitness costs, worse performance, or mortality when a species does not meet its critical resources in a certain time window. Timing mismatches can result from phenological changes of players of various trophic levels and can be related to climate change.

Phenology: temporal occurrence and/or succession of lifecycle events over the course of a year; for example, dispersal, mating, or entry into dormancy.


Phenotypic plasticity: capacity of a single genotype to produce multiple phenotypes. This includes irreversible developmental changes (e.g., solitary versus gregarious locust morphs), but also reversible changes (e.g., seasonal increases in cold tolerance that decrease later).

Seasonality: habitats, for example, in temperate and polar regions are characterized by changing environmental conditions throughout a year. Organisms must adjust their lifecycles to biotic and abiotic resources to increase fitness and avoid harmful consequences.

Sensitive developmental stage: diapause induction generally occurs in species-specific lifecycle stages where

Box 2. Forest management, climate change, and diapause

Forests capture and store carbon in tree biomass, deliver societal requirements for timber, and harbor and conserve biodiversity. As 3D, long-term, and complex ecosystems, forests vary significantly from agricultural systems focused on annual row cropping. Tree canopies buffer low and high temperature extremes by filtering incoming solar radiation and regulating evapotranspiration, and the structural complexity of vegetation creates multiple microclimates that deviate from the background atmosphere (Figure 1) [103,104]. Many organisms including forest insects do not experience climate at a coarse scale, and instead live in microclimates where conditions can vary dramatically from those recorded at weather stations [105,106]. Forest microclimates among the soil, understory vegetation, tree bole, and canopy harbor different functional groups of insects and often different life stages of the same species. For example, eggs of the pine processionary moth (Box 3) are frequently deposited on lower pine branches, neonate larvae feed gregariously for a few weeks, then more mature larvae spin a silken tent and feed in the canopy for several months, after which pupae overwinter in the soil, a well buffered environment [107]. Logging in closed canopy forests removes leaf density and creates gaps, thereby changing thermal and solar radiation levels, resulting in hotter, drier, and more variable microclimates in the understory [108,109]. Mean air temperatures are higher in newly created forest-edge microclimates relative to forest interiors and soil temperatures can increase more dramatically than air following tree removal due to solar radiation [108,110]. Because even slight changes in microclimate can influence changes in individual and population-level traits [111], forest management can greatly impact insect fitness and population success. Specifically, diapause induction and termination, which are often regulated by temperature and photoperiod [1], might be altered by changes in temperature and solar radiation [112] that can occur following logging. Effects on diapause will be species- and life stage-specific as timing of diapause induction, termination, and postdiapause development vary across species. Not all microclimates will be affected equally due to the spatial and structural complexity of forest ecosystems. Tree removal is a form of localized climate change and species living in postmanagement forests face the combined forces of global climate change on top of management-induced changes in microclimate [109,113]. Microclimate regulation should be considered an important ecosystem service that is incorporated into forest management strategies in a changing climate.



Trends in Ecology & Evolution

Figure 1. (A) Mountainous conifer forest in Central Europe, consisting of Norway spruce and European larch. Forest management after infestations of the spruce bark beetle *Ips typographus*. Clear cuts result in an immediate change of abiotic conditions (Gstatterboden, Austria). (B) Trees and associated forest insects in urban and suburban landscapes face extreme abiotic conditions (e.g., drought and heat) and are exposed to artificial light at night (Rennes, France). (C) Deciduous forests differ from conifer forests in environmental conditions over various spatiotemporal scales and provide specific microclimates for forest insects (Vienna, Austria). (D) Natural, unmanaged forest ecosystems have a high structural variety with multiple microhabitats showing complexity of microclimates on a small scale (National Park Bavarian Forest, Germany). (E) Plantation of bluegum, *Eucalyptus globulus*. Environmental conditions in young stands of these highly managed forests strongly differ from buffered, heterogeneous natural forest ecosystems (Green Triangle Region, South Australia, Australia). (F) Unmanaged *Pinus ponderosa* forest in Argentina. Multiple pine species were introduced to various regions in the Southern Hemisphere where they grow in both managed plantations and unmanaged ecosystems (Patagonia, Argentina). (G) Natural stand of acacia and eucalypt where trees are intermingled with clearings exploited for grazing. The area is exposed to frequent droughts that are affecting phenology and diapause of insects (New South Wales, Australia). (H) Managed mountainous forest, consisting of western larch (*Larix occidentalis*) and other conifer species, showing the effect of seasonality on leaf fall (evergreen versus deciduous trees), which ultimately affects the climatic conditions of forest stands (Lolo National Forest, Montana, USA). (I) Managed Scots pine forest in Northern Europe during winter. Boreal forest ecosystems are often characterized by high and long-lasting snow cover, which has implications on the microclimate of insects hibernating belowground (Öby, Finland). Photos: (A–D) by Martin Schebeck, (E) by Dianne Patzel, (F) by Victoria Lantschner and Juan C. Corley, (G) by Andrea Battisti, (H) by Barbara J. Bentz, (I) by Philipp Lehmann.

an insect is perceptive to internal or external cues (see also token stimulus) to start the diapause program.


Synchronization: intrinsic or extrinsic processes that unify lifecycles so that most individuals of a population are simultaneously active and available to time their occurrence with critical resources; for example, mating partners or food.

Token stimulus: external environmental signal that is strongly correlated with a future, unfavorable seasonal event. The token stimulus itself is not adverse, but it programs an insect to enter diapause. Common token stimuli for insect diapause are photoperiod or temperature.

Voltinism: number of generations produced per year. Univoltine: one generation/year; bivoltine: two generations/year; multivoltine: more than two/year; semivoltine: lifecycles that are completed in two years or longer.

Wood ants: eusocial insects belonging to the hymenopteran family Formicidae, and the genus *Formica*. In particular, wood ants comprise species of the *Formica rufa* species group that are widely distributed in the Holarctic. The majority of species inhabit forest ecosystems and are characterized by their mound-building behavior. They fulfill pivotal functions in forests, ranging from soil bioturbation, to seed dispersal, predation of other animals, etc.

beetles of the Northern Hemisphere show eruptive population outbreaks under suitable environmental conditions, resulting in large-scale tree death [22,23]. Additionally, larval feeding of many cerambycid and buprestid beetles (many saproxylic species) initially occurs in the phloem before they proceed into woody tissues as development progresses. The third group includes **wood ants**, which are key elements of forests across the Northern Hemisphere. These social insects



Trends in Ecology & Evolution

Figure 1. Spatial and temporal microclimatic variations across contrasted forest microhabitats. Simplified conceptual overview showing (A) insects overwintering in three different microhabitats. Note that some species (e.g., soil-overwintering moth species) are restricted to a single microhabitat, while others (e.g., the spongy moth *Lymantria dispar* or the spruce bark beetle *Ips typographus*) can overwinter in multiple microhabitats. (B) Thermal variation over a certain time-span (e.g., a day) is shown as a function of tree microhabitat. This microclimatic variation can have significant impact on insect performance. (C) Effect of forest structure and type on thermal variation. (D) Highlights of some important landscape impacts on microclimate.

are abundant in multiple ecosystems and perform pivotal functions including predation on insects, bioturbation of soil or decomposition of plant biomass [24,25].

Lifecycle regulation and diapause of forest insects

Lifecycles of insects are strongly shaped by abiotic factors. Temperature affects many life-history traits, including developmental time, reproduction, survival, and **voltinism** [26]. Additionally, lifecycles of numerous forest insects are affected by dormancy, particularly diapause. Although diapause expression in canopy-dwelling, trunk-inhabiting, and soil/litter-inhabiting insects is phenotypically diverse, there are common strategies and overlapping tactics across taxa.

Both facultative and obligate diapause occur among forest insects. For example, obligate diapause is expressed in the spongy moth *Lymantria dispar* (native to Eurasia and invasive in North America) [27]. This common species of Holarctic broadleaf forests enters obligate diapause as a fully developed first-instar larva still inside its eggshell and hibernates in egg batches on trunks (or branches). Larvae hatch and begin growth the following spring when temperatures allow for development and food is available [27]. Likewise, the emerald ash borer *Agrilus planipennis* (native to Asia and now highly invasive in parts of North America and Eurasia) enters obligate diapause as late-instar larva to survive unfavorable cold conditions [28].

Furthermore, the ecologically significant group of wood ants enter an endogenously regulated dormancy during winter, regardless of environmental conditions [29,30]. Queens stop laying eggs in late summer/early fall and prepare for overwintering. Only adult females (queens and

workers) overwinter in belowground nests prior to reproduction the following spring. Although this dormancy resembles an obligate reproductive diapause, it is difficult to put these long-lived insects (particularly queens) in the standard framework of diapause classification because they enter dormancy repeatedly during their lifecycles.

Many forest insects from the aforementioned functional groups enter environmentally sensitive, facultative diapause. The most common cues for diapause induction are photoperiod (i.e., day length) and temperature, or a combination of both. Photosensitive species respond to short day-lengths in late summer/early fall to prepare for the upcoming winter and to synchronize lifecycles with the favorable growing season [1]. For example, bark beetles in the genera *Ips* and *Pityogenes* that are common in Northern Hemisphere forests enter a photoperiod-regulated diapause [3,31,32]. Defoliators of both Northern and Southern Hemisphere forests, such as the eastern spruce budworm *Choristoneura fumiferana* [33], the autumn gum moth *Mnesampela privata* [34], and some hymenopteran sawflies of the genera *Diprion* and *Neodiprion* [35] also enter diapause in response to short photoperiods.

Photoperiod changes predictably across the year, allowing organisms to anticipate seasonal low temperatures and adjust resistance, or to enter the unfavorable season in a specific developmental stage that withstands cold. Thus, photoperiod is a reliable intra- and interannual indicator for upcoming adversity. Temperature-regulated diapause is common in *Dendroctonus* bark beetles, which express facultative prepupal diapause cued by thermoperiod (seasonal and daily fluctuations of temperature) [36,37]. Diapause-driven developmental delays ensure that the most cold-resistant ontogenetic stage enters winter. Induction by temperature provides more flexibility compared with photoperiodic diapause, potentially allowing plasticity in voltinism that might track changing climates across latitudes and altitudes [38].

The process of diapause termination determines the duration of diapause and is often temperature-sensitive; many forest insects require a certain duration of cold exposure to successfully terminate diapause [31]. Diapause often ends by mid-winter, when conditions are still adverse, and is followed by postdiapause quiescence. In this state, an overwintering insect can rapidly resume development, activity, and reproduction as soon as conditions become favorable. This combined diapause/quiescence strategy is very effective at synchronizing lifecycles so that most of a population is simultaneously active and available for reproduction in spring.

Beyond low temperatures, some forest insects enter diapause to mitigate heat or drought. For example, the bag shelter moth *Ochrogaster lunifer*, feeding on leaves of acacias and eucalypts in Australia, undergoes prepupal diapause to survive the dry season [39,40]. Conifer-associated aphids of the family Adelgidae (native to the Northern Hemisphere) enter diapause as so-called sistentes, which have morphological adaptations, like heavy sclerotization, to overcome hot and dry periods [41]. For example, the hemlock woolly adelgid *Adelges tsugae* (native to Asia and Western North America, and invasive in Eastern North America where it causes mortality on eastern hemlock) prefers cool conditions in spring and fall and expresses diapause during summer to mitigate high temperatures [41,42].

Some forest insect species also show remarkable forms of diapause with implications for lifecycle regulation. For example, in several species part of a population expresses prolonged diapause beyond one year, helping to spread risks across years within a population [43–45] (Box 3). Additionally, there are examples where both facultative and obligate diapause occur within a single species [3,31,36], a strategy to respond to diverse environmental conditions across a wide geographic range.

Climate change and alterations of diapause-related environmental factors

Climate and anthropogenic change can have substantial effects on insect dormancy. In temperate regions, some of the general trends for climate change are a shortening of the duration of winter, an increase in night-time temperatures across the year, and an expansion of seasonally warm conditions [46]. Climatic variability is also increasing, particularly in autumn and spring, with potential negative effects on the timing of entry into or exit from diapause [46].

Photoperiodic cues are well known to induce diapause [1,6]. Due to climate-change-induced shifts in the phenology of some tree species, photoperiodic cues that once signaled the onset or the end of unfavorable conditions and the timing of bud break might no longer accurately reflect the timing of the onset or end of winter [47]. The duration of photoperiod that cues diapause can rapidly evolve to track changes in **seasonality** of climatic variables [47,48]. However, the extent to which rapid adaptation in photoperiodic cues for diapause will allow forest insects to mitigate the effects of climate change is a multi-faceted problem and we lack generalizable rules for which species have more or less capacity for rapid adaptation [49,50].


Adapting to novel climatic conditions is especially challenging for species with range expansions into novel habitats where they encounter new thermal and photoperiodic conditions; for example, due to introductions or release from previous low-temperature constraints during winter [51,52]. While temperatures can rapidly change due to ongoing warming, photoperiods do not, leading to asymmetries among drivers that can challenge seasonal adaptation and potentially lead to **phenological traps** [38].

Urbanization might interact with changing climates to influence seasonality and dormancy cues in forest insects in urban and suburban landscapes [53,54]. These landscapes tend to be warmer than surrounding rural areas and form urban heat islands, where trees and insects are also exposed to an excess of artificial light at night [55–57]. Forest insects inhabiting urban/suburban habitats therefore experience seasonality very differently than their rural counterparts, leading to

Box 3. Prolonged diapause in forest insects – the pine processionary moth as an outstanding example

Prolonged diapause is a long-term, bet-hedging strategy by which insects can mitigate risks posed by year-to-year variation in conditions, from temperature and rainfall to food resources or disease. The pine processionary moth (*Thaumetopoea pityocampa*) is a pine-defoliating insect typical of the Mediterranean region, although it is currently expanding north and retracting from some of the southernmost areas of its range [51]. Prolonged diapause occurs in the pupal stage. Each year, some fraction of individuals from a population enter multiyear diapause that can last as long as 7–9 years in the soil before emergence [43,44]. During prolonged diapause, metabolism is maintained at a minimum level except for short peaks evidenced by an increase of oxygen consumption and body temperature that are observed every year in spring, ~1.5 months before the expected emergence of adults [114]. After each peak, some individuals continue to ramp up metabolism and emerge as adults while others suppress their metabolism again until the following year. The mechanisms behind the ultimate duration of prolonged diapause are unknown, although fat body reserves might be a factor.

Considering the high inter-individual variation in the number of years spent in prolonged diapause within populations, including siblings of a single colony, prolonged diapause likely allows some larvae to escape lethal temperatures in the case of extreme events and mitigate local population crashes during unfavorably hot or cold years [67,107]. Larvae are gregarious and occur on trees from late summer to spring, depending on local populations and climatic conditions. Larvae spend the day inside a silk tent and feed outside the tent at night (Figure 1). Temperature is the major driver of activity as larvae feed only when they experience a temperature >9°C during day, and when night-time temperatures are above 0°C, but only if both conditions are met concurrently. If daytime and night-time temperature thresholds are not met, larvae do not feed and starve to death (although they can survive up to four weeks without feeding). Extreme cold or warm winter temperatures were found to be the best predictors of the probability of individuals entering prolonged diapause, indicating that prolonged diapause could be a general bet-hedging, or risk-spreading, mechanism in unpredictable climates [67]. The physiological mechanisms driving prolonged diapause and the associated ecological and management consequences in *T. pityocampa* and other forest insects remain poorly understood, highlighting the need for future research on this topic.



Trends in Ecology & Evolution

Figure 1. Overview of (A) canopies in a pine stand in Southern Bulgaria showing how pine processionary moth (*Thaumetopoea pityocampa*) nests are spun to be well exposed to radiation. (B) Eggs deposited on pine needles, which can be exposed to intense heat in summer, but scales are assumed to help thermoregulating/buffering the microenvironment. (C) Pine with nests. (D) Procession of larvae crawling down their host tree (band heading to suitable habitats). (E) Larvae looking for pupation and subsequent diapause sites in the soil. (F) After burying down at the end of their procession, larvae spin a silken cocoon where they pupate and undergo obligate diapause. (G) Pupa as well as its former larval exuvia without the cocoon. Photos: (A, B) by Mathieu Laparie, (C–G) by Andrea Battisti.

different sets of selective forces on seasonal dormancy [58–60]. Thus, urbanization can combine with climate change to have unexpected effects on forest insect dormancy, from urban forest insects foregoing dormancy, having truncated dormancy or undergoing more intense or longer dormancies [61–63].

Effects of climate change on performance of diapausing insects

Forests can buffer environmental variation in some microhabitats, but changing climates are likely to influence diapause performance [64]. Three central dormancy-related traits influenced by climate change are: (i) timing mismatches; (ii) direct temperature-related stress; and (iii) energetic stress.

Effects on seasonal timing have been discussed earlier and can result in mismatched timing of entry into and exit from dormancy [47,48]. Depending on the life stage, diapause includes a block [1] or a heavy suppression [65] of morphogenic and/or reproductive development. The block or suppression of development is a key feature of diapause because it protects against premature development during the favorable season (late summer and fall for winter-diapausing insects), and acts to synchronize postdormancy emergence in local populations [66]. Because accumulation of a threshold quantity of low temperature is a common driver of diapause termination, warming winters are expected to reduce the ability of some insect species or populations to accumulate adequate amounts of cold for a timely diapause termination and subsequent resumption of development [1,67]. This can lead to emergence mismatches, and will likely have multiple thus-far poorly understood effects later in the season [47].

Most dormant insects, including forest-dwelling species, increase abiotic stress tolerance as part of the diapause phenotype [4]. This includes increased resistance to low and high temperature, ice formation, and desiccation [66]. Mounting these stress responses is often costly. Therefore, insects tend to show resistance in relation to the level of stress found in a region, including a buffer or safety margin between their tolerance limits and local conditions [68,69]. Climate change can thus lead to maladapted responses through costly energy drain during dormancy, for example, when winter temperatures are warmer (see later). Snow cover also affects the dormancy performance of insects overwintering in microhabitats that are below snow. Snow cover buffers direct **abiotic stressors**, and reductions in snow depth or complete absence of snow cover can expose dormant insects to thermal variability and extreme thermal events [70]. Insects overwintering in the trunk or the canopy are less affected by changes in snow cover than are insects dormant in the leaf litter or soil; for example, wood ants, some bark beetles, and moth pupae [3,32,71]. Because forest type influences snow cover accumulation across microhabitats (forest edges vs. inner parts of stands, or continuous versus fragmented forests), overwintering in deciduous, coniferous, or mixed forests each has distinct implications for the nature of the cold stress dormant insects experience (Figure 1).

Changing climates can increase energy drain in overwintering insects because they generally do not feed during winter, regardless of the dormant stage [72,73]. Thus, the dormant phase uses energy stored prior to the onset of dormancy [72], and most insects display substantial metabolic suppression during diapause [74–76]. While many insects show a reduction in the thermal sensitivity of metabolism in their dormant state [70,77], higher temperatures nevertheless will lead to greater metabolic demand. Thus, increasing temperatures are expected to lead to severe energy drain and to reduced post-dormancy fitness [78,79]. Reduction of snow depth/cover might further exacerbate energy drain by decreasing thermal buffering [71]. Such extra energy expenditures due to warmer conditions could become particularly significant in insects undergoing prolonged diapause spanning multiple seasons and years [43,67].

Effects of climate change and diapause on phenology and voltinism

While advancing spring phenology is one of the most well-documented responses to warming climates [80], phenology is not advancing in all insects. In some species there is little evidence of any phenological change [81], whereas in others, fall phenology is altered rather than spring phenology [82]. Distinct geographic populations of the same species can differ in the directionality or magnitude of changes in seasonal timing [31,83]. Thus, the phenological alterations of forest insects with climate change can be a complex mosaic of responses [84]. Given the central role of dormancy in forest insect lifecycles, further understanding of how each of the components of dormancy responds to changing microclimates could provide generalizable rules that can be used to untangle this mosaic of responses more clearly and more accurately predict how forest insect populations will be affected by climate and other sources of anthropogenic change. Further, the small-scale microclimatic variation in forests contrasts with other terrestrial ecosystems that are less buffered; for example, agricultural systems. Comparing habitat conditions and insect responses across multiple ecosystems will be needed to develop generalizable concepts of how dormancy affects population processes across habitats.

Often acting as a central mediator of lifecycle timing, diapause can modulate phenology in two nonmutually exclusive ways: (i) there is **phenotypic plasticity** in diapause regulation with respect to temperature, thus changing temperatures affect the onset, duration and end of diapause; and (ii) heritable variation in diapause regulation evolves in response to changing climates. Documented and predicted responses of forest insect phenology to climate change are largely based on responses to temperature, though diapause is known to be highly variable among populations and rapidly evolving in insects [85–87].

Climate-change-related temperature increases can have a range of effects depending on the different phases of diapause in which thermal increases occur, consequently affecting phenology, synchrony, and voltinism [84]. Induction of photoperiod-regulated diapause can be modified by temperature, for example, in the bark beetle *Ips typographus* [3,32,88]. Temperature increase in late summer/early fall, when **sensitive developmental stages** perceive the **token stimulus**, can postpone diapause induction resulting in an extended reproductive season. Together with warmer spring conditions and an earlier onset of activity, this lengthening of the growing season and shrinking of the dormant season will likely increase voltinism. For example, the bark beetle *Dendroctonus rufipennis* has a temperature-regulated facultative prepupal diapause that is induced by low temperatures in late summer [3,37]. Diapause can be averted under warm conditions which ultimately results in a shorter lifecycle and a shift from semivoltinism to univoltinism [89].

Photoperiod-mediated induction or termination of diapause often occurs over a small developmental window, a sensitive developmental stage, and changes in phenology caused by changing temperatures can disrupt developmental synchrony of the sensitive stage from the photoperiodic cue, resulting in altered phenological timing and consequently changes in voltinism [1,3]. Because many forest insects must experience some period of cold to terminate diapause, warmer winters might postpone diapause termination. Therefore, warming temperatures could actually delay spring phenology, or possibly lead to no change if the delay is offset by more rapid post-diapause development [90].

As mentioned earlier, climate change can cause phenological mismatches. For example, the timing of emergence from larval diapause does not change with increasing temperature in the spruce budworm *C. fumiferana*, possibly facilitating a shift from the primary host to another secondary host tree because budburst is temperature sensitive in the primary host [91].

Defoliating moths including the winter moth (*Operophtera brumata*) experience changes in phenology, population dynamics, and geographic range due to changes in winter diapause termination and synchrony with food sources the following spring [92–94]. The larch bud moth (*Zeiraphera griseana*) requires **synchronization** between hatching of overwintering eggs and flushing of the larch needles they feed on [95,96]. Because the lack of cold during winter is altering the timing of egg hatch more than leaf flush, larch bud moth populations are shifting to elevations where synchronization still is possible [95]. Once the tree line has been reached, however, finding suitable trees will be less likely, unless a shift in the tree line to higher elevations also occurs. The latter will take decades, and therefore local extinctions of the larch bud moth seem likely. Other voltinism-related phenological mismatches can occur when certain developmental stages that are not suitable to cold enter the winter season [38,46]. For example, although an extended reproductive season increases voltinism in the bark beetle *I. typographus*, preimaginal stages (which are less resistant to cold than adults) that do not become adults in time will suffer from cold-related mortality over a wide part of the range of the species [3,32]. Similarly, *Dendroctonus* bark beetles synchronize their lifecycles such that the most cold-hardy developmental stage enters winter. Thus, climate change-related phenological shifts can result in increased overwinter mortality if conditions deteriorate while the insect is in a less hardy stage [3,97].

Finally, a few forest insect species exhibit potentially heritable, and therefore evolvable variation in diapause strategies. For example, populations of the bark beetles *I. typographus*, *Dendroctonus simplex*, *Dendroctonus ponderosae*, and *D. rufipennis* vary across a range of environmental sensitivity from effectively obligate to highly facultative diapause [31,36,98]. An extended favorable season for reproduction and development due to higher temperatures might increase the proportion of facultative diapausing lineages within and across populations over obligate diapausing conspecifics [1]. This could result in evolutionary changes in environmental sensitivity of dormancy, possibly accompanied by alterations in voltinism.

Concluding remarks

Forest insect dormancies have both plastic and adaptive potential and are responding to changing environmental conditions, with significant impacts on forest ecosystems, including carbon sequestration [99]. Insects are a tremendously successful group of animals occupying all continents on Earth and show remarkable responses for adapting to climate change. Probably due to the strong selection imposed by mortality in individuals making poor timing decisions in seasonal environments, evolutionary pressures on entry into and exit from diapause include some of the most famous cases of rapid adaptation known, including responses to climate change and urbanization [60,100]. Large-scale range shifts are also becoming ever clearer, with strong associations between dormancy type and extent of range expansion [101,102]. In addition, local adaptation in physiological traits, including metabolic suppression and tolerance to low temperatures, are apparent in forest insects; indicating high potential for rapid evolution and strong selection on dormancy-related traits [64,87]. As such, insects clearly possess the capacity to respond to changing seasonality and an understanding of how diapause tactics are affected by environmental parameters is needed to predict potential impacts of insects on forests under future climates (see Outstanding questions).

Acknowledgments

M.S. was financially supported by the Austrian Ministry of Agriculture, Forestry, Regions and Water Management ('Waldfonds' projects 'NewIPS', grant no. 101686, 'IpsEMAN', grant no. 101687). P.L. was funded by the Swedish Research Council (grants no. 2017-04159 and 2022-03343), the Research Council Formas (grant no. 2017-00965) and the Carl Tryggers Stiftelse (grant no. CTS20-242). D.A.H. was supported by Florida Agricultural Experiment Station (Hatch project FLA-ENY-005943). A.B. was supported by Bolzano/Bozen district grant 'Forest Insects' and UNIPD DOR 2344584.

Outstanding questions

Although still difficult to collect, would microclimatic data with fine spatiotemporal resolution improve our estimates of dormancy performance; for example, phenology or winter survival, of forest insects?

Climate change can result in an asymmetry in abiotic conditions regulating diapause, for example, temperatures in a given habitat might change while photoperiod does not. Do photoperiod-regulated traits evolve faster than temperature responses in forest insects? Can we use data on the asymmetry of different drivers to predict range changes?

Will information on phenotypic plasticity and adaptive potential of diapause responses allow us to understand the effects of seasonality on insect performance in a changing climate?

Whether urban or rural, managed or unmanaged forests, insects face a broad spectrum of diapause-regulating conditions and diapause responses might consequently vary depending on forest type. Can we use information on diapause tactics and diapause responses of insects across habitats to get insights into possible adaptations to different environments?

Some species do not fit into the standard framework of insect diapause; for example, eusocial insects, like ant queens of some *Formica* species that live >20 years and go through dormant and reproductive cycles multiple times during their life. How are diapause and/or dormancy regulated in long-lived eusocial insects and how does this affect the response of perennially long-lived insects to seasonally changing environments?

Our current understanding of forest insect dormancy is based on economically relevant species from the Northern Hemisphere. Can we infer generalizable rules and concepts about forest insect diapause by studying species from numerous taxa and functional groups across a diversity of ecosystems worldwide?

Declaration of interests

No interests are declared.

References

- Denlinger, D.L. (2022) *Insect Diapause*, Cambridge University Press
- Pureswaran, D.S. *et al.* (2018) Forest insects and climate change. *Curr. For. Rep.* 4, 35–50
- Schebeck, M. *et al.* (2017) Diapause and overwintering of two spruce bark beetle species. *Physiol. Entomol.* 42, 200–210
- Wilsterman, K. *et al.* (2021) A unifying, eco-physiological framework for animal dormancy. *Funct. Ecol.* 35, 11–31
- Kostal, V. (2006) Eco-physiological phases of insect diapause. *J. Insect Physiol.* 52, 113–127
- Numata, H. and Shintani, Y. (2023) Diapause in univoltine and semivoltine life cycles. *Annu. Rev. Entomol.* 68, 257–276
- Poitou, L. *et al.* (2020) Diapause regulation in newly invaded environments: termination timing allows matching novel climatic constraints in the Box Tree Moth, *Cydalima perspectalis* (Lepidoptera: Crambidae). *Insects* 11, 629
- De Frenne, P. *et al.* (2021) Forest microclimates and climate change: importance, drivers and future research agenda. *Glob. Chang. Biol.* 27, 2279–2297
- Richter, R. *et al.* (2022) Tree species matter for forest microclimate regulation during the drought year 2018: disentangling environmental drivers and biotic drivers. *Sci. Rep.* 12, 17559
- Parker, G.G. and Brown, M.J. (2000) Forest canopy stratification – is it useful? *Am. Nat.* 155, 473–484
- Parker, G.G. (1995) Structure and microclimate of forest canopies. In *Forest Canopies* (Lowman, M.D. and Nadkarni, N.M., eds), pp. 73–106, Academic Press
- Didham, R.K. and Ewers, R.M. (2014) Edge effects disrupt vertical stratification of microclimate in a temperate forest canopy. *Pac. Sci.* 68, 493–508
- Ulyshen, M.D. (2011) Arthropod vertical stratification in temperate deciduous forests: implications for conservation-oriented management. *For. Ecol. Manag.* 261, 1479–1489
- Geiger, R. *et al.* (1995) Forest climatology. In *The Climate Near the Ground* (Geiger, R. *et al.*, eds), pp. 259–325, Vieweg+Teubner Verlag Wiesbaden
- May, M.L. (1979) Insect thermoregulation. *Annu. Rev. Entomol.* 24, 313–349
- Röhrig, E. and Ulrich, B. (1991) *Temperate Deciduous Forests*, Elsevier
- Nadkarni, N.M. *et al.* (2004) The nature of forest canopies. In *Forest Canopies* (Lowman, M. and Rinker, H.B., eds), pp. 3–23, Elsevier Academic Press
- Decocq, G. *et al.* (2004) Plant diversity in a managed temperate deciduous forest: understorey response to two silvicultural systems. *J. Appl. Ecol.* 41, 1065–1079
- Brockhoff, E.G. *et al.* (2017) Forest biodiversity, ecosystem functioning and the provision of ecosystem services. *Biodivers. Conserv.* 26, 3005–3035
- Crespo-Pérez, V. *et al.* (2020) The importance of insects on land and in water: a tropical view. *Curr. Opin. Insect Sci.* 40, 31–38
- Boyd, I.L. *et al.* (2013) The consequence of tree pests and diseases for ecosystem services. *Science* 342, 1235–1237
- Biedermann, P.H.W. *et al.* (2019) Bark beetle population dynamics in the Anthropocene: challenges and solutions. *Trends Ecol. Evol.* 34, 10
- Hlásny, T. *et al.* (2021) Bark beetle outbreaks in Europe: state of knowledge and ways forward for management. *Curr. For. Rep.* 7, 138–165
- Trigos-Peral, G. *et al.* (2021) Wood ants as biological control of the forest pest beetles *Ips* spp. *Sci. Rep.* 11, 17931
- Stockan, J.A. and Robinson, E.J. (2016) *Wood Ant Ecology and Conservation*, Cambridge University Press
- Harvey, J.A. *et al.* (2020) Climate change-mediated temperature extremes and insects: from outbreaks to breakdowns. *Glob. Chang. Biol.* 26, 6685–6701
- Boukouvaa, M.C. *et al.* (2022) *Lymantria dispar* (L.) (Lepidoptera: Erebidae): current status of biology, ecology, and management in Europe with notes from North America. *Insects* 13, 854
- Duan, J.J. *et al.* (2021) Effects of low temperature exposure on diapause, development, and reproductive fitness of the emerald ash borer (Coleoptera: Buprestidae): implications for voltinism and laboratory rearing. *J. Econ. Entomol.* 114, 201–208
- Lopatina, E.B. (2018) Structure, diversity and adaptive traits of seasonal cycles and strategies in ants. In *The Complex World of Ants* (Shields, V., ed.), pp. 7–49, InTech
- Kipyatkov, V.E. (2001) Seasonal life cycles and the forms of dormancy in ants (Hymenoptera: Formicoidea). *Acta Soc. Zool. Bohem.* 65, 211–238
- Schebeck, M. *et al.* (2022) Facultative and obligate diapause phenotypes in populations of the European spruce bark beetle *Ips typographus*. *J. Pest. Sci.* 95, 889–899
- Schebeck, M. *et al.* (2023) Evolutionary ecology of the bark beetles *Ips typographus* and *Pityogenes chalcographus*. *Bull. Entomol. Res.* 113, 1–10
- Marshall, K.E. and Roe, A.D. (2021) Surviving in a frozen forest: the physiology of eastern spruce budworm overwintering. *Physiology* 36, 174–182
- Steinbauer, M.J. *et al.* (2004) Modelling a forest lepidopteran: phenological plasticity determines voltinism which influences population dynamics. *Forest Ecol. Manag.* 198, 117–131
- Davis, J.S. *et al.* (2023) Identification, biology, and management of conifer sawflies (Hymenoptera: Diprioninae) in eastern North America. *J. Integr. Pest. Manag.* 14, 13
- Bentz, B.J. and Hansen, E.M. (2018) Evidence for a prepupal diapause in the Mountain Pine Beetle (*Dendroctonus ponderosae*). *Environ. Entomol.* 47, 175–183
- Hansen, E.M. *et al.* (2011) Prepupal diapause and instar IV developmental rates of the spruce beetle, *Dendroctonus rufipennis* (Coleoptera: Curculionidae, Scolytinae). *J. Insect Physiol.* 57, 1347–1357
- Van Dyck, H. *et al.* (2015) The lost generation hypothesis: could climate change drive ectotherms into a developmental trap? *Oikos* 124, 54–61
- Floater, G.J. and Zalucki, M.P. (1999) Life tables of the processionary caterpillar *Ochrogaster lunifer* Herrich-Schaffer (Lepidoptera: Thaumetopoeidae) at local and regional scales. *Aust. J. Entomol.* 38, 330–339
- Floater, G.J. (2001) Habitat complexity, spatial interference, and “minimum risk distribution”: a framework for population stability. *Ecol. Monogr.* 71, 447–468
- Havill, N.P. and Footitt, R.G. (2007) Biology and evolution of Adelgidae. *Annu. Rev. Entomol.* 52, 325–349
- Salom, S.M. *et al.* (2001) Evaluation of aestival diapause in hemlock woolly adelgid (Homoptera: Adelgidae). *Environ. Entomol.* 30, 877–882
- Salman, M.H.R. *et al.* (2016) Prolonged pupal diapause drives population dynamics of the pine processionary moth (*Thaumetopoea pityocampa*) in an outbreak expansion area. *Forest Ecol. Manag.* 361, 375–381
- Halperin, J. (1990) Life history of *Thaumetopoea* spp. (Lep., Thaumetopoeidae) in Israel. *J. Appl. Entomol.* 110, 1–6
- Corley, J.C. and Villacide, J.M. (2012) Population dynamics of *Sirex noctilio*: influence of diapause, spatial aggregation and flight potential on outbreaks and spread. In *The Sirex Woodwasp and Its Fungal Symbiont: Research and Management of a Worldwide Invasive Pest* (Slippers, B. *et al.*, eds), pp. 51–64, Springer
- Marshall, K.E. *et al.* (2020) Evolutionary impacts of winter climate change on insects. *Curr. Opin. Insect Sci.* 41, 54–62
- Visser, M.E. and Gienapp, P. (2019) Evolutionary and demographic consequences of phenological mismatches. *Nat. Ecol. Evol.* 3, 879–885
- Ittonen, M. *et al.* (2022) Local adaptation to seasonal cues at the fronts of two parallel, climate-induced butterfly range expansions. *Ecol. Lett.* 25, 2022–2033
- Hoffmann, A.A. and Sgrò, C.M. (2011) Climate change and evolutionary adaptation. *Nature* 470, 479–485

50. Franks, S.J. and Hoffmann, A.A. (2012) Genetics of climate change adaptation. *Annu. Rev. Genet.* 46, 185–208
51. Bourougaoui, A. *et al.* (2021) Has North Africa turned too warm for a Mediterranean forest pest because of climate change? *Clim. Chang.* 165, 46
52. Tobin, P.C. *et al.* (2014) The influence of climate change on insect invasions in temperate forest ecosystems. In *Challenges and Opportunities for the World's Forests in the 21st Century* (Fenning, T., ed.), pp. 267–293, Springer
53. Lehmann, P. *et al.* (2020) Complex responses of global insect pests to climate warming. *Front. Ecol. Environ.* 18, 141–149
54. Backe, K. *et al.* (2021) Human health risks of invasive caterpillars increase with urban warming. *Landsc. Ecol.* 36, 1475–1487
55. Frank, S.D. and Backe, K.M. (2023) Effects of urban heat islands on temperate forest trees and arthropods. *Curr. For. Rep.* 9, 48–57
56. Lahr, E.C. *et al.* (2018) Getting ahead of the curve: cities as surrogates for global change. *Proc. R. Soc. B Biol. Sci.* 285, 20180643
57. Boyes, D.H. *et al.* (2021) Is light pollution driving moth population declines? A review of causal mechanisms across the life cycle. *Insect Conserv. Divers.* 14, 167–187
58. Deitsch, J.F. and Kaiser, S.A. (2023) Artificial light at night increases top-down pressure on caterpillars: experimental evidence from a light-naïve forest. *Proc. R. Soc. B Biol. Sci.* 290, 20230153
59. Lockett, M.T. *et al.* (2022) Artificial light at night promotes bottom-up changes in a woodland food chain. *Environ. Pollut.* 310, 119803
60. Merckx, T. *et al.* (2021) Urbanization extends flight phenology and leads to local adaptation of seasonal plasticity in Lepidoptera. *Proc. Natl. Acad. Sci. U. S. A.* 118, e2106006118
61. Merckx, T. *et al.* (2023) Dim light pollution prevents diapause induction in urban and rural moths. *J. Appl. Ecol.* 60, 1022–1031
62. Mukai, A. *et al.* (2021) Urban warming and artificial light alter dormancy in the flesh fly. *R. Soc. Open Sci.* 8, 210866
63. Fyfe, L.R. *et al.* (2021) Artificial light at night alters the seasonal responses of biting mosquitoes. *J. Insect Physiol.* 129, 104194
64. Williams, C.M. *et al.* (2015) A cross-seasonal perspective on local adaptation: metabolic plasticity mediates responses to winter in a thermal-generalist moth. *Funct. Ecol.* 29, 549–561
65. Dowie, E.J. *et al.* (2020) Genome-wide variation and transcriptional changes in diverse developmental processes underlie the rapid evolution of seasonal adaptation. *Proc. Natl. Acad. Sci. U. S. A.* 117, 23960–23969
66. Tougeron, K. (2019) Diapause research in insects: historical review and recent work perspectives. *Entomol. Exp. Appl.* 167, 27–36
67. Salzman, M.H.R. *et al.* (2019) Winter temperature predicts prolonged diapause in pine processionary moth species across their geographic range. *PeerJ* 7, e6530
68. Bale, J.S. and Hayward, S.A.L. (2010) Insect overwintering in a changing climate. *J. Exp. Biol.* 213, 980–994
69. Sinclair, B.J. *et al.* (2015) An invitation to measure insect cold tolerance: methods, approaches, and workflow. *J. Therm. Biol.* 53, 180–197
70. Roberts, K.T. and Williams, C.M. (2022) The impact of metabolic plasticity on winter energy use models. *J. Exp. Biol.* 225, jeb243422
71. Williams, C.M. *et al.* (2015) Cold truths: how winter drives responses of terrestrial organisms to climate change. *Biol. Rev.* 90, 214–235
72. Short, C.A. and Hahn, D.A. (2023) Fat enough for the winter? Does nutritional status affect diapause? *J. Insect Physiol.* 145, 104488
73. Roberts, K.T. *et al.* (2023) Seasonal energetics: are insects constrained by energy during dormancy? *J. Exp. Biol.* 226, jeb245782
74. Leberzon, J.E. *et al.* (2022) Reversible mitophagy drives metabolic suppression in diapausing beetles. *Proc. Natl. Acad. Sci. U. S. A.* 119, e2201089119
75. Hahn, D.A. and Denlinger, D.L. (2011) Energetics of insect diapause. *Annu. Rev. Entomol.* 56, 103–121
76. Hahn, D.A. and Denlinger, D.L. (2007) Meeting the energetic demands of insect diapause: nutrient storage and utilization. *J. Insect Physiol.* 53, 760–773
77. Powell, T.H.Q. *et al.* (2020) A rapidly evolved shift in life-history timing during ecological speciation is driven by the transition between developmental phases. *J. Evol. Biol.* 33, 1371–1386
78. Lindestad, O. *et al.* (2020) Variation in butterfly diapause duration in relation to voltinism suggests adaptation to autumn warmth, not winter cold. *Funct. Ecol.* 34, 1029–1040
79. Nielsen, M.E. *et al.* (2022) Longer and warmer prewinter periods reduce post-winter fitness in a diapausing insect. *Funct. Ecol.* 36, 1151–1162
80. Forister, M.L. and Shapiro, A.M. (2003) Climatic trends and advancing spring flight of butterflies in lowland California. *Glob. Chang. Biol.* 9, 1130–1135
81. Thackeray, S.J. *et al.* (2016) Phenological sensitivity to climate across taxa and trophic levels. *Nature* 535, 241–245
82. Karlsson, B. (2014) Extended season for northern butterflies. *Int. J. Biometeorol.* 58, 691–701
83. Kerr, N.Z. *et al.* (2020) Developmental trap or demographic bonanza? Opposing consequences of earlier phenology in a changing climate for a multivoltine butterfly. *Glob. Chang. Biol.* 26, 2014–2027
84. Bentz, B.J. *et al.* (2022) Complexities in predicting mountain pine beetle and spruce beetle response to climate change. In *Bark Beetle Management, Ecology, and Climate Change* (Gandhi, J.K. and Hofstetter, R.W., eds), pp. 31–54, Academic Press
85. Ragland, G.J. *et al.* (2019) Evolutionary and functional genetics of insect diapause: a call for greater integration. *Curr. Opin. Insect Sci.* 36, 74–81
86. Armbruster, P.A. (2016) Photoperiodic diapause and the establishment of *Aedes albopictus* (Diptera: Culicidae) in North America. *J. Med. Entomol.* 53, 1013–1023
87. Batz, Z.A. *et al.* (2020) Rapid adaptive evolution of the diapause program during range expansion of an invasive mosquito. *Evolution* 74, 1451–1465
88. Dolezal, P. and Sehna, F. (2007) Effects of photoperiod and temperature on the development and diapause of the bark beetle *Ips typographus*. *J. Appl. Entomol.* 131, 165–173
89. Hansen, E.M. *et al.* (2001) Physiological basis for flexible voltinism in the spruce beetle (Coleoptera: Scolytidae). *Can. Entomol.* 133, 805–817
90. Forrest, J.R.K. (2016) Complex responses of insect phenology to climate change. *Curr. Opin. Insect Sci.* 17, 49–54
91. Pureswaran, D.S. *et al.* (2019) Phenological synchrony between eastern spruce budworm and its host trees increases with warmer temperatures in the boreal forest. *Ecol. Evol.* 9, 576–586
92. Ammoun, T. *et al.* (2012) Invading and resident defoliators in a changing climate: cold tolerance and predictions concerning extreme winter cold as a range-limiting factor. *Ecol. Entomol.* 37, 212–220
93. Jepsen, J.U. *et al.* (2008) Climate change and outbreaks of the geometrids *Operophtera brumata* and *Epirrita autumnata* in subarctic birch forest: evidence of a recent outbreak range expansion. *J. Anim. Ecol.* 77, 257–264
94. van Dis, N.E. *et al.* (2024) Temperature has an overriding role compared to photoperiod in regulating the seasonal timing of winter moth egg hatching. *Oecologia* 204, 734–750
95. Rozenberg, P. *et al.* (2020) Direct and indirect analysis of the elevational shift of larch budmoth outbreaks along an elevation gradient. *Front. For. Glob. Chang.* 3, 86
96. Saulnier, M. *et al.* (2017) Spatiotemporal heterogeneity of larch budmoth outbreaks in the French Alps over the last 500 years. *Can. J. For. Res.* 47, 667–680
97. Bentz, B.J. and Powell, J.A. (2014) Mountain Pine Beetle seasonal timing and constraints to bivoltinism (a comment on Mitton and Ferrenberg, "Mountain Pine Beetle develops an unprecedented summer generation in response to climate warming"). *Am. Nat.* 184, 787–796
98. McKee, F.R. and Aukema, B.H. (2015) Successful reproduction by the eastern larch beetle (Coleoptera: Curculionidae) in the absence of an overwintering period. *Can. Entomol.* 147, 602–610
99. Albrich, K. *et al.* (2023) From sink to source: changing climate and disturbance regimes could tip the 21st century carbon balance of an unmanaged mountain forest landscape. *Forestry* 96, 399–409
100. Bradshaw, W.E. and Holzapfel, C.M. (2001) Genetic shift in photoperiodic response correlated with global warming. *Proc. Natl. Acad. Sci. U. S. A.* 98, 14509–14511

101. Pöyry, J. *et al.* (2009) Species traits explain recent range shifts of Finnish butterflies. *Glob. Chang. Biol.* 15, 732–743
102. Sunde, J. *et al.* (2023) Century-long butterfly range expansions in northern Europe depend on climate, land use and species traits. *Commun. Biol.* 6, 601
103. Williams, J.J. and Newbold, T. (2020) Local climatic changes affect biodiversity responses to land use: a review. *Divers. Distrib.* 26, 76–92
104. von Arx, G. *et al.* (2013) Microclimate in forests with varying leaf area index and soil moisture: potential implications for seedling establishment in a changing climate. *J. Ecol.* 101, 1201–1213
105. Moise, E.R.D. *et al.* (2023) Suboptimal host tree benefits the overwintering of a destructive forest insect pest. *Basic Appl. Ecol.* 71, 72–84
106. Pincebourde, S. and Woods, H.A. (2020) There is plenty of room at the bottom: microclimates drive insect vulnerability to climate change. *Curr. Opin. Insect Sci.* 41, 63–70
107. Battisti, A. *et al.* (2005) Expansion of geographic range in the pine processionary moth caused by increased winter temperatures. *Ecol. Appl.* 15, 2084–2096
108. Tuff, K.T. *et al.* (2016) A framework for integrating thermal biology into fragmentation research. *Ecol. Lett.* 19, 361–374
109. Boyle, M.J.W. *et al.* (2021) Localised climate change defines ant communities in human-modified tropical landscapes. *Funct. Ecol.* 35, 1094–1108
110. Kautz, M. *et al.* (2013) The "sun-effect": microclimatic alterations predispose forest edges to bark beetle infestations. *Eur. J. Forest Res.* 132, 453–465
111. Greiser, C. *et al.* (2022) Microclimatic variation affects developmental phenology, synchrony and voltinism in an insect population. *Funct. Ecol.* 36, 3036–3048
112. Axelsen, J.A. *et al.* (1997) Global solar radiation as the factor controlling induction of diapause in the pod midge (*Dasyneura brassicae* Winn). *Oecologia* 111, 178–182
113. Barahona-Segovia, R.M. *et al.* (2019) Anthropogenic thermal gradient in managed landscapes determines physiological performance and explains the edge-biased distribution of ectothermic arthropods. *Forest. Ecol. Manag.* 440, 147–157
114. Salman, M.H.R. *et al.* (2019) Termination of pupal diapause in the pine processionary moth. *Physiol. Entomol.* 44, 53–59