

Pantropical tree rings show small effects of drought on stem growth

Pieter A. Zuidema^{1*}, Peter Groenendijk², Mizanur Rahman³, Valerie Trouet^{4,5}, Abrham Abiyu⁶, Rodolfo Acuña-Soto^{7,8}, Eduardo Adenesky-Filho⁹, Raquel Alfaro-Sánchez^{10,11}, Claudio Roberto Anholetto Jr¹², José Roberto Vieira Aragão^{13,2}, Gabriel Assis-Pereira^{14,15}, Claudia C. Astudillo-Sánchez¹⁶, Ana Carolina Barbosa¹⁷, Giovanna Battipaglia¹⁸, Hans Beeckman¹⁹, Paulo Cesar Botosso²⁰, Nils Bourland¹⁹, Achim Bräuning²¹, Roel Brienen²², Matthew Brookhouse²³, Supaporn Buajan²⁴, Brendan M. Buckley²⁵, J. Julio Camarero²⁶, Artemio Carrillo-Parra²⁷, Gregório Ceccantini²⁸, Librado R. Centeno-Erguera²⁹, Julián Cerano-Paredes²⁹, Rosalinda Cervantes-Martínez³⁰, Wirong Chanthorn³¹, Ya-Jun Chen³², Bruno Barçante Ladavat Cintra^{33,34}, Eladio Heriberto Cornejo-Oviedo³⁰, Otoniel Cortés-Cortés³⁰, Clayane Matos Costa³⁵, Camille Couralet¹⁹, Doris Bianca Crispín-De-La-Cruz^{2,36}, Rosanne D'arrigo²⁵, Diego A. David³⁷, Maaïke De Ridder¹⁹, Jorge Ignacio Del Valle³⁷, Mário Dobner Jr³⁸, Jean-Louis Doucet³⁹, Oliver Dünisch⁴⁰, Brian J. Enquist^{41,42}, Karin Esemann-Quadros^{43,44}, Gerardo Esquivel-Arriaga²⁹, Ze-Xin Fan³², Adeline Fayolle^{45,46}, Tatiele Anete Bergamo Fenilli⁹, M. Eugenia Ferrero^{47,36}, Esther Fichtler⁴⁸, Patrick M. Finnegan⁴⁹, Claudia Fontana¹⁴, Kainana S. Francisco⁵⁰, Pei-Li Fu³², Franklin Galvão⁵¹, Aster Gebrekirstos⁵², Jorge A. Giraldo⁵³, Emanuel Gloor²², Milena Godoy-Veiga²⁸, Daniela Granato-Souza⁵⁴, Anthony Guerra⁵⁵, Kristof Haneca⁵⁶, Grant Logan Harley⁵⁷, Ingo Heinrich^{58,59}, Gerhard Helle⁶⁰, Bruna Hornink^{2,14}, Wannes Hubau⁶¹, Janet G. Inga³⁶, Mahmuda Islam^{3,62}, Yu-Mei Jiang⁶³, Mark Kaib⁶⁴, Zakia Hassan Khamisi⁴, Marcin Koprowski^{65,66}, Eva Layme^{67,68}, A. Joshua Leffler⁶⁹, Gauthier Ligot³⁹, Claudio Sergio Lisi³⁵, Neil J. Loader⁷⁰, Francisco De Almeida Lobo⁷¹, Giuliano Maselli Locosselli^{28,72}, Tomaz Longhi-Santos⁷³, Lidio Lopez⁴⁷, María I. López-Hernández³⁰, José Luís Penetra Cerveira Lousada⁷⁴, Rubén D. Manzanedo^{75,76}, Amanda K. Marcon⁵¹, Justin T. Maxwell⁷⁷, Omar N. Mendoza-Villa⁷⁸, Ítallo Romany Nunes Menezes^{35,79}, Mulugeta Mokria^{21,52}, Valdiney Ribeiro Montóia⁸⁰, Eddy Moors^{81,82}, Miyaer Moreno⁸³, Miguel Angel Muñoz-Castro⁸⁴, Cristina Nabais⁸⁵, Anuttara Nathalang⁸⁶, Justine Ngoma⁸⁷, Francisco De Carvalho Nogueira Jr⁸⁸, Juliano Moraes Oliveira^{89,90}, Gabriela Morais Olmedo⁹¹, Daigard Ricardo Ortega-Rodriguez¹⁴, Carmen Eugenia Rodríguez Ortiz⁹², Mariana Alves Pagotto³⁵, Shankar Panthi³², Kathelyn Paredes-Villanueva^{93,94}, Gonzalo Pérez-de-Lis⁹⁵, Laura Patricia Ponce Calderón⁹⁶, Leif Armando Portal-Cahuana^{97,14}, Darwin Alexander Pucha-Cofrep⁹⁸, Nathsuda Pumijumnong²⁴, Paulo Quadri⁹⁹, Jorge Andrés Ramírez¹⁰⁰, Edilson Jimmy Requena-Rojas³⁶, Judith Reyes-Flores³⁰, Adauto de Souza Ribeiro³⁵, Iain Robertson⁷⁰, Fidel Alejandro Roig^{47,101}, José Guilherme Roquette¹⁰², Ernesto Alonso Rubio-Camacho¹⁰³, Raúl Sánchez-Salguero¹⁰⁴, Ute Sass-Klaassen^{1,105}, Jochen Schöngart¹⁰⁶, Marcelo Callegari Scipioni³⁸, Paul R. Sheppard⁴, Lucas C.R. Silva¹⁰⁷, Franziska Slotta¹⁰⁸, Leroy Soria-Díaz¹⁰⁹, Luciana K.V.S. Sousa¹¹⁰, James H. Speer¹¹¹, Matthew D. Therrell¹¹², Ginette Ticse-Otarola^{36,113}, Mario Tomazello-Filho¹⁴, Max C.A. Torbenson¹¹⁴, Pantana Tor-Ngern^{115,116}, Ramzi Touchan⁴, Jan Van den Bulcke⁶¹, Lorenzo Vázquez-Selem¹¹⁷, Adin H. Velázquez-Pérez¹¹⁸, Alejandro Venegas-González¹¹⁹, Ricardo Villalba⁴⁷, Jose Villanueva-Díaz²⁹, Mart Vlam¹⁰⁵, George Vourlitis¹²⁰, Christian Wehenkel²⁷, Tommy Wils^{121,122}, Erika S. Zavaleta¹²³, Eshetu Asfaw Zewdu¹²⁴, Yong-Jiang Zhang¹²⁵, Zhe-Kun Zhou³², Flurin Babst^{126,4}

Increasing drought pressure under anthropogenic climate change may jeopardize the potential of tropical forests to capture carbon in woody biomass and act as a long-term carbon dioxide sink. To evaluate this risk, we assessed drought impacts in 483 tree-ring chronologies from across the tropics and found an overall modest stem growth decline (2.5% with a 95% confidence interval of 2.2 to 2.7%) during the 10% driest years since 1930. Stem growth declines exceeded 10% in 25% of cases and were larger at hotter and drier sites and for gymnosperms compared with angiosperms. Growth declines generally did not outlast drought years and were partially mitigated by growth stimulation in wet years. Thus, pantropical forest carbon sequestration through stem growth has hitherto shown drought resilience that may, however, diminish under future climate change.

Tropical forests and woodlands are key components of the global carbon cycle. They represent much of the carbon stocks in terrestrial vegetation, contribute strongly to the land carbon sink, and are a key driver of the interannual variation in this sink (1, 2). The long-term capture and storage of carbon in tropical woody biomass has a high potential to contribute to nature-based solutions to climate change by acting as a CO₂ sink (3, 4). However, the increasing incidence and intensity of droughts may fundamentally alter these services (5–8), temporarily shifting tropical vegetation to a net carbon source (5, 7, 9) and thus jeopardizing its role in climate change mitigation (10). Droughts can be caused by low precipitation (P), high atmospheric water demand (vapor pressure deficit, VPD), or both simultaneously, leading to a strong climatic water deficit (CWD), calculated as water demand minus supply. Each of these drought types may reduce tropical tree growth.

To understand and predict the risks that droughts pose for the long-term capture of carbon in tropical vegetation, a pantropical assessment of drought effects on stem growth is needed. However, studies of drought effects on tree growth are scarce and poorly replicated spatially (11–13). This situation limits the ability to generalize and contextualize local empirical findings and to validate terrestrial biosphere models, which currently poorly represent the formation (14, 15) and climate sensitivity of tropical woody biomass (16). These modeling

uncertainties are bound to persist unless they are constrained by extensive empirical studies.

Here, we leveraged the recent expansion of tropical tree-ring studies (17, 18) to assess drought effects on stem growth at an annual resolution and over multidecadal time scales. We assembled the most extensive pantropical tree-ring network to date, extending across all tropical climates, to answer the following questions: (i) To what extent is stem growth reduced during drought years and does this differ across drought types (i.e., low P, high VPD, and high CWD), seasons (i.e., dry versus wet), and major clades (i.e., angiosperms versus gymnosperms)? (ii) Is there evidence for lagged drought effects on stem growth and, if so, how strong or persistent are these effects? (iii) To what extent does growth stimulation during wet extremes compensate for stem growth reduction during droughts? (iv) Does the magnitude of drought impacts depend on local climatic conditions?

On the basis of current knowledge about growth-limiting factors and results from extratropical tree-ring studies, we hypothesize drought effects to be stronger for CWD droughts than for P and VPD droughts and for gymnosperms than for angiosperms [question (i) (19, 20)]; lag effects to exist, especially for gymnosperms [question (ii) (19, 21)]; drought-induced growth declines to be partially compensated for by growth stimulation during wet extremes [question (iii) (22, 23)]; and drought effects to be stronger in more arid regions [question (iv) (19, 23)].

Pantropical tree-ring network

We compiled a network of 483 tree-ring width chronologies from >10,000 trees spanning tropical and subtropical latitudes (30°N to 30°S). This dataset comprises 163 species and 33 plant families, with similar proportions of angiosperm and gymnosperm chronologies (Fig. 1A and fig. S1). Chronologies were redeveloped from >20,000 time series of raw tree-ring width measurements using a single, flexible detrending procedure that retains short-term growth responses to climate extremes. The resulting ring-width index (RWI) chronologies represent a relative, population-level measure of tree-growth variability. For each site, we identified the 10% driest years since 1930 in terms of P (lowest values), VPD (highest values), and CWD (highest values)

in either the wet or dry season. CWD was calculated as cumulative abs(P-PET) for months during the season when the potential evapotranspiration (PET) is greater than P. We studied droughts at the seasonal level because the climate sensitivity of tropical tree growth differs among seasons (17). We quantified the impacts of droughts (and wet extremes) on RWI using superposed epoch analysis (SEA) (24), a technique that compares the mean RWI anomaly in extreme years with that of random draws from all “normal” years in the chronology and tests significant deviations (21). We estimated lag effects during the first and second postdrought years. RWI anomalies provide robust, site-level estimates of drought impacts and can be interpreted as proportional reductions in stem diameter growth (fig. S2); therefore, we

¹Forest Ecology & Forest Management Group, Wageningen University, Wageningen, Netherlands. ²Department of Plant Biology, Institute of Biology, University of Campinas (UNICAMP), Campinas, SP, Brazil. ³Department of Forestry and Environmental Science, Shahjalal University of Science and Technology, Sylhet, Bangladesh. ⁴Laboratory of Tree-Ring Research, University of Arizona, Tucson, AZ, USA. ⁵Belgian Climate Centre, Uccle, Belgium. ⁶World Agroforestry Centre (ICRAF), Addis Ababa, Ethiopia. ⁷Department of Microbiology and Parasitology, Universidad Nacional Autónoma de México, Mexico City, Mexico. ⁸Colegio de Geografía, Facultad de Filosofía y Letras, Universidad Nacional Autónoma de México, Mexico City, Mexico. ⁹Department of Forest Engineering, Universidade Regional de Blumenau, Santa Catarina, Brazil. ¹⁰Northern Forestry Centre, Canadian Forest Service, Natural Resources Canada, Edmonton, AB, Canada. ¹¹Higher Technical School of Agricultural and Forestry Engineering and Biotechnology, University of Castilla-La Mancha, Albacete, Spain. ¹²Forest Management Program, Mamirauá Institute for Sustainable Development, Tefé, Brazil. ¹³Department of Plant Biology, Federal University of Ceará (UFC), Fortaleza, CE, Brazil. ¹⁴Department of Forest Sciences, Luiz de Queiroz College of Agriculture, University of São Paulo, Piracicaba, Brazil. ¹⁵Instituto de Pesquisas Ambientais do Estado de São Paulo, Assis, São Paulo, Brazil. ¹⁶Facultad de Ingeniería y Ciencias, Universidad Autónoma de Tamaulipas, Ciudad Victoria, Tamaulipas, Mexico. ¹⁷Tree-Ring Laboratory, Forest Science Department, Federal University of Lavras, Lavras, MG, Brazil. ¹⁸University of Campania L. Vanvitelli, Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, Caserta, Italy. ¹⁹Wood Biology Service, Royal Museum for Central Africa, Tervuren, Belgium. ²⁰Brazilian Agricultural Research Corporation (Embrapa), Embrapa Forestry, Colombo, PR, Brazil. ²¹Institute of Geography, Friedrich-Alexander-University Erlangen-Nuremberg, Erlangen, Germany. ²²School of Geography, University of Leeds, Leeds, UK. ²³Fenner School of Environment and Society, Australian National University, Canberra, Australian Capital Territory, Australia. ²⁴Faculty of Environment and Resource studies, Mahidol University, Salaya, Phutthamonthon, Nakhon Pathom, Thailand. ²⁵Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY, USA. ²⁶Instituto Pirenaico de Ecología (IPE-CSIC), Zaragoza, Spain. ²⁷Instituto de Silvicultura e Indústria da Madeira, Universidade Juárez del Estado de Durango, Durango, Mexico. ²⁸Department of Botany, Institute of Biosciences, University of São Paulo, São Paulo, Brazil. ²⁹Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias (INIFAP), Centro Nacional de Investigación Disciplinaria en Relación Agua-Suelo-Planta-Atmósfera (CENID-RASPA), Durango, Mexico. ³⁰Departamento Forestal, Universidad Autónoma Agraria Antonio Narro, Saltillo, Coahuila, Mexico. ³¹Department of Environmental Technology and Management, Faculty of Environment, Kasetsart University, Bangkok, Thailand. ³²CAS Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, Yunnan, China. ³³School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, UK. ³⁴Birmingham Institute of Forest Research, University of Birmingham, Birmingham, UK. ³⁵Laboratory of Plant Anatomy and Dendrochronology, Department of Biology, Universidade Federal de Sergipe - UFS, Aracaju, Brazil. ³⁶Laboratório de Dendrocronologia, Universidade Continental, Huancayo, Perú. ³⁷Department of Forest Sciences, Universidad Nacional de Colombia - Sede Medellín, Colombia. ³⁸Departamento de Agricultura, Biodiversidade e Florestas, Universidade Federal de Santa Catarina - UFSC, Curitiba, Santa Catarina, Brazil. ³⁹TERRA Teaching and Research Centre, Gembloux Agro-Bio Tech, University of Liège, Gembloux, Belgium. ⁴⁰Master School for Carpentry and Cabinetmaking, Ebern, Germany. ⁴¹Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, USA. ⁴²Santa Fe Institute, Santa Fe, NM, USA. ⁴³Department of Biological Sciences, University of Joazeiro Region - UNIVILLE, Joinville, Brazil. ⁴⁴Postgraduate Program in Forestry, Regional University of Blumenau - FURB, Blumenau, SC, Brazil. ⁴⁵UR Forêts et Sociétés du CIRAD, Montpellier cedex 5, France. ⁴⁶Gestion des Ressources Forestières de Gembloux Agro-Bio Tech, ULiège, Gembloux, Belgium. ⁴⁷Laboratório de Dendrocronologia e História Ambiental, INIA/CLA, CCT-CONICET-Mendoza, Mendoza, Argentina. ⁴⁸Department of Crop Sciences, Tropical Plant Production and Agricultural Systems Modelling, Göttingen University, Göttingen, Germany. ⁴⁹School of Biological Sciences, University of Western Australia, Perth, Australia. ⁵⁰USDA Forest Service, Pacific Southwest Research Station, Institute of Pacific Islands Forestry, Hilo, HI, USA. ⁵¹Department of Forest Sciences, Federal University of Paraná, Curitiba, PR, Brazil. ⁵²World Agroforestry Centre (ICRAF), Nairobi, Kenya. ⁵³Facultad de Ingeniería, Tecnológico de Antioquia, Medellín, Colombia. ⁵⁴Natural Resources and Environmental Sciences Department, Forestry, Ecology and Wildlife Program, Alabama A&M University, Huntsville, AL, USA. ⁵⁵Programa de Pós-Graduação em Agronomia/Fisiologia Vegetal, Departamento de Biologia, Instituto de Ciências Naturais, Universidade Federal de Lavras, Lavras, MG, Brazil. ⁵⁶Flanders Heritage Agency, Brussels, Belgium. ⁵⁷Department of Earth and Spatial Sciences, University of Idaho, Moscow, ID, USA. ⁵⁸German Archaeological Institute DAI, Department of Natural Sciences, Berlin, Germany. ⁵⁹Geography Department, Humboldt-University Berlin, Berlin, Germany. ⁶⁰Section Geomorphology, GFZ Helmholtz Centre for Geosciences, Telegrafenberg, Potsdam, Germany. ⁶¹Faculty of Bioscience Engineering, Department of Environment, Laboratory of Wood Technology (UGent-Woodlab), Ghent University, Ghent, Belgium. ⁶²School of Biological Sciences, Washington State University, Pullman, WA, USA. ⁶³Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Prague, Czech Republic. ⁶⁴US Fish and Wildlife Service, Albuquerque, NM, USA. ⁶⁵Department of Ecology and Biogeography, Faculty of Biological and Veterinary Sciences, Nicolaus Copernicus University, Toruń, Poland. ⁶⁶Centre for Climate Change Research, Nicolaus Copernicus University, Toruń, Poland. ⁶⁷Instituto Nacional de Innovación Agraria, Programa Nacional de Investigación Forestal, Huancayo, Perú. ⁶⁸CITEforestal Pucallpa, Pucallpa, Perú. ⁶⁹Department of Natural Resource Management, South Dakota State University, Brookings, SD, USA. ⁷⁰Department of Geography, Swansea University, Singleton Park, Swansea, UK. ⁷¹Department of Soils and Rural Engineering, Faculty of Agronomy and Animal Science, Federal University of Mato Grosso - UFMT, Cuiabá, Mato Grosso, Brazil. ⁷²Center of Nuclear Energy in Agriculture, University of São Paulo, São Paulo, Brazil. ⁷³Department of Animal Science, Federal University of Paraná, Curitiba, PR, Brazil. ⁷⁴CITAB - Department of Forestry Sciences and Landscape (CIFAP), University of Trás-os-Montes and Alto Douro, Vila Real, Portugal. ⁷⁵Plant Ecology, Institute of Integrative Biology, D-USYS, ETH-Zürich, Zürich, Switzerland. ⁷⁶Institute for Plant Sciences (IPS), University of Bern, Bern, Switzerland. ⁷⁷Department of Geography, Indiana University, Bloomington, IN, USA. ⁷⁸Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro, Querétaro, Mexico. ⁷⁹Graduate Program in Ecology and Conservation, Federal University of Sergipe - UFS, São Cristóvão, Sergipe, Brazil. ⁸⁰Brazilian Agricultural Research Corporation (Embrapa), Embrapa Amazônia Ocidental, Manaus, AM, Brazil. ⁸¹IHE Delft, Delft, Netherlands. ⁸²VU University Amsterdam, Amsterdam, Netherlands. ⁸³Posgrado en Bosques y Conservación Ambiental, Universidad Nacional de Colombia, Medellín, Colombia. ⁸⁴Centro Universitario de Ciencias Biológicas y Agropecuarias, Universidad de Guadalajara, Zapopan, Jalisco, Mexico. ⁸⁵Centre for Functional Ecology, Department of Life Sciences, Faculty of Sciences and Technology, University of Coimbra, Coimbra, Portugal. ⁸⁶National Biobank of Thailand, National Science and Technology Development Agency, Pathum Thani, Thailand. ⁸⁷Department of Biomaterials Science and Technology, School of Natural Resources, The Copperbelt University, Kitwe, Zambia. ⁸⁸Laboratory of Ecology and Dendrology, Federal Institute of Sergipe IFS, São Cristóvão, Sergipe, Brazil. ⁸⁹Laboratory of Plant Ecology, Universidade do Vale do Rio dos Sinos (UNISINOS), São Leopoldo, RS, Brazil. ⁹⁰Instituto Biodiversa, Porto Alegre, RS, Brazil. ⁹¹Nuclear Energy in Agriculture, University of São Paulo, Piracicaba, Brazil. ⁹²Department of Botany and Ecology, Institute of Biosciences, Federal University of Mato Grosso (UFMT), Cuiabá, Mato Grosso, Brazil. ⁹³Tree-Ring Laboratory, Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY, USA. ⁹⁴Carrera de Ingeniería Forestal, Universidad Autónoma Gabriel René Moreno, Facultad de Ciencias Agrícolas, El Vallecito, Santa Cruz, Bolivia. ⁹⁵BIOAPLIC, Departamento de Botánica, Universidade de Santiago de Compostela, EPSE, Campus Terra, Lugo, Spain. ⁹⁶El Colegio de la Frontera Sur, Departamento de Sociedad y Cultura, San Cristóbal de Las Casas, Chiapas, Mexico. ⁹⁷Escuela Profesional de Ingeniería Forestal de la Universidad Nacional Toribio Rodríguez de Mendoza de Amazonas, Chachapoyas, Perú. ⁹⁸Laboratorio de Dendrocronología, Carrera de Ingeniería Forestal, Universidad Nacional de Loja, Loja, Ecuador. ⁹⁹Department of Environmental Studies, University of California Santa Cruz, Santa Cruz, CA, USA. ¹⁰⁰Facultad de Ciencias Agrarias, Universidad del Cauca, Popayán, Colombia. ¹⁰¹Hémera Centro de Observación de la Tierra, Escuela de Ingeniería Forestal, Facultad de Ciencias, Universidad Mayor, Santiago, Chile. ¹⁰²Centro de Apoio Técnico à Execução Ambiental, Ministério Público do Estado de Mato Grosso, Mato Grosso, Brazil. ¹⁰³Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias (INIFAP), Centro de Investigación Regional Pacífico Centro - Campo Experimental, Centro Altos de Jalisco, Jalisco, Mexico. ¹⁰⁴DendrOlive, Departamento de Sistemas Físicos, Químicos y Naturales, Universidad Pablo de Olavide, Sevilla, Spain. ¹⁰⁵Forest and Nature Management, Van Hall Larenstein University of Applied Sciences, Velp, Netherlands. ¹⁰⁶National Institute for Amazon Research, Manaus, Amazonas, Brazil. ¹⁰⁷Environmental Studies, Department of Biology, Institute of Ecology and Evolution, University of Oregon, Eugene, OR, USA. ¹⁰⁸Department of Earth Sciences, Freie Universität Berlin, Berlin, Germany. ¹⁰⁹Instituto de Ecología Aplicada, Universidad Autónoma de Tamaulipas, Ciudad Victoria, Tamaulipas, Mexico. ¹¹⁰Institute of Biodiversity and Forests, Federal University of Western Pará - UFOPA, Santarém, Brazil. ¹¹¹Department of Earth and Environmental Systems, Indiana State University, Terre Haute, IN, USA. ¹¹²Department of Geography, University of Alabama, Tuscaloosa, AL, USA. ¹¹³Programa de Investigación de Ecología y Biodiversidad, Asociación ANDINUS, Sicaya, Huancayo, Junín, Perú. ¹¹⁴Department of Geography, Johannes Gutenberg University, Mainz, Germany. ¹¹⁵Department of Environmental Science, Faculty of Science, Chulalongkorn University, Bangkok, Thailand. ¹¹⁶Water Science and Technology for Sustainable Environment Research Unit, Chulalongkorn University, Bangkok, Thailand. ¹¹⁷Instituto de Geografía, Universidad Nacional Autónoma de México, Ciudad Universitaria, Mexico City, Mexico. ¹¹⁸Facultad de Ciencias Forestales, Colegio de Postgraduados, Montecillos, Texcoco, Mexico. ¹¹⁹Instituto de Ciencias Agroalimentarias, Animales y Ambientales (ICA3), Universidad de O'Higgins, San Fernando, Chile. ¹²⁰Biological Sciences Department, California State University, San Marcos, CA, USA. ¹²¹Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, Netherlands. ¹²²School of Teacher Training for Secondary Education Tilburg, Fontys University of Applied Sciences, Tilburg, Netherlands. ¹²³Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, Santa Cruz, CA, USA. ¹²⁴School of Earth Sciences, Addis Ababa University, Addis Ababa, Ethiopia. ¹²⁵School of Biology and Ecology, University of Maine, Orono, ME, USA. ¹²⁶School of Natural Resources and the Environment, University of Arizona, Tucson, AZ, USA. *Corresponding author. Email: pieter.zuidema@wur.nl




Fig. 1. Pantropical growth anomalies during drought years obtained from a tree-ring network. (A) Geographical distribution of 483 tropical tree-ring width chronologies from gymnosperms ($n = 273$, triangles) and angiosperms ($n = 210$, circles). The background color is tree cover percentage ($>10\%$). (B) Pantropical median anomalies of annual tree growth (whiskers indicate bootstrapped 95% CIs) during the 10% years with the lowest P, highest VPD, or highest CWD. Results are shown for droughts occurring during the dry season (filled symbols, bold text) and wet season (transparent symbols, normal text) and for angiosperms and gymnosperms combined ("all") or separately. Results of weighted Mann-Whitney U tests between seasons are shown (nonsignificant, NS: $P > 0.05$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$). No significant differences in growth anomalies were found between drought types for any combination of species groups and season (Kruskal-Wallis tests, $P > 0.05$). Percentage values represent the proportion of significant negative anomalies in SEA analyses ($P < 0.05$). Sample size per symbol is provided in table S1B.

refer to them as "growth anomalies" (or "growth reductions" when negative). To account for the higher abundance of tree-ring studies at high elevations and in arid climates (fig. S3), we present pantropical medians weighted by climatic representativeness.

The drought intensity of the selected 10% driest years (expressed as the number of standard deviations away from the long-term mean) was comparable across the three drought types (fig. S4). Overlap in event years across drought types was small, except for dry-season extremes in P and CWD (fig. S5), suggesting that CWD droughts are more driven by very low P than by very high VPD. Across drought types and seasons, 7% of the droughts occurred during 2 consecutive years, and 0.3% lasted for 3 years. Since 1930, the frequency of droughts has increased for all drought types (fig. S6A), whereas the intensity has increased for some (VPD and CWD) but decreased for others (P) (fig. S6B).

Variable but overall modest drought effects

Growth anomalies during the 10% driest years were predominantly negative (63% of incidences), with strong growth reductions ($>10\%$) occurring in 25% of the chronologies. However, growth anomalies varied across the network from strongly negative to moderately positive (fig. S7A), and growth anomaly variations were stronger for individual drought years (fig. S6C) than for averages across drought years.

This wide variety of growth anomalies during individual drought years fully captures previously reported responses of tropical trees to episodic droughts, ranging from stem growth reductions of 20 to 50% (25, 26) to undetectable change (13, 27) or growth increases [for small trees (28)]. The latter may result from reduced cloudiness, enhanced leaf production, and/or competitive release in humid regions (28, 29).

Despite the large variation in site-specific drought responses, bootstrapped 95% confidence intervals (CIs) of the weighted median responses were narrow (Fig. 1B). Overall, the magnitude of growth declines during drought years was modest (Fig. 1B). The pantropical growth reduction for all drought types and seasons was 2.5% (weighted median, 95% CI: 2.2 to 2.7%, with angiosperms and gymnosperms combined). Growth reduction was significantly stronger for dry-season droughts (3.1%; 95% CI: 2.8 to 3.5%) than wet-season droughts (1.8%; 95% CI: 1.6 to 2.1%) (Mann-Whitney U test, $P < 0.001$). Our assessments focus on the pantropical level because continental-level comparisons are hampered by differences in climatic representation (fig. S1D) and local-scale assessments by high local tree diversity (fig. S1E).

One fourth of the growth anomalies during dry-season droughts and 15% during wet-season droughts were significant ($P < 0.05$; Fig. 1B

and fig. S7A). SEAs of individual ring-width series revealed smaller proportions of significant growth declines: 10% during the dry season and 7% during the wet season (fig. S8). These lower percentages likely reflect a larger role of nonclimatic factors in inducing growth variation of individual trees (e.g., canopy disturbance and carryover effects) than variation at the population level.

Applying a more stringent selection of drought years (i.e., the 5% driest years) led to a somewhat stronger, but still modest, mean reduction of 3.2% (95% CI: 2.7 to 3.5%). This modest drought response is comparable in magnitude to that from multisite assessments in tropical forests, which reported 6 to 9% reductions in diameter growth or woody biomass productivity (11, 30, 31). The stronger growth reductions in those studies may be explained by the selection of more extreme droughts.

Drought responses differed markedly between angiosperms and gymnosperms (Fig. 1B). Dry-season droughts induced significantly stronger growth reductions for gymnosperms (4.4%; 95% CI: 4.0 to 5.1%) than for angiosperms (1.1%; 95% CI: 0.5 to 1.9%) (table S1). By contrast, wet-season droughts resulted in stronger growth declines for angiosperms (2.7%; 95% CI: 2.2 to 3.1%) than gymnosperms (1.3%; 95% CI: 0.8 to 1.6%) (table S1). Stronger drought responses of gymnosperms are broadly consistent with the results of extratropical tree-ring studies (19, 21, 22, 32). These taxonomic differences cannot be explained

by the drought intensity, which was similar for angiosperms and gymnosperms (fig. S4). Possible explanations of stronger drought effects for gymnosperms include a dominant evergreen leaf phenology (89% of gymnosperm chronologies), an inherently stronger growth variability (21), a stronger vulnerability to cavitation for *Pinaceae* (80% of our chronologies) (33), and lower concentrations of nonstructural carbohydrate (NSC) reserves (34). The marked differences in drought responses and hydraulic architecture between angiosperms and gymnosperms call for separate drought analyses of these major clades.

In contrast to our hypothesis about question (i), growth reductions were not larger during CWD droughts compared with P or VPD droughts (Fig. 1B and table S1). This may partially be explained by the high coincidence of CWD and P droughts (fig. S5) resulting in similar effects on tree growth.

Rapid postdrought recovery

Contrary to our hypothesis about question (ii), drought analyses revealed no evidence for strong or long-lasting lag effects for any drought type or season (Fig. 2). Compared with drought years, postdrought years exhibited significantly smaller negative growth anomalies, and anomalies often shifted to positive. Such significant changes were found in 80% of the 36 possible comparisons (of clades, seasons, and drought types) between drought and postdrought years. During postdrought years, the 95% CIs of growth anomalies included 0 or were fully >0 for 70% of the 36 combinations of drought types, seasons, and clades. Postdrought recovery was similar for angiosperms and gymnosperms (Fig. 2) and cannot be explained by postdrought climatic conditions, which were close to normal or only slightly wetter (fig. S4, B and C).

Robustness tests revealed that these drought responses do not shift when applying more rigid detrending methods (fig. S9), alternative climate products (figs. S10 and S11), or more stringent drought selection criteria (fig. S12, A and B) or when selecting 2-year droughts (fig. S12C).

Combined, our results point to a rapid recovery of tree growth to predrought levels and provide no evidence for the existence of strong or long lag effects at the pantropical scale. These results are consistent with the short (<18 months) lag effects in tree growth after droughts observed in tropical dendrometer studies (11, 35) and extratropical tree-ring studies (22). However, our results contrast with the strong lagged drought responses for gymnosperms observed in a tree-ring study across extratropical sites (19), likely due to differences in site aridity, drought definitions, detrending methods, and species selection (22).

Drought effects are partially mitigated by wet extremes

To contextualize drought-induced growth anomalies, we compared the overall effect of the 10% driest years on tree growth with growth changes during the 10% wettest years (SEA analysis). As hypothesized for question (iii), these wettest years caused pantropical tree growth to increase (median 1.8%; 95% CI: 1.5 to 2.0%) (fig. S7B), a magnitude that is comparable to the 2.5% drought-induced growth reduction (Fig. 1B). Similar to drought responses, growth anomalies after wet extreme years were short-lived (fig. S13).

As a result of the growth increases during extreme wet years, net growth anomalies of the 10% driest and wettest years combined were overall very small (-0.4% ; 95% CI: -0.2 to -0.5%) and not significantly different from 0 for most drought types (Fig. 3). However, wet extremes mitigated only half (47%; 95% CI: 43 to 52%) of the growth reduction during drought years (across seasons and drought types and for angiosperms and gymnosperms combined; Fig. 3). With increasing frequency or severity of droughts (fig S6, A and B) and in more arid conditions (23), this wet-year compensation may be reduced.

Drought impacts increase with aridity

We evaluated the climatic drivers of observed drought effects by interpolating growth anomalies across climate space (mean annual precipitation and temperature). We found that wet-season droughts reduced growth in angiosperms more strongly in hotter and more arid climates (Fig. 4A and figs. S14A and S15A), whereas dry-season droughts had stronger impacts on gymnosperms at more arid sites (Fig. 4A and figs. S14B and S15B). Path analyses yielded consistent results (fig. S16) and also revealed that the expected role of first-order growth autocorrelation in shaping drought effects (22) was very small. Our finding of aggravating drought effects with increasing aridity is consistent with those from earlier studies on tropical (36) and extratropical (19, 20, 23, 37) tree species. Strong local variability in drought responses is common (35), likely due to interspecific differences in drought resistance (38), spatial variation in soil parameters, rooting depth, access to water (39), and variations in the timing, intensity, and duration of droughts (37).

We projected the climate-space patterns for growth anomalies during low-precipitation years (i.e., models with highest R^2) into geographic space (Fig. 4B). For angiosperms, this model suggests that the strongest drought-induced growth reductions occur in dry-forest biomes




Fig. 2. Drought effects on pantropical tree growth are short-lived. Growth anomalies during and after years with low P, high VPD, or high CWD. Responses are shown for droughts occurring during the dry season and wet season and for angiosperms and gymnosperms combined ("all") or separately. Bands indicate bootstrapped 95% CIs of medians. Different letters indicate a significant difference between years (Mann-Whitney U tests; $P < 0.05$) per season (dry | wet). Sample size per line is provided in table S1B.




Fig. 3. Wet extremes partially mitigate drought effects on tropical tree growth. Pantropical medians (and bootstrapped 95% CIs) of the net growth anomaly of dry and wet extreme years. Results are shown for extremes occurring in the dry or wet seasons and for angiosperms and gymnosperms combined (“all”) or separately. Droughts (and wet extremes) were identified as the 10% years with the lowest (highest) P, highest (lowest) VPD, or highest (lowest) CWD. The net anomaly was calculated as the average of the anomalies of the dry and wet extreme years. Percentages denote the growth loss during drought years that is mitigated during wet extremes per season (**dry** | **wet**). Sample size per symbol is provided in table S1B.




Fig. 4. Climatic and spatial distribution of drought-induced anomalies of tropical tree growth. (A) Growth anomalies during years with low precipitation during the wet season (angiosperms, $n = 210$) and dry season (gymnosperms, $n = 273$) interpolated across climate space. The interpolated climate space was restricted to 95% of the climatic ranges of precipitation and temperature. (B) Projections of the interpolated values from (A) into geographic space. The geographic distribution of gymnosperms was restricted to elevations >700 m above sea level (i.e., the 10th percentile of their elevational distribution). For Africa, the gymnosperm distribution was further restricted by a species distribution model of the only African gymnosperm in our network (*Juniperus procera*).

in the Americas, Africa, South Asia, and eastern Australia. Modest or positive drought effects on stem growth are estimated in warm, humid zones such as the Amazon region. For gymnosperms, negative growth anomalies due to dry-season droughts are strong throughout high-elevation regions in the Americas, Asia, and eastern Africa. Despite large local variability in drought responses, the bootstrapped uncertainty of these maps is low (fig. S17). Thus, across major climatic gradients at the subcontinental scale, predicted distributions of drought responses likely hold. These results offer opportunities to benchmark and constrain simulated drought responses of woody biomass production in tropical forests using terrestrial biosphere models (16, 40).

Drought responses across components of productivity
The drought effects on stem growth are smaller than those on leaf-level photosynthesis (31) and forest-level gross primary productivity (GPP) (13) but of comparable magnitude to effects on leaf fall, flushing, and mature leaf area (35). In terms of duration, drought impacts on stem growth tend to last somewhat longer (4 to 18 months) (27, 35) than effects on GPP (typically 4 to 6 months) (41, 42). These comparisons suggest more direct and stronger drought effects on carbon uptake through photosynthesis than on

its sequestration in woody biomass, consistent with the decoupling of these processes observed in extratropical forests (43).

One of the plausible explanations of the observed drought resilience of tropical tree growth is the mobilization of NSCs. Tropical tree species store large amounts of starch and soluble sugars in stems, branches, roots, and leaves (44) that are available during dry episodes for osmoregulation, leaf flushing, and stem growth (13, 34, 45, 46). NSC mobilization may buffer reductions in stem growth and explain the decoupling of canopy and stem responses to droughts, but this remains poorly quantified. Other mechanisms, including shifts in leaf phenology and stem hydraulics (47), may also contribute to drought resilience. Long-term, well-replicated field studies measuring functional and productivity changes are needed.

Limitations, mortality risks, and climate change

We acknowledge several limitations of our study that should be addressed in future studies. First, major sampling gaps exist in Africa and in extreme arid and humid climates (figs. S1 and S3) (18). Although we statistically account for biases, data scarcity can only be solved through new studies. Second, our chronologies only include cross-dated RWI series, and our study thus may have excluded individuals that did not cross-date due to extraordinarily strong (e.g., missing rings) or weak drought responses. The latter group of individuals may produce progeny for drought-resilient next generations if weak responses have a genetic basis. Third, our study species represent a small fraction of tropical tree richness, and their drought responses may somewhat deviate from that of the “average” tropical species. Finally, we do not quantify drought responses at the stand level. Upscaling from the tree level to the forest level would require comparing tree-ring-based responses with those from community-wide dendrometer or plot data.

Episodic droughts increase the mortality of tropical trees (6, 48), and drought-induced stem growth reductions may be associated with elevated mortality. To provide a first estimate of the mortality risks associated with the growth reductions reported here for angiosperms, we used significant growth-mortality associations from 10 tropical forest field studies (49) (fig. S18, A and B). The resulting estimate is a first indication of the order of magnitude of additional tree mortality. We estimate this to be 0.1% per drought year (95% CI: 0.08 to 0.15%) on top of a 1% year⁻¹ baseline (fig. S18C). The resulting carbon loss of this additional mortality may be substantial at the pantropical scale and likely recovers slowly (50).

Anthropogenic climate change has intensified drought stress (fig. S6, A and B), resulting in stronger drought-induced growth reductions for all drought types (fig. S6C). Future climate change will further increase the frequency of droughts in tropical forests (51, 52). Our climate space interpolation and path analyses suggest that future warming and amplified climatic variability will cause stronger stem-growth declines during wet-season droughts in angiosperm-dominated tropical lowland forests and woodlands. For gymnosperm-dominated high-elevation forests, the stronger impacts of dry-season droughts are expected in regions where climate change would reduce precipitation. These shifts are unlikely to be mitigated by concurrent CO₂ rise (9) because no consistent CO₂-induced growth stimulation has been observed in tree-ring studies (53).

Stronger and more extensive droughts occurring under future climate change may shift the modest drought responses observed here for the past decades toward considerably larger and more widespread declines in tropical wood productivity. Such shifts may have implications for the dynamics and residence time of carbon in tropical forests, especially when stronger growth reductions are accompanied by increasing tree mortality.

REFERENCES AND NOTES

1. Y. Pan et al., *Science* **333**, 988–993 (2011).
2. P. Friedlingstein et al., *Earth Syst. Sci. Data* **11**, 1783–1838 (2019).

3. A. Koch, J. O. Kaplan, *Nat. Clim. Chang.* **12**, 279–283 (2022).
4. B. Buma et al., *Nat. Clim. Chang.* **14**, 402–406 (2024).
5. N. G. McDowell et al., *Science* **368**, eaaz9463 (2020).
6. P. M. Brando et al., *Annu. Rev. Earth Planet. Sci.* **47**, 555–581 (2019).
7. D. M. Lapola et al., *Science* **379**, eabp8622 (2023).
8. W. Yuan et al., *Sci. Adv.* **5**, eaax1396 (2019).
9. J. Peñuelas et al., *Nat. Ecol. Evol.* **1**, 1438–1445 (2017).
10. W. R. L. Anderegg et al., *Science* **368**, eaaz7005 (2020).
11. S. W. Rifai et al., *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **373**, 20170410 (2018).
12. P. M. Brando et al., *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **363**, 1839–1848 (2008).
13. C. E. Doughty et al., *Nature* **519**, 78–82 (2015).
14. P. A. Zuidema, B. Poulter, D. C. Frank, *Trends Plant Sci.* **23**, 1006–1015 (2018).
15. F. Babst et al., *Trends Plant Sci.* **26**, 210–219 (2021).
16. X. Xu et al., *Glob. Chang. Biol.* **30**, e17075 (2024).
17. P. A. Zuidema et al., *Nat. Geosci.* **15**, 269–276 (2022).
18. P. Groenendijk et al., *Quat. Sci. Rev.* **355**, 109233 (2025).
19. W. R. L. Anderegg et al., *Science* **349**, 528–532 (2015).
20. A. Gazol, J. J. Camarero, W. R. L. Anderegg, S. M. Vicente-Serrano, *Glob. Ecol. Biogeogr.* **26**, 166–176 (2017).
21. A. Gazol et al., *J. Ecol.* **108**, 2473–2484 (2020).
22. S. Klesse et al., *J. Ecol.* **111**, 1188–1202 (2022).
23. M. P. Dannenberg, E. K. Wise, W. K. Smith, *Sci. Adv.* **5**, eaaw0667 (2019).
24. J. M. Lough, H. C. Fritts, *Clim. Change* **10**, 219–239 (1987).
25. M. Rahman, M. Islam, A. Bräuning, *For. Ecol. Manage.* **433**, 487–496 (2019).
26. D. B. Clark, D. A. Clark, S. F. Oberbauer, *Glob. Chang. Biol.* **16**, 747–759 (2010).
27. J. A. Hogan et al., *Biotropica* **51**, 139–154 (2019).
28. A. C. Bennett, N. G. McDowell, C. D. Allen, K. J. Anderson-Teixeira, *Nat. Plants* **1**, 15139 (2015).
29. L. Rowland et al., *Nature* **528**, 119–122 (2015).
30. T. R. Feldpausch et al., *Global Biogeochem. Cycles* **30**, 964–982 (2016).
31. T. Janssen, K. Fleischer, S. Luyssaert, K. Naudts, H. Dolman, *Biogeosciences* **17**, 2621–2645 (2020).
32. X. Li et al., *Nat. Ecol. Evol.* **4**, 1075–1083 (2020).
33. B. Choat et al., *Nature* **491**, 752–755 (2012).
34. J. Martínez-Vilalta et al., *Ecol. Monogr.* **86**, 495–516 (2016).
35. T. Janssen et al., *Biogeosciences* **18**, 4445–4472 (2021).
36. D. Bauman et al., *Glob. Chang. Biol.* **28**, 1414–1432 (2022).
37. X. C. Wu et al., *Earths Futur.* **10**, e2021EF002530 (2022).
38. R. S. Oliveira et al., *New Phytol.* **230**, 904–923 (2021).
39. N. B. Schwartz et al., *Front. For. Glob. Change* **3**, 596256 (2020).
40. S. Klesse et al., *Global Biogeochem. Cycles* **32**, 1226–1240 (2018).
41. Z. Zhang, W. Ju, Y. Zhou, X. Li, *Glob. Chang. Biol.* **28**, 3620–3635 (2022).
42. J. Yang et al., *Glob. Chang. Biol.* **24**, 1919–1934 (2018).
43. A. Cabon et al., *Science* **376**, 758–761 (2022).
44. M. K. Würth, S. Peláez-Riedl, S. J. Wright, C. Körner, *Oecologia* **143**, 11–24 (2005).
45. L. T. Dickman et al., *Plant Cell Environ.* **42**, 1705–1714 (2019).
46. C. Signori-Müller et al., *Nat. Commun.* **12**, 2310 (2021).
47. J. V. Tavares et al., *Nature* **617**, 111–117 (2023).
48. W. M. Hammond et al., *Nat. Commun.* **13**, 1761 (2022).
49. S. E. Russo et al., *Nat. Ecol. Evol.* **5**, 174–183 (2021).
50. J. P. Wigneron et al., *Sci. Adv.* **6**, eaay4603 (2020).
51. C. G. Xu et al., *Nat. Clim. Chang.* **9**, 948–953 (2019).
52. S. Saatchi et al., *One Earth* **4**, 988–1003 (2021).
53. A. P. Walker et al., *New Phytol.* **229**, 2413–2445 (2021).
54. P. A. Zuidema et al., Data and code for: Pantropical tree rings show small effects of drought on stem growth. Dryad (2025); <https://doi.org/10.5061/dryad.hx3ffbgq4>.
55. National Centers for Environmental Information, National Oceanic and Atmospheric Administration, “International tree-ring data bank (ITRDB)” (NOAA, 2025); <https://www.ncei.noaa.gov/products/paleoclimatology/tree-ring>.

ACKNOWLEDGMENTS

We thank the Smithsonian Tropical Research Institute-Panama and S. Bernal (RaAS) for support and assistance. H. Muller-Landau and S. Joseph Wright (RaAS) supervised the project. COOMFLONA - FLONA TAPAJÓS, Universidade do Oeste do Pará (BH), Sutó Company, and A. Chavez at Consultora Forestal Bosques e Industria (KPV) and the logging company AMATA (DROR) provided fieldwork support. R. Franklin, G. Guada, Q. Hakkaert, A. Nijmeijer, and P. van der Sleen (KPV) provided laboratory support. **Funding:** This work was supported by AECID grant 11-CAP2-1730 (J.J.C.); Agencia Nacional de Promoción Científica y Tecnológica, Argentina grant PICT 2014-2797 (M.E.F.); Agencia Nacional de Promoción Científica y Tecnológica, Argentina grant PICT 2019-01336 (M.E.F.); BBVA Foundation (J.J.C.); Belspo BRAIN grant BR/143/A3/HERBAXYLAREDD (H.B.); CAPES - Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (T.L.S.); CAPES grant 88887199858/2018-00 (G.A.P.); CAPES grant 88887495294/2020-00 (B.H.); CAPES (P.G.); CAPES/PDSE grant 15011/13-5 (M.A.P.); CGIAR (M.M.); CNPq grant 140849/2015-7 (A.K.M.); CNPq ENV grant FRG 0339638 (O.D.); CNPq grant 1009/4785031-2 (G.C.); CNPq grant 140277/2024-2 (D.B.C.C.); CNPq grant 311247/2021-0 (J.S.); CNPq grant 405923/2021-0 (M.C.S.); CNPq grant 406062/2023-4 (M.C.S.); CNPq grant 44181/2020-5 (J.S.); CNPq grant PQ 313129/2022-3 (A.C.B.); CONACYT Consejo Nacional de Ciencia y Tecnología México master

fellowship (M.I.L.H.); CONACYT grant CB2016-283134 (J.V.D.); CONACYT grant CB-2016-283134 (L.R.C.E.); CONAFOR-CONACYT grant C01-234547 (J.C.P.); CONAFOR-CONACYT grant CONAFOR-2014 (L.R.C.E.); CONCYTEC Peru and World Bank grant 043-2019-FONDECYT-BM-INC.INV (J.G.I.); CONICET grant PIP-11220200102929CO (M.E.F.); Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (E.A.F.); Copel Geração e Transmissão S.A. grant PD-06491-0405-2015 (A.K.M.); CUOMO Foundation (M.M.); DAAD (M.I.); DFG grant BR 1895/15-1 (A.B.); DFG grant BR 1895/23-1 (A.B.); DFG grant BR 1895/29-1 (A.B.); Dirección de Investigación de la Universidad Nacional de Loja (D.P.C.); Dirección General de Asuntos del Personal Académico of the UNAM (Mexico) (R.B.); Estudios dendrocronológicos en las Sierras Madre Occidental, Oriental y del Sur de México, grant 38111-4251030012346 (E.C.O.); FACEPE grant IBPG-1418-5.00/21 (D.B.C.C.); FAPEAM grant 01.02.016301.02630/2022-76 (J.S.); FAPEMAT (F.A.L.); FAPEMIG grant APQ-01544-22 (A.C.B.); FAPEMIG grant APQ-02541-14 (G.A.P.); FAPESC grant 2019TR65 (T.A.B.F.); FAPESP grant 12/50457-4 (G.C.); FAPESP grant 2009/53951-7 (M.T.F.); FAPESP grant 2017/50085-3 (G.C., G.M.L., and D.R.O.R.); FAPESP grant 2018/01847-0 (P.G.); FAPESP grant 2018/07632-6 (M.G.V.); FAPESP grant 2018/22914-8 (D.R.O.R.); FAPESP grant 2018/24514-7 (J.R.V.A.) FAPESP grant 2019/08783-0 (G.M.L.); FAPESP grant 2019/09813-0 (M.G.V.); FAPESP grant 2020/04608-7 (D.R.O.R.); FAPESP-NERC grant 18/50080-4 (G.C.); FAPITEC (C.S.L. and A.S.R.); FAPITEC/SE/FUNTEC grant 01/2011 (M.A.P.); FCT - Portuguese Foundation for Science and Technology grant UIDB/04033/2020 (J.L.P.C.L.); FONDECYT grant BM-INC.INV 039-2019 (M.E.F.); Fulbright Fellowship (B.J.E.); German Academic Exchange Service (DAAD) (M.R.); German Research Council (M.M.); HELVETAS Swiss Intercooperation (M.E.F.); IAI-SGP-CRA grant 2047 (J.V.D.); IFS grant D/5466-1 (J.H.S.); IFS grant D/5466-1 (J.N.); Inter-American Institute for Global Change Research IAI (F.A.R.); ITTO Fellowship Award 046/12S (E.J.R.R.); Lamont Climate Center (B.M.B.); Mahidol University grant FRB660042/0185 (N.P.); María Zambrano postdoctoral research program MZ2021 (RaAS); Minciencias grant 1118-714-51372 (I.R.N.M.); Ministerio Ciencia grant TED2021-129770B-C22 (R.S.S.); National Geographic Global Exploration Fund grant GEFNE80-13 (I.R.); National Natural Science Foundation of China grant 31870591 (P.F.); National Research Council of Thailand (NRCT) grant N42A660392 (P.T.); NSF grant AGS-1501321 (D.G.S.); NSF (B.M.B.); NSF CREST grant 0833211 (K.S.F.); NSF grant IBN-9801287 (A.J.L.); NSF grant AGS-1501321 (G.A.P.); NSF grant AGS-2102888 (J.M.); NSF grant AGS-2102938 (G.L.H.); NSF grant GER 9553623 (B.J.E.); NSF postdoctoral fellowship (B.J.E.); NSF-FAPESP PIRE grant 2017/50085-3 (C.F. and M.T.F.); NSF-FAPESP PIRE grant 2019/27110-7 (C.F.); NUFFIC (J.H.S. and J.N.); NUFFIC-NICHE (E.M.);

PIRE-CREATE grant 2017/50085-3 (M.G.V.); PROCAD-AM grant 88887.625854/2021-00 (B.H.); Proyectos de generacion de conocimiento grant PID2021-1236750B-C44 (R.S.S.); Schlumberger Foundation (J.H.S. and J.N.); short-term fellowship from the Smithsonian Tropical Research Institute (RaAS); Sigma Xi (A.J.L.); Spanish Agency for International Development Cooperation (K.P.V.); Thailand Science Research and Innovation Fund Chulalongkorn University (P.T.); The Copperbelt University (J.H.S. and J.N.); The National Institute of Science and Technology in Plant Physiology under Stress Conditions grant 406455/2022-8 (F.A.L.); UK NERC grant NE/K01353X/1 (E.G.); UKRI grant EP/X025098/1 (N.J.L.); UKRI grant NE/B501504 (N.J.L.); UNAM-PAPIIT grant IN110223 (L.V.S.); Universidad Autónoma Agraria Antonio Narro (E.C.O.); World Wildlife Fund (WWF) (K.P.V.); Xunta de Galicia grant ED431C 2023/19 (G.P.L.); and Xunta de Galicia grant ED481D 2023/012 (G.P.L.).

Author contributions: Conceptualization: P.A.Z., F.B., P.G., M.R., VT; Data analysis: P.A.Z., F.B., M.R., P.G.; Data contribution: all authors except F.B.; Writing – original draft: P.A.Z. and F.B. with important contributions from V.T., M.R., P.G.; Writing – review & editing: all authors.

Competing interests: The authors declare no competing interests. **Data and materials availability:** All tree growth anomalies during and after drought years from the 483 tree-ring chronologies included in the study are provided as data S1. All 483 chronologies (from 1930 onward) are publicly available at Dryad (54). Raw tree-ring width measurements for all 483 chronologies are publicly available free of charge and for all research purposes on the International Tree-Ring Data Bank (ITRDB) (55). Links to the raw data of all 483 chronologies available on the ITRDB are included in data S1 and in the file metadata_FINAL.csv on Dryad (54). All code to build chronologies from raw tree-ring data, process climate data, conduct superposed epoch analyses, analyze statistics, and produce figures has been deposited on Dryad (54). **License information:** Copyright © 2025 the authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original US government works. <https://www.science.org/about/science-licenses-journal-article-reuse>

SUPPLEMENTARY MATERIALS

science.org/doi/10.1126/science.adq6607
Materials and Methods; Figs. S1 to S18; Table S1; References (56–75); MDAR Reproducibility Checklist; Data S1

Submitted 7 June 2024; accepted 16 April 2025

10.1126/science.adq6607