



# Forest restoration treatments increase native plant diversity but open the door to invasion in the Colorado Front Range<sup>☆</sup>

Adam L. Mahood<sup>a,\*</sup>, Camille S. Stevens-Rumann<sup>b,c</sup>, Allison E. Rhea<sup>c</sup>, Scott M. Ritter<sup>b</sup>, Kevin Barrett<sup>c</sup>, Paula J. Fornwalt<sup>d</sup>, David M. Barnard<sup>a</sup>

<sup>a</sup> Agricultural Research Service, United States Department of Agriculture, 2150 Centre Ave Building D, Fort Collins, CO 80526, USA

<sup>b</sup> Forest and Rangeland Stewardship, Colorado State University, 400 University Ave, Fort Collins, CO 80523, USA

<sup>c</sup> Colorado Forest Restoration Institute, Colorado State University, 1001 Amy Van Dyken Way, Fort Collins, CO 80521, USA

<sup>d</sup> Rocky Mountain Research Station, Forest Service, United States Department of Agriculture, 240 West Prospect Road, Fort Collins, CO 80526, USA

## ARTICLE INFO

### Keywords:

Forest restoration treatments  
Understory diversity  
Ponderosa pine  
Non-native plants

## ABSTRACT

Forest restoration treatments in dry conifer forests of the western United States are often done with objectives to move current forest structure toward historical conditions and, in turn, increase the system's resilience to future wildfires. But little is known about their effects on understory plant composition, particularly over the long-term. This is especially true in the Colorado Front Range (CFR). We used a before/after control/impact study design to assess the effects of mechanical forest restoration treatments on forest structure, surface fuels and understory plant composition 1, 5, and 10 years after treatment. Five and 10 years after treatment, treated areas had lower basal area and tree density, and higher quadratic mean diameter, consistent with treatment objectives. Treated areas also had higher native understory plant cover and richness, and higher graminoid cover. Species accumulation curves showed treatment-wide increases in native richness at 5 years post-treatment, and further increases at 10 years, with a net increase of 47 native species in treated areas compared to 11 in untreated controls. However, treatments also had higher relative numbers of non-native plant species, and increased probability of non-native plant invasion, though the overall level of invasion was low. Native richness, native cover and graminoid cover were more strongly associated with stand structure and topography, while non-native invasions were more strongly associated with moisture deficit. This suggests that climate played a greater role in enabling the introduction of species than changes in stand structure affected by treatment. In the CFR, forest restoration treatments can benefit the native understory, but also provide an opening for invasion in relatively uninvaded sites.

## 1. Introduction

Fire regimes are changing globally, with widespread documentation of increases in burned area (Abatzoglou and Williams, 2016), burn severity (Cansler and McKenzie, 2014; Parks, 2016), and damage to infrastructure (Mietkiewicz, 2020; Cook, 2021; Higuera, 2023). In dry conifer ecosystems in the western United States, the effects of fire exclusion (Kreider, 2024) and timber harvesting (Naficy, 2010) have exacerbated severe wildfire risk by altering the quantity and spatial arrangement of fuels. Land management agencies are implementing programs to mitigate this risk, in particular via forest restoration

treatments that move stands toward more open conditions similar to those that existed historically (Agee and Skinner, 2005; Stephens, 2020; Hurteau, 2024).

In western US dry conifer forest types, forest restoration treatments are commonly implemented via tree thinning (Hessburg, 2021). Despite the benefits of prescribed fire, implementation challenges continue to limit the scale of its application, leaving forest managers to largely rely on thinning-only treatments. In addition to benefits for wildfire hazard, thinning treatments can enhance tree growth, reduce drought vulnerability, and promote biodiversity (Knapp, 2021; McCauley, 2022; Zald, 2022; Rodman, 2024). Over the past decade managers have also

<sup>☆</sup> Open Research Statement: All data and code to fully reproduce the analysis will be made public at [www.github.com/admahood/thinned\\_comms](https://www.github.com/admahood/thinned_comms) and given a permanent digital object identifier upon acceptance.

\* Corresponding author.

E-mail address: [admahood@gmail.com](mailto:admahood@gmail.com) (A.L. Mahood).

<https://doi.org/10.1016/j.foreco.2025.122881>

Received 7 April 2025; Received in revised form 21 May 2025; Accepted 29 May 2025

Available online 7 June 2025

0378-1127/Published by Elsevier B.V.

increasingly embraced thinning approaches that aim to create highly heterogeneous canopy structures through the deliberate creation of isolated trees, discrete tree groups, and non-treed openings (Larson and Churchill, 2012; Addington, 2018), while reducing tree density, retaining large, fire-resistant trees, and increasing mean canopy base height. These approaches result in stands that better reflect wildfire-resilient historical forest structures (Jeronimo, 2019; Chamberlain, 2023). Enhancing the heterogeneity of the horizontal canopy structure also increases the diversity of the understory light environment which may have implications for tree regeneration and cover and biodiversity of understory plant communities.

In forested ecosystems, understory plant species make up most of the diversity and are crucial components of ecosystem functions such as decomposition, water balance and nutrient cycling (Balandier, 2022). After thinning-based forest restoration treatments, the understory plant community can be affected by physical disturbance during operations (Korb et al., 2007; Nikooy, 2020; Labelle, 2022), increased surface fuels (Wolk and Rocca, 2009), and increased canopy openness (McConnell and Smith, 1970). A more open canopy increases light availability (Riegel et al., 1992), belowground resources (Riegel et al., 1992; Neal, 2007), and soil water content (Zou, 2008). The increase in resource availability after a forest restoration treatment can ultimately lead to a bolstering of the understory community as newly available niche space is filled. Native species that are adapted to open conditions can be among the primary beneficiaries, particularly if tree canopy cover is reduced to 30–50 % or less (Abella and Springer, 2015) or if basal area is reduced to less than 20 m ha<sup>-1</sup> (Demarest, 2023). Increases in native plant abundance are commonly observed by the time treatments are ~3–5 years old (Metlen and Fiedler, 2006; Abella and Springer, 2015; Willms, 2017; Hood et al., 2024), as are increases in plant diversity (McConnell and Smith, 1970; Nelson et al., 2008; McGlone et al., 2009; Abella and Springer, 2015; Vernon, 2023; Springer, 2024).

Despite the observed benefits, there are concerns about forest restoration treatments having unintended consequences, including the invasion by non-native species and the initiation of human-grass-fire cycles (Kerns, 2020; Fusco, 2021). There are non-native species capable of contributing to ecosystem transformation present across much of the United States (Fusco, 2019), consistent observations of increases in non-native plant abundance after treatments (Nelson et al., 2008; Willms, 2017), and consistent observations of graminoids increasing with lower tree density (McConnell and Smith, 1970; Naumburg and DeWald, 1999; Fornwalt, 2017). The increased resource availability that benefits native plants in the understory also provides colonization opportunities for non-native plants (Davis et al., 2000). While non-native plants may remain at low levels of abundance following treatment (Hood et al., 2024; Springer, 2024), there are also cases in which non-native abundance was minimal immediately after treatment, but then increased dramatically (McGlone et al., 2009). Non-native plant invasions often proceed in stages, for example: stage 1) uninvaded; stage 2) present at low cover and having little ecological impact; stage 3) present and abundant enough to impact ecosystem function; stage 4) dominant and capable of causing ecosystem transformation (Brooks, 2004). Most analyses of thinning treatments on plant understories commonly report estimates of absolute cover or species richness of non-native plants to assess non-native invasion (Abella and Springer, 2015; Willms, 2017). While these metrics can characterize the impact of invasions, they are scale-dependent (Gotelli and Colwell, 2001; Thompson and Withers, 2003) and have limited comparability among habitats (Catford, 2012). Occurrence of any non-native species can be used to differentiate from stage 1 (uninvaded), but there is no information on relative impact. Differentiating among stages 2 through 4 can be assessed in a way that is comparable across regions and systems with invasion rate, defined as the percentage of non-native species, and invasion impact, defined as the relative cover of non-native species (Catford, 2012).

Regional differences in long-term effects of forest restoration

treatments on the understory (Schwilk, 2009) underscore the importance of understanding the effects of treatment on a regional level, rather than making broad generalizations across the entire western US.

Dry conifer ecosystems range from mesic to semi-arid to arid regions. Productivity is lower, masting events are less frequent (Wion, 2023), and seedling mortality is higher (Rother et al., 2015; Davis, 2019) in more arid locations. Understory plant communities can also remain more stable after treatment in the long term (> 10 years) (Laughlin, 2008; Springer, 2024) (Jang, 2021; Hood et al., 2024). The Colorado Front Range lies in the semi-arid part of this range, where there have only been a handful of studies on the effects of thinning on understory plant communities, particularly over the long term (Miller and Seastedt, 2009; Briggs et al., 2017; Fornwalt, 2017; Demarest, 2023). Additionally, there is little information available about how long forest restoration treatments remain effective in these systems, i.e. how long the canopy will remain open, and the rate of surface fuel accumulation (Fialko et al., 2020).

Here, we studied the decadal effects of forest restoration treatments in the CFR on surface fuels, forest structure and understory plant communities across two thinning projects using a before/after/control/impact (BACI) framework. The treatments were in ponderosa pine dominated, dry conifer forests that were historically characterized by relatively frequent low and mixed severity fires (Battaglia, 2018). Measurements were taken at 47 plots 1–2 years pre-, 1–2 years post-, 5 years post- and 10 years post-treatment. Our aim was to investigate three research questions. 1) What were the immediate and long-term changes to surface fuels and forest structure as a result of treatments, 2) What were the impacts of thinning on the overall plant functional group composition, diversity, and susceptibility to invasion of the understory, and 3) How did climate, topography and stand structure interact to influence understory plant diversity and functional group composition in the long-term?


## 2. Methods

### 2.1. Study area

This study focuses on two sites: Estes Valley (2109–2573 m) and Phantom Creek (2511–2771 m). These sites, and the plots within them, were established by Briggs et al. (2017) in support of the Colorado Front Range Collaborative Forest Landscape Restoration Program's monitoring effort. The sites are in the montane zone and are dominated by ponderosa pine (*P. ponderosa*) and increasingly Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) at higher elevations. Estes Valley includes three paired treatment and control units that were treated in 2011–2012 (Fig. 1b); Phantom Creek includes three paired treatment and control units that were treated in 2011 (Fig. 1c). All treatments were mechanical thinning, where hand crews and/or specialized machinery were used to remove small diameter trees (< 20 cm at breast height), increase tree spatial heterogeneity, and increase ponderosa pine dominance (Briggs et al., 2017; Cannon, 2018; Barrett, 2021). Estes Valley units are located on Arapaho - Roosevelt National Forest land in both the Big Thompson and St. Vrain watersheds. Phantom Creek units are in the Upper South Platte watershed on the Pike - San Isabel National Forest. On average, Estes Valley receives 514 mm of annual precipitation and Phantom Creek receives 465 mm of annual precipitation (PRISM Climate Group, 2024). Moisture deficit during the study period for each site are shown in Fig. 1d.

### 2.2. Field data collection

This study leverages previously described data collected in 2011 (~1 y pre-treatment), 2012–2013 (~1 year post-treatment) and 2017 (~5 years post-treatment; (Briggs et al., 2017; Demarest, 2023), and builds upon them by adding data collected in 2022–2023 (~10 y post-treatment). Collectively, the data span 47 plots and 4 timesteps,



**Fig. 1.** The study area. Panel a shows the broader location in Colorado. Colored outlines of treatment and control units and points for monitoring locations are in panels b (Estes Valley, Arapaho - Roosevelt National Forest) and c (Phantom Creek, Pike - San Isabel National Forest). panel d is the annual cumulative climatic water deficit for monitoring locations, extracted for each plot, averaged by site (error bars are  $\pm 2$  standard deviations). the vertical dashed lines show the year of treatments (bold red), and years of sampling (black).

totaling 268 observations (Fig. 1). Plots were randomly located and permanently marked within each treatment and control unit. From plot center, all overstory trees (height  $> 1.37$  m and diameter at breast height (DBH)  $> 12.7$  cm) were surveyed within variable radius plots with a  $10 \text{ ft ac}^{-1}$  ( $2.3 \text{ m ha}^{-1}$ ) basal area factor prism. Tree species, status (live/dead), and DBH were recorded for each tree. Tree seedlings ( $< 1.37$  m tall) were inventoried within a 20 m circular subplot around plot center. For untreated controls, tree measurements were not conducted for the 1 year post-treatment timestep as they were assumed to be approximately identical to the pre-treatment timestep. Understory plant cover, as well as fuel cover (fine wood (1-, 10-, and 100-hour fuel), coarse wood (1000-hour fuel), and litter and duff), was quantified with the line-point intercept method. All plant species and fuels were identified every 7.62 cm along four permanently marked 7.5 m transects oriented from plot center in cardinal directions ( $0^\circ$ ,  $90^\circ$ ,  $180^\circ$ ,  $270^\circ$ ). We also searched the entire plot to identify all understory plant species present but not documented along the transects.

Six additional variables were measured only at 10 years after treatment. These were fine wood, measured separately as 1-, 10-, and 100-

hour fuel loads, coarse wood, measured as 1000-hour fuel load, litter and duff fuel load, sapling density, shrub height, and tree canopy cover. Fine wood fuel loading was estimated using the photoload technique (Keane and Dickinson 2007) in 1 square meter quadrats located at plot center, and 7.5 m from plot center in each of the cardinal directions. Coarse wood was measured in a 0.004 ha circular plot around plot center. The diameters of both ends of the log were measured to the nearest 0.25 cm, and the length was measured to the nearest 0.25 cm. Litter and duff depth was measured at 3.05, 6.10, and 9.15 ft along the cardinal directions to the nearest 9.64 cm. Density of tree saplings ( $> 1.37$  m tall, but  $< 12.7$  cm DBH) was measured in a 0.002 ha subplot around plot center. Average shrub height was measured for each of the transects using the LPI method described above for understory plants. Canopy cover was collected at 75 evenly spaced intervals along a 22.86 m N-S transect using a densitometer.

### 2.3. Aggregation of field data

From our field-collected data, we calculated 6 stand structure metrics

that could plausibly impact understory vegetation at each plot for each timestep: crown base height (CBH), quadratic mean diameter (QMD), basal area (BA), tree density, seedling density and cover of fine wood. We derived 10 metrics to characterize aspects of understory composition: the Shannon-Wiener diversity index (Shannon 1948), total cover, total species richness, native cover, native species richness, invasion rate (non-native richness / total richness \* 100), invasion impact (non-native cover / total cover \* 100), forb cover, graminoid cover, and shrub cover.

## 2.4. Ancillary data

We acquired climate and topography data to incorporate into our statistical models (below). For climate at the scale of the treatment unit, we calculated the 30 year (1991–2020) normal of CWD ( $CWD_{norm}$ ), the Z-scores of CWD for the spring (March - June) of sampling ( $CWD_s$ ), and the Z-scores of CWD for the year of treatment ( $CWD_{trt}$ ) for each plot.  $CWD_{norm}$  and  $CWD_{trt}$  were extracted from TopoTerra (Hoecker, 2025). TopoTerra is derived from 4 km resolution TerraClimate (Abatzoglou, 2018) and downscaled to 220 m via the gradient-plus-inverse distance squared method (Flint and Flint, 2012) using Topofire (Holden, 2019) as a template to capture fine-scale spatial variability in climatic water balance. Topofire is derived from a process-based model driven by topographic features (aspect, slope angle, topographic position, soil water holding capacity), and observations of biophysical variables (temperature, precipitation, insolation, and cloud cover). Because TopoTerra is not yet available for 2023, the last year of sampling, we derived  $CWD_s$  Z-scores from TerraClimate and extracted those for each plot at each timestep. We considered 4 km to be sufficient in this case because we expected the interannual variability in CWD to be much greater than the among-site variation within a given year, and furthermore TerraClimate at 4 km has been used in other studies (Wion, 2025) to represent interannual climate variability.

Because our CWD data was at the scale of the treatment unit, we used finer scale topographic data to account for topographic variation within pixels. In addition to in situ measurements of slope, aspect and elevation, we acquired 30 m digital elevation models (DEM) from the shuttle radar topography mission using the *elevatr* R package (Hollister, 2023) in order to calculate topographic indexes that we expected to be associated with understory vegetation, as well as to augment any field measurements that were missed. We used the DEMs to calculate heat load index (HLI) (McCune and Keon, 2002) using the *spatialEco* R package (Evans and Murphy, 2023), and folded aspect and the topographic wetness index (TWI) (Beven and Kirkby, 1979) using the R packages *topmodel* (Buytaert, 2022) and *topomicro* (Mahood, 2024).

## 2.5. Statistical analysis

### 2.5.1. Estimated marginal means

Estimated marginal means and confidence intervals for each metric of stand structure and understory composition at each timestep for control and treated plots were calculated using the *emmeans* package (Lenth, 2025). These were estimated from linear mixed models created using the R package *glmmTMB* (Brooks, 2017). Each model had site (Estes Valley or Phantom Creek), treatment, and timestep as fixed effects, and plot nested within site as a random effect. Site was included as a fixed rather than random effect because random effects with less than six levels are known to produce biased estimates (Oberpriller et al., 2022).

### 2.5.2. Change from pre-treatment

Having a BACI experimental design allowed us to calculate the difference from pre-treatment values for each metric we analyzed, and then test both whether the change from pre-treatment was significant, as well as whether treatment had a significant effect on the change from pre-treatment. Statistical tests on the difference from pre-treatment values rather than raw values account for the possibility of large differences in

means among sites that can obscure treatment effects when only looking at means by timestep. We used one-sided Wilcoxon Rank Sum tests (Bauer, 1972) to test whether a given treatment by post-treatment timestep combination changed significantly from pre-treatment. Treatment effects for changes from pre-treatment were calculated from linear mixed models that were fit with difference from pre-treatment as the response variable, and the same fixed and random effects as above.

For the six aspects of stand structure that were measured only at the 10-year timestep (fuel loads of coarse wood, fine wood, litter/duff, sapling density, tree canopy cover, and shrub height), we created generalized linear mixed models for each variable with site and treatment as fixed effects and plot as a random effect, estimated the marginal means, and used Wald Chi-square tests to assess significance (Fox and Weisberg, 2019).

### 2.5.3. Species accumulation curves

In order to assess how forest restoration treatments affected the regional species pools, we created species accumulation curves for each timestep and treatment combination, and estimated the total species pool using asymptotic richness estimators (Smith and Van Belle, 1984) using the *vegan* R package (Oksanen, 2022). While species richness measured at the plot scale is a commonly used metric of alpha diversity, it has the problem of scale-dependency. Species accumulation curves are a way to overcome these issues (Gotelli and Colwell, 2001; Thompson and Withers, 2003). Extrapolating the curve to its asymptote yields an estimated species pool, or gamma diversity. The slope is analogous to beta diversity, and a gradual ascent to the asymptote indicates a high proportion of rare species.

### 2.5.4. Compositional change

We conducted two permutational multivariate analysis of variance (PERMANOVA) (Anderson 2001) tests to assess the change in community composition in response to treatments. For the first test we excluded the 5 and 10 year post-treatment timesteps and tested the differences between the pre- and 1 year post-treatment phases. For the second test, we used all data and tested the difference between the pre- and 1 year post-treatment timesteps grouped, and the 5 and 10 year post-treatment timesteps grouped. For both models, we had site and treatment as additional independent variables, and had 999 permutations per model.

### 2.5.5. Modeling the effect of treatments on understory plants

We used a mixed modelling approach to explore how much the treatment itself affects understory plant communities, versus how much the changes in forest structure resulting from thinning affect understory plant communities. In order to explore how the changes in structure that resulted from treatments affected the understory, we created mixed effects models for those understory metrics that had treatment effects 5 and 10 years after treatment. These were P(invasion), invasion rate, graminoid cover, forb cover, native richness and native cover (see results). The model for P(invasion) had a binomial error distribution, the models for native cover, graminoid cover and invasion rate had beta error distributions, and the model for native species richness had a Poisson error distribution. For each response variable, we created two models. The first model was a 'naive' model with treatment, timestep and their interaction as predictors. The second model was an 'informed' model that used AIC-based model selection procedures (Zuur, 2009) to select the strongest predictors from metrics of stand structure (BA, QMD, tree density, seedling density, and fine wood), topography (HLI, TWI, slope, and elevation) and water balance ( $CWD_{norm}$ ,  $CWD_{trt}$  and  $CWD_s$ ). Plot identity was used as a random effect in all models. Each candidate model was fit with maximum likelihood, candidate models were compared with AIC and diagnosed using tools from the DHARMA and performance packages to check for violations of assumptions, including normality of residuals, normality of random effects, linear relationship, homogeneity of variance, and multicollinearity (Lüdecke, 2021; Hartig, 2024), and the final model was refit with restricted maximum

likelihood. Significance of the predictors for each final model was assessed using Wald Chi-squared tests using the R package *car* (Fox and Weisberg, 2019). We then compared the marginal  $R^2$  (the variance explained by fixed effects (Nakagawa and Schielzeth, 2013)) for both sets of models to evaluate how well each set of predictors influenced each metric of understory plant composition.

### 3. Results

#### 3.1. Stand structure and fuels changes

Treatment effects on changes from pre-treatment were statistically significant for QMD, BA and tree density at all post-treatment timesteps, and all three metrics remained near immediate post-treatment values 5 and 10 years after treatment (Fig. 2). Mean BA in treated plots was 18–22  $\text{m}^2 \text{ha}^{-1}$  post-treatment, which was 6.5  $\text{m}^2 \text{ha}^{-1}$  below pre-treatment values on average. Mean tree density in treated plots ranged from 268 to 368  $\text{trees ha}^{-1}$  after treatment, an average of 305  $\text{trees ha}^{-1}$  lower than pre-treatment values. Mean QMD in treated plots was 31.6–33.6 cm post-treatment, 3.3 cm higher than pre-treatment values, on average. QMD also increased in treated plots from an average of 31.6 cm 1 year post-treatment to 33.6 cm ten years after treatment, an increase of 0.3 cm  $\text{year}^{-1}$  ( $p < 0.05$ ). Cover of fine wood in treated plots

was approximately 3 % higher than pre-treatment values 1 and 5 years after treatment but was not different from pre-treatment at year 10 (Fig. 2). Crown base height did not have significant treatment effects, and was only significantly higher than pre-treatment in year 5.

Seedling density had significant treatment effects for years 5 and 10 (Fig. 2; Table 1), but this was caused by a decrease in control plot post-treatment values during these later years. Untreated controls had an average of 7420 seedlings  $\text{ha}^{-1}$  before treatments and 8930 seedlings  $\text{ha}^{-1}$  in the year after treatments, and this declined to 1830 and 1740 seedlings  $\text{ha}^{-1}$  5 and 10 years after treatment, respectively (Table 2). For those metrics that were measured only in year ten, fuel loads of coarse wood and fine wood were higher in treated plots, while tree canopy cover and sapling density were lower than control plots (Fig. 4).

#### 3.2. Understory changes after treatments

One year after treatment, there was a decrease in total species richness for both control and treated plots and a decrease in shrub cover and total cover in treated plots (Fig. 4; Table 2). Five years after treatment, there were increases in graminoid cover, total cover, total richness and invasion rate in the treated plots that were significantly greater than any increases that occurred in the control plots, and these increases remained in year 10. Total cover and total richness were also




Fig. 2. Change from pre-treatment for six stand structure metrics. Black outlines indicate that the difference from pre-treatment values was significant ( $p < 0.05$ ) according to a one-sample Wilcoxon rank sum test (each timestep by treatment combination was tested individually). stars indicate a significant treatment effect in a given timestep according to a linear mixed model. Y-axis units are indicated in panel titles. Note that basal area, crown base height, quadratic mean diameter, seedling density, and tree density values are not available for control plots 1 year post-treatment because tree measurements were not conducted during this timestep.

**Table 1**

Estimated marginal means for stand structure metrics at each timestep, according to a generalized linear mixed model. Numbers in parenthesis are low and high 95 % confidence intervals.

| Response                                       | Treatment | Timestep               |                        |                   |                     |
|------------------------------------------------|-----------|------------------------|------------------------|-------------------|---------------------|
|                                                |           | Pre-Treatment          | 1 Year Post            | 5 Year Post       | 10 Year Post        |
| Tree Density (trees ha <sup>-1</sup> )         | Control   | 609 (500, 718)         | 607 (497, 716)         | 749 (640, 859)    | 618 (508, 727)      |
|                                                | Treatment | 583 (478, 688)         | 340 (235, 445)         | 369 (263, 475)    | 271 (165, 376)      |
| Quadratic Mean Diameter (cm)                   | Control   | 29.3 (27, 32)          | 29.3 (27, 32)          | 29.2 (27, 31)     | 30.0 (28, 32)       |
|                                                | Treatment | 29.2 (27, 31)          | 31.6 (29, 34)          | 32.3 (30, 35)     | 33.6 (31, 36)       |
| Basal Area (m <sup>2</sup> ha <sup>-1</sup> )  | Control   | 25.6 (22, 29)          | 25.4 (22, 29)          | 31.2 (28, 34)     | 26.3 (23, 29)       |
|                                                | Treatment | 26.4 (23, 29)          | 18.8 (16, 22)          | 22.2 (19, 25)     | 18.1 (15, 21)       |
| Fine Wood (%)                                  | Control   | 14 (11, 17)            | 8 (5, 11)              | 11 (8, 14)        | 9 (6, 12)           |
|                                                | Treatment | 11 (8, 14)             | 15 (12, 18)            | 14 (12, 17)       | 9 (7, 12)           |
| Crown Base Height (m)                          | Control   | 42 (36, 48)            | 42 (36, 48)            | 46 (40, 52)       | 44 (38, 51)         |
|                                                | Treatment | 43 (37, 49)            | 45 (39, 51)            | 47 (41, 54)       | 43 (37, 50)         |
| Seedling Density (seedlings ha <sup>-1</sup> ) | Control   | 7410<br>(4140, 10,678) | 8930<br>(5665, 12,204) | 1830<br>(0, 5101) | 1740<br>(0, 5006)   |
|                                                | Treatment | 3550<br>(419, 6690)    | 3270<br>(83, 6448)     | 2800<br>(0, 5989) | 3520<br>(337, 6706) |

**Table 2**

Estimated marginal means of probability of invasion, invasion rate, invasion impact, total cover, total richness, graminoid cover, forb cover and shrub cover at each of the four timesteps in treated and control plots. Upper and lower ends of 95 % confidence intervals are in parentheses.

| Response               | Treatment | Timestep           |                    |                   |                   |
|------------------------|-----------|--------------------|--------------------|-------------------|-------------------|
|                        |           | Pre-Treatment      | 1 Year Post        | 5 Year Post       | 10 Year Post      |
| P(Invasion)            | Control   | 0.24 (0.082, 0.53) | 0.1 (0.022, 0.33)  | 0.52 (0.25, 0.77) | 0.86 (0.62, 0.96) |
|                        | Treatment | 0.18 (0.059, 0.43) | 0.13 (0.039, 0.37) | 0.89 (0.66, 0.97) | 0.97 (0.82, 0.99) |
| Invasion Rate (%)      | Control   | 1.58 (0.46, 2.7)   | 0.96 (0, 2.1)      | 2.65 (1.5, 3.8)   | 3.47 (2.4, 4.6)   |
|                        | Treatment | 1.09 (0.014, 2.2)  | 0.88 (0, 2)        | 6.66 (5.6, 7.8)   | 5.86 (4.8, 7)     |
| Invasion Impact (%)    | Control   | 0.27 (0, 1.9)      | 0.68 (0, 2.3)      | 1.25 (0, 2.9)     | 1.81 (0.18, 3.4)  |
|                        | Treatment | 1.82 (0.26, 3.4)   | 0.16 (0, 1.7)      | 1.99 (0.4, 3.6)   | 1.13 (0, 2.7)     |
| Total Cover (%)        | Control   | 20.47 (15, 26)     | 18.93 (13, 25)     | 32.36 (27, 38)    | 19.51 (14, 25)    |
|                        | Treatment | 18.56 (13, 24)     | 11.05 (5.7, 16)    | 37.79 (32, 43)    | 24.47 (19, 30)    |
| Total Richness (count) | Control   | 35.26 (32, 39)     | 32.43 (29, 36)     | 36.04 (33, 40)    | 39.69 (36, 43)    |
|                        | Treatment | 31.88 (28, 35)     | 30.12 (27, 34)     | 38.76 (35, 42)    | 41.01 (38, 44)    |
| Graminoid Cover (%)    | Control   | 3.77 (1.4, 6.1)    | 3.75 (1.4, 6.1)    | 8.46 (6.1, 11)    | 4.09 (1.7, 6.4)   |
|                        | Treatment | 2.78 (0.52, 5)     | 2.87 (0.61, 5.1)   | 13.18 (11, 15)    | 7.16 (4.9, 9.4)   |
| Forb Cover (%)         | Control   | 2.75 (0.87, 4.6)   | 2.37 (0.49, 4.2)   | 5.08 (3.2, 7)     | 2.98 (1.1, 4.9)   |
|                        | Treatment | 1.93 (0.12, 3.7)   | 1.6 (0, 3.4)       | 7.88 (6, 9.7)     | 4.04 (2.2, 5.9)   |
| Shrub Cover (%)        | Control   | 9.46 (6.4, 12)     | 8.8 (5.8, 12)      | 12.01 (9, 15)     | 7.92 (4.9, 11)    |
|                        | Treatment | 8.69 (5.8, 12)     | 5.33 (2.4, 8.2)    | 13.53 (11, 16)    | 9.94 (7, 13)      |

significantly increased from pre-treatment in treated plots 5 and 10 years after treatment, and were significantly higher than control plots.

Probability of invasion in both the control and treatment plots ranged from 0.2 or less in the pre- and 1 year post-treatment timesteps, and increased to 0.47–0.97 in the 5 and 10 year timesteps (Table 2). In treated plots, the differences between pre- and 1 year post-treatment timesteps and the differences between 5 and 10 year post-treatment timesteps were not significant, but all others were (Table S1). In control plots, the only significant differences were among the pre-treatment and 10 year post-treatment timesteps, and among the 1 year post-treatment and the 10 year post-treatment timesteps (Table S1).

There was little difference in the regional species pool estimations immediately before and after treatment, but native and non-native species pools increased after 5 years (Fig. 5). Ten years after treatment, there were an estimated 209 species (from 149 pre-treatment) in treated areas, compared to 177 (from 165) in control plots (Fig. 5,

Table 3). The non-native species pool increased from an estimated 5–15 in treated areas, and from 7 to 10 species in control plots increased. Native species increased from 151 to 162 species in control plots, and from 140 to 187 in treated areas.

### 3.2.1. Compositional change

Our PERMANOVA models indicated that the composition remained largely unchanged between pre-treatment and 1 year post-treatment timesteps, but changed significantly after 5 years (Table 4) but with a small effect size. Site (Estes Valley or Phantom Creek) was significant in both models and explained 11.6–12.8 percent of the variance. In the comparison of pre-treatment and 1 year post-treatment timesteps, both treatment and timestep were not significant. In the comparison of 5 and 10 year timesteps against pre- and 1 year post-treatment, both treatment and timestep were significant and explained 1.8 and 9.9 percent of the variance, respectively.

**Table 3**

Estimates of the regional species pool based on extrapolating the species accumulation curve to its asymptote. Standard errors are in parentheses.

|            | Treatment | Pre-Treatment | 1 Year Post | 5 Year Post | 10 Year Post | 10 year change |
|------------|-----------|---------------|-------------|-------------|--------------|----------------|
| Non-native | Control   | 7 (1)         | 6 (1)       | 6 (1)       | 9 (1)        | + 2            |
|            | Treatment | 5 (1)         | 5 (1)       | 14 (1)      | 13 (1)       | + 8            |
| Native     | Control   | 151 (6)       | 144 (8)     | 145 (5)     | 162 (7)      | + 11           |
|            | Treatment | 140 (6)       | 130 (6)     | 161 (8)     | 187 (11)     | + 47           |
| Total      | Control   | 165 (7)       | 154 (9)     | 153 (6)     | 176 (8)      | + 11           |
|            | Treatment | 149 (7)       | 140 (6)     | 181 (9)     | 208 (12)     | + 59           |

**Table 4**

PERMANOVA results. In the top table, community data was filtered to only pre- and 1 year post-treatment timesteps. In the bottom table, all timesteps were used, and the difference between both 5 and 10 year post-treatment timesteps against pre- and 1 year post-treatment were tested.

| Comparison                                                      | Term                      | Df  | Sum Of Squares | R <sup>2</sup> | F     | Pr (>F) |
|-----------------------------------------------------------------|---------------------------|-----|----------------|----------------|-------|---------|
| Pre-Treatment versus 1 Year Post-Treatment                      | Treatment                 | 1   | 0.217          | 0.00878        | 0.948 | 0.454   |
|                                                                 | Site                      | 1   | 3.17           | 0.128          | 13.9  | 0.001   |
|                                                                 | 1 year post-treatment     | 1   | 0.248          | 0.0101         | 1.09  | 0.345   |
|                                                                 | Residual                  | 92  | 21             | 0.853          |       |         |
|                                                                 | Total                     | 95  | 24.7           | 1              |       |         |
| Pre- & 1 Year Post-Treatment versus 5 & 10 Years Post-Treatment | Treatment                 | 1   | 0.948          | 0.0178         | 4.33  | 0.001   |
|                                                                 | Site                      | 1   | 6.14           | 0.116          | 28    | 0.001   |
|                                                                 | 5–10 years post-treatment | 1   | 5.24           | 0.0988         | 23.9  | 0.001   |
|                                                                 | Residual                  | 185 | 40.5           | 0.763          |       |         |
|                                                                 | Total                     | 189 | 53.1           | 1              |       |         |

### 3.2.2. Associations of stand structure, topography and climate with understory vegetation

For all response variables, the variance explained by fixed effects ( $R_m^2$ ) for the models informed by stand structure, topography and climate was greater than or equal to the variance explained by the naive models with treatment and timestep as predictors. The difference in  $R_m^2$  values for the two model sets ranged from 0 % to 40 % of the variance explained. The informed model of native cover had an  $R_m^2$  of 0.35 while that of the naive model was 0.35. Native cover was positively associated with fine wood, QMD, slope and TWI, and negatively associated with  $CWD_s$  (Fig. 6a, Table 5). The informed model of native species richness had an  $R_m^2$  of 0.41 while that of the naive model was 0.09. Native species richness was negatively associated with  $CWD_s$ , HLI and tree density, positively associated with total vegetation cover, and had a hump-shaped relationship with  $CWD_{norm}$  (Fig. 6b, Table 5). Forb cover had an  $R_m^2$  of 0.76 for the informed model compared to 0.34 for the naive model. Forb cover was positively associated with total vegetation cover, negatively associated with tree density,  $CWD_s$  and HLI, and its relationship with  $CWD_{norm}$  was hump-shaped (Fig. 6c, Table 5). The informed model of graminoid cover had an  $R_m^2$  of 0.54 while that of the naive model was 0.42. Graminoid cover was negatively associated with  $CWD_s$ , HLI and tree density, positively associated with BA, and had a hump-shaped relationship with  $CWD_{norm}$  (Fig. 6d, Table 5). For P(invasion), the informed model had an  $R_m^2$  of 0.56 while the naive model had an  $R_m^2$  of 0.30. P(invasion) was negatively associated with  $CWD_s$  and  $CWD_{trt}$ , positively associated with total vegetation cover, and had a hump-shaped relationship with  $CWD_{norm}$  (Fig. 6e, Table 5). For invasion rate, the informed model had an  $R_m^2$  of 0.86 while that of the naive model was 0.58. Invasion rate was negatively associated with  $CWD_s$ ,  $CWD_{trt}$  and tree density, and positively associated with total vegetation cover (Fig. 6f, Table 5).

## 4. Discussion

This work adds to the growing consensus that forest restoration treatments can benefit native understory plant communities (Vernon, 2023; Springer, 2024), while also increasing the probability of non-native plant invasion (Willms, 2017). While the immediate effects of treatments on native understory plant cover and diversity were minimal, after 5 years largely beneficial impacts were observed, and these benefits remained unchanged or increased after 10 years (Figs. 4–5). These outcomes were associated with greater QMD and lower BA and tree density (Fig. 6)—common goals of restoration treatments (Larson and Churchill, 2012; Addington, 2018). However, the treatments also resulted in invasions of non-native plants. These invasions became detectable after 5 years, but likely had a low level of

**Table 5**

Significance of generalized linear model coefficients as tested by a wald chi-squared test.

| Response        | Treatment + Timestep Model Term | X <sup>2</sup> | Stand Structure, Topography, Climate Model |                |
|-----------------|---------------------------------|----------------|--------------------------------------------|----------------|
|                 |                                 |                | Term                                       | X <sup>2</sup> |
| Forb Cover      | Timestep                        | 169.97***      | Total Cover                                | 371.52***      |
|                 | Treatment                       | 0.45           | CWD Normal                                 | 10.23**        |
|                 | Timestep x Treatment            | 31.06***       | CWD(z) Sample Year                         | 5.94*          |
|                 |                                 |                | Heat Load                                  | 14.39***       |
| Graminoid Cover | Timestep                        | 225.08***      | Tree Density                               | 8.13**         |
|                 |                                 |                | CWD Normal                                 | 6.09*          |
|                 | Treatment                       | 1.99           | CWD(z) Sample Year                         | 10.89***       |
|                 | Timestep x Treatment            | 32.79***       | Heat Load                                  | 5.49*          |
| Invasion Rate   |                                 |                | Tree Density                               | 13.85***       |
|                 | Timestep                        | 56.82***       | Basal Area                                 | 4.44*          |
|                 | Treatment                       | 3.97*          | Tree Density                               | 7.62**         |
|                 | Timestep x Treatment            | 10.84*         | CWD(z) Treatment Year                      | 22.67***       |
| Native Cover    |                                 |                | CWD(z) Sample Year                         | 17.89***       |
|                 | Timestep                        | 210.98***      | Total Cover                                | 14.95***       |
|                 | Treatment                       | 0              | CWD(z) Sample Year                         | 25.21***       |
|                 | Timestep x Treatment            | 37.1***        | Fine Wood Cover                            | 3.03.          |
| Native Richness |                                 |                | Quadratic Mean Diameter                    | 5.09*          |
|                 | Timestep                        | 32.89***       | Slope                                      | 4.25*          |
|                 | Treatment                       | 0.3            | Topographic Wetness                        | 4.01*          |
|                 | Timestep x Treatment            | 3.5            | CWD Normal                                 | 26.87***       |
| P(Invasion)     |                                 |                | CWD(z) Sample Year                         | 9.09**         |
|                 | Timestep                        | 34.5***        | Heat Load                                  | 27.98***       |
|                 | Treatment                       | 2.2            | Tree Density                               | 17.28***       |
|                 | Timestep x Treatment            | 4.24           | Total Cover                                | 13.76***       |
|                 |                                 |                | CWD(z) Sample Year                         | 26.73***       |
|                 |                                 |                | CWD Normal                                 | 12.47**        |
|                 |                                 |                | CWD(z)                                     | 24.21***       |
|                 |                                 |                | Treatment Year                             |                |
|                 |                                 |                | Total Cover                                | 8.21**         |

p < 0.1, \* p < 0.05, \*\* p < 0.01, \*\*\* p < 0.001

ecological impact. Probability of invasion and invasion rate were less strongly associated with stand structure metrics than they were with climate (Fig. 6) or with treatment per se, indicating that forest structure resulting from treatments may have less influence over non-native plant invasions than regional climate and the physical disturbance and potential for propagule introduction due to treatment.

### 4.1. Treatment effects on forest structure persisted for 10 years

Mechanical treatment at Estes Valley and Phantom Creek resulted in lower tree densities, lower BAs, and higher QMDs in treated plots relative to untreated controls. These values remained near their immediate post-treatment values 5 and 10 years after treatment (Fig. 2), indicating that treatments showed minimal signs of returning to pre-treatment conditions. While treated plots remained significantly different from pre-treatment conditions, they still had higher residual tree density and BA, but similar QMD, than dendrochronological reconstructions of forest structure as it was just before Euro-American settlement (Battaglia, 2018). Plots treated here may have had higher targeted basal areas and densities than the regional averages, which may be due to the treatment projects being situated on the higher end of the productivity gradient. It

is also possible that these treatments, which were among the first to occur during the 10-year lifetime of the Colorado Front Range Collaborative Forest Landscape Restoration Program, were constrained by decision documents that predated emerging research on historical forest structures and adaptively developing forest restoration objectives (Barrett, 2021). In the more arid areas of western US dry mixed conifer forests, forest restoration treatments have been observed to make trees more resilient against drought (McCauley, 2022), and enhance tree-level growth across sites spanning ponderosa pine's climatic niche, even during extreme drought (Knapp, 2021; Zald, 2022; Rodman, 2024), and perhaps even increase carbon storage in the long-term (McCauley, 2019; Doughty, 2021). Our observations of increased QMD through time after treatment suggest that these findings likely hold true in the CFR as well.

The primary mechanism by which treatments lose their effectiveness is for seedlings, saplings and shrubs to grow in the newly opened canopies and create ladder fuels, or for surface fuels to accumulate (Wasserman, 2022). We did not detect any signs of increased ladder fuels. There were minimal differences in seedling density (Fig. 2) and shrub cover (Fig. 4), as well as lower sapling density in treated plots at year 10 (Fig. 3). Seedling density in untreated control plots was much higher than in treated plots in pre- and 1 year post-treatment, but not in years 5 and 10 (Table 1). Climatic water deficit was much higher than normal in the year of treatment, and there was only one wetter than normal year (2015) within 5 years of treatment (Fig. 1d), so it is possible that there was a masting event in some of the control plots at or before the start of the pre-treatment monitoring, followed by a dieoff (Wion, 2020). Prior research has found that warming temperatures and drought may inhibit regeneration in *P. ponderosa* and *P. menziesii* forests (Rother et al., 2015) (Davis, 2019; Stevens-Rumann and Morgan, 2019). Lack of seedling and sapling regeneration may increase the longevity of treatment effectiveness. The only evidence we found of declining treatment effectiveness for these plots was increased surface fuel load. While the cover of fine wood returned to pre-treatment values by year 10 (Fig. 2), the fuel load of fine and coarse wood was significantly higher in treated plots than in untreated controls 10years post-treatment (Fig. 3).

#### 4.2. Native species richness and cover were higher in treated plots

Native understory plant richness was largely unchanged immediately after treatment (also see (Briggs et al., 2017), but then increased in diversity in years 5 and 10 (Figs. 4, 5, Table S3). Composition followed the same pattern (Table 4). The contrast between species accumulation curves increasing from year 5 to 10 and plot-level richness remaining constant is an artifact of the scale-dependency of species richness (Gotelli and Colwell, 2001; Thompson and Withers, 2003). While similar numbers of species were encountered at each plot at timesteps 5 and 10, the particular species encountered at each plot were more likely to be different by year 10. The gradual curve to the asymptote in the native species accumulation curves (Fig. 4) suggest that plots are diversifying by adding new species (Gotelli and Colwell, 2001; Thompson and Withers, 2003). Therefore, the increased heterogeneity created by the treatments, along with spatial variation along resource gradients, may be creating suitable habitat for a wider range of species. Facilitation among native species allows the community to continue to assemble and diversify through time (Dovčiak and Halpern, 2010). The increase in plot-level native species richness may be explained by increased above- and below-ground resources available for utilization by the understory (McConnell and Smith, 1970; Riegel et al., 1992; Zou, 2008). In addition to spatial variation in resource gradients, some differences in observed species diversity were partially explained by temporal variation in water availability (Fig. 1d). However, if this were the main driver we would also expect to see differences between timesteps 5 and 10 for changes in native richness. Furthermore, while native richness had a negative association with CWD<sub>s</sub>, it had stronger associations with HLI, total vegetation cover, CWD<sub>norm</sub> and tree density (Fig. 6), suggesting that higher richness values were more strongly influenced by the changes in stand structure than sample year climate.

Native cover and total vegetation cover were higher in treated units than untreated controls in years 5 and 10. While the effect size associated with CWD<sub>s</sub> was smaller than those of the stand structure and topographic variables, the wettest CWD<sub>s</sub> values were associated with a 75 % increase in native cover over the driest year, and it had stronger statistical support (Table 5). So while these site and stand structure





Fig. 3. Mean comparisons for variables measured only ten years after treatment. Stars indicate significant differences from a linear mixed model.



**Fig. 4.** Understory changes from pre-treatment. dark outlines indicate that the change from pre-treatment values was significant ( $p < 0.05$ ) according to a one-sample Wilcoxon rank sum test (each timestep by treatment combination was tested individually). stars indicate a significant treatment effect in a given timestep according to a linear mixed model. Y-axis units are indicated in panel titles.

metrics strongly influence the potential cover of native plants, there is high year-to-year variation in cover, and this is to be expected in semi-arid systems with high interannual climate variability. Graminoid cover was higher than untreated controls in years 5 and 10 (Fig. 4). Higher graminoid cover may be desirable in cases where a return to historical frequent fire is desirable (Brown and Smith, 2000; Veblen et al., 2000;

Schoennagel et al., 2011). Most of the forb and graminoid species encountered here were native, and non-native cover was very low in all cases. Forb cover was more strongly associated with aspect and total vegetation cover, while graminoid cover was comparatively more strongly associated with stand metrics. They both had negative relationships with CWD<sub>s</sub> that were weak, but with strong statistical




Fig. 5. Species accumulation curves. Vertical bars represent the standard error of the estimated species count.

support (Table 5), indicating that, like native cover, stand structure and topography are broad determinants of niche space, and interannual variability in abundance is associated with yearly fluctuations in temperature and moisture. There was no sign of a return to pre-treatment cover as has been found in more mesic conifer systems (Jang, 2021).

#### 4.3. Non-native species richness remained constant after 5 years


Taken together, our data indicate that forest restoration treatments in the CFR can result in introductions of non-native species in previously uninvaded areas. These new invasions were mostly detectable after 5 years and remained at very low values of relative cover (Table 2; Fig. 4) indicating low ecological impact (stage 2 from the model of (Brooks, 2004)). There were minimal changes in probability of invasion immediately after treatment, but it was higher in treated plots than it was in untreated controls in years 5 and 10 (Table 2). Probability of invasion was strongly associated with  $CWD_s$ ,  $CWD_{trt}$  and  $CWD_{norm}$ , as well as total vegetation cover. The strongest effects with the most statistical support were  $CWD_{trt}$  and  $CWD_a$ . The negative association with  $CWD_z$  suggests that annual fluctuations in climate drive interannual variability in abundance, and since these invasions were all at very low levels of impact (Table 2), many species may have been at very low population levels or remained dormant, or ungerminated in the seed bank (Faist et al., 2015). The negative association with  $CWD_{trt}$  was strong, however at all the sites were treated during the same year, which had high  $CWD_z$  scores. So while the negative association between  $CWD_{trt}$  suggests that treatments conducted during abnormally hot and dry years will have a lower chance of being invaded, these results should be interpreted with caution, and more research across a broader range of treatment conditions is needed. The stand structure and climate model explained more variation than just the naive model, indicating the role that climate played in constraining the introduction of species was stronger than the act of treatment alone. Control plots also had higher risk of invasion in the 5 and 10 year timesteps, but less so than treatments (Table 2). Many

of the non-native species we encountered were common across the western US (Table S2). Over the broader spatial scale of the western US, many of these species have a residence time of decades if not centuries, and may be in the midst of an ongoing invasion to fill out their potential range (Richardson and Pyšek, 2006; Wilson, 2007; Rouget, 2016).

Analysis of species accumulation curves and extrapolated species pools yielded a regional species pool estimated at 10 non-native species across control plots and 15 in treated areas (Fig. 5, Table 3). The overlap in species accumulation curves for years 5 and 10 suggests that colonization by non-natives via the newly available resources had mostly taken place by the 5 year mark. After treatment there are newly available resources, including increased light and water availability from the opened canopy (McConnell and Smith, 1970; Riegel et al., 1992), and increased bare soil from physical soil disturbance (Leffler, 2016).

#### 4.4. Management implications

It has long been recognized that monitoring and control of non-native plants need to be part of any forest restoration treatment prescription (Schwilke, 2009). While there are long-term observations of non-native populations remaining stable after 20 years from similar treatment experiments in the northern Rockies (Hood et al., 2024) and southwest (Springer, 2024), there are also observations of abrupt increases in non-native populations within 10 years of being at low-impact levels like those observed here (McGlone et al., 2009; Wion, 2024) that proceeded in the stages outlined by Brooks (2004). In a *P. ponderosa* forest in Northern Arizona, McGlone et al. (2009) observed *B. tectorum* at low cover (stage 2) before and three years after treatment, followed by dominant levels of cover (stage 3) 6 years after treatment without a subsequent disturbance. In another example from a *P. ponderosa* forest in New Mexico, *Bromus inermis* L., which was present in our plots and is common in the CFR, was observed to increase rapidly from low levels like those observed here (stage 2) in 1997 to become the dominant understory species in 2008 (stage 3) (Wion, 2024). When those plots



**Fig. 6.** Partial effects of GLMMs. for each panel, the bold line represents the mean estimate of the partial effect, and the colored area surrounding the line is the 95 % confidence interval.

subsequently burned in 2011, non-native grass cover, which was mostly *B. inermis*, was associated with higher burn severity, suggesting a transition to stage 4.

Most of the observed changes in understory composition took place after 5 years and remained steady 10 years after treatment. This is consistent with studies from other regions, where the first 5 years after treatment saw changes in composition, and then the composition remained stable for up to 23 years (Jang, 2021; Vernon, 2023). The time period between 5 and 10 years after treatment is likely the most effective time to monitor and control non-native plants, as it is a long enough period of time for the understory to reach the new equilibrium in productivity, and for non-native populations to become detectable. It is also short enough that non-native populations that establish after treatment are likely to still be small and manageable (Schuurman, 2020). After 10

years, it is unknown whether populations of introduced plants will continue to grow or remain stable without further impacts.

It is important to understand the efficacy and impact of thinning-only forest restoration prescriptions. Literature reviews have largely supported the notion that forest restoration treatments reduce subsequent wildfire severity, however the available literature consistently shows greater severity reductions when thinning is paired with prescribed fire (Kallies and Yocom Kent, 2016; Davis, 2023). However, in the CFR prescribed fires are less tractable due to high topographic complexity and relief, and the high prevalence of low-density residential development in forested areas (Ryan et al., 2013). Lower tree density and BA, and higher QMD were associated with desirable outcomes of higher native cover, graminoid cover and native species richness following mechanical thinning (Fig. 6). Here, desirable understory attributes were

associated with stand structure attributes that are often targets for fire behavior goals, and invasion probability and impact were lower when treatment year climate was hotter and drier.

### CRediT authorship contribution statement

**Scott M. Ritter:** Writing – review & editing, Writing – original draft, Methodology, Investigation, Funding acquisition, Conceptualization. **David M. Barnard:** Writing – review & editing, Writing – original draft, Supervision, Resources, Project administration, Investigation, Funding acquisition, Conceptualization. **Camille S. Stevens-Rumann:** Writing – review & editing, Writing – original draft, Supervision, Project administration, Investigation, Funding acquisition, Conceptualization. **Allison E. Rhea:** Writing – review & editing, Writing – original draft, Visualization, Investigation, Funding acquisition, Conceptualization. **Adam L. Mahood:** Writing – review & editing, Writing – original draft, Visualization, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization. **Kevin Barrett:** Conceptualization, Data curation, Methodology, Writing – review & editing. **Paula J. Fornwalt:** Conceptualization, Methodology, Writing – review & editing.

### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Acknowledgements

This project was funded by Colorado Forest Restoration Institute appropriations funding, and Joint Fire Science Program grant #23–201–10: Using landscape features, plot measurements, and remote sensing data to improve predictions of fuels treatment longevity in the Colorado Front Range. We are grateful to Tyler Hoecker and Jeffrey Chandler for assistance with access to the TopoTerra data product, and field crews from the US Geological Survey, the Rocky Mountain Research Station, and the Colorado Forest Restoration Institute for field data collection.

### Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at [doi:10.1016/j.foreco.2025.122881](https://doi.org/10.1016/j.foreco.2025.122881).

### Data availability

Data and code are shared via DOI

### References

- Abatzoglou, J.T., et al., 2018. TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958–2015 (Available at). *Sci. Data* 5 (1), 170191. <https://doi.org/10.1038/sdata.2017.191>.
- Abatzoglou, J.T., Williams, A.P., 2016. Impact of anthropogenic climate change on wildfire across Western US forests (Available at). *Proc. Natl. Acad. Sci.* 113 (42), 11770–11775. <https://doi.org/10.1073/pnas.1607171113>.
- Abella, S.R., Springer, J.D., 2015. Effects of tree cutting and fire on understory vegetation in mixed conifer forests (Available at). *For. Ecol. Manag.* 335, 281–299. <https://doi.org/10.1016/j.foreco.2014.09.009>.
- Addington, R.N. et al. (2018) *Principles and practices for the restoration of ponderosa pine and dry mixed-conifer forests of the Colorado Front Range*. RMRS-GTR-373. Ft. Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, p. RMRS-GTR-373. Available at: <https://doi.org/10.2737/RMRS-GTR-373>.
- Agee, J.K., Skinner, C.N., 2005. Basic principles of forest fuel reduction treatments (Available at). *For. Ecol. Manag.* 211 (1–2), 83–96. <https://doi.org/10.1016/j.foreco.2005.01.034>.
- Balandier, P., et al., 2022. Tamm review: forest understorey and overstorey interactions: so much more than just light interception by trees (Available at). *For. Ecol. Manag.* 526, 120584. <https://doi.org/10.1016/j.foreco.2022.120584>.
- Barrett, K.J., et al., 2021. Effects of collaborative monitoring and adaptive management on restoration outcomes in dry conifer forests (Available at). *For. Ecol. Manag.* 488, 119018. <https://doi.org/10.1016/j.foreco.2021.119018>.
- Battaglia, M.A., et al., 2018. Changes in forest structure since 1860 in ponderosa pine dominated forests in the Colorado and Wyoming front range, USA (Available at). *For. Ecol. Manag.* 422, 147–160. <https://doi.org/10.1016/j.foreco.2018.04.010>.
- Bauer, D.F., 1972. Constructing confidence sets using rank statistics (Available at). *J. Am. Stat. Assoc.* 67 (339), 687–690. <https://doi.org/10.1080/01621459.1972.10481279>.
- Beven, K.J., Kirkby, M.J., 1979. A physically based, variable contributing area model of basin hydrology / un modèle à base physique de zone d'appel variable de l'hydrologie du bassin versant (Available at). *Hydrol. Sci. Bull.* 24 (1), 43–69. <https://doi.org/10.1080/02626667909491834>.
- Briggs, J.S., Fornwalt, P.J., Feinstein, J.A., 2017. Short-term ecological consequences of collaborative restoration treatments in ponderosa pine forests of Colorado (Available at). *For. Ecol. Manag.* 395, 69–80. <https://doi.org/10.1016/j.foreco.2017.03.008>.
- Brooks, M., E., et al., 2017. glmmTMB balances speed and flexibility among packages for Zero-inflated generalized linear mixed modeling (Available at). *R. J.* 9 (2), 378. <https://doi.org/10.32614/RJ-2017-066>.
- Brooks, M.L., et al., 2004. Effects of invasive alien plants on fire regimes (Available at). *BioScience* 54 (7), 677. [https://doi.org/10.1641/0006-3568\(2004\)054\[0677:EOIAP0\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0677:EOIAP0]2.0.CO;2).
- Brown, J.K. and Smith, J.K. (2000) *Wildland fire in ecosystems: effects of fire on flora*. RMRS-GTR-42-V2. Ft. Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, p. RMRS-GTR-42-V2. Available at: <https://doi.org/10.2737/RMRS-GTR-42-V2>.
- Buytaert, W. (2022) *topmodel: Implementation of the Hydrological Model TOPMODEL in R*. Available at: <https://CRAN.R-project.org/package=topmodel>.
- Cannon, J.B., et al., 2018. Collaborative restoration effects on forest structure in ponderosa pine-dominated forests of Colorado (Available at). *For. Ecol. Manag.* 424, 191–204. <https://doi.org/10.1016/j.foreco.2018.04.026>.
- Cansler, C.A., McKenzie, D., 2014. Climate, fire size, and biophysical setting control fire severity and spatial pattern in the Northern cascade range, USA (Available at). *Ecol. Appl.* 24 (5), 1037–1056. <https://doi.org/10.1890/13-1077.1>.
- Catford, J.A., et al., 2012. Quantifying levels of biological invasion: towards the objective classification of invaded and invasional ecosystems (Available at). *Glob. Change Biol.* 18 (1), 44–62. <https://doi.org/10.1111/j.1365-2486.2011.02549.x>.
- Chamberlain, C.P., et al., 2023. Consistently heterogeneous structures observed at multiple spatial scales across fire-intact reference sites (Available at). *For. Ecol. Manag.* 550, 121478. <https://doi.org/10.1016/j.foreco.2023.121478>.
- Cook, M.C. (2021) *Drivers of wildfire-related home loss across the western United States 2001–2018*. University of Colorado. Available at: <https://www.proquest.com/openview/a27424a020ca5e959fbd3f047fcc0229/1?pq-origsite=gscholar&cbl=18750&diss=y>.
- Davis, K.T., et al., 2019. Wildfires and climate change push low-elevation forests across a critical climate threshold for tree regeneration (Available at). *Proc. Natl. Acad. Sci.* 201815107. <https://doi.org/10.1073/pnas.1815107116>.
- Davis, K.T., et al., 2023. Reduced fire severity offers near-term buffer to climate-driven declines in conifer resilience across the Western United States (Available at). *Proc. Natl. Acad. Sci.* 120 (11), e2208120120. <https://doi.org/10.1073/pnas.2208120120>.
- Davis, M.A., Grime, J.P., Thompson, K., 2000. Fluctuating resources in plant communities: a general theory of invasibility. *J. Ecol.* 88 (3), 528–534.
- Demarest, A.B., et al., 2023. Mechanical forest restoration treatments stimulate understory plants in the Colorado front range (Available at). *For. Ecol. Manag.* 548, 121322. <https://doi.org/10.1016/j.foreco.2023.121322>.
- Doughty, C.E., et al., 2021. Forest thinning in ponderosa pines increases carbon use efficiency and energy flow from primary producers to primary consumers. e2020JG005947. Available at: *J. Geophys. Res. Biogeosciences* 126 (3). <https://doi.org/10.1029/2020JG005947>.
- Dovčiak, M., Halpern, C.B., 2010. Positive diversity–stability relationships in forest herb populations during four decades of community assembly (Available at). *Ecol. Lett.* 13 (10), 1300–1309. <https://doi.org/10.1111/j.1461-0248.2010.01524.x>.
- Evans, J.S. and Murphy, M.A. (2023) *spatialEco*. Available at: <https://github.com/jeffrejevans/spatialEco>.
- Faist, A., Stone, H., Tripp, E., 2015. Impacts of mastication: soil seed bank responses to a forest thinning treatment in three Colorado (USA) conifer forest types (Available at). *Forests* 6 (9), 3060–3074. <https://doi.org/10.3390/f6093060>.
- Fialko, K., Ex, S., Wolk, B.H., 2020. Ecological niches of tree species drive variability in conifer regeneration abundance following fuels treatments (Available at). *For. Ecol. Manag.* 476, 118475. <https://doi.org/10.1016/j.foreco.2020.118475>.
- Flint, L.E., Flint, A.L., 2012. Downscaling future climate scenarios to fine scales for hydrologic and ecological modeling and analysis (Available at). *Ecol. Process.* 1 (1), 2. <https://doi.org/10.1186/2192-1709-1-2>.
- Fornwalt, P.J., et al., 2017. Mulching fuels treatments promote understory plant communities in three Colorado, USA, coniferous forest types (Available at). *For. Ecol. Manag.* 385, 214–224. <https://doi.org/10.1016/j.foreco.2016.11.047>.
- Fox, J., Weisberg, S., 2019. *An R Companion to Applied Regression*, Third. Sage, Thousand Oaks CA. (<https://www.john-fox.ca/Companion/>). (Available at).
- Fusco, E.J., et al., 2019. Invasive grasses increase fire occurrence and frequency across US ecoregions (Available at). *Proc. Natl. Acad. Sci.* 116 (47), 23594–23599. <https://doi.org/10.1073/pnas.1908253116>.
- Fusco, E.J., et al., 2021. The human–grass–fire cycle: how people and invasives co-occur to drive fire regimes (Available at). *Front. Ecol. Environ.* 2432. <https://doi.org/10.1002/fee.2432>.

- Gotelli, N.J., Colwell, R.K., 2001. Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness (Available at). *Ecol. Lett.* 4 (4), 379–391. <https://doi.org/10.1046/j.1461-0248.2001.00230.x>.
- Hartig, F. (2024) *DHARMA: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models*. Available at: <https://CRAN.R-project.org/package=DHARMA>.
- Hessburg, P.F., et al., 2021. Wildfire and climate change adaptation of Western north American forests: a case for intentional management (Available at). *Ecol. Appl.* 31 (8). <https://doi.org/10.1002/eap.2432>.
- Higuera, P.E. et al. (2023) Shifting social-ecological fire regimes explain increasing structure loss from Western wildfires. *PNAS Nexus*. Edited by J. Liu, 2(3), p. pgad005. Available at: <https://doi.org/10.1093/pnasnexus/pgad005>.
- Hoecker, T. et al. (2025) TopoTerra. OSF. Available at: <https://doi.org/10.17605/OSF.IO/W6JVK>.
- Holden, Z.A., et al., 2019. TOPOFIRE: a topographically resolved wildfire danger and drought monitoring system for the conterminous United States (Available at). *Bull. Am. Meteorol. Soc.* 100 (9), 1607–1613. <https://doi.org/10.1175/BAMS-D-18-0178.1>.
- Hollister, J. et al. (2023) *elevation: Access Elevation Data from Various APIs*. Available at: <https://doi.org/10.5281/zenodo.8335450>.
- Hood, S.M., Crotteau, J.S., Cleveland, C.C., 2024. Long-term efficacy of fuel reduction and restoration treatments in northern rocky dry forests (Available at). *Ecol. Appl.* 34 (2), e2940. <https://doi.org/10.1002/eap.2940>.
- Hurteau, M.D., et al., 2024. Managing fire-prone forests in a time of decreasing carbon carrying capacity (Available at). *Front. Ecol. Environ.* 22 (10), e2801. <https://doi.org/10.1002/fee.2801>.
- Jang, W., et al., 2021. Native and non-native understory vegetation responses to restoration treatments in a dry conifer forest over 23 years (Available at). *For. Ecol. Manag.* 481, 118684. <https://doi.org/10.1016/j.foreco.2020.118684>.
- Jeronimo, S.M.A., et al., 2019. Forest structure and pattern vary by climate and landform across active-fire landscapes in the montane Sierra Nevada (Available at). *For. Ecol. Manag.* 437, 70–86. <https://doi.org/10.1016/j.foreco.2019.01.033>.
- Kalies, E.L., Yocom Kent, L.L., 2016. Tamm review: are fuel treatments effective at achieving ecological and social objectives? A systematic review (Available at). *For. Ecol. Manag.* 375, 84–95. <https://doi.org/10.1016/j.foreco.2016.05.021>.
- Kerns, B.K., et al., 2020. Invasive grasses: a new perfect storm for forested ecosystems? (Available at). *For. Ecol. Manag.* 463, 117985. <https://doi.org/10.1016/j.foreco.2020.117985>.
- Knapp, E.E., et al., 2021. Variable thinning and prescribed fire influence tree mortality and growth during and after a severe drought (Available at). *For. Ecol. Manag.* 479, 118595. <https://doi.org/10.1016/j.foreco.2020.118595>.
- Korb, J.E., Fulé, P.Z., Gideon, B., 2007. Different restoration thinning treatments affect level of soil disturbance in ponderosa pine forests of Northern Arizona, USA (Available at). *Ecol. Restor.* 25 (1), 43–49. <https://doi.org/10.3368/er.25.1.43>.
- Kreider, M.R., et al., 2024. Fire suppression makes wildfires more severe and accentuates impacts of climate change and fuel accumulation (Available at). *Nat. Commun.* 15 (1), 2412. <https://doi.org/10.1038/s41467-024-46702-0>.
- Labelle, E.R., et al., 2022. Strategies to mitigate the effects of soil physical disturbances caused by forest machinery: a comprehensive review (Available at). *Curr. For. Rep.* 8 (1), 20–37. <https://doi.org/10.1007/s40725-021-00155-6>.
- Larson, A.J., Churchill, D., 2012. Tree spatial patterns in fire-frequent forests of Western North America, including mechanisms of pattern formation and implications for designing fuel reduction and restoration treatments (Available at). *For. Ecol. Manag.* 267, 74–92. <https://doi.org/10.1016/j.foreco.2011.11.038>.
- Laughlin, D.C., et al., 2008. Restoring plant species diversity and community composition in a ponderosa pine-bunchgrass ecosystem (Available at). *Plant Ecol.* 197 (1), 139–151. <https://doi.org/10.1007/s11258-007-9367-9>.
- Leffler, A.J., et al., 2016. Importance of soil and plant community disturbance for establishment of bromus tectorum in the intermountain west, USA (Available at). *Neobiota* 30, 111–125. <https://doi.org/10.3897/neobiota.30.7119>.
- Lenth, R.V. (2025) *emmeans: Estimated Marginal Means, aka Least-Squares Means*. Available at: <https://CRAN.R-project.org/package=emmeans>.
- Lüdtke, D., et al., 2021. Performance: an r package for assessment, comparison and testing of statistical models (Available at). *J. Open Source Softw.* 6 (60), 3139. <https://doi.org/10.21105/joss.03139>.
- Mahood, A. (2024) *topomicro: Topography and Microclimate Tools*. Available at: [www.github.com/admahood/topomicro](https://www.github.com/admahood/topomicro).
- McCauley, L.A., et al., 2019. Large-scale forest restoration stabilizes carbon under climate change in southwest United States (Available at). *Ecol. Appl.* 29 (8). <https://doi.org/10.1002/eap.1979>.
- McCauley, L.A., et al., 2022. Landscape-scale forest restoration decreases vulnerability to drought mortality under climate change in southwest USA ponderosa forest (Available at). *For. Ecol. Manag.* 509, 120088. <https://doi.org/10.1016/j.foreco.2022.120088>.
- McConnell, B.R., Smith, J.G., 1970. Response of understory vegetation to ponderosa pine thinning in eastern Washington (Available at). *J. Range Manag.* 23 (3), 208. <https://doi.org/10.2307/3896390>.
- McCune, B., Keon, D., 2002. Equations for potential annual direct incident radiation and heat load. *J. Veg. Sci.* 13 (4), 603–606.
- McGlone, C.M., Springer, J.D., Laughlin, D.C., 2009. Can pine forest restoration promote a diverse and abundant understory and simultaneously resist nonnative invasion? (Available at). *For. Ecol. Manag.* 258 (12), 2638–2646. <https://doi.org/10.1016/j.foreco.2009.09.024>.
- Metlen, K.L., Fiedler, C.E., 2006. Restoration treatment effects on the understory of ponderosa pine/Douglas-fir forests in Western Montana, USA (Available at). *For. Ecol. Manag.* 222 (1–3), 355–369. <https://doi.org/10.1016/j.foreco.2005.10.037>.
- Mietkiewicz, N., et al., 2020. In the line of fire: consequences of Human-Ignited wildfires to homes in the U.S. (1992–2015) (Available at). *Fire* 3 (3), 50. <https://doi.org/10.3390/fire3030050>.
- Miller, E.M., Seastedt, T.R., 2009. Impacts of woodchip amendments and soil nutrient availability on understory vegetation establishment following thinning of a ponderosa pine forest. *For. Ecol. Manag.* 258.
- Naficy, C., et al., 2010. Interactive effects of historical logging and fire exclusion on ponderosa pine forest structure in the Northern Rockies (Available at). *Ecol. Appl.* 20 (7), 1851–1864. <https://doi.org/10.1890/09-0217.1>.
- Nakagawa, S. and Schielzeth, H. (2013) A general and simple method for obtaining  $R^2$  from generalized linear mixed-effects models, *Methods in Ecology and Evolution*. Edited by R.B. OHara, 4(2), pp. 133–142. Available at: <https://doi.org/10.1111/j.2041-210x.2012.00261.x>.
- Naumburg, E., DeWald, L.E., 1999. Relationships between pinus ponderosa forest structure, light characteristics, and understory graminoid species presence and abundance (Available at). *For. Ecol. Manag.* 124 (2–3), 205–215. [https://doi.org/10.1016/S0378-1127\(99\)00067-5](https://doi.org/10.1016/S0378-1127(99)00067-5).
- Neal, S.M., 2007. Tree thinning treatments alter soil properties, arbuscular mycorrhizal fungi and understory plant communities. *Flagstaff AZ North. Univ.* 73.
- Nelson, C.R., Halpern, C.B., Agee, J.K., 2008. THINNING AND BURNING RESULT IN LOW-LEVEL INVASION BY NONNATIVE PLANTS BUT NEUTRAL EFFECTS ON NATIVES (Available at). *Ecol. Appl.* 18 (3), 762–770. <https://doi.org/10.1890/07-0474.1>.
- Nikooy, M., et al., 2020. Soil impacts and residual stand damage from thinning operations (Available at). *Int. J. For. Eng.* 31 (2), 126–137. <https://doi.org/10.1080/14942119.2020.1744954>.
- Oberpriller, J., De Souza Leite, M., Pichler, M., 2022. Fixed or random? On the reliability of mixed-effects models for a small number of levels in grouping variables (Available at). *Ecol. Evol.* 12 (7), e9062. <https://doi.org/10.1002/ecs3.9062>.
- Oksanen, J. et al. (2022) *vegan: Community Ecology Package*. Available at: <https://CRAN.R-project.org/package=vegan>.
- Parks, S.A., et al., 2016. How will climate change affect wildland fire severity in the Western US? (Available at). *Environ. Res. Lett.* 11 (3), 035002. <https://doi.org/10.1088/1748-9326/11/3/035002>.
- PRISM Climate Group (2024) *PRISM Gridded Climate Data*. Available at: <https://prism.oregonstate.edu>.
- Richardson, D.M., Pyšek, P., 2006. Plant invasions: merging the concepts of species invasiveness and community invasibility (Available at). *Prog. Phys. Geogr. Earth Environ.* 30 (3), 409–431. <https://doi.org/10.1191/0309133306pp490pr>.
- Riegel, G.M., Miller, R.F., Krueger, W.C., 1992. Competition for resources between understory vegetation and overstory pinus ponderosa in northeastern Oregon (Available at). *Ecol. Appl.* 2 (1), 71–85. <https://doi.org/10.2307/1941890>.
- Rodman, K.C., et al., 2024. Restoration treatments enhance tree growth and alter climatic constraints during extreme drought (Available at). *Ecol. Appl.*, e3072. <https://doi.org/10.1002/eap.3072>.
- Rother, M.T., Veblen, T.T., Furman, L.G., 2015. A field experiment informs expected patterns of conifer regeneration after disturbance under changing climate conditions (Available at). *Can. J. For. Res.* 45 (11), 1607–1616. <https://doi.org/10.1139/cjfr-2015-0033>.
- Rouget, M. et al. (2016) Invasion debt – quantifying future biological invasions, *Diversity and Distributions*. Edited by I. Kühn, 22(4), pp. 445–456. Available at: <https://doi.org/10.1111/ddi.12408>.
- Ryan, K.C., Knapp, E.E., Varner, J.M., 2013. Prescribed fire in north American forests and woodlands: history, current practice, and challenges (Available at). *Front. Ecol. Environ.* 11 (s1). <https://doi.org/10.1890/120329>.
- Schoenagel, T., Sherriff, R.L., Veblen, T.T., 2011. Fire history and tree recruitment in the Colorado front range upper montane zone: implications for forest restoration (Available at). *Ecol. Appl.* 21 (6), 2210–2222. <https://doi.org/10.1890/10-1222.1>.
- Schuerman, G. et al. (2020) *Resist-accept-direct (RAD)—a framework for the 21st-century natural resource manager*. National Park Service. Available at: <https://doi.org/10.36967/nrr-2283597>.
- Schwillk, D.W., et al., 2009. The National fire and fire surrogate study: effects of fuel reduction methods on forest vegetation structure and fuels (Available at). *Ecol. Appl.* 19 (2), 285–304. <https://doi.org/10.1890/07-1747.1>.
- Smith, E.P., Van Belle, G., 1984. Nonparametric estimation of species richness (Available at). *Biometrics* 40 (1), 119. <https://doi.org/10.2307/2530750>.
- Springer, J.D., et al., 2024. Increases in understory plant cover and richness persist following restoration treatments in pinus ponderosa forests (Available at). *J. Appl. Ecol.* 61 (1), 25–35. <https://doi.org/10.1111/1365-2664.14538>.
- Stephens, S.L., et al., 2020. Forest restoration and fuels reduction: convergent or divergent? (Available at). *BioScience* baa134. <https://doi.org/10.1093/biosci/baa134>.
- Stevens-Rumann, C.S., Morgan, P., 2019. Tree regeneration following wildfires in the Western US: a review (Available at). *Fire Ecol.* 15 (1), 15. <https://doi.org/10.1186/s42408-019-0032-1>.
- Thompson, G.G., Withers, P.C., 2003. Effect of species richness and relative abundance on the shape of the species accumulation curve (Available at). *Austral Ecol.* 28 (4), 355–360. <https://doi.org/10.1046/j.1442-9993.2003.01294.x>.
- Veblen, T.T., Kitzberger, T., Donnell, J., 2000. CLIMATIC AND HUMAN INFLUENCES ON FIRE REGIMES IN PONDEROSA PINE FORESTS IN THE COLORADO FRONT RANGE (Available at). *Ecol. Appl.* 10 (4), 1178–1195. [https://doi.org/10.1890/1051-0761\(2000\)010\[1178:CAHIOF\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[1178:CAHIOF]2.0.CO;2).
- Vernon, M.J., et al., 2023. Mechanical thinning restores ecological functions in a seasonally dry ponderosa pine forest in the inland Pacific Northwest, USA (Available at). *For. Ecol. Manag.* 546, 121371. <https://doi.org/10.1016/j.foreco.2023.121371>.

- Wasserman, T.N., et al., 2022. Natural regeneration responses to thinning and burning treatments in ponderosa pine forests and implications for restoration (Available at). *J. For. Res.* 33 (3), 741–753. <https://doi.org/10.1007/s11676-021-01404-x>.
- Willms, J., et al., 2017. The effects of thinning and burning on understory vegetation in North America: a meta-analysis (Available at). *For. Ecol. Manag.* 392, 184–194. <https://doi.org/10.1016/j.foreco.2017.03.010>.
- Wilson, J.R.U., et al., 2007. Residence time and potential range: crucial considerations in modelling plant invasions (Available at). *Divers. Distrib.* 13 (1), 11–22. <https://doi.org/10.1111/j.1366-9516.2006.00302.x>.
- Wion, A.P., et al., 2020. Aridity drives spatiotemporal patterns of masting across the latitudinal range of a dryland conifer (Available at). *Ecography* 43 (4), 569–580. <https://doi.org/10.1111/ecog.04856>.
- Wion, A.P., et al., 2023. Masting is shaped by tree-level attributes and stand structure, more than climate, in a rocky mountain conifer species (Available at). *For. Ecol. Manag.* 531, 120794. <https://doi.org/10.1016/j.foreco.2023.120794>.
- Wion, A.P., et al., 2024. Multidecadal vegetation transformations of a new Mexico ponderosa pine landscape after severe fires and aerial seeding (Available at). *Ecol. Appl.* 34 (6), e3008. <https://doi.org/10.1002/eap.3008>.
- Wion, A.P., et al., 2025. Mast hindcasts reveal pervasive effects of extreme drought on a foundational conifer species (Available at). *N. Phytol. nph.20321*. <https://doi.org/10.1111/nph.20321>.
- Wolk, B., Rocca, M.E., 2009. Thinning and chipping small-diameter ponderosa pine changes understory plant communities on the Colorado front range (Available at). *For. Ecol. Manag.* 257 (1), 85–95. <https://doi.org/10.1016/j.foreco.2008.08.014>.
- Zald, H.S.J., et al., 2022. Tree growth responses to extreme drought after mechanical thinning and prescribed fire in a Sierra Nevada mixed-conifer forest, USA (Available at). *For. Ecol. Manag.* 510, 120107. <https://doi.org/10.1016/j.foreco.2022.120107>.
- Zou, C.B., et al., 2008. Soil water dynamics under low- versus high-ponderosa pine tree density: ecohydrological functioning and restoration implications (Available at). *Ecohydrology* 1 (4), 309–315. <https://doi.org/10.1002/eco.17>.
- Zuur, A.F., et al., 2009. Mixed effects models and extensions in ecology with r. Springer New York (Statistics for Biology and Health), New York, NY. <https://doi.org/10.1007/978-0-387-87458-6>.