

Review

Integrating Habitat Suitability in Urban Forest Ecosystem Service Assessments: Reflections from i-Tree Wildlife

Susannah B. Lerman ^{1,*}, Corinne G. Bassett ², Daniel E. Crane ^{3,†}, David J. Nowak ^{3,‡}, Alexis Ellis ³ and Jason Henning ⁴

¹ US Forest Service Northern Research Station, 160 Holdsworth Way, Amherst, MA 01003, USA

² US Forest Service Northern Research Station, 5523 Research Park Drive Ste 350, Baltimore, MD 21228, USA; corinne.bassett@usda.gov

³ US Forest Service Northern Research Station, 5 Moon Library, SUNY-ESF, 1 Forestry Drive, Syracuse, NY 13210, USA; decrane64@gmail.com (D.E.C.); alexis.ellis@usda.gov (A.E.)

⁴ The Davey Institute, Philadelphia, PA 19103, USA; jason.henning@davey.com

* Correspondence: susannah.b.lerman@usda.gov

† Retired.

‡ Emeritus Senior Scientist.

Abstract

Urban forests support wildlife populations across North America and the world. Yet, challenges remain for research and practice to integrate wildlife habitat as a core component of the myriad objectives that urban foresters manage. Ecosystem services have been adopted as a dominant paradigm in urban forestry for both advocacy and management, yet accounting for contributions to wildlife habitat does not fit squarely within typical ecosystem service frameworks. The i-Tree program, a suite of urban forest ecosystem service models and tools developed by the US Forest Service, presented an opportunity to link widely used urban forest assessment field protocols with indicators of suitable habitat. In this reflection piece, we demonstrate how the i-Tree Wildlife project assessed whether urban forest structural assessment methods could be applied to assess wildlife habitat provision, operationalizing the fundamental question “How do urban forests support wildlife?” We describe the development process for integrating bird habitat suitability models for 12 species present in the northeastern US, ten native and two non-native birds, into the flagship i-Tree Eco tool. We offer reflections, challenges, and opportunities from this process. Ultimately, the improvement of ecosystem assessment tools like i-Tree can assist practitioners who aim to manage healthy and productive urban forests that benefit people and wildlife.

Keywords: bird habitat; decision support tool; ecosystem services; i-Tree; urban wildlife

Academic Editors: Le Li and
Alessandro Paletto

Received: 7 April 2026

Revised: 8 May 2026

Accepted: 15 May 2026

Published: 20 May 2026

Copyright: © 2026 by the authors.
Licensee MDPI, Basel, Switzerland.
This article is an open access article
distributed under the terms and
conditions of the [Creative Commons
Attribution \(CC BY\)](https://creativecommons.org/licenses/by/4.0/) license.

1. Introduction

Urban forests are complex social and ecological systems that provide a multitude of irreplaceable benefits to both city dwellers and populations beyond city limits [1–4]. These benefits range from processes that regulate our environment, like cleaning air and water, and providing shade, to more tangible items like food production and wood products, to the ways in which their functions support the foundations of the healthy ecosystems upon which biodiversity relies [5–7]. The science and practice of urban forestry have evolved over the last several decades in parallel with the modern concept of ecosystem services.

The concept of ecosystem services was developed to quantify and communicate the many essential benefits that natural ecosystems contribute to human populations, which can easily be overlooked in economic analyses or management objectives [8–10]. Certain ecosystem services have been much easier to develop quantification methods for, such as wood products or carbon sequestration, while others have been more difficult, such as support of wildlife habitat and biodiversity [11–13]. Indeed, many have argued that some of the most foundational benefits of nature elude quantification entirely—such as cultural and spiritual connections [14]. Nonetheless, assessing the structure, function, and service of ecosystems provides an analytical framework to simplify the complexity of ecosystems and to work towards addressing questions like “How do urban forests support wildlife?” Importantly, this framework can also serve as a link between forestry and wildlife biology, traditionally separated by strong disciplinary boundaries.

In urban forestry, during the 1990s to early 2000s, US Forest Service scientists and partners began developing models and associated tools to quantify the ecosystem services that trees provide to urban areas [15,16]. In 2006, these efforts were brought together, in partnership with industry leaders in the US, and the i-Tree program was launched [17]. The i-Tree program advances science to model the ecosystem services of trees and operationalizes this science in a suite of urban forest ecosystem service assessment tools for diverse user groups. The core models encompass an array of ecosystem services, including air pollution removal and associated health effects, carbon storage and sequestration, avoided storm water runoff, and effects on building energy use [17]. Since its inception, i-Tree’s urban forest assessment tools have been used in over 185 countries [17]. Because of i-Tree’s global reach and user base, and its foundation on a robust urban forest structure data collection protocol, it is in many ways an ideal platform to assess diverse functions and services of urban forests, including as habitats for wildlife.

Understanding how urban forests support wildlife and biodiversity more broadly is a pressing question for managers, scientists, and communities [18–20], and yet, it is often left out of ecosystem service assessment models like i-Tree because of the complexity of factors required for consideration. Urban areas support biodiversity in many critical ways—from serving as stopover sites for migrating wildlife to encompassing sensitive habitat areas [20–24]. The benefits that urban residents receive from supporting biodiversity in their cities also vary and are highly context-specific, from the benefits of connecting with nature on an individual level to all of the benefits received from living within healthy ecosystems [25–29], given the various ecosystem services many wildlife species provide (e.g., seed dispersal, insect control, and consuming carrion [30]). In addition, the wildlife recreation industry has tremendous economic benefits. The most recent State of the Birds report estimates that 37% of the US adult population identifies as a birder, and birdwatching contributes approximately \$279 billion to the US economy each year [31]. The majority of self-identified birders (70%) live in urban and suburban areas, and thus, a sizable proportion of the economic contribution likely reflects the urban wildlife recreation industry.

Because of these benefits to people, and an interest in contributing to biodiversity conservation, many practitioners and stakeholders seek to protect and promote wildlife and their habitats in urban forests [32–34]. However, strategies and tools can vary, and research methods require extensive expertise and are difficult to share among geographies [35,36]. Overcoming these disciplinary divisions is key to ensuring that biodiversity and wildlife are included in advancing the field of urban forestry [37]. Connecting urban forest structural variables with their subsequent impact on wildlife, even when highly simplified, promises a chance to provide practitioners with actionable recommendations. In this review, we present lessons learned from this endeavor to use i-Tree’s urban forest assessment methods to understand the contributions of urban forests to wildlife habitat and diversity (Figure 1). The initial wildlife habitat models focused on birds because of their

role as bioindicators, coupled with the wealth of information available about their natural history [38] and available i-Tree data from the northeastern US. In doing so, we share reflections from the theoretical to the practical on connecting methods and datasets across disciplines and integration with an applied, operational program like i-Tree.




Figure 1. Theory of change for how i-Tree Wildlife addressed the fundamental research question “How do urban forests support wildlife?” (Credit: flaticon.com).

2. Background of Wildlife and the Urban Forest

The seminal 1973 US Forest Service publication “Invite wildlife to your backyard” [39] marked a shift from managing game species in wildlands to recognizing the importance of managing nongame species in suburban and urban ecosystems. The publication also inspired a new field of wildlife and biodiversity research. From the mid-1970s to the 2000s, urban wildlife and biodiversity research focused on distribution patterns along urban gradients, using distance to the urban center or the amount of impervious surface as urban metrics, often from a single region and predominantly conducted in forest patches and public spaces [40–42]. This approach primarily engaged traditional ecological methodology and was largely observational to assess the ecological structure of habitats and communities and the function of these systems. Studies from different ecoregions in North America, Europe, and Australia concluded that in more built-up and urban areas, bird diversity decreased, yet density (i.e., number of individuals) increased, while in rural areas, the opposite patterns occurred [43,44]. These results supported the emphasis on urbanization as a negative disturbance and urban landscapes as non-habitat areas. However, as indicated in the 1973 USDA Forest Service publication, elements within the urban matrix, ranging from individual trees to small residential yards, in addition to large intact forest patches, have the potential to support wildlife and biodiversity [45,46].

By the mid-2000s, urban wildlife research broadened to examine how urban wildlife and biodiversity enhance quality of life and other ecosystem services for urban and suburban (human) residents, and in turn, the integral role humans play in shaping urban ecosystems [5]. The two urban National Science Foundation-funded Long Term Ecological Research (LTER) sites (Phoenix and Baltimore) spear-headed this shift in urban ecological research (e.g., [47,48]). In light of the potential of greenspaces broadly and biodiversity specifically to support ecosystem services for the millions of residents who live in

urban and suburban areas, some urban wildlife research shifted towards identifying how to better manage urban greenspaces as wildlife habitats [49–52]. Within any type of greenspace, there is considerable variation in management decision criteria (e.g., tree removal, native plantings [53–55]) and management goals (i.e., regarding aesthetics, ease of maintenance, and safety [56]) that can have far-reaching implications on habitat quality for wildlife. For example, modifying tree-pruning schedules to benefit cavity-nesting birds and leaving dead trees for habitat, when they pose minimal or no risk, can support wildlife populations [53]. These various management practices result in heterogeneous patches of urban greenspace that support a range of both generalist and specialist native species, as well as non-native and invasive species [45,57–60].

Urban forests and urban greenspaces consist of a diversity of plant communities, and their configuration can vary from an individual tree to forest patches. Plants include ornamental, native, non-native, and invasive species [61], with birds responding to these different plants in unique ways. For example, research from the Washington DC metropolitan area demonstrated that Carolina chickadees (*Poecile carolinensis*) spent more time foraging in neighborhoods with at least 70% native plant biomass because of the strong relationship between native plants and their preferred insect prey items [54,62]. Denser and larger street trees in Los Angeles attracted a higher percentage of foraging birds, with strong preferences for both native and non-native species [63]. When urban forests structurally resemble the wildlands they replace (i.e., similar heights and abundance of different types of vegetation), they can support native bird communities [64,65]. And finally, different land-use types throughout the urban matrix (e.g., residential yards, urban parks, and street trees) support distinct species assemblages [60,66,67], though their importance might shift across seasons [68]. These and other studies have documented how birds and other wildlife species respond to the density and structure of the urban forest and provide support for how the urban forest provides wildlife habitat.

The vast majority of urban bird studies have been limited in scope, either focusing on a single species (e.g., [54]), or within a single city (e.g., [59,69], but see [70–72]). Studies conducted at continental scales that investigate multiple taxa are necessary to address the generalizability of biodiversity patterns [73] and to build management recommendations. Empirical studies allow for replication across ecosystems (e.g., deserts and forests) and consider how several factors, such as the amount of impervious surface, tree canopy cover, and management practices, shape bird communities [74]. However, the amount of coordination needed for continental or even global studies is frequently prohibitive. Thus, developing predictive models that can assess existing bird habitat quality for multiple species while identifying opportunities for improving habitat conditions represents an important contribution to habitat conservation. Given the strong relationships between birds and the urban forest, particularly the features captured in the i-Tree Eco plot data collection protocol, it is possible to use urban forest structural datasets to calculate the overall bird habitat suitability for a city or a delineated urban forest assessment project, and the habitat suitability for different types of land-use (e.g., residential, commercial, and parkland) for each bird species [38]. Bird habitat suitability indices based on local data inputs could enable researchers and managers to create and quantify targeted assessments and detailed predictions, and guide land management decisions that support bird habitats. They could also illustrate trade-offs among wildlife species as urban forest structure changes.

3. Developing Models and Operationalizing the Habitat Suitability Index

In 2014, Lerman and colleagues [38] developed habitat suitability index (HSI) models for nine species and integrated them into the publicly accessible urban forest assessment tool, i-Tree Eco (Figure 2). In 2025, the i-Tree Wildlife module expanded to include three additional HSI models for a total of 12 species observed throughout the northeastern and midwestern United States (see Supplementary Materials for three additional species accounts). We selected these species based on specific criteria so they can serve as representatives for other species with similar life history traits or conservation status. We categorized birds into different foraging and nesting guilds, focusing on guilds reliant on forests [41]. The species list includes cavity and open cup nesters, species nesting high in the forest canopy, and those reliant on shrubs and early successional habitat, including species foraging throughout different canopy heights, on bark, and on the ground. Some species specialize in insects, while others have a more generalized diet. We included species with a Partners in Flight (PIF) conservation designation, which ranks a species' conservation vulnerability based on global population, population trends, and threats to breeding populations. Species with a high PIF score have targeted conservation initiatives [75]. We also included two non-native species: house sparrow (*Passer domesticus*) and European starling (*Sturnus vulgaris*). Although all species have unique and sometimes nuanced habitat requirements, we can generalize some aspects and use the habitat suitability indices for some of our 12 species to assess habitat suitability for other forest-dependent species. For example, the red-bellied woodpecker (*Melanerpes carolinus*) requires some deadwood for excavating nests, similar to other woodpeckers like the downy woodpecker (*Picoides pubescens*) and hairy woodpecker (*Leuconotopicus villosus*), and thus, parts of the red-bellied woodpecker HSI could be extended to these other cavity-nesting species. By including two non-native species that might outcompete native species [57], the model outputs could also provide guidance on how to deter some species based on their habitat requirements. For more details about the model building, validation, and uncertainty quantification, please refer to [38].




Figure 2. Approach to developing the habitat suitability index [38] and operationalization for practitioner users in the i-Tree Eco tool.

To date, the original nine habitat suitability models have been incorporated into the i-Tree Eco tool. Although the geographic scope for building the models was limited to the northeastern US, the selected species have a broad geographic range, and thus, the models can be and have been applied in different regions (e.g., Michigan; Table 1) when i-Tree urban forest data exist. The modeling approach follows an established protocol developed by Tirpak and colleagues from the midwestern US [76–78], where they integrated natural history information with the US Forest Service Forest Inventory & Analysis program’s Nation-wide Forest Inventory. Since the publication of the i-Tree bird habitat models, other urban bird studies have followed similar habitat assessments (e.g., [54]). In Santo Domingo, Dominican Republic, research linked city-wide bird surveys with i-Tree plots to expand the geographic scope of the tool (Arendt et al., in review). These and other examples demonstrate the utility and feasibility of replicating the modeling process in other regions using the same i-Tree urban forest assessment metrics. Integrating additional bird species, as well as other taxa (e.g., insect pollinators and mammals), has potential, given their strong associations with urban forests [45,79], coupled with existing natural history information.

Table 1. “How well does my urban forest support birds?” Example output report for an i-Tree project in Grand Rapids, Michigan, USA, from 2011. The suitability index (SI) ranges between 0 and 1, whereby a score of 0 indicates an i-Tree plot has unsuitable habitat conditions with or without trees, i.e., a strong likelihood the species is not present, whereas a score of 1 indicates the habitat conditions have a strong likelihood of supporting the species. The Index Change Due to Trees calculates the relative change (absolute value divided by SI with trees) and the absolute change when a plot includes trees.

Species	Suitability Index		Index Change Due to Trees	
	With Trees	Without Trees	Relative (%)	Absolute
American Robin	0.552	0.549	0.558	0.003
Baltimore Oriole	0.288	0.052	82.122	0.237
Black-capped Chickadee	0.223	0.062	72.051	0.161
European Starling	0.271	0.268	1.085	0.003
Northern Cardinal	0.272	0.289	−6.384	−0.017
Red-bellied Woodpecker	0.239	0.02	91.694	0.22
Scarlet Tanager	0.012	0	97.005	0.011
Wood Thrush	0.074	0	99.547	0.074

The inclusion of assessing wildlife habitat suitability in a practitioner-level tool such as i-Tree Eco demonstrated the feasibility of linking forest management tools with bird conservation planning. i-Tree Eco was designed as a quantitative tool, allowing users to conduct analyses and estimate multiple benefits using data on their own urban forest. Adding bird suitability data to this urban forest assessment tool could enable users to develop quantitative urban forest management targets, rendering plans that prioritize biodiversity more actionable [18]. The inclusion of a wildlife module in i-Tree Eco also expands the scope of the tool by assessing another output or beneficiary of the urban forest, in this case, birds. Having detailed information about bird species supported alongside ecosystem service estimates, such as pollutants removed, carbon dioxide absorbed, and stormwater intercepted, can provide science-based information for enhancing urban biodiversity and addressing other urban greening goals [23].

To fully operationalize the habitat suitability model outputs, framing the models within the context of how urban foresters operate is essential. For urban foresters, outputs can answer the question “How well does my urban forest support birds?” (Table 1). To answer this question, the i-Tree models calculate a suitability index that includes tree variables for a particular species, and the suitability index that excludes tree variables for that

species (i.e., “with” and “without” trees), to capture the contributions of trees, regardless of their structure or composition, to the habitat for a particular bird species within a city. Based on these indices, the assessment calculates the benefit of trees for each species at each plot/stratum by calculating a relative change (%) and absolute index change due to trees being present. For example, trees in Grand Rapids, Michigan, US, improve Baltimore oriole (*Icterus galbula*) habitat by 82% compared with locations without trees (Table 1). Though this approach aligns with other modeling efforts in the i-Tree tool (e.g., calculating the tree ecosystem services related to stormwater runoff), this framing is not ideal for decision-making when the interest focuses on *how* to improve habitat.

A question closer to the decision-making context asks, “What are specific urban forest structure goals for supporting birds?” (Table 2). To address this question, linking suitability indices of different land-use strata with optimal suitability, practitioners can set targets in management plans, such as how much to increase tree canopy cover or how many trees with deadwood to conserve (Table 2). For example, based on the HSI for the red-bellied woodpecker, increasing canopy cover from 20% to 35% increases the suitability of the urban forest threefold (from 0.3 to 0.9; Table 2), and provides guidance for other species with similar tree canopy requirements [38]. At its core, i-Tree Eco is a science delivery tool, and thus, having intentionality on how results are shared can increase the likelihood of tool use. Although Table 2 addresses this issue, it is not included in the current i-Tree Eco reports. Adding these types of results would increase the tool’s utility in creating actionable information from the HSI models and forest inventory data. As a caveat, the habitat suitability models demonstrate the potential habitat conditions for supporting a particular species by providing a threshold, and not a guarantee of occurrence. The habitat variables might not represent the most informative variable for suitability, yet they still contribute to habitat selection. Plus, the i-tree variables can be manipulated by urban foresters. Thus, the HSI models are best used for guidance and conservation planning.

Table 2. “What are specific urban forest structure goals for supporting birds?” An example of the suitability indices for the red-bellied woodpecker, at the city-level, from two different cities (Philadelphia, PA, USA, and New York, NY, USA) and within two different strata (i.e., residential and park land-uses) for each habitat variable. The shaded row represents the optimal values for each variable, i.e., when suitability is 1. Asterisks indicate the specific habitat conditions for a variable that falls within optimal levels. Variables that fall below optimal values, e.g., canopy % for residential lands in both cities, provide a starting point for developing management goals. For example, targeting an increase of ~25% canopy cover in residential areas in Philadelphia is one part of improving habitat suitability for the species.

Red-Bellied Woodpecker	Suitability Index	Canopy % ¹ (0.04 ha)	Large Tree Density ² (0.04 ha)	Basal Area ³ (m ² /ha)	Deadwood Abundance ⁴ (0.04 ha)
Study Area: Philadelphia	0.39	17.6	2.41	12.62 *	4.75
Residential land-use	0.20	12.5	0.68	1.91	1.26
Park land-use	0.70	75.5 *	5.11	10.91 *	9.06
Optimal		35%–62%	>6	8–14	1–3
Study Area: New York	0.17	23.2	1.03	5.51	0.49
Residential land-use	0.15	19.5	0.63	1.47	0.3
Park land-use	0.30	36.0 *	2.12	4.57	0.85

¹ Canopy % is the percentage of the i-Tree plot covered by tree canopy. ² Large tree is the total number of trees >23 cm DBH within the i-tree plot. ³ Basal area calculates the basal area of all trees >6 cm DBH. ⁴ Deadwood is the total number of trees within the i-Tree plot classified as “Fair”, “Poor”, “Dying”, or “Dead”.

4. Challenges and Opportunities

Integrating wildlife habitat models into i-Tree Eco presented an opportunity to use a platform familiar to urban foresters who make management decisions that have implications for wildlife habitat. However, even after developing the models, challenges exist, e.g., balancing the complexity of the models with the simplicity of reporting, and how to expand the number of species included and the geographical scope.

Building the wildlife habitat module extended the reporting capacity of i-Tree Eco for US users. However, there were several limitations to the model-building process. First, developing a habitat suitability index for each species is time-consuming, taking approximately 80 h per species. The process included an extensive literature search to identify studies that quantified habitat relationships with a specific metric, e.g., nest success. Fortunately, birds are very well-studied, and extensive information exists. However, this also represented a challenge because of the excessive amount of information to search. Furthermore, the literature often focuses on a single species, with habitat relationships for a particular species (e.g., [80]). Although this provides detailed information for a particular species, it translates to building one habitat suitability index at a time. Once a study has been identified as having habitat information linked to a metric of success, the specific habitat features and the scale in which they were collected might not match the urban forest structural variables or plot sizes (in our case, an 11.2 m radius i-Tree Eco plot); e.g., see studies on the wood thrush (*Hylocichla mustelina*) home range and roosting areas [80,81]. Habitat measurements include vegetation density and forest structure using LiDAR to address the specific hypothesis, but do not easily translate to field-collected structural variables. This challenge is compounded by the need to identify multiple studies per habitat variable to account for variation, and the need to replicate the process for multiple habitat variables.

Augmenting the scarcity of published datasets with estimated values can fill these data challenges. For example, when building the HSIs for i-Tree, iterative values were supplemented, which improved the predictability of the habitat models. These iterative values were then reviewed by a panel of experts, which allowed for the models to be revised according to recommendations [38]. Soliciting experts can also address the labor-intensive process, which is difficult to scale. For example, in Australia, the Biodiversity Sensitive Urban Design (BSUD) framework solicited expert knowledge to assist with urban planning. Experts helped estimate habitat and connectivity requirements for woodland birds. The baseline data was integrated into urban planning actions, ultimately benefiting biodiversity [82].

Once the studies had been identified, variables selected, and the habitat suitability models developed, the models then required validation [38]. Validating the habitat suitability models required datasets that included bird survey data co-located with i-Tree plots. At the time, to our knowledge, these data only existed in Baltimore, MD, as part of the Baltimore Ecosystem Study Long-Term Ecological Research (BES LTER) program [38,83,84]. Harnessing support from birders [31] to conduct additional surveys to help build and validate models holds potential. However, coordinating data generated by thousands of citizen scientists requires considerable resources and expertise. Although robust models depend upon the underlying data used for building habitat relationships, the unvalidated models still have value [85], and can serve as a starting point for understanding how birds might respond to different urban forest management practices.

Geographical challenges also existed with the model-building process. The i-Tree user (e.g., urban forester, urban planner, and the public) is focused on local information about a specific urban forest or urban trees. Because of the challenges of matching the ornithological literature with urban forest structural variables, the studies available for building the habitat suitability models might have a geographical mismatch. For example, the red-bellied

woodpecker habitat suitability model relied on studies from Ontario, the Midwest, and southwestern Virginia [86–88], though they were applied to ten northeastern US cities [38]. Although it is likely that the urban forest composition and configuration amongst these different regions have some similarities [89], it is unclear how consistent the habitat relationships are across a species' breeding range. Ornithologists do not have a standardized way to collect habitat data, and oftentimes the variables are particular to a specific species or research question [90,91]. However, establishing a set of consistent variables to collect, e.g., using variables from established urban forest monitoring protocols for habitat assessments and habitat relationships, can further address these challenges [92,93].

5. Conclusions

The i-Tree Wildlife module demonstrated how to model habitat suitability based on urban forest structure. Additional opportunities exist for further integration of other wildlife data with urban forest datasets and tools. This integration can increase our understanding of habitat suitability relationships alongside tree and urban forest assessments. For example, integrating landscape-scale habitat variables, e.g., the US Forest Service Tree Canopy Cover datasets [94] or the U.S. Geological Survey's National Land Cover Database [95] with citizen science wildlife monitoring programs such as eBird (a Cornell Lab of Ornithology citizen science repository for birders' species list [96]) and iNaturalist, has the potential to uncover coarse relationships between available habitat and bird presence at a continental scale [97,98]. The InVEST (Integrated Valuation of Ecosystem Services and Trade-offs) tool also assesses the ecosystem services in urban systems. Integrating land-cover data with pollinator habitat studies, InVEST developed a pollination model to identify areas that would benefit from habitat improvement efforts. However, their models are based on spatial greenspace availability data, not forest structure and composition. The US Forest Service's Urban Forest Inventory and Analysis (FIA) program and other urban forest inventories with structural variables have strong potential for informing additional HSIIs [61]. Finally, i-Tree's other ecosystem service models have been integrated into other public tools, e.g., the American Forests' Tree Equity Score (www.treeequityscore.org), as well as industry tools by private companies in the US and abroad.

Future research that builds from the understanding of bird habitat and urban forest relationships can further address how to improve decision support tools [99]. Addressing the challenges of geography, labor intensiveness of model building, capacity issues, input data, and how to generate results that users can operationalize are key factors to consider. Participatory evaluation of future user interfaces and reporting outputs by both urban foresters and wildlife managers could also aid in the effective use of the results in practice [100]. Here, we shared generalizable findings and key takeaways from the i-Tree Wildlife endeavor. Ultimately, the improvement of tools, e.g., i-Tree, can assist practitioners who aim to manage healthy and productive urban forests that benefit people and wildlife.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/f17050620/s1>.

Author Contributions: Conceptualization, S.B.L. and C.G.B.; writing—original draft preparation, S.B.L. and C.G.B.; writing—review and editing, D.E.C., D.J.N., A.E., and J.H. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: No new data were created for this article.

Acknowledgments: We thank Mark Majewsky for his guidance during the development of the additional i-Tree habitat suitability models. The findings and conclusions in this publication are those of the authors and should not be construed to represent any official USDA or U.S. Government determination or policy.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Vogt, J. A Comprehensive Framework for Understanding Urban Forests as Social-Ecological Systems. *Arboric. Urban For. (AUF)* **2024**, *50*, 427–469. <https://doi.org/10.48044/jauf.2024.021>.
2. Tyrväinen, L.; Pauleit, S.; Seeland, K.; De Vries, S. Benefits and Uses of Urban Forests and Trees. In *Urban Forests and Trees*; Konijnendijk, C., Nilsson, K., Randrup, T., Schipperijn, J., Eds.; Springer Berlin Heidelberg: Berlin/Heidelberg, Germany, 2005; pp. 81–114, ISBN 978-3-540-25126-2.
3. O'Brien, L.E.; Urbanek, R.E.; Gregory, J.D. Ecological Functions and Human Benefits of Urban Forests. *Urban For. Urban Green.* **2022**, *75*, 127707.
4. Dwyer, J.F.; McPherson, E.G.; Schroeder, H.W.; Rowntree, R.A. Assessing the Benefits and Costs of the Urban Forest. *J. Arboric.* **1992**, *18*, 227–234.
5. Grimm, N.B.; Faeth, S.H.; Golubiewski, N.E.; Redman, C.L.; Wu, J.; Bai, X.; Briggs, J.M. Global Change and the Ecology of Cities. *Science* **2008**, *319*, 756–760. <https://doi.org/10.1126/science.1150195>.
6. Ziter, C. The Biodiversity–Ecosystem Service Relationship in Urban Areas: A Quantitative Review. *Oikos* **2016**, *125*, 761–768. <https://doi.org/10.1111/oik.02883>.
7. Alvey, A.A. Promoting and Preserving Biodiversity in the Urban Forest. *Urban For. Urban Green.* **2006**, *5*, 195–201.
8. Daily, G.C. *Nature's Services: Societal Dependence on Natural Ecosystems.*; Island Press: Washington, DC, USA, 1997; ISBN 978-1-55963-476-2.
9. Leemans, R.; de Groot, R.S. *Millennium Ecosystem Assessment: Ecosystems and Human Well-Being: A Framework for Assessment*; Island Press: Washington, DC, USA, 2003; ISBN 978-1-55963-402-1.
10. Bolund, P.; Hunhammar, S. Ecosystem Services in Urban Areas. *Ecol. Econ.* **1999**, *29*, 293–301. [https://doi.org/10.1016/S0921-8009\(99\)00013-0](https://doi.org/10.1016/S0921-8009(99)00013-0).
11. Mace, G.M.; Norris, K.; Fitter, A.H. Biodiversity and Ecosystem Services: A Multilayered Relationship. *Trends Ecol. Evol.* **2012**, *27*, 19–26. <https://doi.org/10.1016/j.tree.2011.08.006>.
12. Boerema, A.; Rebelo, A.J.; Bodi, M.B.; Esler, K.J.; Meire, P. Are Ecosystem Services Adequately Quantified? *J. Appl. Ecol.* **2017**, *54*, 358–370. <https://doi.org/10.1111/1365-2664.12696>.
13. Endreny, T.; Santagata, R.; Perna, A.; Stefano, C.D.; Rallo, R.F.; Ulgiati, S. Implementing and Managing Urban Forests: A Much Needed Conservation Strategy to Increase Ecosystem Services and Urban Wellbeing. *Ecol. Model.* **2017**, *360*, 328–335. <https://doi.org/10.1016/j.ecolmodel.2017.07.016>.
14. Chan, K.M.A.; Guerry, A.D.; Balvanera, P.; Klain, S.; Satterfield, T.; Basurto, X.; Bostrom, A.; Chuenpagdee, R.; Gould, R.; Halpern, B.S.; et al. Where Are Cultural and Social in Ecosystem Services? A Framework for Constructive Engagement. *BioScience* **2012**, *62*, 744–756. <https://doi.org/10.1525/bio.2012.62.8.7>.
15. Nowak, D.J. Atmospheric Carbon Reduction by Urban Trees. *J. Environ. Manag.* **1993**, *37*, 207–217. <https://doi.org/10.1006/jema.1993.1017>.
16. Nowak, D.J.; Crane, D.E. The Urban Forest Effects (UFORE) Model: Quantifying Urban Forest Structure and Functions. In *Integrated Tools for Natural Resources Inventories in the 21st Century*; Hansen, M., Burk, T., Eds.; Gen. Tech. Rep. NC-212; Department of Agriculture, Forest Service, North Central Forest Experiment Station: St. Paul, MN, USA, 2000; pp. 714–720.
17. Nowak, D.J. *Understanding I-Tree: 2023 Summary of Programs and Methods*; General Technical Report NRS-200-2023; Department of Agriculture, Forest Service, Northern Research Station: Madison, WI, USA, 2024; 103p.
18. Nilon, C.H.; Aronson, M.F.J.; Cilliers, S.S.; Dobbs, C.; Frazee, L.J.; Goddard, M.A.; O'Neill, K.M.; Roberts, D.; Stander, E.K.; Werner, P.; et al. Planning for the Future of Urban Biodiversity: A Global Review of City-Scale Initiatives. *BioScience* **2017**, *67*, 332–342.
19. Lambert, M.R.; Donihue, C.M. Urban Biodiversity Management Using Evolutionary Tools. *Nat. Ecol. Evol.* **2020**, *4*, 903–910. <https://doi.org/10.1038/s41559-020-1193-7>.
20. Kowarik, I. Novel Urban Ecosystems, Biodiversity, and Conservation. *Environ. Pollut.* **2011**, *159*, 1974–1983. <https://doi.org/10.1016/j.envpol.2011.02.022>.

21. Aronson, M.F.; Lepczyk, C.A.; Evans, K.L.; Goddard, M.A.; Lerman, S.B.; MacIvor, J.S.; Nilon, C.H.; Vargo, T. Biodiversity in the City: Key Challenges for Urban Green Space Management. *Front. Ecol. Environ.* **2017**, *15*, 189–196. <https://doi.org/10.1002/fee.1480>.
22. Lerman, S.B.; Larson, K.L.; Narango, D.L.; Goddard, M.A.; Marra, P.P. Humanity for Habitat: Residential Yards as an Opportunity for Biodiversity Conservation. *BioScience* **2023**, *73*, 671–689.
23. Spotswood, E.N.; Beller, E.E.; Grossinger, R.; Grenier, J.L.; Heller, N.E.; Aronson, M.F.J. The Biological Deserts Fallacy: Cities in Their Landscapes Contribute More than We Think to Regional Biodiversity. *BioScience* **2021**, *71*, 148–160. <https://doi.org/10.1093/biosci/biaa155>.
24. Beninde, J.; Veith, M.; Hochkirch, A. Biodiversity in Cities Needs Space: A Meta-Analysis of Factors Determining Intra-Urban Biodiversity Variation. *Ecol. Lett.* **2015**, *18*, 581–592. <https://doi.org/10.1111/ele.12427>.
25. Carrus, G.; Scopelliti, M.; Laforteza, R.; Colangelo, G.; Ferrini, F.; Salbitano, F.; Agrimi, M.; Portoghesi, L.; Semenzato, P.; Sanesi, G. Go Greener, Feel Better? The Positive Effects of Biodiversity on the Well-Being of Individuals Visiting Urban and Peri-Urban Green Areas. *Landsc. Urban Plan.* **2015**, *134*, 221–228. <https://doi.org/10.1016/j.landurbplan.2014.10.022>.
26. Taylor, L.; Hochuli, D.F. Creating Better Cities: How Biodiversity and Ecosystem Functioning Enhance Urban Residents' Well-being. *Urban Ecosyst.* **2015**, *18*, 747–762.
27. Marselle, M.R.; Hartig, T.; Cox, D.T.C.; de Bell, S.; Knapp, S.; Lindley, S.; Triguero-Mas, M.; Böhning-Gaese, K.; Braubach, M.; Cook, P.A.; et al. Pathways Linking Biodiversity to Human Health: A Conceptual Framework. *Environ. Int.* **2021**, *150*, 106420. <https://doi.org/10.1016/j.envint.2021.106420>.
28. Cox, D.T.C.; Shanahan, D.F.; Hudson, H.L.; Plummer, K.E.; Siriwardena, G.M.; Fuller, R.A.; Anderson, K.; Hancock, S.; Gaston, K.J. Doses of Neighborhood Nature: The Benefits for Mental Health of Living with Nature. *Bioscience* **2017**, *67*, 147–155. <https://doi.org/10.1093/biosci/biw173>.
29. Soga, M.; Gaston, K.J. The Ecology of Human–Nature Interactions. *Proc. R. Soc. B.* **2020**, *287*, 20191882. <https://doi.org/10.1098/rspb.2019.1882>.
30. Whelan, C.J.; Wenny, D.G.; Marquis, R.J. Ecosystem Services Provided by Birds. *Ann. N. Y. Acad. Sci.* **2008**, *1134*, 25–60.
31. U.S. Department of the Interior; U.S. Fish and Wildlife Service. *Birding in the United States—A Demographic and Economic Analysis, Addendum to the 2022 National Survey of Fishing, Hunting, and Wildlife-Associated Recreation*; U.S. Department of the Interior: Washington, DC, USA, 2024.
32. Martin, A.J.F.; Almas, A. Arborists and Urban Foresters Support for Urban Wildlife and Habitat Sustainability: Results of an Urban Ecology-Focused Survey of Arborists. *Sustainability* **2022**, *14*, 15962. <https://doi.org/10.3390/su142315962>.
33. Jimenez, M.F.; Pejchar, L.; Reed, S.E.; McHale, M.R. The Efficacy of Urban Habitat Enhancement Programs for Conserving Native Plants and Human-Sensitive Animals. *Landsc. Urban Plan.* **2022**, *220*, 104356. <https://doi.org/10.1016/j.landurbplan.2022.104356>.
34. Dunster, J.A. The Role of Arborists in Providing Wildlife Habitat and Landscape Linkages throughout the Urban Forest. *J. Arboric.* **1998**, *24*, 160–167.
35. Bassett, C.G.; Gilpin, R.; Donohue, K. Lessons Learned from Developing Best Management Practices for Urban Tree Care and Wildlife. *Arboric. Urban For. (AUF)* **2022**, *48*, 1–8. <https://doi.org/10.48044/jauf.2022.001>.
36. Hutt-Taylor, K.; Bassett, C.G.; Kinnunen, R.P.; Frei, B.; Ziter, C.D. Existing Evidence on the Effect of Urban Forest Management in Carbon Solutions and Avian Conservation: A Systematic Literature Map. *Environ. Évid.* **2024**, *13*, 23. <https://doi.org/10.1186/s13750-024-00344-3>.
37. O'Herrin, K.; Bassett, C.G.; Day, S.D.; Ries, P.D.; Wiseman, P.E. Borrowed Credentials and Surrogate Professional Societies: A Critical Analysis of the Urban Forestry Profession. *Arboric. Urban For. (AUF)* **2023**, *49*, 107–136. <https://doi.org/10.48044/jauf.2023.009>.
38. Lerman, S.B.; Nislow, K.H.; Nowak, D.J.; DeStefano, S.; King, D.I.; Jones-Farrand, D.T. Using Urban Forest Assessment Tools to Model Bird Habitat Potential. *Landsc. Urban Plan.* **2014**, *122*, 29–40. <https://doi.org/10.1016/j.landurbplan.2013.10.006>.
39. Thomas, J.W.; Brush, R.O.; DeGraaf, R.M. Invite Wildlife to Your Backyard. *Natl. Wildl.* **1973**, *11*, 5–16.
40. Chace, J.F.; Walsh, J.J. Urban Effects on Native Avifauna: A Review. *Landsc. Urban Plan.* **2006**, *74*, 46–69.
41. Degraaf, R.M.; Wentworth, J.M. Avian Guild Structure and Habitat Associations in Suburban Bird Communities. *Urban Ecol.* **1986**, *9*, 399–412.
42. Emlen, J.T. Urban Bird Community in Tucson, Arizona—Derivation, Structure, Regulation. *Condor* **1974**, *76*, 184–197.
43. Blair, R.B. Land Use and Avian Species Diversity along an Urban Gradient. *Ecol. Appl.* **1996**, *6*, 506–519. <https://doi.org/10.2307/2269387>.
44. McKinney, M.L. Urbanization as a Major Cause of Biotic Homogenization. *Biol. Conserv.* **2006**, *127*, 247–260.

45. Lepczyk, C.A.; Aronson, M.F.J.; Evans, K.L.; Goddard, M.A.; Lerman, S.B.; MacIvor, J.S. Biodiversity in the City: Fundamental Questions for Understanding the Ecology of Urban Green Spaces for Biodiversity Conservation. *BioScience* **2017**, *67*, 799–807. <https://doi.org/10.1093/biosci/bix079>.
46. Vasquez, A.V.; Wood, E.M. Urban Parks Are a Refuge for Birds in Park-Poor Areas. *Front. Ecol. Evol.* **2022**, *10*, 1048.
47. Hope, D.; Gries, C.; Zhu, W.X.; Fagan, W.F.; Redman, C.L.; Grimm, N.B.; Nelson, A.L.; Martin, C.; Kinzig, A. Socioeconomics Drive Urban Plant Diversity. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 8788–8792.
48. Kinzig, A.P.; Warren, P.; Martin, C.; Hope, D.; Katti, M. The Effects of Human Socioeconomic Status and Cultural Characteristics on Urban Patterns of Biodiversity. *Ecol. Soc.* **2005**, *10*, 23.
49. Belaire, J.A.; Whelan, C.J.; Minor, E.S. Having Our Yards and Sharing Them Too: The Collective Effects of Yards on Native Bird Species in an Urban Landscape. *Ecol. Appl.* **2014**, *24*, 2132–2143. <https://doi.org/10.1890/13-2259.1>.
50. Goddard, M.A.; Dougill, A.J.; Benton, T.G. Scaling up from Gardens: Biodiversity Conservation in Urban Environments. *Trends Ecol. Evol.* **2010**, *25*, 90–98. <https://doi.org/10.1016/j.tree.2009.07.016>.
51. van Heezik, Y.M.; Dickinson, K.J.M.; Freeman, C. Closing the Gap: Communicating to Change Gardening Practices in Support of Native Biodiversity in Urban Private Gardens. *Ecol. Soc.* **2012**, *17*, 34. <https://doi.org/10.5751/ES-04712-170134>.
52. Apfelbeck, B.; Snep, R.P.H.; Hauck, T.E.; Ferguson, J.; Holy, M.; Jakoby, C.; Scott MacIvor, J.; Schär, L.; Taylor, M.; Weisser, W.W. Designing Wildlife-Inclusive Cities That Support Human-Animal Co-Existence. *Landsc. Urban Plan.* **2020**, *200*, 103817. <https://doi.org/10.1016/j.landurbplan.2020.103817>.
53. Kane, B.; Warren, P.S.; Lerman, S.B. A Broad Scale Analysis of Tree Risk, Mitigation and Potential Habitat for Cavity-Nesting Birds. *Urban For. Urban Green.* **2015**, *14*, 1137–1146. <https://doi.org/10.1016/j.ufug.2015.10.012>.
54. Narango, D.L.; Tallamy, D.W.; Marra, P.P. Native Plants Improve Breeding and Foraging Habitat for an Insectivorous Bird. *Biol. Conserv.* **2017**, *213*, 42–50. <https://doi.org/10.1016/j.biocon.2017.06.029>.
55. Faeth, S.H.; Bang, C.; Saari, S. Urban Biodiversity: Patterns and Mechanisms. In *Year in Ecology and Conservation Biology*; Ostfeld, R., Schlesinger, W., Eds.; Blackwell Science Publ: Oxford, UK, 2011; Volume 1223, pp. 69–81, ISBN 978-1-57331-833-4.
56. Larson, K.L.; Nelson, K.C.; Samples, S.R.; Hall, S.J.; Bettez, N.; Cavender-Bares, J.; Groffman, P.M.; Grove, M.; Heffernan, J.B.; Hobbie, S.E.; et al. Ecosystem Services in Managing Residential Landscapes: Priorities, Value Dimensions, and Cross-Regional Patterns. *Urban Ecosyst.* **2016**, *19*, 95–113. <https://doi.org/10.1007/s11252-015-0477-1>.
57. Shochat, E.; Lerman, S.B.; Anderies, J.M.; Warren, P.S.; Faeth, S.H.; Nilon, C.H. Invasion, Competition, and Biodiversity Loss in Urban Ecosystems. *BioScience* **2010**, *60*, 199–208. <https://doi.org/10.1525/bio.2010.60.3.6>.
58. Aronson, M.F.J.; La Sorte, F.A.; Nilon, C.H.; Katti, M.; Goddard, M.A.; Lepczyk, C.A.; Warren, P.S.; Williams, N.S.G.; Cilliers, S.; Clarkson, B.; et al. A Global Analysis of the Impacts of Urbanization on Bird and Plant Diversity Reveals Key Anthropogenic Drivers. *Proc. R. Soc. B-Biol. Sci.* **2014**, *281*, 20133330. <https://doi.org/10.1098/rspb.2013.3330>.
59. Goddard, M.A.; Dougill, A.J.; Benton, T.G. Why Garden for Wildlife? Social and Ecological Drivers, Motivations and Barriers for Biodiversity Management in Residential Landscapes. *Ecol. Econ.* **2013**, *86*, 258–273. <https://doi.org/10.1016/j.ecolecon.2012.07.016>.
60. Baldock, K.C.R.; Goddard, M.A.; Hicks, D.M.; Kunin, W.E.; Mitschunas, N.; Morse, H.; Osgathorpe, L.M.; Potts, S.G.; Robertson, K.M.; Scott, A.V.; et al. A Systems Approach Reveals Urban Pollinator Hotspots and Conservation Opportunities. *Nat. Ecol. Evol.* **2019**, *3*, 363. <https://doi.org/10.1038/s41559-018-0769-y>.
61. Zipperer, W.C. Species Composition and Structure of Regenerated and Remnant Forest Patches within an Urban Landscape. *Urban Ecosyst.* **2002**, *6*, 271–290. <https://doi.org/10.1023/B:UECO.0000004827.12561.d4>.
62. Narango, D.L.; Tallamy, D.W.; Marra, P.P. Nonnative Plants Reduce Population Growth of an Insectivorous Bird. *Proc. Natl. Acad. Sci. USA* **2018**, *115*, 201809259. <https://doi.org/10.1073/pnas.1809259115>.
63. Wood, E.M.; Esaian, S. The Importance of Street Trees to Urban Avifauna. *Ecol. Appl.* **2020**, *30*, e02149. <https://doi.org/10.1002/eap.2149>.
64. Lerman, S.B.; Warren, P.S. The Conservation Value of Residential Yards: Linking Birds and People. *Ecol. Appl.* **2011**, *21*, 1327–1339. <https://doi.org/10.1890/10-0423.1>.
65. Smallwood, N.L.; Wood, E.M. The Ecological Role of Native-Plant Landscaping in Residential Yards to Birds during the Non-breeding Period. *Ecosphere* **2023**, *14*, e4360. <https://doi.org/10.1002/ecs2.4360>.
66. Baldock, K.C.R.; Goddard, M.A.; Hicks, D.M.; Kunin, W.E.; Mitschunas, N.; Osgathorpe, L.M.; Potts, S.G.; Robertson, K.M.; Scott, A.V.; Stone, G.N.; et al. Where Is the UK's Pollinator Biodiversity? The Importance of Urban Areas for Flower-Visiting Insects. *Proc. R. Soc. B-Biol. Sci.* **2015**, *282*, 20142849. <https://doi.org/10.1098/rspb.2014.2849>.
67. Threlfall, C.G.; Mata, L.; Mackie, J.A.; Hahs, A.K.; Stork, N.E.; Williams, N.S.G.; Livesley, S.J. Increasing Biodiversity in Urban Green Spaces through Simple Vegetation Interventions. *J. Appl. Ecol.* **2017**, *54*, 1874–1883. <https://doi.org/10.1111/1365-2664.12876>.

68. Pollock, H.S.; Van Riper, C.J.; Goodson, D.J.; Lerman, S.B.; Hauber, M.E. The Public–Private Divide and Seasonal Variation Shape Bird Diversity in Greenspaces of Two Neighboring Midwestern USA Cities. *Landsc. Urban Plan.* **2024**, *247*, 105060. <https://doi.org/10.1016/j.landurbplan.2024.105060>.
69. Wood, E.M.; Esaian, S.; Benitez, C.; Ethington, P.J.; Longcore, T.; Pomara, L.Y. Historical Racial Redlining and Contemporary Patterns of Income Inequality Negatively Affect Birds, Their Habitat, and People in Los Angeles, California. *Ornithol. Appl.* **2024**, *126*, duad044. <https://doi.org/10.1093/ornithapp/duad044>.
70. Ibáñez-Álamo, J.D.; Morelli, F.; Benedetti, Y.; Rubio, E.; Jokimäki, J.; Pérez-Contreras, T.; Sprau, P.; Suhonen, J.; Tryjanowski, P.; Kaisanlahti-Jokimäki, M.-L.; et al. Biodiversity within the City: Effects of Land Sharing and Land Sparing Urban Development on Avian Diversity. *Sci. Total Environ.* **2020**, *707*, 135477. <https://doi.org/10.1016/j.scitotenv.2019.135477>.
71. Tratalos, J.; Fuller, R.A.; Evans, K.L.; Davies, R.G.; Newson, S.E.; Greenwood, J.J.; Gaston, K.J. Bird Densities Are Associated with Household Densities. *Glob. Change Biol.* **2007**, *13*, 1685–1695.
72. Campbell, C.E.; Jones, D.N.; Awasthy, M.; Castley, J.G.; Chauvenet, A.L. Big Changes in Backyard Birds: An Analysis of Long-Term Changes in Bird Communities in Australia’s Most Populous Urban Regions. *Biol. Conserv.* **2022**, *272*, 109671.
73. Heffernan, J.B.; Soranno, P.A.; Angilletta, M.J.; Buckley, L.B.; Gruner, D.S.; Keitt, T.H.; Kellner, J.R.; Kominoski, J.S.; Rocha, A.V.; Xiao, J.; et al. Macrosystems Ecology: Understanding Ecological Patterns and Processes at Continental Scales. *Front. Ecol. Environ.* **2014**, *12*, 5–14. <https://doi.org/10.1890/130017>.
74. Lerman, S.B.; Narango, D.L.; Avolio, M.L.; Bratt, A.R.; Engebretson, J.M.; Groffman, P.M.; Hall, S.J.; Heffernan, J.B.; Hobbie, S.E.; Larson, K.L.; et al. Residential Yard Management and Landscape Cover Affect Urban Bird Community Diversity across the Continental USA. *Ecol. Appl.* **2021**, *31*, e02455. <https://doi.org/10.1002/eap.2455>.
75. Rich, T.D.; Beardmore, C.J.; Berlanga, H.; Blancher, P.J.; Bradstreet, M.S.W.; Butcher, G.S.; Demarest, D.W.; Dunn, E.H.; Hunter, W.C.; Iñigo-Elias, E.E.; et al. *Partners in Flight North American Landbird Conservation Plan*; Cornell Lab of Ornithology: Ithaca, NY, USA, 2004.
76. Tirpak, J.M.; Jones-Farrand, D.T.; Thompson, F.R.; Twedt, D.J.; Baxter, C.K.; Fitzgerald, J.A.; Uihlein, W.B. Assessing Ecoregional-Scale Habitat Suitability Index Models for Priority Landbirds. *J. Wildl. Manag.* **2009**, *73*, 1307–1315. <https://doi.org/10.2193/2008-125>.
77. Tirpak, J.M.; Jones-Farrand, D.T.; Thompson, F.R., III; Twedt, D.J.; Nelson, M.D.; Uihlein, W.B., III. *Predicting Bird Habitat Quality from a Geospatial Analysis of FIA Data*; U.S. Government: Washington, DC, USA, 2009.
78. Jones-Farrand, D.T.; Fearer, T.M.; Thogmartin, W.E.; Iii, F.R.T.; Nelson, M.D.; Tirpak, J.M. Comparison of Statistical and Theoretical Habitat Models for Conservation Planning: The Benefit of Ensemble Prediction. *Ecol. Appl.* **2011**, *21*, 2269–2282. <https://doi.org/10.1890/10-1047.1>.
79. Buron, R.; Hostetler, M.E.; Andreu, M. Urban Forest Fragments vs Residential Neighborhoods: Urban Habitat Preference of Migratory Birds. *Landsc. Urban Plan.* **2022**, *227*, 104538. <https://doi.org/10.1016/j.landurbplan.2022.104538>.
80. Hoover, J.; Brittingham, M. Nest-Site Selection and Nesting Success of Wood Thrushes. *Wilson Bull.* **1998**, *110*, 375–383.
81. Jirinec, V.; Varian, C.P.; Smith, C.J.; Leu, M. Mismatch between Diurnal Home Ranges and Roosting Areas in the Wood Thrush (*Hylocichla Mustelina*): Possible Role of Habitat and Breeding Stage. *Auk* **2016**, *133*, 1–12. <https://doi.org/10.1642/AUK-15-76.1>.
82. Courtney Jones, S.K.; O’Loughlin, L.S.; Starrs, D.; Humphrey, J.E.; Pulsford, S.A.; Allan, H.; Beitzel, M.; Birgen, K.; Bond, S.; Bounds, J.; et al. Quantifying Taxon-Specific Habitat Connectivity Requirements of Urban Wildlife Using Structured Expert Judgement. *Biol. Conserv.* **2025**, *305*, 110991. <https://doi.org/10.1016/j.biocon.2025.110991>.
83. Nilon, C.H.; Warren, P.S.; Wolf, J. Urban Habitats: Baltimore Birdscape Study: Identifying Habitat and Land-Cover Variables for an Urban Bird-Monitoring Project. *Urban Habitats* **2011**, *6*. url: https://www.urbanhabitats.org/v06n01/baltimore_full.html
84. Grove, J.M.; Cadenasso, M.L.; Pickett, S.T.; Machlis, G.E.; Burch, W.R. *The Baltimore School of Urban Ecology: Space, Scale, and Time for the Study of Cities*; Yale University Press: New Haven, CT, USA, 2015; ISBN 978-0-300-10113-3.
85. Brooks, R.P. Improving Habitat Suitability Index Models. *Wildl. Soc. Bull.* **1997**, *25*, 163–167. <https://doi.org/10.2307/3783299>. url: <https://www.jstor.org/stable/3783299>
86. Conner, R.N. Foraging Habitats of Woodpeckers in Southwestern Virginia. *J. Field Ornithol.* **1980**, *51*, 119–127.
87. Adkins Giese, C.L.; Cuthbert, F.J. Woodpecker Nest Tree Characteristics in Upper Midwestern Oak Forests. *Can. Field-Nat.* **2005**, *119*, 367–376.
88. Straus, M.A.; Bavrlic, K.; Nol, E.; Burke, D.M.; Elliott, K.A. Reproductive Success of Cavity-Nesting Birds in Partially Harvested Woodlots. *Can. J. For. Res.* **2011**, *41*, 1004–1017. <https://doi.org/10.1139/X11-012>.
89. Groffman, P.M.; Cavender-Bares, J.; Bettez, N.D.; Grove, J.M.; Hall, S.J.; Heffernan, J.B.; Hobbie, S.E.; Larson, K.L.; Morse, J.L.; Neill, C.; et al. Ecological Homogenization of Urban USA. *Front. Ecol. Environ.* **2014**, *12*, 74–81. <https://doi.org/10.1890/120374>.

90. Newhouse, M.J.; Marra, P.P.; Johnson, L.S. Reproductive Success of House Wrens in Suburban and Rural Landscapes. *Wilson J. Ornithol.* **2008**, *120*, 99–104. <https://doi.org/10.1676/06-156.1>.
91. Stagoll, K.; Manning, A.D.; Knight, E.; Fischer, J.; Lindenmayer, D.B. Using Bird–Habitat Relationships to Inform Urban Planning. *Landsc. Urban Plan.* **2010**, *98*, 13–25. <https://doi.org/10.1016/j.landurbplan.2010.07.006>.
92. Roman, L.A.; McPherson, E.G.; Scharenbroch, B.C.; Bartens, J. Identifying Common Practices and Challenges for Local Urban Tree Monitoring Programs across the United States. *Arboric. Urban For. (AUF)* **2013**, *39*, 292–299. <https://doi.org/10.48044/jauf.2013.038>.
93. Lepczyk, C.A.; La Sorte, F.A.; Aronson, M.F.J.; Goddard, M.A.; MacGregor-Fors, I.; Nilon, C.H.; Warren, P.S. Global Patterns and Drivers of Urban Bird Diversity. In *Ecology and Conservation of Birds in Urban Environments*; Murgui, E., Hedblom, M., Eds.; Springer International Publishing: Cham, Switzerland, 2017; pp. 13–33, ISBN 978-3-319-43314-1.
94. USDA Forest Service Tree Canopy Cover (TCC) Product Suite 2025.url: <https://data.fs.usda.gov/geodata/rastergateway/treecanopycover/#table1> Accessed May 18, 2026
95. U.S. Geological Survey Annual NLCD Collection 1 Science Products 2025. <https://doi.org/10.5066/P94UXNTS>
96. Sullivan, B.L.; Aycrigg, J.L.; Barry, J.H.; Bonney, R.E.; Bruns, N.; Cooper, C.B.; Damoulas, T.; Dhondt, A.A.; Dietterich, T.; Farnsworth, A.; et al. The eBird Enterprise: An Integrated Approach to Development and Application of Citizen Science. *Biol. Conserv.* **2014**, *169*, 31–40. <https://doi.org/10.1016/j.biocon.2013.11.003>.
97. Callaghan, C.T.; Bino, G.; Major, R.E.; Martin, J.M.; Lyons, M.B.; Kingsford, R.T. Heterogeneous Urban Green Areas Are Bird Diversity Hotspots: Insights Using Continental-Scale Citizen Science Data. *Landsc. Ecol.* **2019**, *34*, 1231–1246.
98. Li, E.; Parker, S.S.; Pauly, G.B.; Randall, J.M.; Brown, B.V.; Cohen, B.S. An Urban Biodiversity Assessment Framework That Combines an Urban Habitat Classification Scheme and Citizen Science Data. *Front. Ecol. Evol.* **2019**, *7*, 277.
99. Łopucki, R.; Kiersztyn, A. Urban Green Space Conservation and Management Based on Biodiversity of Terrestrial Fauna—A Decision Support Tool. *Urban For. Urban Green.* **2015**, *14*, 508–518. <https://doi.org/10.1016/j.ufug.2015.05.001>.
100. Murveit, A.M.; Gerlak, A.K.; Kaibara, M.D.; Hovis, M.; Minde, J.; Colella, T. Geospatial Public Participation Tool Adoption in U.S. Natural Resource Agencies. *Soc. Nat. Resour.* **2025**, *38*, 587–605. <https://doi.org/10.1080/08941920.2025.2450693>.
101. Dykstra, B.L.; Rumble, M.A.; Flake, L.D. Effects of Timber Harvesting on Birds in the Black Hills of South Dakota and Wyoming, USA. In Proceedings of the In: Cook, James E.; Oswald, Brian P., comps. First biennial North American forest ecology workshop: June 24–26, 1997, North Carolina St. Univ., Raleigh, NC.[Raleigh, NC: North Carolina State University]: 16–26; 1997.
102. Lehmkuhl, J.F.; Burger, E.D.; Drew, E.K.; Lindsey, J.P.; Haggard, M.; Woodruff, K.Z. Breeding Birds in Riparian and Upland Dry Forests of the Cascade Range. *J Wildl Manag* **2007**, *71*, 2632–2643, doi:10.2193/2007-004.
103. Rodewald, A.D.; Abrams, M.D. Floristics and Avian Community Structure: Implications for Regional Changes in Eastern Forest Composition. *Forest Science* **2002**, *48*, 267–272.
104. Rodewald, P.G.; Smith, K.G. Short-Term Effects of Understory and Overstory Management on Breeding Birds in Arkansas Oak-Hickory Forests. *The Journal of wildlife management* **1998**, 1411–1417.
105. Garrison, B.A. Avian Population Densities, Habitat Use, and Foraging Ecology in Thinned and Unthinned Hardwood Forests in Southwestern Virginia. PhD Thesis, Virginia Polytechnic Institute and State University, 1986.
106. Mills, T.R.; Rumble, M.A.; Flake, L.D. Habitat of Birds in Ponderosa Pine and Aspen/Birch Forest in the Black Hills, South Dakota. *Journal of Field Ornithology* **2000**, *71*, 187–206.
107. Annand, E.M.; Thompson, F.R. Forest Bird Response to Regeneration Practices in Central Hardwood Forests. *The Journal of Wildlife Management* **1997**, *61*, 159–171, doi:10.2307/3802425.
108. Greenberg, C.H.; Waldrop, T.A.; Tomcho, J.; Phillips, R.J.; Simon, D. Bird Response to Fire Severity and Repeated Burning in Upland Hardwood Forest. *Forest Ecology and Management* **2013**, *304*, 80–88.
109. Greenberg, C.H.; Lanham, J.D. Breeding Bird Assemblages of Hurricane-Created Gaps and Adjacent Closed Canopy Forest in the Southern Appalachians. *Forest Ecology and Management* **2001**, *154*, 251–260.
110. Meiklejohn, B.A.; Hughes, J.W. Bird Communities in Riparian Buffer Strips of Industrial Forests. *The American midland naturalist* **1999**, *141*, 172–184.
111. Chalfoun, A.D.; Ratnaswamy, M.J.; Thompson, F.R. Songbird Nest Predators in Forest–Pasture Edge and Forest Interior in a Fragmented Landscape. *Ecological Applications* **2002**, *12*, 858–867, doi:10.1890/1051-0761(2002)012[0858:SNPIFP]2.0.CO;2.
112. Pennington, D.N.; Blair, R.B. Habitat Selection of Breeding Riparian Birds in an Urban Environment: Untangling the Relative Importance of Biophysical Elements and Spatial Scale. *Diversity and Distributions* **2011**, *17*, 506–518, doi:10.1111/j.1472-4642.2011.00750.x.

113. Tarvin, K.A.; Smith, K.G. Microhabitat Factors Influencing Predation and Success of Suburban Blue Jay *Cyanocitta Cristata* Nests. *Journal of Avian Biology* 1995, 296–304.
114. Chávez-Zichinelli, C.A.; MacGregor-Fors, I.; Rohana, P.T.; Valdéz, R.; Romano, M.C.; Schondube, J.E. Stress Responses of the House Sparrow (*Passer Domesticus*) to Different Urban Land Uses. *Landscape and Urban Planning* 2010, 98, 183–189.
115. Williamson, R.D.; DeGraaf, R.M. Habitat Associations of Ten Bird Species in Washington, DC. *Urban Ecology* 1981, 5, 125–136, doi:10.1016/0304-4009(81)90005-X.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.