

10190

Great Basin Pinyon-Juniper Woodland

BpS Model/Description Version: Aug. 2020

Modelers		Reviewers	
Jim Hurja	jhurja@fs.fed.us	None	None
None	None	None	None
None	None	None	None

Vegetation Type

Forest and Woodland

Map Zones

15

Geographic Range

This ecological system occurs on dry mountain ranges of the Great Basin region and eastern foothills of the Sierra Nevada.

Biophysical Site Description

System typically found from 5,500-8,000ft above the blackbrush (*Coleogyne ramosissima*) zone. This type generally occurred on most soil types and landforms, including fire-safe sites of steep and rocky slopes. Severe climatic events occurring during the growing season, such as frosts and drought, are thought to limit the distribution of pinyon-juniper woodlands to relatively narrow altitudinal belts on mountainsides. Soils supporting this system vary in texture ranging from stony, cobbly, gravelly sandy loams to clay loam or clay.

Vegetation Description

Woodlands dominated by a mix of *Pinus monophylla* and *Juniperus osteosperma*, pure or nearly pure occurrences of *Pinus monophylla*, or woodlands dominated solely by *Juniperus osteosperma* comprise this system. *Cercocarpus ledifolius* is a common associate. Understory layers are variable. Associated species include shrubs such as *Arctostaphylos patula*, *Arctostaphylos pungens*, *Artemisia nova*, *Artemisia tridentata*, *Cercocarpus ledifolius*, *Cercocarpus intricatus*, *Coleogyne ramosissima*, *Purshia stansburiana*, *Ceanothus greggii*, *Symphoricarpos oreophilus*, *Garrya falvescens*, *Yucca baccata*, and bunch grasses *Pseudoroegneria spicata*, *Achnatherum hymenoides*, *Elymus elymoides*, and *Poa fendleriana*. *Quercus gambelii* and *Quercus turbinella* may be present.

Since disturbance was uncommon to rare in this ecological system and the overstory conifers may have lived for several hundred years, patches were primarily composed of later seral stages (D & E; see below) that did not occur as extensive woodlands, and that should be distinguished from shrubland ecological sites encroached by pinyon or juniper during the last 150yrs. The age structure may vary from uneven to even aged. The overstory cover is normally > 25%, although it can sometimes be higher (up to 60%) where pinyon occurs.

BpS Dominant and Indicator Species

Symbol	Scientific Name	Common Name
PIMO	<i>Pinus monophylla</i>	Singleleaf pinyon
JUOS	<i>Juniperus osteosperma</i>	Utah juniper
CELE3	<i>Cercocarpus ledifolius</i>	Curl-leaf mountain mahogany
PUST	<i>Purshia stansburiana</i>	Stansbury cliffrose
ARPA	<i>Arabis parishii</i>	Dwarf rockcress
ARTR2	<i>Artemisia tridentata</i>	Big sagebrush

Species names are from the NRCS PLANTS database. Check species codes at <http://plants.usda.gov>.

Disturbance Description

Uncertainty exists about the fire frequencies of this ecological system, especially since this ecological system groups different types of pinyon-juniper communities for different slopes, exposures, and elevations. Fire occurrence may be influenced by fires spreading from shrub and grassland dominated vegetation of lower and higher altitudinal zones. Replacement fires were uncommon to rare and occurred primarily during extreme fire behavior conditions and during long droughts. Mixed severity fire was characterized as a mosaic of replacement and surface fires distributed through the patch at a fine scale (<0.1ac). There is limited evidence for surface fires (Gruell 1994; Bauer and Weisberg, unpublished data), which likely occurred only in the more productive sites during years where understory grass cover was high, providing adequate fuel. Although fire scars are only rarely found in pinyon-juniper of the Colorado Plateau and elsewhere (Baker and Shinneman 2004, Eisenhart 2004), ongoing studies in the central Great Basin are observing fire-scarred trees, suggesting that surface fires historically occurred at low frequency. Limited evidence to date suggests that while lightning ignitions in this biophysical setting may have been common, the resulting fires only rarely spread to affect more than a few trees.

Prolonged weather-related stress (mostly drought), insects, and tree pathogens are coupled disturbances that thin trees to varying degrees and kills small patches with greater frequency in more closed stands. The most common disturbance in this type is very small scale--either single tree, or small groups. If the conditions are right, it will have replacement fires that burn stands up to a maximum of 1000s acres. This type may also have mixed severity fires of 10-100s of acres.

Fire Frequency

Severity	Avg FI	Percent of All Fires	Min FI	Max FI
Replacement	479	33		
Moderate (Mixed)	338	47		
Low (Surface)	771	20		
All Fires	158	100		

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is the central tendency modeled. Percent of all fires is the percent of all fires modeled in that severity class. Minimum and Maximum FIs show the relative range of fire intervals as estimated by model contributors, if known.

Scale Description

10,000ac possible, more likely 1,000ac.

Adjacency or Identification Concerns

This system occurs at lower elevations than Colorado Plateau Pinyon-Juniper Woodland (BpS 1016) where sympatric. In modern days, surrounding matrix shrubland vegetation (blackbrush

and sagebrush) has changed somewhat to young-mid-aged woodlands. Two major modern issues, climate change and invasive plant species (especially annual grasses red brome and cheatgrass), lead to non-equilibrium vegetation dynamics for this ecological system, making it difficult to categorize and usefully apply natural disturbance regimes. Sites with an important annual grass component in the understory experience greater fire frequency, and result in more intense fire and greater spread. Moreover, fire from adjacent BpS invaded by annual grasses will spread more frequently into in BpS 1019, which is fire sensitive.

Issues or Problems

There is much uncertainty in model parameters, particularly the fire regime. Quantitative data are lacking and research is on-going. The literature for this ecological system's fire history is based on the chronologies from other pines species that are better fire recorders, growing under conditions that may not represent fire environments typical of infrequent-fire pinyon and juniper communities. For example, surface fire, which leaves scars on these other pine species (but not generally on fire-sensitive pinyon or juniper), has no effect on the dynamics of the model, although surface fire maintains the open structure of classes D and E by thinning younger trees.

Further study is needed to better elucidate the independent and interactive effects of fire, insects, pathogens, climate, grazing and anthropogenic impacts on historical and current vegetation dynamics in the Great Basin Pinyon-Juniper Woodland type.

Native Uncharacteristic Conditions

Cover > 60% is uncharacteristic.

Comments

Succession Classes

Mapping Rules

Upper Layer Lifeform	Height (m)	Canopy Cover (%)									
		0-10	11-20	21-30	31-40	41-50	51-60	61-70	71-80	81-90	91-100
Herb	0-0.5	A	A	A	A	A	A	A	A	A	A
Herb	0.5-1.0	A	A	A	A	A	A	A	A	A	A
Herb	>1.0	A	A	A	A	A	A	A	A	A	A
Shrub	0-0.5	B	B	B	B	B	B	UN	UN	UN	UN
Shrub	0.5-1.0	B	B	B	B	B	B	UN	UN	UN	UN
Shrub	1.0-3.0	B	B	B	B	B	B	UN	UN	UN	UN
Shrub	>3.0	B	B	B	B	B	B	UN	UN	UN	UN
Tree	0-5	C	C	D	D	D	D	UN	UN	UN	UN
Tree	5-10	D	D	D	D	D	D	UN	UN	UN	UN
Tree	10-25	D	D	D	D	D	D	UN	UN	UN	UN
Tree	25-50	D	D	D	D	D	D	UN	UN	UN	UN
Tree	>50	D	D	D	D	D	D	UN	UN	UN	UN

Succession class letters A-E are described in the Succession Class Description section. Some classes use a leafform distinction where a qualifier is added to the class letter: Brdl (broadleaf), Con (conifer), or Mix (mixed conifer and broadleaf). UN refers to uncharacteristic native or a combination of height and cover that would not be expected under the reference condition. NP refers to not possible or a combination of height and cover which is not physiologically possible for the species in the BpS.

Description

Class A	2	Early Development 1 - Open
----------------	----------	-----------------------------------

Indicator Species

Symbol	Scientific Name	Common Name	Canopy Position
ELEL5	Elymus elymoides	Squirreltail	Mid-Upper
SPHAE	Sphaeralcea	Globemallow	Upper
ZIPA2	Zigadenus paniculatus	Foothill deathcamas	Mid-Upper
POFE	Poa fendleriana	Muttongrass	Mid-Upper

Description

Initial post-fire community dominated by annual grasses and forbs. Later stages of this class contain greater amounts of perennial grasses and forbs. Evidence of past fires (burnt stumps and charcoal) should be observed.

Maximum Tree Size Class

No Data

Class B	7	Mid Development 1 - Open
----------------	----------	---------------------------------

Indicator Species

Symbol	Scientific Name	Common Name	Canopy Position
ARTR2	Artemisia tridentata	Big sagebrush	Upper
ARPU5	Arctostaphylos pungens	Pointleaf manzanita	Upper
PIMO	Pinus monophylla	Singleleaf pinyon	Mid-Upper
JUOS	Juniperus osteosperma	Utah juniper	Mid-Upper

Description

Dominated by shrubs, perennial forbs and grasses. Tree seedlings starting to establish on favorable microsites. Total cover remains low due to shallow unproductive soil. It is important to note that replacement fire at this stage does not eliminate perennial grasses. Mixed severity fire thins the woody vegetation.

Maximum Tree Size Class

No data

Class C	27	Mid Development 2 - Open
----------------	-----------	---------------------------------

Indicator Species

Symbol	Scientific Name	Common Name	Canopy Position
PIMO	Pinus monophylla	Singleleaf pinyon	Upper

Replacement Fire	Mid1:OPN	Early1:OPN	0.005	200	Yes	0
Insects or Disease	Mid2:OPN	Mid1:OPN	0.001	1000	Yes	0
Replacement Fire	Mid2:OPN	Early1:OPN	0.004	250	Yes	0
Mixed Fire	Mid2:OPN	Mid2:OPN	0.005	200	No	0
Replacement Fire	Late1:OPN	Early1:OPN	0.001	1000	Yes	0
Insects or Disease	Late1:OPN	Mid2:OPN	0.002	500	Yes	0
Insects or Disease	Late1:OPN	Mid1:OPN	0.002	500	Yes	0
Surface Fire	Late1:OPN	Late1:OPN	0.002	500	No	0
Mixed Fire	Late1:OPN	Late1:OPN	0.002	500	No	0

References

- Alexander, R.R. and F. Ronco, Jr. 1987. Classification of the forest vegetation on the National Forests of Arizona and New Mexico. Res. Note RM-469. Fort Collins, CO. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station. 10 pp.
- Anderson, H.E. 1982. Aids to Determining Fuel Models For Estimating Fire Behavior. GTR-INT-122. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station. 22 pp.
- Arno, S.F. 2000. Fire in western forest ecosystems. In: J.K. Brown and J Kapler-Smith, eds. Wildland fire in ecosystems: Effects of fire on flora. RMRS-GTR-42-vol. 2. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station: 97-120.
- Baker, W.L. and D.J. Shinneman. 2004. Fire and restoration of pinyon-juniper woodlands in the western United States. A review. Forest Ecology and Management 189: 1-21.
- Barney, M.A. and N.C. Frischknecht. 1974. Vegetation changes following fire in the Pinyon-Juniper type of West-Central Utah. Jour. Range Manage. 27: 91-96.
- Bradley, A.F., N.V. Noste and W.C. Fischer. 1992. Fire Ecology of Forests and Woodlands in Utah. Gen.Tech. Rep. GTR-INT-287. Ogden, UT: USDA Forest Service, Intermountain Research Station. 127 pp.
- Brown, J.K. and J. Kapler-Smith, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station. 257 pp.
- Despain, D.W. and J.C. Mosely. 1990. Fire history and stand structure of a pinyon-juniper woodland at Walnut Canyon National Monument, Arizona. USDI National Park Service

Technical Report No. 34. Cooperative National Park Resources Studies Unit, University of Arizona, Tucson AZ. 27 pp.

Eisenhart, K.S. 2004. Historic range of variability and stand development in pinon-juniper woodlands of western Colorado. Boulder, Colorado: University of Colorado. Dissertation.

Erdman, J.A. 1970. Pinyon-juniper succession after natural fires on residual soils of Mesa Verde, Colorado. Science Bulletin, Biological Series --Volume XI, No. 2. Brigham Young University, Provo, UT. 26 pp.

Everett, R.L. and K. Ward. 1984. Early Plant Succession on Pinyon-Juniper Controlled Burns. Northwest Science 58: 57-68.

Eyre, F.H., ed. 1980. Forest cover types of the United States and Canada. Washington, DC: Society of American Foresters. 148 pp.

Goodrich, S. and B. Barber. 1999. Return Interval for Pinyon-Juniper Following Fire in the Green River Corridor, Near Dutch John, Utah. In: USDA Forest Service Proceedings RMRS-P-9.

Gruell, G.E. 1999. Historical and modern roles of fire in pinyon-juniper. In: Monsen, Stephen B.; Stevens, Richard, compilers. Proceedings: ecology and management of pinyon-juniper communities within the Interior West: Sustaining and restoring a diverse ecosystem; 1997 September 15-18; Provo, UT. Proceedings RMRS-P-9. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station: 24-28.

Gruell, G.E., L.E. Eddleman and R. Jaindl. 1994. Fire History of the Pinyon-Juniper Woodlands of Great Basin National Park. Technical Report NPS/PNROSU/NRTR-94/01. U.S. Department of Interior, National Park Service, Pacific Northwest Region. 27 pp.

Hardy, C.C., K.M. Schmidt, J.P. Menakis and R.N. Samson. 2001. Spatial data for national fire planning and fuel management. Int. J. Wildland Fire. 10(3&4): 353-372.

Hessburg, P.F., B.G. Smith, R.B. Salter, R.D. Ottmar and E. Alvarado. 2000. Recent changes (1930s-1990s) in spatial patterns of interior northwest forests, USA. Forest Ecology and Management 136: 53-83.

Kilgore, B.M. 1981. Fire in ecosystem distribution and structure: western forests and scrublands. Pages 58-89 in: H.A. Mooney et al., technical coordinators. Proceedings: Conference on Fire Regimes and Ecosystem Properties, Honolulu, 1978. Gen. Tech. Rep. WO-GTR-26.

Kuchler, A.W. 1964. Potential Natural Vegetation of the Conterminous United States. American Geographic Society Special Publication No. 36. 116 pp.

NatureServe. 2004. International Ecological Classification Standard: Terrestrial Ecological Classifications. Terrestrial ecological systems of the Great Basin US: DRAFT legend for Landfire project. NatureServe Central Databases. Arlington, VA. Data current as of 4 November 2004.

NatureServe. 2007. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. Data current as of 10 February 2007.

Ogle, K. and V. DuMond. 1997. Historical Vegetation on National Forest Lands in the Intermountain Region. USDA Forest Service, Intermountain Region, Ogden, UT. 129 pp.

Ott, J.E., E.D. McArthur and S.C. Sanderson. 2001. Plant Community Dynamics of Burned and Unburned Sagebrush and Pinyon-Juniper Vegetation in West-Central Utah. In: Proceedings, USDA Forest Service RMRS-P-9. 177-190.

Romme, W.H., L. Floyd-Hanna and D. Hanna. 2002. Ancient Pinyon-Juniper forests of Mesa Verde and the West: A cautionary note for forest restoration programs. In: Conference Proceedings – Fire, Fuel Treatments, and Ecological Restoration: Proper Place, Appropriate Time, Fort Collins, CO, April 2002. 19 pp.

Rondeau, R. 2001. Ecological System Viability Specifications for Southern Rocky Mountain Ecoregion. Colorado Natural Heritage Program. 181 pp.

Schmidt, K.M., J.P. Menakis, C.C. Hardy, W.J. Hann and D.L. Bunnell. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: USDA Forest Service, Rocky Mountain Research Station. 41 pp. + CD.

Soule', P.T. and P.A. Knapp. 1999. Western juniper expansion on adjacent disturbed and near-relict sites. *Journal of Range Management* 52: 525-533.

Soule', P.T. and P.A. Knapp. 2000. *Juniperus occidentalis* (western juniper) establishment history on two minimally disturbed research natural areas in central Oregon. *Western North American Naturalist* (60)1: 26-33.

Stein, S.J. 1988. Fire History of the Paunsaugunt Plateau in Southern Utah. *Great Basin Naturalist*. 48: 58-63.

Tausch, R.J., N.E. West and A.A. Nabi. 1981. Tree Age and Dominance Patterns in Great Basin Pinyon-Juniper Woodlands. *Jour. Range. Manage.* 34: 259-264.

Tausch, R.J. and N.E. West. 1987. Differential Establishment of Pinyon and Juniper Following Fire. *The American Midland Naturalist* 119(1). 174-184.

USDA Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (2002, December). Fire Effects Information System, [Online]. Available: <http://www.fs.fed.us/database/feis/> [Accessed: 11/15/04].

Ward, K.V. 1977. Two-Year Vegetation Response and Successional Trends for Spring Burns in the Pinyon-Juniper Woodland. M.S. Thesis, University of Nevada, Reno. 54 pp.

Wright, H.A., L.F. Neuenschwander and C.M. Britton. 1979. The role and use of fire in Sagebrush Grass and Pinyon-Juniper Plant Communities. Gen. Tech. Rep. INT-GTR-58. Ogden, UT: USDA Forest Service, Intermountain Research Station. 48 pp.

Young, J.A., and R.A. Evans. 1978. Population Dynamics after Wildfires in Sagebrush Grasslands. *Journal of Range Management* 31: 283-289.

Young, J.A. and R.A. Evans. 1981. Demography and Fire History of a Western Juniper Stand. *Journal of Range Management* 34: 501-505.