

10560

Rocky Mountain Subalpine Mesic-Wet Spruce-Fir Forest and Woodland

BpS Model/Description Version: Aug. 2020

Modelers		Reviewers	
Chris Thomas	cthomas@fs.fed.us	Elena Contreras	econtreras@tnc.org
Dennis Sandbak	dsandbak@fs.fed.us	None	None
Paul Mock	pmock@fs.fed.us	None	None

Vegetation Type

Forest and Woodland

Map Zones

29

Geographic Range

Northern Rockies, including MT, ID and WY. In MZ29, this occurs in the Bighorn Mountains, and there is some spruce/fir in the Laramie Peak Range. Sections M331B and M331I (Cleland et al. 2007).

Biophysical Site Description

Upper subalpine zone and mesic sites. Occurrences are typically found in locations with cold-air drainage or ponding, or where snowpacks linger late into the summer, such as north-facing slopes and high-elevation ravines. They can extend down in elevation below the subalpine zone in places where cold-air ponding occurs; northerly and easterly aspects predominate. These forests are found on gentle to very steep mountain slopes, high-elevation ridgetops and upper slopes, plateau-like surfaces, basins, alluvial terraces, well-drained benches, and inactive stream terraces. Reviewers for MZ20 felt that the preceding sentence does not align with this BpS shrub understory component, as this BpS has moist-driven shrubs and may occur near riparian areas.

Vegetation Description

Engelmann spruce and subalpine fir dominate on most aspects with lodgepole pine comprising a greater component on dryer sites or earlier successional stages. Early succesional vegetation contains *Eurybia conspicua* and *Carex geyeri*. *Vaccinium scoparium* is a common understory associate in later successional stages.

Mesic understory shrubs include *Menziesia ferruginea*, *Vaccinium membranaceum*, *Rubus parviflorus*, and *Ledum glandulosum*. Herbaceous species include *Actaea rubra*, *Maianthemum stellatum*, *Cornus canadensis*, *Erigeron engelmannii*, *Saxifraga bronchialis*, *Lupinus argenteus* ssp. *subalpinus*, *Valeriana sitchensis*, and graminoids such as *Carex generii* and *Calamagrostis canadensis*.

BpS Dominant and Indicator Species

Symbol	Scientific Name	Common Name
PIEN	<i>Picea engelmannii</i>	Engelmann spruce

ABLA	<i>Abies lasiocarpa</i>	Subalpine fir
PICO	<i>Pinus contorta</i>	Lodgepole pine

Species names are from the NRCS PLANTS database. Check species codes at <http://plants.usda.gov>.

Disturbance Description

Fire Regime Group V or IV; primarily long-interval stand replacement fires. In some areas, spruce beetle can influence successional stage, species composition and stand density. Spruce beetle may act to accelerate succession.

It has been suggested that this system is not outside of its HRV for fire frequency and severity. Fire interval could be greater than 400yrs at times (Romme, personal correspondence; Veblen et al. 1991, 1994), or between 335-400yrs (Bradley et al. 1992), and there is no equilibrium achieved in this system, as it fluctuates widely normally in each class.

Fire Frequency

Severity	Avg FI	Percent of All Fires	Min FI	Max FI
Replacement	346	75	100	600
Moderate (Mixed)	1061	25		
Low (Surface)				
All Fires	261	100		

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is the central tendency modeled. Percent of all fires is the percent of all fires modeled in that severity class. Minimum and Maximum FIs show the relative range of fire intervals as estimated by model contributors, if known.

Scale Description

Fires could range from 1,000s-10,000s of acres. Variability of climate, topography and other site factors can result in a wide range of representation of successional stages on the landscape. Equilibrium landscapes are not likely to develop in areas <500,000ac, or perhaps not in areas greater than that either (Romme, personal correspondence). Reviewers state that typical fires in this system are small clumps of stand replacing fires with spots to adjacent clumps.

Adjacency or Identification Concerns

Adjacent to drier, lower subalpine forests (lodgepole-spruce-fir) and to krummholz and alpine vegetation. This system typically has more precipitation and longer winters than lower subalpine types.

This BpS 1056 will be difficult to distinguish from the 1050/1055 system in MZ29. Note that some of the Englemann spruce/subalpine fir should be keyed to 1056 as well in MZ29.

Climate (severely dry conditions) is the primary driver of fire regimes in this system. Long-term changes in climate as well as inter-annual climate variability will affect the frequency of fire in this system.

This BpS corresponds to the following habitat types (Pfister et al. 1977): ABLA/ALSI, ABLA/CAGE, ABLA/VASC, TSME/XETE, TSME/MEFE, TSME/CLUN, PICEA/GART, PICEA/LIBO and PICEA/PHMA.

It has been suggested that this system is not outside of its historic range of variability for proportions of seral stages (Romme, personal correspondence; Veblen et al. 1991, 1994), and there is no equilibrium achieved in this system, as it fluctuates widely normally in each class.

Issues or Problems

Currently, balsam bark beetle is killing subalpine fir trees throughout the Rocky Mountain subalpine mesic spruce fir region.

Reviewers state that since 1990, subalpine fir mortality complex has increased.

Native Uncharacteristic Conditions

Comments

Succession Classes

Mapping Rules

Upper Layer Lifeform	Height (m)	Canopy Cover (%)									
		0-10	11-20	21-30	31-40	41-50	51-60	61-70	71-80	81-90	91-100
Herb	0-0.5	UN	UN	UN	UN	UN	UN	UN	UN	UN	UN
Herb	0.5-1.0	UN	UN	UN	UN	UN	UN	UN	UN	UN	UN
Herb	>1.0	UN	UN	UN	UN	UN	UN	UN	UN	UN	UN
Shrub	0-0.5	UN	UN	UN	UN	UN	UN	UN	UN	UN	UN
Shrub	0.5-1.0	UN	UN	UN	UN	UN	UN	UN	UN	UN	UN
Shrub	1.0-3.0	UN	UN	UN	UN	UN	UN	UN	UN	UN	UN
Shrub	>3.0	UN	UN	UN	UN	UN	UN	UN	UN	UN	UN
Tree	0-5	A	A	A	A	A	A	A	A	A	A
Tree	5-10	C	C	C	C	B	B	B	B	B	B
Tree	10-25	D	D	D	D	D	E	E	E	E	E
Tree	25-50	D	D	D	D	D	E	E	E	E	E
Tree	>50	D	D	D	D	D	E	E	E	E	E

Succession class letters A-E are described in the Succession Class Description section. Some classes use a leafform distinction where a qualifier is added to the class letter: Brdl (broadleaf), Con (conifer), or Mix (mixed conifer and broadleaf). UN refers to uncharacteristic native or a combination of height and cover that would not be expected under the reference condition. NP refers to not possible or a combination of height and cover which is not physiologically possible for the species in the BpS.

Description

Class A	8	Early Development 1 - All Structures
----------------	----------	---

Indicator Species

Symbol	Scientific Name	Common Name	Canopy Position
PIEN	<i>Picea engelmannii</i>	Engelmann spruce	Upper
PICO	<i>Pinus contorta</i>	Lodgepole pine	Upper
ABLA	<i>Abies lasiocarpa</i>	Subalpine fir	Mid-Upper

Description

Symbol	Scientific Name	Common Name	Canopy Position
--------	-----------------	-------------	-----------------

PIEN	<i>Picea engelmannii</i>	Engelmann spruce	Upper
ABLA	<i>Abies lasiocarpa</i>	Subalpine fir	Upper
PICO	<i>Pinus contorta</i>	Lodgepole pine	Upper

Description

Poles (five inches DBH+) and larger diameter moderately shade tolerant conifer species in small to moderate size patches. Patches would include subalpine fir seedlings.

This stage occupies 15-50% of the landscape.

Maximum Tree Size Class

Very Large >33"DBH

Class E 47 Late Development 1 - Closed

Indicator Species

Symbol	Scientific Name	Common Name	Canopy Position
PIEN	<i>Picea engelmannii</i>	Engelmann spruce	Upper
ABLA	<i>Abies lasiocarpa</i>	Subalpine fir	Upper
PICO	<i>Pinus contorta</i>	Lodgepole pine	Upper

Description

Pole and larger diameter moderately to shade tolerant conifer species, in moderate to large size patches, all aspects.

Maximum Tree Size Class

Very Large >33"DBH

Model Parameters

Deterministic Transitions

From Class	Begins at (yr)	Succeeds to	After (years)
Early1:ALL	0	Mid1:CLS	29
Mid1:OPN	30	Late1:OPN	100
Mid1:CLS	30	Late1:CLS	100
Late1:OPN	101	Late1:OPN	999
Late1:CLS	101	Late1:CLS	999

Probabilistic Transitions

Disturbance Type	Disturbance occurs In	Moves vegetation to	Disturbance Probability	Return Interval (yrs)	Reset Age to New Class Start Age After Disturbance?	Years Since Last Disturbance
Alternative Succession	Early1:ALL	Mid1:OPN	0.001	1000	Yes	0
Replacement Fire	Early1:ALL	Early1:ALL	0.002	500	Yes	0
Replacement Fire	Mid1:OPN	Early1:ALL	0.002	500	Yes	0

Wind or Weather or Stress	Mid1:CLS	Mid1:OPN	0.001	1000	Yes	0
Replacement Fire	Mid1:CLS	Early1:ALL	0.002	500	Yes	0
Insects or Disease	Late1:OPN	Mid1:OPN	0.001	1000	Yes	0
Replacement Fire	Late1:OPN	Early1:ALL	0.0025	400	Yes	0
Alternative Succession	Late1:OPN	Late1:CLS	0.004	250	Yes	0
Insects or Disease	Late1:CLS	Late1:OPN	0.001	1000	Yes	0
Mixed Fire	Late1:CLS	Late1:OPN	0.002	500	Yes	0
Replacement Fire	Late1:CLS	Early1:ALL	0.0035	286	Yes	0

References

Alexander, R.R., G.R. Hoffman and J.M. Wirsing. 1986. Forest vegetation of the Medicine Bow National Forest in southeastern Wyoming: a habitat type classification. Research Paper RM-271. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station. On file at MBR-TB SO, Laramie, WY.

Alexander, R.R. 1986. Silvicultural systems and cutting methods for old-growth spruce-fir forests in the central and southern Rocky Mountains. General Technical Report RM-126. Fort Collins, CO: Rocky Mountain Forest and Range Experiment Station.

Alexander, R.R. 1988. Forest vegetation on national forests in the Rocky Mountain and Intermountain regions: habitat types and community types. General Technical Report RM-162. Fort Collins, CO: Rocky Mountain Forest and Range Experiment Station.

Alexander, R.R. and O. Engelby. 1983. Engelmann spruce - subalpine fir. In: Silvicultural systems for the major forest types of the United States. Agriculture Handbook 445. Washington DC: US Dept. of Agriculture.

Arno, S.F. 2000. Fire in western forest ecosystems. Pages 97-120 in: J.K. Brown and J. Kapler-Smith, eds. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station. 257 pp.

Arno, S.F. 2000. Fire in western forest ecosystems. Pages 97-120 in: J.K. Brown and J. Kapler-Smith, eds. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station. 257 pp.

Baker, W.L. and T.T. Veblen. 1990. Spruce beetles and fires in the nineteenth-century subalpine forests of western Colorado, U.S.A. Arctic and Alpine Research 22(1): 65-80.

Baker, W.L. 1994. Landscape structure measurements for watersheds in the Medicine Bow National Forest using GIS analysis. Department of Geography and Recreation, Univ. of Wyoming. Prepared under agreement with the USDA Forest Service, Medicine Bow NF. On file at Medicine Bow-Routt NFs and Thunder Basin NG Supervisor's Office, Laramie, WY.

Barrett, S.W. 1994. Fire regimes on andesitic mountain terrain in northeastern Yellowstone National Park. *International Journal of Wildland Fire* 4: 65-76.

Bradley, A.F., W.C. Fischer and N.V. Noste. 1992. Fire ecology of the forest habitat types of eastern Idaho and western Wyoming. Gen. Tech. Rep. INT-290. Ogden, UT: USDA Forest Service, Intermountain Research Station. 92 pp.

Buechling, A. and W.L. Baker. 2004. A fire history from tree rings in a high-elevation forest of Rocky Mountain National Park. *Canadian Journal of Forest Research* 34: 1259-1273.

Cleland, D.T.; Freeouf, J.A.; Keys, J.E.; Nowacki, G.J.; Carpenter, C.A.; and McNab, W.H. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. Gen. Tech. Report WO-76D [Map on CD-ROM] (A.M. Sloan, cartographer). Washington, DC: U.S. Department of Agriculture, Forest Service, presentation scale 1:3,500,000; colored

Coleman, M.D., T.M. Hinckley, G. McNaughton and B.A. Smit. 1992. Root cold hardiness and native distribution of sub-alpine conifers. *Canadian Journal of Forest Research* 22(7): 932-938.

Crane, M.F. 1982. Fire ecology of Rocky Mountain region forest habitat types. Report prepared under contract to USDA Forest Service Region 2.

Despain, D.G. 1973. Vegetation of the Big Horn Mountains in relation to substrate and climate. *Ecological Monographs* 43:329-355.

Despain, D.G. and R.E. Sellers. 1977. Natural fire in Yellowstone National Park. *Western Wildlands*, summer.

Dillon, G.K., D. Knight and C. Meyer. 2003. Historic variability for upland vegetation in the Medicine Bow National Forest. Department of Botany, Univ. of Wyoming: prepared under agreement with the USDA Forest Service Medicine Bow NF 1102-0003-98-043.

Griggs, R.F. 1938. Timberlines in the northern Rocky Mountains. *Ecology* 19(4): 548-564.

Griggs, R.F. 1946. The timberlines of Northern America and their interpretation. *Ecology* 27(4): 275-289.

Hinds, T.E., F.G. Hawksworth and R.W. Davidson. 1965. Beetle-killed Engelmann spruce: Its deterioration in Colorado. *J. For.* 63(7): 536-542.

Jenkins, M.J., C.A. Dicus and E.G. Hebertson. 1998. Post-fire succession and disturbance interactions on an intermountain subalpine spruce-fir forest. Pages 219-229 in: T.L. Pruden and

L.A. Brennan, eds. Proceedings, Symposium: Fire in Ecosystem Management: Shifting the paradigm from suppression to prescription. Tall Timbers Fire Ecology Conference Proceedings, No. 20. Tall Timbers Research Station, Tallahassee, FL.

Jones, G.P. and S.M. Ogle. 2000. Characterization abstracts for vegetation types on the Bighorn, Medicine Bow, and Shoshone National Forests. Laramie. Prepared for USDA Forest Service, Region 2, by George Jones and Steve Ogle, WYNDD, UW, Laramie WY.

Kane, T.L., B.G. Brown and R. Sharman. 1999. A preliminary climatology of upper level turbulence reports. Preprints pages 363-367 in: 8th Conf. on Aviation, Range and Aerospace Meteorology, 10-15 January. Dallas, TX: American Meteorology Society.

KipfmueLLer, K.F. and W.L. Baker. 2000. A fire history of a subalpine forest in southeastern Wyoming, USA. *Journal of Biogeography* 27: 71-85.

KipfmueLLer, K.F. 1997. A fire history of a subalpine forest in southeastern Wyoming. Thesis. University of Wyoming. Laramie, WY.

KipfmueLLer, K.F. and W.L. Baker. 1998a. A comparison of three techniques to date stand-replacing fires in lodgepole pine forests. *Forest Ecology and Management* 104(1998): 171-177.

KipfmueLLer, K.F. and W.L. Baker. 2000. A fire history of a subalpine forest in southeastern Wyoming, USA. *Journal of Biogeography* 27: 71-85.

Knight, D.H. 1987. Ecosystem studies in the subalpine coniferous forests of Wyoming. In: Management of subalpine forests: building on 50 years of research: proceedings of a technical conference. General Technical Report RM-149. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station.

Knight, D.H. 1994. Mountains and Plains, The Ecology of Wyoming Landscapes. New Haven, CT: Yale University Press.

Knight, D.H. and W.A. Reiners. 2000. Natural patterns in southern Rocky Mountain landscapes and their relevance to forest management. Pages 15-30 in: R. Knight, F.W. Smith, W.H. Romme and W.L. Baker, eds. Forest fragmentation in the Southern Rocky Mountains. Boulder, CO: University Press of Colorado.

Knight, D.H., A.D. Anderson, G.T. Baxter, K.L. Diem, M. Parker, P.A. Rechard, P.C. Singleton, J.F. Thilenius, A.L. Ward and R.W. Weeks. 1975. Final report: the Medicine Bow ecology project: the potential sensitivity of various ecosystem components to winter precipitation management in the Medicine Bow Mountains, Wyoming. Prepared for the Division of Atmospheric Water Resources Management, Bureau of Reclamation, USDI, Denver, CO by the Rocky Mountain Forest and Range Experiment Station, USDA Forest Service and the Wyoming Water Resource Research Institute.

Logan, J.A., J.M. Schmid and M.S. Mehl. 1980. A computer program to calculate susceptibility of spruce-fir stands to spruce beetle outbreaks. USDA Forest Service Research Note RM-303. Fort Collins, CO: Rocky Mountain Forest and Range Experiment Station.

Loope, L.L. and G.E. Gruell. 1973. The ecological role of fire in the Jackson Hole area, northwestern Wyoming. *Quaternary Research* 3(3): 425-443.

Mehl, M. 1992. Old-growth descriptions for the major forest cover types in the Rocky Mountain region. In: Old-growth forests in the Southwest and Rocky Mountain regions. Proceedings of a workshop. Gen. Tech. Report RM-213. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station.

Merrill, E.H., T.W. Kohley, M.E. Herdendorf, W.A. Reiners, K.L. Driese, R.W. Marrs and S.A. Anderson. 1996. Wyoming GAP analysis project final report. University of Wyoming Department of Physiology and Department of Botany, Wyoming Cooperative Fish and Wildlife Research Unit and USGS Biological Resources Division. Available: <http://www.sdvc.uwyo.edu/wbn/abstract.html>.

Meyer, C.B. and D.H. Knight. 2001. Historic variability of upland vegetation in the Bighorn National Forest, Wyoming. Draft report, November 30, 2001.

Mielke, J.L. 1950. Rate of Deterioration of beetle-killed Engelmann spruce. *J. For.* 48(12): 882-888.

Moir, W.H. 1992. Ecological concepts in old-growth forest definition. In: Old-growth forests in the Southwest and Rocky Mountain regions. Proceedings of a workshop. Gen. Tech. Report RM-213. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station.

NatureServe. 2006. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA, U.S.A. Data current as of 18 July 2006.

NatureServe. 2007. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. Data current as of 10 February 2007.

Pennanen, J. 2002. Forest age distribution under mixed-severity fire regimes - A simulation-based analysis for middle boreal Fennoscandia. *Silva Fennica: quarterly issues*: 36(1): 213-231.

Pfister, R.D., B.L. Kovalchik, S.F. Arno and R.C. Presby. 1977. Forest habitat types of Montana. General Technical Report INT-34. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station. 174 pp.

Schmid, J.M. and R.H. Frye. 1977. Spruce beetle in the Rockies. General Technical Report RM 49. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station. 38 pp.

Schmid, J.M. and R.C. Beckwith. 1977. The Spruce Beetle. Pest Leaflet 127. USDA Forest Service 7 pp.

Schmid, J.M. and R.H. Frye. 1976. Stand ratings for spruce beetles. Research Note RM-309. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station.

Schmid, J.M. and T.E. Hinds. 1974. Development of spruce-fir stands following spruce beetle outbreaks. Research Paper RM-131. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station.

Schmidt, K.M., J.P. Menakis, C.C. Hardy, W.J. Hann and D.L. Bunnell. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: USDA Forest Service, Rocky Mountain Research Station. 41 pp. + CD.

Sibold, J. 2001. The forest fire regime of an upper montane and subalpine forest, Wild Basin, Rocky Mountain National Park. M.S. Thesis, University of Colorado, Boulder, CO.

Stahelin, R. 1943. Factors influencing the natural restocking of high altitude gurns by coniferous trees in the central Rocky Mountains. Ecology 24: 19-30.

Veblen, T.T., K.S. Hadley and M.S. Reid. 1991. Disturbance and stand development of a Colorado subalpine forest. Journal of Biogeography 18: 707-716.

Veblen, T.T., K.S. Hadley, E.M. Nel, T. Kitzberger, M.S. Reid and R. Villalba. 1994. Disturbance regime and disturbance interactions in a Rocky Mountain subalpine forest. Journal of Ecology 82: 125-135.

Veblen, T.T., K.S. Hadley, M.S. Reid and A.J. Rebertus. 1989. Blowdown and stand development in a Colorado subalpine forest. Canadian Journal of Forest Resources 19: 1218-1225.

Veblen, T.T. and T. Kitzberger. 2002. Inter-hemispheric comparison of fire history: The Colorado Front Range, U.S.A. and the Northern Patagonian Andes, Argentina. Plant Ecology, in press.