

10760

Chihuahuan Stabilized Coppice Dune and Sand Flat Scrub

BpS Model/Description Version: Aug. 2020

None

Modelers		Reviewers	
John Karges	jkarges@tnc.org	John Morlock	John_Morlock@nps.gov
Lee Elliott	lelliott@tnc.org	None	None
None	None	None	None

Vegetation Type

Shrubland

Map Zone

26

Geographic Range

Chihuahuan Desert dunes and sandsheets.

Biophysical Site Description

Coppice dunes and sandsheets are found in the Chihuahuan Desert. Most moisture occurs during intense, short-duration, late-summer thunderstorms and some winter rains. Another very important process driving this system is wind. This type may also occur in sandy basins.

Vegetation Description

Dominated by mesquite (*Prosopis glandulosa*) but includes fourwing saltbush (*Atriplex canescens*), Torrey's jointfir (*Ephedra torreyana*), Mexican tea (*E. trifurca*), rosemary mint (*Poliomintha incana*), and littleleaf sumac (*Rhus microphylla*) coppice sand scrub with 10-30% total vegetation cover. Soap tree yucca (*Yucca elata*), broom snakeweed (*Gutierrezia sarothrae*), and mesa dropseed (*Sporobolus flexuosus*) are commonly present. Mesquites are scattered, of various ages, and characterized by occurring on pedestaled soil with interstices wind-scoured.

BpS Dominant and Indicator Species

Symbol	Scientific Name	Common Name
PRGL2	<i>Prosopis glandulosa</i>	Honey mesquite
ATCA2	<i>Atriplex canescens</i>	Fourwing saltbush
EPTO	<i>Ephedra torreyana</i>	Torrey's jointfir
EPTR	<i>Ephedra trifurca</i>	Longleaf jointfir
POIN3	<i>Poliomintha incana</i>	Frosted mint

Species names are from the NRCS PLANTS database. Check species codes at <http://plants.usda.gov>.

Disturbance Description

Eolian movement of sparsely vegetated areas between coppice dunes. Fire plays little to no role in this Biophysical Setting (BpS)

Fire Frequency

Severity	Avg FI	Percent of All Fires	Min FI	Max FI
Replacement				
Moderate (Mixed)				

Low (Surface)				
All Fires				

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is the central tendency modeled. Percent of all fires is the percent of all fires modeled in that severity class. Minimum and Maximum FIs show the relative range of fire intervals as estimated by model contributors, if known.

Scale Description

Occurred in small patches, historically.

Adjacency or Identification Concerns

Issues or Problems

Heavy grazing in late 1800s and early 1900s may have caused mesquite to increase. Naturally occurring coppice dunes may have been limited to areas peripheral to active dunes. Coppice dunes are currently more extensive, resulting from sand movement due to degradation of grasslands.

Native Uncharacteristic Conditions

Comments

Succession Classes

Mapping Rules

Upper Layer Lifeform	Height (m)	Canopy Cover (%)									
		0-10	11-20	21-30	31-40	41-50	51-60	61-70	71-80	81-90	91-100
Herb	0-0.5	A	A	A	A	A	A	A	A	A	A
Herb	0.5-1.0	A	A	A	A	A	A	A	A	A	A
Herb	>1.0	A	A	A	A	A	A	A	A	A	A
Shrub	0-0.5	A	A	A	UN	UN	UN	UN	UN	UN	UN
Shrub	0.5-1.0	A	A	A	UN	UN	UN	UN	UN	UN	UN
Shrub	1.0-3.0	A	A	A	UN	UN	UN	UN	UN	UN	UN
Shrub	>3.0	A	A	A	UN	UN	UN	UN	UN	UN	UN
Tree	0-5	A	A	A	UN	UN	UN	UN	UN	UN	UN
Tree	5-10	UN	UN	UN	UN	UN	UN	UN	UN	UN	UN
Tree	10-25	UN	UN	UN	UN	UN	UN	UN	UN	UN	UN
Tree	25-50	UN	UN	UN	UN	UN	UN	UN	UN	UN	UN
Tree	>50	UN	UN	UN	UN	UN	UN	UN	UN	UN	UN

Succession class letters A-E are described in the Succession Class Description section. Some classes use a leafform distinction where a qualifier is added to the class letter: Brdl (broadleaf), Con (conifer), or Mix (mixed conifer and broadleaf). UN refers to uncharacteristic native or a combination of height and cover that would not be expected under the reference condition. NP refers to not possible or a combination of height and cover which is not physiologically possible for the species in the BpS.

Description

Class A	100	Mid Development 1 - Open
---------	-----	--------------------------

Indicator Species

Symbol	Scientific Name	Common Name	Canopy Position
PRGL2	Prosopis glandulosa	Honey mesquite	All
ATCA2	Atriplex canescens	Fourwing saltbush	None
POIN3	Poliomintha incana	Frosted mint	None

Description

This class occurs as a mosaic of open, unvegetated sand substrate and dunes vegetated with scattered multi-branched, recumbent mesquites. Significant amounts of bare soil and dune conditions result in low percent canopy closure. Wind movement of this BpS maintains the open aspect. Fire plays little to no role.

Maximum Tree Size Class

None

Model Parameters

Deterministic Transitions

From Class	Begins at (yr)	Succeeds to	After (years)
Mid1:OPN	0	Mid1:OPN	999

Probabilistic Transitions

Disturbance Type	Disturbance occurs In	Moves vegetation to	Disturbance Probability	Return Interval (yrs)	Reset Age to New Class Start Age After Disturbance?	Years Since Last Disturbance
Wind or Weather or Stress	Mid1:OPN	Mid1:OPN	1	1	No	0

References

Bowers, J.E. 1982. The plant ecology of inland dunes in western North America. Journal of Arid Environments 5: 199-220.

Bowers, J.E. 1984. Plant geography of southwestern sand dunes. Desert Plants 6(1):31-42, 51-54.

Campbell, R.S. 1929. Vegetation succession in the Prosopis sand dunes of southern New Mexico. Ecology 10: 392-398.

Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, K. Snow and J. Teague. 2003. Ecological systems of the United States: A working classification of U.S. terrestrial systems. NatureServe, Arlington, VA.

Dick-Peddie, W.A. 1993. New Mexico vegetation: Past, present, and future. University of New Mexico Press, Albuquerque. 244 pp.

Muldavin, E., Y. Chauvin and G. Harper. 2000b. Vegetation of White Sands Missile Range, New Mexico: Volume I Handbook of vegetation communities. Final Report to White Sands Missile Range by New Mexico Natural Heritage Program, University of New Mexico, NM. 192 pp.

NatureServe. 2007. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. Data current as of 10 February 2007.

NatureServe. 2006. NatureServe Explorer: An online encyclopedia of life [web application]. Version 5.0. NatureServe, Arlington, Virginia. Available <http://www.natureserve.org/explorer>. (Accessed: August 21, 2006).

Schussman, H. and D. Gori. 2004. An ecological assessment of the Bureau of Land Management's current fire management plans: Materials and recommendations for future fire planning. The Nature Conservancy, Arizona Chapter.

Steinberg, P. 2001. *Prosopis glandulosa*. in Fire Effects Information System, [Online]. USDA Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis/> [2006, August 21].