

11080

Sonora-Mojave Semi-Desert Chaparral

BpS Model/Description Version: Aug. 2020

Modelers		Reviewers	
Matt Brooks	matt_brooks@usgs.gov	Jan Beyers	jbeyers@fs.fed.us
None	None	Richard Halsey	richardhalsey@sbcglobal.net
None	None	Hugh Safford/Dave Schmidt	hughsafford@fs.fed.us

Vegetation Type

Shrubland

Map Zones

4, 5, 13, 14, 15, 24, 25

Geographic Range

This ecological system is composed of evergreen shrublands on sideslopes transitioning from low-elevation desert landscapes up into woodlands of the western Mojave and Sonoran Deserts. It extends from northeast Kern County, CA, into Baja California Norte. This vegetation type is also found on the northern slopes of the Transverse Ranges and eastern slopes of Peninsular and coast ranges.

Biophysical Site Description

This system includes chaparral on sideslopes transitioning from low-elevation desert landscapes up into pinyon-juniper and ponderosa pine woodlands of the western and central Great Basin, between 4000-7000ft (including north and east slopes of Transverse and Peninsular ranges in southern California).

Vegetation Description

Associated species include *Quercus john-tuckeri*, *Quercus cornelius-mulleri*, *Quercus berberidifolia*, *Arctostaphylos patula*, *Arctostaphylos pungens*, *Arctostaphylos glauca*, *Rhus ovata*, *Juniperus californica*, *Cercocarpus montanus* var. *glaber* (= *Cercocarpus betuloides*), *Ceanothus greggii*, *Garrya flavescens* and *Nolina parryi*. Stands tend to be on steep slopes embedded in larger stands of blackbrush, pinyon, juniper, and sagebrush.

BpS Dominant and Indicator Species

Symbol	Scientific Name	Common Name
QUJO3	<i>Quercus john-tuckeri</i>	Tucker oak
QUCO7	<i>Quercus cornelius-mulleri</i>	Muller oak
QUBE5	<i>Quercus berberidifolia</i>	Scrub oak
ARPA6	<i>Arctostaphylos patula</i>	Greenleaf manzanita
ARPU5	<i>Arctostaphylos pungens</i>	Pointleaf manzanita
ARGL4	<i>Arctostaphylos glauca</i>	Bigberry manzanita
CEMOG	<i>Cercocarpus montanus</i> var. <i>glaber</i>	Birchleaf mountain mahogany

GAFL2	<i>Garrya flavescens</i>	Ashy silktassel
-------	--------------------------	-----------------

Species names are from the NRCS PLANTS database. Check species codes at <http://plants.usda.gov>.

Disturbance Description

Typical fire regime in these systems varies with the amount of organic accumulation and is largely related to fire regimes of surrounding vegetation. The only significant disturbance to the system is stand-replacing fire occurring every 100+ yrs on average. Shrubs resprout rapidly after fire or regenerate from soil-stored seeds.

Fire Frequency

Severity	Avg FI	Percent of All Fires	Min FI	Max FI
Replacement	81	100	75	100
Moderate (Mixed)				
Low (Surface)				
All Fires	81	100		

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is the central tendency modeled. Percent of all fires is the percent of all fires modeled in that severity class. Minimum and Maximum FIs show the relative range of fire intervals as estimated by model contributors, if known.

Scale Description

Vegetation found in small patches of 10s-100s of acres.

Adjacency or Identification Concerns

BPS 1108 will be difficult to distinguish from BPS 1103 (Great Basin Semi-Desert Chaparral) and BPS 1104 (Mogollon Semi-Desert Chaparral). Smaller stand size than BpS 1104 and longer fire return intervals.

Issues or Problems

Comments

Succession Classes

Mapping Rules

Upper Layer Lifeform	Height (m)	Canopy Cover (%)									
		0-10	11-20	21-30	31-40	41-50	51-60	61-70	71-80	81-90	91-100
Herb	0-0.5	A	A	A	A	A	A	A	A	A	A
Herb	0.5-1.0	A	A	A	A	A	A	A	A	A	A
Herb	>1.0	A	A	A	A	A	A	A	A	A	A
Shrub	0-0.5	A	A	A	A	A	A	A	A	A	A
Shrub	0.5-1.0	A	A	A	A	A	A	A	A	A	A
Shrub	1.0-3.0	B	B	B	B	B	B	B	B	B	B
Shrub	>3.0	B	B	B	B	B	B	B	B	B	B
Tree	0-5	B	B	B	B	B	UN	UN	UN	UN	UN
Tree	5-10	B	B	B	B	B	UN	UN	UN	UN	UN
Tree	10-25	B	B	B	B	B	UN	UN	UN	UN	UN
Tree	25-50	B	B	B	B	B	UN	UN	UN	UN	UN
Tree	>50	B	B	B	B	B	UN	UN	UN	UN	UN

Succession class letters A-E are described in the Succession Class Description section. Some classes use a leafform distinction where a qualifier is added to the class letter: Brdl (broadleaf), Con (conifer), or Mix (mixed conifer and broadleaf). UN refers to uncharacteristic native or a combination of height and cover that would not be expected

under the reference condition. NP refers to not possible or a combination of height and cover which is not physiologically possible for the species in the BpS.

Description

Class A 21 Early Development 1 - All Structures

Indicator Species

Symbol	Scientific Name	Common Name	Canopy Position
ARPA6	Arctostaphylos patula	Greenleaf manzanita	Upper
ARPU5	Arctostaphylos pungens	Pointleaf manzanita	Upper
QUJO3	Quercus john-tuckeri	Tucker oak	Upper
CEMOG	Cercocarpus montanus var. glaber	Birchleaf mountain mahogany	Upper

Description

After fire, shrubs resprout strongly from roots or from the base of plants; a few species regenerate from soil-stored seed. Shrubs can cause stands to become impenetrable.

Maximum Tree Size Class

None

Class B 79 Late Development 1 - All Structures

Indicator Species

Symbol	Scientific Name	Common Name	Canopy Position
ARPA6	Arctostaphylos patula	Greenleaf manzanita	Upper
ARPU5	Arctostaphylos pungens	Pointleaf manzanita	Upper
QUJO3	Quercus john-tuckeri	Tucker oak	Upper
CEMOG	Cercocarpus montanus var. glaber	Birchleaf mountain mahogany	Upper

Description

Dense shrubs with grasses present in the few openings.

Maximum Tree Size Class

None

Model Parameters

Deterministic Transitions

From Class	Begins at (yr)	Succeeds to	After (years)
Early1:ALL	1	Late1:ALL	20
Late1:ALL	21	Late1:ALL	999

Probabilistic Transitions

Disturbance Type	Disturbance occurs In	Moves vegetation to	Disturbance Probability	Return Interval (yrs)	Reset Age to New Class Start Age After Disturbance?	Years Since Last Disturbance
Replacement Fire	Early1:ALL	Early1:ALL	0.01	100	Yes	0
Replacement Fire	Late1:ALL	Early1:ALL	0.013	77	Yes	0

References

Barbour, M.G. and J. Major, eds. 1988. Terrestrial vegetation of California: New expanded edition. Special Publication 9. Sacramento, CA: California Native Plant Society. 1030 pp.

Brooks, M.L. and J.R. Matchett. 2003. Plant community patterns in unburned and burned blackbrush (*Coleogyne ramosissima* Torr.) shrublands in the Mojave Desert. *Western North American Naturalist* 63(3): 283-298.

Brooks, M.L, T.C. Esque and T. Duck, 2003. Fuels and fire regimes in creosotebush, blackbrush, and Interior chaparral shrublands. Report for the Southern Utah Demonstration Fuels Project. MT: USDA Forest Service. Rocky Mountain Research station. 18 pp.

Brooks, M.L., C.M. D'Antonio, D.M. Richardson, J.B. Grace, J.E. Keeley, J.M. DiTomaso, R.J. Hobbs, M. Pellant and D. Pyke. 2004. Effects of invasives alien plants on fire regimes. *BioSciences* 54: 677-688.

Brown, D.E., ed. 1982. Biotic communities of the American Southwest-United States and Mexico. *Desert Plants Special Issue* 4(1-4): 1-342.

Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, K. Snow and J. Teague. 2003. Ecological systems of the United States: A working classification of U.S. terrestrial systems. Arlington, VA: NatureServe.

Holland, V.L. and D.J. Keil. 1995. California vegetation. Dubuque, IA: Kendall/Hunt Publishing Company. 516 pp.

MacMahon, J.A. 1988. Warm deserts. In: Barbour, M.G. and W. D. Billings, eds. *North American terrestrial vegetation*. NY: Cambridge University Press. 232-264

NatureServe. 2007. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. Data current as of 10 February 2007.

Thomas, K., J. Franklin, T. Keeler-Wolf and P. Stine. 2003. Mojave Desert Ecosystem Program: Central Mojave Vegetation Mapping Project. USGS Open-file Report. In press.

