

11160

Madrean Juniper Savanna

BpS Model/Description Version: Aug. 2020

Modelers		Reviewers	
Mike Babler	mbabler@tnc.org	Dave Anderson	danderson@wsmr.army.mil
Phil Smith	phil_smith@blm.gov	Tim Christiansen	tchristiansen@tnc.org
Steve Bumgarner	steve_bumgarner@blm.gov	None	None
		Keith Schulz	Keith_Schulz@natureserve.org
		Reese Lolley	rlolley@fs.fed.us

Reviewed by: Tim Christiansen

Vegetation Type

Steppe/Savanna

Map Zones

15, 25

Model Splits or Lumps

This Biophysical Setting (BpS) is lumped with: 2510250 and 1510250.

Geographic Range

Sierra Madre Occidentale and Sierra Madre Oriental in Mexico, Trans-Pecos Texas, southern New Mexico, southern Arizona (south of the Mogollan Rim), and southwestern New Mexico.

Biophysical Site Description

This BpS typically is found at elevations between 1,400-2,200m. Madrean juniper savannas occur at the lower altitudinal limits for foothill tree species, below the pinyon-juniper (PJ) woodlands, but at or above semi-desert grassland where soil moisture limits cover of montane woody plants. At higher and therefore moister elevations, woodlands grade into Madrean encinal or, less frequently, montane pine-oak woodlands characterized by taller and denser vegetation than these woodlands. Savannas and woodlands are found on many and varied topographic positions, including low- to mid-elevation mountain slopes, hills, plateaus, basins, and flats.

Vegetation Description

The upper vegetation canopy is composed of open to moderately dense tree layer dominated by *Pinus cembroides*, *Pinus discolor*, *Pinus edulis*, *Juniperus depeanna*, and/or *Juniperus monosperma*. Madrean oaks such as *Quercus arizonica*, *Q. emoryi*, *Q. grisia*, *Q. oblongifolia*, or *Q. mohriana* may be present to co-dominant with pinyon and/or juniper trees. In southwest New Mexico, *Q. grisia* may be dominant. *Pinus engelmannii* occurs infrequently, and *P. ponderosa* and *P. arizonica* are absent. *Juniperus monosperma* is often present to dominant on the Gila National Forest. Understory layers are variable and may be dominated by shrubs such as manzanita (*Arctostaphylos pungens*, *A. pringlei*), cliffrose (*Cowania mexicana*), Apache plume (*Fallugia paradoxa*), or barberry (*Berberis* spp.). Graminoids may form dense (savanna) to sparse canopy. Common species include sideoats grama (*Bouteloua curtipendula*), cane bluestem (*Bothriochloa barbinodis*), muhly grasses (*Muhlenbergia emersleyi*, *M. torreyi*, *M. porteri*). Graminoids decrease in cover and biomass with increasing cover of woody plants.

BpS Dominant and Indicator Species

Symbol	Scientific Name	Common Name
JUDE	<i>Juncus debilis</i>	Weak rush
PIED	<i>Pinus edulis</i>	Twoneedle pinyon
PIDI3	<i>Pinus discolor</i>	Border pinyon
QUAR	<i>Quercus arizonica</i>	Arizona white oak
QUEM	<i>Quercus emoryi</i>	Emory oak
QUGR3	<i>Quercus grisea</i>	Gray oak
QUOB	<i>Quercus oblongifolia</i>	Mexican blue oak
PILE	<i>Pinus leiophylla</i>	Chihuahuan pine

Species names are from the NRCS PLANTS database. Check species codes at <http://plants.usda.gov>.

Disturbance Description

The fire regime of this ecological system is almost completely unknown. There are essentially no data about fire frequency, fire history, or fire behavior. The occasional paper that addresses fire includes considerably more speculation than data. It would seem that fire occurrence was determined primarily by fire occurrence in the surrounding matrix vegetation and was ignited by lightning during early summer. However, even this information is speculative and undocumented; based on contemporary ecological knowledge, models that assume specific fire regimes are little more than wild guesses. Such a model follows, in the sincere hope that it will be ignored or improved upon.

This system is likely predisposed to stand-replacement fires during the earliest stage of stand development. Replacement fires are assumed to have occurred every century or so and mixed-severity fires slightly less frequently. Significant drought occurs about every 60yrs and, in combination with herbivory from invertebrates, causes disproportional mortality of large, old trees.

Fire Frequency

Severity	Avg FI	Percent of All Fires	Min FI	Max FI
Replacement	136	34	20	1000
Moderate (Mixed)	208	22	20	1000
Low (Surface)	107	44		
All Fires	46	100		

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is the central tendency modeled. Percent of all fires is the percent of all fires modeled in that severity class. Minimum and Maximum FIs show the relative range of fire intervals as estimated by model contributors, if known.

Scale Description

Pinyon-juniper woodland usually was distributed across the landscape in patches between 100s-1,000s of acres in size. In particularly dissected topography, this type may have occurred in smaller patches.

Adjacency or Identification Concerns

This system generally is found at higher elevations and more mesic sites than semi-desert grassland. Typically, it is bordered at higher elevations by Madrean Pine-Oak Woodlands.

Cover and density of juniper and pinyon trees at lower elevations in this type doubtless have increased as a result of fire suppression (possibly as mitigated by livestock grazing). This phenomenon is characteristic of BpS 1116 (juniper savanna), with which BpS 1025 (pinyon-juniper woodland) has been lumped for map zone (MZ)15. Bordered at low elevation by Madrean Encinal and Desert Grasslands on Gila NF.

Issues or Problems

Virtually no components of the fire regimes are known with any certainty. Fire scars are rare, and trees in this system cannot be aged with conventional dendrochronological techniques. Information about fire regimes is extrapolated from adjacent systems, and extreme caution is warranted when interpreting these models. Fire season can be inferred more reliably than fire frequency; the former likely is equally or more important than the latter.

This ecological system was originally described assuming evergreen oak species are present to co-dominant in the understory or canopy in MZ15 (Schulz, personal communication). This system also includes pinyon-juniper woodlands and savannas with understories dominated by other shrubs or a grass layer and lacks evergreen oaks, which may have a different fire behavior.

Native Uncharacteristic Conditions

Comments

Lower moisture levels due to drought may cause this BpS to decline in area since it occurs in a mesic habitat. Fire behavior and fire intervals will increase because of loss of understory fuels.

Succession Classes

Mapping Rules

Upper Layer Lifeform	Height (m)	Canopy Cover (%)									
		0-10	11-20	21-30	31-40	41-50	51-60	61-70	71-80	81-90	91-100
Herb	0-0.5	A	A	A	A	A	A	A	A	A	A
Herb	0.5-1.0	A	A	A	A	A	A	A	A	A	A
Herb	>1.0	A	A	A	A	A	A	A	A	A	A
Shrub	0-0.5	A	A	A	A	A	A	A	A	A	A
Shrub	0.5-1.0	A	A	A	A	A	A	A	A	A	A
Shrub	1.0-3.0	A	A	A	A	A	A	A	A	A	A
Shrub	>3.0	A	A	A	A	A	A	A	A	A	A
Tree	0-5	B	B	B	B	B	B	B	B	B	B
Tree	5-10	B	B	B	B	B	B	B	B	B	B
Tree	10-25	C	C	C	C	C	C	C	C	C	C
Tree	25-50	C	C	C	C	C	C	C	C	C	C
Tree	>50	C	C	C	C	C	C	C	C	C	C

Succession class letters A-E are described in the Succession Class Description section. Some classes use a leafform distinction where a qualifier is added to the class letter: Brdl (broadleaf), Con (conifer), or Mix (mixed conifer and broadleaf). UN refers to uncharacteristic native or a combination of height and cover that would not be expected under the reference condition. NP refers to not possible or a combination of height and cover which is not physiologically possible for the species in the BpS.

Description

Class A	5	Early Development 1 - All Structures
----------------	----------	---

Indicator Species

Symbol	Scientific Name	Common Name	Canopy Position
BOCU	Bouteloua curtipendula	Sideoats grama	Upper
BOBA3	Bothriochloa barbinodis	Cane bluestem	Upper
QUAR	Quercus arizonica	Arizona white oak	Upper
QUEM	Quercus emoryi	Emory oak	Upper

Description

Initial post-fire community dominated by perennial caespitose grasses. Evidence of past fires may be observed, including charcoal and resprouting woody plants.

Maximum Tree Size Class

None

Class B	57	Mid Development 1 - Closed
----------------	-----------	-----------------------------------

Indicator Species

Symbol	Scientific Name	Common Name	Canopy Position
JUDE	Juncus debilis	Weak rush	Upper
PIED	Pinus edulis	Twoneedle pinyon	Upper
QUAR	Quercus arizonica	Arizona white oak	Upper
QUEM	Quercus emoryi	Emory oak	Upper

Description

Community dominated by young to mature alligator juniper and evergreen oak trees of various ages.

Maximum Tree Size Class

Pole 5-9" DBH

Class C	38	Late Development 1 - Closed
----------------	-----------	------------------------------------

Indicator Species

Symbol	Scientific Name	Common Name	Canopy Position
JUDE	Juncus debilis	Weak rush	Upper
PIED	Pinus edulis	Twoneedle pinyon	Upper
QUAR	Quercus arizonica	Arizona white oak	Upper
QUEM	Quercus emoryi	Emory oak	Upper

Description

Site dominated by relatively dense old alligator juniper and evergreen oak trees.

Maximum Tree Size Class

Medium 9-21" DBH

Model Parameters

Deterministic Transitions

From Class	Begins at (yr)	Succeeds to	After (years)
Early1:ALL	0	Mid1:CLS	9
Mid1:CLS	10	Late1:CLS	84
Late1:CLS	85	Late1:CLS	984

Probabilistic Transitions

Disturbance Type	Disturbance occurs In	Moves vegetation to	Disturbance Probability	Return Interval (yrs)	Reset Age to New Class Start Age After Disturbance?	Years Since Last Disturbance
Replacement Fire	Early1:ALL	Early1:ALL	0.1	10	Yes	0
Replacement Fire	Mid1:CLS	Early1:ALL	0.003	333	Yes	0
Mixed Fire	Mid1:CLS	Mid1:CLS	0.005	200	No	0
Replacement Fire	Late1:CLS	Early1:ALL	0.003	333	Yes	0
Mixed Fire	Late1:CLS	Late1:CLS	0.005	200	No	0
Wind or Weather or Stress	Late1:CLS	Mid1:CLS	0.015	67	Yes	0
Surface Fire	Late1:CLS	Late1:CLS	0.025	40	No	0

References

Alexander, R.R., F. Ronco, Jr. 1987. Classification of the forest vegetation on the National Forests of Arizona and New Mexico. Res. Note RM-469. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station. 10 pp.

Anderson, H.E. 1982. Aids to Determining Fuel Models For Estimating Fire Behavior. Gen. Tech. Rep. INT-122. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station. 22 pp.

Arno, S.F. 2000. Fire in western forest ecosystems. Pages 97-120 in: J.K. Brown and J. Kapler-Smith, eds. Wildland fire in ecosystems: Effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station.

Baker, W.L. and D.J. Shinneman. 2004. Fire and restoration of piñon-juniper woodlands in the western United States. A review. Forest Ecology and Management 189: 1-21.

Barton, A.M. 2002. Intense wildfire in southeastern Arizona: transformation of a Madrean oak-pine forest to oak woodland. *Forest Ecology and Management* 165: 205-212.

Barton, A.M. 1999. Pines versus oaks: effects of fire on the composition of Madrean forests in Arizona. *Forest Ecology and Management* 120: 143-156.

DeBano, L.F., P.F. Ffolliott, A. Ortega-Rubio, G.J. Gottfried, R.H. Hamre and C.B. Edminster, technical coordinators. 1995. Biodiversity and Management of the Madrean Archipelago: The Sky Islands of the Southwestern United States and Northern Mexico. USDA Forest Service Rocky Mountain Experiment Station General Technical Report RM-264.

Dick-Peddie, W.A. 1993. New Mexico vegetation: Past, present, and future. Albuquerque, NM: University of New Mexico Press. 244 pp.

Ffolliott, P.F. et al., technical coordinators. 1996. Effects of Fire on Madrean Province Ecosystems: A Symposium Proceedings. USDA Forest Service General Technical Report RM-GTR-289. 277 pp.

Gruell, G.E. Historical and Modern Roles of Fire in Pinyon-Juniper. Pages 24-28 in: Proceedings, USDA Forest Service RMRS-P-9.

Hardy, C.C., K.M. Schmidt, J.P. Menakis and R.N. Samson. 2001. Spatial data for national fire planning and fuel management. *Int. J. Wildland Fire*. 10(3&4): 353-372.

Haworth, K. and G.R. McPherson. 1994. Effects of *Quercus emoryi* on herbaceous vegetation in a semi-arid savanna. *Vegetatio* 112: 153-159.

Kuchler, A.W. 1964. Potential Natural Vegetation of the Conterminous United States. American Geographic Society Special Publication No. 36. 116 pp.

McClaran, M.P. and G.R. McPherson. 1999. Oak savanna of the American Southwest. Pages 275-287 in R.C. Anderson, J.S. Fralish and J. Baskin, editors. *Savannas, Barrens, and Rock Outcrop Plant Communities of North America*. Cambridge University Press, Cambridge, England.

McPherson, G.R. and J.F. Weltzin. 1998. Response of understory to overstory removal in southwestern oak woodlands. *Journal of Range Management* 51: 674-678.

McPherson, G.R. and J.F. Weltzin. 2000. The role and importance of disturbance and climate change in U.S./Mexico borderlands: a state-of-the-knowledge review. USDA Forest Service Rocky Mountain Research Station General Technical Report RMRS-GTR-50.

NatureServe. 2007. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. Data current as of 10 February 2007.

Romme, W.H., L. Floyd-Hanna and D. Hanna. 2002. Ancient Pinyon-Juniper forests of Mesa Verde and the West: A cautionary note for forest restoration programs. In: Conference Proceedings – Fire, Fuel Treatments, and Ecological Restoration: Proper Place, Appropriate Time, Fort Collins, CO, April 2002. 19 pp.

Schmidt, K.M., J.P. Menakis, C.C. Hardy, W.J. Hann and D.L. Bunnell. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: USDA Forest Service, Rocky Mountain Research Station. 41 pp. + CD.

Tausch, R.J. and N.E. West. 1987. Differential Establishment of Pinyon and Juniper Following Fire. *The American Midland Naturalist* 119(1). 174-184.

USDA Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (2002, December). Fire Effects Information System, [Online]. Available: <http://www.fs.fed.us/database/feis/> [Accessed: 11/15/04].

Webster, G.L. and C.J. Bahre, editors. 2001. *Changing Plant Life of La Frontera: Observations on Vegetation in the United States/Mexico Borderlands*. University of New Mexico Press, Albuquerque. 260 pp.

West, N.E. 1999. Juniper pinon savanna and woodlands of Western North America. Pages 288-308 in R.C. Anderson, J.S. Fralish, and J. Baskin, editors. *Savannas, Barrens, and Rock Outcrop Plant Communities of North America*. Cambridge University Press, Cambridge England.

Wright, H.A., L.F. Neuenschwander and C.M. Britton. 1979. The role and use of fire in Sagebrush-Grass and Pinyon-Juniper Plant Communities. Gen. Tech. Rep. INT-GTR-58. Ogden, UT: USDA Forest Service, Intermountain Research Station. 48 pp.