

LANDFIRE Biophysical Setting Model

Biophysical Setting 2911480

Western Great Plains Sand Prairie

- ☐ This BPS is lumped with:
☐ This BPS is split into multiple models:

General Information

Contributors (also see the Comments field)

Date 4/6/2006

Modeler 1 Peter Lesica	peter.lesica@mso.umn.edu	Reviewer Steve Cooper	scooper@mt.gov
Modeler 2 Vinita Shea	vinita_shea@blm.gov	Reviewer Jim Von Loh	jvonloh@e2m.net
Modeler 3 Ben Pratt	ben_pratt@fws.gov	Reviewer	

Vegetation Type

Upland
Grassland/Herbaceous

Dominant Species

CALO
SCSC

Map Zone

29

Model Zone

- | | |
|--------------------------------------|--|
| ☐ Alaska | ☐ Northern Plains |
| ☐ California | ☐ N-Cent. Rockies |
| ☐ | ☒ |
| ☐ Great Basin | ☐ Pacific Northwest |
| ☐ Great Lakes | ☐ South Central |
| ☐ Hawaii | ☐ Southeast |
| ☐ Northeast | ☐ S. Appalachians |
| | ☐ Southwest |

General Model Sources

- | | |
|---|--------|
| ☒ Literature | BOGR2 |
| ☒ Local Data | HECO26 |
| ☒ Expert Estimate | SPCR |

Geographic Range

Predominantly the eastern portion of MZ20. Also found in scattered pockets elsewhere throughout the zone. It probably occurs on the Charles Russel National Wildlife Refuge. In 331Kf, this might occur.

In MZs 29 and 30, more of this type than in MZ20 because more sandstone and sandy soils. Mostly only in MZ30. About 30K acres (guesstimate) maybe found in the southern portion of MZ29. Occurs around Broadus and Ekalaka. Medicine Rocks State Park almost all sand prairie. 331Kf, in 331Gf and d. Occurs in Little Missouri Grasslands in Dakotas.

There are good classification data and local descriptions of this type for Theodore Roosevelt National Park near Medora, ND as part of the USGS-NPS Vegetation Mapping Program (Jim Von Loh, pers comm).

Biophysical Site Description

This BpS would be found in NRCS's sand type or the Sandy Ecological site description. Occurs around sandstone outcrops.

Lower productivity on these sandy sites versus the mixedgrass prairie sites.

Vegetation Description

Dominant vegetation includes prairie sandreed (*Calamovilfa longifolia*), little bluestem (*Schizachyrium scoparium*), blue grama (*Bouteloua gracilis*), needle-and-thread grass (*Stipa comata*) and sand dropseed (*Sporobolus cryptandrus*). Shrubs seen may include horizontal juniper (*Juniperus horizontalis*), silver sagebrush (*Artemisia cana*) and skunkbrush (*Rhus trilobata*). Further east (not in MZ20), BOHI2 and

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

ANHA might occur.

It's uncommon to find Wyoming big sagebrush, and when you do - it's usually Wyoming big sagebrush with bluebunch wheatgrass or needle-and-thread grass, that you would find on a sandy soil. The sagebrush in this type is usually silver sagebrush. It would be unusual to have more than 10-15% shrub cover except in the case of *Juniperus horizontalis*, where cover can go up to 80% or more.

Disturbance Description

Fire, grazing and drought were the primary disturbances. Disturbances were cyclic with the earliest and latest seral stages fluctuating widely in accordance with changes in climate.

The principal large grazer of the system was most likely bison (*Bison bison*) which, when occurring in large numbers, would have locally disturbed large areas due both to grazing impact and physical disturbances such as trampling and wallowing. Grazing impacts are more pronounced near water and removed from steep, rough terrain. Overall the whole system would have been frequently impacted by large ungulate grazers.

Prairie dogs might have been a very minor component of the system. Where they occurred, prairie dogs grazed vegetation close to the ground which provided a local firebreak. It is questionable, however, as to whether prairie dogs prefer sandy soil and actually occurred here. It is thought that prairie dogs would not occur on these sandy sites and rather they usually occur on fine textured soils.

Fire was a frequent and widespread occurrence. The most extensive fires are likely to have occurred in years with wet springs followed by hot, dry summers when grazing pressure was low. Wet springs would have resulted in more productive and more continuous plant cover (ie, fuel) that would have supported and expanded fires ignited under dry conditions occurring later in the season. In addition, litter accumulation over several fire-free years would also have supported widespread fire, in any conditions. The litter component, a determining factor in fire size and frequency, is correlated with seral stage. Three to five fire-free years produce enough litter to carry another fire. Post-fire shifts in species composition depend on the timing and condition of fire. It is also speculated that native burning might have been an influence in this BpS.

Fire regime similar to adjacent grassland.

Extended periods of severe drought is likely to have affected both species composition and the stability of the sandy soil, particularly when compounded by wind and heavy grazing. Droughts could affect the entire region.

Adjacency or Identification Concerns

Northwestern Plains Mixed Grass Prairie systems are often found nearby, especially in the western portion of the zone. The sand prairie, however, occurs on the sandy sites. You should identify this system by sandy/soil types.

Pine savanna is sometimes at top of these sandy sites. Trees on northerly or easterly slopes, might be looking at sandy outcrop. Portion of upper slope might be associated with sand prairie. Top will be mapped to pine savanna or woodland.

The disturbance regime has been drastically changed since European settlement. Agriculture replaced

****Fire Regime Groups are:** I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

bison and fires have been effectively suppressed.

Bromus much less prevalent on sandy soils than on mixedgrass prairie, but does occur. There aren't any invasives for this system that are particularly an issue.

It is uncommon to find Wyoming big sagebrush, and when you do - it's usually Wyoming big sagebrush with bluebunch wheatgrass or needle-and-thread, that you would find on a sandy soil. The sagebrush in this type is usually silver sagebrush. It would be unusual to have more than 10-15% shrub cover except in the case of Juniperus horizontalis, where cover can go up to 80% or more.

This system is much less departed than 1141. This system is probably not very far departed from the HRV, if at all.

Native Uncharacteristic Conditions

Over 60% cover herbaceous would be uncharacteristic... probably wouldn't be a sandy site anymore. Herb height also wouldn't be over one meter.

Scale Description

This is generally a patch that occurs within the larger northwestern Great Plains mixed grass. Size probably varies widely, but is generally going to be 10s of thousands of acres in MZ29 at the large side versus hundreds of acres or less in MZ20.

In terms of disturbance impact, in MZ20, entire patches are going to be impacted, whereas it will vary to an unknown level in MZ29.

Issues/Problems

Very little data are available from presettlement times.

Comments

This model for MZs 29 and 30 was adapted from the same BpS from MZ20 created by Peter Lesica, Vinita Shea, Ben Pratt and reviewed by Steve Barrett and Brian Martin.

This model for MZ20 was adapted from the Rapid Assessment model R4NESP Nebraska Sandhills Prairie created by Tom Bragg (tbragg@mail.unomaha.edu), Mary Lata (mlata@fs.fed.us) and Dave Shadis (dshadis@fs.fed.us) and reviewed by John Ortmann (jortmann@tnc.org). Major descriptive and quantitative changes were made so that the model more appropriately represented MT, instead of NE.

Vegetation Classes											
Class A	20 %	Indicator Species and Canopy Position									
Early Development 1 Open		SPCR									
Upper Layer Lifeform		Upper									
☒ Herbaceous		HECO26									
☐ Shrub		Lower									
☐ Tree	Fuel Model	SCSC									
		Upper									
		CALO									
		Structure Data (for upper layer lifeform)									
		<table><thead><tr><th></th><th>Min</th><th>Max</th></tr></thead><tbody><tr><td>Cover</td><td>0 %</td><td>40 %</td></tr><tr><td>Height</td><td>Herb 0m</td><td>Herb 0.5m</td></tr></tbody></table>		Min	Max	Cover	0 %	40 %	Height	Herb 0m	Herb 0.5m
	Min	Max									
Cover	0 %	40 %									
Height	Herb 0m	Herb 0.5m									
		Tree Size Class									
☐ Upper layer lifeform differs from dominant lifeform.											

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Description

Upper

Class A represents immediate to three year post disturbance conditions. Vegetation consists of resprouting and seedling grass and forbs. Total bare soil is greater than before the disturbance particularly on less productive sites. The vigor of new growth and the specific species affected depend on the season of the disturbance and on pre- and post-disturbance environmental conditions (eg, available soil moisture). Litter is low initially but increases until, by year three, there is enough to support fire under average burning conditions. Fire was therefore modeled as occurring somewhat less frequently than in class B. In uplands, where soil-type is dominated by coarse-grained sands with low water-holding capacity, post-disturbance primary production initially decreases thus fire may only carry under ideal conditions. Under these conditions, grazing is likely to be light. In lowlands, with finer-textured soils, primary production is determined largely by moisture availability.

HECO26 was used as an indicator species for MZs 29 and 30 instead of BOGR2.

ARCA13 can resprout immediately after fire, so it could be present in this stage as well. It could, however, be killed following intensive fires. But since not much litter in these sites, possibility of intense fire reduced.

It was originally suggested that there would be a prairie-dog influenced stage at approximately two percent of the landscape. However, there was some disagreement as to whether this class should exist or not for this system, as it is thought that these sandy sites might have been unlikely to have prairie dog towns. It was only distinguished from A by different species (Buchloe dactyloides - only in the extreme southeast portion of the stage and Bouteloua gracilis, and Agropyron dasystachyum). Canopy cover was 0-20%. This (very unlikely) prairie dog influenced class was therefore merged into the early successional stage, class A. It is doubtful that prairie dogs would colonize very sandy sites; most prairie dog sites have fairly fine-textured soils.

Repeated grazing of these areas will prevent succession to class B. Grazing occurs with a probability of 0.05. Prairie dog grazing was modeled as optional 1, with a very unlikely probability of 0.0007. Both of these will set succession back to the beginning.

Replacement fires occur every 40yrs (due to less litter until the last years of this class, they were modeled as occurring less frequently than in class B. (20yr vs 40yr FRI does not change percentages in each class.)

Class B 80 %		<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>												
Late Development 1 Closed		CALO	<table><tr><td></td><td>Min</td><td>Max</td></tr><tr><td>Cover</td><td>41 %</td><td>60 %</td></tr><tr><td>Height</td><td>Herb 0m</td><td>Herb 1.0m</td></tr><tr><td>Tree Size Class</td><td colspan="2"></td></tr></table>		Min	Max	Cover	41 %	60 %	Height	Herb 0m	Herb 1.0m	Tree Size Class		
	Min	Max													
Cover	41 %	60 %													
Height	Herb 0m	Herb 1.0m													
Tree Size Class															
<u>Upper Layer Lifeform</u>		Upper	☐ Upper layer lifeform differs from dominant lifeform.												
☒ Herbaceous		SCSC													
☐ Shrub		Upper													
☐ Tree		BOGR2													
<u>Fuel Model</u>		Lower													
1		ARCA13													
<u>Description</u>		Upper													

This system was originally modeled as a four-box model with a mid and late stage; however, it was changed to a three-box model, combining the mid and late stages, since species and structural info was very similar, as were disturbances. It was then combined into a two-box model because of the lack of a prairie dog stage.

This mid-to-late seral stage would persist three years after a fire. The maximum cover height for grasses

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

would be approximately 60%, even though in other mapzones, cover might be much higher.

Other species indicators could be JUHO2 and SPCR - in the later part of this stage. Various sprouting shrubs may be established. The shrubs are as tall or taller than the grasses, but they would not be dominant; shrubs might occupy approximately 10% of the area. Some of the shrubs include Juniper horizontalis and skunkbush sumac (Rhus trilobata). Other woody species such as chokecherry (Prunus virginiana) and snowberry (Symphoricarpos occidentalis) may also be established.

This stage includes moderate grazing by native ungulates and insects.

Litter accumulates providing continuous fuel for fires thereby increasing the probability of larger fires. However, for the model, fires were attributed similarly between classes A and B.

Prairie dogs might impact this class with a very unlikely probability of 0.0007, bringing this class back to class A.

Other native grazing occurs with a probability of 0.01, bringing the class back to an early seral state, or with a probability of 0.15, maintaining this stage.

Replacement fires occur every 20yrs.

Class C	0 %	<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>												
[Not Used]	[Not Used]		<table><tr><td></td><td><i>Min</i></td><td><i>Max</i></td></tr><tr><td><i>Cover</i></td><td>%</td><td>%</td></tr><tr><td><i>Height</i></td><td></td><td></td></tr><tr><td><i>Tree Size Class</i></td><td colspan="2"></td></tr></table>		<i>Min</i>	<i>Max</i>	<i>Cover</i>	%	%	<i>Height</i>			<i>Tree Size Class</i>		
	<i>Min</i>	<i>Max</i>													
<i>Cover</i>	%	%													
<i>Height</i>															
<i>Tree Size Class</i>															
<u>Upper Layer Lifeform</u>			☐ Upper layer lifeform differs from dominant lifeform.												
☐ Herbaceous															
☐ Shrub															
☐ Tree	<u>Fuel Model</u>														

Description

Class D	0 %	<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>												
[Not Used]	[Not Used]		<table><tr><td></td><td><i>Min</i></td><td><i>Max</i></td></tr><tr><td><i>Cover</i></td><td>%</td><td>%</td></tr><tr><td><i>Height</i></td><td></td><td></td></tr><tr><td><i>Tree Size Class</i></td><td colspan="2"></td></tr></table>		<i>Min</i>	<i>Max</i>	<i>Cover</i>	%	%	<i>Height</i>			<i>Tree Size Class</i>		
	<i>Min</i>	<i>Max</i>													
<i>Cover</i>	%	%													
<i>Height</i>															
<i>Tree Size Class</i>															
<u>Upper Layer Lifeform</u>			☐ Upper layer lifeform differs from dominant lifeform.												
☐ Herbaceous															
☐ Shrub															
☐ Tree	<u>Fuel Model</u>														

Description

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Class E 0 %

[Not Used] [Not Used]

**Indicator Species and
Canopy Position**

Structure Data (for upper layer lifeform)

	Min	Max
Cover	%	%
Height		
Tree Size Class		

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Fuel Model

☐ Upper layer lifeform differs from dominant lifeform.

Description

Disturbances

Fire Regime Group:** II

Historical Fire Size (acres)

Avg
Min
Max

Fire Intervals

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
Replacement	20	2	100	0.05	100
Mixed					
Surface					
All Fires	20			0.05002	

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class.

Sources of Fire Regime Data

- ☒ Literature
☐ Local Data
☒ Expert Estimate

Additional Disturbances Modeled

- ☐ Insects/Disease ☒ Native Grazing ☒ Other (optional 1) Prairie Dogs
☐ Wind/Weather/Stress ☐ Competition ☐ Other (optional 2)

References

Auld, T.D. and R.A. Bradstock. 1996. Soil temperatures after the passage of a fire: Do they influence the germination of buried seeds? Australian Journal of Ecology 21: 106-109

Bragg, T.B. 1986. Fire history of a North American sandhills prairie. Page 99 in: Program of the Ivth International Congress of Ecology, Syracuse University, Syracuse, NY. 10-16 August 1986.

Bragg, T.B. 1997. Response of a North American sandhills grassland to spring, summer, and fall burning: Community resistance to disturbance (1984-1996). In: B.J. McKaige, R.J. Williams and W.M. Waggitt, editors. Bushfires 97 Proceedings. Parks Australia North and CSIRO Tropical Ecosystems Research Centre, Darwin, Northern Territory, Australia.

Bragg, T.B. 1998. Fire in the Nebraska sandhills prairie. Pages 179-194 in: Fire in ecosystem management: Shifting the paradigm from suppression to prescription, T.L. Pruden and L.A. Brennan, editors. Tall Timbers Fire Ecology Conference Proceedings No. 20, Tall Timbers Research Station, Tallahassee, FL.

Cheney, P and A. Sullivan. 1997. Grassfires: fuel, weather and fire behaviour. CSIRO Publishing, Australia

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Clark, O.R. 1940. Interception of rainfall by prairie grasses, weeds, and certain crop plants Ecological Monographs 10(2): 243-277

Lindvall, M. 2000. Evaluation of the Suitability of Habitat at Valentine National Wildlife Refuge for Prairie Dog Introduction. Draft for Review sent out in 2000.

NatureServe. 2006. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA, U.S.A. Data current as of 18 July 2006.

NatureServe. 2007. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. Data current as of 10 February 2007.

Pfeiffer, K.E. and A.A. Steuter. 1994. Preliminary response of Sandhills prairie to fire and bison grazing. J. Range Manage 47: 395-397.

Steuter, A.A., E.M. Steinauer, G.L. Hill, P.A. Bowers and L.L. Tieszen. 1995. Distribution and diet of bison and pocket gophers in a sandhills prairie. Ecological Applications, 5(3):756-766.

Swinehart, J. 2005. Personal communication at Rapid Assessment Northern Plains workshop.

USDA-NRCS. 2003. eFOTG: Electronic Field Office Technical Guide. Available at: <http://www.nrcs.usda.gov/technical/efotg/>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.