

LANDFIRE Biophysical Setting Model

Biophysical Setting 3214220

Southern Blackland Tallgrass Prairie

☐ This BPS is lumped with:

☐ This BPS is split into multiple models:

General Information

Contributors (also see the Comments field) **Date** 4/18/2007

Modeler 1 Jim Eidson jeidson@tnc.org

Reviewer

Modeler 2 Roger Mangham rmangham@tnc.org

Reviewer

Modeler 3

Reviewer

Vegetation Type

Upland
Grassland/Herbaceous

Dominant Species

SCHIZ4
SONU2

Map Zone

32

Model Zone

☐ Alaska
California

☐ Northern Plains
N-Cent. Rockies

General Model Sources

- ☒ Literature
☐ Local Data
☒ Expert Estimate

ANGE
TRIPS
PAVI2
SPSI2
TRST2
PAPL3

☐
☐ Great Basin
☐ Great Lakes
☐ Hawaii
☐ Northeast

☐
☐ Pacific Northwest
☒ South Central
☐ Southeast
☐ S. Appalachians
☐ Southwest

Geographic Range

This BpS extends through north central TX from the Red River to near the southern Gulf coast bordered by the Coastal Prairie (Kuchler: bluestem-sacahuista), to the east bordering and mingling with oak-hickory forest (savanna), in central portions bounded by eastern and western Cross Timbers, to the west bordered by the mesquite-buffalograss and bluestem-grama vegetation types (Kuchler 1964). [Text from Masters 2004]

Biophysical Site Description

The main belt of the Blackland Prairie is divided into four narrow, geomorphic areas aligned in a north south direction. These include-- from west to east-- the Eagle Ford Prairie, the White Rock Cuesta, the Taylor Black Prairie and the Eastern Marginal Prairie (Montgomery 1993). The soils of the Eagle Ford and Taylor Black Prairies are primarily clays of the order vertisol, while the soils of the White Rock Cuesta are mollisols and the Eastern Marginal Prairie of the order alfisol. Alfisols are the important soil order in the San Antonio prairie, while both Alfisols and Vertisols are important in the Fayette prairie. Microtopography such as gilgai on vertisols and mima mounds on alfisols are important microhabitats. Gilgai are shallow microdepressions one to several meters across formed by pedoturbation of montmorillonitic clays. Mima mounds are small circular hills which are variable in size but may be more than a meter high and 1-14m across. The origins of mima mounds are not clear and are probably of variable origin (Diamond and Smeins 1993). The climate is warm temperate to subtropical and humid. Precipitation ranges from 762mm on the western edge to 1016mm on the east (Eidson and Smeins 2001).

Vegetation Description

Little bluestem (*Schizachyrium scoparium*) and Indiangrass (*Sorghastrum nutans*) are frequently dominants on Blackland Prairie alfisols and vertisols. Big bluestem (*Andropogon gerardii*) is of variable

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

importance on vertisols and is frequently a dominant on Blackland Prairie mollisols. Gamagrass-switchgrass (*Tripsacum dactyloides*-*Panicum virgatum*) prairies are associated with bottomland sites throughout the region, and are also found on upland sites of the northern main belt vertisols where they are especially associated with gilgai microtopography. Silveanus dropseed- mead's sedge (*Sporobolus silveanus*-*Carex meadii*) prairies are found over low pH soils of the northern main belt. Little bluestem-brownseed paspalum (*S. scoparium*-*Paspalum plicatulum*) prairie is associated with Fayette Prairie alfisols. Each community differs further in secondary florae. For example, eastern forb species such as prairie blazing star (*Liatris pycnostachya*) and largeflower tickseed (*Coreopsis grandiflora*) are largely limited to the alfisols of the Eastern Marginal prairies, while grasses such as hairy grama (*Bouteloua hirsuta*) and seep muhly (*Muhlenbergia reverchonii*), as well as a diversity of species in the genus *Dalea* (prairie clover) are generally found on the mollisols of the White Rock Cuesta (Eidson and Smeins 2001).

Disturbance Description

The Blackland Prairie was a disturbance maintained system. Prior to European settlement (pre-1825 for the southern and pre-1845 for the northern half) important natural landscape-scale disturbances included fire and periodic grazing by large herbivores, primarily bison and, to a lesser extent, pronghorn (*Antilocapra americana*). Infrequent but intense fire combined with short duration grazing suppressed woody species and invigorated herbaceous prairie species. The latter were adapted to fire and grazing by virtue of maintaining perenniating tissues below ground. It has been suggested that second only to climate, fire has been the most important determinant of the spread and maintenance of grasslands (Anderson 1990). Fire frequency in the presettlement blackland prairie is unclear, but may have occurred at intervals of 1-3yrs (Pyne 1982). The majority of fires in grasslands were stand-replacement fires (top-killing greater than 75% of the grass), with surface fires and mixed fires occurring more frequently in woody stands. Both natural (i.e. lightning strike) and anthropogenic ignition sources are recognized. Bison were probably extirpated from the region by the 1850s (Eidson and Smeins 2001).

Adjacency or Identification Concerns

The modern landscape has been converted to croplands, tame pasture and urban areas. Natural Heritage surveys suggest only two percent of this BpS cover may have survived to the 21st century. Some pastures may contain native species or warm season grasses that give similar signatures from remote sensing.

Native Uncharacteristic Conditions

Scale Description

Stand replacement fires identified by early settlers are described as ranging over wide areas. It is worth noting that bison impacts to fuelbeds would have been eliminated by those mid-1800s descriptions, allowing for more widespread growth of fires. One could estimate this dominant fire type (in the grasslands) to regularly spread from thousands to tens of thousands of acres, moving through uplands between riparian areas. Grazing disturbances likely varied widely with short duration, high impact bison herds moving through thousands of acres at a time, but less frequently than in other areas of the Great Plains. Numerous other grazers were noted, including deer and pronghorn, that would have had more widespread but less intense impacts. Fires through grazed areas would not have spread extensively, unless able to break out into heavier, ungrazed prairie fuel.

Issues/Problems

There is an eastern extension of blackland prairie that occurs in southern AR and in MS into AL. This eastern blackland type is in a higher rainfall area and is smaller in contiguous extent and adjacent to southern woodland cover types. This model type focuses on the contiguous TX prairies.

****Fire Regime Groups are:** I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Comments

For MZs 32 and 35 this model was developed using the RA model R5PRBL -- Blackland Prairie by Blane Heumann (bheumann@tnc.org) and reviewed by Douglas Zollner and Maria Melnechuck. Changes to the description and the model resulted in a change in modelership. Suggested reviewers for MZs 32 and 35 include Dave Diamond (MORAP, Columbia, MO).

Vegetation Classes

Class A 90 %

Early Development 1 All Structure

Upper Layer Lifeform

- ☒ Herbaceous
☐ Shrub
☐ Tree

Fuel Model

3

Indicator Species and Canopy Position

SCHIZ4
Upper
ANGE
Upper
SONU2
Upper
SPAS
Upper

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	100 %
Height	Herb 0m	Herb 1.0m
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.

Description

This stage will persist indefinitely with disturbance. Post replacement herbaceous vegetation with open structure and minimal thatch. Diverse expression of forbs and annual species in the open herbaceous structure. Dominant and characteristic species vary across major soil types. Replacement fire and grazing are the dominant disturbance types (MFRI = 2.5yrs and grazing rotation = seven years) in this class. Drought is also an important disturbance factor that can replace this class (50yr return interval). Without disturbance for 10yrs, this class will succeed to class B.

Class B 5 %

Mid Development 1 Open

Upper Layer Lifeform

- ☐ Herbaceous
☒ Shrub
☐ Tree

Fuel Model

3

Indicator Species and Canopy Position

PRRI
Upper
CRATA
Upper
ULCR
Upper
CELA
Upper

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	40 %
Height	Shrub 0.6m	Shrub 3.0m
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.

Description

This class develops if fire does not occur in class A for 10yrs. Sugarberry (*C. laevigata*), cedar elm (*U. crassifolia*), honey locust (*Gleditsia triacanthos*), creek plum (*P. rivularis*) and hawthorn (*Crataegus* spp). (Note: these are tree species but are in their shrubby form). Replacement fire is the dominant disturbance type (surface MFRI = three years, mixed MFRI = 15yrs, replacement MFRI = 50yrs) in this class. Drought is also an important disturbance factor but it does not replace this class (15yr return interval).

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Class C 5 %

Late Development 1 Closed

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☒ Tree

Fuel Model

9

Indicator Species and Canopy Position

CELA
 Upper
 ULCR
 Upper
 GLTR
 Upper
 SMBO
 Lower

Structure Data (for upper layer lifeform)

	Min	Max
Cover	41 %	100 %
Height	Tree 5.1m	Tree 10m
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.
Description

Woodland to forest community. Forms closed mottes scattered throughout the grassland. Replacement fire is a rare event (every 50yrs). Mixed and low intensity fires are more likely at 15yrs and three years, respectively. Drought is also an important disturbance type in this system, with regular drought every 15yrs maintaining this class and replacement drought occurring every 50yrs.

Class D 0 %

[Not Used] [Not Used]

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Fuel Model**Indicator Species and Canopy Position****Structure Data (for upper layer lifeform)**

	Min	Max
Cover	%	%
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.
Description**Class E 0 %**

[Not Used] [Not Used]

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Fuel Model**Indicator Species and Canopy Position****Structure Data (for upper layer lifeform)**

	Min	Max
Cover	%	%
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.
Description**Disturbances**

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Fire Regime Group:** II

Historical Fire Size (acres)

Avg 2000

Min 100

Max 100000

Sources of Fire Regime Data

- ☒ Literature
- ☒ Local Data
- ☒ Expert Estimate

Additional Disturbances Modeled

- ☐ Insects/Disease
- ☒ Native Grazing
- ☐ Other (optional 1)
- ☒ Wind/Weather/Stress
- ☐ Competition
- ☐ Other (optional 2)

Fire Intervals

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
Replacement	3			0.33333	91
Mixed	170			0.00588	2
Surface	35			0.02857	8
All Fires	3			0.36779	

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class.

References

Anderson, R.C. 1990. The historic role of fire in the North American Grassland. In: L. Wallace and S. Collins [eds.], Fire in tallgrass prairie ecosystem. University of Oklahoma Press, Norman, OK. 8–18.

Diamond, D.D. and F.E. Smeins. 1993. The native plant communities of the Blackland Prairie. In: M.R. Sharpless and J.C. Yelderman, eds. The Texas Blackland Prairie, land, history, and culture. Baylor Univ. Program for Regional Studies, Waco, TX. 66-81.

Eidson, J. and F. Smeins. 2001. Terrestrial ecoregions of North America: a conservation assessment. Island Press, Washington D.C.

Kuchler, A.W. 1964. Potential Natural Vegetation of the Conterminous United States, American Geographical Society, Special Publication No. 36.

Masters, Ron. 2004. Fire Regime Condition Class (FRCC) Interagency Handbook Reference Conditions. Blackland Prairie (Kuchler Type 76 and 88-Fayette Prairie and also San Antonio Prairie which was not mapped by Kuchler) - PRAR6. Available online:
http://www.frcc.gov/docs/PNVG/EastNew/PRAR6_Description.pdf.

NatureServe. 2007. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA, U.S.A. Data current as of 10 February 2007.

Packard, S. and C.F. Mutel. 1997. The Tallgrass Restoration Handbook: For Prairies, Savannahs, and Woodlands. Island Press, Washington, D.C./Covelo, California, USA.

Pyne, S.J. 1982. Fire in America: A cultural history of wildland and rural fire. Princeton, NJ: Princeton University Press.

Texas Parks and Wildlife. 1978. Ecological Regions in Texas. From LBJ School of Public Affairs, Preserving Texas' Natural Heritage. LBJ School of Public Affairs Policy Research Report #31.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Texas Parks and Wildlife. Post Oak Savannah and Blackland Prairie Wildlife Management. 4200 Smith School Road, Austin, TX.

United States Department of Agriculture. Dec. 1981. Alabama, Mississippi, and Arkansas Blackland Prairie. Land Resource Regions and Major Land Resource Areas of the United States. USDA Soil Conservation Service Handbook 296. Pages 97-98.

United States Department of Agriculture. Dec. 1981. Texas Blackland Prairie. Land Resource Regions and Major Land Resource Areas of the United States. USDA Soil Conservation Service Handbook 296. Pages 62-63.

World Wildlife Fund. 2001. Texas blackland prairies (NA0814). Terrestrial ecoregions of North America: a conservation assessment. Island Press, Washington D.C.

****Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.**