

LANDFIRE Biophysical Setting Model

Biophysical Setting 3511320

Central Mixedgrass Prairie

- ☐ This BPS is lumped with:
☐ This BPS is split into multiple models:

General Information

Contributors (also see the Comments field)

Date 1/10/2007

Modeler 1 Delbert M. Bassett

dmbassett@neo.tamu.edu

Reviewer Sam Fuhlendorf

sam.fuhlendorf@okstate.edu

Modeler 2 Lee Elliott

lelliott@tnc.org

Reviewer

Modeler 3

Reviewer

Vegetation Type

Upland
Grassland/Herbaceous

Dominant Species

BOBA3
SCSC

Map Zone

35

Model Zone

☐ Alaska
☐ California

☐ Northern Plains
☐ N-Cent. Rockies

General Model Sources

- ☒ Literature
☒ Local Data
☒ Expert Estimate

NALE3
ERIN
BOGR2
BOCU
SONU2
PRGL2

☐
☐ Great Basin
☐ Great Lakes
☐ Hawaii
☐ Northeast

☐
☐ Pacific Northwest
☒ South Central
☐ Southeast
☐ S. Appalachians
☐ Southwest

Geographic Range

This type historically occurs in western KS, western NE, eastern CO, northeastern NM, eastern TX panhandle into central TX, and along the western portion of OK proper. This BpS comprises the eastern half of MZ34 in ECOMAP sections 332F and 315C. This type occurs in the western half of MZ32 and in the northern half of MZ35 (west of Interstate-35).

Biophysical Site Description

This type occurs on sandy loam, loamey, or clayey upland sites of the southern Great Plains. In Texas and Oklahoma, elevations range from 915-275m (Wright and Bailey 1982). Precipitation ranges from 20-28in (Wright and Bailey 1982), although mixedgrass can occur on sites with up to 32in, and occurs mostly from May through September.

Vegetation Description

Historically, vegetation was generally dominated by midgrass, with some areas occupied by tallgrasses or short grasses, as an effect of grazing, edaphic and topographic conditions. (Species in order of dominance in boxes.) Dominant species include mostly midgrasses with some mix of tall and short grass - side oats grama (*Bouteloua curtipendula*), needlegrasses, little bluestem (*Schizachyrium scoparium*), *Bothriochloa barbinodis*, *Nassella leucotricha* and *Eragrostis intermedia*. Tallgrasses such as yellow indiagrass (*Sorghastrum nutans*), big bluestem (*Andropogon gerardii*) and switchgrass (*Panicum virgatum*) also occur. Blue grama (*B. gracilis*) can be prevalent, especially following significant grazing and on poor sites. Forbs present can include purple prairie clover (*Dalea purpurea*), vervain (*Verbena bipinnatifida*), dotted gayfeather (*Liatris punctata*). Mesquite (*Prosopis glandulosa*), Lotebush (*Ziziphus obtusifolia*), Soapweed yucca (*Yucca glauca*), Juniper species and prickly pear (*Opuntia* spp.) are present and may

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

dominate large portions of the landscape under current conditions.

Shrubs included four-wing saltbush (*Atriplex canescens*), winterfat (*Krascheninnikovia lanata*), with lesser amounts of rabbitbrush (*Chrysothamnus nauseosus*), broom snakeweed (*Gutierrezia sarothrae*), fringed sage (*Artemisia frigida*) and sunsedge (*Carex inops* ssp. *heliophila*) are present mostly in MZ34. In MZs 32 and 35 dominant shrub species include broom snakeweed (*Gutierrezia sarothrae*), shinnery oak (*Quercus havardii*), sand sage brush, sumac (*Rhus glabra*) and chickasaw plum (*Prunus angustifolia*).

Disturbance Description

Historically, fire return intervals were probably approximately 4-7yrs - slightly shorter return interval (more frequent fire) than shortgrass prairie due to higher fuel loads.

Episodic disturbance caused by insect infestation (grasshoppers, range caterpillars, Mormon crickets). Large herds of bison went through this system - as well as deer and pronghorn. Grazing impact would be greatest in recently burned areas because grasses are more palatable and nutritious.

This is a drought tolerant system. However, extended drought (over 3-4yrs) will reduce cover. Drought, grazing and fire were probably most important disturbances historically.

Adjacency or Identification Concerns

This system could be confused with shortgrass prairie to the west and tallgrass prairie to the east. Production is less in shortgrass versus mixedgrass prairie. Grasses are taller in mixedgrass than shortgrass prairie. These systems are intermixed, with the shorter grasses further west with less precipitation (other than the foothills areas).

Naturally occurring shrub dominated areas on sandy soils, on deposits associated with riparian areas can be found throughout the mixedgrass prairie type. It is hard to distinguish whether these shrub areas should be considered as part of the mixedgrass or if they should be considered sand sagebrush or sand shinnery oak. At finer scales, these types are easily distinguished and likely can be mapped at 30m resolution as a shrub community rather than grass.

What adds to the confusion is that current overgrazing has transitioned large portions of the region from mixed grass to short grass. In addition, the areas that have transitioned to a short grass prairie have also been invaded by woody shrubs, mainly honey mesquite (*Prosopis glandulosa*) and eastern redcedar (*Juniperus virginiana*).

What has historically been a mixedgrass prairie has been converted to row crops/ag. The mixedgrass prairie is primarily in rural areas so there is not as much residential development; however, urbanization is coming at an alarming rate. Mixedgrass prairie is now extremely rare as a result of land conversion to agriculture and woody species invasion.

Current condition is a mesquite or mixed shrub dominated landscape, resulting from spread of mesquite and loss of fire in the system.

Native Uncharacteristic Conditions

Mesquite (*P. glandulosa*), eastern redcedar (*Juniperus virginiana*) and Ashe juniper (*J. ashei*) have expanded well beyond their expected historical range and/or density. The introduction of cattle and their eating of the mesquite bean has promoted the expansion of the woody shrub. Also, overgrazing of the grasses has also allowed the mesquite to expand. Fire suppression has allowed eastern redcedar to encroach

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

into grasslands.

Scale Description

This is a matrix community - small to large patches. Disturbances can also occur within a matrix - small to large, huge patches. Driving variable is climate (drought, low rainfall, etc), grazing, and to a lesser extent fire.

Issues/Problems

A reviewer for MZs 32 and 35 indicated that both class A and B could start immediately after a fire (i.e. age zero) and that grazing could be removed as a disturbance from the model because it is the grazing/fire interaction, which could be represented by the fire alone, that causes the transition between states in the model. The regional lead could not change the start age of classes A and B to zero because LANDFIRE rules only allow one class to start at age zero. However, the regional lead did run the model with out grazing, as suggested by the reviewer, and found that it did not change the class percents or the fire return interval. The regional lead chose to leave grazing in the model to maintain consistency with MZ34 but it should be recognized that the same results can be achieved if grazing is removed from the model. Furthermore, it should be recognized that there is an interaction between fire and grazing in this BpS even though the two disturbances are represented independently in the VDDT model.

Comments

For MZ35 this model was adopted without changes from the same BpS for MZ32.

Suggested reviewers for MZ34 include Dr. Fred Smeins (TAMU, RLEM), Dr. Mark Simmons (Wildflower Center, Austin, TX), Dr. Steve Windhager (Wildflower Center, Austin, TX).

For MZs 35 and 32 this model was adopted from the same BpS in MZ34 modeled by Delber Bassett. Lee Elliott reviewed the model for MZ34 and altered it for MZs 35, 34 and 32. The same description and VDDT model now represent this BpS in MZs 35, 34 and 32. Sam Fulendorf reviewed this system for MZs 32 and 35.

For MZ34 this model was developed from the MZs 27 and 33 model, developed by Dan Nosal (dan.nosal@co.usda.gov), Rich Sterry (richard_sterry@fws.gov) and Terri Schulz (tschulz@tnc.org), as a starting point. Both the description and the model were significantly altered resulting in a change in modelership.

Vegetation Classes															
Class A	50 %	<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>												
Early Development 1 Open		BOGR2	<table><thead><tr><th></th><th>Min</th><th>Max</th></tr></thead><tbody><tr><td>Cover</td><td>0 %</td><td>30 %</td></tr><tr><td>Height</td><td>Herb 0m</td><td>Herb 0.5m</td></tr><tr><td>Tree Size Class</td><td colspan="2">None</td></tr></tbody></table>		Min	Max	Cover	0 %	30 %	Height	Herb 0m	Herb 0.5m	Tree Size Class	None	
	Min	Max													
Cover	0 %	30 %													
Height	Herb 0m	Herb 0.5m													
Tree Size Class	None														
<u>Upper Layer Lifeform</u>		Upper													
☒ Herbaceous		BUDA													
☐ Shrub		Upper													
☐ Tree		AMPS													
	<u>Fuel Model</u>	Upper	☐ Upper layer lifeform differs from dominant lifeform.												
	1	AMDR													
<u>Description</u>		Upper													

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Class A is early succession stage. Grasses that exert more dominance in the early stages include western wheatgrass (*Pascopyrum smithii*) and tumble windmill grass (*Chloris verticillata*). Indicator grasses, such as blue grama (*Bouteloua gracilis*) and buffalograss (*Buchloe dactyloides*) will still be present, but in lesser amounts than the climax community. Annual forbs such as broomweed (*Amphiachyris dracunculoides*), ragweed (*Ambrosia psilostachya*) and false mesquite (*Calliandra eriophylla*) will typically be present early.

In this early stage, grasses will be more nutritious and palatable, grazing by bison, antelope, deer, rodents and lagomorphs would be common and maintain this class (modeled as native grazing with a probability of 0.1) Replacement fire might occur every 15yrs.

Class B	40 %	<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>
Late Development 1 Closed		BOBA3	
<u>Upper Layer Lifeform</u>		Upper	
☒ Herbaceous		SCSC	
☐ Shrub		Upper	
☐ Tree	<u>Fuel Model</u>	BOCU	
	3	Lower	
		ZIOB	
<u>Description</u>		Upper	
			☐ Upper layer lifeform differs from dominant lifeform.
			Scattered shrubs may be present (up to 10%, maybe up to one meter) - honey mesquite, Ashe juniper (<i>Juniperus ashei</i>), and lotebush (<i>Ziziphus obtusifolia</i>), yucca (sandy sites) and elbowbush (<i>Forestiera pubescens</i>).

This is the historic climax plant community with big bluestem (*Andropogon gerardii*), little bluestem (*Schizachyrium scoparium*), switchgrass (*Panicum virgatum*), yellow indiangrass (*Sorghastrum nutans*), blue grama (*Bouteloua gracilis*), buffalograss (*Buchloe dactyloides*), sideoats grama (*Bouteloua curtipendula*) and hairy grama (*Bouteloua hirsuta*).

There would also be scatterings of black grama (*B. eriopoda*) and vine mesquite (*P. obtusum*) on heavier soils.

Replacement fires occur more often in this class, every 10yrs. Native grazing maintains this class (modeled as native grazing with a probability of 0.1). Replacement drought also could occur every 40yrs. Without disturbance for 12yrs, this class will succeed to class C.

Class C	10 %	<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>
Late Development 2 Closed		ZIOB	
		Upper	
<u>Upper Layer Lifeform</u>		JUAS	
☐ Herbaceous		Upper	
☒ Shrub		ZIOB	
☐ Tree	<u>Fuel Model</u>	Upper	
	3	PRGL2	
		Lower	
<u>Description</u>			
			☐ Upper layer lifeform differs from dominant lifeform.

Shrubs - honey mesquite, Ashe juniper (*Juniperus ashei*), lotebush (*Ziziphus obtusifolia*), yucca (sandy sites) and elbowbush (*Forestiera pubescens*). Replacement fires occur in this class every 10yrs; mixed fire maintains this class every 15yrs. Native grazing maintains this class (modeled as native grazing with a

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

probability of 0.1). Drought occurring at approximately every 30yrs could drive this class to B.

Class D	0 %	<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>
[Not Used]	[Not Used]		
<u>Upper Layer Lifeform</u>			
☐ Herbaceous			
☐ Shrub			
☐ Tree	<u>Fuel Model</u>		☐ Upper layer lifeform differs from dominant lifeform.

Description

Class E	0 %	<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>
[Not Used]	[Not Used]		
<u>Upper Layer Lifeform</u>			
☐ Herbaceous			
☐ Shrub			
☐ Tree	<u>Fuel Model</u>		☐ Upper layer lifeform differs from dominant lifeform.

Description

Disturbances							
<u>Fire Regime Group**:</u>	II	<u>Fire Intervals</u>	<i>Avg FI</i>	<i>Min FI</i>	<i>Max FI</i>	<i>Probability</i>	<i>Percent of All Fires</i>
<u>Historical Fire Size (acres)</u>		<i>Replacement</i>	6			0.16667	95
		<i>Mixed</i>	125			0.008	5
		<i>Surface</i>					
		<i>All Fires</i>	6			0.17468	
<u>Sources of Fire Regime Data</u>		<u>Fire Intervals (FI):</u> Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class.					
☒ Literature							
☒ Local Data							
☒ Expert Estimate							
<u>Additional Disturbances Modeled</u>							
☐ Insects/Disease	☒ Native Grazing	☐ Other (optional 1)					
☒ Wind/Weather/Stress	☐ Competition	☐ Other (optional 2)					

References
Brown, James K. and Jane Kapler Smith, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station. 257 pp.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

- Dick-Peddie, W.A. 1993. New Mexico vegetation, past, present and future. Albuquerque, NM: Univ. New Mexico Press. Xxxii, 244 pp.
- Ford, P.L. 1999. Response of buffalograss (*Buchloe dactyloides*) and blue grama (*Bouteloua gracilis*) to fire. *Great Plains Research* 9: 261-276.
- Fueldorf, S.D. and D. Engle. 2004. Application of the fire grazing interaction to restore a shifting mosaic on tallgrass prairie. *Journal of applied ecology* 41: 604-614.
- Fueldorf, S.D. and D. Engle. 2001. Restoring heterogeneity on rangelands: ecosystem management based on evolutionary grazing patterns. *Bioscience* 51: 625-632.
- Howard, Janet L. 1995. *Buchloe dactyloides*. In: Fire Effects Information System, [Online]. USDA Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis/> [2005, May 4].
- Miller, Greg et al. 1993. Terrestrial Ecosystem Survey of the Santa Fe National Forest. USDA Forest Service Southwestern Region.
- NatureServe. 2007. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA, U.S.A. Data current as of 10 February 2007.
- NatureServe. 2005. NatureServe Explorer: An online encyclopedia of life [web application]. Version 4.4. NatureServe, Arlington, Virginia. Available <http://www.natureserve.org/explorer>. (Accessed: May 4, 2005).
- Wright, H.A. and A.W. Bailey. 1982. Fire Ecology, United States and Southern Canada. John Wiley & Sons, Inc. New York.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.