

LANDFIRE Biophysical Setting Model

Biophysical Setting 6716942

Alaska Arctic Tussock Tundra - Infrequent Fire

☒ This BPS is lumped with: see below

☒ This BPS is split into multiple models: Alaska Arctic Tussock Tundra, Alaska Arctic Shrub-Tussock Tundra and Alaska Arctic Tussock-Lichen Tundra were lumped and then split into two models based on differences in the MFRI.

General Information

Contributors (also see the Comments field)

Date 9/30/2008

Modeler 1 Coleen Ryan

colleen_ryan@tnc.org

Reviewer Janet Jorgenson

Janet_Jorgenson@fws.gov

Modeler 2 Kori Blankenship

kblankenship@tnc.org

Reviewer

Modeler 3 Keith Boggs

Ankwb@uaa.alaska.edu

Reviewer

Vegetation Type

Upland Shrubland

Map Zone

67

Model Zone

- | | |
|--|--|
| ☒ Alaska | ☐ N-Cent.Rockies |
| ☐ California | ☐ Pacific Northwest |
| ☐ Great Basin | ☐ South Central |
| ☐ Great Lakes | ☐ Southeast |
| ☐ Northeast | ☐ S. Appalachians |
| ☐ Northern Plains | ☐ Southwest |

Dominant Species*

ERVA4 SAPU15
VAUL LEPAD
CABI5 EMNI
BENA

General Model Sources

- ☒ Literature
☐ Local Data
☒ Expert Estimate

Geographic Range

This BpS is found in arctic AK except within ecoregions (Nowakii et al. 2001) 4, 5, 7 and south of the Brooks Range in ecoregion 3.

Biophysical Site Description

This type is common in valleys and slopes. These sites are cold, poorly drained, and underlain by mesic, silty mineral soils with a shallow surface organic layer surrounding the tussocks (Viereck et al. 1992). Permafrost is present.

Vegetation Description

Eriophorum vaginatum is the primary tussock-former in most sites, but Carex bigelowii may dominate some sites. Calamagrostis canadensis, Arctagrostis latifolia and Chamerion latifolium may be common. Shrubs include Betula nana, Salix pulchra, Ledum palustre ssp. decumbens, and Vaccinium spp. (e.g. Vaccinium vitis-idaea and Vaccinium uliginosum) and Empetrum nigrum. Mosses (Sphagnum spp., Polytrichum strictum and Hylocomium splendens) may form a nearly continuous mat between tussocks. Lichens are more common on the drier tussock tundra sites in western Alaska and may include Flavocetraria cucullata, Cetraria islandica, Cladonia spp., Cladina rangiferina and Thamnia subuliformis.

Acidic sites have more ericaceous shrubs and Sphagnum and less Eriophorum spp, Betula nana and Carex bigelowii. Acidic sites also have more organic matter buildup and the tussocks tend to be larger.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Disturbance Description

There is little information about fire regimes in the arctic region of Alaska (Viereck and Schandelmeier 1980). Mean fire return interval estimates for tussock tundra ecosystems include:

- 50-300yrs (personal communication FRCC experts' workshop March 2004)
- According to lake-core records, the fire-return interval is approximately 240yrs on the Seward Peninsula, and 1000yrs+ on the Beaufort Coastal Plain (Jennifer Allen pers. comm.).
- 260yr (s.d. 170) fire return interval for past 1500yrs for Noatak National Preserve (preliminary data from Higuera et al. 2008)
- 612yrs for Noatak River watershed (all vegetation types) (Racine et al 1983)
- 611yr fire rotation for Noatak River watershed (all vegetation below 600m which is predominantly tundra) (Racine et al 1985)
- Rapid recovery following fire makes fire frequency difficult to determine (Wein 1971)
- The fire regime of tundra systems are likely quite variable from one region to another making generalizations difficult (Viereck and Schandelmeier 1980).

Frost action, polygonal ground and thermokarst dynamics are widespread, small-scale and continuous disturbances within the Tussock Tundra type. Change in the arctic and subarctic climate is another source of disturbance that is currently affecting tundra ecosystems.

Adjacency or Identification Concerns

This type can be found adjacent to most other BpS in the arctic.

Native Uncharacteristic Conditions

Scale Description

This type can be found in large or small patches and is sometimes matrix forming.

Issues/Problems

The modelers were not sure what percent of the landscape should be in each succession class.

Comments

This model was based on input from the experts who attended the LANDFIRE Fairbanks, Arctic (April 08) modeling meeting and was refined by Colleen Ryan, Kori Blankenship and Keith Boggs.

Vegetation Classes

Class A 5%

Early Development 1 All Structures

Upper Layer Lifeform

- ☒ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and Canopy Position

ERVA4 Upper
CABI5 Upper
CACA4 Upper
ARCTA Upper

Structure Data (for upper layer lifeform)

	Min	Max
Cover	Herbaceous	Herbaceous
Height	Herbaceous	Herbaceous
Tree Size Class	None	

☐ Upper layer lifeform differs from dominant lifeform.

Classes A and C should be mapped based on EVT if possible. Class C represents Alaska Arctic Tussock-Lichen Tundra.

Description

0-14yrs

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

This class is characterized by tussock forming sedges. Common species include *Eriophorum vaginatum*, *Carex bigelowii* and *Rubus chamaemorus*. Cottongrass and sedges regrow from rhizomes. Grasses, including *Calamagrostis canadensis*, *Arctagrostis latifolia* and *Chamerion latifolium* (= *Epilobium latifolium*), may be common, especially for the first 5-10yrs. Mosses (*Sphagnum* spp., *Polytrichum strictum* and *Hylocomium splendens*) may form a nearly continuous mat between tussocks.

Succession to class B. Replacement MFRI = 1000yrs.

Class B 50 %

Structure Data (for upper layer lifeform)

Late Development 1 All Structures

Upper Layer Lifeform

- ☐ Herbaceous
☒ Shrub
☐ Tree

Indicator Species* and

Canopy Position

BENA Upper
 SALIX Upper
 ERVA4 Upper
 CABI5 Upper

	Min	Max
Cover	Open Shrub (25-74% shrub cover)	Closed Shrub (> 75% shrub cover)
Height	Low Shrub (20cm to 1.5m)	Low Shrub (20cm to 1.5m)
Tree Size Class	None	

☐ Upper layer lifeform differs from dominant lifeform.

Description

15yrs+

Low shrub-tussock. Tussock shrub tundra has >35% cover of sedges in a tussock growth form, and the combined cover of dwarf and low shrubs is >25% (Boggs et al. 2008). Tussocks are dominated by *Eriophorum* (cottongrass) and *Carex* spp. (sedges). Common shrub species include *Betula nana*, *Salix* spp., *Ledum palustre* ssp. *decumbens*, *Vaccinium vitis-idaea* and *V. uliginosum*. Lichens begin to re-establish but do not reach former abundance until 50-120yrs following fire. Former abundances of all species are typically reached 50-120yrs post fire. Lichens, if present, have <25% cover.

This class can persist in the absence of disturbance or follow an alternate succession pathway to class C (probability = 0.001). Replacement MFRI = 1000yrs.

Class C 45 %

Structure Data (for upper layer lifeform)

Late Development 2 All Structures

Upper Layer Lifeform

- ☒ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and

Canopy Position

CLADO3 Upper
 ERVA4 Upper
 CABI5 Upper
 Lower

	Min	Max
Cover	Herbaceous	Herbaceous
Height	Herbaceous	Herbaceous
Tree Size Class	None	

☐ Upper layer lifeform differs from dominant lifeform.

Classes A and C should be mapped based on EVT if possible. Class C represents Alaska Arctic Tussock-Lichen Tundra.

Description

80yrs+

This class is characterized by >35% cover of sedges in a tussock growth form, <25% shrub cover, and >25% lichen cover. Dwarf-shrubs may be common. *Eriophorum vaginatum* is the primary tussock-former in most

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

stands, but *Carex bigelowii* may dominate some sites. Lichens may include *Flavocetraria cucullata* (= *Cetraria cucullata*), *Cetraria islandica*, *Cladonia* spp., *Cladonia rangiferina* and *Thamnolia subuliformis*. This class tends to occur on drier sites. Permafrost is present, and sphagnum is an important component of the ground cover on these acidic sites.

This class can persist in the absence of disturbance. Replacement MFRI = 1000yrs.

Class D 0 %

[Not Used] [Not Used]

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover		
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.

Description

Class E 0 %

[Not Used] [Not Used]

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover		
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.

Description

Disturbances

Fire Regime Group:** V

Historical Fire Size (acres)

Avg 0

Min 0

Max 0

Sources of Fire Regime Data

- ☐ Literature
☐ Local Data
☒ Expert Estimate

Additional Disturbances Modeled

- ☐ Insects/Disease ☐ Native Grazing ☐ Other (optional 1)
☐ Wind/Weather/Stress ☐ Competition ☐ Other (optional 2)

Fire Intervals

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
Replacement	1023			0.00098	98
Mixed					
Surface					
All Fires	1021			0.001	

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class.

References

Boggs et al. 2008. International Ecological Classification Standard: Terrestrial Ecological Classifications. Draft Ecological Systems Description for the Alaska Arctic Region.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Higuera, P.E., M. Chipman, J. Allen, S. Rupp and F.S. Hu. 2008. Preliminary data provided by Jenifer Allen from a study of tundra fire regimes in the Noatak National Preserve since 6000 years before present.

Nowacki, G., P. Spencer, M. Fleming, T. Brock and T. Jorgenson. 2001. Unified ecoregions of Alaska. U.S. Department of the Interior, U.S. Geological Survey. Open file-report 02-297. 2 page map.

Payette, S., C. Morneau, L. Sirois and M. Despons. 1989. Recent fire history of the northern Quebec biomes. *Ecology*. 70:656-673.

Personal communication experts' workshop, March 2-4 2004. Fire Regime Condition Class (FRCC) interagency experts' workshop to develop and review Potential Natural Vegetation (PNV) groups for Alaska. Anchorage, Alaska.

Racine, C.H., W.A. Patterson III and J.G. Dennis. 1983. Permafrost thaw associated with tundra fires in northwest Alaska. Pages 1024-1029 in *Proceedings, Permafrost, Fourth International Conference*. National Academy Press, Washington, D.C.

Racine, C.H., J.G. Dennis and W.A. Patterson III. 1985. Tundra fire regimes in the Noatak River Watershed, Alaska: 1956-83. *Arctic* 38:194-200.

Racine, C.H., L.A. Johnson and L.A. Viereck. 1987. Patterns of vegetation recovery after tundra fires in northwestern Alaska, USA. *Arctic and Alpine Research*. 19:461-469.

Viereck, L.A., C.T. Dyrness, A.R. Batten, K.J. Wenzlick. 1992. The Alaska vegetation classification. Pacific Northwest Research Station, USDA Forest Service, Portland, OR. Gen. Tech. Rep. PNW-GTR286. 278 p.

Viereck, L.A. and L.A. Schandelmeier. 1980. Effects of fire in Alaska and adjacent Canada--a literature review. BLM-Alaska Tech. Rep. No. 6. Anchorage, Alaska: U.S. Department of the Interior, Bureau of Land Management. 124 p

Wein, R.W. 1971. Panel discussion: In: Slaughter, C.W., Barney, Richard J., and Hansen, G.M. (editors). 1971. *Fire in the Northern Environment – A Symposium*. Sponsors: Alaska Forest Fire Council and Society of American Foresters. Held at: The University of Alaska, Fairbanks. April 13-14, 1971. Published by: Pacific.

Wein, R.W. 1976. Frequency and characteristics of arctic tundra fires. *Arctic*, 29,213-222.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.