

LANDFIRE Biophysical Setting Model

Biophysical Setting 6816141

**Western North American Boreal Montane
Floodplain Forest and Shrubland - Boreal**

☐ This BPS is lumped with:

☒ This BPS is split into multiple models: Western North American Boreal Montane Floodplain Forest and Shrubland was split into a Boreal and Sub-boreal variant for BpS modeling so that regional differences could be represented.

General Information

Contributors (also see the Comments field)

Date 4/7/2008

Modeler 1 Michelle Schuman	michelle.schuman@ak.usda.gov	Reviewer Tom Paragi	tom.paragi@alaska.gov
Modeler 2 Karen Murphy	karen_a_murphy@fws.gov	Reviewer	
Modeler 3 Evie Witten	ewitten@tnc.org	Reviewer	

Vegetation Type

Forest and Woodland

Map Zone

68

Model Zone

☒ Alaska	☐ N-Cent.Rockies
☐ California	☐ Pacific Northwest
☐ Great Basin	☐ South Central
☐ Great Lakes	☐ Southeast
☐ Northeast	☐ S. Appalachians
☐ Northern Plains	☐ Southwest

Dominant Species*

POBAB2 SHCA
PIGL ROAC
ALNUS VIED
SALIX EQUIS

General Model Sources

☒ Literature
☐ Local Data
☒ Expert Estimate

Geographic Range

This type occurs on floodplains throughout the boreal region of AK. It may also be found in arctic AK.

Biophysical Site Description

The following information was taken from the draft Boreal Ecological Systems description (NatureServe 2008):

The substrate is typically well-drained sand or cobble, although finer silts and clays can be found on higher terraces, in ponds, on distal floodplains, and in lower energy systems. Permafrost is usually absent. Oxbows and other wet depressions commonly form on the floodplains, and these sites commonly develop into wetlands.

Vegetation Description

Primary succession on floodplains begins when new alluvial surfaces are colonized by tree, shrub, and herbaceous species. Common early seral woody species include *Populus balsamifera* (seedlings), *Picea glauca* (seedlings), *Alnus viridis* ssp. *sinuata*, *Alnus incana* ssp. *tenuifolia*, *Salix barclayi* and *Salix alaxensis* (Boggs 2000, Scott 1974, Shephard 1995, Thilenius 1990, Viereck 1966). Common early seral herbaceous species may include *Lupinus* spp., *Hedysarum* spp., and *Equisetum* spp. (NatureServe 2008). The next seral stage includes communities dominated by *Populus balsamifera* and/or *Picea glauca* with an understory of *Alnus viridis* ssp. *sinuata*, *Salix* spp., and bryophytes. On dry sites the shrub layer may be dominated by *Shepherdia canadensis*, *Dryas octopetala*, *D. integrifolia* and fruticose lichens

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

(*Steroucaulon* spp.) (NatureServe 2008). The tall shrub component of the early successional stages diminishes rapidly, probably because of decreased light from the dense tree overstory. *Populus balsamifera* does not regenerate in the understory and consequently, *Picea glauca* gains dominance in the overstory within 150yrs (NatureServe 2008). *Rosa acicularis* and *Viburnum edule* are common understory shrubs on older surfaces.

Disturbance Description

The following information was taken from the draft Boreal Ecological Systems description (NatureServe 2008):

Flooding is the primary disturbance in this system. Flooding can be caused by snowmelt, precipitation, ice jams and glacial runoff. Different rivers or portions of rivers may be more prone to certain types of flooding. Frequent flooding and channel migration create a pattern of gravel bars and early successional stages across the valley bottom. Sediment deposition raises the surface of the floodplain over time. As the terrace becomes farther removed from the channel, flooding becomes less frequent. Water availability on terraces plays a major role in community structure and composition. Water inputs are from overbank flow (flooding), ground water, and precipitation. Deposits with high permeability become progressively drier as they are vertically and horizontally removed from the active channels.

Vegetation succession on gravel bars can be represented by the following seral stages: barren or herbaceous, willow or willow-alder, alder, poplar or spruce poplar, spruce. Oxbows and other wet depressions commonly form on the floodplains, and develop into wetlands. Succession and species composition is, however, variable due to diverse environmental conditions such as water depth, substrate and nutrient input.

The following information was taken from the Riparian Spruce Hardwood Potential Natural Vegetation Group (PNVG) model (Murphy and Witten 2006):

Estimates of mean fire return intervals include:

- 200yrs+ (200-300yr range) (Viereck 1973, Barney 1971)
- 300yrs (Rowe et al 1974) (for alluvial white spruce MacKenzie River Valley)
- 300yrs (Heinselman 1981)
- 300yrs (Duchesne and Hawkes 2000)
- 300yrs (personal communication experts' workshop March 2004)

Small, relatively infrequent, mixed severity fires characterize this PNVG due to the sites' proximity to rivers, which act as fire breaks (Viereck 1973, Barney 1971, Foote 1983). High moisture content of the vegetation, high percentage of deciduous species, and high relative humidity also contribute to making fires less frequent in the Riparian Spruce Hardwood PNVG than in typically adjacent PNVGs. In interior Alaska the oldest white spruce stands (350+ yrs) are commonly found on islands of floodplains where they are protected from fire (Viereck 1973).

Adjacency or Identification Concerns

This model applies to forest and shrub systems in the active and inactive portions of the floodplain, but not abandoned floodplains. Oxbows and other wet depressions commonly form on the floodplains. Floodplain Wetlands are a separate ecological system and a separate BpS.

Native Uncharacteristic Conditions

Scale Description

Linear.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Issues/Problems

Wetlands that occur on the floodplain are not considered in this model.

Comments

This model was created for the AK boreal region and did not receive review for the arctic region.

This model was based on the FRCC Guidebook PNVG model for Riparian Spruce Hardwood (RSH; Murphy and Witten 2006) and input from the experts who attended the LANDFIRE Fairbanks (Nov. 07) modeling meeting. It was refined by Michelle Schuman. The relative age function used in the RSH model was not used in any class except A to comply with LANDFIRE modeling rules and the 10,000 year replacement fire was removed from class D. These changes did not change the model results. Because changes to the VDDT model were relatively minor Karen Murphy and Evie Witten were retained as modelers and Michelle Schuman's name was added.

Vegetation Classes

Class A 5%

Early Development 1 All Structures

Upper Layer Lifeform

- ☐ Herbaceous
☒ Shrub
☐ Tree

Indicator Species* and Canopy Position

LUPIN Upper
HEDYS Upper
EQUIS Upper
SALIX Upper

Structure Data (for upper layer lifeform)

	Min	Max
Cover	Open Shrub (25-74% shrub cover)	Open Shrub (25-74% shrub cover)
Height	Dwarf Shrub (< 20 m)	Tall Shrub (>1.5m)
Tree Size Class	Seedling/Sapling <5"	

☐ Upper layer lifeform differs from dominant lifeform.

Description

0-4yrs

This class is characterized by post disturbance regeneration (herbs, shrub regeneration, seedlings). Silt is deposited on the inside of river meanders following flood events. Flooding deposits seeds which germinate and take root. Lupinus spp., Hedysarum spp., Equisetum spp. and Salix spp. colonize in the first year. Within five years Salix spp and balsam poplar seedlings are abundant. Plant cover is 1-2% first year. Shrub cover increases up to 40% by the fifth year, with a diverse herbaceous layer underneath. Occasionally white spruce will germinate in large numbers on mineral soil after flooding, resulting in a dense, even-aged stand and eventual succession to class E (via class B).

Succession to class B. Flooding (probability = 0.05), represented by Option 1 in the model, resets the age of this class.

Class B 20%

Mid Development 1 Closed

Upper Layer Lifeform

- ☐ Herbaceous
☒ Shrub
☐ Tree

Indicator Species* and Canopy Position

SALIX Upper
ALNUS Upper
POBAB2 Upper
ROAC Lower

Structure Data (for upper layer lifeform)

	Min	Max
Cover	Closed Shrub (> 75% shrub cover)	Closed Shrub (> 75% shrub cover)
Height	Low Shrub (20cm to 1.5m)	Tall Shrub (>1.5m)
Tree Size Class	Seedling/Sapling <5"	

☐ Upper layer lifeform differs from dominant lifeform.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Description

5-29yrs

Tall shrubs (*Salix* spp., *Alnus* spp.) and saplings with a closed canopy (>60%). Saplings may consist of balsam poplar with white spruce in the understory (succession to class C), or saplings may consist of pure, even-aged spruce (succession to class E). Saplings overtop shrubs at 20-40yrs, when shade-intolerant pioneer shrub species decline and shade-tolerant shrubs (*Rosa acicularis* (prickly rose), *Viburnum edule* (high bushcranberry)) become more common and have a canopy cover of 10%.

Succession to class C or E. The deterministic succession pathway is to class C. The alternate succession pathway to E (probability = 0.01) represents possibility that white spruce will germinate in large numbers on mineral soil after flooding, resulting in a dense, even-aged stand. Flooding (probability = 0.03), represented by Option 1 in the model, causes a transition to class A.

Class C 40 %**Structure Data (for upper layer lifeform)**

Mid Development 1 Open

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☒ Tree

Indicator Species* and Canopy Position

POBAB2 Upper
 PIGL Middle
 ROAC Lower
 VIED Lower

Min		Max
Cover	Closed (60-100% tree cover)	Closed (60-100% tree cover)
Height	Tree (> 3m)	Tree (> 3m)
Tree Size Class	Pole 5-9" (swd)/5-11" (hwd)	

☐ Upper layer lifeform differs from dominant lifeform.

For mapping, absence of spruce in the overstory distinguishes Class C from Class E.

Description

30-149yrs

Closed balsam poplar forest. Balsam poplar is the dominant overstory species but white spruce is commonly in the understory. Shade-tolerant shrub species persist in the understory. If spruce is present, at approximately 100-150yrs the transition from balsam poplar to white spruce dominance begins (succession to class D). If white spruce is not present poplar persists, the stand ages and individual trees are lost to wind, disease or rot. Shrub cover commonly increases as the overstory canopy declines.

Succession to class D. Flooding (probability = 0.01), represented by Option 1 in the model, causes a transition to class A. Mixed fire (MFRI = 400yrs) maintains this class.

Class D 25 %**Structure Data (for upper layer lifeform)**

Late Development 1 Open

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☒ Tree

Indicator Species* and Canopy Position

PIGL Upper
 ROAC Lower
 VIED Lower
 Lower

Min		Max
Cover	Open (25-59% tree cover)	Open (25-59% tree cover)
Height	Tree (> 3m)	Tree (> 3m)
Tree Size Class	Pole 5-9" (swd)/5-11" (hwd)	

☐ Upper layer lifeform differs from dominant lifeform.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Description

150yrs+

Open white spruce forest. Spruce gains dominance over poplar and a mixed age, open stand develops. If enough young spruce establishes as poplar declines, the canopy closes again (modeled as alternate succession to class E). Alternatively, the stand may remain open with shrubs in the understory.

This class can persist in the absence of disturbance or follows an alternate succession pathway to E (probability = 0.005). Flooding (probability = 0.005), represented by Option 1 in the model, causes a transition to class A. Mixed fire (MFRI = 155yrs) causes a transition to class C.

Class E 10%

Structure Data (for upper layer lifeform)

Late Development 1 Closed

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☒ Tree

Indicator Species* and Canopy Position

PIGL Lower
ROAC Lower
VIED Lower
Lower

	Min	Max
	Cover	Cover
Cover	Closed (60-100% tree cover)	Closed (60-100% tree cover)
Height	Tree (> 3m)	Tree (> 3m)
Tree Size Class	Med. 9–20" (swd)/11–20" (hwd)	

- ☐ Upper layer lifeform differs from dominant lifeform.

For mapping, dominance of spruce in the overstory distinguishes Class E from Class C.

Description

30yrs+

Closed white spruce forest. These stands may be even-aged (resulting from spruce establishment on mineral soil after a flood event (succession from class B) or mixed age (succession from class D). If succession is from class D, occasional mature balsam poplar may persist in the overstory. As the spruce canopy closes, feathermoss becomes dominant on the forest floor, reaching 80% cover. *Rosa acicularis*, *Viburnum edule* and *Alnus* spp. may be scattered in the stand. A low shrub and herb layer may also occupy the forest floor.

This class will persist in the absence of disturbance. Flooding (probability = 0.002), represented by Option 1 in the model, causes a transition to class A. Mixed fire (MFRI = 200yrs) causes a transition to class D. Replacement fire (MFRI = 667yrs) causes a transition to class A. Insects/disease (probability = 0.02) can cause a transition to class D.

Disturbances

Fire Regime Group:** V

Historical Fire Size (acres)

Avg 0
Min 0
Max 0

Sources of Fire Regime Data

- ☒ Literature
☐ Local Data
☒ Expert Estimate

Fire Intervals

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
Replacement	5000			0.0002	6
Mixed	312			0.00321	94
Surface					
All Fires	294			0.00342	

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Additional Disturbances Modeled

- ☒ Insects/Disease ☐ Native Grazing ☒ Other (optional 1) Flooding
☐ Wind/Weather/Stress ☐ Competition ☐ Other (optional 2)

References

Barney, R.J. 1971. Wildfires in Alaska – some historical and projected effects and aspects. In: Fire in the Northern Environment – a symposium [Fairbanks, Alaska]. P. 51-59.

Boggs, K. 2000. Classification of community types, successional sequences, and landscapes of the Copper River Delta, Alaska. Gen Tech. Rep. PNW-GTR-469. Portland, OR: US Dept. Ag. Forest Service, Pacific Northwest Research Station. 244p.

Duchesne L.C. and B.C.Hawkes. 2000. Fire in northern ecosystems. In: Brown, J.K. and J.K Smith (eds.) Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol 2. Ogdon, UT: USDA Forest Service, Rocky Mountain Research Station. 257 p.

Heinselman, M.L. 1981. Fire and succession in the conifer forests of northern North America. In: West, D.C., H.H. Shugart, and D.B. Botkin. Forest succession: concepts and application. Springer-Verlag, New York. Chapter 23.

Murphy, K.A. and E. Witten. 2006. Riparian Spruce Hardwood. In Fire Regime Condition Class (FRCC) Interagency Guidebook Reference Conditions. Available at www.frcc.gov.

NatureServe. 2008. International Ecological Classification Standard: Terrestrial Ecological Classifications. Draft Ecological Systems Description for Alaska Boreal and Sub-boreal Regions.

Personal communication experts' workshop, March 2-4 2004. Fire Regime Condition Class (FRCC) interagency experts' workshop to develop and review Potential Natural Vegetation (PNV) groups for Alaska. Anchorage, AK.

Rowe, J.S., Bergsteinsson, J. L. , Padbury, G. A., and R. Hermesh. 1974. Fire studies in the Mackenzie Valley. ALUR Report 73-74-61. Arctic Land Use Research Program, Department of Indian Affairs and Human Development, Ottawa, Canada. 123 p.

Scott, R.W. 1974. Successional patterns on moraines and outwash of the Frederika Glacier, Alaska. In: Bushnell, V.C.; Marcus, M.G., eds. Icefield ranges research project scientific results. New York: American Geographical Society: 319-329. Vol. 4.

Shephard, M. E. 1995. Plant community ecology and classification of the Yakutat Foreland, Alaska. R10-TP-56.

Thilenius, J.F. 1990. Woody plant succession on earthquake-uplifted coastal wetlands of the Copper River Delta, Alaska. Forest Ecology and Management 33/34: 439-462.

Viereck L.A. 1966. Plant succession and soil development on gravel outwash on the Muldrow Glacier, Alaska. Ecological Monographs. 36(3):181-199.

Viereck, L. A. 1973. Ecological effects of river flooding and forest fires on permafrost in the taiga of

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Alaska. In: Permafrost - - The North American Contribution to the Second International Conference.
National Academy of Sciences, Washington, DC. 60-67 p.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.
**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-
year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity