

LANDFIRE Biophysical Setting Model

Biophysical Setting 7516590

**Alaskan Pacific Maritime Mountain Hemlock
Peatland**

☒ This BPS is lumped with: see below

☐ This BPS is split into multiple models: Alaskan Pacific Maritime Dwarf-Shrub-Sphagnum Peatland. Dwarf Shrub Sphagnum Peatland and Mountain Hemlock Peatland often occur in a mosaic especially at higher elevations.

General Information

Contributors (also see the Comments field)

Date 9/15/2008

Modeler 1 Rick Turner rlturmer@fs.fed.us

Reviewer Tom DeMeo tdemeo@fs.fed.us

Modeler 2

Reviewer Tina Boucher antvb@uaa.alaska.edu

Modeler 3

Reviewer

Vegetation Type

Wetlands/Riparian

Map Zone

75

Model Zone

- | | |
|--|--|
| ☒ Alaska | ☐ N-Cent.Rockies |
| ☐ California | ☐ Pacific Northwest |
| ☐ Great Basin | ☐ South Central |
| ☐ Great Lakes | ☐ Southeast |
| ☐ Northeast | ☐ S. Appalachians |
| ☐ Northern Plains | ☐ Southwest |

Dominant Species*

TSME ANPO
VAUL SPHAG2
ELPY NECR2
EMNI CAAN10

General Model Sources

- ☒ Literature
☐ Local Data
☒ Expert Estimate

Geographic Range

This system is found from Kenai Fjords through southeast AK.

Biophysical Site Description

This peatland system typically occurs on sloping terrain (10-50% slopes) and may develop on fairly steep sideslopes in areas with very high rainfall and low permeability (such as Prince William Sound and Kenai Fjords), but can also occur in basins or on flat terrain (NatureServe 2008). Soils are acidic and are usually saturated throughout the growing season (NatureServe 2008). This system occurs from sea level to treeline in the northern portion of the range (Kenai Fjords to Glacier Bay) and at higher elevations (usually above 500m, Banner et al. 1988) in the southern portion of its range (southeast AK and BC).

Vegetation Description

This type is often a mosaic of forested and non-forested peatlands. Stunted forests of *Tsuga mertensiana* (more common), *Cupressus nootkatensis* or *Picea sitchensis*, with *Elliottia pyroliflora* and *Vaccinium ovalifolium* are interspersed with dwarf shrub and herbaceous associations featuring *Empetrum nigrum*, *Vaccinium uliginosum*, *Vaccinium caespitosum*, *Phyllodoce glanduliflora*, *Andromeda polifolia*, *Kalmia polifolia* and *Oxycoccus microcarpus*. Common herbaceous species include *Nephrophyllidium crista-galli*, *Caltha leptosepala*, *Trichophorum caespitosum*, *Dodecatheon pulchellum*, *Drosera* spp., *Geum calthifolium*, *Cornus canadensis*, *Carex pauciflora*, *C. anthoxanthea*, *C. livida*, *C. pluriflora*, *C. sitchensis* and *Eriophorum angustifolium*. *Sphagnum* spp. (especially *S. fuscum*) is usually abundant in the ground layer (DeVelice et al. 1999, Boggs et al. in prep.). Other bryophytes include *Rhytidiadelphus loreus*, *Dicranum* spp. and *Pleurozium schreberi*.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Disturbance Description

The successional dynamics of this system are unknown but it is likely that this system is stable over long time periods (Vioreck et al. 1992, II.A.2.b). Fluctuations in the water table may affect this system over time. An occasional wind event may affect this system but its effect would be minimal.

Adjacency or Identification Concerns

This BpS may be found adjacent to ponds with floating, emergent vegetation or sedge dominated fens. It is also found with in the forest mosaic.

Native Uncharacteristic Conditions

Scale Description

Small patch

Issues/Problems

Comments

This system was created for the maritime region and did not receive review for other regions in the state.

This model was based on input from experts who attended the LANDFIRE Juneau Modeling Meeting (Feb. 08) and refined by Rick Turner. Review comments resulted in minor edits to the Biophysical Site and Vegetation Description sections to encompass the expression of this system in the northern part of its range.

Vegetation Classes

Class A 100 %

Mid Development 1 All Structures

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☒ Tree

Indicator Species* and Canopy Position

TSME Upper
TRCA30 Lower
NECR2 Lower
SPHAG2 Lower

Structure Data (for upper layer lifeform)

	Min	Max
Cover	Woodland (10-24% tree cover)	Open (25-59% tree cover)
Height	Dwarf Tree (< 3 m)	Tree (> 3 m)
Tree Size Class	Pole 5–9" (swd)/5–11" (hwd)	

☐ Upper layer lifeform differs from dominant lifeform.

Description

Zero years plus

This class represents the Mountain Hemlock Peatland BpS. See the Vegetation Description for a list of common species.

This community will persist under appropriate hydrological conditions. No disturbances were modeled.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Class B 0 %

[Not Used] [Not Used]

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and Canopy Position**Structure Data (for upper layer lifeform)**

	Min	Max
Cover		
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.**Description****Class C 0 %**

[Not Used] [Not Used]

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and Canopy Position**Structure Data (for upper layer lifeform)**

	Min	Max
Cover		
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.**Description****Class D 0 %**

[Not Used] [Not Used]

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and Canopy Position**Structure Data (for upper layer lifeform)**

	Min	Max
Cover		
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.**Description****Class E 0 %**

[Not Used] [Not Used]

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and Canopy Position**Structure Data (for upper layer lifeform)**

	Min	Max
Cover		
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.**Description****Disturbances**

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Fire Regime Group:** NA

Historical Fire Size (acres)

Avg 0

Min 0

Max 0

Sources of Fire Regime Data

- ☐ Literature
☐ Local Data
☒ Expert Estimate

Additional Disturbances Modeled

- ☐ Insects/Disease ☐ Native Grazing ☐ Other (optional 1)
☐ Wind/Weather/Stress ☐ Competition ☐ Other (optional 2)

Fire Intervals

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
Replacement					
Mixed					
Surface					
All Fires					

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class.

References

Banner, A., Hebda, R.J., Oswald, E.T., Pojar, J. and Trowbridge, R. 1988. Wetlands of Pacific Canada. In: Wetlands of Canada. National Wetlands Working Group. Ecological Land Classification Series No. 24. Environment Canada, Ottawa, and Polyscience Publications Inc., Montreal. 452 pp.

Boggs, K. et al. (in preparation). Kenai Fjords User's Guide.

NatureServe. 2008. International Ecological Classification Standard: Terrestrial Ecological Classifications. Draft Ecological Systems Description for the Alaska Maritime Region.

DeVelice, R., C. Hubbard, K. Boggs, S. Boudreau, M. Potkin, T. Boucher and C. Wertheim. 1999. Plant community types of the Chugach National Forest: Southcentral Alaska. U.S.D.A., Forest Service: Chugach National Forest, Alaska Region, Anchorage, Alaska. Tech. Pub. R10-TP-76. 375 pp.

Viereck, L., Dryness, C., Batten A. and K. Wenzlick. 1992. The Alaska vegetation classification. General Technical Report PNW-GTR-286. Portland, Oregon. USDA Forest Service, Pacific Northwest Research Station. 278 pp.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.