

LANDFIRE Biophysical Setting Model

Biophysical Setting 7716810

**Alaskan Pacific Maritime Poorly Drained
Conifer Woodland**

- ☐ This BPS is lumped with:
☐ This BPS is split into multiple models:

General Information

Contributors (also see the Comments field)

Date 7/30/2008

Modeler 1 Tom DeMeo

tdemeo@fs.fed.us

Reviewer Paul Alaback

paul.alaback@umontana.edu

Modeler 2

Reviewer Sheila Spores

sspores@fs.fed.us

Modeler 3

Reviewer Sue Kesti

skesti@fs.fed.us

Vegetation Type

Forest and Woodland

Map Zone

77

Model Zone

- | | |
|--|--|
| ☒ Alaska | ☐ N-Cent.Rockies |
| ☐ California | ☐ Pacific Northwest |
| ☐ Great Basin | ☐ South Central |
| ☐ Great Lakes | ☐ Southeast |
| ☐ Northeast | ☐ S. Appalachians |
| ☐ Northern Plains | ☐ Southwest |

Dominant Species*

TSHE NECR2
TSME ELPY
THPL LYAM3
CHNO VAOV

General Model Sources

- ☒ Literature
☐ Local Data
☒ Expert Estimate

Geographic Range

This type is found in the subpolar and perhumid rainforest regions (defined by Alaback 1991, 1995 of the AK maritime regions from Kenai Fjords and Prince William Sound to Yakutat, and south through southeastern AK.

Biophysical Site Description

In the perhumid rainforest region this type is common on rolling terrain, benches, and gentle slopes with restricted drainage. Soils are poorly drained and usually have a thick organic layer. In the subpolar rainforest region this system occurs on low to mid elevations on poorly drained sites. Soils may be shallow to deep and usually have at least some peat development. In some places, stands are often a fine mosaic of peatlands and better-drained inclusions. These are low-productivity sites that are intermediate between shore pine or mountain hemlock peatland sites and productive western hemlock - western Red-cedar sites.

The poorly drained conifer community includes the concept of Western hemlock small tree old-growth (see Caouette and DeGayner 2008). The Western Hemlock BpS represents the western hemlock large tree old-growth found on more productive sites.

Vegetation Description

Open canopy (typically <45% canopy cover) conifer stands tend to dominate these low productivity sites. Trees often show signs of stress such as spike-top or chlorotic foliage (especially spruce) and standing dead trees are common. Average overstory tree height ranges from 18-21m (DeMeo et al. 1992) and dbh averages 25-50cm (Viereck et al. 1992). Overstory trees may include several of the following species:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Tsuga heterophylla, *T. mertensiana*, *Thuja plicata* (found only in the southern most portion of the range of this BpS) and *Chamaecyparis nootkatensis* (= *Cupressus nootkatensis*). *Picea sitchensis* and *Pinus contorta* may also be present, but are not dominant and are nearly always stunted. *T. mertensiana* becomes the dominant overstory species in the northern portion of the maritime region (subpolar rainforest) while most other species, except *Picea sitchensis*, drop out. Common understory species include *Nephrophyllidium crista-galli*, *Elliottia pyroliflorus*, *Gaultheria shallon* (found only in the southern most portion of the range of this BpS), *Lysichiton americanus*, *Vaccinium ovalifolium*, (Demeo et al. 1992, Martin et al. 1995, Shephard et al. 1995). Further north where Mt. hemlock dominates common understory species include *Vaccinium ovalifolium*, *Elliottia pyroliflora*, *Nephrophyllidium crista-galli*, *Thelypteris quelpaertensis*, *Phegopteris connectilis*, *Trichophorum caespitosum*, *Carex anthoxantha*, *C. pluriflora*, *C. stylosa*, *Eriophorum* spp., *Lysichiton americanus* and *Sphagnum* spp.

Disturbance Description

According to Viereck et al. (1992) secondary succession in this type is largely unknown because disturbance is rare. However, Paul Hennon offered the following information related to the successional dynamics of this system:

Short term succession occurs from single tree mortality from a variety of causes [Note: this is not included in the VDDT model because these small scale dynamics can not be mapped into seral stages by LANDFIRE; therefore, the model is represented by a single seral stage]. Tree regeneration frequently occurs on raised organic microsites on the remains of previous trees. Tree growth is generally very slow. Longer term succession is probably influenced by climatic patterns that dictate drainage, either favoring poorer drainage, increased tree mortality, and more open canopy; or, improved drainage, greater tree growth, and a more closed canopy. These patterns can also favor individual tree species based on their tolerance or intolerance of wet soils.

There is some anecdotal evidence of anthropogenic fire in southern southeast AK but it remains largely unstudied. Fire may become a more important factor in this type as climate warms [Note: a reviewer commented that the effects of climate warming are still uncertain and that the assumption of increased fire activity in some areas, particularly Prince William Sound, may not be well founded]. The widespread yellow-cedar decline, which covers 200,000 ha of this type in southeast AK, is an example of a climate-induced tree death that has resulted in a composition shift away from yellow-cedar due to this mortality (Hennon et al. 2008).

Adjacency or Identification Concerns

This type is found adjacent to low productivity shore pine peatland and higher productivity western hemlock-western red cedar communities.

This BpS is referred to as mixed conifer in the SE AK plant association guides.

Native Uncharacteristic Conditions

This type is sensitive to changes in climate and hydrology. Neiland (1971) documents the shift between forest and bog noting that climate and surface physical features determine the vegetation on a site at a particular time.

Scale Description

Matrix

Issues/Problems

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Comments

This model was developed by Tom DeMeo for southeast AK. Review comments resulted in minor descriptive changes to the general description and expanding the description to include poorly drained Mt. hemlock sites that occur in the northern portion of the maritime region (subpolar rainforest). Comments made by Paul Hennon related to the successional dynamics of this system were added to the Disturbance Description. Patricia Krosse (pkrosse@fs.fed.us) also provided review comments on this model.

Vegetation Classes

Class A 100 %

Mid Development 1 All Structures

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☒ Tree

Indicator Species* and Canopy Position

TSHE Upper
TSME Upper
THPL Upper
CHNO Upper

Structure Data (for upper layer lifeform)

	Min	Max
Cover	Open (25-59% tree cover)	Open (25-59% tree cover)
Height	Dwarf Tree (< 3 m)	Tree (> 3 m)
Tree Size Class	Med. 9–20" (swd)/11–20" (hwd)	

☐ Upper layer lifeform differs from dominant lifeform.

Description

Zero plus years

This community is characterized by an open canopy and may include several of the following species: Tsuga heterophylla, T. mertensiana, Thuja plicata and Cupressus nootkatensis. Picea sitchensis and Pinus contorta may also be present, but are not dominant.

No disturbances were modeled.

Class B 0 %

[Not Used] [Not Used]

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover		
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.

Description

Class C 0 %

[Not Used] [Not Used]

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover		
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Description

Class D 0 %

[Not Used] [Not Used]

Structure Data (for upper layer lifeform)

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and Canopy Position

	Min	Max
Cover		
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.

Description

Class E 0 %

[Not Used] [Not Used]

Structure Data (for upper layer lifeform)

Upper Layer Lifeform

- ☐ Herbaceous
☐ Shrub
☐ Tree

Indicator Species* and Canopy Position

	Min	Max
Cover		
Height		
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.

Description

Disturbances

Fire Regime Group:** NA

Historical Fire Size (acres)

Avg 0
Min 0
Max 0

Sources of Fire Regime Data

- ☒ Literature
☐ Local Data
☒ Expert Estimate

Additional Disturbances Modeled

- ☐ Insects/Disease ☐ Native Grazing ☐ Other (optional 1)
☐ Wind/Weather/Stress ☐ Competition ☐ Other (optional 2)

Fire Intervals

Avg FI	Min FI	Max FI	Probability	Percent of All Fires
Replacement				
Mixed				
Surface				
All Fires				

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class.

References

Alaback, P.B. 1991. Comparative ecology of temperate rainforests of the Americas along analogous climatic gradients. Rev. Chil. Hist. Nat. 64: 399–412.

Alaback, P.B. 1995. Biodiversity patterns in relation to climate and a genetic base for the rainforests of the west coast of North America. In R. Lawford, P.B. Alaback, and E.R. Fuentes, eds. 1995. High Latitude Rain Forests of the West Coast of the Americas: Climate, Hydrology, Ecology and Conservation. Berlin: Springer-Verlag.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity

Caouette, J.P. and E.J. DeGayner. 2008. Broad-Scale Classification and Mapping of Tree Size and Density Attributes in Productive Old-Growth Forests in Southeast Alaska's Tongass National Forest. *Western Journal of Applied Forestry*. 23(2): 106-112.

DeMeo, T., J. Martin and R.A. West. 1992. Forest plant association management guide, Ketchikan Area, Tongass National Forest. United States Dept. of Agriculture, Forest Service, Alaska Region. R10-MB-210. 405p.

Hennon, P.E.; D'Amore, D.; Wittwer, D.; Caouette, J. 2008. Yellow-cedar decline: conserving a climate-sensitive tree species as Alaska warms. In: Deal, R., ed. Integrated restoration of forested ecosystems to achieve multiresource benefits: proceedings of the 2007 national silviculture workshop. Gen. Tech. Rep. PNW-GTR-733. Portland, OR: USDA Forest Service, Pacific Northwest Research Station: 233-245.

Martin, Jon R., Susan J. Trull, Ward W. Brady, Randolph A. West and Jim M. Downs. 1995. Forest plant association management guide, Chatham Area, Tongass National Forest. USDA Forest Service, Alaska Region. R10-TP-57.

NatureServe. 2008. International Ecological Classification Standard: Terrestrial Ecological Classifications. Draft Ecological Systems Description for the Alaska Maritime Region.

Neiland, Bonita J. 1971. The forest-bog complex of southeast Alaska. *Vegetatio*. 22: 1-64.

Shephard, M.E. 1995. Plant community ecology and classification of the Yakutat Foreland, Alaska. Technical Report R10-TP-56. Juneau, AK: USDA Forest Service, Alaska Region. 206 p.

Viereck, L., Dryness, C., Batten A. and K. Wenzlick. 1992. The Alaska vegetation classification. General Technical Report PNW-GTR-286. Portland, Oregon. U.S. Forest Service, Pacific Northwest Research Station. 278 p.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100-year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.